

Mathematical Physics

B.D.Gupta

Scanned Version

- 1.1. Introduction
- 1.2. Representation of vectors
- 1.3. Kinds of vectors
- 1.4. Addition of vectors
- 1.5. Subtraction of vectors
- 1.6. Multiplication of a vector by a scalar
- 1.7. Vector space or linear space
- 1.8. Conditions for a physical quantity to be representable by a vector
- 1.9. Resolution of vectors
- 1.10. Linear combination of vectors
- 1.11. Product of two vectors
- 1.12. Triple products of vectors
- 1.13. Product of four vectors
- 1.14. Reciprocal system of vectors
- 1.15. Vector equations
- 1.16. Simple applications of vectors to mechanics
- 1.17. Differentiation of vectors
- 1.18. Some rules for differentiation
- 1.19. Partial differentiation of vectors
- 1.20. Rules for partial differentiation of vectors
- 1.21. The scalar and vector fields
- 1.22. Directional derivatives
- 1.23. Level surfaces
- 1.24. The gradient of a scalar field
- 1.25. The gradient of a scalar-point function
- 1.26. The gradient of sum of two scalar-point functions
- 1.27. The gradient of product of two scalar-point functions
- 1.28. The divergence of a vector-point function
- 1.29. The divergence of sum of two vector functions
- 1.30. The divergence of product
- 1.31. The curl or rotation of a vector-point function
- 1.32. Curl of the sum of two vector-point functions
- 1.33. Curl of the product of two vector-point functions
- 1.34. To express gradient of scalar product in terms of curl

- 1.35. To express divergence of vector products in terms of curl
 - 1.36. Curvilinear co-ordinates
 - 1.37. Orthogonal curvilinear co-ordinates.
 - 1.38. Condition for orthogonality
 - 1.39. Reciprocal sets of two triads of mutually orthogonal vectors
 - 1.40. Gradient in terms of orthogonal curvilinear coordinates
 - 1.41. Divergence in terms of orthogonal curvilinear coordinates
 - 1.42. Curl in terms of orthogonal curvilinear coordinates
 - 1.43. Laplacian (∇^2) in terms of orthogonal curvilinear coordinates
 - 1.44. Equivalent expressions for $\nabla \phi$, $\nabla \cdot F$ and $\nabla \times F$ in rectangular coordinates
 - 1.45. Cylindrical coordinates as a special curvilinear system
 - 1.46. Spherical polar coordinates as a special curvilinear system
 - 1.47. Integration of a vector
 - 1.48. The line integral
 - 1.49. The surface integrals
 - 1.50. The volume integrals
 - 1.51. Gauss' divergence theorem
 - 1.52. Deductions from Gauss' theorem
 - 1.53. Physical interpretation of Gauss' divergence theorem
 - 1.54. Gauss' theorem
 - 1.55. Two-Green's identities
 - 1.56. Green's theorem in the plane
 - 1.57. Vector forms of Green's theorem in the plane
 - 1.58. Green's formula
 - 1.59. Poisson's equation with its solution
 - 1.60. Laplace's equation with its solution
 - 1.61. Stoke's theorem in space
 - 1.62. Some theorems
 - 1.63. The classification of vector fields
 - 1.64. Helmholtz's theorem
- Additional Miscellaneous Problems

2. **MATRICES**

- 2.1. Definitions and notations
- 2.2. Equality of matrices
- 2.3. Addition of matrices
- 2.4. Properties of matrix-addition
- 2.5. Multiplication of matrices
- 2.6. Properties of matrix-multiplication
- 2.7. Partitioning of matrices
- 2.8. Products of matrices by partitioning
- 2.9. Special matrices with their properties
- 2.10. Rank of a matrix
- 2.11. Some theorems on rank

- 2.12. Solutions of linear equations
- 2.13. Methods for solving linear equations
- 2.14. Characteristic matrix and characteristic equation of a matrix
- 2.15. Sub-spaces and null spaces
- 2.16. Transformations
- 2.17. Hermitian forms
- 2.18. Characteristic roots and vectors of a matrix (Eigen values and Eigen vectors)
- 2.19. Quadratic forms and their reductions
- 2.20. Differentiation and integration of matrices
- 2.21. Solution of a system of linear equations using differentiation of matrices
- 2.22. Other element methods of matrix inversion
- 2.23. The eigen-value problem
- Additional Miscellaneous Problems

3. TENSORS

- 3.1. Introduction
- 3.2. Transformation of coordinates
- 3.3. The summation convention and Kronecker delta symbol
- 3.4. Tensors as classification of transformation laws
- 3.5. Symmetric and anti-symmetric tensors
- 3.6. Invariant tensors
- 3.7. Rules which govern tensor analysis
- 3.8. The fundamental tensors
- 3.9. Associated tensors: Raising and lowering of suffixes
- 3.10. Length of a vector, (*i.e.*, a tensor of rank one), angle between two vectors and orthogonality of vectors
- 3.11. Metric tensor, Riemannian spaces
- 3.12. Christoffel's 3-index symbols
- 3.13. Equations of a geodesic
- 3.14. Law of transformation for Christoffel's symbols
- 3.15. Parallel displacement of vectors
- 3.16. Covariant derivative of a vector
- 3.17. Covariant derivative of a tensor
- 3.18. The curvature, tensor (Riemann-Christoffel tensor)
- 3.19. Riemann-Christoffel tensor or covariant curvature tensor
- 3.20. Some important results
- 3.21. Tensor forms of operators
- 3.22. Einstein gravitational equation
- Additional Miscellaneous Problems

4. GROUP THEORY

- 4.1. Introduction to sets, mappings and binary operations
- 4.2. Groups

- 4.3. Elementary properties of a group
- 4.4. Some definitions
- 4.5. The centre of a group
- 4.6. Cosets or cosets of a subgroup
- 4.7. Cyclic group
- 4.8. Permutation or transformation
- 4.9. Homomorphism and isomorphism of groups
- 4.10. Normals and conjugate subgroups
- 4.11. Complexes and kernel
- 4.12. Groups of isometries
- 4.13. Some special groups with linear operators
- 4.14. Permutational representations
- Additional Miscellaneous Problems

5. COMPLEX VARIABLES

- 5.1. Introduction
- 5.2. Definitions
- 5.3. Operation of fundamental laws of algebra on complex numbers
- 5.4. Graphical representation (Argand diagram)
- 5.5. Properties of moduli and arguments and geometry of complex numbers
- 5.6. Regular functions
- 5.7. Elementary functions and mapping by them
- 5.8. Complex integration
- 5.9. Cauchy's theorem
- 5.10. Cauchy's integral formula
- 5.11. Some results based on Cauchy's integral formula
- 5.12. Taylor's expansion
- 5.13. Laurent's expansion
- 5.14. Residue and contour integration
- 5.15. Cauchy's residue theorem
- 5.16. Another important theorem
- 5.17. Computation of residue
- 5.18. Integration round the unit circle
- 5.19. Evaluation of $\int_{-\infty}^{+\infty} f(x) dx$
- 5.20. Jordan's lemma to evaluate infinite integrals
- 5.21. Analytic continuation
- 5.22. Conformal mapping
- 5.23. Convergence of integrals, series and products
- Additional Miscellaneous Problems

6. BETA, GAMMA AND ERROR FUNCTIONS

- 6.1. Definitions
- 6.2. Fundamental property of gamma functions
- 6.3. The value of $\Gamma \frac{1}{2}$ and graph of the gamma function

- 6.4. Transformation of gamma function
- 6.5. To show that $\beta(m, n) = \beta(n, m)$
- 6.6. Different forms of Beta function
- 6.7. To find the relation between beta and gamma functions
- 6.8. Reduction of definite integrals to gamma functions
- 6.9. Miscellaneous propositions
- 6.10. Error function or probability integral
- 6.11. Factorial function
- 6.12. Some allied functions
- 6.13. Orthogonal sets of functions
- Additional Miscellaneous Problems

7. DIFFERENTIAL EQUATIONS

- 7.1. Introduction
- 7.2. Solution in simple cases of ordinary differential equations of second order
- 7.3. Simple problems from physics
- 7.4. Partial differential equations
- 7.5. Special types of differential equations arising in physics
- 7.6. Finite differences
- 7.7. Polynomial interpolation
- 7.8. Interpolation with equal intervals
(i.e. when function-values are given at equidistant intervals)
- 7.9. Interpolation with unequal intervals
- 7.10. Central difference formulae
- 7.11. Approximate integration
- 7.12. Perturbation method
- Additional Miscellaneous Problems

8. HARMONICS (WITH SPECIAL FUNCTIONS)

- 8.1. Introduction
- 8.2. Methods of integration in series
- 8.3. Legendre differential equation, functions and polynomials with properties
- 8.4. Bessel's equation, functions and polynomials
- 8.5. Hypergeometric or Gauss equation and functions
- 8.6. Confluent hypergeometric equations and functions
- 8.7. Hermite equation, functions and polynomials
- 8.8. Laguerre equation and polynomials with properties
- 8.9. Riccati's differential equation
- 8.10. The Dirac-delta function with its formal properties
- 8.11. Riemann-Zeta function
- Additional Miscellaneous Problems

9. FOURIER'S SERIES, INTEGRALS AND TRANSFORMS

- 9.1. Definition and expansion of a function of x
- 9.2. Dirichlet's conditions
- 9.3. Assumptions for the validity of Fourier's series expansion, with allied theorems
- 9.4. Complex representation of a Fourier's series
- 9.5. Parseval's identity for Fourier series
- 9.6. Fourier's integral
- 9.7. Different forms of Fourier's integrals
- 9.8. A remark on convergence of Fourier's series
- 9.9. Physical applications of Fourier's series
- 9.10. The Fourier transforms
- 9.11. Applications of Fourier transforms to boundary value problems
- Additional Miscellaneous Problems

10. THE LAPLACE TRANSFORMS

- 10.1. Introduction
- 10.2. Definition of the integral transform
- 10.3. Deduction of the definition of the Laplace transform from that of the integral transform
- 10.4. Definition of the Laplace transform
- 10.5. Functions of exponential order
- 10.6. Piecewise or sectional continuity
- 10.7. A function of class A
- 10.8. Sufficient conditions for the existence of Laplace transforms
- 10.9. Some properties of Laplace transforms
- 10.10. Some methods for finding Laplace transforms
- 10.11. The Laplace transforms of some special functions
- 10.12. Evaluation of integrals with the help of Laplace transforms
- 10.13. Laplace transform of the Laplace transform
- 10.14. The inverse Laplace transform
- 10.15. Some properties of inverse Laplace transform
- 10.16. Evaluation of some integrals by inverse Laplace transforms
- 10.17. Some methods for finding inverse Laplace transform
- 10.18. Applications of Laplace transforms to differential equations
- 10.19. Applications of Laplace transforms to boundary value problems
- 10.20. The complex inversion formula or inversion theorem
- Additional Miscellaneous Problems

11. HANKEL TRANSFORMS

- 11.1. Introduction
- 11.2. Definition of infinite Hankel transform
- 11.3. Inverse formula for Hankel transform
- 11.4. Parseval's theorem for Hankel transforms
- 11.5. Linearity property
- 11.6. Hankel transform of the derivatives of a function

- 11.7. Hankel transforms of $\frac{d^2 f}{dx^2}$, $\frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx}$ and $\frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} f$ under certain conditions
- 11.8. Applications to boundary value problems
- 11.9. Finite Hankel transforms
- 11.10. Finite Hankel transform of $\frac{df}{dx}$, p being root of $J_n(p) = 0$
- 11.11. Finite Hankel transform of $\frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx}$
- 11.12. Finite Hankel transform of $\frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} f(x)$ p being a root of $J_n(pa) = 0$
- Additional Miscellaneous Problems

12. DIFFUSION, WAVE AND LAPLACE'S EQUATIONS

- 12.1. Introduction
- 12.2. Diffusion equation of Fourier equation of heat flow
- 12.3. One-dimensional diffusion equation $\frac{\partial \mathbf{u}}{\partial t} = h^2 \frac{\partial^2 \mathbf{u}}{\partial x^2} = k \frac{\partial^2 \mathbf{u}}{\partial x^2}$
- 12.4. Two-dimensional diffusion equation *i.e.*, $\frac{\partial \mathbf{u}}{\partial t} = h^2 \left(\frac{\partial^2 \mathbf{u}}{\partial x^2} + \frac{\partial^2 \mathbf{u}}{\partial y^2} \right)$
- 12.5. Three-dimensional diffusion equation
- 12.6. Laplace's equation
- 12.7. Two-dimensional Laplace's equation (steady flow of heat)
- 12.8. Three-dimensional of Laplace's equation
- 12.9. General properties of Harmonic functions
- 12.10. The wave equations
- 12.11. Green's functions for the wave equation
- 12.12. Homogeneous and inhomogeneous wave equations
- 12.13. Theory of wave guides
- 12.14. Solution of one-dimensional wave equation
- 12.15. Square wave
- 12.16. Solution of two-dimensional wave equation
- 12.17. Solution of three-dimensional wave equation
- Additional Miscellaneous Problems

13. MAXWELL'S ELECTROMAGNETIC FIELD EQUATIONS

- 13.1. Introduction
- 13.2. Maxwell's equations for electromagnetic field
- 13.3. To show $\nabla \cdot \mathbf{D} = 4\pi p$
- 13.4. To show $\text{div } \mathbf{B} = 0$
- 13.5. To show $\text{curl } \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t}$

- 13.6. Physical Interpretation of Maxwell's Equations
- 13.7. Decaying of free charge
- 13.8. Poynting vector
- 13.9. Poynting theorem
- Additional Miscellaneous Problems

14. SPECIAL THEORY OF RELATIVITY

- 14.1. Introduction
- 14.2. Basic postulates of special theory of relativity
- 14.3. Lorentz transformations
- 14.4. The Kinematic effects of the Lorentz transformations
- 14.5. Energy-Momentum relation
- Additional Miscellaneous Problems

15. STATISTICAL PROBABILITY

- 15.1. Introduction
- 15.2. Definitions of probability
- 15.3. Events
- 15.4. Theorem of total probability, *i.e.*, additive law of probability
- 15.5. Theorem of compound probability or multiplicative law of probability
- 15.6. Probability: A new standpoint
- 15.7. Repeated trials: Binomial and multinomial expansions
- 15.8. Measures of central tendency
- 15.9. Measures of dispersion (or variation)
- 15.10. Moments
- 15.11. Expectation
- 15.12. Continuous univariate probability distribution
- 15.13. Tchebycheff's Inequality
- 15.14. Characteristic function
- 15.15. Fourier's inversion theorem
- 15.16. Theoretical distributions
- 15.17. Other distributions
- 15.18. The principle of least squares
- 15.19. Correlation and regression
- 15.20. Sampling distributions
- 15.21. Theory of errors
- Additional Miscellaneous Problems

APPENDICES

- A. Some formulated results in basic mathematics
- B. Asymptotic expansion of error function
- C. Character tables in group theory
- D. Approximate solutions of algebraic and transcendental equations

INDEX

CHAPTER 1

VECTORS

1.1. INTRODUCTION

It is generally observed that there exist two types of physical measurements in applied mathematics, physics and mechanics : one involving *only magnitude and no direction* in the space of three dimensions, such as volume, mass, length, speed, temperature, potential, electric charge etc., and the other involving *a definite direction in space associated with their magnitudes* such as velocity, acceleration, momentum, force, electric or magnetic field intensities etc., the former being called *scalar quantities* or simply *scalars* and the latter, *vector quantities* or simply *Vectors*.

A little consideration will exhibit that the complete characterization of a scalar quantity requires length and support, *i.e.*, a specified unit and a number stating how many times that unit is contained in that quantity, while the complete characterization of a vector quantity requires length, support and sense, *i.e.*, a specified unit, a number stating how many times that unit is contained in that quantity and the statement of the direction.

Stating in a precise manner a *vector* means '*a directed line segment*'. In other words we can state that a vector is a quantity having direction as well as magnitude. In Astronomy a vector means an imaginary straight line that joins a planet moving round a centre (generally the focus of an elliptic orbit) to that centre.

1.2. REPRESENTATION OF VECTORS

Since a vector is the result of abstraction, its magnitude and direction may be represented by a line OP directed from the *initial point* O to the *terminal point* P and denoted by

$\overrightarrow{OP}$. Here the length of vector OP denoted

by $|OP|$ is called *magnitude* or *module* or *modulus* of the vector and the direction in space is indicated by an arrow head on the line. In Fig. 1.1, the




Fig. 1.1

vector OP has been shown by $\overrightarrow{V}$ (or in Clarendon print by $\mathbf{V}$) while its scalar magnitude is stated by V . Thus OP is the length of the vector $\mathbf{V}$, while the line of indefinite length of which the directed

line segment OP is only a part is the support of $\mathbf{V}$ and the sense is from O to P .

It should be noted that formulation of a law of physics in terms of vectors is however independent of the choice of axes of reference, *i.e.*, the vector representation has a physical content without reference to any coordinate system.

1.3. KINDS OF VECTORS

Equal vectors. Two given vectors may be equal only when they have the same magnitude and the same direction, *i.e.*, the given two vectors are equal if and only if they have the same or parallel support with equal length and the same sense. For example in Fig. 1.1, we have

$$\vec{V} (=OP) = \vec{V}_1 (=O'P') = -\vec{V}_2 (=O''P'')$$

where $\vec{V}_1$ and $\vec{V}_2$ have the same scalar magnitude as $\vec{V}$ but $\vec{V}_1$ has the same and $\vec{V}_2$ the opposite sense to that of $\vec{V}$.

Null vector. A vector having the initial and the terminal points coincident is termed as a *zero vector* or a *null vector*. Thus a null vector has its module zero.

Unit vector. A vector having its modulus as unity is called a *unit vector*.

If $\vec{a}$ is a vector and ' a ' its modulus, then unit vector $\hat{a}$ is denoted by $\hat{a}$ (read as ' a hat' or ' a caret') defined as

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|} = \frac{\vec{a}}{a}$$

Polar vectors. The line vectors representing the quantities like force, velocity etc., in which merely a linear action in a particular direction is involved, are termed as *polar vectors*.

Axial vectors. The line vectors representing the quantities like angular velocity, angular acceleration etc., in which some rotational action is involved about an axis and which are drawn parallel to the axis of rotation in order that the magnitude of the quantity is determined by the length of the vector and the direction by the rule of right handed screw (*i.e.* rotation being considered in clockwise direction), are termed as *axial vectors*.

Free vector. Evidently a vector can be represented by an infinite number of equal vectors by drawing parallel supports. Such a vector which can be transported from place to place such that it remains of the same magnitude and keeps up the same direction is termed as a *free vector*. In fact a free vector is assumed to remain the same through transportation, irrespective of its position in space.

Localised or Line vector. We have defined that the value of a free vector depends only on its length and direction, but if it depends also on its position in space, *i.e.*, if a vector is restricted to pass through a given origin, then it is termed as a *localised vector*.

Collinear vectors. The vectors parallel to the same line, regardless of their magnitudes and sense of directions are termed as *collinear vectors*. In other words the vectors having the same or parallel supports are

known as collinear vectors. Such vectors are parallel to each other and they may coincide in a special case. As such there exists a scalar ratio say λ between any two collinear vectors $\vec{a}$ and $\vec{b}$ of the form

$$\vec{b} = \lambda \vec{a}$$

which follows that one of the two collinear vectors can be expressed as the scalar multiple of the other.

Non-collinear vectors. The vectors whose directions are neither parallel nor coincident are said to be *non-collinear*.

Like vectors or co-directional vectors. The vectors which are collinear and have the same sense of directions *i.e.*, the vectors directed in the same sense irrespective of their magnitudes are termed as *like vectors*.

Unlike vectors. The vectors which are collinear but have opposite sense of directions from each other are termed as *unlike vectors*.

Coplanar vectors. A system of vectors lying in the parallel planes or which can be made to lie in the same plane are said to be *coplanar vectors*. Evidently any two vectors are always coplanar.

Non-coplanar vectors. A system of vectors consisting of three or more vectors which cannot be made to lie in the same plane are called *non-coplanar vectors*.

Reciprocal vector. Any vector having its direction the same as that of a given vector $\vec{a}$, but its magnitude as the reciprocal of the magnitude of $\vec{a}$ is termed as the *reciprocal vector* of $\vec{a}$ and written as $\vec{a}^{-1}$ or $\frac{1}{a}$. As such

$$\vec{a}^{-1} = \frac{1}{a} \hat{a} = \frac{\vec{a}}{a^2} \hat{a} = \frac{\vec{a}}{a^2} \quad (\text{by definition of a unit vector}).$$

In this connection it is notable that the magnitude and so the reciprocal of the magnitude of a unit vector being unity, the unit vector is reciprocal to itself and it is said to be *self-reciprocal*.

Negative vector. The vector having the same magnitude as the vector $\vec{a}$ but opposite direction, is known as the *negative* of $\vec{a}$ and written as $-\vec{a}$.

Position vector. If a vector $\vec{OP}$ specifies the position of a point relative to an arbitrarily chosen point O , then $\vec{OP}$ is called the *Position vector* of P with respect to O , the origin of vectors.

Problem 1. If $\{\vec{a}, \vec{b}, \vec{c}\}$ is a right handed set, which of the following sets are right handed?

- (i) $\vec{a}, \vec{c}, \vec{b}$; (ii) $\vec{b}, \vec{c}, \vec{a}$; (iii) $\vec{b}, \vec{a}, \vec{c}$; (iv) $\vec{c}, \vec{a}, \vec{b}$; (v) $\vec{c}, \vec{b}, \vec{a}$

It is clear that the sets (ii) and (iv), *i.e.*, $\vec{b}, \vec{c}, \vec{a}$; and $\vec{c}, \vec{a}, \vec{b}$ are right handed.

Problem 2. Discuss the geometrical significance of $a\vec{A} + b\vec{B} = \vec{0}$. We have $a\vec{A} + b\vec{B} = \vec{0}$, a, b being scalars

This can be written as $\vec{A} = -\frac{b}{a}\vec{B}$

$$= \lambda \vec{B} \text{ if } \lambda = -\frac{b}{a}$$

i.e. $\vec{A}$ is expressible as a scalar multiple of $\vec{B}$ so that the vectors $\vec{A}$ and $\vec{B}$ are parallel or collinear.

1.4. ADDITION OF VECTORS

The characterisation of process of summation is inherited in the composition of two or more displacements of a point. Suppose that we have two vectors $\vec{a}$ and $\vec{b}$ acting at a point O as shown in Fig 1.2

Let $\vec{OA} = \vec{a}$ and $\vec{OB} = \vec{b}$.

Clearly the resultant effect of the vectors $\vec{a}$ and $\vec{b}$ is the same as that of their vector sum $\vec{v}$ obtained by setting off the vector $\vec{b}$ at the end of $\vec{a}$ and then joining the beginning of $\vec{a}$ to the end of $\vec{b}$. This geometrical construction utilised to find the vector sum of two vectors $\vec{a}$ and $\vec{b}$ is known as the *parallelogram law of addition of vectors*.




Fig. 1.2

A similar result follows by starting with $\vec{b}$ and setting off the vector $\vec{a}$ on $\vec{b}$, i.e.

$$\vec{v} = \vec{OC} = \vec{OB} + \vec{BC} = \vec{b} + \vec{a} \quad \dots (2)$$

Conclusively the result of adding two co-initial vectors is the vector represented by the diagonal of the parallelogram having the two given vectors as its adjacent sides.

From (1) and (2) it follows that

$$\vec{a} + \vec{b} = \vec{b} + \vec{a}$$

i.e., the two vectors obey the **commutative law of addition**, according to which the vector sum of two vectors is independent of their order.

We now propose to find the sum of three vectors say $\vec{a}$, $\vec{b}$, $\vec{c}$. Let $\vec{OA} = \vec{a}$, $\vec{AB} = \vec{b}$, $\vec{BC} = \vec{c}$ as shown in Fig. 1.3. Then

$$\vec{v} = \vec{OC} = \vec{OA} + \vec{AB} + \vec{BC} = \vec{a} + \vec{b} + \vec{c} \quad \dots (3)$$

Also $\vec{v} = \vec{OC} = \vec{OB} + \vec{BC}$




Fig. 1.3

$$\begin{aligned} &= (\vec{OA} + \vec{AB}) + (\vec{BC}) \\ &= (\vec{a} + \vec{b}) + \vec{c} \end{aligned} \quad \dots (4)$$

Similarly $\vec{v} = \vec{a} + (\vec{b} + \vec{c}) \quad \dots (5)$

and $\vec{v} = (\vec{a} + \vec{c}) + \vec{b} \quad \dots (6)$

It follows from (3), (4), (5) and (6) that

$$\vec{v} = \vec{a} + \vec{b} + \vec{c} = (\vec{a} + \vec{b}) + \vec{c} = \vec{a} + (\vec{b} + \vec{c}) = (\vec{a} + \vec{c}) + \vec{b}$$

i.e. the three vectors obey the **associative law of addition**, according to which the vector sum of three vectors is independent of the mode in which component vectors are associated in different groups.

In general, if there are n vectors $\vec{a}$, $\vec{b}$, $\vec{c}$, ..., $\vec{n}$, then their vector sum $\vec{v}$ is given by

$$\vec{v} = \vec{a} + \vec{b} + \vec{c} + \dots + \vec{n}$$

1.5. SUBTRACTION OF VECTORS

If there are two vectors $\vec{a}$ and $\vec{b}$, then

$$\vec{a} - \vec{b} = \vec{a} + (-\vec{b}),$$

i.e. the subtraction of $\vec{b}$ from $\vec{a}$ may be regarded as the addition of $-\vec{b}$ to $\vec{a}$. Thus to subtract $\vec{b}$ from $\vec{a}$, reverse the direction of $\vec{b}$ and add to $\vec{a}$, (Fig. 1.4).




Fig. 1.4

1.6. MULTIPLICATION OF A VECTOR BY A SCALAR

If $\vec{a}$ be any given vector and s a given scalar, then $s\vec{a}$ or $\vec{a}s$ is defined as a vector whose magnitude is $|s|$ times the magnitude of the vector $\vec{a}$, i.e. $|s|$ times the length of $\vec{a}$, the support being the same or parallel to that of $\vec{a}$ and direction being the same or opposite to that of $\vec{a}$, according as s is positive or negative. We thus have

(i) $s(-\vec{a}) = (-s)\vec{a} = -s\vec{a}$.

(ii) $(-s)(-\vec{a}) = s\vec{a}$.

(iii) $(s+t)\vec{a} = s\vec{a} + t\vec{a}$, t being another scalar.

(iv) $(st)\vec{a} = s(t\vec{a}) = t(s\vec{a})$.

(v) $0\vec{a} = \vec{0}$, $\vec{0}$ being the null vector.

(vi) If two non-zero vectors $\vec{a}$ and $\vec{b}$ are collinear, then there exists a non-zero scalar m , such that

$$\vec{a} = m\vec{b}.$$

Conversely the relation of this type implies that $\vec{b}$ is parallel or collinear to $\vec{a}$.

(vii) If $\hat{a}$ is the unit vector co-directional with $\vec{a}$ while $a = |\vec{a}|$, then

$$\vec{a} = a\hat{a} \text{ or } s\vec{a} = s(a\hat{a}) = sa\hat{a}.$$

Also as $\hat{a} = \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{\mathbf{a}}{a}$ and if $\mathbf{b}$ is parallel to $\mathbf{a}$, then $\mathbf{b} = \pm b \frac{\mathbf{a}}{a}$

according as $\mathbf{b}$ and $\mathbf{a}$ have the same or opposite directions.

Note. Division of a vector $\mathbf{a}$ by a non-zero scalar s is regarded as the multiplication of the vector $\mathbf{a}$ by a scalar $1/s$.

Problem 3. If there are two vectors $\mathbf{a}$ and $\mathbf{b}$ represented by the sides AB and BC of a triangle, then show that their resultant is represented by the third side AC . Why is this method equivalent to the parallelogram law of addition?




Fig. 1.5

As shown in Fig. 1.5, the vectors $\mathbf{a}$ and $\mathbf{b}$ are represented by the sides AB and

BC of the triangle. Here $\vec{AC}$ is a vector drawn between the initial point of $\mathbf{a}$ and terminal point of $\mathbf{b}$ and thus may be obtained by parallelogram law of addition, for if we complete the parallelo-

gram $ABCD$, then $\vec{AC}$ represents a vector along the diagonal of the parallelogram and passing through the common point of the adjacent sides AB and AD representing the vectors $\mathbf{a}$ and $\mathbf{b}$. As such the vector addition obeys the parallelogram law of forces.

Problem 4. What vector must be added to the two vectors $\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$ and $2\mathbf{i} + \mathbf{j} - \mathbf{k}$, so that the resultant may be a unit vector along the x -axis?

Suppose that

$$\mathbf{a} = \mathbf{i} - 2\mathbf{j} + 2\mathbf{k} \text{ and } \mathbf{b} = 2\mathbf{i} + \mathbf{j} - \mathbf{k}.$$

Then

$$\mathbf{a} + \mathbf{b} = 3\mathbf{i} - \mathbf{j} + \mathbf{k}.$$

Hence, in order that the resultant of $\mathbf{a}$ and $\mathbf{b}$, i.e. $\mathbf{a} + \mathbf{b}$ be $\mathbf{i}$, we have to add a vector.

$$\mathbf{i} - (3\mathbf{i} - \mathbf{j} + \mathbf{k}), \text{ i.e., } -2\mathbf{i} + \mathbf{j} - \mathbf{k}.$$

1.7. VECTOR SPACE OR LINEAR SPACE

A vector space (or linear space) over a field F is a set V of elements called vectors which may be combined by two primary operations—addition and scalar multiplication; such that

(A) (i) If the vectors $\mathbf{a}$ and $\mathbf{b}$ belong to V , then $\mathbf{a} + \mathbf{b}$ also belongs to V . (This is known as *closure property*.)

(ii) The vector sum of two vectors $\mathbf{a}$ and $\mathbf{b}$ belonging to V , is commutative, i.e.

$$\mathbf{a} + \mathbf{b} = \mathbf{b} + \mathbf{a}.$$

(iii) The vector sum of three vectors $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$ belonging to V , is associative, i.e.

$$\mathbf{a} + \mathbf{b} + \mathbf{c} = (\mathbf{a} + \mathbf{b}) + \mathbf{c} = (\mathbf{a} + \mathbf{c}) + \mathbf{b} = \mathbf{a} + (\mathbf{b} + \mathbf{c}).$$

(iv) In vector addition there exists an *additive identity vector* known as *null vector*, such that

$$\mathbf{a} + \mathbf{0} = \mathbf{a}.$$

(v) To every vector $\mathbf{a}$ in V , there corresponds a vector $-\mathbf{a}$ known as *additive inverse vector*, such that

$$\mathbf{a} + (-\mathbf{a}) = \mathbf{0}.$$

(B) (i) If m, n are any two scalars and $\mathbf{a}$ is a vector in V , then distributive law holds, i.e.,

$$(m+n)\mathbf{a} = m\mathbf{a} + n\mathbf{a}.$$

(ii) If m is any scalar and $\mathbf{a}, \mathbf{b}$ are two vectors belonging to V , then distributive law of scalar multiplication holds, i.e.

$$(\mathbf{a} + \mathbf{b})m = \mathbf{a}m + \mathbf{b}m.$$

(iii) If m, n are any two scalars and $\mathbf{a}$ is a vector belonging to V , then associative law holds, i.e.

$$m(n\mathbf{a}) = (mn)\mathbf{a} = n(m\mathbf{a}).$$

(iv) To every vector $\mathbf{a}$ in V , there corresponds a *multiplicative identity scalar*, such that

$$1\mathbf{a} = \mathbf{a}.$$

Note. In case of scalar quantities m, n, p , we have the following laws of combination:

(i) The addition is commutative, i.e.

$$m+n = n+m.$$

(ii) The addition is associative, i.e.

$$m+n+p = (m+n)+p = (m+p)+n = m+(n+p).$$

(iii) There exists an *additive identity scalar* 0 , which when added to another scalar, leaves it unchanged, such as

$$m+0 = m.$$

(iv) To every scalar m , there corresponds an *additive inverse scalar* $-m$, such that

$$m+(-m) = 0.$$

In fact m and $-m$ are inverse of each other as their sum is zero (identity scalar).

(v) The multiplication is distributive, i.e.

$$m.(n+p) = m.n + m.p.$$

(vi) The multiplication is commutative, i.e.,

$$m.n = n.m.$$

(vii) The multiplication is associative, i.e.,

$$m.(n.p) = (m.n).p = n.(m.p).$$

(viii) There exists a *multiplicative identity scalar* 1 , such that

$$m.1 = m.$$

(ix) To every non-zero scalar m , there corresponds a *multiplicative inverse scalar* $\frac{1}{m}$, such that

$$m \cdot \frac{1}{m} = 1 \text{ (the identity scalar).}$$

Interpretation : Due to directional properties of vector these laws cannot be applied to vectors and laws for vectors are consistent with the physical problems in which vector quantities occur.

1.8. CONDITIONS FOR A PHYSICAL QUANTITY TO BE REPRESENTABLE BY A VECTOR

We have already mentioned that the vector sum of two or more vectors is inherited in the combination of two or more displacements of a point in flat space. Besides displacements there are other many more physical quantities which enter into combination in accordance with the same invariance properties as displacements. Such quantities are also representable as vectors. Precisely a physical quantity representable by a vector must satisfy the two conditions: (i) It must obey the parallelogram law of addition, and (ii) It must have a magnitude as well as direction independent of any choice of co-ordinate axes.

Examples of physical quantities representable by a vector are: velocity, acceleration, electric field intensity and magnetic fields, etc.

Note. It should be carefully noted that all quantities having magnitude and direction are not necessarily vectors. For example, consider the rotation of a rigid body about an axis fixed in space. It has got the magnitude as the angle of rotation and the direction as the direction of the axis. But two or more such rotations do not obey the parallelogram law of addition as they cannot be combined according to the vector law of addition, unless the angles of rotation are vanishingly small. Hence the finite rotations cannot be represented as vectors as may be seen by experimental verification.

Problem 5. Classify which of the following physical measurements are vectors and scalars :

Volume, velocity, mass, acceleration, length, speed, temperature, momentum, force, power, pressure of a gas, temperature gradient, displacement, work, potential, kinetic energy, electric charge, electric or magnetic intensities, magnetic moment.

In the light of the discussion made in §1.8 and the definitions of vectors and scalars so far introduced the above measurements may be classified as follows :

Scalars	Vectors
Volume	Velocity
Mass	Acceleration
Length	Momentum
Speed	Force
Temperature	Power
Pressure of a gas	Temperature gradient
Work	Displacement

Potential	Electric and magnetic field intensities
Kinetic energy	Magnetic moment
Electric charge	

Problem 6. Which of the following have representations as vectors?

(a) Weight, (b) Specific heat, (c) Momentum, (d) Energy, (e) Speed, (f) Velocity, (g) Magnetic field intensity, (h) Gravitational force, (i) Kinetic energy, (j) Age, (k) Flux.

Applying the conditions for a physical quantity to be representable as vector as mentioned in §1.8, we observe that the following quantities have representations as vectors :

(i) Momentum, (ii) Velocity, (iii) Magnetic field Intensity, (iv) Gravitational force.

1.9. RESOLUTION OF VECTORS

(i) **Coplanar vectors.** If there are two non-collinear vectors **a** and **b**, then a third vector **r** which is non-collinear with neither of **a** and **b** but can be made to lie in the same plane in which **a** and **b** lie, can be uniquely expressed in terms of **a** and **b** in the manner

$$\mathbf{r} = m\mathbf{a} + n\mathbf{b}$$

where *m*, *n* are scalars.

As shown in Fig. 1.6, consider two coplanar vectors $\vec{OP}$ and $\vec{OQ}$ such that $\vec{OP} = \mathbf{a}$ and $\vec{OQ} = \mathbf{b}$. Now take another vector $\vec{OC} = \mathbf{r}$ coplanar with $\vec{OP}$ and $\vec{OQ}$ such that $\vec{OC} = \mathbf{r}$. Now take points *A* and *B* on $\vec{OP}$ and $\vec{OQ}$

respectively, such that $\vec{OA} = m \cdot \vec{OP}$ and $\vec{OB} = n \cdot \vec{OQ}$; *m*, *n* being scalars.

It then follows from the parallelogram law of vectors, that

$$\begin{aligned} \vec{r} = \vec{OC} &= \vec{OA} + \vec{AC} \\ &= \vec{OA} + \vec{OB} \\ &= m\vec{OP} + n\vec{OQ} \\ &= m\mathbf{a} + n\mathbf{b} \end{aligned}$$

(ii) **Non-coplanar vectors.** If **a**, **b**, **c** be three non-coplanar vectors, then any vector **r** can be uniquely expressed as

$$\mathbf{r} = m\mathbf{a} + n\mathbf{b} + p\mathbf{c},$$

where *m*, *n*, *p* are scalars.




Fig. 1.6

Choosing a point O as origin of vectors, let OA, OB, OC be three non-coplanar lines, such that




Fig. 1.7

$$\vec{OA} = \mathbf{a}, \vec{OB} = \mathbf{b}, \vec{OC} = \mathbf{c}.$$

Take any point P , such that $\vec{OP} = \mathbf{r}$, in the space of three dimensions. Through P draw planes parallel to three planes BOC, COA and AOB meeting OA, OB, OC in L, M and N respectively. We thus get a parallelepiped with OP as one of its diagonals. Then

$$\vec{r} = \vec{OP} = \vec{OL} + \vec{LP}$$

$$= \vec{OL} + \vec{LN'} + \vec{NP'}$$

$$= \vec{OL} + \vec{OM} + \vec{ON}$$

$$= m\mathbf{a} + n\mathbf{b} + p\mathbf{c},$$

where m, n, p are scalars and $\vec{OL}, \vec{OM}, \vec{ON}$ being collinear with $\vec{OA}, \vec{OB}, \vec{OC}$ respectively, we have

$$\vec{OL} = m\mathbf{a}, \vec{OM} = n\mathbf{b}, \vec{ON} = p\mathbf{c}.$$

(iii) **Components of a vector in three mutually perpendicular directions.** Let OX, OY, OZ be three non-coplanar lines such that each line is perpendicular to the plane containing the other two. The system of axes so chosen form a right handed co-ordinate system such that if OX is turned towards OY about OZ through a small angle, a right handed screw would advance along the positive direction of OZ .

Let $\vec{OP}$ represent the given vector $\mathbf{r}$ and let the length of orthogonal projections of OP along the axes be x, y, z respectively. It is conventional to take $\mathbf{i}, \mathbf{j}, \mathbf{k}$ as unit vectors along OX, OY, OZ respectively, so that

$$\vec{OA} = x\mathbf{i}, \vec{OB} = y\mathbf{j}, \vec{OC} = z\mathbf{k},$$

where OA, OB, OC are collinear with OX, OY, OZ respectively,

$$\text{Now } \vec{r} = \vec{OP} = \vec{OA} + \vec{AP}$$

$$= \vec{OA} + \vec{AQ} + \vec{QP}$$




Fig. 1.8

$$\begin{aligned} &= \vec{OA} + \vec{OB} + \vec{OC} \\ &= x\mathbf{i} + y\mathbf{j} + z\mathbf{k}. \end{aligned}$$

Here $x\mathbf{i}, y\mathbf{j}, z\mathbf{k}$ are called component vectors of $\mathbf{r}$ along the directions of $\mathbf{i}, \mathbf{j}, \mathbf{k}$ and are the orthogonal vector projections of $\mathbf{r}$ along these directions. The scalar projections x, y, z are the rectangular cartesian co-ordinates of the point P referred to O as origin and OX, OY, OZ as axes of reference.

Note 1. Modulus (magnitude) of a vector. Modulus of $\mathbf{r}$, i.e., $|\vec{OP}|$ is given by

$$\sqrt{OA^2 + OB^2 + OC^2}, \text{ i.e., } \sqrt{x^2 + y^2 + z^2}.$$

Thus if $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$, then

$$|\mathbf{r}| = \sqrt{x^2 + y^2 + z^2} = |\vec{OP}|.$$

Note 2. Direction Cosines. If OP makes angles α, β, γ with OX, OY, OZ , respectively, then $x = OP \cos \alpha, y = OP \cos \beta, z = OP \cos \gamma$, but from note 1, $OP = \sqrt{x^2 + y^2 + z^2}$; therefore we have

$$\cos \alpha = \frac{x}{\sqrt{x^2 + y^2 + z^2}}, \cos \beta = \frac{y}{\sqrt{x^2 + y^2 + z^2}}$$

$$\cos \gamma = \frac{z}{\sqrt{x^2 + y^2 + z^2}}.$$

The quantities $\cos \alpha, \cos \beta, \cos \gamma$ introduced and defined in this manner are called the *direction cosines* of the line OP with the axes OX, OY, OZ respectively and are usually denoted by l, m, n , respectively, i.e., $l = \cos \alpha, m = \cos \beta, n = \cos \gamma$

It is obvious that

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1, \\ l^2 + m^2 + n^2 = 1.$$

i.e.

Note 3. Distance between two points. Let $\mathbf{a}, \mathbf{b}$ be the position vectors of two points A and B whose cartesian coordinates are (x_1, y_1, z_1) and (x_2, y_2, z_2) respectively, then

$$\mathbf{a} = x_1\mathbf{i} + y_1\mathbf{j} + z_1\mathbf{k}; \mathbf{b} = x_2\mathbf{i} + y_2\mathbf{j} + z_2\mathbf{k}.$$

$$\therefore \vec{AB} = \mathbf{b} - \mathbf{a} = (x_2 - x_1)\mathbf{i} + (y_2 - y_1)\mathbf{j} + (z_2 - z_1)\mathbf{k},$$

$$\text{so that } AB = |\vec{AB}| = \sqrt{[(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2]}$$

Problem 7. If $\mathbf{a} = 5\mathbf{i} + 6\mathbf{j} - 4\mathbf{k}$ and $\mathbf{b} = 2\mathbf{i} + 3\mathbf{j}$, find

(i) Magnitudes of $\mathbf{a}$ and $\mathbf{b}$,

(ii) the direction cosines of $\mathbf{a}$ and $\mathbf{b}$.

(i) Here $|\mathbf{a}| = \sqrt{(5)^2 + (6)^2 + (-4)^2} = \sqrt{77},$

$$|\mathbf{b}| = \sqrt{(2)^2 + (3)^2 + (0)^2} = \sqrt{13}.$$

(ii) Direction cosines of $\mathbf{a}$ and $\mathbf{b}$ are respectively

$$\frac{5}{\sqrt{77}}, \frac{6}{\sqrt{77}}, -\frac{4}{\sqrt{77}} \text{ and } \frac{2}{\sqrt{13}}, \frac{3}{\sqrt{13}}, 0. \quad \text{Ans.}$$

Problem 8. If $\mathbf{A} = 4\mathbf{i} + 6\mathbf{j} + 2\mathbf{k}$ and $\mathbf{B} = \mathbf{i} + 6\mathbf{j} + \mathbf{k}$, find the magnitudes and direction cosines of $(\mathbf{A} + \mathbf{B})$ and $(\mathbf{A} - \mathbf{B})$.

Given $\mathbf{A} = 4\mathbf{i} + 6\mathbf{j} + 2\mathbf{k}$ and $\mathbf{B} = \mathbf{i} + 6\mathbf{j} + \mathbf{k}$,

$$(\mathbf{A} + \mathbf{B}) = 4\mathbf{i} + 6\mathbf{j} + 2\mathbf{k} + \mathbf{i} + 6\mathbf{j} + \mathbf{k}$$

$$= 5\mathbf{i} + 12\mathbf{j} + 3\mathbf{k}.$$

$$\therefore \text{magnitude of } (\mathbf{A} + \mathbf{B}) = \sqrt{(5^2 + 12^2 + 3^2)},$$

$$|\mathbf{A} + \mathbf{B}| = \sqrt{(25 + 144 + 9)} = \sqrt{178}$$

$\therefore$ direction cosines are given by

$$\cos \alpha = \frac{|\mathbf{A} + \mathbf{B}|_x}{|\mathbf{A} + \mathbf{B}|} = \frac{5}{\sqrt{178}},$$

$$\cos \beta = \frac{|\mathbf{A} + \mathbf{B}|_y}{|\mathbf{A} + \mathbf{B}|} = \frac{12}{\sqrt{178}},$$

$$\cos \gamma = \frac{|\mathbf{A} + \mathbf{B}|_z}{|\mathbf{A} + \mathbf{B}|} = \frac{3}{\sqrt{178}},$$

where $\cos \alpha$, $\cos \beta$ and $\cos \gamma$ are direction cosines along x , y and z axes respectively.

$$\mathbf{A} - \mathbf{B} = 4\mathbf{i} + 6\mathbf{j} + 2\mathbf{k} - (\mathbf{i} + 6\mathbf{j} + \mathbf{k})$$

$$= 4\mathbf{i} + 6\mathbf{j} + 2\mathbf{k} - \mathbf{i} - 6\mathbf{j} - \mathbf{k}$$

$$= 3\mathbf{i} + \mathbf{k}.$$

$$\therefore \text{magnitude of } (\mathbf{A} - \mathbf{B}) = |\mathbf{A} - \mathbf{B}| = \sqrt{(3^2 + 1^2)} = \sqrt{10}.$$

Direction cosines of $(\mathbf{A} - \mathbf{B})$ are given by

$$\cos \alpha = \frac{|\mathbf{A} - \mathbf{B}|_x}{|\mathbf{A} - \mathbf{B}|} = \frac{3}{\sqrt{10}},$$

$$\cos \beta = \frac{|\mathbf{A} - \mathbf{B}|_y}{|\mathbf{A} - \mathbf{B}|} = 0,$$

$$\cos \gamma = \frac{|\mathbf{A} - \mathbf{B}|_z}{|\mathbf{A} - \mathbf{B}|} = \frac{1}{\sqrt{10}}.$$

Problem 9. A person travelling eastwards at a rate of 3 m.p.h., finds that the wind seems to blow from the north. On doubling the speed it appears to come from north east. Find the true velocity of the wind.




Fig. 1.9

Let $\mathbf{i}$, $\mathbf{j}$ be the unit vectors along east and south, so that by Fig. 1.9, we have

$$\vec{OA} = 3\mathbf{i}, \vec{OB} = 6\mathbf{i}.$$

The relative velocity of wind is along

$$\vec{AC} = y\mathbf{j} \text{ (say)}$$

Actual velocity of the wind

$$\begin{aligned} &= \vec{OA} + \vec{AC} = \vec{OC} \\ &= 3\mathbf{i} + y\mathbf{j} \end{aligned}$$

Again, the relative velocity of the wind is along BC , such that

$$\angle ABC = 45^\circ, \angle BAC = 90^\circ \text{ and so } \angle ACB = 45^\circ,$$

$$\therefore AB = AC = 3 \text{ units} = y.$$

$\therefore$ Actual velocity of the wind $= 3\mathbf{i} + 3\mathbf{j}$ and so

$$|\vec{OC}| = \sqrt{(9+9)} = 3\sqrt{2} \text{ m.p.h.}$$

Also $\vec{OC}$ makes equal angles with OE and OS , i.e., the wind is blowing from north-west at $3\sqrt{2}$ m.p.h.

Problem 10. The vectors of magnitudes a , $2a$, $3a$ meet in a point and their directions are along the diagonals of three adjacent faces of a cube. Determine their resultant.

Consider a cube of unit length and let $\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$ be the unit vectors along the adjacent edges OA , OB , OC as shown in Fig. 1.10.

Also let $\vec{OA}$, $\vec{OQ}$, $\vec{OS}$ be the vectors of magnitudes a , $2a$, $3a$ along the three diagonals, OM , OL , OD of three adjacent faces.

$$\text{Here } \vec{OL} = \vec{OA} + \vec{AL} = \mathbf{i} + \mathbf{j}$$

$$\text{so that } \vec{OL} = \frac{\vec{OL}}{|\vec{OL}|} = \frac{\mathbf{i} + \mathbf{j}}{\sqrt{2}}.$$

Similarly unit vectors along OM

$$\text{and } OD \text{ are } \frac{\mathbf{i} + \mathbf{k}}{\sqrt{2}}, \frac{\mathbf{j} + \mathbf{k}}{\sqrt{2}}.$$

Now

$$\vec{OP} = |\vec{OP}| \vec{OM} = \frac{a}{\sqrt{2}} (\mathbf{k} + \mathbf{i}).$$




Fig. 1.10

$$\text{Similarly } \vec{OQ} = \frac{2a}{\sqrt{2}} (\mathbf{i} + \mathbf{j}) \text{ and } \vec{OS} = \frac{3a}{\sqrt{2}} (\mathbf{j} + \mathbf{k}).$$

If $\mathbf{R}$ be the required resultant, then

$$\mathbf{R} = \vec{OP} + \vec{OQ} + \vec{OS} = \frac{3a}{\sqrt{2}} \mathbf{i} + \frac{5a}{\sqrt{2}} \mathbf{j} + \frac{4a}{\sqrt{2}} \mathbf{k}.$$

$$\text{Its magnitude } = |\mathbf{R}| = \sqrt{\left\{ \left(\frac{3a}{\sqrt{2}} \right)^2 + \left(\frac{5a}{\sqrt{2}} \right)^2 + \left(\frac{4a}{\sqrt{2}} \right)^2 \right\}} = 5a \text{ which}$$

is inclined at angles $\cos^{-1} \frac{3}{5\sqrt{2}}$, $\cos^{-1} \frac{1}{\sqrt{2}}$, $\cos^{-1} \frac{4}{5\sqrt{2}}$ with the edges OA , OB , OC respectively.

Problem 11. If the resultant of two forces is equal in magnitude to one of the components and perpendicular to it in direction, find the other component.




Fig. 1.11

Let P, Q be two forces inclined at an angle θ and let their magnitudes be P and Q respectively. Also let i, j be the unit vectors along the direction of P and in a direction perpendicular to it. According to the question the resultant of P and Q is P in a direction perpendicular to that of P .

Here

$P = Pi$ and $Q = Q \cos \theta i + Q \sin \theta j$,
so that $P + Q = P$ gives

$$Pi + Q \cos \theta i + Q \sin \theta j = Pj,$$

i.e. $(P + Q \cos \theta) i + (Q \sin \theta - P) j = 0$.

Equating the coefficients of like vectors on either side, we get

$$P + Q \cos \theta = 0 \text{ and } Q \sin \theta - P = 0$$

i.e. $Q = -\frac{P}{\cos \theta}$ and $Q = \frac{P}{\sin \theta}$.

Dividing, $\tan \theta = -1$, i.e., $\theta = 135^\circ$.

Hence $Q = \frac{P}{\sin 135^\circ} = P\sqrt{2}$.

Problem 12. The base BC of a triangle ABC is divided at G so that $mBG = nGC$. Show that $mAB^2 + nAC^2 = mBG^2 + nCG^2 + (m+n)AG^2$.

Taking A as origin of vectors, let position vectors of B and C be b and c respectively.

Since $mBG = nGC$

i.e. $\frac{BG}{GC} = \frac{n}{m}$

or $BG : GC = n : m$,




Fig. 1.12

therefore the position vector of G is given by $\vec{AG} = \frac{nc + mb}{n+m}$.

Now

$$\begin{aligned} mBG^2 + nCG^2 + (m+n)AG^2 &= m(BG)^2 + n(CG)^2 + (m+n)(AG)^2 \\ &= m[BA + AG]^2 + n[CA + AG]^2 + (m+n)(AG)^2 \\ &= m\left[-b + \frac{nc + mb}{n+m}\right]^2 + n\left[-c + \frac{nc + mb}{n+m}\right]^2 + (m+n)\left(\frac{nc + mb}{n+m}\right)^2 \end{aligned}$$

$$= \frac{1}{(m+n)^2} [mn^2(c-b)^2 + nm^2(b-c)^2 + (m+n)(nc+mb)^2]$$

$$= \frac{1}{(m+n)^2} [mn(m+n)(b-c)^2 + (m+n)(nc+mb)^2]$$

$$= \frac{1}{m+n} [mn b^2 + mnc^2 - 2mn b \cdot c + n^2 c^2 + m^2 b^2 + 2mn b \cdot c],$$

$$= \frac{1}{m+n} [mb^2(n+m) + nc^2(n+m)] = mb^2 + nc^2$$

$$\therefore b^2 = b^2 \text{ and } c^2 = c^2$$

$$= mAB^2 + nAC^2 \text{ (by properties of dot product)}$$

Problem 13. The line AB is bisected in P_1 , P_1B in P_2 , P_2B in P_3 and so on ad infinitum; and the particles of masses $m, \frac{m}{2}, \frac{m}{2^2}, \dots$ etc., are placed at the points $P_1, P_2, P_3, \dots$ etc., respectively. Prove that the distance of their centre of mass from A is equal to one-third of the distance from B to A .

Taking A as origin let the position vector of B be b . The position vectors of the points $P_1,$




Fig. 1.13

$P_2, P_3, \dots$ etc. are $\frac{b}{2}, \frac{b}{2^2}, \frac{b}{2^3}, \dots$

respectively. Let G be the required centre of mass; then

$$\begin{aligned} \vec{AG} &= \frac{m \frac{b}{2} + \frac{m}{2} \cdot \frac{b}{2^2} + \frac{m}{2^2} \cdot \frac{b}{2^3} + \dots}{m + \frac{m}{2} + \frac{m}{2^2} + \dots} \\ &= b \left\{ \frac{\frac{1}{2} + \frac{1}{2^3} + \frac{1}{2^5} + \dots}{1 + \frac{1}{2} + \frac{1}{2^2} + \dots} \right\} = b \cdot \frac{\frac{1}{2}}{1 - \frac{1}{2}} = \frac{b}{3}, \end{aligned}$$

i.e. $AG = \frac{1}{3}$ rd of the distance from B to A .

Problem 14. Prove that

(a) The internal bisectors of the angles of a triangle are concurrent.

(b) The medians of a triangle meet in a point of trisection of each other. (Nagpur, 1965)

(a) Consider a triangle ABC , the position vectors of whose vertices A, B, C are p, q, r respectively. Let a, b, c be the lengths of the sides BC, CA, AB respectively.

If AD be the internal bisector of the angle A , then by geometry,

$$\begin{aligned} BD : DC &= AB : AC \\ &= c : b. \end{aligned}$$




Fig. 1.14

The symmetry of this result follows that the point I also lies on the other two internal bisectors, namely BE and CF .

Hence the three internal bisectors of a triangle are concurrent.

(b) Consider a triangle ABC , the position vectors of whose vertices are $\mathbf{a}, \mathbf{b}, \mathbf{c}$ respectively. Let D, E, F be the mid. points of the sides BC, CR, AB respectively; then their position vectors are

$$\frac{\mathbf{b}+\mathbf{c}}{2}, \frac{\mathbf{c}+\mathbf{a}}{2}, \frac{\mathbf{a}+\mathbf{b}}{2}$$

respectively.

The position vector of a point G dividing the median AD in the ratio $2:1$ is

$$\frac{2 \cdot \frac{\mathbf{b}+\mathbf{c}}{2} + \mathbf{a}}{2+1} = \frac{\mathbf{a}+\mathbf{b}+\mathbf{c}}{3}$$

The symmetry of the result follows that the point G also lies on the other two medians, namely BE and CF . Hence the medians of a triangle meet in a point of trisection of each other.

Problem 15. Forces P, Q act at O and have a resultant R . If any transversal cuts their lines of action at A, B, C respectively, show that

$$\frac{P}{OA} + \frac{Q}{OB} = \frac{R}{OC}.$$




Fig. 1.16

Taking O as origin, let $\mathbf{a}, \mathbf{b}, \mathbf{c}$ be the position vectors of A, B, C respectively; then

$$\hat{\mathbf{a}} = \frac{\mathbf{a}}{OA}, \hat{\mathbf{b}} = \frac{\mathbf{b}}{OB}, \hat{\mathbf{c}} = \frac{\mathbf{c}}{OC}.$$

so that

$$\mathbf{P} = P\hat{\mathbf{a}} = \frac{P}{OA}\mathbf{a}, \mathbf{Q} = \frac{Q}{OB}\mathbf{b},$$

$\therefore$ The position vector of

$$D = \frac{b\mathbf{q} + c\mathbf{r}}{b+c}.$$

Now the position of a point I dividing AD in the ratio $b+c:a$ is

$$\begin{aligned} & \frac{(b+c) \cdot \frac{b\mathbf{q} + c\mathbf{r}}{b+c} + a\mathbf{p}}{b+c+a} \\ &= \frac{a\mathbf{p} + b\mathbf{q} + c\mathbf{r}}{a+b+c}. \end{aligned}$$




Fig. 1.15

$$\mathbf{R} = \frac{R}{OC}\mathbf{c}.$$

$$\text{Since } \mathbf{P} + \mathbf{Q} = \mathbf{R}, \therefore \frac{P}{OA}\mathbf{a} + \frac{Q}{OB}\mathbf{b} = \frac{R}{OC}\mathbf{c}$$

$$\text{or } \frac{P}{OA}\mathbf{a} + \frac{Q}{OB}\mathbf{b} - \frac{R}{OC}\mathbf{c} = 0.$$

The points A, B, C being collinear, we must have

$$\frac{P}{OA} + \frac{Q}{OB} - \frac{R}{OC} = 0, \text{ i.e. } \frac{P}{OA} + \frac{Q}{OB} = \frac{R}{OC}.$$

Problem 16. If the force $F_1, F_2, \dots, F_n$ acting in a plane at O are in equilibrium and any transversal cuts their lines of action in points $L_1, L_2, \dots, L_n$ and a length OL is positive when in the same direction OF , then prove that $\sum \frac{F}{OL} = 0$.

Let AB be the given transversal such that the force $F_1, F_2, \dots, F_n$ make angles $\theta_1, \theta_2, \dots, \theta_n$ with it.

If p be the length of the perpendicular from O to AB ,

$$\text{then } \sin \theta_1 = \frac{p}{OL_1},$$

$$\sin \theta_2 = \frac{p}{OL_2}, \dots, \sin \theta_n = \frac{p}{OL_n}.$$




Fig. 1.17

Let $\mathbf{i}, \mathbf{j}$ be the unit vectors along and perpendicular to AB and F_1, F_2, F_3 etc., be the magnitudes of the forces, then

$$\mathbf{F}_1 = F_1 \cos \theta_1 \mathbf{i} + F_1 \sin \theta_1 \mathbf{j},$$

$$\mathbf{F}_2 = F_2 \cos \theta_2 \mathbf{i} + F_2 \sin \theta_2 \mathbf{j},$$

$$\dots \dots \dots$$

$$\mathbf{F}_n = F_n \cos \theta_n \mathbf{i} + F_n \sin \theta_n \mathbf{j}.$$

The system being in equilibrium, we have $\mathbf{F}_1 + \mathbf{F}_2 + \dots + \mathbf{F}_n = 0$ i.e. $(F_1 \cos \theta_1 + F_2 \cos \theta_2 + \dots + F_n \cos \theta_n) \mathbf{i} + (F_1 \sin \theta_1 + F_2 \sin \theta_2 + \dots + F_n \sin \theta_n) \mathbf{j} = 0$. Equating the coefficients of $\mathbf{j}$ on either side, we get $F_1 \sin \theta_1 + F_2 \sin \theta_2 + \dots + F_n \sin \theta_n = 0$

$$\text{i.e. } F_1 \frac{p}{OL_1} + F_2 \frac{p}{OL_2} + \dots + F_n \frac{p}{OL_n} = 0 \text{ i.e. } \sum \frac{F}{OL} = 0.$$

Problem 17. A boy runs 3 miles on a road towards east. It then turns towards north and runs 4 miles before stopping. Find the resultant distance covered by the boy.

Taking x and y -axes of co-ordinate system along east and north respectively, let A and B be the two successive displacements given by

$$A = 3i.$$

$$B = 4j.$$

So that their resultant,

$$r = A + B = 3i + 4j$$

$$\therefore |r| = \sqrt{(3^2 + 4^2)} = \sqrt{(25)} = 5 \text{ miles making an angle } \theta \text{ with } x\text{-axis where}$$

$$\tan \theta = \frac{\text{coeff. of } j}{\text{coeff. of } i} = \frac{4}{3}.$$

$$\text{Giving } \cos \theta = \frac{3}{5}, \text{ i.e. } \theta = \cos^{-1} \frac{3}{5}.$$

Problem 18. A car is driven eastward for a distance of 5 miles, then northward for 3 miles and then in a direction 30° east of north for 10 miles. Draw the vector diagram and determine the total displacement of the car from its starting point.

Taking x and y -axes towards east and north respectively, and z -axis along the vertical, let the displacements along these axes be a, b, c respectively. Then the resultant displacement r is given by

$$r = a + b + c.$$

But according to the given problem, $a = 5i, b = 3j$

$$\text{and } c = 10 \sin 30^\circ i + 10 \cos 30^\circ j = 5i + 5\sqrt{3}j.$$

$$\therefore r = 5i + 3j + 5i + 5\sqrt{3}j = 10i + (5\sqrt{3} + 3)j.$$

$$\begin{aligned} \text{So that } |r| = r &= \sqrt{\{10^2 + (5\sqrt{3} + 3)^2\}} \\ &= \sqrt{(100 + 75 + 9 + 30\sqrt{3})} \\ &= (100 + 75 + 9 + 52) = \sqrt{(236)} \\ &= 15.35 \text{ miles} \end{aligned}$$

$$\text{and } \cos \theta = \frac{10}{\sqrt{(236)}} \text{ or } \theta = \cos^{-1} \frac{10}{\sqrt{(236)}}.$$

Hence the resultant displacement has magnitude equal to 15.35 miles making an angle $\cos^{-1} \frac{10}{\sqrt{(236)}}$ north of east, i.e. with x -axis.




Fig. 1.18




Fig. 1.19

Problem 19. The projection velocity of a rocket is expressed as $v = 5i + 7j + 9k$

where i, j, k are unit vectors along east, north and vertical direction respectively. Calculate the magnitude of horizontal and vertical components of the velocity. Also deduce the change in angle of projection if the vertical component is doubled. (Agra, B.Sc. 1969)

The velocity vector is given to be expressed as

$$v = 5i + 7j + 9k$$

where i, j and k are unit vectors along east, north and vertical directions respectively, which are taken as axes of reference.

Clearly the vertical component of velocity vector $= 9k$.

$\therefore$ the magnitude of vertical component of velocity $= |9k| = 9$.

Horizontal component of velocity vector $= 5i + 7j$.

$\therefore$ magnitude of horizontal component of velocity $= \sqrt{(5^2 + 7^2)} = \sqrt{(25 + 49)} = \sqrt{(74)}$.

In case the rocket is projected making an angle α with east, then

$$\begin{aligned} \text{we have } \cos \alpha &= \frac{|5i|}{|5i + 7j + 9k|} = \frac{5}{\sqrt{(5^2 + 7^2 + 9^2)}} \\ &= \frac{5}{\sqrt{(25 + 49 + 81)}} = \frac{5}{\sqrt{(155)}} = \frac{5}{12.45} \\ &= .4016 \end{aligned}$$

$$\therefore \alpha = 66^\circ 17' \text{ (by tables of cosine).}$$

If the vertical component is doubled, then the velocity vector becomes $v_1 = 5i + 7j + 18k$.

As such, the angle of projection of rocket with east is given by

$$\begin{aligned} \cos \beta &= \frac{|5i|}{|5i + 7j + 18k|} = \frac{5}{\sqrt{(5^2 + 7^2 + 18^2)}} = \frac{5}{(398)} = \frac{5}{19.95} = .2507 \\ \text{or } \beta &= 75^\circ 29'. \end{aligned}$$

Hence the change in angle $= \beta - \alpha = 75^\circ 29' - 66^\circ 17' = 9^\circ 12'$.

1.10. LINEAR COMBINATION OF VECTORS

A vector v is termed as a linear combination of a set of n vectors $v_1, v_2, v_3, \dots, v_n$, if it is expressible as

$$v = k_1 v_1 + k_2 v_2 + k_3 v_3 + \dots + k_n v_n$$

where $k_1, k_2, k_3, \dots, k_n$ are scalars.

The set of n vectors $v_1, v_2, \dots, v_n$ is said to be linearly dependent if there exists a set of n scalars $k_1, k_2, \dots, k_n$ such that all of them are not zero i.e., at least one of them is non-zero, satisfying the relation,

$$k_1 v_1 + k_2 v_2 + \dots + k_n v_n = 0$$

If all the scalars $k_1, k_2, \dots, k_n$ are zero i.e. $k_1 = k_2 = \dots = k_n = 0$ then the set of n vectors $v_1, v_2, \dots, v_n$ combined as

$$k_1 v_1 + k_2 v_2 + \dots + k_n v_n = 0$$

is said to be linearly independent.

Note 1. It is evident that if a set m ($m < n$) out of n vectors linearly dependent, then the whole set containing n vectors is linearly dependent.

Note 2. If the set of n vectors is linearly independent, then clearly any subset of these n vectors will also be linearly independent.

Problem 20. If a set of n (> 1) vectors is linearly dependent then at least one of these n vectors can be expressed as a linear combination of the remaining $(n-1)$ vectors.

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ be a set of n vectors which are linearly dependent. This set being linearly dependent, we must have a linear combination of the type

$$k_1 \mathbf{v}_1 + k_2 \mathbf{v}_2 + \dots + k_n \mathbf{v}_n = \mathbf{0}$$

where all of the scalars $k_1, k_2, \dots, k_n$ are not zero.

Suppose $k_r \neq 0$. Then, we can write

$$\mathbf{v}_r = - \left(\frac{k_1}{k_r} \mathbf{v}_1 + \frac{k_2}{k_r} \mathbf{v}_2 + \dots + \frac{k_n}{k_r} \mathbf{v}_n \right)$$

which can be expressed as

$$\mathbf{v}_r = k_1' \mathbf{v}_1 + k_2' \mathbf{v}_2 + \dots + k_n' \mathbf{v}_n$$

thereby proving the proposition.

Problem 21. Show that any set of n vectors containing the null vector is linearly dependent.

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ be a set of n vectors, of which $\mathbf{v}_n$ is a null vector i.e. $\mathbf{v}_n = \mathbf{0}$. Then by note 1 of § 1.10, if the set $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{n-1}$ of $(n-1)$ vectors is linearly dependent, the set of n vectors containing these $(n-1)$ vectors and the null vector will also be linearly dependent.

In the case when the set $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{n-1}$ of $(n-1)$ vectors is linearly independent, then by definition there exists a set of $(n-1)$ scalars $k_1, k_2, \dots, k_{n-1}$, all of which being zero, such that

$$k_1 \mathbf{v}_1 + k_2 \mathbf{v}_2 + \dots + k_{n-1} \mathbf{v}_{n-1} = \mathbf{0}.$$

Assuming that $k_n \neq 0$ and $\mathbf{v}_n = \mathbf{0}$, this relation will still hold in the form

$$k_1 \mathbf{v}_1 + k_2 \mathbf{v}_2 + \dots + k_{n-1} \mathbf{v}_{n-1} + k_n \mathbf{v}_n = \mathbf{0}.$$

Where all the scalars $k_1, k_2, \dots, k_n$ are not zero, as $k_n \neq 0$.

This follows that the set of n vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is linearly dependent thereby proving the proposition.

Problem 22. Show that the set of vectors $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3$ given by

$$\mathbf{r}_1 = 2\mathbf{a} - 3\mathbf{b} + \mathbf{c}$$

$$\mathbf{r}_2 = 3\mathbf{a} - 5\mathbf{b} + 2\mathbf{c}$$

$\mathbf{r}_3 = 4\mathbf{a} - 5\mathbf{b} + \mathbf{c}$, $\mathbf{a}, \mathbf{b}, \mathbf{c}$ being non-zero and non-coplanar vectors, is linearly dependent.

The vectors $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3$ will be linearly dependent if there exists a set of scalars k_1, k_2, k_3 not all zero, such that

$$k_1 \mathbf{r}_1 + k_2 \mathbf{r}_2 + k_3 \mathbf{r}_3 = \mathbf{0} \quad \dots (1)$$

$$k_1 (2\mathbf{a} - 3\mathbf{b} + \mathbf{c}) + k_2 (3\mathbf{a} - 5\mathbf{b} + 2\mathbf{c}) + k_3 (4\mathbf{a} - 5\mathbf{b} + \mathbf{c}) = \mathbf{0}$$

$$(2k_1 + 3k_2 + 4k_3) \mathbf{a} - (3k_1 + 5k_2 + 5k_3) \mathbf{b} + (k_1 + 2k_2 + k_3) \mathbf{c} = \mathbf{0}$$

$\mathbf{a}, \mathbf{b}, \mathbf{c}$ being non-zero and non-coplanar, this relation will hold only if the coefficients of $\mathbf{a}, \mathbf{b}, \mathbf{c}$ separately vanish, i.e., if

$$2k_1 + 3k_2 + 4k_3 = 0 \quad \dots (2)$$

$$3k_1 + 5k_2 + 5k_3 = 0 \quad \dots (3)$$

$$k_1 + 2k_2 + k_3 = 0 \quad \dots (4)$$

Multiplying (4) by 5 and then subtracting from (3), we find

$2k_1 + 5k_2 = 0$ which is satisfied for $k_1 = 5, k_2 = -2$ (non-zero values) and then (4) yields $k_3 = -1$.

These values of k_1, k_2, k_3 also satisfy (2) and hence the relation (1) is expressible in the form

$$5\mathbf{r}_1 - 2\mathbf{r}_2 - \mathbf{r}_3 = \mathbf{0} \quad \text{or} \quad \mathbf{r}_3 = 5\mathbf{r}_1 - 2\mathbf{r}_2$$

showing that the set of vectors $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3$ is linearly dependent.

Problem 23. Show that the set of vectors $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3$ given by

$$\mathbf{r}_1 = \mathbf{j} - 2\mathbf{k}, \mathbf{r}_2 = \mathbf{i} - \mathbf{j} + \mathbf{k}, \mathbf{r}_3 = \mathbf{i} + 2\mathbf{j} + \mathbf{k}$$

is linearly independent.

The vectors $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3$ will be linearly independent if there exists a set of scalars k_1, k_2, k_3 all being zero, such that

$$k_1 \mathbf{r}_1 + k_2 \mathbf{r}_2 + k_3 \mathbf{r}_3 = \mathbf{0} \quad \dots (1)$$

$$k_1 (\mathbf{j} - 2\mathbf{k}) + k_2 (\mathbf{i} - \mathbf{j} + \mathbf{k}) + k_3 (\mathbf{i} + 2\mathbf{j} + \mathbf{k}) = \mathbf{0}$$

$$(k_2 + k_3) \mathbf{i} + (k_1 - k_2 + 2k_3) \mathbf{j} + (-2k_1 + k_2 + k_3) \mathbf{k} = \mathbf{0} \quad \dots (2)$$

Now $\mathbf{i}, \mathbf{j}, \mathbf{k}$ being non-coplanar, this relation will hold if

$$k_2 + k_3 = 0 \quad \dots (3)$$

$$k_1 - k_2 + 2k_3 = 0 \quad \dots (4)$$

$$-2k_1 + k_2 + k_3 = 0 \quad \dots (5)$$

Solving (4) and (5),

$$\frac{k_1}{-1-2} = \frac{k_2}{-4-1} = \frac{k_3}{1-2}$$

$$\frac{k_1}{3} = \frac{k_2}{5} = \frac{k_3}{1} = \lambda \quad (\text{say})$$

These equations give,

$$k_1 = 3\lambda, k_2 = 5\lambda, k_3 = \lambda$$

Substituting these values in (3) we find,

$$6\lambda = 0 \quad \text{i.e., } \lambda = 0$$

As such, we have

$$k_1 = 0 = k_2 = k_3$$

showing that the given set of vectors is linearly independent.

Problem 24. Show that a necessary and sufficient condition for the

$$\mathbf{r}_1 = x_1 \mathbf{i} + y_1 \mathbf{j} + z_1 \mathbf{k}, \mathbf{r}_2 = x_2 \mathbf{i} + y_2 \mathbf{j} + z_2 \mathbf{k}, \mathbf{r}_3 = x_3 \mathbf{i} + y_3 \mathbf{j} + z_3 \mathbf{k}$$

to be linearly independent is that the determinant

$$\begin{vmatrix} x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \\ x_3 & y_3 & z_3 \end{vmatrix} \text{ be different from zero.}$$

The set of given vectors will be linearly independent if there exist a set of scalars k_1, k_2, k_3 all being zero, such that

$$k_1 \mathbf{r}_1 + k_2 \mathbf{r}_2 + k_3 \mathbf{r}_3 = \mathbf{0}$$

$$\text{i.e. } k_1 (x_1 \mathbf{i} + y_1 \mathbf{j} + z_1 \mathbf{k}) + k_2 (x_2 \mathbf{i} + y_2 \mathbf{j} + z_2 \mathbf{k}) + k_3 (x_3 \mathbf{i} + y_3 \mathbf{j} + z_3 \mathbf{k}) = \mathbf{0}$$

$$\text{or } (x_1 k_1 + x_2 k_2 + x_3 k_3) \mathbf{i} + (y_1 k_1 + y_2 k_2 + y_3 k_3) \mathbf{j} + (z_1 k_1 + z_2 k_2 + z_3 k_3) \mathbf{k} = \mathbf{0}$$

But $\mathbf{i}, \mathbf{j}, \mathbf{k}$ being non-coplanar, this result will hold only if

$$\left. \begin{aligned} x_1 k_1 + x_2 k_2 + x_3 k_3 &= 0 \\ y_1 k_1 + y_2 k_2 + y_3 k_3 &= 0 \\ z_1 k_1 + z_2 k_2 + z_3 k_3 &= 0 \end{aligned} \right\} \dots (1)$$

Elimination of k_1, k_2, k_3 from these equations with the help of determinant yields

$$\begin{vmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{vmatrix} = 0 \dots (2)$$

But the given set of vectors being linearly independent, we have

$$k_1 = k_2 = k_3 = 0$$

in which case equations (1) are not solvable in the form (2) showing that the given set of vectors will be linearly independent if and only if

$$\begin{vmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{vmatrix} \neq 0$$

otherwise the scalars k_1, k_2, k_3 will be different from zero.

1.11. PRODUCT OF TWO VECTORS

A careful observation of the ways in which two vector quantities enter into combinations in various branches of mathematics and machine leads us to define two well marked and distinct kinds of products, one being called *scalar or dot product* and other being called *vector or cross product*. The former yields a number (scalar) while the latter, vector quantity. In either case the product is jointly proportional to the modules (moduli) of the two vectors.

Conventionally, the scalar or dot product of two vectors $\mathbf{a}$ and $\mathbf{b}$ denoted by $\mathbf{a} \cdot \mathbf{b}$ or $(\mathbf{a}, \mathbf{b})$ and their vector or cross product by $\mathbf{a} \times \mathbf{b}$ or $[\mathbf{a} \mathbf{b}]$.

(1) The Scalar or Dot Product of Two Vectors

Definition. The scalar or dot product of two vectors $\mathbf{a}$ and $\mathbf{b}$, with modules a and b respectively and their directions being inclined at an angle θ , is defined to be the real number $ab \cos \theta$, i.e.

$$\mathbf{a} \cdot \mathbf{b} = ab \cos \theta.$$

Characteristics of dot product. (i) The dot product of two vectors $\mathbf{a}$ and $\mathbf{b}$ is independent of their order

$$\mathbf{a} \cdot \mathbf{b} = ab \cos \theta = \mathbf{b} \cdot \mathbf{a}$$

(ii) The dot product of two vectors $\mathbf{a}$ and $\mathbf{b}$ may be expressed as the product of two numbers, one being the length of one vector and the other resolute of the second in the direction of the first, i.e.

$$\begin{aligned} \mathbf{a} \cdot \mathbf{b} &= (\text{length of } \mathbf{a}) \text{ times (scalar projection of } \mathbf{b} \text{ onto } \mathbf{a}) \\ &= (\text{length of } \mathbf{b}) \text{ times (scalar projection of } \mathbf{a} \text{ onto } \mathbf{b}). \end{aligned}$$

(iii) If $\mathbf{a} \cdot \mathbf{b} = 0$, then either of the two vectors is a null vector or the vectors $\mathbf{a}$ and $\mathbf{b}$ are mutually perpendicular, i.e.

$$\mathbf{a} = \mathbf{0} \text{ or } \mathbf{b} = \mathbf{0} \text{ or } \theta = \frac{1}{2}\pi.$$

In particular $\mathbf{i} \cdot \mathbf{j} = \mathbf{j} \cdot \mathbf{k} = \mathbf{k} \cdot \mathbf{i} = 0$; $\mathbf{i}, \mathbf{j}, \mathbf{k}$ being mutually perpendicular unit vectors.

(iv) The vectors $\mathbf{a}$ and $\mathbf{b}$ are parallel if $\theta = 0$ or π , i.e., if $\mathbf{a} \cdot \mathbf{b} = \pm ab$, where a, b are modules of $\mathbf{a}$ and $\mathbf{b}$ respectively.

(v) The scalar product of two equal vectors $\mathbf{a}$, $\mathbf{a}$ is given by

$$\mathbf{a} \cdot \mathbf{a} = a \cdot a \cos 0^\circ = a^2, \text{ since then } \theta = 0$$

In case $\mathbf{a}$ is a unit vector i.e., $\mathbf{a} = \hat{\mathbf{a}}$ then $|\hat{\mathbf{a}}| = 1$ so that $\hat{\mathbf{a}} \cdot \hat{\mathbf{a}} = 1^2 = 1$

In particular $\mathbf{i} \cdot \mathbf{i} = \mathbf{j} \cdot \mathbf{j} = \mathbf{k} \cdot \mathbf{k} = 1$.

(vi) The scalar product of two unit vectors $\hat{\mathbf{a}}, \hat{\mathbf{b}}$ is given by

$$\hat{\mathbf{a}} \cdot \hat{\mathbf{b}} = \cos \theta \text{ since then } |\hat{\mathbf{a}}| = 1 = |\hat{\mathbf{b}}|$$

(vii) The scalar product is associative i.e. if $\mathbf{a}, \mathbf{b}$ be any two vectors and m, n be any two scalars, then

$$(m\mathbf{a}) \cdot (n\mathbf{b}) = mn (\mathbf{a} \cdot \mathbf{b}) = mn \mathbf{a} \cdot \mathbf{b} = \mathbf{a} \cdot mn\mathbf{b} = na \cdot mb$$

(viii) The scalar product being a number, can occur as the numerical coefficient of any vector, e.g. $(\mathbf{a} \cdot \mathbf{b}) \mathbf{c}$ represents a vector parallel to $\mathbf{c}$ and whose module is $(\mathbf{a} \cdot \mathbf{b})$ times that of $\mathbf{c}$.

(ix) In the case of scalar product, the distributive law of multiplication holds i.e. if $\mathbf{a}, \mathbf{b}, \mathbf{c}$ be three vectors, then

$$\mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$$

Referred to Fig. 1.20, let $\vec{OA} = \mathbf{a}$, $\vec{OB} = \mathbf{b}$, $\vec{OC} = \mathbf{c}$ and projections of OB and BC on OA be respectively OM and MN , so that $ON = OM + MN$.

It is also clear that

$$\vec{OC} = \vec{OB} + \vec{BC} = \mathbf{b} + \mathbf{c}$$

$$\text{Now, } \mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \vec{OC} = (\text{length of } \mathbf{a})$$

times (scalar projection of OC onto $\mathbf{a}$)

$$= a(ON), a \text{ being module of } \mathbf{a}$$

$$= a(OM + MN)$$




Fig. 1.20

$$=a(OM)+a(MN)$$

$$=a \cdot b + a \cdot c \text{ by (ii)}$$

In general, we have

$$(a+b+c \dots) \cdot (l+m+n \dots) = a \cdot l + a \cdot m + \dots + b \cdot l + b \cdot m + \dots$$

and in particular, $(a \pm b) \cdot (a \pm b) = (a \pm b)^2$

$$=a^2 \pm 2a \cdot b + b^2$$

and

$$(a+b) \cdot (a-b) = a^2 - b^2.$$

(x) If θ be the angle between two vectors a and b whose orthogonal projections (components) in the directions of axes of x, y, z are (a_1, a_2, a_3) and (b_1, b_2, b_3) respectively and if i, j, k be the unit vectors along the axes, then

$$a = a_1 i + a_2 j + a_3 k$$

$$b = b_1 i + b_2 j + b_3 k,$$

$$\text{so that } \cos \theta = \frac{a \cdot b}{ab} = \frac{(a_1 i + a_2 j + a_3 k) \cdot (b_1 i + b_2 j + b_3 k)}{|(a_1 i + a_2 j + a_3 k)| \cdot |(b_1 i + b_2 j + b_3 k)|}$$

$$= \frac{a_1 b_1 + a_2 b_2 + a_3 b_3}{\sqrt{(a_1^2 + a_2^2 + a_3^2)} \sqrt{(b_1^2 + b_2^2 + b_3^2)}}.$$

(2) The Vector or Cross Product of Two Vectors

Definition. Given two vectors a and b whose directions are inclined at an angle θ , their vector product is defined to be the vector r , whose module is $ab \sin \theta$ and whose direction is perpendicular to both a and b , being positive relative to a rotation from a to b , i.e.,

$$r = a \times b = |a| |b| \sin \theta \hat{e}$$

$$= ab \sin \theta \hat{e}$$

where $\hat{e}$ is a unit vector perpendicular to the plane of a and b , and has the same direction as is obtained by the motion of a right-handed screw due to rotation from a to b , and a, b are the modules of a and b respectively.

Fig. 1.21


Characteristics of vector product. (i) The vector product is not commutative, i.e., by reversing the order of the factors, the sign of the product is reversed, e.g.

$$b \times a = ba \sin (-\theta) \hat{e} = -ab \sin \theta \hat{e} = -a \times b$$

(ii) The magnitude of the vector product $a \times b$ is equal to the area of the parallelogram of which a and b are the adjacent sides i.e.

$$|a \times b| = |ab \sin \theta \hat{e}| = ab \sin \theta, \text{ as } |\hat{e}| = 1$$

= OA multiplied by the perpendicular distance of OA from B.
= Area of the parallelogram OACB.



(iii) The vector product is associative, i.e., if m be a scalar and a, b be two vectors, then

$$(ma) \times b = a \times (mb) = m(a \times b) = m(ab \sin \theta \hat{e})$$

(iv) The vectors a and b are parallel, if the angle θ included between their directions is 0 or π i.e. if $\theta = 0$ or π , so that

$$a \times b = 0 \text{ as } \sin \theta = 0 \text{ for } \theta = 0 \text{ or } \pi$$

which follows that the vector product of two parallel vector is a null vector.

(v) The vector product of two equal vectors a, a is given by

$$a \times a = 0.$$

Since the two vectors are equal if they are either collinear or parallel. So that the angle θ between them being 0 or π , $\sin \theta = 0$ and hence the result follows.

In particular, if i, j, k are the unit vectors along the principal axes, then

$$i \times i = j \times j = k \times k = 0.$$

(vi) The two vectors a and b with modules a and b respectively, will be mutually perpendicular if the angle θ between their directions is 90° , so that $\sin \theta = 1$.

As such if a, b are at right angles, then $a \times b = ab \hat{e}$, $\hat{e}$ being a unit vector normal to the plane containing a and b .

In case a and b are unit vectors, then $|\hat{a}| = 1$ and $|\hat{b}| = 1$, therefore $\hat{a} \times \hat{b} = \hat{e}$, which shows that the cross product of two mutually perpendicular unit vectors $\hat{a}$ and $\hat{b}$ is a unit vector $\hat{e}$ normal to the plane of $\hat{a}$ and $\hat{b}$.

Hence, in particular if i, j, k be the unit vectors along the principal axes, then

$$i \times j = k = -j \times i$$

$$j \times k = i = -k \times j$$

$$k \times i = j = -i \times k$$

(vii) The vector product of two unit vectors $\hat{a}$ and $\hat{b}$ is given by

$$\hat{a} \times \hat{b} = \sin \theta \hat{e} \text{ as } |\hat{a}| = |\hat{b}| = 1,$$

where θ is the angle between their directions and $\hat{e}$ is the unit vector normal to the plane of $\hat{a}$ and $\hat{b}$.

(viii) The distributive law holds, i.e. in case of vector product if a, b, c are three vectors, then




Fig. 1.22

$$\mathbf{a} \times (\mathbf{b} + \mathbf{c}) = \mathbf{a} \times \mathbf{b} + \mathbf{a} \times \mathbf{c}.$$

Let the components of $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$ along the principal axes be (a_1, a_2, a_3) ; (b_1, b_2, b_3) and (c_1, c_2, c_3) respectively. Then if $\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$ be the unit vectors along the axes, we have

$$\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$$

$$\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$$

$$\mathbf{c} = c_1\mathbf{i} + c_2\mathbf{j} + c_3\mathbf{k}$$

So that, $\mathbf{b} + \mathbf{c} = (b_1 + c_1)\mathbf{i} + (b_2 + c_2)\mathbf{j} + (b_3 + c_3)\mathbf{k}$

$$\therefore \mathbf{a} \times (\mathbf{b} + \mathbf{c}) = (a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}) \times \{(b_1 + c_1)\mathbf{i} + (b_2 + c_2)\mathbf{j} + (b_3 + c_3)\mathbf{k}\}$$

$$= \{a_2(b_3 + c_3) - a_3(b_2 + c_2)\}\mathbf{i}$$

$$+ \{a_3(b_1 + c_1) - a_1(b_3 + c_3)\}\mathbf{j}$$

$$+ \{a_1(b_2 + c_2) - a_2(b_1 + c_1)\}\mathbf{k}$$

$$\text{as } \mathbf{i} \times \mathbf{j} = \mathbf{k} \text{ etc.}$$

$$= \{(a_2b_3 - a_3b_2)\mathbf{i} + (a_3b_1 - a_1b_3)\mathbf{j} + (a_1b_2 - a_2b_1)\mathbf{k}\}$$

$$+ \{a_2c_3 - a_3c_2\}\mathbf{i} + \{a_3c_1 - a_1c_3\}\mathbf{j} + \{a_1c_2 - a_2c_1\}\mathbf{k}$$

$$= \mathbf{a} \times \mathbf{b} + \mathbf{a} \times \mathbf{c}.$$

$$[\text{Since } \mathbf{a} \times \mathbf{b} = (a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}) \times (b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k})]$$

$$= (a_2b_3 - a_3b_2)\mathbf{i} + (a_3b_1 - a_1b_3)\mathbf{j} + (a_1b_2 - a_2b_1)\mathbf{k}$$

$$\text{Similarly } \mathbf{a} \times \mathbf{c} = (a_2c_3 - a_3c_2)\mathbf{i} + (a_3c_1 - a_1c_3)\mathbf{j} + (a_1c_2 - a_2c_1)\mathbf{k}$$

$$\text{In general, } (\mathbf{a} + \mathbf{b} + \mathbf{c} + \dots) \times (\mathbf{l} + \mathbf{m} + \mathbf{n} + \dots)$$

$$= \mathbf{a} \times \mathbf{l} + \mathbf{a} \times \mathbf{m} + \dots + \mathbf{b} \times \mathbf{l} + \mathbf{b} \times \mathbf{m} + \dots + \dots$$

(ix) *Vector product in terms of components.* Consider two vectors $\mathbf{a}$ and $\mathbf{b}$ whose components are (a_1, a_2, a_3) , (b_1, b_2, b_3) respectively along the principal axes. Then if $\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$ are the unit vectors along these axes, we have

$$\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$$

$$\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$$

$$\therefore \mathbf{a} \times \mathbf{b} = (a_2\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}) \times (b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k})$$

$$= (a_2b_3 - a_3b_2)\mathbf{i} + (a_3b_1 - a_1b_3)\mathbf{j} + (a_1b_2 - a_2b_1)\mathbf{k}$$

$$\text{as } \mathbf{i} \times \mathbf{j} = \mathbf{k} = -\mathbf{j} \times \mathbf{i} \text{ etc.}$$

$$= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

(Agra, 1954)

Now if θ be the angle between the directions of $\mathbf{a}$ and $\mathbf{b}$ and $\hat{\epsilon}$, a unit vector normal to the plane of $\mathbf{a}$ and $\mathbf{b}$, then

$$(\mathbf{a} \times \mathbf{b})^2 = (ab \sin \theta \hat{\epsilon})^2 = \{(a_2b_3 - a_3b_2)\mathbf{i} + (a_3b_1 - a_1b_3)\mathbf{j} + (a_1b_2 - a_2b_1)\mathbf{k}\}^2$$

$$\text{i.e., } a^2b^2 \sin^2 \theta = (a_2b_3 - a_3b_2)^2 + (a_3b_1 - a_1b_3)^2 + (a_1b_2 - a_2b_1)^2 \text{ as } \hat{\epsilon}^2 = 1$$

$$\sin^2 \theta = \frac{(a_2b_3 - a_3b_2)^2 + (a_3b_1 - a_1b_3)^2 + (a_1b_2 - a_2b_1)^2}{(a_1^2 + a_2^2 + a_3^2)(b_1^2 + b_2^2 + b_3^2)}$$

$$\text{as } a^2 = a^2 = a_1^2 + a_2^2 + a_3^2 \text{ etc.}$$

(x) *Cross product in terms of dot product.* By definition,

$$(\mathbf{a} \times \mathbf{b})^2 = (ab \sin \theta \hat{\epsilon})^2$$

$$= a^2b^2 \sin^2 \theta,$$

$$= a^2b^2 (1 - \cos^2 \theta)$$

$$= a^2b^2 - a^2b^2 \cos^2 \theta$$

$$= a^2b^2 - (\mathbf{a} \cdot \mathbf{b})^2, \because a^2 = a^2, b^2 = b^2, \mathbf{a} \cdot \mathbf{b} = ab \cos \theta$$

Problem 25. If $\mathbf{a}$ and $\mathbf{b}$ are unit vectors and θ is the angle between them, show that

$$\sin \frac{\theta}{2} = \frac{1}{2} |\mathbf{a} - \mathbf{b}|.$$

$$\text{We have } |\mathbf{a} - \mathbf{b}|^2 = (\mathbf{a} - \mathbf{b})^2$$

$$= a^2 + b^2 - 2\mathbf{a} \cdot \mathbf{b}, \because a^2 = a^2 = 1 \text{ and } b^2 = b^2 = 1$$

$$= 2 - 2 \cos \theta$$

$$= 4 \sin^2 \theta/2,$$

$$\text{i.e., } |\mathbf{a} - \mathbf{b}| = 2 \sin \theta/2,$$

$$\text{so that } \sin \theta/2 = \frac{1}{2} |\mathbf{a} - \mathbf{b}|.$$

Problem 26. From the relations (Lorentz transformation equations in theory of relativity),

$$\begin{cases} \mathbf{r}' = \mathbf{r} + \left[\frac{\gamma - 1}{v^2} \mathbf{v} \cdot \mathbf{r} - \gamma t \right] \mathbf{v}, \\ t' = \gamma \left[t - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2} \right], \end{cases}$$

where $\gamma = \frac{c}{\sqrt{c^2 - v^2}}$, prove the reciprocal relations

$$\begin{cases} \mathbf{r} = \mathbf{r}' + \left[\frac{\gamma - 1}{v^2} \mathbf{v} \cdot \mathbf{r}' + \gamma t' \right] \mathbf{v}, \\ t = \gamma \left[t' + \frac{\mathbf{v} \cdot \mathbf{r}'}{c^2} \right] \end{cases}$$

$$\text{Here, } \mathbf{r}' + \left[\frac{\gamma - 1}{v^2} \mathbf{v} \cdot \mathbf{r}' + \gamma t' \right] \mathbf{v}$$

$$= \mathbf{r} + \left[\frac{\gamma - 1}{v^2} \mathbf{v} \cdot \mathbf{r} - \gamma t \right] \mathbf{v} + \left[\frac{\gamma - 1}{v^2} \mathbf{v} \cdot \left\{ \mathbf{r} + \left(\frac{\gamma - 1}{v^2} \mathbf{v} \cdot \mathbf{r} - \gamma t \right) \mathbf{v} \right\} \right. \\ \left. + \gamma^2 \left(t - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2} \right) \right] \mathbf{v}$$

$$\begin{aligned}
&= \mathbf{r} + \frac{\gamma-1}{v^2} (\mathbf{v} \cdot \mathbf{r}) \mathbf{v} - \gamma t \mathbf{v} + \frac{\gamma-1}{v^2} (\mathbf{v} \cdot \mathbf{r}) \mathbf{v} + \left(\frac{\gamma-1}{v^2} \right)^2 (\mathbf{v} \cdot \mathbf{r}) v^2 \mathbf{v} \\
&\quad - \frac{\gamma-1}{v^2} \gamma t v^2 \mathbf{v} + \gamma^2 t \mathbf{v} - \frac{\gamma^2}{c^2} \mathbf{v} \cdot \mathbf{r} \mathbf{v} \\
&= \mathbf{r} + 2 \frac{(\gamma-1)}{v^2} (\mathbf{v} \cdot \mathbf{r}) \mathbf{v} - \gamma t \mathbf{v} + \frac{(\gamma-1)^2}{v^2} (\mathbf{v} \cdot \mathbf{r}) \mathbf{v} - (\gamma-1) \gamma t \mathbf{v} \\
&\quad + \gamma^2 t \mathbf{v} - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2 - v^2} \mathbf{v} \left[\because \gamma^2 = \frac{c^2}{c^2 - v^2} \right] \\
&= \mathbf{r} + \frac{\gamma-1}{v^2} (\mathbf{v} \cdot \mathbf{r}) \mathbf{v} (2 + \gamma - 1) - \gamma t \mathbf{v} - \gamma^2 t \mathbf{v} + \gamma t \mathbf{v} + \gamma^2 t \mathbf{v} - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2 - v^2} \mathbf{v} \\
&= \mathbf{r} + \frac{\gamma^2 - 1}{v^2} (\mathbf{v} \cdot \mathbf{r}) \mathbf{v} - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2 - v^2} \mathbf{v} \\
&= \mathbf{r} + \frac{1}{v^2} \left(\frac{c^2}{c^2 - v^2} - 1 \right) (\mathbf{v} \cdot \mathbf{r}) \mathbf{v} - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2 - v^2} \mathbf{v} \\
&= \mathbf{r} + \frac{\mathbf{v} \cdot \mathbf{r}}{c^2 - v^2} \mathbf{v} - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2 - v^2} \mathbf{v} = \mathbf{r}.
\end{aligned}$$

Again $\gamma \left[t' + \frac{\mathbf{v} \cdot \mathbf{r}'}{c^2} \right]$

$$\begin{aligned}
&= \gamma \left[\gamma \left(t - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2} \right) + \frac{1}{c^2} \mathbf{v} \cdot \int \mathbf{r} + \left(\frac{\gamma-1}{v^2} \mathbf{v} \cdot \mathbf{r} - \gamma t \right) \mathbf{v} \right] \\
&= \gamma^2 t - \gamma^2 \frac{\mathbf{v} \cdot \mathbf{r}}{c^2} + \gamma \frac{\mathbf{v} \cdot \mathbf{r}}{c^2} + \frac{\gamma}{c^2} \cdot \frac{\gamma-1}{v^2} (\mathbf{v} \cdot \mathbf{r}) v^2 - \frac{\gamma^2 t}{c^2} v^2 \\
&= \frac{c^2 t}{c^2 - v^2} - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2 - v^2} + \gamma \frac{\mathbf{v} \cdot \mathbf{r}}{c^2} + \frac{\gamma^2}{c^2} \mathbf{v} \cdot \mathbf{r} - \gamma \frac{\mathbf{v} \cdot \mathbf{r}}{c^2} - \frac{v^2 t}{c^2 - v^2} \\
&= \frac{(c^2 - v^2) t}{c^2 - v^2} - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2 - v^2} + \frac{\mathbf{v} \cdot \mathbf{r}}{c^2 - v^2} \left[\text{since } \frac{\gamma^2}{c^2} = \frac{1}{c^2 - v^2} \right] \\
&= t.
\end{aligned}$$

Problem 27. What is the meaning of $(\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} - \mathbf{b})$ for the case where $\mathbf{a}^2 = \mathbf{b}^2$?

Here $(\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} - \mathbf{b}) = \mathbf{a}^2 - \mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{a} - \mathbf{b}^2$
 $= \mathbf{a}^2 - \mathbf{b}^2$ as $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$
 $= 0$ as $\mathbf{a}^2 = \mathbf{b}^2$.

This shows that either $\mathbf{a} + \mathbf{b} = 0$, $\mathbf{a} - \mathbf{b} = 0$ or the vectors $\mathbf{a} + \mathbf{b}$ and $\mathbf{a} - \mathbf{b}$ are mutually at right angles.

In the former case when $\mathbf{a} + \mathbf{b} = 0$, $\mathbf{a} - \mathbf{b} = 0$, we have $\mathbf{a} = 0$ and $\mathbf{b} = 0$, i.e. both the vectors $\mathbf{a}$ and $\mathbf{b}$ are null vectors.

Conclusively, either both the vectors $\mathbf{a}$ and $\mathbf{b}$ are null vectors or the angle between the vectors $\mathbf{a} + \mathbf{b}$ and $\mathbf{a} - \mathbf{b}$ is $\frac{1}{2}\pi$.

Problem 28. What is the unit vector perpendicular to each of the vectors $2\mathbf{i} - \mathbf{j} + \mathbf{k}$ and $3\mathbf{i} + 4\mathbf{j} - \mathbf{k}$? Calculate the sine of the angle between these vectors.

Let $\mathbf{a} = 2\mathbf{i} - \mathbf{j} + \mathbf{k}$ and $\mathbf{b} = 3\mathbf{i} + 4\mathbf{j} - \mathbf{k}$.

If $\hat{\mathbf{e}}$ be a unit vector perpendicular to the plane of $\mathbf{a}$ and $\mathbf{b}$, then since $\mathbf{a} \times \mathbf{b}$ is also a vector perpendicular to the plane of $\mathbf{a}$ and $\mathbf{b}$, we have

$$\hat{\mathbf{e}} = \frac{\mathbf{a} \times \mathbf{b}}{|\mathbf{a} \times \mathbf{b}|} \quad \dots (1)$$

Now $\mathbf{a} \times \mathbf{b} = (2\mathbf{i} - \mathbf{j} + \mathbf{k}) \times (3\mathbf{i} + 4\mathbf{j} - \mathbf{k})$

$$= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -1 & 1 \\ 3 & 4 & -1 \end{vmatrix} = -3\mathbf{i} + 5\mathbf{j} + 11\mathbf{k}$$

$\therefore |\mathbf{a} \times \mathbf{b}| = |-3\mathbf{i} + 5\mathbf{j} + 11\mathbf{k}| = \sqrt{(9+25+121)} = \sqrt{(155)}$

Hence $\hat{\mathbf{e}} = \frac{\mathbf{a} \times \mathbf{b}}{|\mathbf{a} \times \mathbf{b}|} = \frac{-3\mathbf{i} + 5\mathbf{j} + 11\mathbf{k}}{\sqrt{(155)}}$

Again $|\mathbf{a}| = |2\mathbf{i} - \mathbf{j} + \mathbf{k}| = \sqrt{(4+1+1)} = \sqrt{6}$

$|\mathbf{b}| = |3\mathbf{i} + 4\mathbf{j} - \mathbf{k}| = \sqrt{(9+16+1)} = \sqrt{26}$

Thus if θ is the angle between the directions of $\mathbf{a}$ and $\mathbf{b}$, then

$$\mathbf{a} \times \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \sin \theta \hat{\mathbf{e}},$$

i.e., $\sin \theta = \frac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}| |\mathbf{b}|} = \frac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}| |\mathbf{b}|} \text{ by (1)}$
 $= \frac{\sqrt{(155)}}{\sqrt{6}\sqrt{26}} = \sqrt{\left(\frac{155}{156}\right)}$

Problem 29. If $\mathbf{a} = 3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k}$ and $\mathbf{b} = -\mathbf{i} + 2\mathbf{j} + 6\mathbf{k}$, then calculate

- the module of each,
- the scalar product $\mathbf{a} \cdot \mathbf{b}$,
- the vector sum and difference $\mathbf{a} + \mathbf{b}$ and $\mathbf{a} - \mathbf{b}$.

(i) We have $|\mathbf{a}| = |3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k}|$
 $= \sqrt{(9+16+25)} = 5\sqrt{2}$
 and $|\mathbf{b}| = |-\mathbf{i} + 2\mathbf{j} + 6\mathbf{k}|$
 $= \sqrt{(1+4+36)} = \sqrt{41}$.

(ii) $\mathbf{a} \cdot \mathbf{b} = (3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k}) \cdot (-\mathbf{i} + 2\mathbf{j} + 6\mathbf{k})$
 $= 3(-1) + 4 \cdot 2 + (-5) \cdot 6$
 $= -3 + 8 - 30 = -25$.

$$\begin{aligned}
 \text{(iii) } \mathbf{a} + \mathbf{b} &= 3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k} + (-\mathbf{i} + 2\mathbf{j} + 6\mathbf{k}) \\
 &= 2\mathbf{i} + 6\mathbf{j} + \mathbf{k}, \\
 \mathbf{a} - \mathbf{b} &= 3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k} - (-\mathbf{i} + 2\mathbf{j} + 6\mathbf{k}) = 4\mathbf{i} + 2\mathbf{j} - 11\mathbf{k}.
 \end{aligned}$$

Problem 30. Show that $\mathbf{a}$ is perpendicular to $\mathbf{b}$ if $|\mathbf{a} + \mathbf{b}| = |\mathbf{a} - \mathbf{b}|$.
We have $|\mathbf{a} + \mathbf{b}| = |\mathbf{a} - \mathbf{b}|$.

Squaring both sides, we get

$$\mathbf{a}^2 + \mathbf{b}^2 + 2\mathbf{a} \cdot \mathbf{b} = \mathbf{a}^2 + \mathbf{b}^2 - 2\mathbf{a} \cdot \mathbf{b}$$

$$\text{or } 4\mathbf{a} \cdot \mathbf{b} = 0$$

$$\text{or } \mathbf{a} \cdot \mathbf{b} = 0,$$

which shows that $\mathbf{a}$ and $\mathbf{b}$ are mutually at right angles.

Problem 31. Two particles emitting from a source have displacements $\mathbf{r}_1 = 4\mathbf{i} + 3\mathbf{j} + 8\mathbf{k}$ and $\mathbf{r}_2 = 2\mathbf{i} + 10\mathbf{j} + 5\mathbf{k}$ at any time. Find the displacement of second particle relative to first.

$$\begin{aligned}
 \text{Required displacement} &= \mathbf{r}_2 - \mathbf{r}_1 \\
 &= 2\mathbf{i} + 10\mathbf{j} + 5\mathbf{k} - (4\mathbf{i} + 3\mathbf{j} + 8\mathbf{k}) \\
 &= -2\mathbf{i} + 7\mathbf{j} - 3\mathbf{k}.
 \end{aligned}$$

Problem 32. Find the scalar and vector products of the vectors $\mathbf{A}$ and $\mathbf{B}$, where $\mathbf{A} = 2\mathbf{i} + \mathbf{j} + \mathbf{k}$ and $\mathbf{B} = 4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$. Also find the angle between $\mathbf{A}$ and $\mathbf{B}$.

Given

$$\mathbf{A} = 2\mathbf{i} + \mathbf{j} + \mathbf{k},$$

$$\mathbf{B} = 4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}.$$

Scalar product

$$= \mathbf{A} \cdot \mathbf{B}$$

$$= (2\mathbf{i} + \mathbf{j} + \mathbf{k}) \cdot (4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k})$$

$$= 8 + 2 - 3 = 7.$$

Vector product

$$= \mathbf{A} \times \mathbf{B}$$

$$= (2\mathbf{i} + \mathbf{j} + \mathbf{k}) \times (4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k})$$

$$= 4\mathbf{k} + 6\mathbf{j} - 4\mathbf{k} - 3\mathbf{i} + 4\mathbf{j} - 2\mathbf{i}$$

$$= -5\mathbf{i} + 10\mathbf{j}.$$

... (1)

According to definition of scalar product,

$$\mathbf{A} \cdot \mathbf{B} = AB \cos \theta,$$

... (2)

where θ is the angle between $\mathbf{A}$ and $\mathbf{B}$.

$$A = |\mathbf{A}| = \sqrt{(A_x^2 + A_y^2 + A_z^2)}$$

$$= \sqrt{(4 + 1 + 1)} = \sqrt{6},$$

$$B = |\mathbf{B}| = \sqrt{(B_x^2 + B_y^2 + B_z^2)} = \sqrt{(16 + 4 + 9)}$$

$$= \sqrt{29}.$$

Substituting values of $(\mathbf{A} \cdot \mathbf{B})$, A and B in equation (2), we get

$$7 = \sqrt{29} \sqrt{6} \cos \theta.$$

$$\therefore \cos \theta = \frac{7}{\sqrt{(29 \times 6)}} = \frac{7}{\sqrt{(174)}}.$$

i.e.,

$$\theta = \cos^{-1} \frac{7}{\sqrt{(174)}}.$$

Problem 33. Prove that an angle inscribed in a semi-circle is a right angle.

Referred to Fig. 1.23 ACB is a semi-circle with AB as bounding diameter and C any point on its circumference. Let O be its

centre and r the radius. Also let $\vec{AO} = \vec{a} = \vec{OB}$ and $\vec{OC} = \vec{b}$.




Fig. 1.23

Now

$$\vec{AC} = \vec{AO} + \vec{OC} = \vec{a} + \vec{b}$$

and

$$\vec{CB} = \vec{CO} + \vec{OB} = -\vec{b} + \vec{a}.$$

$$\begin{aligned}
 \therefore \vec{AC} \cdot \vec{CB} &= (\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) \\
 &= \mathbf{a}^2 - \mathbf{b}^2
 \end{aligned}$$

$$= AO^2 - OC^2 \text{ as } \mathbf{a}^2 = AO^2 = AO^2 \text{ etc.}$$

$$= r^2 - r^2$$

$$= 0.$$

But from the definition of dot product,

$$\cos \angle ACB = \frac{\vec{AC} \cdot \vec{CB}}{|\vec{AC}| |\vec{CB}|} = 0.$$

$$\angle ACB = \frac{1}{2}\pi.$$

Problem 34. Prove that the area of a triangle whose two sides are $\mathbf{A}$ and $\mathbf{B}$ is given by $\frac{1}{2} |\mathbf{A} \times \mathbf{B}|$. Also find the direction-cosines of normal to this area.




Fig. 1.24

Area of the $\triangle XYZ$

$$= \frac{1}{2} A \cdot h$$

$$= \frac{1}{2} AB \sin \theta$$

$$= \frac{1}{2} |\mathbf{A} \times \mathbf{B}|.$$

The vector area is perpendicular to the plane containing $\mathbf{A}$ and $\mathbf{B}$.

And magnitude of area

$$= \frac{1}{2} |\mathbf{A} \times \mathbf{B}|.$$

$$\begin{aligned}
 \therefore \text{Vector area} &= \frac{1}{2} (\mathbf{A} \times \mathbf{B}) = \frac{1}{2} (iA_x + jA_y + kA_z) \times (iB_x + jB_y + kB_z) \\
 &= \frac{1}{2} i(A_y B_z - A_z B_y) + \frac{1}{2} j(A_z B_x - A_x B_z) + \frac{1}{2} k(A_x B_y - A_y B_x).
 \end{aligned}$$

Thus, direction-cosines of the normal to the given area are given by

$$\cos \alpha = \frac{A_y B_z - A_z B_y}{2 |\mathbf{A} \times \mathbf{B}|}, \cos \beta = \frac{A_z B_x - A_x B_z}{2 |\mathbf{A} \times \mathbf{B}|} \text{ and } \cos \gamma = \frac{A_x B_y - A_y B_x}{2 |\mathbf{A} \times \mathbf{B}|}$$

Note. Area of the parallelogram with sides **A** and **B** is double of the above area, i.e., $(\mathbf{A} \times \mathbf{B})$.

Problem 35. Show that $\mathbf{a} \times (\mathbf{b} + \mathbf{c}) + \mathbf{b} \times (\mathbf{c} + \mathbf{a}) + \mathbf{c} \times (\mathbf{a} + \mathbf{b}) = \mathbf{0}$.

We have $\mathbf{a} \times (\mathbf{b} + \mathbf{c}) + \mathbf{b} \times (\mathbf{c} + \mathbf{a}) + \mathbf{c} \times (\mathbf{a} + \mathbf{b}) = \mathbf{a} \times \mathbf{b} + \mathbf{a} \times \mathbf{c} + \mathbf{b} \times \mathbf{c} + \mathbf{b} \times \mathbf{a} + \mathbf{c} \times \mathbf{a} + \mathbf{c} \times \mathbf{b}$
 $= \mathbf{a} \times \mathbf{b} - \mathbf{c} \times \mathbf{a} + \mathbf{b} \times \mathbf{c} - \mathbf{a} \times \mathbf{b} + \mathbf{c} \times \mathbf{a} - \mathbf{b} \times \mathbf{c} = \mathbf{0}$.

1.12. TRIPLE PRODUCTS OF VECTORS

We have stated that the vector product of two vectors **b** and **c** is a vector quantity. So this product $(\mathbf{b} \times \mathbf{c})$ may be multiplied scalarly or vectorially with a third vector **a** to give two triple products namely $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$ and $\mathbf{a} \times (\mathbf{b} \times \mathbf{c})$. The former being a scalar quantity is termed as *scalar triple product* and the latter being a vector quantity is called *vector triple product*.

(1) The Scalar Triple Product

Definition. Let **a**, **b**, **c** be three vectors. Then the scalar product of any of these vectors with the vector product of the other two such as $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$ is called scalar triple product of the vectors **a**, **b**, **c** and denoted by $[\mathbf{abc}]$ or $[a, b, c]$. Obviously, this type of triple product is a scalar quantity.

Note. The scalar triple product is sometimes known as **Box Product**.

Characteristics of Scalar Triple Product. (i) Geometrically interpreted as, below the scalar triple product of three vectors **a**, **b**, **c** represents the volume of a parallelepiped which has for its three coterminus edges the vectors **a**, **b**, **c**.




Fig. 1.25

the area of the parallelogram **OBEC**.

Volume of the parallelepiped with coterminus edges **OA**, **OB**, **OC** = (area of the parallelogram of the base **OBEC**) multiplied by the perpendicular distance of the plane **OBEC** from the point **A** = (area of the parallelogram **OBEC**) multiplied by (the scalar projection of **OA** on **ON**)
 $= |\mathbf{n}| (OA \cos \theta)$, where $\angle AON = \theta$

Construct a parallelepiped with coterminus edges **OA**, **OB** and **OC**, such that $\vec{OA} = \mathbf{a}$, $\vec{OB} = \mathbf{b}$, $\vec{OC} = \mathbf{c}$.

Suppose that $\mathbf{n} = \mathbf{b} \times \mathbf{c}$ and its direction is **ON**, which is perpendicular to the plane **OBEC** whose adjacent sides are **b** and **c**. The direction of the vector **n** is positive in the sense of rotation from **OB** to **OC**.

From the property of vector product of two vectors, it follows that $|\mathbf{n}|$ measures

$= |\mathbf{n}| |\mathbf{a}| \cos \theta$, since $OA = |\vec{OA}| = |\mathbf{a}|$
 $= \mathbf{a} \cdot \mathbf{n}$ from the definition of dot product
 $= \mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = [\mathbf{abc}] = V$ (say)

where **V** measures the volume of the parallelepiped.

The product is regarded as positive or negative according as θ is acute or obtuse. It is easy to show that

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \mathbf{b} \cdot (\mathbf{c} \times \mathbf{a}) = \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b}) = \pm V,$$

$$\text{i.e. } [\mathbf{abc}] = [\mathbf{bca}] = [\mathbf{cab}] = \pm V.$$

This follows that if the cyclic order of the occurrence of the vectors **a**, **b**, **c** is maintained, the position of cross and dot may be interchanged without changing the value of the product.

Again since $\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a}$ etc., therefore

$$[\mathbf{bac}] = [\mathbf{cba}] = -[\mathbf{abc}] = \mp V.$$

(ii) If the vectors **a**, **b**, **c** are coplanar, then their scalar triple product is zero, i.e. $[\mathbf{abc}] = 0$.

Since the volume of the parallelepiped, so formed with coplanar vectors **a**, **b**, **c** as coterminus edges will be zero.

$$\text{As such } [\mathbf{aab}] = [\mathbf{abb}] = [\mathbf{cbc}] \text{ etc.} = 0.$$

(iii) The scalar triple product may be expressed in terms of components.

Let **a**, **b**, **c** be three vectors whose magnitudes in a right handed system of unit vectors **i**, **j**, **k** are (a_1, a_2, a_3) , (b_1, b_2, b_3) and (c_1, c_2, c_3) respectively. Then

$$\mathbf{a} = a_1 \mathbf{i} + a_2 \mathbf{j} + a_3 \mathbf{k},$$

$$\mathbf{b} = b_1 \mathbf{i} + b_2 \mathbf{j} + b_3 \mathbf{k},$$

$$\mathbf{c} = c_1 \mathbf{i} + c_2 \mathbf{j} + c_3 \mathbf{k},$$

$$\text{we have, } \mathbf{b} \times \mathbf{c} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = (b_2 c_3 - b_3 c_2) \mathbf{i} + (b_3 c_1 - b_1 c_3) \mathbf{j} + (b_1 c_2 - b_2 c_1) \mathbf{k}.$$

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = (a_1 \mathbf{i} + a_2 \mathbf{j} + a_3 \mathbf{k}) \cdot \{(b_2 c_3 - b_3 c_2) \mathbf{i} + (b_3 c_1 - b_1 c_3) \mathbf{j} + (b_1 c_2 - b_2 c_1) \mathbf{k}\},$$

$$\text{i.e. } [\mathbf{abc}] = a_1 (b_2 c_3 - b_3 c_2) + a_2 (b_3 c_1 - b_1 c_3) + a_3 (b_1 c_2 - b_2 c_1)$$

$$= \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}.$$

$$\text{In particular, } [\mathbf{ijk}] = [\mathbf{jki}] = [\mathbf{kij}] = 1$$

$$\text{and } [\mathbf{ikj}] = [\mathbf{kji}] = [\mathbf{jik}] = -1.$$

In general if the three vectors **a**, **b**, **c** are resolved in terms of three non-coplanar vectors **l**, **m**, **n**, then

$$\mathbf{a} = a_1 \mathbf{l} + a_2 \mathbf{m} + a_3 \mathbf{n},$$

$$\mathbf{b} = b_1 \mathbf{l} + b_2 \mathbf{m} + b_3 \mathbf{n},$$

$$\mathbf{c} = c_1 \mathbf{l} + c_2 \mathbf{m} + c_3 \mathbf{n}.$$

We have $(\mathbf{b} \times \mathbf{c}) = (b_2c_3 - b_3c_2) \mathbf{m} \times \mathbf{n} + (b_3c_1 - b_1c_3) \mathbf{n} \times \mathbf{l} + (b_1c_2 - b_2c_1) \mathbf{l} \times \mathbf{m}$

$$\therefore \mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = a_1 (b_2c_3 - b_3c_2) [\mathbf{lmn}] + a_2 (b_3c_1 - b_1c_3) [\mathbf{mnl}] + a_3 (b_1c_2 - b_2c_1) [\mathbf{nml}]$$

$$= \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} [\mathbf{lmn}] \quad \text{as } [\mathbf{lmn}] = [\mathbf{mnl}] = [\mathbf{nml}]$$

(2) The Vector Triple Product.

Definition. The product of the type $\mathbf{a} \times (\mathbf{b} \times \mathbf{c})$ is called the vector triple product of given three vectors $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$ and is expressed as

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \cdot \mathbf{c}) \mathbf{b} - (\mathbf{a} \cdot \mathbf{b}) \mathbf{c}.$$

Suppose that $\mathbf{q} = \mathbf{a} \times (\mathbf{b} \times \mathbf{c})$.

Then, $\mathbf{q}$ being a vector product of two vectors $\mathbf{a}$ and $(\mathbf{b} \times \mathbf{c})$ represents a vector perpendicular to both $\mathbf{a}$ and $(\mathbf{b} \times \mathbf{c})$ and therefore by the property of dot product, we have

$$\mathbf{q} \cdot \mathbf{a} = 0 \text{ and } \mathbf{q} \cdot (\mathbf{b} \times \mathbf{c}) = 0.$$

But the product $\mathbf{b} \times \mathbf{c}$ being a vector product of two vectors $\mathbf{b}$ and $\mathbf{c}$ is itself a vector perpendicular to both $\mathbf{b}$ and $\mathbf{c}$, i.e. a vector normal to the plane of $\mathbf{b}$ and $\mathbf{c}$.

It follows that $\mathbf{q}$ lies in the plane of $\mathbf{b}$ and $\mathbf{c}$, so that $\mathbf{q}$ is expressible in terms of $\mathbf{b}$ and $\mathbf{c}$.

Suppose that $\mathbf{q} = s\mathbf{b} + t\mathbf{c}$, where s and t are scalars.

Multiplying both sides scalarly by $\mathbf{a}$, we get

$$\mathbf{q} \cdot \mathbf{a} = s(\mathbf{a} \cdot \mathbf{b}) + t(\mathbf{a} \cdot \mathbf{c}),$$

$$\text{i.e.,} \quad \frac{s}{\mathbf{a} \cdot \mathbf{c}} = \frac{-t}{\mathbf{a} \cdot \mathbf{b}} = \lambda \text{ (say)} \quad (\text{since } \mathbf{q} \cdot \mathbf{a} = 0)$$

$$\text{Then } \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{q} = \lambda [(\mathbf{a} \cdot \mathbf{c}) \mathbf{b} - (\mathbf{a} \cdot \mathbf{b}) \mathbf{c}]. \quad \dots (1)$$

In order to find λ , let us introduce an orthogonal right handed system of three unit vectors $\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$ such that $\mathbf{i}$ is along $\mathbf{a}$ and $\mathbf{j}$ perpendicular to it in the plane of $\mathbf{a}$ and $\mathbf{b}$, the direction of $\mathbf{k}$ is automatically decided because, $\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$ form a right handed system of vectors. Then

$$\mathbf{a} = a_1\mathbf{i}, \mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j}, \mathbf{c} = c_1\mathbf{i} + c_2\mathbf{j} + c_3\mathbf{k}$$

$$\text{Now } \mathbf{b} \times \mathbf{c} = b_2c_3\mathbf{i} - b_1c_3\mathbf{j} + (b_1c_2 - b_2c_1)\mathbf{k}$$

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (a_1b_2c_3 - a_1b_1c_2)\mathbf{j} - a_1b_1c_3\mathbf{k}$$

so that

$$\text{Also } (\mathbf{a} \cdot \mathbf{c}) \mathbf{b} = (a_1b_1c_1 + a_1b_2c_2)\mathbf{j} \text{ and } (\mathbf{a} \cdot \mathbf{b}) \mathbf{c} = a_1b_1c_1\mathbf{i} + a_1b_1c_2\mathbf{j} + a_1b_1c_3\mathbf{k}$$




Fig. 1.26

Putting these values in (1), we get, $\lambda = 1$.

Substituting this value of λ in (1), we find

$$\mathbf{q} = \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \cdot \mathbf{c}) \mathbf{b} - (\mathbf{a} \cdot \mathbf{b}) \mathbf{c} \quad (\text{Vikram, 1969})$$

Characteristics of Vector Triple Product.

The vector triple product is not associative, i.e.,

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) \neq (\mathbf{a} \times \mathbf{b}) \times \mathbf{c}.$$

$$\text{Since } \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \cdot \mathbf{c}) \mathbf{b} - (\mathbf{a} \cdot \mathbf{b}) \mathbf{c}$$

$$\text{and } (\mathbf{a} \times \mathbf{b}) \times \mathbf{c} = -\mathbf{c} \times (\mathbf{a} \times \mathbf{b})$$

$$= -[(\mathbf{c} \cdot \mathbf{b}) \mathbf{a} - (\mathbf{c} \cdot \mathbf{a}) \mathbf{b}]$$

$$= (\mathbf{a} \cdot \mathbf{c}) \mathbf{b} - (\mathbf{b} \cdot \mathbf{c}) \mathbf{a}.$$

which follows that the product $\mathbf{a} \times (\mathbf{b} \times \mathbf{c})$ represents a vector coplanar with $\mathbf{b}$ and $\mathbf{c}$ while the product $(\mathbf{a} \times \mathbf{b}) \times \mathbf{c}$ represents a vector lying in the plane of $\mathbf{a}$ and $\mathbf{b}$. Hence the two products do not represent the same vector quantity, i.e.,

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) \neq (\mathbf{a} \times \mathbf{b}) \times \mathbf{c}.$$

Problem 36. If $\mathbf{A} = 4\mathbf{i} - 5\mathbf{j} + 3\mathbf{k}$, $\mathbf{B} = 2\mathbf{i} - 10\mathbf{j} - 7\mathbf{k}$ and $\mathbf{C} = 5\mathbf{i} + 7\mathbf{j} - 4\mathbf{k}$ deduce the values of

$$(i) (\mathbf{A} \times \mathbf{B}) \cdot \mathbf{C} \text{ and } (ii) \mathbf{A} \times (\mathbf{B} \times \mathbf{C}). \quad (\text{Agra, 1969})$$

$$(i) \text{ Given } \mathbf{A} = 4\mathbf{i} - 5\mathbf{j} + 3\mathbf{k}$$

$$\mathbf{B} = 2\mathbf{i} - 10\mathbf{j} - 7\mathbf{k}$$

$$\mathbf{C} = 5\mathbf{i} + 7\mathbf{j} - 4\mathbf{k}.$$

$$\begin{aligned} \mathbf{A} \times \mathbf{B} &= (4\mathbf{i} - 5\mathbf{j} + 3\mathbf{k}) \times (2\mathbf{i} - 10\mathbf{j} - 7\mathbf{k}) \\ &= -40\mathbf{k} + 28\mathbf{j} + 10\mathbf{k} + 35\mathbf{i} + 6\mathbf{j} + 30\mathbf{i} \\ &= 65\mathbf{i} + 34\mathbf{j} - 30\mathbf{k}. \end{aligned}$$

$$\begin{aligned} (\mathbf{A} \times \mathbf{B}) \cdot \mathbf{C} &= (65\mathbf{i} + 34\mathbf{j} - 30\mathbf{k}) \cdot (5\mathbf{i} + 7\mathbf{j} - 4\mathbf{k}) \\ &= 325 + 238 + 120 \\ &= 683. \end{aligned}$$

$$\begin{aligned} (ii) \quad (\mathbf{B} \times \mathbf{C}) &= (2\mathbf{i} - 10\mathbf{j} - 7\mathbf{k}) \times (5\mathbf{i} + 7\mathbf{j} - 4\mathbf{k}) \\ &= 14\mathbf{k} + 8\mathbf{j} + 50\mathbf{k} + 40\mathbf{i} - 35\mathbf{j} + 49\mathbf{i} \\ &= 89\mathbf{i} - 27\mathbf{j} + 64\mathbf{k}. \end{aligned}$$

$$\begin{aligned} \mathbf{A} \times (\mathbf{B} \times \mathbf{C}) &= (4\mathbf{i} - 5\mathbf{j} + 3\mathbf{k}) \times (89\mathbf{i} - 27\mathbf{j} + 64\mathbf{k}) \\ &= -108\mathbf{k} - 256\mathbf{j} + 445\mathbf{k} - 320\mathbf{i} + 267\mathbf{j} + 81\mathbf{i} \\ &= -239\mathbf{i} + 11\mathbf{j} + 337\mathbf{k}. \end{aligned}$$

Problem 37. Show that $[\mathbf{a} \times \mathbf{b}, \mathbf{b} \times \mathbf{c}, \mathbf{c} \times \mathbf{a}] = [\mathbf{abc}]$

$$= \begin{vmatrix} \mathbf{a} \cdot \mathbf{a} & \mathbf{a} \cdot \mathbf{b} & \mathbf{a} \cdot \mathbf{c} \\ \mathbf{b} \cdot \mathbf{a} & \mathbf{b} \cdot \mathbf{b} & \mathbf{b} \cdot \mathbf{c} \\ \mathbf{c} \cdot \mathbf{a} & \mathbf{c} \cdot \mathbf{b} & \mathbf{c} \cdot \mathbf{c} \end{vmatrix} \text{ by means of determinant.}$$

Let the components of the vectors $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$ in the directions of the axes of x , y , z along which $\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$ are the unit vectors, be (a_1, a_2, a_3) , (b_1, b_2, b_3) and (c_1, c_2, c_3) respectively. Then

$$\left. \begin{aligned} \mathbf{a} &= a_1 \mathbf{i} + a_2 \mathbf{j} + a_3 \mathbf{k} \\ \mathbf{b} &= b_1 \mathbf{i} + b_2 \mathbf{j} + b_3 \mathbf{k} \\ \mathbf{c} &= c_1 \mathbf{i} + c_2 \mathbf{j} + c_3 \mathbf{k} \end{aligned} \right\},$$

so that

$$\left. \begin{aligned} \mathbf{a} \cdot \mathbf{a} &= a_1^2 + a_2^2 + a_3^2 = \Sigma a_1^2 \text{ (say)}, \quad \mathbf{b} \cdot \mathbf{b} = \Sigma b_1^2, \quad \mathbf{c} \cdot \mathbf{c} = \Sigma c_1^2 \\ \mathbf{a} \cdot \mathbf{b} &= \Sigma a_1 b_1, \quad \mathbf{b} \cdot \mathbf{c} = \Sigma b_1 c_1 \text{ and } \mathbf{c} \cdot \mathbf{a} = \Sigma c_1 a_1 \end{aligned} \right\}$$

Now, $[\mathbf{a}\mathbf{b}\mathbf{c}] = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$ from (1)

$$\begin{aligned} \therefore [\mathbf{a}\mathbf{b}\mathbf{c}]^2 &= \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ &= \begin{vmatrix} \Sigma a_1^2 & \Sigma a_1 b_1 & \Sigma c_1 a_1 \\ \Sigma a_1 b_1 & \Sigma b_1^2 & \Sigma b_1 c_1 \\ \Sigma c_1 a_1 & \Sigma b_1 c_1 & \Sigma c_1^2 \end{vmatrix} = \begin{vmatrix} \mathbf{a} \cdot \mathbf{a} & \mathbf{a} \cdot \mathbf{b} & \mathbf{a} \cdot \mathbf{c} \\ \mathbf{b} \cdot \mathbf{a} & \mathbf{b} \cdot \mathbf{b} & \mathbf{b} \cdot \mathbf{c} \\ \mathbf{c} \cdot \mathbf{a} & \mathbf{c} \cdot \mathbf{b} & \mathbf{c} \cdot \mathbf{c} \end{vmatrix} \text{ from (2)} \end{aligned}$$

Again, Let $\Delta \equiv \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = [\mathbf{a}\mathbf{b}\mathbf{c}]$.

Then, if $A_1, B_1, C_1 \dots$ etc. be the cofactors of $a_1, b_1, c_1 \dots$ etc. we have

$$\left. \begin{aligned} a_1 A_1 + a_2 A_2 + a_3 A_3 &= \Delta \\ a_1 B_1 + a_2 B_2 + a_3 B_3 &= 0 \\ a_1 C_1 + a_2 C_2 + a_3 C_3 &= 0 \text{ etc.} \end{aligned} \right\}$$

where $A_1 = b_2 c_3 - b_3 c_2$ etc.

Suppose that $\Delta' \equiv \begin{vmatrix} A_1 & A_2 & A_3 \\ B_1 & B_2 & B_3 \\ C_1 & C_2 & C_3 \end{vmatrix}$

$$\begin{aligned} \therefore \Delta \Delta' &= \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \begin{vmatrix} A_1 & A_2 & A_3 \\ B_1 & B_2 & B_3 \\ C_1 & C_2 & C_3 \end{vmatrix} \\ &= \begin{vmatrix} \Delta & 0 & 0 \\ 0 & \Delta & 0 \\ 0 & 0 & \Delta \end{vmatrix} \text{ from (3)} \end{aligned}$$

$$= \Delta^3,$$

i.e. $\Delta' = \Delta^2$

or $[\mathbf{a}\mathbf{b}\mathbf{c}]^2 = \begin{vmatrix} A_1 & A_2 & A_3 \\ B_1 & B_2 & B_3 \\ C_1 & C_2 & C_3 \end{vmatrix}$.

But $\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = (a_2 b_3 - a_3 b_2) \mathbf{i} + (b_1 a_3 - b_3 a_1) \mathbf{j} + (a_1 b_2 - a_2 b_1) \mathbf{k}$.

Similarly $\mathbf{b} \times \mathbf{c} = (b_2 c_3 - b_3 c_2) \mathbf{i} + (c_1 b_3 - c_3 b_1) \mathbf{j} + (b_1 c_2 - b_2 c_1) \mathbf{k}$
 $\mathbf{c} \times \mathbf{a} = (c_2 a_3 - c_3 a_2) \mathbf{i} + (a_1 c_3 - a_3 c_1) \mathbf{j} + (c_1 a_2 - c_2 a_1) \mathbf{k}$.

$$[\mathbf{a} \times \mathbf{b}, \mathbf{b} \times \mathbf{c}, \mathbf{c} \times \mathbf{a}]$$

$$= \begin{vmatrix} a_2 b_3 - a_3 b_2 & b_1 a_3 - b_3 a_1 & a_1 b_2 - a_2 b_1 \\ b_2 c_3 - b_3 c_2 & c_1 b_3 - c_3 b_1 & b_1 c_2 - b_2 c_1 \\ c_2 a_3 - c_3 a_2 & a_1 c_3 - a_3 c_1 & c_1 a_2 - c_2 a_1 \end{vmatrix}$$

$$= \begin{vmatrix} C_1 & C_2 & C_3 \\ A_1 & A_2 & A_3 \\ B_1 & B_2 & B_3 \end{vmatrix} \text{ when } A_1, B_1, C_1 \dots \text{ etc. are the cofactors in } \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

$$= \begin{vmatrix} A_1 & A_2 & A_3 \\ B_1 & B_2 & B_3 \\ C_1 & C_2 & C_3 \end{vmatrix}$$

$$= [\mathbf{a}\mathbf{b}\mathbf{c}]^2 \text{ from (4).}$$

Problem 38. Show that $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) + \mathbf{b} \times (\mathbf{c} \times \mathbf{a}) + \mathbf{c} \times (\mathbf{a} \times \mathbf{b}) = 0$.

We have $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \cdot \mathbf{c}) \mathbf{b} - (\mathbf{a} \cdot \mathbf{b}) \mathbf{c}$
 $\mathbf{b} \times (\mathbf{c} \times \mathbf{a}) = (\mathbf{b} \cdot \mathbf{a}) \mathbf{c} - (\mathbf{b} \cdot \mathbf{c}) \mathbf{a}$
 $\mathbf{c} \times (\mathbf{a} \times \mathbf{b}) = (\mathbf{c} \cdot \mathbf{b}) \mathbf{a} - (\mathbf{c} \cdot \mathbf{a}) \mathbf{b}$.

Adding all together, we get

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) + \mathbf{b} \times (\mathbf{c} \times \mathbf{a}) + \mathbf{c} \times (\mathbf{a} \times \mathbf{b}) = 0.$$

Problem 39. Find the volume of a parallelepiped whose three extremities edges are described by the vectors $\mathbf{i} + 2\mathbf{j}$, $4\mathbf{j}$ and $\mathbf{j} + 3\mathbf{k}$.

Volume of the required parallelepiped

= scalar triple product of the vectors $\mathbf{i} + 2\mathbf{j}$, $4\mathbf{j}$ and $\mathbf{j} + 3\mathbf{k}$

$$= (\mathbf{i} + 2\mathbf{j}) \cdot \{4\mathbf{j} \times (\mathbf{j} + 3\mathbf{k})\}$$

$$= (\mathbf{i} + 2\mathbf{j}) \cdot 12\mathbf{i}$$

$$= 12 \quad \mathbf{i} \cdot \mathbf{i} = 1 \text{ and } \mathbf{j} \cdot \mathbf{i} = 0.$$

Problem 40. Show that the law of refraction of light passing from a medium of refractive index μ into one of index μ' is expressed by the equation $\mu \mathbf{a} \times \mathbf{n} = \mu' \mathbf{c} \times \mathbf{n}$,

where $\mathbf{n}$, $\mathbf{a}$, $\mathbf{c}$ are the unit vectors perpendicular to the boundary, along the incident and along the refracted ray respectively.

Also find the law of reflection, if $\mathbf{b}$ be a unit vector in the direction of the reflected ray.

Let i and r be the angles of incidence and refraction respectively. Then the laws of refraction are

$$\frac{\mu}{\mu'} = \frac{\sin r}{\sin i} \quad \dots (1)$$

and $\mathbf{n}$, $\mathbf{a}$, $\mathbf{c}$ are coplanar. $\dots (2)$

Now $\mathbf{a} \times \mathbf{n} = 1 \cdot 1 \sin i \hat{\epsilon}$, i.e.,

$$\sin i = \frac{\mathbf{a} \times \mathbf{n}}{\hat{\epsilon}}$$

where $\hat{\epsilon}$ is a unit vector normal to the plane containing $\mathbf{n}$, $\mathbf{a}$, $\mathbf{c}$.




Fig. 1.27

Similarly,

$$\sin r = \frac{\mathbf{c} \times \mathbf{n}}{\hat{\mathbf{e}}}$$

From (1), we have $\mu \sin i = \mu' \sin r$,

i.e.,

$$\mu \frac{\mathbf{a} \times \mathbf{n}}{\hat{\mathbf{e}}} = \mu' \frac{\mathbf{c} \times \mathbf{n}}{\hat{\mathbf{e}}}$$

or

$$\mu \mathbf{a} \times \mathbf{n} = \mu' \mathbf{c} \times \mathbf{n}.$$

Again, the angle of incidence = angle of reflection, gives

$$\mathbf{a} \times \mathbf{n} = \mathbf{b} \times \mathbf{n}.$$

Problem 41. Decompose a vector $\mathbf{r}$ as a linear combination of vector $\mathbf{a}$, another vector perpendicular to $\mathbf{a}$, and coplanar with $\mathbf{r}$ and $\mathbf{a}$.

We know that $\mathbf{a} \times (\mathbf{a} \times \mathbf{r})$ is a vector which is coplanar with $\mathbf{a}$ and is perpendicular to $\mathbf{a}$. Let us therefore suppose that

$$\mathbf{r} = l\mathbf{a} + m\mathbf{a} \times (\mathbf{a} \times \mathbf{r}). \quad \dots (1)$$

Premultiplying both sides of (1) scalarly and vectorially with $\mathbf{a}$ we get

$$\mathbf{a} \cdot \mathbf{r} = l\mathbf{a} \cdot \mathbf{a}, \text{ i.e., } l = \frac{\mathbf{a} \cdot \mathbf{r}}{\mathbf{a} \cdot \mathbf{a}}$$

and

$$\mathbf{a} \times \mathbf{r} = m\mathbf{a} \times [\mathbf{a} \times (\mathbf{a} \times \mathbf{r})]$$

$$= m\mathbf{a} \times [(\mathbf{a} \cdot \mathbf{r})\mathbf{a} - (\mathbf{a} \cdot \mathbf{a})\mathbf{r}] = -m(\mathbf{a} \cdot \mathbf{a})\mathbf{a} \times \mathbf{r},$$

i.e.,

$$m = \frac{1}{\mathbf{a} \cdot \mathbf{a}}$$

Hence from (1), (2) and (3), we have

$$\mathbf{r} = \frac{\mathbf{a} \cdot \mathbf{r}}{\mathbf{a} \cdot \mathbf{a}} \mathbf{a} - \frac{1}{\mathbf{a} \cdot \mathbf{a}} [\mathbf{a} \times (\mathbf{a} \times \mathbf{r})].$$

Problem 42. Prove that $[\mathbf{l}\mathbf{m}\mathbf{n}] [\mathbf{l}'\mathbf{m}'\mathbf{n}'] = \begin{vmatrix} \mathbf{l} \cdot \mathbf{l}' & \mathbf{l} \cdot \mathbf{m}' & \mathbf{l} \cdot \mathbf{n}' \\ \mathbf{m} \cdot \mathbf{l}' & \mathbf{m} \cdot \mathbf{m}' & \mathbf{m} \cdot \mathbf{n}' \\ \mathbf{n} \cdot \mathbf{l}' & \mathbf{n} \cdot \mathbf{m}' & \mathbf{n} \cdot \mathbf{n}' \end{vmatrix},$

where $\mathbf{l}, \mathbf{m}, \mathbf{n}; \mathbf{l}', \mathbf{m}', \mathbf{n}'$ are any vectors.

Suppose that $\mathbf{l}' \times \mathbf{m}' = \mathbf{p}$ and consider the four vectors $\mathbf{l}, \mathbf{m}, \mathbf{n}$, which can be connected linearly as

$$[\mathbf{l}\mathbf{m}\mathbf{n}] \mathbf{p} = [\mathbf{m}\mathbf{n}\mathbf{p}] \mathbf{l} - [\mathbf{l}\mathbf{n}\mathbf{p}] \mathbf{m} + [\mathbf{l}\mathbf{m}\mathbf{p}] \mathbf{n}.$$

Here $[\mathbf{m}\mathbf{n}\mathbf{p}] = \mathbf{m} \times \mathbf{n} \cdot \mathbf{p} = (\mathbf{m} \times \mathbf{n}) \cdot (\mathbf{l}' \times \mathbf{m}') = \begin{vmatrix} \mathbf{m} \cdot \mathbf{l}' & \mathbf{m} \cdot \mathbf{m}' \\ \mathbf{n} \cdot \mathbf{l}' & \mathbf{n} \cdot \mathbf{m}' \end{vmatrix},$

$$[\mathbf{l}\mathbf{n}\mathbf{p}] = \mathbf{l} \times \mathbf{n} \cdot \mathbf{p} = (\mathbf{l} \times \mathbf{n}) \cdot (\mathbf{l}' \times \mathbf{m}') = \begin{vmatrix} \mathbf{l} \cdot \mathbf{l}' & \mathbf{l} \cdot \mathbf{m}' \\ \mathbf{n} \cdot \mathbf{l}' & \mathbf{n} \cdot \mathbf{m}' \end{vmatrix}$$

and $[\mathbf{l}\mathbf{m}\mathbf{p}] = \mathbf{l} \times \mathbf{m} \cdot \mathbf{p} = (\mathbf{l} \times \mathbf{m}) \cdot (\mathbf{l}' \times \mathbf{m}') = \begin{vmatrix} \mathbf{l} \cdot \mathbf{l}' & \mathbf{l} \cdot \mathbf{m}' \\ \mathbf{m} \cdot \mathbf{l}' & \mathbf{m} \cdot \mathbf{m}' \end{vmatrix}$

With these substitutions (1) becomes

$$[\mathbf{l}\mathbf{m}\mathbf{n}] \mathbf{p} = [\mathbf{l}\mathbf{m}\mathbf{n}] \mathbf{l}' \times \mathbf{m}' = \begin{vmatrix} \mathbf{m} \cdot \mathbf{l}' & \mathbf{m} \cdot \mathbf{m}' \\ \mathbf{n} \cdot \mathbf{l}' & \mathbf{n} \cdot \mathbf{m}' \end{vmatrix} \mathbf{l} - \begin{vmatrix} \mathbf{l} \cdot \mathbf{l}' & \mathbf{l} \cdot \mathbf{m}' \\ \mathbf{n} \cdot \mathbf{l}' & \mathbf{n} \cdot \mathbf{m}' \end{vmatrix} \mathbf{m} + \begin{vmatrix} \mathbf{l} \cdot \mathbf{l}' & \mathbf{l} \cdot \mathbf{m}' \\ \mathbf{m} \cdot \mathbf{l}' & \mathbf{m} \cdot \mathbf{m}' \end{vmatrix} \mathbf{n}.$$

Multiplying scalarly both sides by $\mathbf{n}'$, we get

$$[\mathbf{l}\mathbf{m}\mathbf{n}] \mathbf{l}' \times \mathbf{m}' \cdot \mathbf{n}' = \begin{vmatrix} \mathbf{m} \cdot \mathbf{l}' & \mathbf{m} \cdot \mathbf{m}' \\ \mathbf{n} \cdot \mathbf{l}' & \mathbf{n} \cdot \mathbf{m}' \end{vmatrix} \mathbf{l} \cdot \mathbf{n}' - \begin{vmatrix} \mathbf{l} \cdot \mathbf{l}' & \mathbf{l} \cdot \mathbf{m}' \\ \mathbf{n} \cdot \mathbf{l}' & \mathbf{n} \cdot \mathbf{m}' \end{vmatrix} \mathbf{m} \cdot \mathbf{n}' + \begin{vmatrix} \mathbf{l} \cdot \mathbf{l}' & \mathbf{l} \cdot \mathbf{m}' \\ \mathbf{m} \cdot \mathbf{l}' & \mathbf{m} \cdot \mathbf{m}' \end{vmatrix} \mathbf{n} \cdot \mathbf{n}'$$

or $[\mathbf{l}\mathbf{m}\mathbf{n}] [\mathbf{l}'\mathbf{m}'\mathbf{n}'] = \begin{vmatrix} \mathbf{l} \cdot \mathbf{l}' & \mathbf{l} \cdot \mathbf{m}' & \mathbf{l} \cdot \mathbf{n}' \\ \mathbf{m} \cdot \mathbf{l}' & \mathbf{m} \cdot \mathbf{m}' & \mathbf{m} \cdot \mathbf{n}' \\ \mathbf{n} \cdot \mathbf{l}' & \mathbf{n} \cdot \mathbf{m}' & \mathbf{n} \cdot \mathbf{n}' \end{vmatrix}.$

Note: If $\mathbf{l} = \mathbf{l}' = \mathbf{a},$
 $\mathbf{m} = \mathbf{m}' = \mathbf{b},$
 $\mathbf{n} = \mathbf{n}' = \mathbf{c},$ } then $[\mathbf{a}\mathbf{b}\mathbf{c}]^2 = \begin{vmatrix} \mathbf{a} \cdot \mathbf{a} & \mathbf{a} \cdot \mathbf{b} & \mathbf{a} \cdot \mathbf{c} \\ \mathbf{b} \cdot \mathbf{a} & \mathbf{b} \cdot \mathbf{b} & \mathbf{b} \cdot \mathbf{c} \\ \mathbf{c} \cdot \mathbf{a} & \mathbf{c} \cdot \mathbf{b} & \mathbf{c} \cdot \mathbf{c} \end{vmatrix}$

and **Problem 43.** Prove that

$$(\mathbf{b} \times \mathbf{c}) \cdot (\mathbf{a} \times \mathbf{d}) + (\mathbf{c} \times \mathbf{a}) \cdot (\mathbf{b} \times \mathbf{d}) + (\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = 0$$

and deduce that

$$\sin(A+B) \sin(A-B) = \sin^2 A - \sin^2 B = \frac{1}{2} (\cos 2B - \cos 2A).$$

Here

$$(\mathbf{b} \times \mathbf{c}) \cdot (\mathbf{a} \times \mathbf{d}) + (\mathbf{c} \times \mathbf{a}) \cdot (\mathbf{b} \times \mathbf{d}) + (\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d})$$

$$= \begin{vmatrix} \mathbf{b} \cdot \mathbf{a} & \mathbf{b} \cdot \mathbf{d} \\ \mathbf{c} \cdot \mathbf{a} & \mathbf{c} \cdot \mathbf{d} \end{vmatrix} + \begin{vmatrix} \mathbf{c} \cdot \mathbf{b} & \mathbf{c} \cdot \mathbf{d} \\ \mathbf{a} \cdot \mathbf{b} & \mathbf{a} \cdot \mathbf{d} \end{vmatrix} + \begin{vmatrix} \mathbf{a} \cdot \mathbf{c} & \mathbf{a} \cdot \mathbf{d} \\ \mathbf{b} \cdot \mathbf{c} & \mathbf{b} \cdot \mathbf{d} \end{vmatrix}$$

$$= (\mathbf{b} \cdot \mathbf{a})(\mathbf{c} \cdot \mathbf{d}) - (\mathbf{b} \cdot \mathbf{d})(\mathbf{c} \cdot \mathbf{a}) + (\mathbf{c} \cdot \mathbf{b})(\mathbf{a} \cdot \mathbf{d}) - (\mathbf{c} \cdot \mathbf{d})(\mathbf{a} \cdot \mathbf{b}) + (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \cdot \mathbf{d})(\mathbf{b} \cdot \mathbf{c})$$

$$= 0 \text{ as } \mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a} \text{ etc.}$$

For the second part, let $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ be four coplanar vectors and $\hat{\mathbf{e}}$ be a unit vector in the direction perpendicular to the plane containing $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$.

Let the angles between the directions of $\mathbf{a}$ and $\mathbf{b}$, $\mathbf{b}$ and $\mathbf{c}$, $\mathbf{c}$ and $\mathbf{d}$ be $\theta_1, \theta_2, \theta_3$ respectively.

Since $\mathbf{b} \times \mathbf{c} = bc \sin \theta_2 \hat{\mathbf{e}}$ etc., a, b, c, d being modules of $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ respectively,

$$(\mathbf{b} \times \mathbf{c}) \cdot (\mathbf{a} \times \mathbf{d}) + (\mathbf{c} \times \mathbf{a}) \cdot (\mathbf{b} \times \mathbf{d}) + (\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = 0 \text{ gives}$$

$$(bc \sin \theta_2 \hat{\mathbf{e}}) \cdot (ad \sin (\theta_1 + \theta_2 + \theta_3) \hat{\mathbf{e}})$$

$$+ [-ca \sin (\theta_1 + \theta_2) \hat{\mathbf{e}}] \cdot [bd \sin (\theta_2 + \theta_3) \hat{\mathbf{e}}]$$

$$+ (ab \sin \theta_1 \hat{\mathbf{e}}) \cdot (cd \sin \theta_3 \hat{\mathbf{e}}) = 0$$

$$\text{or } \sin \theta_2 \sin (\theta_1 + \theta_2 + \theta_3) - \sin (\theta_1 + \theta_2) \sin (\theta_2 + \theta_3) + \sin \theta_1 \sin \theta_3 =$$

$$[\because \hat{\mathbf{e}} \cdot \hat{\mathbf{e}} = 1].$$

Putting $\theta_1 = B, \theta_2 = A$ and $\theta_3 = -B$, this gives

$$\sin^2 A - \sin(A+B) \sin(A-B) - \sin^2 B = 0,$$




Fig. 1.28

$$\begin{aligned} \text{i.e., } \sin(A+B) \sin(A-B) &= \sin^2 A - \sin^2 B \\ &= \frac{1}{2}(1 - \cos 2A) - \frac{1}{2}(1 - \cos 2B) \\ &= \frac{1}{2}(\cos 2B - \cos 2A). \end{aligned}$$

Problem 44. Prove that $2(\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) = \begin{vmatrix} -\mathbf{a} & -\mathbf{b} & \mathbf{c} & \mathbf{d} \\ a_1 & b_1 & c_1 & d_1 \\ a_2 & b_2 & c_2 & d_2 \\ a_3 & b_3 & c_3 & d_3 \end{vmatrix}$,

where $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$ etc.

Since $(\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) = [\mathbf{abd}] \mathbf{c} - [\mathbf{abc}] \mathbf{d}$

$$= \begin{vmatrix} a_1 & b_1 & d_1 \\ a_2 & b_2 & d_2 \\ a_3 & b_3 & d_3 \end{vmatrix} \mathbf{c} - \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} \mathbf{d} \dots$$

Also $(\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) = [\mathbf{acd}] \mathbf{b} - [\mathbf{bcd}] \mathbf{a}$

$$= \begin{vmatrix} a_1 & c_1 & d_1 \\ a_2 & c_2 & d_2 \\ a_3 & c_3 & d_3 \end{vmatrix} \mathbf{b} - \begin{vmatrix} b_1 & c_1 & d_1 \\ b_2 & c_2 & d_2 \\ b_3 & c_3 & d_3 \end{vmatrix} \mathbf{a} \dots$$

Adding (1) and (2), we get

$$2(\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) = \begin{vmatrix} -\mathbf{a} & -\mathbf{b} & \mathbf{c} & \mathbf{d} \\ a_1 & b_1 & c_1 & d_1 \\ a_2 & b_2 & c_2 & d_2 \\ a_3 & b_3 & c_3 & d_3 \end{vmatrix}$$

Problem 45. If OX, OY, OZ and $O'X', O'Y', O'Z'$ are two sets of rectangular co-ordinate axes and $l_1, m_1, n_1; l_2, m_2, n_2; l_3, m_3, n_3$ denote the direction cosines of the members of either set with respect to other

then

$$\begin{vmatrix} l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \\ l_3 & m_3 & n_3 \end{vmatrix} = 1.$$

Let $\mathbf{i}, \mathbf{j}, \mathbf{k}; \mathbf{i}', \mathbf{j}', \mathbf{k}'$ be unit vectors along the two sets of axes.

Then using the adjoining scheme of transformation, we have

$$\begin{aligned} \mathbf{i}' &= l_1\mathbf{i} + m_1\mathbf{j} + n_1\mathbf{k}, \\ \mathbf{j}' &= l_2\mathbf{i} + m_2\mathbf{j} + n_2\mathbf{k}, \\ \mathbf{k}' &= l_3\mathbf{i} + m_3\mathbf{j} + n_3\mathbf{k}. \end{aligned}$$

	$\mathbf{i}$	$\mathbf{j}$	$\mathbf{k}$
$\mathbf{i}'$	l_1	m_1	n_1
$\mathbf{j}'$	l_2	m_2	n_2
$\mathbf{k}'$	l_3	m_3	n_3

Now $\mathbf{j}' \times \mathbf{k}' = (l_2\mathbf{i} + m_2\mathbf{j} + n_2\mathbf{k}) \times (l_3\mathbf{i} + m_3\mathbf{j} + n_3\mathbf{k})$
 $= (n_2m_3 - m_2n_3) \mathbf{j} \times \mathbf{k} + (n_2l_3 - n_3l_2) \mathbf{k} \times \mathbf{i}$
 $+ (l_2m_3 - l_2m_2) \mathbf{i} \times \mathbf{j}$

Multiplying scalarly by $\mathbf{i}'$, we get

$$\mathbf{i}' \cdot \mathbf{j}' \times \mathbf{k}' = (l_1\mathbf{i} + m_1\mathbf{j} + n_1\mathbf{k}) \cdot [(n_2m_3 - m_2n_3) \mathbf{j} \times \mathbf{k} + (n_2l_3 - n_3l_2) \mathbf{k} \times \mathbf{i} + (l_2m_3 - l_2m_2) \mathbf{i} \times \mathbf{j}]$$

$$\begin{aligned} \mathbf{i}' \cdot \mathbf{j}' \times \mathbf{k}' &= l_1(n_2m_3 - m_2n_3) [\mathbf{i} \cdot \mathbf{j} \times \mathbf{k}] + m_1(n_2l_3 - n_3l_2) [\mathbf{j} \cdot \mathbf{k} \times \mathbf{i}] \\ &\quad + n_1(l_2m_3 - l_3m_2) [\mathbf{k} \cdot \mathbf{i} \times \mathbf{j}] \end{aligned}$$

$$= \begin{vmatrix} l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \\ l_3 & m_3 & n_3 \end{vmatrix} [\mathbf{i} \cdot \mathbf{j} \times \mathbf{k}].$$

Since $[\mathbf{i}' \mathbf{j}' \mathbf{k}'] = [\mathbf{i} \mathbf{j} \mathbf{k}]$, we have $\begin{vmatrix} l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \\ l_3 & m_3 & n_3 \end{vmatrix} = 1.$

1.13. PRODUCT OF FOUR VECTORS

(1) **Scalar product of four vectors.** If $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ are four vectors then the product of the type $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d})$ is called scalar product of four vectors. In fact this being a scalar product of two vectors $(\mathbf{a} \times \mathbf{b})$ and $(\mathbf{c} \times \mathbf{d})$, is a scalar quantity. We can treat this product as a scalar triple product of three vectors $\mathbf{a}, \mathbf{b}$ and $(\mathbf{c} \times \mathbf{d})$.

Since the dot and cross may be interchanged in a scalar triple product, we have

$$\begin{aligned} (\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) &= \mathbf{a} \cdot \mathbf{b} \times (\mathbf{c} \times \mathbf{d}) \\ &= \mathbf{a} \cdot [(\mathbf{b} \cdot \mathbf{d}) \mathbf{c} - (\mathbf{b} \cdot \mathbf{c}) \mathbf{d}] \\ &= (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \cdot \mathbf{d})(\mathbf{b} \cdot \mathbf{c}) \\ &= \begin{vmatrix} \mathbf{a} \cdot \mathbf{c} & \mathbf{b} \cdot \mathbf{c} \\ \mathbf{a} \cdot \mathbf{d} & \mathbf{b} \cdot \mathbf{d} \end{vmatrix}. \end{aligned}$$

Problem 46. Show that

$$(\mathbf{b} \times \mathbf{c}) \cdot (\mathbf{a} \times \mathbf{d}) + (\mathbf{c} \times \mathbf{a}) \cdot (\mathbf{b} \times \mathbf{d}) + (\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = 0,$$

We have $(\mathbf{b} \times \mathbf{c}) \cdot (\mathbf{a} \times \mathbf{d}) = \begin{vmatrix} \mathbf{b} \cdot \mathbf{a} & \mathbf{c} \cdot \mathbf{a} \\ \mathbf{b} \cdot \mathbf{d} & \mathbf{c} \cdot \mathbf{d} \end{vmatrix}$
 $= (\mathbf{b} \cdot \mathbf{a})(\mathbf{c} \cdot \mathbf{d}) - (\mathbf{b} \cdot \mathbf{d})(\mathbf{c} \cdot \mathbf{a}).$

Similarly $(\mathbf{c} \times \mathbf{a}) \cdot (\mathbf{b} \times \mathbf{d}) = (\mathbf{c} \cdot \mathbf{b})(\mathbf{a} \cdot \mathbf{d}) - (\mathbf{c} \cdot \mathbf{d})(\mathbf{a} \cdot \mathbf{b})$

and $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \cdot \mathbf{d})(\mathbf{b} \cdot \mathbf{c}).$

Adding all together, we get

$$(\mathbf{b} \times \mathbf{c}) \cdot (\mathbf{a} \times \mathbf{d}) + (\mathbf{c} \times \mathbf{a}) \cdot (\mathbf{b} \times \mathbf{d}) + (\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = 0.$$

(2) **Vector product of four vectors.** If $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ are four vectors then the product of the type $(\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d})$ is called vector product of four vectors. The value of this vector product is a vector which is at right angles to the vectors $(\mathbf{a} \times \mathbf{b})$ and $(\mathbf{c} \times \mathbf{d})$ both, and therefore is coplanar with $\mathbf{a}, \mathbf{b}$ and also $\mathbf{c}, \mathbf{d}$. Conclusively this vector is parallel to the line of intersection of a plane parallel to $\mathbf{a}$ and $\mathbf{b}$ with another plane parallel to $\mathbf{c}$ and $\mathbf{d}$.

The value of this vector product may be obtained in two ways:

(i) If we put $\mathbf{c} \times \mathbf{d} = \mathbf{p}$, then

$$\begin{aligned} (\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) &= (\mathbf{a} \times \mathbf{b}) \times \mathbf{p} \\ &= (\mathbf{a} \cdot \mathbf{p}) \mathbf{b} - (\mathbf{b} \cdot \mathbf{p}) \mathbf{a} \\ &= (\mathbf{a} \cdot \mathbf{c} \times \mathbf{d}) \mathbf{b} - (\mathbf{b} \cdot \mathbf{c} \times \mathbf{d}) \mathbf{a} \\ &= [\mathbf{a} \mathbf{c} \mathbf{d}] \mathbf{b} - [\mathbf{b} \mathbf{c} \mathbf{d}] \mathbf{a} \end{aligned}$$

(ii) If we put $\mathbf{a} \times \mathbf{b} = \mathbf{q}$, then

$$\begin{aligned} (\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) &= \mathbf{q} \times (\mathbf{c} \times \mathbf{d}) \\ &= (\mathbf{q} \cdot \mathbf{d}) \mathbf{c} - (\mathbf{q} \cdot \mathbf{c}) \mathbf{d} \\ &= (\mathbf{a} \times \mathbf{b} \cdot \mathbf{d}) \mathbf{c} - (\mathbf{a} \times \mathbf{b} \cdot \mathbf{c}) \mathbf{d} \\ &= [\mathbf{a} \mathbf{b} \mathbf{d}] \mathbf{c} - [\mathbf{a} \mathbf{b} \mathbf{c}] \mathbf{d} \end{aligned}$$

Linear relationship between four vectors.

The two results namely (i) and (ii), on subtraction, yield

$$0 = [\mathbf{b} \mathbf{c} \mathbf{d}] \mathbf{a} - [\mathbf{a} \mathbf{c} \mathbf{d}] \mathbf{b} + [\mathbf{a} \mathbf{b} \mathbf{d}] \mathbf{c} - [\mathbf{a} \mathbf{b} \mathbf{c}] \mathbf{d}$$

Rearranging, we get

$$\mathbf{d} = \frac{[\mathbf{d} \mathbf{b} \mathbf{c}] \mathbf{a} + [\mathbf{d} \mathbf{c} \mathbf{a}] \mathbf{b} + [\mathbf{d} \mathbf{a} \mathbf{b}] \mathbf{c}}{[\mathbf{a} \mathbf{b} \mathbf{c}]} \text{ provided } [\mathbf{a} \mathbf{b} \mathbf{c}] \neq 0.$$

This may also be expressed as

$$\mathbf{d} = \mathbf{d} \cdot \frac{\mathbf{b} \times \mathbf{c}}{[\mathbf{a} \mathbf{b} \mathbf{c}]} \mathbf{a} + \mathbf{d} \cdot \frac{\mathbf{c} \times \mathbf{a}}{[\mathbf{a} \mathbf{b} \mathbf{c}]} \mathbf{b} + \mathbf{d} \cdot \frac{\mathbf{a} \times \mathbf{b}}{[\mathbf{a} \mathbf{b} \mathbf{c}]} \mathbf{c}.$$

Note. As an alternative, the linear relationship between four non-coplanar vectors $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ may be found as follows:

Suppose $\mathbf{d} = \lambda \mathbf{a} + \mu \mathbf{b} + \nu \mathbf{c}$, where λ, μ, ν are scalars.

Multiplying both sides scalarly by $(\mathbf{b} \times \mathbf{c})$, we get

$$[\mathbf{d} \mathbf{b} \mathbf{c}] = \lambda [\mathbf{a} \mathbf{b} \mathbf{c}], \text{ other terms vanishing as } [\mathbf{b} \mathbf{b} \mathbf{c}] = 0 = [\mathbf{c} \mathbf{b} \mathbf{c}]$$

$$\text{or } \lambda = \frac{[\mathbf{d} \mathbf{b} \mathbf{c}]}{[\mathbf{a} \mathbf{b} \mathbf{c}]}$$

$$\text{Similarly } \mu = \frac{[\mathbf{d} \mathbf{c} \mathbf{a}]}{[\mathbf{a} \mathbf{b} \mathbf{c}]} \text{ and } \nu = \frac{[\mathbf{d} \mathbf{a} \mathbf{b}]}{[\mathbf{a} \mathbf{b} \mathbf{c}]}.$$

Substituting the values of λ, μ, ν , we have

$$\mathbf{d} = \frac{[\mathbf{d} \mathbf{b} \mathbf{c}] \mathbf{a} + [\mathbf{d} \mathbf{c} \mathbf{a}] \mathbf{b} + [\mathbf{d} \mathbf{a} \mathbf{b}] \mathbf{c}}{[\mathbf{a} \mathbf{b} \mathbf{c}]}.$$

Problem 47. Prove that $\mathbf{d} \cdot [\mathbf{a} \times \{\mathbf{b} \times (\mathbf{c} \times \mathbf{d})\}] = (\mathbf{b} \cdot \mathbf{d}) [\mathbf{a} \mathbf{c} \mathbf{d}]$.

Here $\mathbf{d} \cdot [\mathbf{a} \times \{\mathbf{b} \times (\mathbf{c} \times \mathbf{d})\}] = \mathbf{d} \cdot [\mathbf{a} \times \{(\mathbf{b} \cdot \mathbf{d}) \mathbf{c} - (\mathbf{b} \cdot \mathbf{c}) \mathbf{d}\}]$

$$\begin{aligned} &= \mathbf{d} \cdot [(\mathbf{b} \cdot \mathbf{d}) (\mathbf{a} \times \mathbf{c}) - (\mathbf{b} \cdot \mathbf{c}) (\mathbf{a} \times \mathbf{d})] \\ &= (\mathbf{b} \cdot \mathbf{d}) (\mathbf{a} \times \mathbf{c} \cdot \mathbf{d}) \text{ as } \mathbf{a} \times \mathbf{d} \cdot \mathbf{d} = [\mathbf{a} \mathbf{d} \mathbf{d}] = 0 \\ &= (\mathbf{b} \cdot \mathbf{d}) [\mathbf{a} \mathbf{c} \mathbf{d}]. \end{aligned}$$

Problem 48. Prove the identity

$$\mathbf{a} \times [\mathbf{b} \times (\mathbf{c} \times \mathbf{d})] = (\mathbf{a} \times \mathbf{c}) (\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \times \mathbf{d}) (\mathbf{b} \cdot \mathbf{c}).$$

L.H.S. $= \mathbf{a} \times [\mathbf{b} \times (\mathbf{c} \times \mathbf{d})]$

$$\begin{aligned} &= \mathbf{a} \times [(\mathbf{b} \cdot \mathbf{d}) \mathbf{c} - (\mathbf{b} \cdot \mathbf{c}) \mathbf{d}] \\ &= (\mathbf{b} \cdot \mathbf{d}) (\mathbf{a} \times \mathbf{c}) - (\mathbf{b} \cdot \mathbf{c}) (\mathbf{a} \times \mathbf{d}) \\ &= \text{R.H.S.} \end{aligned}$$

1.14. RECIPROCAL SYSTEM OF VECTORS

If there are two sets of non-coplanar vectors $\mathbf{a}, \mathbf{b}, \mathbf{c}$ and $\mathbf{a}', \mathbf{b}', \mathbf{c}'$ such that

VECTORS

$$\mathbf{a}' = \frac{\mathbf{b} \times \mathbf{c}}{[\mathbf{a} \mathbf{b} \mathbf{c}]}, \mathbf{b}' = \frac{\mathbf{c} \times \mathbf{a}}{[\mathbf{a} \mathbf{b} \mathbf{c}]}, \mathbf{c}' = \frac{\mathbf{a} \times \mathbf{b}}{[\mathbf{a} \mathbf{b} \mathbf{c}]},$$

then $\mathbf{a}, \mathbf{b}, \mathbf{c}$ and $\mathbf{a}', \mathbf{b}', \mathbf{c}'$ are said to be *Reciprocal systems* of vectors. It is so called, because

$$\mathbf{a} \cdot \mathbf{a}' = \mathbf{b} \cdot \mathbf{b}' = \mathbf{c} \cdot \mathbf{c}' = 1 \text{ since } \mathbf{a} \cdot \frac{\mathbf{b} \times \mathbf{c}}{[\mathbf{a} \mathbf{b} \mathbf{c}]} = \frac{[\mathbf{a} \mathbf{b} \mathbf{c}]}{[\mathbf{a} \mathbf{b} \mathbf{c}]} = 1 \text{ etc.,}$$

so that

$$\mathbf{a}' = \mathbf{a}^{-1}, \mathbf{b}' = \mathbf{b}^{-1}, \mathbf{c}' = \mathbf{c}^{-1}.$$

In fact the two systems of vectors, i.e. $\mathbf{a}, \mathbf{b}, \mathbf{c}$ and $\mathbf{a}', \mathbf{b}', \mathbf{c}'$ are mutually reciprocal as

$$\mathbf{a} = \frac{\mathbf{b}' \times \mathbf{c}'}{[\mathbf{a}' \mathbf{b}' \mathbf{c}']}, \mathbf{b} = \frac{\mathbf{c}' \times \mathbf{a}'}{[\mathbf{a}' \mathbf{b}' \mathbf{c}']}, \mathbf{c} = \frac{\mathbf{a}' \times \mathbf{b}'}{[\mathbf{a}' \mathbf{b}' \mathbf{c}']}.$$

In particular if $\mathbf{i}, \mathbf{j}, \mathbf{k}$ be the unit vectors along the principal axes and $\mathbf{i}', \mathbf{j}', \mathbf{k}'$ their reciprocals, then

$$\mathbf{i}' = \mathbf{i}, \mathbf{j}' = \mathbf{j} \text{ and } \mathbf{k}' = \mathbf{k} \text{ as } [\mathbf{i} \mathbf{j} \mathbf{k}] = 1.$$

This is called *self-reciprocal system*.

Note With these notations, the linear relationship between four vectors $\mathbf{a}, \mathbf{b}, \mathbf{c}$ and $\mathbf{d}$ may be expressed as

$$\mathbf{d} = (\mathbf{d} \cdot \mathbf{a}') \mathbf{a} + (\mathbf{d} \cdot \mathbf{b}') \mathbf{b} + (\mathbf{d} \cdot \mathbf{c}') \mathbf{c}.$$

Problem 49. Prove that $(\mathbf{a} \times \mathbf{a}') + (\mathbf{b} \times \mathbf{b}') + (\mathbf{c} \times \mathbf{c}') = 0$, where $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are vectors and $\mathbf{a}', \mathbf{b}', \mathbf{c}'$ their reciprocals.

We have

$$\mathbf{a}' = \frac{\mathbf{b} \times \mathbf{c}}{[\mathbf{a} \mathbf{b} \mathbf{c}]}, \mathbf{b}' = \frac{\mathbf{c} \times \mathbf{a}}{[\mathbf{a} \mathbf{b} \mathbf{c}]}, \mathbf{c}' = \frac{\mathbf{a} \times \mathbf{b}}{[\mathbf{a} \mathbf{b} \mathbf{c}]}.$$

$$\mathbf{a} \times \mathbf{a}' = \mathbf{a} \times \frac{(\mathbf{b} \times \mathbf{c})}{[\mathbf{a} \mathbf{b} \mathbf{c}]} = \frac{1}{[\mathbf{a} \mathbf{b} \mathbf{c}]} [(\mathbf{a} \cdot \mathbf{c}) \mathbf{b} - (\mathbf{a} \cdot \mathbf{b}) \mathbf{c}].$$

Similarly

$$\mathbf{b} \times \mathbf{b}' = \frac{1}{[\mathbf{a} \mathbf{b} \mathbf{c}]} [(\mathbf{b} \cdot \mathbf{a}) \mathbf{c} - (\mathbf{b} \cdot \mathbf{c}) \mathbf{a}]$$

and

$$\mathbf{c} \times \mathbf{c}' = \frac{1}{[\mathbf{a} \mathbf{b} \mathbf{c}]} [(\mathbf{c} \cdot \mathbf{b}) \mathbf{a} - (\mathbf{c} \cdot \mathbf{a}) \mathbf{b}].$$

Adding all together, we get

$$(\mathbf{a} \times \mathbf{a}') + (\mathbf{b} \times \mathbf{b}') + (\mathbf{c} \times \mathbf{c}') = \frac{1}{[\mathbf{a} \mathbf{b} \mathbf{c}]} [0] = 0.$$

1.15. VECTOR EQUATIONS

Here below some methods for solving the vector equations are explained with the help of examples.

Problem 50. Solve the vector equation $\mathbf{x} \times \mathbf{a} = \mathbf{b}$.

Given equation is $\mathbf{x} \times \mathbf{a} = \mathbf{b}$ (1)

We know that $\mathbf{a} \times \mathbf{b}$ is a vector perpendicular to both $\mathbf{a}$ and $\mathbf{b}$, therefore the vectors $\mathbf{a}, \mathbf{b}$ and $\mathbf{a} \times \mathbf{b}$ are three non-coplanar vectors. Let us assume that the solution of the given equation is of the form

$$\mathbf{x} = \lambda \mathbf{a} + \mu \mathbf{b} + \nu (\mathbf{a} \times \mathbf{b}), \quad \dots (2)$$

where λ, μ, ν are scalars.

Since (2) is a solution of (1), therefore substituting in (1) the value of $\mathbf{x}$ from (2), we get

$$\begin{aligned} & \{\lambda \mathbf{a} + \mu \mathbf{b} + \nu (\mathbf{a} \times \mathbf{b})\} \times \mathbf{a} = \mathbf{b}, \\ \text{or } & \mu (\mathbf{b} \times \mathbf{a}) + \nu \{(\mathbf{a} \times \mathbf{b}) \times \mathbf{a}\} = \mathbf{b} \text{ as } \mathbf{a} \times \mathbf{a} = 0 \\ \text{or } & -\mu (\mathbf{a} \times \mathbf{b}) + \nu \{(\mathbf{a} \cdot \mathbf{a}) \mathbf{b} - (\mathbf{b} \cdot \mathbf{a}) \mathbf{a}\} = \mathbf{b} \\ \text{or } & -\mu (\mathbf{a} \times \mathbf{b}) + \nu \{a^2 \mathbf{b} - (\mathbf{b} \cdot \mathbf{a}) \mathbf{a}\} = \mathbf{b}, \text{ where } |\mathbf{a}| = a \end{aligned}$$

Equating the coefficients of like vectors on either side, we get

$$\begin{aligned} & -\mu = 0, \nu a^2 = 1, -\nu \mathbf{b} \cdot \mathbf{a} = 0, \\ \text{i.e. } & \mu = 0, \nu = 1/a^2 \text{ and } \mathbf{a} \cdot \mathbf{b} = 0 \text{ as } \nu \neq 0. \end{aligned}$$

Substituting in (2), these values of μ and ν , the general solution of the given equation is

$$\mathbf{x} = \lambda \mathbf{a} - 1/a^2 (\mathbf{a} \times \mathbf{b})$$

and the condition for the existence of this solution is

$$\mathbf{a} \cdot \mathbf{b} = 0,$$

i.e. the vectors $\mathbf{a}$ and $\mathbf{b}$ are mutually at right angles.

Problem 51. Solve the simultaneous equations

$$\mathbf{x} \times \mathbf{b} = \mathbf{a} \times \mathbf{b}, \mathbf{x} \cdot \mathbf{c} = 0 \text{ provided } \mathbf{b} \cdot \mathbf{c} \neq 0.$$

The given equations are

$$\mathbf{x} \times \mathbf{b} = \mathbf{a} \times \mathbf{b} \quad \dots (1)$$

$$\text{and } \mathbf{x} \cdot \mathbf{c} = 0. \quad \dots (2)$$

The equation (1) can be written as

$$(\mathbf{x} - \mathbf{a}) \times \mathbf{b} = 0,$$

which follows that $(\mathbf{x} - \mathbf{a})$ and $\mathbf{b}$ are parallel, i.e.

$$\mathbf{x} - \mathbf{a} = t\mathbf{b}, \text{ where } t \text{ is a scalar}$$

$$\text{or } \mathbf{x} = \mathbf{a} + t\mathbf{b}. \quad \dots (3)$$

Substituting this value of $\mathbf{x}$ in (2), we get

$$(\mathbf{a} + t\mathbf{b}) \cdot \mathbf{c} = 0.$$

$$\text{Giving } t = -\frac{\mathbf{a} \cdot \mathbf{c}}{\mathbf{b} \cdot \mathbf{c}} \text{ when } \mathbf{b} \cdot \mathbf{c} \neq 0 \text{ (given).}$$

Hence the required solution is obtained by putting the value of t in (3) and that is

$$\mathbf{x} = \mathbf{a} - \frac{\mathbf{a} \cdot \mathbf{c}}{\mathbf{b} \cdot \mathbf{c}} \mathbf{b}.$$

Problem 52. Solve the simultaneous equations

$$s\mathbf{x} + t\mathbf{y} = \mathbf{a}, \mathbf{x} \times \mathbf{y} = \mathbf{b} \text{ provided } \mathbf{a} \cdot \mathbf{b} = 0.$$

The given equations are

$$s\mathbf{x} + t\mathbf{y} = \mathbf{a}, \quad \dots (1)$$

$$\mathbf{x} \times \mathbf{y} = \mathbf{b}. \quad \dots (2)$$

Multiplying (1) vectorially by $\mathbf{x}$, we get

$$\mathbf{x} \times (s\mathbf{x} + t\mathbf{y}) = \mathbf{x} \times \mathbf{a}$$

$$\text{or } t\mathbf{x} \times \mathbf{y} = \mathbf{x} \times \mathbf{a} \text{ as } \mathbf{x} \times \mathbf{x} = 0$$

$$\text{or } \mathbf{x} \times \mathbf{a} = t\mathbf{b} \text{ from (2).} \quad \dots (3)$$

Multiplying (3) vectorially by $\mathbf{a}$, we have

$$\mathbf{a} \times (\mathbf{x} \times \mathbf{a}) = t (\mathbf{a} \times \mathbf{b})$$

$$(\mathbf{a} \cdot \mathbf{a}) \mathbf{x} - (\mathbf{a} \cdot \mathbf{x}) \mathbf{a} = t (\mathbf{a} \times \mathbf{b})$$

$$\mathbf{x} - \lambda \mathbf{a} = t \frac{(\mathbf{a} \times \mathbf{b})}{a^2}$$

where λ is a scalar parameter,

$$\mathbf{x} = \lambda \mathbf{a} + \frac{t}{a^2} (\mathbf{a} \times \mathbf{b})$$

which is the general solution for $\mathbf{x}$. Similar procedure will yield the solution for $\mathbf{y}$.

1.16. SIMPLE APPLICATIONS OF VECTORS TO MECHANICS

(1) **Concurrent forces.** It is found experimentally that the resultant of two concurrent forces is equivalent to a single force acting at the same point. The single force is represented by their vector sum. In general a system of forces acting at a point and represented by the vectors $\mathbf{P}_1, \mathbf{P}_2, \mathbf{P}_3, \dots, \mathbf{P}_n$ is equivalent to a single resultant force $\mathbf{F}$ acting at the same point such that

$$\mathbf{F} = \mathbf{P}_1 + \mathbf{P}_2 + \mathbf{P}_3 + \dots + \mathbf{P}_n.$$

It can be obtained by constructing a vector polygon of which

$$\overrightarrow{AB} = \mathbf{P}_1, \overrightarrow{BC} = \mathbf{P}_2, \dots, \overrightarrow{MN} = \mathbf{P}_n.$$

The resultant is represented by $\overrightarrow{AN} = \mathbf{F}$, which is drawn to close up the polygon opposite to that in which the sides have been drawn. The polygon does not necessarily lie in a plane as the forces, may not be coplanar.




Fig. 1.29

COLLARY. Lami's theorem. If three forces acting at a point be in equilibrium, then each is proportional to the sine of the angle between the other two.

In the case of three concurrent forces the closed polygon will be a triangle, the forces will be coplanar and each side is proportional to the sine of the opposite angle and hence each force will be proportional to the sine of the angle between the other two.

(2) **Work done by a force.** We know that a force acting on a particle does work when the displacement of the particle takes place in the direction which is not perpendicular to the direction of the force. The work done is measured by the product of the force and the resolved part of the displacement in the direction of the force. Hence if the vectors representing the force and the displacement be respectively $\mathbf{F}$ and $\mathbf{d}$ inclined at an angle θ and whose respective modules are F and d , then the work done $Fd \cos \theta = \mathbf{F} \cdot \mathbf{d}$ (by the definition of dot product).

In case $\mathbf{d}$ is perpendicular to $\mathbf{F}$, i.e. $\theta = 90^\circ$, the work done is zero.

COROLLARY. Rate of doing work. If a particle acted upon by a force $\mathbf{F}$ is moving with a velocity $\mathbf{v}$, then the rate of work done, $\frac{dW}{dt}$ is given by

$$\frac{dW}{dt} = \mathbf{F} \cdot \mathbf{v}.$$

(3) **Vector moment or torque of a force.** The vector moment or torque of a force about a point O is a vector quantity related to an axis through O perpendicular to the plane containing O and the line of action of the force $\mathbf{F}$. The magnitude of the vector moment is jointly proportional to the force and the perpendicular distance ON upon the line of the force.




Fig. 1.30

Take a point P on the line of action of the force. Let the position vector of P be $\mathbf{r}$. The moment of the force $\mathbf{F}$ about O is represented by a vector perpendicular to the plane of $\mathbf{r}$ and $\mathbf{F}$. Since $\mathbf{r} \times \mathbf{F}$ is a vector perpendicular to the plane of $\mathbf{r}$ and $\mathbf{F}$, therefore the vector representing the moment of $\mathbf{F}$ about O is $\mathbf{r} \times \mathbf{F}$. Hence if $\mathbf{M}$ be the moment vector, then

$$\mathbf{M} = \mathbf{r} \times \mathbf{F}.$$

(4) **Force on a particle in a magnetic field.** Let $\mathbf{F}$ be the force on a point charge in a magnetic field of intensity $\mathbf{B}$. Then $\mathbf{F}$ is proportional to the component of $\mathbf{B}$ perpendicular to the velocity of the charge. If $\mathbf{v}$ be the velocity of the charge then the vector product $\mathbf{v} \times \mathbf{B}$ represents a vector normal to the plane of $\mathbf{v}$ and $\mathbf{B}$ and hence may be regarded as the component of $\mathbf{B}$ along the perpendicular to the velocity $\mathbf{v}$. Thus if q be the charge on the particle and c the speed of light then the above relation existing between the vectors $\mathbf{F}$, $\mathbf{B}$ and $\mathbf{v}$ is expressed as

$$\mathbf{F} = \frac{q}{c} \mathbf{v} \times \mathbf{B} \text{ in Gaussian units}$$

or

$$\mathbf{F} = q\mathbf{v} \times \mathbf{B} \text{ in MKS units.}$$

(5) **Force on a charged particle.** If a particle of charge q is in an electric field of intensity $\mathbf{E}$ at rest, then force on the charged particle due to electric field $\mathbf{F}_e = q\mathbf{E}$ (in e.s.u.)




Fig. 1.31

But if the charged particle is moving with velocity $\mathbf{v}$ relative to an observer, the magnetic field is produced. The force experienced by a moving charged particle due to magnetic field $\mathbf{F}_m = q\mathbf{v} \times \mathbf{B}$ (in e.m.u.), where $\mathbf{B}$ is the intensity of magnetic field. This is known as *magnetic force*.

Total force on the moving charged particle is the sum of electrostatic and magnetic forces and is given by

$$\mathbf{F} = \mathbf{F}_e + \mathbf{F}_m = q\mathbf{E} + q(\mathbf{v} \times \mathbf{B}),$$

$$\mathbf{F} = q[\mathbf{E} + \mathbf{v} \times \mathbf{B}]. \quad \dots (1)$$

This is known as *Lorentz force*.

If $\mathbf{E}$ is in e.s.u. and $\mathbf{B}$ in gauss, the equation (1) can be written as

$$\mathbf{F} = q \left[\mathbf{E} + \frac{\mathbf{v} \times \mathbf{B}}{c} \right], \quad \dots (2)$$

where c is speed of light in vacuum.

(a) **Circular motion.** Let us consider a particle moving in a circle of radius r with angular velocity ω . Let O be the centre of the circular path and A the starting point. Join O to A and draw OB perpendicular to OA . Let $\mathbf{i}, \mathbf{j}$ be the unit vectors along OA and OB taken as axes of x and y respectively. Let at an instant of time t , P be the position of the particle such that its position vector referred to O as origin is $\mathbf{r}$.




Fig. 1.32

Assuming $\angle AOP = \theta$, we have $\theta = \omega t$. $\dots (1)$

Since the radius r of a circular orbit is constant and the unit vector $\mathbf{i}$ rotates at a constant rate, the equation of the circular orbit can be given as

$$\mathbf{r}(t) = r\hat{\mathbf{r}}(t). \quad \dots (2a)$$

Now draw PM perpendicular to OA . Then

$$\begin{aligned} \vec{OP} &= \vec{OM} + \vec{MP} \\ &= OP \cos \theta \mathbf{i} + OP \sin \theta \mathbf{j}, \end{aligned}$$

$$\mathbf{r}(t) = r\hat{\mathbf{r}}(t) = r \cos \omega t \mathbf{i} + r \sin \omega t \mathbf{j} \text{ as } OP = r. \quad \dots (2b)$$

$$\hat{\mathbf{r}}(t) = \cos \omega t \mathbf{i} + \sin \omega t \mathbf{j}. \quad \dots (3)$$

In particular if $\theta = \frac{\pi}{4}$, then $\hat{\mathbf{r}} = \frac{1}{\sqrt{2}} \mathbf{i} + \frac{1}{\sqrt{2}} \mathbf{j}$

and if $\theta = \frac{\pi}{2}$, then $\hat{\mathbf{r}} = \mathbf{j}$

and if $\theta = 0$, then $\hat{\mathbf{r}} = \mathbf{i}$.

Now the velocity, $\mathbf{v}$ of the particle is given by (differentiation of vectors being defined in §1.17)

$$\mathbf{v} = \frac{d\mathbf{r}}{dt} = r \frac{d\hat{\mathbf{r}}}{dt} = r \left(\mathbf{i} \frac{d}{dt} \cos \omega t + \mathbf{j} \frac{d}{dt} \sin \omega t \right) \text{ from (2)}$$

$$= \omega r (-\sin \omega t \mathbf{i} + \cos \omega t \mathbf{j}),$$

which follows that the magnitude of the velocity

$$|\mathbf{v}| = \omega r \text{ say } v = \omega r.$$

Again the acceleration of the particle is given by

$$\mathbf{a} = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{r}}{dt^2}$$

$$= \omega r \frac{d}{dt} (-\sin \omega t \mathbf{i} + \cos \omega t \mathbf{j})$$

$$= -\omega^2 r (\cos \omega t \mathbf{i} + \sin \omega t \mathbf{j})$$

$$= -\omega^2 \mathbf{r} \text{ from (3).}$$

$\therefore$ magnitude of the acceleration, say a

$$= |\mathbf{a}| = |-\omega^2 \mathbf{r}|,$$

i.e.,

$$a = \omega^2 r.$$

Elimination of ω between (4) and (5) yields

$$a = \frac{v^2}{r}.$$

The acceleration given by the expression (6) is known as **Centripetal** (i.e. centre seeking) acceleration.

Now we know that if f be the frequency of the particle, then

$$2\pi f = \omega,$$

and the time period T of the motion is given by

$$T = \frac{2\pi}{\omega} = \frac{1}{f} \text{ from (7).}$$

(7) **Angular velocity of a rigid body about a fixed axis.** Consider the motion of a rigid body rotating about a fixed axis ON at the rate of ω radians per second. Then the angular velocity of the body




Fig. 1.35

and $\mathbf{r}$ and its magnitude is given by

$$v = \omega r \sin \theta.$$

is specified by the vector $\vec{\omega}$ whose modulus is ω and whose direction is parallel to the axis and in the positive sense relative to rotation.

Let O be a point on the fixed axis; P a point fixed in the body, $\mathbf{r}$ the position vector of P referred to O as origin, and PN perpendicular to the axis of rotation. The part at P is moving in a circular path of radius $PN = r \sin \theta$ about N as centre. Its velocity vector $\mathbf{v}$ is at right angles to the plane of

$$\vec{\mathbf{v}} = \vec{\omega} \times \mathbf{r}.$$

Problem 53. A particle, acted on by constant forces $6\mathbf{i} + \mathbf{j} - 3\mathbf{k}$ and $(1 - \mathbf{j}) - \mathbf{k}$ is displaced from the point $\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ to the point $5\mathbf{i} + 4\mathbf{j} + \mathbf{k}$. Find the total work done by the forces.

Let $\mathbf{F}_1 = 6\mathbf{i} + \mathbf{j} - 3\mathbf{k}$, $\mathbf{F}_2 = \mathbf{i} - \mathbf{j} - \mathbf{k}$

and the displacement $\mathbf{r} = 5\mathbf{i} + 4\mathbf{j} + \mathbf{k} - (\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$

$$= 4\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}.$$

Work done by force $\mathbf{F}_1 = \mathbf{F}_1 \cdot \mathbf{r}$

$$= (4\mathbf{i} + \mathbf{j} - 3\mathbf{k}) \cdot (4\mathbf{i} + 2\mathbf{j} - 2\mathbf{k})$$

$$= 4 \cdot 4 + 1 \cdot 2 - 3 \cdot (-2) = 24 \text{ units.}$$

Work done by the force $\mathbf{F}_2$

$$= \mathbf{F}_2 \cdot \mathbf{r} = (\mathbf{i} - \mathbf{j} - \mathbf{k}) \cdot (4\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}) = 16 \text{ units.}$$

Total work done $= 24 + 16 = 40$ units.

Problem 54. A rigid body is spinning with an angular velocity of 4 radians per second about an axis parallel to $3\mathbf{j} - \mathbf{k}$ passing through the point $\mathbf{i} + 3\mathbf{j} - \mathbf{k}$. Find the velocity of the particle at the point $\mathbf{i} - 3\mathbf{j} + \mathbf{k}$.

Let $\mathbf{r}$ be the position vector of the point relative to the given point on the axis, then

$$\mathbf{r} = 4\mathbf{i} - 2\mathbf{j} + \mathbf{k} - (\mathbf{i} + 3\mathbf{j} - \mathbf{k})$$

$$= 3\mathbf{i} - 5\mathbf{j} + 2\mathbf{k}.$$

Angular velocity of the particle is given by

$$\vec{\omega} = 4 \cdot \frac{(3\mathbf{j} - \mathbf{k})}{|3\mathbf{j} - \mathbf{k}|} = \frac{4}{\sqrt{10}} (3\mathbf{j} - \mathbf{k}).$$

Hence the velocity of the particle

$$\vec{\omega} \times \mathbf{r} = \frac{2}{\sqrt{10}} (3\mathbf{j} - \mathbf{k}) \times (3\mathbf{i} - 5\mathbf{j} + 2\mathbf{k})$$

$$= \frac{4}{\sqrt{10}} (\mathbf{i} - 3\mathbf{j} - 9\mathbf{k}).$$

Its magnitude $= \frac{4}{\sqrt{10}} |\mathbf{i} - 3\mathbf{j} - 9\mathbf{k}| = 4 \sqrt{\frac{91}{10}} = 12$ approx.

Problem 55. Find the torque about the point $10\mathbf{j}$ of a force represented by $-3\mathbf{i} + \mathbf{j} + 5\mathbf{k}$ acting through the point $7\mathbf{i} + 3\mathbf{j} + \mathbf{k}$.

Let $\mathbf{F} = -3\mathbf{i} + \mathbf{j} + 5\mathbf{k}$ and $\mathbf{r}$ be the vector from the given point to the point of application of the force, then

$$\mathbf{r} = 10\mathbf{j} - (7\mathbf{i} + 3\mathbf{j} + \mathbf{k}) = -7\mathbf{i} + 7\mathbf{j} - \mathbf{k}.$$

$$\text{Reqd. Torque} = \mathbf{r} \times \mathbf{F} = (-7\mathbf{i} + 7\mathbf{j} - \mathbf{k}) \times (-3\mathbf{i} + \mathbf{j} + 5\mathbf{k})$$

$$= 36\mathbf{i} - 38\mathbf{j} - 14\mathbf{k}.$$

Problem 56. A particle is moving in a circular orbit of radius 10 cms. Its frequency of motion is 60 cycles per sec., find the time period, velocity and acceleration of the particle.

Given frequency $f=60$ cycle/sec.; radius of the orbit = 10 cms.

Let T be the time period, v the velocity and a the acceleration of the particle. Then

$$T = \frac{1}{f} = \frac{1}{60} = 0.017 \text{ sec. nearly.}$$

Now $\omega = 2\pi f = 2 \times \frac{22}{7} \times 60 = \frac{2640}{7} = 377$ radians/sec. approx.

$\therefore v = \omega r = 377 \times 10 = 3.8 \times 10^3$ sec. nearly

and $a = \omega^2 r = 377 \times 377 \times 10 = 10^6$ cm./sec. nearly.

Problem 57. Calculate the force in dynes acting (1) on a proton, (2) on an electron, in an electric field of intensity 1000 volts per cm. Given charge on electron = 4.8×10^{-10} e.s.u.

The intensity of electric field = 1000 volts/cm.

$$= \frac{1000}{300} \text{ e.s.u. of volts/cm.}$$

$$= \frac{10}{3} \text{ stat. volts/cm.}$$

The force on the proton = qE
 $= 4.8 \times 10^{-10} \times \frac{10}{3} = 1.6 \times 10^{-9}$ dynes.

The force on the electron = $qE = -1.6 \times 10^{-9}$ dynes.

Problem 58. Calculate the force on the proton in dynes in a magnetic field of intensity 100 gauss directed along z-axis:

(i) when the proton moves with velocity 10^8 along x-axis.

(ii) when the proton is at rest.

(i) Given $B = 100$ gauss along z-axis = $100\mathbf{k}$,
 $v = 10^8 \mathbf{i}$.

$\therefore$ the force on the proton due to magnetic field

$$= \frac{q}{c} (\mathbf{v} \times \mathbf{B})$$

$$= \frac{4.8 \times 10^{-10}}{3 \times 10^{10}} [10^8 \mathbf{i} \times (100\mathbf{k})]$$

$$= 1.6 \times 10^{20} [-10^{10} \mathbf{j}]$$

$$= -1.6 \times 10^{10} \mathbf{j} \text{ dynes,}$$

i.e., the force of magnitude 1.6×10^{10} is acting along negative direction of y-axis.

(ii) When the proton is at rest, there is no magnetic field and hence no force.

Problem 59. Using a right-handed system, the electric field $\mathbf{E}$, and velocity $\mathbf{v}$ of a particle of charge q e.s.u. are given by

$$\mathbf{E} = 2\mathbf{i} \text{ e.s.u.}$$

$$\mathbf{B} = (3\mathbf{i} + 4\mathbf{j}) \text{ e.m.u.}$$

$$\mathbf{v} = 9\mathbf{i} + 3\mathbf{j} + 4\mathbf{k} \text{ cm./sec.}$$

Calculate the electrostatic and magnetic force on the charge.

Electrostatic force on the charge $q = qE = q \cdot 2\mathbf{i} = 2q\mathbf{i}$ dynes, q is e.s.u.

Therefore electrostatic force has magnitude $2q$ along positive x-axis.

Magnetic force on charge $q = \frac{q}{c} (\mathbf{v} \times \mathbf{B})$, $\mathbf{B}$ being in e.m.u.

$$= \frac{q}{3 \times 10^{10}} [(2\mathbf{i} + 3\mathbf{j} + 4\mathbf{k}) \times (3\mathbf{i} + 4\mathbf{j})]$$

$$= \frac{q}{3 \times 10^{10}} [8\mathbf{k} - 9\mathbf{k} + 12\mathbf{j} - 16\mathbf{i}]$$

$$= \frac{q}{3 \times 10^{10}} [-16\mathbf{i} + 12\mathbf{j} - \mathbf{k}].$$

$$\text{magnitude of magnetic force} = \frac{q}{3 \times 10^{10}} \sqrt{16^2 + 12^2 + 1^2}$$

$$= \frac{q}{3 \times 10^{10}} [\sqrt{401}]$$

$$\approx 7 \times 10^{-19} q \text{ dynes.}$$

(q is in e.s.u.)

Direction cosines of Lorentz force are

$$\left[\frac{-16}{7 \times 10^{-10} q}, \frac{12}{7 \times 10^{-10} q}, \frac{-1}{7 \times 10^{-10} q} \right].$$

Problem 60. A proton is moving with velocity 10^{10} cm./sec. along x-axis in an electric field of intensity 3×10^4 volts along x-axis and magnetic field of intensity 3000 gauss along y-axis. Calculate the magnitude and direction of total force.

Charge on the proton $q = 4.8 \times 10^{-10}$ e.s.u.

Intensity of electric field $\mathbf{E} = 3 \times 10^4 \mathbf{i}$ volts

$$= \frac{3 \times 10^4}{300} \mathbf{i} \text{ stat-volts}$$

$$\mathbf{E} = 100 \mathbf{i} \text{ stat-volts.}$$

Intensity of magnetic field $\mathbf{B} = 3000 \mathbf{j}$

and velocity of proton $\mathbf{v} = 10^{10} \mathbf{k}$.

$$\text{total force on the proton} = q \left[\mathbf{E} + \frac{\mathbf{v} \times \mathbf{B}}{c} \right]$$

$$= 4.8 \times 10^{-10} \left[100\mathbf{i} + \frac{1}{c} \{10^{10} \mathbf{k} \times 3000 \mathbf{j}\} \right]$$

$$= 4.8 \times 10^{-10} \left[100\mathbf{i} + \frac{1}{3 \times 10^{10}} \{-3 \times 10^{13} \mathbf{i}\} \right]$$

$$= 4.8 \times 10^{-10} [100\mathbf{i} - 1000\mathbf{i}]$$

$$= 4.8 \times 10^{-10} [-900\mathbf{i}]$$

$$= -4.32 \times 10^{-7} \mathbf{i} \text{ dynes.}$$

Thus the total force has magnitude 4.32×10^{-7} dynes along the negative direction of x-axis.

1.17. DIFFERENTIATION OF VECTORS

Vector Function of a Single Scalar Variable. $\mathbf{F}$ is a vector function of a single scalar variable t , if to each t of the range of values of t there corresponds a vector $\mathbf{F}$ and is written as

$$\mathbf{F} = \mathbf{F}(t).$$

The vector $\mathbf{F}$ can also be expressed in components form, such as

$$\mathbf{F} = f_1(t) \mathbf{i} + f_2(t) \mathbf{j} + f_3(t) \mathbf{k}$$

where $f_1(t)$, $f_2(t)$, $f_3(t)$ are components of $\mathbf{F}(t)$ defined for the range of values of t and $\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$ are unit vectors in the directions of the principal axes.

Illustration: If a moving particle undergoes a displacement in a manner such that at any time t , its position is at a point P whose position vector relative to any fixed origin O is $\mathbf{r}$, then the vector $\mathbf{r}$ is the function of scalar variable t , i.e., $\mathbf{r} = \mathbf{F}(t)$.

Derivative of a vector function of a scalar variable. If $\mathbf{F}(t)$ represents a vector function of a scalar variable t , over the interval $a \leq t \leq b$ and if $\lim_{\delta t \rightarrow 0} \frac{\mathbf{F}(t+\delta t) - \mathbf{F}(t)}{\delta t}$ exists, then this limit is called as the derivative or differential coefficient of $\mathbf{F}(t)$ at t . The process of finding out differential coefficient is called *differentiation*.

By convention the method of denoting a derivative is

$$\mathbf{F}'(t) \text{ or } \frac{d}{dt} \mathbf{F}(t).$$




Fig. 1.34

curve $\mathbf{r} = \mathbf{F}(t)$. Assuming that an increment δt in t produces an increment $\delta \mathbf{r}$ in $\mathbf{r}$, we have

$$\mathbf{r} + \delta \mathbf{r} = \mathbf{F}(t + \delta t)$$

It is apparent from Fig. 1.34 that $\overrightarrow{PQ} = \delta \mathbf{r}$

Obviously the quotient $\frac{\delta \mathbf{r}}{\delta t}$ is a vector, since $\mathbf{r}$ is a vector and t is a scalar and it gives the average rate of change of $\mathbf{r}$ with t .

As δt tends to take zero value, the point Q moves up to coincide with P , so that the chord PQ coincides with tangent at P to the curve.

Time-derivative of a vector. Let us suppose that vector $\mathbf{r}$ is a continuous single valued function of a scalar variable t , i.e.

$$\mathbf{r} = \mathbf{F}(t)$$

At an instant of time t , let P be the position of a particle whose position vector referred to a fixed origin O , be $\mathbf{r}$. After an interval of time δt , let Q be the position of the moving particle along the curve.

Thus the vector $\frac{\delta \mathbf{r}}{\delta t}$ as $\delta t \rightarrow 0$ is along the direction of tangent at P in the plane for increasing t . The limiting value of this quotient when exists, is the derivative or differential coefficient of $\mathbf{r}$ with respect to t (time), and denoted by $\frac{d\mathbf{r}}{dt}$ or $\dot{\mathbf{r}}$ i.e.

$$\lim_{\delta t \rightarrow 0} \frac{\delta \mathbf{r}}{\delta t} = \frac{d\mathbf{r}}{dt} = \dot{\mathbf{r}}.$$

But we know that velocity $\mathbf{v}$ of a particle is a vector and this is the instantaneous rate of change of the position of the particle, therefore

$$\mathbf{v} = \frac{d\mathbf{r}}{dt},$$

the first time derivative of position of a particle gives its velocity and its magnitude i.e. $v = |\mathbf{v}|$ is known as the speed of the particle. Clearly the speed is a scalar quantity.

Further the derivative $\frac{d\mathbf{r}}{dt}$ is also in general a function of time t , and may itself possess a derivative, which is called the second derivative of $\mathbf{r}$ and is denoted by $\frac{d^2\mathbf{r}}{dt^2}$.

But the acceleration $\mathbf{a}$ of a particle is a vector and this is the time rate of change of the velocity of the particle, therefore

$$\mathbf{a} = \frac{d^2\mathbf{r}}{dt^2}$$

the second time-derivative of the position of a particle gives its acceleration.

Note 1. The second derivative $\frac{d^2\mathbf{r}}{dt^2}$ is also a vector function of t and it possesses a derivative $\frac{d^3\mathbf{r}}{dt^3}$ known as the third derivative of $\mathbf{r}$. Similarly the existence of fourth, fifth, sixth...derivatives can be established.

Note 2. The derivative of a constant vector $\mathbf{c}$ is zero i.e. if $\mathbf{c}$ is a constant vector, then $\frac{d\mathbf{c}}{dt} = 0$; for, then the increment in δt produces no change in $\mathbf{c}$.

SOME RULES FOR DIFFERENTIATION

Rule 1. The derivative of the sum of two differentiable vector functions, is equal to the sum of their derivatives.

Let $\mathbf{r}_1$ and $\mathbf{r}_2$ be two differentiable vector functions of a scalar variable t . Suppose that

$$\mathbf{r} = \mathbf{r}_1 + \mathbf{r}_2$$

Then if a change δt in t corresponds a change $\delta \mathbf{r}$ in $\mathbf{r}$, $\delta \mathbf{r}_1$ in $\mathbf{r}_1$, $\delta \mathbf{r}_2$ in $\mathbf{r}_2$, we have $\mathbf{r} + \delta \mathbf{r} = (\mathbf{r}_1 + \delta \mathbf{r}_1) + (\mathbf{r}_2 + \delta \mathbf{r}_2)$
i.e. $\delta \mathbf{r} = \delta \mathbf{r}_1 + \delta \mathbf{r}_2$.

Dividing throughout by δt and proceeding to the limit as $\delta t \rightarrow 0$ we find,

$$\frac{d\mathbf{r}}{dt} = \frac{d\mathbf{r}_1}{dt} + \frac{d\mathbf{r}_2}{dt}.$$

This result may be extended to any number of vectors.

In particular if any vector is expressed as the sum of rectangular component vectors such as

$\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$, where $\mathbf{i}, \mathbf{j}, \mathbf{k}$ are unit vectors along the axes; then the derivative of $\mathbf{r}$ is given by

$$\frac{d\mathbf{r}}{dt} = \frac{dx}{dt} \mathbf{i} + \frac{dy}{dt} \mathbf{j} + \frac{dz}{dt} \mathbf{k}$$

Note. If $\mathbf{r}$ is a differentiable function of a scalar variable s and s is a differentiable function of another variable t , then we can state that

$$\frac{d\mathbf{r}}{dt} = \frac{d\mathbf{r}}{ds} \cdot \frac{ds}{dt}.$$

(2) The differentiation of scalar product of two vectors. The derivative of product of two differential vector functions, is equal to the sum of the quantities found by differentiating one of the factors and leaving the other unchanged and then differentiating the other and leaving the first unchanged i.e., if $\mathbf{r} = \mathbf{r}_1 \cdot \mathbf{r}_2$, where $\mathbf{r}_1, \mathbf{r}_2$ are the vector functions of scalar variable t ,

then
$$\frac{d\mathbf{r}}{dt} = \mathbf{r}_1 \cdot \frac{d\mathbf{r}_2}{dt} + \frac{d\mathbf{r}_1}{dt} \cdot \mathbf{r}_2$$

If a change δt in t corresponds the changes $\delta \mathbf{r}$ in $\mathbf{r}$, $\delta \mathbf{r}_1$ in $\mathbf{r}_1$ and $\delta \mathbf{r}_2$ in $\mathbf{r}_2$, then we have,

$$\begin{aligned} \mathbf{r} + \delta \mathbf{r} &= (\mathbf{r}_1 + \delta \mathbf{r}_1) \cdot (\mathbf{r}_2 + \delta \mathbf{r}_2) \\ &= \mathbf{r}_1 \cdot \mathbf{r}_2 + \mathbf{r}_1 \cdot \delta \mathbf{r}_2 + \delta \mathbf{r}_1 \cdot \mathbf{r}_2 + \delta \mathbf{r}_1 \cdot \delta \mathbf{r}_2 \\ &= \mathbf{r}_1 \cdot \mathbf{r}_2 + \mathbf{r}_1 \cdot \delta \mathbf{r}_2 + \delta \mathbf{r}_1 \cdot \mathbf{r}_2, \text{ neglecting the product } \delta \mathbf{r}_1 \cdot \delta \mathbf{r}_2 \text{ as it is vanishingly small} \end{aligned}$$

or $\delta \mathbf{r} = \mathbf{r}_1 \cdot \delta \mathbf{r}_2 + \delta \mathbf{r}_1 \cdot \mathbf{r}_2$ as $\mathbf{r} = \mathbf{r}_1 \cdot \mathbf{r}_2$.

Dividing throughout by δt and proceeding to the limit as $\delta t \rightarrow 0$ we get

$$\frac{d\mathbf{r}}{dt} = \mathbf{r}_1 \cdot \frac{d\mathbf{r}_2}{dt} + \frac{d\mathbf{r}_1}{dt} \cdot \mathbf{r}_2$$

Note 1. If u is a scalar function of t , and $\mathbf{r} = u\mathbf{r}_1$, then

$$\frac{d\mathbf{r}}{dt} = \frac{du}{dt} \mathbf{r}_1 + u \frac{d\mathbf{r}_1}{dt}.$$

Note 2. If $\mathbf{r} = \mathbf{r}_1 \cdot \mathbf{r}_1$, then $\frac{d\mathbf{r}}{dt} = \frac{d\mathbf{r}_1}{dt} \cdot \mathbf{r}_1 + \mathbf{r}_1 \cdot \frac{d\mathbf{r}_1}{dt}$

$$= 2\mathbf{r}_1 \cdot \frac{d\mathbf{r}_1}{dt} \text{ since dot product is commutative}$$

$$= 2\mathbf{r}_1 \cdot \frac{d\mathbf{r}_1}{dt} \text{ as } \mathbf{r}_1 \cdot \mathbf{r}_1 = r_1^2 = r_1^2$$

which follows that $\mathbf{r}_1 \cdot \frac{d\mathbf{r}_1}{dt} = r_1 \cdot \frac{dr_1}{dt}$.

Note 3. The necessary and sufficient condition for a vector $\mathbf{r}$ to have constant magnitude is $\mathbf{r} \cdot \frac{d\mathbf{r}}{dt} = 0$.

$$\text{Since } \mathbf{r} \cdot \mathbf{r} = |\mathbf{r}|^2, \text{ i.e. } 2\mathbf{r} \cdot \frac{d\mathbf{r}}{dt} = 2|\mathbf{r}| \frac{d}{dt} (|\mathbf{r}|).$$

Therefore $\mathbf{r} \cdot \frac{d\mathbf{r}}{dt} = 0$, if and only if $\frac{d}{dt} |\mathbf{r}| = 0$, i.e. if $|\mathbf{r}|$ is constant.

(3) Differentiation of vector product of two vectors. Suppose that

$$\mathbf{r} = \mathbf{r}_1 \times \mathbf{r}_2,$$

where $\mathbf{r}_1$ and $\mathbf{r}_2$ are differentiable vector functions of a scalar variable t .

If an increment δt corresponds to increments $\delta \mathbf{r}$ in $\mathbf{r}$, $\delta \mathbf{r}_1$ in $\mathbf{r}_1$ and $\delta \mathbf{r}_2$ in $\mathbf{r}_2$, then we have $\mathbf{r} + \delta \mathbf{r} = (\mathbf{r}_1 + \delta \mathbf{r}_1) \times (\mathbf{r}_2 + \delta \mathbf{r}_2)$

$$= \mathbf{r}_1 \times \mathbf{r}_2 + \mathbf{r}_1 \times \delta \mathbf{r}_2 + \delta \mathbf{r}_1 \times \mathbf{r}_2 \text{ neglecting the product } \delta \mathbf{r}_1 \times \delta \mathbf{r}_2 \text{ as it is vanishingly small}$$

$$\delta \mathbf{r} = \mathbf{r}_1 \times \delta \mathbf{r}_2 + \delta \mathbf{r}_1 \times \mathbf{r}_2.$$

Dividing throughout by δt and proceeding to the limit $\delta t \rightarrow 0$, we have

$$\frac{d\mathbf{r}}{dt} = \mathbf{r}_1 \times \frac{d\mathbf{r}_2}{dt} + \frac{d\mathbf{r}_1}{dt} \times \mathbf{r}_2.$$

Note. The necessary and sufficient condition for $\mathbf{r}$ to have constant direction is

$$\mathbf{r} \times \frac{d\mathbf{r}}{dt} = 0.$$

Let r be the magnitude of $\mathbf{r}$ and $\hat{\mathbf{r}}$ be the unit vector in the direction of $\mathbf{r}$. Then

$$\mathbf{r} = r\hat{\mathbf{r}}.$$

$$\frac{d\mathbf{r}}{dt} = r \frac{d\hat{\mathbf{r}}}{dt} + \frac{dr}{dt} \hat{\mathbf{r}},$$

$$\mathbf{r} \times \frac{d\mathbf{r}}{dt} = r\hat{\mathbf{r}} \times \left\{ r \frac{d\hat{\mathbf{r}}}{dt} + \frac{dr}{dt} \hat{\mathbf{r}} \right\}$$

$$= r^2 \hat{\mathbf{r}} \times \frac{d\hat{\mathbf{r}}}{dt} \quad \dots (i)$$

the term vanishing as $\hat{\mathbf{r}} \times \hat{\mathbf{r}} = 0$.

Let us now suppose that $\mathbf{r}$ has constant direction, so that $\hat{\mathbf{r}}$ is a constant vector giving,

$$\frac{d\hat{\mathbf{r}}}{dt} = 0.$$

As such we have from (i)

$$\mathbf{r} \times \frac{d\mathbf{r}}{dt} = 0 \text{ i.e., the condition is necessary.}$$

Now to show that the condition is sufficient, suppose that

$$\mathbf{r} \times \frac{d\mathbf{r}}{dt} = 0$$

$$\text{So that from (i) } r^2 \hat{\mathbf{r}} \times \frac{d\hat{\mathbf{r}}}{dt} = 0 \quad \dots$$

But $\hat{\mathbf{r}}$ is a vector having unity as its length therefore by Note 3,

$$\text{Rule 2 of § 1.17 } \hat{\mathbf{r}} \cdot \frac{d\hat{\mathbf{r}}}{dt} = 0. \quad \dots$$

From (ii) and (iii) we see that $\hat{\mathbf{r}} \neq 0$ therefore $\frac{d\hat{\mathbf{r}}}{dt} = 0$.

$\therefore \hat{\mathbf{r}}$ is a constant vector, i.e. the direction of $\mathbf{r}$ remains the same.

(4) The differentiation of a scalar triple product. Let $\mathbf{v}$ be expressed as the scalar triple product of three vectors $\mathbf{p}, \mathbf{q}, \mathbf{r}$ as

$$\begin{aligned} \mathbf{v} &= [\mathbf{pqr}] \\ &= \mathbf{p} \cdot \mathbf{q} \times \mathbf{r} \end{aligned}$$

where $\mathbf{p}, \mathbf{q}, \mathbf{r}$ are the differentiable vector functions of a scalar variable t . If a change δt in t corresponds a change $\delta \mathbf{v}$ in $\mathbf{v}$, $\delta \mathbf{p}$ in $\mathbf{p}$, $\delta \mathbf{q}$ in $\mathbf{q}$ and $\delta \mathbf{r}$ in $\mathbf{r}$, then we have

$$\begin{aligned} \mathbf{v} + \delta \mathbf{v} &= (\mathbf{p} + \delta \mathbf{p}) \cdot \{(\mathbf{q} + \delta \mathbf{q}) \times (\mathbf{r} + \delta \mathbf{r})\} \\ &= (\mathbf{p} + \delta \mathbf{p}) \cdot \{\mathbf{q} \times \mathbf{r} + \mathbf{q} \times \delta \mathbf{r} + \delta \mathbf{q} \times \mathbf{r} + \delta \mathbf{q} \times \delta \mathbf{r}\} \\ &= (\mathbf{p} + \delta \mathbf{p}) \cdot (\mathbf{q} \times \mathbf{r} + \mathbf{q} \times \delta \mathbf{r} + \delta \mathbf{q} \times \mathbf{r}) \\ &\quad \text{(leaving } \delta \mathbf{q} \times \delta \mathbf{r} \text{ as it is vanishingly small)} \\ &= \mathbf{p} \cdot \mathbf{q} \times \mathbf{r} + \delta \mathbf{p} \cdot \mathbf{q} \times \mathbf{r} + \mathbf{p} \cdot \delta \mathbf{q} \times \mathbf{r} + \mathbf{p} \cdot \mathbf{q} \times \delta \mathbf{r} \\ &\quad \text{(Leaving small value products)} \end{aligned}$$

$$\text{or } \delta \mathbf{v} = \delta \mathbf{p} \cdot \mathbf{q} \times \mathbf{r} + \mathbf{p} \cdot \delta \mathbf{q} \times \mathbf{r} + \mathbf{p} \cdot \mathbf{q} \times \delta \mathbf{r} \dots \text{using (i)}$$

Dividing it throughout by δt and proceeding to the limit as $\delta t \rightarrow 0$ we have

$$\lim_{\delta t \rightarrow 0} \frac{\delta \mathbf{v}}{\delta t} = \lim_{\delta t \rightarrow 0} \left\{ \frac{\delta \mathbf{p}}{\delta t} \cdot \mathbf{q} \times \mathbf{r} + \mathbf{p} \cdot \frac{\delta \mathbf{q}}{\delta t} \times \mathbf{r} + \mathbf{p} \cdot \mathbf{q} \times \frac{\delta \mathbf{r}}{\delta t} \right\}$$

$$\begin{aligned} \text{or } \frac{d\mathbf{v}}{dt} &= \left[\frac{d\mathbf{p}}{dt} \cdot \mathbf{q} \times \mathbf{r} + \mathbf{p} \cdot \frac{d\mathbf{q}}{dt} \times \mathbf{r} + \mathbf{p} \cdot \mathbf{q} \times \frac{d\mathbf{r}}{dt} \right] \\ &= \left\{ \frac{d\mathbf{p}}{dt} \cdot \mathbf{qr} \right\} + \left\{ \mathbf{p} \cdot \frac{d\mathbf{q}}{dt} \times \mathbf{r} \right\} + \left\{ \mathbf{pq} \cdot \frac{d\mathbf{r}}{dt} \right\} \end{aligned}$$

(5) The differentiation of vector triple product. Let $\mathbf{v}$ be expressed as the vector triple product of three vectors $\mathbf{p}, \mathbf{q}, \mathbf{r}$, i.e.,

$$\mathbf{v} = \mathbf{p} \times (\mathbf{q} \times \mathbf{r}) \quad \dots (i)$$

$\mathbf{p}, \mathbf{q}, \mathbf{r}$, being differentiable vector functions of scalar variable t . If a change δt in t corresponds a change $\delta \mathbf{p}$ in $\mathbf{p}$, $\delta \mathbf{q}$ in $\mathbf{q}$, and $\delta \mathbf{r}$ in $\mathbf{r}$, then we have

$$\begin{aligned} \mathbf{v} + \delta \mathbf{v} &= (\mathbf{p} + \delta \mathbf{p}) \times \{(\mathbf{q} + \delta \mathbf{q}) \times (\mathbf{r} + \delta \mathbf{r})\} \\ &= (\mathbf{p} + \delta \mathbf{p}) \times \{\mathbf{q} \times \mathbf{r} + \mathbf{q} \times \delta \mathbf{r} + \delta \mathbf{q} \times \mathbf{r} + \delta \mathbf{q} \times \delta \mathbf{r}\} \\ &= (\mathbf{p} + \delta \mathbf{p}) \times \{\mathbf{q} \times \mathbf{r} + \mathbf{q} \times \delta \mathbf{r} + \delta \mathbf{q} \times \mathbf{r}\} \\ &\quad \text{(neglecting product } \delta \mathbf{q} \times \delta \mathbf{r} \text{ as it is vanishingly small)} \\ &= \mathbf{p} \times (\mathbf{q} \times \mathbf{r}) + \mathbf{p} \times (\mathbf{q} \times \delta \mathbf{r}) + \mathbf{p} \times (\delta \mathbf{q} \times \mathbf{r}) + \delta \mathbf{p} \times (\mathbf{q} \times \mathbf{r}) \\ &\quad + \delta \mathbf{p} \times (\mathbf{q} \times \delta \mathbf{r}) + \delta \mathbf{p} \times (\delta \mathbf{q} \times \mathbf{r}) \\ &= \mathbf{p} \times (\mathbf{q} \times \mathbf{r}) + \mathbf{p} \times (\mathbf{q} \times \delta \mathbf{r}) + \mathbf{p} \times (\delta \mathbf{q} \times \mathbf{r}) + \delta \mathbf{p} \times (\mathbf{q} \times \mathbf{r}) \\ &\quad \text{(neglecting small value products)} \end{aligned}$$

$$\text{or } \delta \mathbf{v} = \mathbf{p} \times (\mathbf{q} \times \delta \mathbf{r}) + \mathbf{p} \times (\delta \mathbf{q} \times \mathbf{r}) + \delta \mathbf{p} \times (\mathbf{q} \times \mathbf{r}) \quad \text{using (i)}$$

Dividing it throughout by δt and proceeding to the limit as $\delta t \rightarrow 0$, we have

$$\begin{aligned} \lim_{\delta t \rightarrow 0} \frac{\delta \mathbf{v}}{\delta t} &= \lim_{\delta t \rightarrow 0} \left\{ \mathbf{p} \times \left(\mathbf{q} \times \frac{\delta \mathbf{r}}{\delta t} \right) \right. \\ &\quad \left. + \mathbf{p} \times \left(\frac{\delta \mathbf{q}}{\delta t} \times \mathbf{r} \right) + \frac{\delta \mathbf{p}}{\delta t} \times (\mathbf{q} \times \mathbf{r}) \right\} \end{aligned}$$

$$\text{or } \frac{d\mathbf{v}}{dt} = \left\{ \mathbf{p} \times \left(\mathbf{q} \times \frac{d\mathbf{r}}{dt} \right) + \mathbf{p} \times \left(\frac{d\mathbf{q}}{dt} \times \mathbf{r} \right) + \frac{d\mathbf{p}}{dt} \times (\mathbf{q} \times \mathbf{r}) \right\}$$

Problem 61. If $\mathbf{F}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}$, where x, y, z are differentiable functions of a scalar variable t , prove that

$$\frac{d\mathbf{F}}{dt} = \frac{dx}{dt} \mathbf{i} + \frac{dy}{dt} \mathbf{j} + \frac{dz}{dt} \mathbf{k}.$$

By the definition of differential coefficient

$$\begin{aligned} \frac{d\mathbf{F}}{dt} &= \lim_{\delta t \rightarrow 0} \frac{\mathbf{F}(t + \delta t) - \mathbf{F}(t)}{\delta t} \\ &= \lim_{\delta t \rightarrow 0} \left\{ \frac{[x(t + \delta t)\mathbf{i} + y(t + \delta t)\mathbf{j} + z(t + \delta t)\mathbf{k}] - [x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}]}{\delta t} \right\} \\ &= \lim_{\delta t \rightarrow 0} \left[\frac{[x(t + \delta t) - x(t)]}{\delta t} \mathbf{i} + \frac{[y(t + \delta t) - y(t)]}{\delta t} \mathbf{j} \right. \\ &\quad \left. + \frac{[z(t + \delta t) - z(t)]}{\delta t} \mathbf{k} \right] \\ &= \lim_{\delta t \rightarrow 0} \left(\frac{\delta x}{\delta t} \mathbf{i} + \frac{\delta y}{\delta t} \mathbf{j} + \frac{\delta z}{\delta t} \mathbf{k} \right) \\ &= \frac{dx}{dt} \mathbf{i} + \frac{dy}{dt} \mathbf{j} + \frac{dz}{dt} \mathbf{k} \end{aligned}$$

Problem 62. If $\mathbf{r} = t^2\mathbf{i} - t^2\mathbf{j} + (2t+1)\mathbf{k}$. Find the value of

$$\left| \frac{d\mathbf{r}}{dt} \right|, \left| \frac{d^2\mathbf{r}}{dt^2} \right|, \left| \frac{d\mathbf{r}}{dt} \right|, \left| \frac{d^2\mathbf{r}}{dt^2} \right| \text{ at } t=0.$$

(Agra, M.Sc., 1966)

Given vector $\mathbf{r}$ is the function of scalar variable t , expressed in the form

$$\mathbf{r} = t^2\mathbf{i} - t^2\mathbf{j} + (2t+1)\mathbf{k}. \quad \dots (1)$$

$$\therefore \frac{d\mathbf{r}}{dt} = 2t\mathbf{i} - 2t\mathbf{j} + 2\mathbf{k} \quad \dots (2)$$

($\because \mathbf{i}, \mathbf{j}, \mathbf{k}$ are constant vectors)

Therefore at $t=0$

$$\frac{d\mathbf{r}}{dt} = -2\mathbf{j} + 2\mathbf{k}.$$

so that $\left| \frac{d\mathbf{r}}{dt} \right| = \sqrt{(-1)^2 + (2)^2}$
 $= \sqrt{5}.$

Again differentiating (2), with respect to t , we get

$$\frac{d^2\mathbf{r}}{dt^2} = 2\mathbf{i}$$

at $t=0, \frac{d^2\mathbf{r}}{dt^2} = 2\mathbf{i}.$

$$\therefore \left| \frac{d^2\mathbf{r}}{dt^2} \right| = |2\mathbf{i}| = 2.$$

Problem 63. If $\mathbf{r} = \mathbf{a} \cos \omega t + \mathbf{b} \sin \omega t$, show that

$$\mathbf{r} \times \frac{d\mathbf{r}}{dt} = \omega \mathbf{a} \times \mathbf{b}$$

and $\frac{d^2\mathbf{r}}{dt^2} = -\omega^2 \mathbf{r}$ $\mathbf{a}, \mathbf{b}$ being constant vectors and ω is also a constant.

Given $\mathbf{r} = \mathbf{a} \cos \omega t + \mathbf{b} \sin \omega t \quad \dots (1)$

$$\therefore \frac{d\mathbf{r}}{dt} = \frac{d}{dt} (\cos \omega t) \mathbf{a} + \frac{d}{dt} (\sin \omega t) \mathbf{b}$$

$$= -\omega \sin \omega t \mathbf{a} + \omega \cos \omega t \mathbf{b}. \quad \dots (2)$$

Differentiating (2) with respect to t , we get

$$\frac{d^2\mathbf{r}}{dt^2} = -\omega \frac{d}{dt} (\sin \omega t) \mathbf{a} + \omega \frac{d}{dt} (\cos \omega t) \mathbf{b}$$

$$= -\omega^2 \cos \omega t \mathbf{a} - \omega^2 \sin \omega t \mathbf{b}$$

$$= -\omega^2 [\cos \omega t \mathbf{a} + \sin \omega t \mathbf{b}]$$

$$= -\omega^2 \mathbf{r} \quad \text{by (1)}$$

i.e. $\frac{d^2\mathbf{r}}{dt^2} = -\omega^2 \mathbf{r}.$

Again, we have

$$\mathbf{r} \times \frac{d\mathbf{r}}{dt} = (\mathbf{a} \cos \omega t + \mathbf{b} \sin \omega t) \times (-\omega \sin \omega t \mathbf{a} + \omega \cos \omega t \mathbf{b})$$

$$= \omega [\mathbf{a} \cos \omega t \times -\sin \omega t \mathbf{a} + \mathbf{a} \times \mathbf{b} \cos^2 \omega t$$

$$- \mathbf{b} \times \mathbf{a} \sin^2 \omega t + \mathbf{b} \times \mathbf{b} \sin \omega t \cos \omega t]$$

$$= (\mathbf{a} \times \mathbf{b}) \{ \cos^2 \omega t + \sin^2 \omega t \} \omega$$

As $\mathbf{a} \times \mathbf{a} = \mathbf{b} \times \mathbf{b} = 0$ and $-\mathbf{b} \times \mathbf{a} = \mathbf{a} \times \mathbf{b}$

$$= (\mathbf{a} \times \mathbf{b}) \omega,$$

i.e. $\mathbf{r} \times \frac{d\mathbf{r}}{dt} = \omega (\mathbf{a} \times \mathbf{b})$

Problem 64. If $\mathbf{r} = \mathbf{a}e^{\omega t} + \mathbf{b}e^{-\omega t}$, show that

$$\frac{d^2\mathbf{r}}{dt^2} - \omega^2 \mathbf{r} = 0; \mathbf{a}, \mathbf{b} \text{ are constant vectors and } \omega \text{ being a}$$

constant.

Given

$$\mathbf{r} = \mathbf{a}e^{\omega t} + \mathbf{b}e^{-\omega t} \quad \dots (1)$$

$$\frac{d\mathbf{r}}{dt} = \omega \mathbf{a}e^{\omega t} - \omega \mathbf{b}e^{-\omega t}$$

Differentiating it w. r. t. t , we get

$$\frac{d^2\mathbf{r}}{dt^2} = \omega^2 \mathbf{a}e^{\omega t} + \omega^2 \mathbf{b}e^{-\omega t}$$

$$= \omega^2 (\mathbf{a}e^{\omega t} + \mathbf{b}e^{-\omega t})$$

$$= \omega^2 \mathbf{r} \text{ from (1)}$$

$$\text{or } \frac{d^2\mathbf{r}}{dt^2} - \omega^2 \mathbf{r} = 0.$$

Problem 65. A particle moves along a curve whose parametric equations are $x = e^{-t}$, $y = 2 \cos 3t$, $z = 2 \sin 3t$, where t is the time.

(a) Determine its velocity and acceleration at any time.

(b) Find the magnitudes of velocity and acceleration at $t=0$.

(a) Since a vector $\mathbf{r}$ can be expressed in terms of rectangular components as $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$,

therefore, the position vector of the moving particle at any time t is

$$\mathbf{r} = e^{-t} \mathbf{i} + 2 \cos 3t \mathbf{j} + 2 \sin 3t \mathbf{k}.$$

Thus, the velocity $\mathbf{v} = \frac{d\mathbf{r}}{dt} = -e^{-t} \mathbf{i} - 6 \sin 3t \mathbf{j} + 6 \cos 3t \mathbf{k}$

and the acceleration is $\frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{r}}{dt^2} = e^{-t} \mathbf{i} - 18 \cos 3t \mathbf{j} - 18 \sin 3t \mathbf{k}.$

(b) At $t=0$, $\frac{d\mathbf{r}}{dt} = -\mathbf{i} + 6\mathbf{k}$ and $\frac{d^2\mathbf{r}}{dt^2} = \mathbf{i} - 18\mathbf{j}.$

∴ magnitude of velocity at $t=0$ is

$$\left| \frac{d\mathbf{r}}{dt} \right| = \sqrt{(-1)^2 + (6)^2} = \sqrt{37}.$$

and magnitude of acceleration at $t=0$ is

$$\left| \frac{d^2\mathbf{r}}{dt^2} \right| = \sqrt{(1)^2 + (-18)^2} = \sqrt{325}.$$

Problem 66. A particle moves along the curve $x=2t^2$, $y=t^2-4t$, $z=3t-5$, where t is the time. Find component of its velocity and acceleration at $t=1$ in the direction $\mathbf{i}-3\mathbf{j}+2\mathbf{k}$.

The position vector of the moving particle at any time t is

$$\mathbf{r} = 2t^2\mathbf{i} + (t^2 - 4t)\mathbf{j} + (3t - 5)\mathbf{k}.$$

$$\therefore \text{velocity} = \frac{d\mathbf{r}}{dt} = 4t\mathbf{i} + (2t - 4)\mathbf{j} + 3\mathbf{k},$$

$$\text{so that } \left[\frac{d\mathbf{r}}{dt} \right]_{\text{at } t=1} = 4\mathbf{i} - 2\mathbf{j} + 3\mathbf{k}$$

$$\text{and acceleration} = \frac{d^2\mathbf{r}}{dt^2} = 4\mathbf{i} + 2\mathbf{j},$$

$$\text{so that } \left[\frac{d^2\mathbf{r}}{dt^2} \right]_{\text{at } t=1} = 4\mathbf{i} + 2\mathbf{j}.$$

Now unit vector along $\mathbf{i}-3\mathbf{j}+2\mathbf{k}$

$$= \frac{\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}}{\sqrt{\{1\}^2 + \{-3\}^2 + \{2\}^2}} = \frac{\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}}{\sqrt{14}}.$$

Hence, component of velocity in the direction $\mathbf{i}-3\mathbf{j}+2\mathbf{k}$ is

$$\frac{(\mathbf{i} - 3\mathbf{j} + 2\mathbf{k})}{\sqrt{14}} \cdot (4\mathbf{i} - 2\mathbf{j} + 3\mathbf{k}) = \frac{16}{\sqrt{14}} = \frac{8}{\sqrt{7}} \sqrt{14}$$

and component of acceleration in that direction

$$= \left(\frac{\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}}{\sqrt{14}} \right) \cdot (4\mathbf{i} + 2\mathbf{j}) \\ = -\frac{2}{\sqrt{14}} = -\frac{1}{\sqrt{7}} \sqrt{14}.$$

Problem 67. Prove the following relations:

$$(a) \mathbf{v} = \mathbf{u} + \mathbf{f}t.$$

$$(b) \mathbf{s} = \mathbf{s}_0 + \mathbf{u}t + \frac{1}{2}\mathbf{f}t^2.$$

$$(c) v^2 = u^2 + 2\mathbf{f} \cdot \mathbf{s}.$$

(a) Velocity is defined as rate of change of displacement,

$$\text{i.e. } \mathbf{v} = \frac{d\mathbf{s}}{dt}. \quad \dots (1)$$

Acceleration is defined as rate of change of velocity, i.e.

$$\mathbf{f} = \frac{d\mathbf{v}}{dt} = \frac{d}{dt} \left(\frac{d\mathbf{s}}{dt} \right) = \frac{d^2\mathbf{s}}{dt^2} \quad \dots (2)$$

$$d\mathbf{v} = \mathbf{f} dt. \quad \dots (3)$$

Let initially at $t=0$, the velocity be $\mathbf{u}$ and after time t , let it be $\mathbf{v}$. Integrating (3) we get

$$\int_{\mathbf{u}}^{\mathbf{v}} d\mathbf{v} = \int_0^t \mathbf{f} dt$$

$$\mathbf{v} - \mathbf{u} = \mathbf{f}t.$$

$$\mathbf{v} = \mathbf{u} + \mathbf{f}t$$

$$(b) \text{ We have from (a), } \frac{d\mathbf{s}}{dt} = \mathbf{u} + \mathbf{f}t.$$

Integrating w.r.t. to t ,

$$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{f}t^2 + \mathbf{B},$$

where $\mathbf{B}$ is constant of integration.

If at $t=0$, $\mathbf{s} = \mathbf{s}_0$, then $\mathbf{B} = \mathbf{s}_0$.

$$\mathbf{s} = \mathbf{s}_0 + \mathbf{u}t + \frac{1}{2}\mathbf{f}t^2.$$

(c) Multiplying scalarly equation (2) by $2 \frac{d\mathbf{s}}{dt}$, we get

$$2 \frac{d\mathbf{s}}{dt} \cdot \frac{d^2\mathbf{s}}{dt^2} = 2\mathbf{f} \cdot \frac{d\mathbf{s}}{dt}.$$

Integrating with respect to t , we get

$$2 \int_0^t \mathbf{v} \cdot \frac{d\mathbf{v}}{dt} dt = \int_0^t 2\mathbf{f} \cdot \frac{d\mathbf{s}}{dt} dt.$$

$$2 \int_{\mathbf{u}}^{\mathbf{v}} \mathbf{v} \cdot d\mathbf{v} = 2 \int_0^s \mathbf{f} \cdot d\mathbf{s}.$$

$$\left[\mathbf{v} \cdot \mathbf{v} \right]_{\mathbf{u}}^{\mathbf{v}} = 2\mathbf{f} \cdot \mathbf{s}.$$

$$v^2 - u^2 = 2\mathbf{f} \cdot \mathbf{s}.$$

$$v^2 = u^2 + 2\mathbf{f} \cdot \mathbf{s}.$$

Problem 68. A particle of 1 kg. moving with initial velocity $(\mathbf{i} + 2\mathbf{k})$ metres/sec. is acted upon by a constant force $(\mathbf{i} + 2\mathbf{j} - 2\mathbf{k})$ newtons. Calculate the distance and velocity after 5 seconds. Also find the time in which the particle reaches the xy -plane.

Mass of the particle = 1 kg.

Force acting on the particle = $(\mathbf{i} + 2\mathbf{j} - 2\mathbf{k})$ newtons.

$$\text{Acceleration } \mathbf{f} = \frac{\text{force}}{\text{mass}} = \frac{\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}}{1} = (\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}) \text{ metres/sec.}$$

Initial velocity $\mathbf{u} = (\mathbf{i} + 2\mathbf{k})$,

$$t = 5 \text{ seconds.}$$

Using $\mathbf{v} = \mathbf{u} + \mathbf{f}t$ we have

$$\begin{aligned}\text{Velocity after 5 seconds} &= (\mathbf{i} + 2\mathbf{k}) + (\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}) \cdot 5 \\ &= \mathbf{i} + 2\mathbf{k} + 5\mathbf{i} + 10\mathbf{j} - 10\mathbf{k} \\ &= 6\mathbf{i} + 10\mathbf{j} - 8\mathbf{k}.\end{aligned}$$

$$\begin{aligned}\text{So that } v &= \sqrt{(6^2 + 10^2 + 8^2)} = \sqrt{200} \\ &= 10\sqrt{2} \text{ metres/sec.}\end{aligned}$$

Distance covered in 5 seconds,

$$\begin{aligned}s &= \mathbf{u}t + \frac{1}{2}\mathbf{f}t^2 = (\mathbf{i} + 2\mathbf{k}) \cdot 5 + \frac{1}{2}(\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}) \cdot 25 \\ &= 5\mathbf{i} + 10\mathbf{k} + \frac{25}{2}\mathbf{i} + 25\mathbf{j} - 25\mathbf{k} \\ &= \frac{35}{2}\mathbf{i} + 25\mathbf{j} - 15\mathbf{k} \\ &= 17.5\mathbf{i} + 25\mathbf{j} - 15\mathbf{k}\end{aligned}$$

$\therefore$

$$\begin{aligned}s &= \sqrt{\{(17.5)^2 + (25)^2 + (15)^2\}} \\ &= \sqrt{(306.25 + 625 + 225)} \\ &= \sqrt{(1156.25)} \\ &= 34 \text{ metres.}\end{aligned}$$

Let t be the time when the particle reaches the xy -plane, i.e., when s is equal to zero.

Using equation $s = \mathbf{u}t + \frac{1}{2}\mathbf{f}t^2$, we have

$$\mathbf{i}s_x + \mathbf{j}s_y + \mathbf{k}s_z = (\mathbf{i} + 2\mathbf{k})t + \frac{1}{2}(\mathbf{i} + 2\mathbf{j} - 2\mathbf{k})t^2$$

Comparing coefficients of $\mathbf{k}$,

$$\begin{aligned}s_z &= 2t - t^2 = 0 \\ t &= 2 \text{ sec.}\end{aligned}$$

or

Problem 69. If $\mathbf{r} = t^2\mathbf{i} - t\mathbf{j} + (2t + 1)\mathbf{k}$ and $\mathbf{s} = (2t - 3)\mathbf{i} + \mathbf{j} - t\mathbf{k}$, find

$$(a) \quad \frac{d}{dt}(\mathbf{r} \cdot \mathbf{s}), \quad (b) \quad \frac{d}{dt}(\mathbf{r} \times \mathbf{s}),$$

$$(c) \quad \frac{d}{dt}(\mathbf{r} + \mathbf{s}), \quad (d) \quad \frac{d}{dt}\left(\mathbf{r} \times \frac{d\mathbf{s}}{dt}\right)_{\text{at } t=1}$$

$$\begin{aligned}(a) \quad \frac{d}{dt}(\mathbf{r} \cdot \mathbf{s}) &= \mathbf{r} \cdot \frac{d\mathbf{s}}{dt} + \frac{d\mathbf{r}}{dt} \cdot \mathbf{s} \\ &= \{t^2\mathbf{i} - t\mathbf{j} + (2t + 1)\mathbf{k}\} \cdot \{2\mathbf{i} - \mathbf{k}\} \\ &\quad + \{2t\mathbf{i} - \mathbf{j} + 2\mathbf{k}\} \cdot \{(2t - 3)\mathbf{i} + \mathbf{j} - t\mathbf{k}\} \\ &= 2t^2 - 2t - 1 + 4t^2 - 6t - 1 - 3t \\ &= 6t^2 - 11t - 1 \\ &= -6, \text{ when } t = 1,\end{aligned}$$

$$\text{i.e. } \frac{d}{dt}(\mathbf{r} \cdot \mathbf{s}) = -6 \text{ at } t = 1.$$

$$\begin{aligned}(b) \quad \frac{d}{dt}(\mathbf{r} \times \mathbf{s}) &= \mathbf{r} \times \frac{d\mathbf{s}}{dt} + \frac{d\mathbf{r}}{dt} \times \mathbf{s} \\ &= \{t^2\mathbf{i} - t\mathbf{j} + (2t + 1)\mathbf{k}\} \times \{2\mathbf{i} - \mathbf{k}\} \\ &\quad + \{2t\mathbf{i} - \mathbf{j} + 2\mathbf{k}\} \times \{(2t - 3)\mathbf{i} + \mathbf{j} - t\mathbf{k}\}\end{aligned}$$

$$\begin{aligned}&= 4t^2\mathbf{j} + 2t\mathbf{k} + t\mathbf{i} + 2(2t + 1)\mathbf{j} + 2t\mathbf{k} + 2t^2\mathbf{j} \\ &\quad + (2t - 3)\mathbf{k} + t\mathbf{i} + 2(2t - 3)\mathbf{j} - 2t\mathbf{k} \\ &= (2t - 2)\mathbf{i} + (3t^2 + 8t - 4)\mathbf{j} + (6t - 3)\mathbf{k} \\ &= 0\mathbf{i} + 7\mathbf{j} + 3\mathbf{k} \text{ at } t = 1.\end{aligned}$$

$$\text{Therefore } \frac{d}{dt}(\mathbf{r} \times \mathbf{s})_{\text{at } t=1} = 7\mathbf{j} + 3\mathbf{k}.$$

$$\begin{aligned}(c) \quad \frac{d}{dt}(\mathbf{r} + \mathbf{s}) &= \frac{d\mathbf{r}}{dt} + \frac{d\mathbf{s}}{dt} \\ &= \frac{d}{dt}[t^2\mathbf{i} - t\mathbf{j} + (2t + 1)\mathbf{k}] + \frac{d}{dt}[(2t - 3)\mathbf{i} + \mathbf{j} - t\mathbf{k}] \\ &= (2t\mathbf{i} - \mathbf{j} + 2\mathbf{k}) + (2\mathbf{i} + 0 - \mathbf{k}) \\ &= (2t + 2)\mathbf{i} - \mathbf{j} + \mathbf{k} \\ &= 4\mathbf{i} - \mathbf{j} + \mathbf{k} \text{ at } t = 1.\end{aligned}$$

$$\text{Thus } \frac{d}{dt}(\mathbf{r} + \mathbf{s}) = 4\mathbf{i} - \mathbf{j} + \mathbf{k} \text{ at } t = 1.$$

$$\begin{aligned}(d) \quad \mathbf{s} &= (2t - 3)\mathbf{i} + \mathbf{j} - t\mathbf{k} \\ \frac{d\mathbf{s}}{dt} &= 2\mathbf{i} - \mathbf{k}.\end{aligned}$$

$$\begin{aligned}\left(\mathbf{r} \times \frac{d\mathbf{s}}{dt}\right) &= \{t^2\mathbf{i} - t\mathbf{j} + (2t + 1)\mathbf{k}\} \times \{2\mathbf{i} - \mathbf{k}\} \\ &= t^2\mathbf{j} + 2t\mathbf{k} + t\mathbf{i} + (2t + 1)2\mathbf{j} \\ &= t\mathbf{i} + (t^2 + 4t - 2)\mathbf{j} + 2t\mathbf{k}.\end{aligned}$$

$$\begin{aligned}\frac{d}{dt}\left(\mathbf{r} \times \frac{d\mathbf{s}}{dt}\right) &= \frac{d}{dt}\{t\mathbf{i} + (t^2 + 4t - 2)\mathbf{j} + 2t\mathbf{k}\} \\ &= \mathbf{i} + (2t + 4)\mathbf{j} + 2\mathbf{k} \\ &= (\mathbf{i} + 6\mathbf{j} + 2\mathbf{k}) \text{ at } t = 1.\end{aligned}$$

$$\text{Thus } \frac{d}{dt}\left(\mathbf{r} \times \frac{d\mathbf{s}}{dt}\right) = \mathbf{i} + 6\mathbf{j} + 2\mathbf{k}.$$

Problem 70. Differentiate $\frac{\mathbf{r} \times \mathbf{a}}{\mathbf{r} \cdot \mathbf{a}}$, $\mathbf{a}$ being a constant vector.

Let

$$\mathbf{v} = \frac{\mathbf{r} \times \mathbf{a}}{\mathbf{r} \cdot \mathbf{a}}$$

$$\begin{aligned}\frac{d\mathbf{v}}{dt} &= \frac{d}{dt}\left(\frac{\mathbf{r} \times \mathbf{a}}{\mathbf{r} \cdot \mathbf{a}}\right) \\ &= \frac{(\mathbf{r} \cdot \mathbf{a})(\dot{\mathbf{r}} \times \mathbf{a}) - (\mathbf{r} \times \mathbf{a})(\dot{\mathbf{r}} \cdot \mathbf{a})}{[\mathbf{r} \cdot \mathbf{a}]^2} \\ &= \frac{(\dot{\mathbf{r}} \times \mathbf{a})}{(\mathbf{r} \cdot \mathbf{a})} - \frac{(\mathbf{r} \times \mathbf{a})(\dot{\mathbf{r}} \cdot \mathbf{a})}{[\mathbf{r} \cdot \mathbf{a}]^2} \text{ as } \frac{d\mathbf{a}}{dt} = 0.\end{aligned}$$

Problem 71. Find the derivative of the product $\mathbf{r} \times \mathbf{s}$ and deduce

$$\frac{d}{dt} \left(\mathbf{r} \times \frac{d\mathbf{r}}{dt} \right) = \mathbf{r} \times \frac{d^2\mathbf{r}}{dt^2}. \quad [\text{Agra, 1963}]$$

For the derivative of $\mathbf{r} \times \mathbf{s}$ see Rule (2) of §1.17

$$\text{i.e.} \quad \frac{d}{dt} (\mathbf{r} \times \mathbf{s}) = \frac{d\mathbf{r}}{dt} \times \mathbf{s} + \mathbf{r} \times \frac{d\mathbf{s}}{dt}. \quad \dots(1)$$

For second part putting $\frac{d\mathbf{r}}{dt}$ in place of $\mathbf{s}$ in (1), we get

$$\begin{aligned} \frac{d}{dt} \left(\mathbf{r} \times \frac{d\mathbf{r}}{dt} \right) &= \frac{d\mathbf{r}}{dt} \times \frac{d\mathbf{r}}{dt} + \mathbf{r} \times \frac{d}{dt} \left(\frac{d\mathbf{r}}{dt} \right) \\ &= \mathbf{r} \times \frac{d^2\mathbf{r}}{dt^2} \text{ as } \frac{d\mathbf{r}}{dt} \times \frac{d\mathbf{r}}{dt} = 0. \end{aligned}$$

1.19. PARTIAL DIFFERENTIATION OF VECTORS

Let $\mathbf{F}$ be a vector function depending upon more than one scalar variables, say x, y, z ; then we write $\mathbf{F} = \mathbf{F}(x, y, z)$ and the partial derivative of $\mathbf{F}$ with respect to x is defined as

$$\frac{\partial \mathbf{F}}{\partial x} = \lim_{\delta x \rightarrow 0} \frac{\mathbf{F}(x + \delta x, y, z) - \mathbf{F}(x, y, z)}{\delta x}$$

if this limit exists.

Similarly partial derivatives of $\mathbf{F}$ with respect to y and z can be written as

$$\frac{\partial \mathbf{F}}{\partial y} = \lim_{\delta y \rightarrow 0} \frac{\mathbf{F}(x, y + \delta y, z) - \mathbf{F}(x, y, z)}{\delta y}$$

$$\frac{\partial \mathbf{F}}{\partial z} = \lim_{\delta z \rightarrow 0} \frac{\mathbf{F}(x, y, z + \delta z) - \mathbf{F}(x, y, z)}{\delta z}$$

provided these limits exist.

Note. If $\mathbf{F} = u(x, y, z, t) \mathbf{i} + v(x, y, z, t) \mathbf{j} + w(x, y, z, t) \mathbf{k}$, then the partial derivatives of $\mathbf{F}$ with respect to x, y, z, t respectively, may be expressed as

$$\frac{\partial \mathbf{F}}{\partial x} = \frac{\partial u}{\partial x} \mathbf{i} + \frac{\partial v}{\partial x} \mathbf{j} + \frac{\partial w}{\partial x} \mathbf{k}$$

$$\frac{\partial \mathbf{F}}{\partial y} = \frac{\partial u}{\partial y} \mathbf{i} + \frac{\partial v}{\partial y} \mathbf{j} + \frac{\partial w}{\partial y} \mathbf{k}$$

$$\frac{\partial \mathbf{F}}{\partial z} = \frac{\partial u}{\partial z} \mathbf{i} + \frac{\partial v}{\partial z} \mathbf{j} + \frac{\partial w}{\partial z} \mathbf{k}$$

$$\frac{\partial \mathbf{F}}{\partial t} = \frac{\partial u}{\partial t} \mathbf{i} + \frac{\partial v}{\partial t} \mathbf{j} + \frac{\partial w}{\partial t} \mathbf{k}.$$

Higher order partial derivatives. The partial derivatives $\frac{\partial \mathbf{F}}{\partial x}, \frac{\partial \mathbf{F}}{\partial y}, \frac{\partial \mathbf{F}}{\partial z}$... being themselves functions of the same set of scalar variables may again be partially differentiated, giving second order partial derivatives such as

$$\frac{\partial}{\partial x} \left(\frac{\partial \mathbf{F}}{\partial x} \right), \frac{\partial}{\partial y} \left(\frac{\partial \mathbf{F}}{\partial y} \right), \frac{\partial}{\partial z} \left(\frac{\partial \mathbf{F}}{\partial z} \right), \frac{\partial}{\partial x} \left(\frac{\partial \mathbf{F}}{\partial y} \right), \frac{\partial}{\partial y} \left(\frac{\partial \mathbf{F}}{\partial x} \right) \dots \text{etc.}$$

$$\text{which are denoted by } \frac{\partial^2 \mathbf{F}}{\partial x^2}, \frac{\partial^2 \mathbf{F}}{\partial y^2}, \frac{\partial^2 \mathbf{F}}{\partial z^2}, \frac{\partial^2 \mathbf{F}}{\partial x \partial y}, \frac{\partial^2 \mathbf{F}}{\partial y \partial x} \dots \text{etc.}$$

Further differentiation of second order partial derivatives may give third and higher order partial derivatives.

Note 1. The two second order partial derivatives, viz.,

$$\frac{\partial^2 \mathbf{F}}{\partial x \partial y} \text{ and } \frac{\partial^2 \mathbf{F}}{\partial y \partial x}$$

are equal, if each of them is a continuous function, i.e.,

$$\frac{\partial^2 \mathbf{F}}{\partial x \partial y} = \frac{\partial^2 \mathbf{F}}{\partial y \partial x}.$$

Note 2. If $\mathbf{r} = \mathbf{F}(x, y)$ and $x = f_1(t, s), y = f_2(t, s)$; then we have

$$\frac{\partial \mathbf{r}}{\partial t} = \frac{\partial \mathbf{F}}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial \mathbf{F}}{\partial y} \frac{\partial y}{\partial t}$$

$$\text{and} \quad \frac{\partial \mathbf{r}}{\partial s} = \frac{\partial \mathbf{F}}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial \mathbf{F}}{\partial y} \frac{\partial y}{\partial s}.$$

These two results are similar to those of the results in ordinary calculus.

Total differentials. If $\mathbf{F}$ is a vector function of scalar variables $x, y, z, \dots$, and we assume that the values of the variables increase from $x, y, z, \dots$ to $x + \delta x, y + \delta y, z + \delta z, \dots$, when the corresponding change in $\mathbf{F}$ is $\delta \mathbf{F}$, then we write

$$\mathbf{F}(x, y, z, \dots) + \delta \mathbf{F} = \mathbf{F}(x + \delta x, y + \delta y, z + \delta z, \dots)$$

$$\delta \mathbf{F} = \mathbf{F}(x + \delta x, y + \delta y, z + \delta z, \dots) - \mathbf{F}(x, y, z, \dots)$$

which may be expressed in the form,

$$\begin{aligned} \delta \mathbf{F} &= \frac{\mathbf{F}(x + \delta x, y + \delta y, z + \delta z, \dots) - \mathbf{F}(x, y + \delta y, z + \delta z, \dots)}{\delta x} \delta x \\ &+ \frac{\mathbf{F}(x, y + \delta y, z + \delta z, \dots) - \mathbf{F}(x, y, z + \delta z, \dots)}{\delta y} \delta y + \dots + \dots \end{aligned}$$

Now if $\delta x, \delta y, \delta z, \dots$ tend to zero then $\delta \mathbf{F}$ will also tend to zero; so that the coefficients of $\delta x, \delta y, \delta z, \dots$ in the above expression tend to the limiting values

$$\frac{\partial \mathbf{F}(x, y, z, \dots)}{\partial x}, \frac{\partial \mathbf{F}(x, y, z, \dots)}{\partial y}, \frac{\partial \mathbf{F}(x, y, z, \dots)}{\partial z}, \dots$$

or simply to

$$\frac{\partial \mathbf{F}}{\partial x}, \frac{\partial \mathbf{F}}{\partial y}, \frac{\partial \mathbf{F}}{\partial z}.$$

As such the above expression can be written as

$$\delta \mathbf{F} = \frac{\partial \mathbf{F}}{\partial x} \delta x + \frac{\partial \mathbf{F}}{\partial y} \delta y + \frac{\partial \mathbf{F}}{\partial z} \delta z + \dots$$

Proceeding to the limit when $\delta F \rightarrow 0$, we have

$$dF = \frac{\partial F}{\partial x} dx + \frac{\partial F}{\partial y} dy + \frac{\partial F}{\partial z} dz + \dots$$

This gives the total differential of F for the scalar variables x, y, z . The total differential dF is given by

$$dF = du\mathbf{i} + dv\mathbf{j} + dw\mathbf{k}$$

When $\mathbf{F} = u(x, y, z)\mathbf{i} + v(x, y, z)\mathbf{j} + w(x, y, z)\mathbf{k}$

$$\text{Where } du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy + \frac{\partial u}{\partial z} dz$$

$$dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy + \frac{\partial v}{\partial z} dz$$

$$dw = \frac{\partial w}{\partial x} dx + \frac{\partial w}{\partial y} dy + \frac{\partial w}{\partial z} dz.$$

1.20. RULES FOR PARTIAL DIFFERENTIATION OF VECTOR

If $\mathbf{r}$ and $\mathbf{s}$ are functions of x, y, z , then

$$(1) \quad \frac{\partial}{\partial x}(\mathbf{r} \cdot \mathbf{s}) = \mathbf{r} \cdot \frac{\partial \mathbf{s}}{\partial x} + \frac{\partial \mathbf{r}}{\partial x} \cdot \mathbf{s}.$$

$$(2) \quad \frac{\partial}{\partial x}(\mathbf{r} \times \mathbf{s}) = \mathbf{r} \times \frac{\partial \mathbf{s}}{\partial x} + \frac{\partial \mathbf{r}}{\partial x} \times \mathbf{s}.$$

$$(3) \quad \frac{\partial^2}{\partial y \partial x}(\mathbf{r} \cdot \mathbf{s}) = \frac{\partial}{\partial y} \left\{ \frac{\partial}{\partial x}(\mathbf{r} \cdot \mathbf{s}) \right\} = \frac{\partial}{\partial y} \left\{ \mathbf{r} \cdot \frac{\partial \mathbf{s}}{\partial x} + \frac{\partial \mathbf{r}}{\partial x} \cdot \mathbf{s} \right\} \\ = \mathbf{r} \cdot \frac{\partial^2 \mathbf{s}}{\partial y \partial x} + \frac{\partial \mathbf{r}}{\partial y} \cdot \frac{\partial \mathbf{s}}{\partial x} + \frac{\partial \mathbf{r}}{\partial x} \cdot \frac{\partial \mathbf{s}}{\partial y} + \frac{\partial^2 \mathbf{r}}{\partial y \partial x} \cdot \mathbf{s}.$$

The proofs of these results are similar to those given in §1.18.

Problem 72. Find $\frac{\partial \mathbf{r}}{\partial x}, \frac{\partial \mathbf{r}}{\partial y}, \frac{\partial^2 \mathbf{r}}{\partial x^2}, \frac{\partial^2 \mathbf{r}}{\partial y^2}, \frac{\partial^2 \mathbf{r}}{\partial x \partial y}$ for the following functions

$$(a) \quad \mathbf{r} = x \cos y\mathbf{i} + x \sin y\mathbf{j} + ae^{my}\mathbf{k}.$$

$$(b) \quad \mathbf{r} = \frac{1}{2}a(x+y)\mathbf{i} + b/2(x-y)\mathbf{j} + \frac{1}{2}xy\mathbf{k}.$$

$$(c) \quad \mathbf{r} = x \cos y\mathbf{i} + x \sin y\mathbf{j} + c \log \{x + \sqrt{(x^2 - c^2)}\} \mathbf{k}.$$

$$(a) \quad \text{Given } \mathbf{r} = x \cos y\mathbf{i} + x \sin y\mathbf{j} + ae^{my}\mathbf{k}$$

$$\therefore \frac{\partial \mathbf{r}}{\partial x} = \cos y\mathbf{i} + \sin y\mathbf{j}$$

$$\frac{\partial \mathbf{r}}{\partial y} = -x \sin y\mathbf{i} + x \cos y\mathbf{j} + ame^{my}\mathbf{k}$$

$$\frac{\partial^2 \mathbf{r}}{\partial x^2} = 0$$

$$\frac{\partial^2 \mathbf{r}}{\partial y^2} = -x \cos y\mathbf{i} - x \sin y\mathbf{j} + am^2 e^{my}\mathbf{k}$$

$$\frac{\partial^2 \mathbf{r}}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial \mathbf{r}}{\partial y} \right) = -\sin y\mathbf{i} + \cos y\mathbf{j}.$$

$$(b) \quad \text{Given } \mathbf{r} = \frac{1}{2}a(x+y)\mathbf{i} + \frac{1}{2}b(x-y)\mathbf{j} + \frac{1}{2}xy\mathbf{k}$$

$$\frac{\partial \mathbf{r}}{\partial x} = \frac{1}{2} [a\mathbf{i} + b\mathbf{j} + y\mathbf{k}]$$

$$\frac{\partial \mathbf{r}}{\partial y} = \frac{1}{2} [a\mathbf{i} + b\mathbf{j} + x\mathbf{k}]$$

$$\frac{\partial^2 \mathbf{r}}{\partial x^2} = 0$$

$$\frac{\partial^2 \mathbf{r}}{\partial y^2} = \frac{1}{2}\mathbf{k}$$

$$\frac{\partial^2 \mathbf{r}}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial \mathbf{r}}{\partial y} \right) = 0.$$

$$(c) \quad \text{Given } \mathbf{r} = x \cos y\mathbf{i} + x \sin y\mathbf{j} + c \log \{x + \sqrt{(x^2 - c^2)}\} \mathbf{k}$$

$$\frac{\partial \mathbf{r}}{\partial x} = \cos y\mathbf{i} + \sin y\mathbf{j} - c \left\{ \frac{[1 + \frac{1}{2}(x^2 - c^2)^{-1/2} \cdot 2x]}{x + \sqrt{(x^2 - c^2)}} \right\} \mathbf{k}$$

$$= \cos y\mathbf{i} + \sin y\mathbf{j} - \frac{c}{\sqrt{(x^2 - c^2)}} \mathbf{k}$$

$$\frac{\partial \mathbf{r}}{\partial y} = -x \sin y\mathbf{i} + x \cos y\mathbf{j}$$

$$\frac{\partial^2 \mathbf{r}}{\partial x^2} = \frac{1}{2}c \frac{\mathbf{k} \cdot 2x}{(x^2 - c^2)^{3/2}}$$

$$= \frac{2x\mathbf{k}}{(x^2 - c^2)^{3/2}}$$

$$\frac{\partial^2 \mathbf{r}}{\partial y^2} = -x \cos y\mathbf{i} - x \sin y\mathbf{j}$$

$$\frac{\partial^2 \mathbf{r}}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial \mathbf{r}}{\partial y} \right) = -\sin y\mathbf{i} + \cos y\mathbf{j}.$$

Problem 73. If $\mathbf{A} = x^2yz\mathbf{i} - 2xz^3\mathbf{j} + xz^3\mathbf{k}$, $\mathbf{B} = 2z\mathbf{i} + y\mathbf{j} - x^2\mathbf{k}$, find

$$\frac{\partial^2}{\partial x \partial y} (\mathbf{A} \times \mathbf{B}) \text{ at } (1, 0, -2).$$

$$\text{We have } \mathbf{A} \times \mathbf{B} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ x^2yz - 2xz^3 & xz^3 & 2z \\ 2z & y & -x^2 \end{vmatrix} \\ = \mathbf{i}(2x^3z^3 - xyz^2) - \mathbf{j}(-x^4yz - 2xz^3) + \mathbf{k}(x^2y^2z + xz^4)$$

$$\therefore \frac{\partial}{\partial y} (\mathbf{A} \times \mathbf{B}) = \mathbf{i}(-xz^2) - \mathbf{j}(-x^4z) + \mathbf{k}(2x^2yz)$$

$$\text{and } \frac{\partial^2}{\partial x \partial y} (\mathbf{A} \times \mathbf{B}) = \frac{\partial}{\partial x} \left\{ \frac{\partial}{\partial y} (\mathbf{A} \times \mathbf{B}) \right\} = -z^2\mathbf{i} + 4xz^3\mathbf{j} + 4xyz\mathbf{k} \\ = -4\mathbf{i} - 8\mathbf{j} \text{ at } (1, 0, -2)$$

Problem 74. If $\phi(x, y, z) = xyz^2z$ and $\mathbf{r} = xz\mathbf{i} - xy^2\mathbf{j} + yz^2\mathbf{k}$, find

$$\frac{\partial^3}{\partial x^2 \partial z} (\phi \mathbf{r}) \text{ at the point } (2, -1, 1).$$

We have $\phi(\mathbf{r}) = (xy^2z)(xzi - xy^2\mathbf{j} + yz^2\mathbf{k})$
 $= x^2y^2z^2\mathbf{i} - x^2y^4\mathbf{j} + xy^3z^2\mathbf{k}$

$$\therefore \frac{\partial(\phi\mathbf{r})}{\partial z} = 2x^2y^2z\mathbf{i} - x^2y^4\mathbf{j} + 3xy^3z^2\mathbf{k}$$

and $\frac{\partial z}{\partial x \partial z}(\phi\mathbf{r}) = \frac{\partial}{\partial x} \left(\frac{\partial \phi\mathbf{r}}{\partial z} \right) = 4xy^2z\mathbf{i} - 2xy^4\mathbf{j} + 3y^3z^2\mathbf{k}$

So that

$$\frac{\partial^3}{\partial x^2 \partial z}(\phi\mathbf{r}) = \frac{\partial}{\partial x} \left\{ \frac{\partial^2(\phi\mathbf{r})}{\partial x \partial z} \right\} = 4y^2z\mathbf{i} - 2y^4\mathbf{j}$$

$$= 4\mathbf{i} - 2\mathbf{j} \text{ at } (2, -1, 1)$$

i.e. at $x=2, y=-1, z=1$.

Problem 75. If $\mathbf{F}$ depends on x, y, z, t where x, y, z depend on t prove that $\frac{d\mathbf{F}}{dt} = \frac{\partial \mathbf{F}}{\partial t} + \frac{\partial \mathbf{F}}{\partial x} \frac{dx}{dt} + \frac{\partial \mathbf{F}}{\partial y} \frac{dy}{dt} + \frac{\partial \mathbf{F}}{\partial z} \frac{dz}{dt}$.

Let us suppose that

$$\mathbf{F} = f_1(x, y, z, t)\mathbf{i} + f_2(x, y, z, t)\mathbf{j} + f_3(x, y, z, t)\mathbf{k} \dots$$

So that $d\mathbf{F} = df_1\mathbf{i} + df_2\mathbf{j} + df_3\mathbf{k} \dots$

$$\text{Now } df_1 = \frac{\partial f_1}{\partial x} dx + \frac{\partial f_1}{\partial y} dy + \frac{\partial f_1}{\partial z} dz + \frac{\partial f_1}{\partial t} dt$$

$$df_2 = \frac{\partial f_2}{\partial x} dx + \frac{\partial f_2}{\partial y} dy + \frac{\partial f_2}{\partial z} dz + \frac{\partial f_2}{\partial t} dt$$

$$df_3 = \frac{\partial f_3}{\partial x} dx + \frac{\partial f_3}{\partial y} dy + \frac{\partial f_3}{\partial z} dz + \frac{\partial f_3}{\partial t} dt$$

Putting the values of, df_1, df_2, df_3 , in (ii) we get

$$d\mathbf{F} = \left(\frac{\partial f_1}{\partial x} dx + \frac{\partial f_1}{\partial y} dy + \frac{\partial f_1}{\partial z} dz + \frac{\partial f_1}{\partial t} dt \right) \mathbf{i} + \dots + \dots$$

$$= \left(\frac{\partial f_1}{\partial x} \mathbf{i} + \frac{\partial f_2}{\partial x} \mathbf{j} + \frac{\partial f_3}{\partial x} \mathbf{k} \right) dx + \left(\frac{\partial f_1}{\partial y} \mathbf{i} + \frac{\partial f_2}{\partial y} \mathbf{j} + \frac{\partial f_3}{\partial y} \mathbf{k} \right) dy$$

$$+ \left(\frac{\partial f_1}{\partial z} \mathbf{i} + \frac{\partial f_2}{\partial z} \mathbf{j} + \frac{\partial f_3}{\partial z} \mathbf{k} \right) dz + \left(\frac{\partial f_1}{\partial t} \mathbf{i} + \frac{\partial f_2}{\partial t} \mathbf{j} + \frac{\partial f_3}{\partial t} \mathbf{k} \right) dt$$

or $\frac{d\mathbf{F}}{dt} = \left(\frac{\partial f_1}{\partial x} \mathbf{i} + \frac{\partial f_2}{\partial x} \mathbf{j} + \frac{\partial f_3}{\partial x} \mathbf{k} \right) \frac{dx}{dt} + \left(\frac{\partial f_1}{\partial y} \mathbf{i} + \frac{\partial f_2}{\partial y} \mathbf{j} + \frac{\partial f_3}{\partial y} \mathbf{k} \right) \frac{dy}{dt}$

$$+ \left(\frac{\partial f_1}{\partial z} \mathbf{i} + \frac{\partial f_2}{\partial z} \mathbf{j} + \frac{\partial f_3}{\partial z} \mathbf{k} \right) \frac{dz}{dt} + \left(\frac{\partial f_1}{\partial t} \mathbf{i} + \frac{\partial f_2}{\partial t} \mathbf{j} + \frac{\partial f_3}{\partial t} \mathbf{k} \right) \frac{dt}{dt}$$

$$= \frac{\partial \mathbf{F}}{\partial x} \frac{dx}{dt} + \frac{\partial \mathbf{F}}{\partial y} \frac{dy}{dt} + \frac{\partial \mathbf{F}}{\partial z} \frac{dz}{dt} + \frac{\partial \mathbf{F}}{\partial t}$$

Problem 76. Prove that $\mathbf{F} = \frac{\mathbf{a} e^{i\omega(t-r/c)}}{r}$ satisfies the partial differential equation

$$\frac{\partial^2 \mathbf{F}}{\partial r^2} + \frac{2}{r} \frac{\partial \mathbf{F}}{\partial r} = \frac{1}{c^2} \frac{\partial^2 \mathbf{F}}{\partial t^2}$$

where $\mathbf{a}$ is a constant vector, ω and c are scalar constants and $i = \sqrt{-1}$.

We are given that

$$\mathbf{F} = \frac{\mathbf{a} e^{i\omega(t-r/c)}}{r}$$

$$\frac{\partial \mathbf{F}}{\partial r} = \frac{[r e^{-i\omega r/c} \times -i\omega/c - e^{i\omega(t-r/c)}]}{r^2}$$

$$= -\frac{i\omega}{cr} \mathbf{a} e^{-i\omega r/c} - \mathbf{a} \frac{e^{i\omega(t-r/c)}}{r^2}$$

$$\frac{\partial^2 \mathbf{F}}{\partial r^2} = -\frac{i\omega}{c} \mathbf{a} \frac{[r e^{-i\omega r/c} \times -i\omega/c - e^{-i\omega r/c}]}{r^2}$$

$$= -\mathbf{a} \frac{[r^2 e^{-i\omega r/c} \times -i\omega/c - 2r e^{i\omega(t-r/c)}]}{r^4}$$

$$= -\frac{\omega^2}{c^2} \mathbf{a} \frac{1}{r} e^{-i\omega r/c} + \frac{i\omega \mathbf{a}}{cr^2} e^{-i\omega r/c} + \frac{i\omega}{cr^2} e^{-i\omega r/c} + \frac{2\mathbf{a}}{r^3} e^{i\omega(t-r/c)}$$

$$= -\frac{\omega^2}{c^2} \mathbf{a} \frac{1}{r} e^{-i\omega r/c} + \frac{2i\omega \mathbf{a}}{cr^2} e^{-i\omega r/c} + \frac{2\mathbf{a}}{r^3} e^{i\omega(t-r/c)}$$

Again $\frac{\partial \mathbf{F}}{\partial t} = \frac{\mathbf{a}}{r} e^{i\omega t} \times i\omega = \frac{i\omega \mathbf{a}}{r} e^{i\omega t}$

and $\frac{\partial^2 \mathbf{F}}{\partial t^2} = \frac{i^2 \omega^2 \mathbf{a}}{r} e^{i\omega t} = -\frac{\omega \mathbf{a}}{r} e^{i\omega t}$

So that

$$\frac{\partial^2 \mathbf{F}}{\partial r^2} = -\frac{\omega^2 \mathbf{a}}{c^2 r} e^{-i\omega r/c} + \frac{2i\omega \mathbf{a}}{cr^2} e^{-i\omega r/c} + \frac{2\mathbf{a}}{r^3} e^{i\omega(t-r/c)} \dots (1)$$

$$\frac{2}{r} \left(\frac{\partial \mathbf{F}}{\partial r} \right) = -\frac{2i\omega \mathbf{a}}{cr^2} e^{-i\omega r/c} - \frac{2\mathbf{a} e^{i\omega(t-r/c)}}{r^3} \dots (2)$$

$$\text{and } \frac{1}{c^2} \left(\frac{\partial^2 \mathbf{F}}{\partial t^2} \right) = -\frac{2\omega \mathbf{a}}{rc^2} e^{i\omega t} \dots (3)$$

From (1), (2) and (3), it is evident that

$$\frac{\partial^2 \mathbf{F}}{\partial r^2} + \frac{2}{r} \frac{\partial \mathbf{F}}{\partial r} = \frac{1}{c^2} \frac{\partial^2 \mathbf{F}}{\partial t^2}$$

11. THE SCALAR AND VECTOR FIELDS

A physical quantity which is expressible as a continuous function and which can assume one or more definite values at each point of a region of space, is said to be a *point function* in that region and the region specifying the physical quantity is called as a *field*. Point functions are of two kinds: *Scalar-point function* and *Vector-point function* according to the nature of the quantity concerned.

(1) **Scalar-point function.** At each point P of a domain or region the function denoted by $f(P)$, or $f(x, y, z)$ is known as a *scalar-point function*. The points of the region R together with the function values $f(P)$ constitute a *scalar-field* over R .

The examples of scalar-fields are the temperature distribution in medium, the gravitational potential of a system of masses, and the electrostatic potential of a system of charges, etc.

(2) **Vector-point function.** If to each point P of a region R , there is associated a vector $\mathbf{F}(P)$, the function $\mathbf{F}(P)$ is known as a *vector-point function*, and points of R together with these vectors constitute a *vector-field* over R .

The examples of vector-fields are the velocity of a moving particle, the electrostatic, the magnetostatic, and gravitational fields, the electric intensity of force etc.

Continuity of Scalar and Vector-point Functions

(1) **Scalar-point function.** A scalar-point function $f(P)$ is continuous at a point $P=P_0$, if

(i) $f(P)$ is defined.

(ii) Given a number $\epsilon > 0$ however small there exists a positive number δ such that

$$\begin{aligned} & |f(P) - f(P_0)| < \epsilon \\ & \text{provided } |P - P_0| < \delta, \text{ where } \delta \text{ depends on } \epsilon \text{ and } P \text{ both.} \\ & \text{i.e. } \delta = \delta(\epsilon, P) \end{aligned}$$

In other words, a scalar point function $f(P)$ is said to be *continuous* at a point P_0 , if

$$\lim_{P \rightarrow P_0} f(P) = f(P_0)$$

The scalar-point function $f(P)$ is continuous in a region if it is continuous at every point of the region.

(2) **Vector-point function.** The continuity of vector-point function is defined similarly as in case of scalar-point functions except that it is replaced by $\mathbf{f}$ for vector-point functions.

1.22. DIRECTIONAL DERIVATIVES

If $f(x, y, z)$ be defined and differentiable at each point (x, y, z) in a certain region of space, then the partial derivative $\frac{\partial f}{\partial x}$ is defined as the rate of change of f at (x, y, z) in the direction of axis of x and is called as the *directional derivative*.

(1) **Scalar-point function.** Consider a line QP drawn in a scalar-field f , the sense of it being positive from Q to P . Let $f(Q)$ and $f(P)$ be two function values of scalar-point function f at Q and P respectively.

Then, if $\lim_{P \rightarrow Q} \frac{f(P) - f(Q)}{QP}$ exists, it




Fig. 1.35

called *directional-derivative* of the scalar-point function at Q along QP .

(2) **Vector-point function.** The directional derivatives for vector-point functions is defined similarly as in case of directional derivative for scalar-point functions except that f is replaced by $\mathbf{f}$ for vector-point function i.e.

$$\lim_{P \rightarrow Q} \frac{\mathbf{f}(P) - \mathbf{f}(Q)}{QP} \text{ exists,}$$

it is called *directional derivative* of a vector-point function at a point along the direction of the vector.

(3) **Directional derivatives of scalar-point function along coordinate axes.** Referred to Fig. 1.36, we have $QP = \delta x$.

The definition of directional derivative for scalar-point function, then leads

$$\begin{aligned} & \lim_{P \rightarrow Q} \frac{f(P) - f(Q)}{QP} \\ &= \lim_{\delta x \rightarrow 0} \frac{f(x + \delta x, y, z) - f(x, y, z)}{\delta x} \\ &= \text{partial derivative of } f \text{ with respect to } x \\ &= \frac{\partial f}{\partial x}. \end{aligned}$$




Fig. 1.36

Proceeding similarly, we find that directional derivative at Q along the positive directions of axis of y and axis of z are $\frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$ respectively.

(4) **Directional derivative of vector-point function along coordinate axes.** Proceeding as in case of (3), we may find that directional derivative of vector-point functions in the positive directions of axes are

$$\frac{\partial \mathbf{f}}{\partial x}, \frac{\partial \mathbf{f}}{\partial y}, \frac{\partial \mathbf{f}}{\partial z} \text{ respectively.}$$

(5) **Directional derivative of scalar-point function along any line.**

If QP be a line in the space in positive sense being from Q to P , and the direction cosines of QP are l, m, n , then the coordinates of the point P such that $QP = r$, are

$$(x + lr, y + mr, z + nr).$$

Thus the definition of directional derivative gives

$$\begin{aligned} & \lim_{P \rightarrow Q} \frac{f(P) - f(Q)}{QP} \\ &= \lim_{r \rightarrow 0} \frac{f(x + lr, y + mr, z + nr) - f(x, y, z)}{r} \end{aligned} \quad \dots (i)$$




Fig. 1.37

If $f(x, y, z)$ possesses first order continuous partial derivatives then $f(x+lr, y+mr, z+nr)$ can be expanded by Taylor's Theorem so that we have

$$\begin{aligned} f(x+lr, y+mr, z+nr) = & f(x, y, z) + lr \frac{\partial f}{\partial x}(x_1, y_1, z_1) \\ & + mr \frac{\partial f}{\partial y}(x_1, y_1, z_1) + nr \frac{\partial f}{\partial z}(x_1, y_1, z_1) \dots \end{aligned}$$

where (x_1, y_1, z_1) is a point on QP .

$$\begin{aligned} \text{Hence } \lim_{P \rightarrow Q} \frac{f(P) - f(Q)}{QP} &= l \frac{\partial f}{\partial x}(x, y, z) \\ &+ m \frac{\partial f}{\partial y}(x, y, z) + n \frac{\partial f}{\partial z}(x, y, z) \quad [\text{from (i) and (ii)}] \\ &= l \frac{\partial f}{\partial x} + m \frac{\partial f}{\partial y} + n \frac{\partial f}{\partial z} \end{aligned}$$

which follows that the directional derivatives along any line can be expressed in terms of those along three coordinate axes.

(6) Directional derivative for vector-point function along any line
Just as in case (5), take the direction cosines of a line as, l, m, n and define a vector-point function $\mathbf{f}$ in the region of line in terms of unit vectors $\mathbf{i}, \mathbf{j}, \mathbf{k}$, as

$$\mathbf{f}(x, y, z) = f_1(x, y, z) \mathbf{i} + f_2(x, y, z) \mathbf{j} + f_3(x, y, z) \mathbf{k}.$$

If $\mathbf{f}$ possesses first order continuous partial derivatives then components f_1, f_2, f_3 will also possess first order continuous partial derivatives.

The definition of directional derivative for vector-point function gives

$$\begin{aligned} \lim_{P \rightarrow Q} \frac{\mathbf{f}(P) - \mathbf{f}(Q)}{QP} &= \mathbf{i} \frac{f_1(P) - f_1(Q)}{QP} + \mathbf{j} \frac{f_2(P) - f_2(Q)}{QP} \\ &+ \mathbf{k} \frac{f_3(P) - f_3(Q)}{QP} \\ &= \mathbf{i} \left[l \frac{\partial f_1}{\partial x} + m \frac{\partial f_1}{\partial y} + n \frac{\partial f_1}{\partial z} \right] + \mathbf{j} \left[l \frac{\partial f_2}{\partial x} + m \frac{\partial f_2}{\partial y} + n \frac{\partial f_2}{\partial z} \right] \\ &+ \mathbf{k} \left[l \frac{\partial f_3}{\partial x} + m \frac{\partial f_3}{\partial y} + n \frac{\partial f_3}{\partial z} \right] \text{ by (1)} \\ &= \left[\mathbf{i} \frac{\partial f_1}{\partial x} + \mathbf{j} \frac{\partial f_2}{\partial x} + \mathbf{k} \frac{\partial f_3}{\partial x} \right] l + \left[\mathbf{i} \frac{\partial f_1}{\partial y} + \mathbf{j} \frac{\partial f_2}{\partial y} + \mathbf{k} \frac{\partial f_3}{\partial y} \right] m \\ &+ \left[\mathbf{i} \frac{\partial f_1}{\partial z} + \mathbf{j} \frac{\partial f_2}{\partial z} + \mathbf{k} \frac{\partial f_3}{\partial z} \right] n \\ &= \frac{\partial \mathbf{f}}{\partial x} l + \frac{\partial \mathbf{f}}{\partial y} m + \frac{\partial \mathbf{f}}{\partial z} n \end{aligned}$$

which is similar to the direction derivative of scalar-point function.

1.3. LEVEL SURFACES

Assuming that f is a continuous point-function, through any point P of the region considered, we can draw a surface such that, at each point on it the function has the same value as that at P . Such a surface is called a level surface of the function.

The examples of level surface are: isothermal surfaces and equipotential surfaces for temperature and potential respectively.

1.4. THE GRADIENT OF A SCALAR FIELD

Consider a scalar function i.e. a function whose value depends upon the values of co-ordinates (x, y, z) . Being a scalar its value is constant at a fixed point in space.

The gradient of any scalar function ϕ is defined as

$$\begin{aligned} \text{grad } \phi &= \mathbf{i} \frac{\partial \phi}{\partial x} + \mathbf{j} \frac{\partial \phi}{\partial y} + \mathbf{k} \frac{\partial \phi}{\partial z} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \phi \\ &= \nabla \phi \end{aligned} \quad \dots (1)$$

The operator ∇ is generally known as 'del' or 'nabla' operator and is called a 'gradient' or 'grad' in short. We have already mentioned that a scalar field is the region in which the scalar point function specifies a scalar physical quantity like temperature, electric potential, etc. It is represented by a continuous scalar function giving the value of the quantity at each point. In scalar field all the points having same value of ϕ can be connected by means of surfaces, which are called equal or level surfaces.

Consider a co-ordinate system with axes such that any level surface is in xy plane while z -axis is along the normal to that level surface. Then the value of ϕ does not change along the level surface, i.e.

$$\frac{\partial \phi}{\partial x} = \frac{\partial \phi}{\partial y} = 0,$$

$$\text{therefore } \text{grad } \phi = \mathbf{k} \frac{\partial \phi}{\partial z}. \quad \dots (2)$$

Clearly $\text{grad } \phi$ is directed along z -axis, i.e., along the normal to the level surface. Therefore equation (2) may be written as

$$\text{grad } \phi = \frac{\partial \phi}{\partial n} \mathbf{n} \quad \dots (3)$$

where $\mathbf{n}$ is unit vector along the normal to the level surface at any point.

From equation (3) we may state, "The magnitude of $\text{grad } \phi$ at any point is rate of change of function ϕ with distance along the normal to the level surface at the point and is directed along unit vector $\mathbf{n}$."

It is to be noted that gradient of any scalar quantity is a vector. (Agra, 1965)

Problem 77. Prove $\nabla r^n = nr^{n-2} \mathbf{r}$.

$$\begin{aligned}
 \text{L.H.S.} &= \nabla r^n = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (r^n) = \mathbf{i} \frac{\partial r^n}{\partial x} + \mathbf{j} \frac{\partial r^n}{\partial y} + \mathbf{k} \frac{\partial r^n}{\partial z} \\
 &= \mathbf{i} n r^{n-1} \frac{\partial r}{\partial x} + \mathbf{j} n r^{n-1} \frac{\partial r}{\partial y} + \mathbf{k} n r^{n-1} \frac{\partial r}{\partial z} \\
 &= n r^{n-1} \left[\mathbf{i} \frac{\partial r}{\partial x} + \mathbf{j} \frac{\partial r}{\partial y} + \mathbf{k} \frac{\partial r}{\partial z} \right] \left[\text{since } r^2 = x^2 + y^2 + z^2; \right. \\
 &\quad \left. \therefore \frac{\partial r}{\partial x} = \frac{x}{r}, \frac{\partial r}{\partial y} = \frac{y}{r}, \frac{\partial r}{\partial z} = \frac{z}{r} \right] \\
 &= n r^{n-2} (\mathbf{i}x + \mathbf{j}y + \mathbf{k}z) = n r^{n-2} \mathbf{r}.
 \end{aligned}$$

Problem 78. If $\mathbf{r}$ is the position vector of a point, deduce the value of $\text{grad}(1/r)$.

As given, $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$.

$$\text{So that } \frac{1}{r} = \frac{1}{(x^2 + y^2 + z^2)^{1/2}}.$$

$$\therefore \text{grad}(1/r) = \nabla(1/r)$$

$$\begin{aligned}
 &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \left\{ \frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right\} \\
 &= \mathbf{i} \frac{\partial}{\partial x} \left[\frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right] + \mathbf{j} \frac{\partial}{\partial y} \left[\frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right] \\
 &\quad + \mathbf{k} \frac{\partial}{\partial z} \left[\frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right] \\
 &= \mathbf{i} \left[-\frac{1}{2} \cdot \frac{2x}{(x^2 + y^2 + z^2)^{3/2}} \right] + \mathbf{j} \left[-\frac{1}{2} \cdot \frac{2y}{(x^2 + y^2 + z^2)^{3/2}} \right] \\
 &\quad + \mathbf{k} \left[-\frac{1}{2} \cdot \frac{2z}{(x^2 + y^2 + z^2)^{3/2}} \right] \\
 &= -\frac{x\mathbf{i} + y\mathbf{j} + z\mathbf{k}}{(x^2 + y^2 + z^2)^{3/2}} = -\frac{\mathbf{r}}{(r^2)^{3/2}} = -\frac{\mathbf{r}}{r^3}.
 \end{aligned}$$

1.25. THE GRADIENT OF A SCALAR-POINT FUNCTION

If $\phi(x, y, z)$ be defined and differentiable at each point (x, y, z) in a certain region of space specified as a scalar field, we have

$$\left(l \frac{\partial \phi}{\partial x} + m \frac{\partial \phi}{\partial y} + n \frac{\partial \phi}{\partial z} \right) = (\mathbf{l} + \mathbf{m}\mathbf{j} + \mathbf{n}\mathbf{k}) \cdot \left(\mathbf{i} \frac{\partial \phi}{\partial x} + \mathbf{j} \frac{\partial \phi}{\partial y} + \mathbf{k} \frac{\partial \phi}{\partial z} \right)$$

Its R.H.S. is the scalar product of two vectors $(\mathbf{l} + \mathbf{m}\mathbf{j} + \mathbf{n}\mathbf{k})$ and $\left(\mathbf{i} \frac{\partial \phi}{\partial x} + \mathbf{j} \frac{\partial \phi}{\partial y} + \mathbf{k} \frac{\partial \phi}{\partial z} \right)$, where the vector $(\mathbf{l} + \mathbf{m}\mathbf{j} + \mathbf{n}\mathbf{k})$ is a unit vector along a line whose direction cosines are l, m, n and the second vector depends only on the point (x, y, z) and not on any direction. We conclude that directional derivative along any line can be obtained by multiplying the vector

$$\mathbf{i} \frac{\partial \phi}{\partial x} + \mathbf{j} \frac{\partial \phi}{\partial y} + \mathbf{k} \frac{\partial \phi}{\partial z}$$

scalarly with the unit vector

$$\mathbf{l} + \mathbf{m}\mathbf{j} + \mathbf{n}\mathbf{k}.$$

The vector function

$$\mathbf{i} \frac{\partial \phi}{\partial x} + \mathbf{j} \frac{\partial \phi}{\partial y} + \mathbf{k} \frac{\partial \phi}{\partial z}$$

is called the *gradient of a scalar-point function* ϕ and is written as $\text{grad } \phi$ or $\nabla \phi$. Thus,

$$\nabla \phi = \text{grad } \phi \equiv \mathbf{i} \frac{\partial \phi}{\partial x} + \mathbf{j} \frac{\partial \phi}{\partial y} + \mathbf{k} \frac{\partial \phi}{\partial z} \equiv \frac{\partial \phi}{\partial x} \mathbf{i} + \frac{\partial \phi}{\partial y} \mathbf{j} + \frac{\partial \phi}{\partial z} \mathbf{k}.$$

It is clear that the gradient of a scalar-point function is a vector.

In case, ϕ is a constant, $\text{grad } \phi = 0$ since $\frac{\partial \phi}{\partial x}, \frac{\partial \phi}{\partial y}, \frac{\partial \phi}{\partial z}$ all will be zero in that case. Its converse is also true.

1.26. THE GRADIENT OF SUM OF TWO SCALAR-POINT FUNCTIONS

If u and v are two differentiable scalar functions of x, y, z , then the gradient of their sum is given by

$$\begin{aligned}
 \nabla(u+v) &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (u+v) \\
 &= \mathbf{i} \frac{\partial}{\partial x} (u+v) + \mathbf{j} \frac{\partial}{\partial y} (u+v) + \mathbf{k} \frac{\partial}{\partial z} (u+v) \\
 &= \mathbf{i} \frac{\partial u}{\partial x} + \mathbf{i} \frac{\partial v}{\partial x} + \mathbf{j} \frac{\partial u}{\partial y} + \mathbf{j} \frac{\partial v}{\partial y} + \mathbf{k} \frac{\partial u}{\partial z} + \mathbf{k} \frac{\partial v}{\partial z} \\
 &= \left(\mathbf{i} \frac{\partial u}{\partial x} + \mathbf{j} \frac{\partial u}{\partial y} + \mathbf{k} \frac{\partial u}{\partial z} \right) + \left(\mathbf{i} \frac{\partial v}{\partial x} + \mathbf{j} \frac{\partial v}{\partial y} + \mathbf{k} \frac{\partial v}{\partial z} \right) \\
 &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) u + \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) v \\
 &= \nabla u + \nabla v.
 \end{aligned}$$

Showing that the gradient of sum of two scalar-point functions is equal to the sum of their gradients.

This rule may be generalized for any number of scalar-point functions.

1.27. THE GRADIENT OF PRODUCT OF TWO SCALAR-POINT FUNCTIONS

If u and v be two differentiable scalar-point functions of x, y, z then the gradient of their product is given by

$$\nabla(uv) = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (uv)$$

$$\begin{aligned}
&= \mathbf{i} \frac{\partial}{\partial x}(uv) + \mathbf{j} \frac{\partial}{\partial y}(uv) + \mathbf{k} \frac{\partial}{\partial z}(uv) \\
&= \mathbf{i} \left(u \frac{\partial v}{\partial x} + v \frac{\partial u}{\partial x} \right) + \mathbf{j} \left(u \frac{\partial v}{\partial y} + v \frac{\partial u}{\partial y} \right) + \mathbf{k} \left(u \frac{\partial v}{\partial z} + v \frac{\partial u}{\partial z} \right) \\
&= u \left[\mathbf{i} \frac{\partial v}{\partial x} + \mathbf{j} \frac{\partial v}{\partial y} + \mathbf{k} \frac{\partial v}{\partial z} \right] + v \left[\mathbf{i} \frac{\partial u}{\partial x} + \mathbf{j} \frac{\partial u}{\partial y} + \mathbf{k} \frac{\partial u}{\partial z} \right] \\
&= u \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) v + v \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) u \\
&= u \nabla v + v \nabla u.
\end{aligned}$$

Showing that the gradient of the product of two scalar-point functions is obtained by the same rule as is valid for derivatives of the algebraic functions.

Problem 79. Find $\nabla \phi$ and $|\nabla \phi|$ for the function.

$$\phi = 2xz^4 - x^2y \text{ at the point } (2, -2, -1).$$

We know by the definition of grad ϕ that

$$\nabla \phi = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \phi$$

Here

$$\phi = 2xz^4 - x^2y$$

$$\therefore \nabla \phi = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (2xz^4 - x^2y)$$

$$\begin{aligned}
&= \mathbf{i} \frac{\partial}{\partial x} (2xz^4 - x^2y) + \mathbf{j} \frac{\partial}{\partial y} (2xz^4 - x^2y) \\
&\quad + \mathbf{k} \frac{\partial}{\partial z} (2xz^4 - x^2y)
\end{aligned}$$

$$= \mathbf{i} [2z^4 - 2xy] + \mathbf{j} [-x^2] + \mathbf{k} [8xz^3]$$

$$= \mathbf{i} [2(-1)^4 - 2(2)(-2)] + \mathbf{j} [-(+2)^2]$$

$$+ \mathbf{k} [8(2)(-1)^3] \text{ at } x=2, y=-2, z=-1$$

$$= \mathbf{i} [2+8] + \mathbf{j} [-4] + \mathbf{k} [-16]$$

$$= 10\mathbf{i} - 4\mathbf{j} - 16\mathbf{k}.$$

$$\text{And } |\nabla \phi| = \sqrt{[(10)^2 + (-4)^2 + (-16)^2]}$$

$$= \sqrt{(372)} = 2\sqrt{93}.$$

Problem 80. If $u = x^2z + e^{y/x}$ and $v = 2z^2y - xy^2$ find

(a) $\nabla(u+v)$, (b) $\nabla(uv)$ at the point $(1, 0, -2)$.

We have

$$\nabla(u+v) = \nabla u + \nabla v$$

$$\nabla(uv) = u\nabla v + v\nabla u$$

Given that $u = x^2z + e^{y/x}$

$$\therefore \nabla u = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (x^2z + e^{y/x})$$

$$= \mathbf{i} \frac{\partial}{\partial x} (x^2z + e^{y/x}) + \mathbf{j} \frac{\partial}{\partial y} (x^2z + e^{y/x}) + \mathbf{k} \frac{\partial}{\partial z} (x^2z + e^{y/x})$$

$$= \mathbf{i} \left[2xz + e^{y/x} \times -\frac{y}{x^2} \right] + \mathbf{j} \left[e^{y/x} \times \frac{1}{x} \right] + \mathbf{k} [x^2]$$

$$= \mathbf{i} \left[2(1)(-2) + e^0 \times -\frac{0}{1} \right] + \mathbf{j} \left[e^0 \times \frac{1}{1} \right] + \mathbf{k} [(1)^2]$$

$$= -4\mathbf{i} + \mathbf{j} + \mathbf{k} \dots (2)$$

$$\text{at } x=1, y=0, z=-2$$

And

$$v = 2z^2y - xy^2$$

$$\nabla v = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (2z^2y - xy^2)$$

$$= \mathbf{i} \frac{\partial}{\partial x} (2z^2y - xy^2) + \mathbf{j} \frac{\partial}{\partial y} (2z^2y - xy^2) + \mathbf{k} \frac{\partial}{\partial z} (2z^2y - xy^2)$$

$$= \mathbf{i} [-y^2] + \mathbf{j} [2z^2 - 2xy] + \mathbf{k} [4zy]$$

$$= \mathbf{i} [0] + \mathbf{j} [2(-2)^2] + \mathbf{k} [0] \text{ at } x=1, y=0, z=-2$$

$$= 8\mathbf{j}.$$

$$\dots (3)$$

Putting the values of ∇u and ∇v from (2) and (3) in (1) we get

$$(a) \nabla(u+v) = \nabla u + \nabla v = (-4\mathbf{i} + \mathbf{j} + \mathbf{k}) + 8\mathbf{j} = -4\mathbf{i} + 9\mathbf{j} + \mathbf{k}$$

$$(b) \nabla(uv) = u\nabla v + v\nabla u = (x^2z + e^{y/x}) 8\mathbf{j} + (2z^2y - xy^2)(-4\mathbf{i} + \mathbf{j} + \mathbf{k})$$

$$= (-1) 8\mathbf{j} = -8\mathbf{j} \text{ at } (1, 0, -2).$$

Problem 81. If $\Lambda = 2x^2\mathbf{i} - 3yz\mathbf{j} + xz^2\mathbf{k}$ and $\phi = 2z - x^3y$ find

(a) $\Lambda \cdot \nabla \phi$ (b) $\Lambda \times \nabla \phi$ at the point $(1, -1, 1)$.

Then

$$\phi = 2z - x^3y$$

$$\nabla \phi = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (2z - x^3y)$$

$$= \mathbf{i} (-3x^2y) + \mathbf{j} (-x^3) + \mathbf{k} (2)$$

$$= 3\mathbf{i} - \mathbf{j} + 2\mathbf{k} \text{ at } x=1, y=-1, z=1$$

$$\Lambda = 2x^2\mathbf{i} - 3yz\mathbf{j} + xz^2\mathbf{k}$$

$$= 2\mathbf{i} + 3\mathbf{j} + \mathbf{k} \text{ at } x=1, y=-1, z=1$$

$$(a) \Lambda \cdot \nabla \phi = (2\mathbf{i} + 3\mathbf{j} + \mathbf{k}) \cdot (3\mathbf{i} - \mathbf{j} + 2\mathbf{k})$$

$$= [6 - 3 + 2] = 5.$$

$$(b) \Lambda \times \nabla \phi = (2\mathbf{i} + 3\mathbf{j} + \mathbf{k}) \times (3\mathbf{i} - \mathbf{j} + 2\mathbf{k})$$

$$= [-2\mathbf{k} - 4\mathbf{j} - 9\mathbf{k} + 6\mathbf{i} + 3\mathbf{i} + \mathbf{j}]$$

$$= 7\mathbf{i} - \mathbf{j} - 11\mathbf{k}.$$

Problem 82. Prove $\nabla \left(\frac{u}{v} \right) = \frac{v \nabla u - u \nabla v}{v^2}$ provided $v \neq 0$.

Then

$$\phi = \frac{u}{v} = (uv^{-1})$$

$$\nabla \phi = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (uv^{-1})$$

$$= \mathbf{i} \frac{\partial}{\partial x} (uv^{-1}) + \mathbf{j} \frac{\partial}{\partial y} (uv^{-1}) + \mathbf{k} \frac{\partial}{\partial z} (uv^{-1})$$

$$\begin{aligned}
 &= \mathbf{i} \left[-\frac{u}{v^2} \frac{\partial v}{\partial x} + v^{-1} \frac{\partial u}{\partial x} \right] + \mathbf{j} \left[\frac{u}{v^2} \frac{\partial v}{\partial z} + v^{-1} \frac{\partial u}{\partial y} \right] \\
 &\quad + \mathbf{k} \left[\frac{u}{v^2} \frac{\partial v}{\partial z} + v^{-1} \frac{\partial u}{\partial z} \right] \\
 &= -\frac{u}{v^2} \left(\mathbf{i} \frac{\partial v}{\partial x} + \mathbf{j} \frac{\partial v}{\partial y} + \mathbf{k} \frac{\partial v}{\partial z} \right) + v^{-1} \left(\mathbf{i} \frac{\partial u}{\partial x} + \mathbf{j} \frac{\partial u}{\partial y} + \mathbf{k} \frac{\partial u}{\partial z} \right) \\
 &= -\frac{u}{v^2} \left[\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right] v \\
 &\quad + \frac{1}{v} \left[\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right] u \\
 &= -\frac{u}{v^2} \nabla v + \frac{1}{v} \nabla u = \frac{1}{v} \nabla u - \frac{u}{v^2} \nabla v = \frac{v \nabla u - u \nabla v}{v^2}
 \end{aligned}$$

Problem 83. If u is differentiable function of x, y, z , prove $\nabla u \cdot d\mathbf{r}$
 Given $v = u(x, y, z)$

i.e. $du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy + \frac{\partial u}{\partial z} dz$

and $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$.

$\therefore d\mathbf{r} = dx\mathbf{i} + dy\mathbf{j} + dz\mathbf{k}$.

Now $\nabla u = \left(\mathbf{i} \frac{\partial u}{\partial x} + \mathbf{j} \frac{\partial u}{\partial y} + \mathbf{k} \frac{\partial u}{\partial z} \right)$.

$$\begin{aligned}
 \therefore \nabla u \cdot d\mathbf{r} &= \left(\mathbf{i} \frac{\partial u}{\partial x} + \mathbf{j} \frac{\partial u}{\partial y} + \mathbf{k} \frac{\partial u}{\partial z} \right) \cdot d\mathbf{r} \\
 &= \left(\mathbf{i} \frac{\partial u}{\partial x} + \mathbf{j} \frac{\partial u}{\partial y} + \mathbf{k} \frac{\partial u}{\partial z} \right) \cdot (dx\mathbf{i} + dy\mathbf{j} + dz\mathbf{k}) \\
 &= \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy + \frac{\partial u}{\partial z} dz = du
 \end{aligned}$$

i.e. $\nabla u \cdot d\mathbf{r} = du$.

1.28. THE DIVERGENCE OF A VECTOR-POINT FUNCTION (Agra, 1952)

If $\mathbf{V}(x, y, z) = V_1\mathbf{i} + V_2\mathbf{j} + V_3\mathbf{k}$ be a continuous differentiable vector point function specified in a vector field, then the divergence of $\mathbf{V}$ is defined as :

$$\mathbf{i} \cdot \frac{\partial \mathbf{V}}{\partial x} + \mathbf{j} \cdot \frac{\partial \mathbf{V}}{\partial y} + \mathbf{k} \cdot \frac{\partial \mathbf{V}}{\partial z}$$

and is written as $\nabla \cdot \mathbf{V}$ or $\text{div } \mathbf{V}$ and read as divergence $\mathbf{V}$.

$$\begin{aligned}
 \therefore \nabla \cdot \mathbf{V} &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (V_1\mathbf{i} + V_2\mathbf{j} + V_3\mathbf{k}) \\
 &= \frac{\partial V_1}{\partial x} + \frac{\partial V_2}{\partial y} + \frac{\partial V_3}{\partial z}
 \end{aligned}$$

(Agra,

which is clearly a scalar quantity.

Note, If $\nabla \cdot \mathbf{V} = 0$ then $\mathbf{V}$ is known as *Solenoidal Vector*.

Interpretation of $\text{div } \mathbf{V}$. Let us consider the motion of a homogenous incompressible fluid particle at an instant of time t and let u, v, w be the components of its velocity parallel to axes of x, y, z respectively.

Construct an elementary rectangular parallelepiped of fluid with sides $\delta x, \delta y, \delta z$ having $P(x, y, z)$ at the centre.

The velocity vector at P is given by

$$\mathbf{V} = u\mathbf{i} + v\mathbf{j} + w\mathbf{k}.$$

Now, components of velocity being the functions of x, y, z , we have

$$u = u(x, y, z), v = v(x, y, z), w = w(x, y, z)$$

the component of velocity parallel to x -axis at a point on the face $A'D'C'B'$ will be

$$= u \left(x + \frac{\delta x}{2}, y, z \right)$$

$$= u(x, y, z) + \frac{\delta x}{2} \left[\frac{\partial u}{\partial x} \right] + \dots \text{ (by Taylor's Theorem).}$$

The mass of the fluid passing out of the face $A'D'C'B'$ in small time δt = density of the fluid $\times$ velocity normal to the face $A'D'C'B'$ $\times$ area of face $A'D'C'B'$ $\times$ time

$$= \rho \left[u + \frac{\delta x}{2} \left(\frac{\partial u}{\partial x} \right) \right] \times \delta y \times \delta z \times \delta t$$

Similarly the mass of fluid that passes through the face $ADCB$

$$= \rho \left[u - \frac{\delta x}{2} \frac{\partial u}{\partial x} \right] \delta y \delta z \delta t.$$

And the mass of the fluid that passes out through the faces $ADCB$ and $A'D'C'B'$

$$\begin{aligned}
 &= \rho \left[u + \frac{\delta x}{2} \frac{\partial u}{\partial x} \right] \delta y \delta z \delta t - \rho \left[u - \frac{\delta x}{2} \frac{\partial u}{\partial x} \right] \delta y \delta z \delta t \\
 &= \rho \frac{\partial u}{\partial x} \delta x \delta y \delta z \delta t.
 \end{aligned}$$

If the other two pairs of faces are considered then we have masses of fluid that moves out as

$$\rho \frac{\partial v}{\partial y} \delta x \delta y \delta z \delta t \text{ and } \rho \frac{\partial w}{\partial z} \delta x \delta y \delta z \delta t.$$




Fig. 1.38

The total mass of fluid that moves out the elementary parallelepiped in time δt

$$= \rho \frac{\partial u}{\partial x} \delta x \delta y \delta z \delta t + \rho \frac{\partial v}{\partial y} \delta x \delta y \delta z \delta t + \rho \frac{\partial w}{\partial z} \delta x \delta y \delta z \delta t$$

$$= \rho \left[\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right] \delta x \delta y \delta z \delta t.$$

The volume of the elementary parallelepiped is $\delta x \delta y \delta z$ and proceeding to the limit when $\delta x, \delta y, \delta z, \delta t$ all tend to zero we find that an amount of fluid mass per unit volume per unit time passes through the point $P(x, y, z)$

$$= \rho \frac{\left[\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right] \delta x \delta y \delta z \delta t}{\delta x \delta y \delta z \delta t}$$

$$= \rho \left[\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right]$$

$$= \rho \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (u\mathbf{i} + v\mathbf{j} + w\mathbf{k})$$

$$= \rho \nabla \cdot \mathbf{V} = \rho \operatorname{div} \mathbf{V}.$$

1.29. THE DIVERGENCE OF SUM OF TWO VECTOR FUNCTIONS

If $\mathbf{U}$ and $\mathbf{V}$ be two vector-point functions expressed as

$$\mathbf{U} = U_1\mathbf{i} + U_2\mathbf{j} + U_3\mathbf{k}$$

$$\mathbf{V} = V_1\mathbf{i} + V_2\mathbf{j} + V_3\mathbf{k}.$$

Then

$$\nabla \cdot (\mathbf{U} + \mathbf{V}) = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot [(U_1 + V_1)\mathbf{i} + (U_2 + V_2)\mathbf{j} + (U_3 + V_3)\mathbf{k}]$$

$$= \frac{\partial}{\partial x} [U_1 + V_1] + \frac{\partial}{\partial y} [U_2 + V_2] + \frac{\partial}{\partial z} [U_3 + V_3]$$

$$= \left(\frac{\partial U_1}{\partial x} + \frac{\partial U_2}{\partial y} + \frac{\partial U_3}{\partial z} \right) + \left(\frac{\partial V_1}{\partial x} + \frac{\partial V_2}{\partial y} + \frac{\partial V_3}{\partial z} \right)$$

$$= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (U_1\mathbf{i} + U_2\mathbf{j} + U_3\mathbf{k})$$

$$+ \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (V_1\mathbf{i} + V_2\mathbf{j} + V_3\mathbf{k})$$

$$= \nabla \cdot \mathbf{U} + \nabla \cdot \mathbf{V}$$

$$= \operatorname{div} \mathbf{U} + \operatorname{div} \mathbf{V}.$$

Showing that the divergence of the sum of two vector functions is equal to the sum of their divergences.

This rule may be generalised for any number of vector functions.

1.30. THE DIVERGENCE OF PRODUCT

If the vector point function $\mathbf{U}$ is expressed as

$$\mathbf{U} = U_1\mathbf{i} + U_2\mathbf{j} + U_3\mathbf{k} \quad \text{and} \quad V \text{ is a scalar point-function.}$$

$$\text{Then } \nabla \cdot (\mathbf{U}V) = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot [(U_1\mathbf{i} + U_2\mathbf{j} + U_3\mathbf{k})V]$$

$$= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot [VU_1\mathbf{i} + VU_2\mathbf{j} + VU_3\mathbf{k}]$$

$$= \frac{\partial}{\partial x} (VU_1) + \frac{\partial}{\partial y} (VU_2) + \frac{\partial}{\partial z} (VU_3)$$

$$= U_1 \frac{\partial V}{\partial x} + V \frac{\partial U_1}{\partial x} + U_2 \frac{\partial V}{\partial y} + V \frac{\partial U_2}{\partial y} + U_3 \frac{\partial V}{\partial z} + V \frac{\partial U_3}{\partial z}$$

$$= \left(U_1 \frac{\partial V}{\partial x} + U_2 \frac{\partial V}{\partial y} + U_3 \frac{\partial V}{\partial z} \right) + V \left(\frac{\partial U_1}{\partial x} + \frac{\partial U_2}{\partial y} + \frac{\partial U_3}{\partial z} \right)$$

$$= \left(\mathbf{i} \frac{\partial V}{\partial x} + \mathbf{j} \frac{\partial V}{\partial y} + \mathbf{k} \frac{\partial V}{\partial z} \right) \cdot (U_1\mathbf{i} + U_2\mathbf{j} + U_3\mathbf{k})$$

$$+ V \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (U_1\mathbf{i} + U_2\mathbf{j} + U_3\mathbf{k})$$

$$= (\nabla V) \cdot \mathbf{U} + V (\nabla \cdot \mathbf{U})$$

$$\operatorname{div} (\mathbf{U}V) = (\operatorname{grad} V) \cdot \mathbf{U} + V \operatorname{div} \mathbf{U}.$$

Problem 84. (a) If $\mathbf{V} = x^2z\mathbf{i} - 2y^2z^2\mathbf{j} + xy^2z\mathbf{k}$

(Agra, 1961, 63)

Find $\nabla \cdot \mathbf{V}$ at the point $(1, -1, 1)$.

(b) If $\mathbf{V} = \frac{x\mathbf{i} + y\mathbf{j}}{x+y}$ find $\nabla \cdot \mathbf{V}$

(c) If $\mathbf{V} = x \cos z\mathbf{i} + y \log x\mathbf{j} - z^2\mathbf{k}$ evaluate $\nabla \cdot \mathbf{V}$

(d) Given $\mathbf{V} = x^2z\mathbf{i} - 2y^2z^2\mathbf{j} + xy^2z\mathbf{k}$.

$$\nabla \cdot \mathbf{V} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (x^2z\mathbf{i} - 2y^2z^2\mathbf{j} + xy^2z\mathbf{k})$$

$$= \frac{\partial}{\partial x} (x^2z) + \frac{\partial}{\partial y} (-2y^2z^2) + \frac{\partial}{\partial z} (xy^2z)$$

$$= 2xz - 6y^2z + xy^2$$

$$= 2 - 6 + 1 = -3 \text{ at } x=1, y=-1, z=1.$$

(e) Given $\mathbf{V} = \frac{x\mathbf{i} + y\mathbf{j}}{x+y}$.

$$\nabla \cdot \mathbf{V} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot \left(\frac{x}{x+y}\mathbf{i} + \frac{y}{x+y}\mathbf{j} \right)$$

$$\begin{aligned}
 &= \frac{\partial}{\partial x} \left(\frac{x}{x+y} \right) + \frac{\partial}{\partial y} \left(\frac{y}{x+y} \right) \\
 &= \frac{(x+y) - x}{(x+y)^2} + \frac{(x+y) - y}{(x+y)^2} \\
 &= \frac{x+y}{(x+y)^2} \\
 &= \frac{1}{x+y}
 \end{aligned}$$

(c) Given $\mathbf{V} = x \cos z \mathbf{i} + y \log x \mathbf{j} - z^2 \mathbf{k}$

$$\begin{aligned}
 \therefore \nabla \cdot \mathbf{V} &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (x \cos z \mathbf{i} + y \log x \mathbf{j} - z^2 \mathbf{k}) \\
 &= \frac{\partial}{\partial x} (x \cos z) + \frac{\partial}{\partial y} (y \log x) + \frac{\partial}{\partial z} (-z^2) \\
 &= \cos z + \log x - 2z.
 \end{aligned}$$

Problem 85. If $\phi = 2x^3y^2z^4$ then find

(a) $\text{div}(\text{grad } \phi)$, and

(b) Show that $\nabla \cdot \nabla \phi = \nabla^2 \phi$ where $\nabla^2 \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$

(a) Given $\phi = 2x^3y^2z^4$.

$$\begin{aligned}
 \therefore \nabla \phi &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (2x^3y^2z^4) \\
 &= (6x^2y^2z^4) \mathbf{i} + (4x^3y^2z^4) \mathbf{j} + (8x^3y^2z^3) \mathbf{k}
 \end{aligned}$$

So that

$$\begin{aligned}
 \nabla \cdot \nabla \phi &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (6x^2y^2z^4 \mathbf{i} + 4x^3y^2z^4 \mathbf{j} + 8x^3y^2z^3 \mathbf{k}) \\
 &= \frac{\partial}{\partial x} (6x^2y^2z^4) + \frac{\partial}{\partial y} (4x^3y^2z^4) + \frac{\partial}{\partial z} (8x^3y^2z^3) \\
 &= 12xy^2z^4 + 4x^3z^4 + 24x^3y^2z^2.
 \end{aligned}$$

$$\begin{aligned}
 (b) \quad \nabla \cdot \nabla \phi &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot \left(\mathbf{i} \frac{\partial \phi}{\partial x} + \mathbf{j} \frac{\partial \phi}{\partial y} + \mathbf{k} \frac{\partial \phi}{\partial z} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{\partial \phi}{\partial x} \right) + \frac{\partial}{\partial y} \left(\frac{\partial \phi}{\partial y} \right) + \frac{\partial}{\partial z} \left(\frac{\partial \phi}{\partial z} \right) \\
 &= \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} + \frac{\partial^2 \phi}{\partial z^2} \\
 &= \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \phi = \nabla^2 \phi
 \end{aligned}$$

Note. $\nabla^2 \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ is known as *Laplacian operator*.

Problem 86. If $u = 3x^2y$, $v = xz^2 - 2y$ then evaluate $\text{grad}[(\text{grad } v)]$.

At given

$$\begin{aligned}
 \nabla u &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (3x^2y) \\
 &= 6xy \mathbf{i} + 3x^2 \mathbf{j}
 \end{aligned}$$

and

$$\begin{aligned}
 \nabla v &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (xz^2 - 2y) \\
 &= z^2 \mathbf{i} - 2 \mathbf{j} + 2xz \mathbf{k}.
 \end{aligned}$$

$$\begin{aligned}
 \nabla u \cdot \nabla v &= (6xy \mathbf{i} + 3x^2 \mathbf{j}) \cdot (z^2 \mathbf{i} - 2 \mathbf{j} + 2xz \mathbf{k}) \\
 &= 6xyz^2 - 2x^2.
 \end{aligned}$$

$$\begin{aligned}
 \nabla(\text{grad } u \cdot \text{grad } v) &= \left[\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right] (6xyz^2 - 2x^2) \\
 &= (6yz^2 - 4x) \mathbf{i} + (6xz^2) \mathbf{j} + (12xyz) \mathbf{k}.
 \end{aligned}$$

Problem 87. Prove $\nabla^2(uv) = u \nabla^2 v + 2 \nabla u \cdot \nabla v + v \nabla^2 u$.

We have

$$\nabla^2(uv) = \nabla \cdot [\nabla(uv)]$$

where,

$$\nabla(uv) = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (uv)$$

$$= \mathbf{i} \frac{\partial}{\partial x} (uv) + \mathbf{j} \frac{\partial}{\partial y} (uv) + \mathbf{k} \frac{\partial}{\partial z} (uv)$$

$$\begin{aligned}
 &= \left[u \frac{\partial v}{\partial x} + v \frac{\partial u}{\partial x} \right] \mathbf{i} + \left[u \frac{\partial v}{\partial y} + v \frac{\partial u}{\partial y} \right] \mathbf{j} \\
 &\quad + \left[u \frac{\partial v}{\partial z} + v \frac{\partial u}{\partial z} \right] \mathbf{k}
 \end{aligned}$$

$$\begin{aligned}
 &= \left[\frac{\partial v}{\partial x} \mathbf{i} + \frac{\partial v}{\partial y} \mathbf{j} + \frac{\partial v}{\partial z} \mathbf{k} \right] + v \left[\frac{\partial u}{\partial x} \mathbf{i} + \frac{\partial u}{\partial y} \mathbf{j} + \frac{\partial u}{\partial z} \mathbf{k} \right] \\
 &= u \nabla v + v \nabla u.
 \end{aligned}$$

$$\nabla^2(uv) = \nabla \cdot (u \nabla v + v \nabla u)$$

$$\begin{aligned}
 &= \nabla \cdot (u \nabla v) + \nabla \cdot (v \nabla u) \\
 &= u \nabla \cdot (\nabla v) + \nabla v \cdot \nabla u + v \nabla \cdot (\nabla u) + \nabla u \cdot \nabla v \\
 &= u \nabla^2 v + \nabla v \cdot \nabla u + v \nabla^2 u + \nabla u \cdot \nabla v \\
 &= u \nabla^2 v + 2 \nabla u \cdot \nabla v + v \nabla^2 u.
 \end{aligned}$$

Problem 88. Prove

$$\nabla \cdot (u \nabla v - v \nabla u) = u \nabla^2 v - v \nabla^2 u.$$

We have, $\nabla \cdot (u \nabla v) = u [\nabla \cdot (\nabla v)] + \nabla v \cdot \nabla u$

$$= u \nabla^2 v + \nabla v \cdot \nabla u \quad \dots (1)$$

and similarly $\nabla \cdot (v \nabla u) = v \nabla^2 u + \nabla u \cdot \nabla v$.

$$\dots (2)$$

Subtracting (2) from (1), we get

$$\begin{aligned}
 \nabla \cdot (u \nabla v - v \nabla u) &= u \nabla^2 v + \nabla v \cdot \nabla u - v \nabla^2 u - \nabla u \cdot \nabla v \\
 &= u \nabla^2 v - v \nabla^2 u.
 \end{aligned}$$

Problem 89. Prove that

$$\nabla^2 \left(\frac{1}{r} \right) = 0$$

$$r^2 = x^2 + y^2 + z^2$$

(Agra, 1957)

We have $\nabla^2\left(\frac{1}{r}\right) = \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}\right) \left[\frac{1}{(x^2+y^2+z^2)^{1/2}}\right]$

But
$$\frac{\partial}{\partial x} \left[\frac{1}{(x^2+y^2+z^2)^{1/2}} \right] = -\frac{x}{(x^2+y^2+z^2)^{3/2}}$$

$$\frac{\partial}{\partial x^2} \left[\frac{1}{(x^2+y^2+z^2)^{1/2}} \right] = \frac{\partial}{\partial x} \left[\frac{-x}{(x^2+y^2+z^2)^{3/2}} \right]$$

$$= \frac{3x^2}{(x^2+y^2+z^2)^{5/2}} - \frac{1}{(x^2+y^2+z^2)^{3/2}}$$

and similarly

$$\frac{\partial^2}{\partial y^2} \left[\frac{1}{(x^2+y^2+z^2)^{1/2}} \right] = \frac{3y^2}{(x^2+y^2+z^2)^{5/2}} - \frac{1}{(x^2+y^2+z^2)^{3/2}}$$

$$\frac{\partial^2}{\partial z^2} \left[\frac{1}{(x^2+y^2+z^2)^{1/2}} \right] = \frac{3z^2}{(x^2+y^2+z^2)^{5/2}} - \frac{1}{(x^2+y^2+z^2)^{3/2}}$$

$$\therefore \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \left[\frac{1}{(x^2+y^2+z^2)^{1/2}} \right]$$

$$= \frac{3(x^2+y^2+z^2)}{(x^2+y^2+z^2)^{5/2}} - \frac{3}{(x^2+y^2+z^2)^{3/2}}$$

$$= \frac{3}{(x^2+y^2+z^2)^{3/2}} - \frac{3}{(x^2+y^2+z^2)^{3/2}}$$

$$= 0$$

i.e. $\nabla^2\left(\frac{1}{r}\right) = 0.$

Note. The equation $\nabla^2\phi=0$ is known as *Laplace's equation*, and hence $\phi = \frac{1}{r}$ is a solution of this equation.

Problem 90. Prove the following propositions:

(a) $\mathbf{V} = 3y^4z^2\mathbf{i} + 4x^3z^2\mathbf{j} - 3x^2y^2\mathbf{k}$ is a solenoidal vector.

(b) $\mathbf{V} = (x+3y)\mathbf{i} + (y-2z)\mathbf{j} + (x+az)\mathbf{k}$ is a solenoidal vector when $a = -2$.

(c) $\mathbf{U} = (2x^2+8xy^2z)\mathbf{i} + (3x^3y-3xy)\mathbf{j} - (4y^2z^2+2x^3z)\mathbf{k}$ is not solenoidal vector.

But vector $\mathbf{V} = xyz^2\mathbf{U}$ is solenoidal.

By note on § 1.28, we know that if the divergence of a vector then the vector is called *solenoidal*.

(a) Given $\mathbf{V} = 3y^4z^2\mathbf{i} + 4x^3z^2\mathbf{j} - 3x^2y^2\mathbf{k}$

$$\therefore \nabla \cdot \mathbf{V} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (3y^4z^2\mathbf{i} + 4x^3z^2\mathbf{j} - 3x^2y^2\mathbf{k})$$

$$= \frac{\partial}{\partial x} (y^4z^2) + \frac{\partial}{\partial y} (x^3z^2) + \frac{\partial}{\partial z} (-3x^2y^2)$$

$$= 0.$$

Hence the given vector $\mathbf{V}$ is solenoidal.

(b) Given $\mathbf{V} = (x+3y)\mathbf{i} + (y-2z)\mathbf{j} + (x+az)\mathbf{k}$

$$\therefore \nabla \cdot \mathbf{V} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot [(x+3y)\mathbf{i} + (y-2z)\mathbf{j} + (x+az)\mathbf{k}]$$

$$= \frac{\partial}{\partial x} [(x+3y)] + \frac{\partial}{\partial y} [(y-2z)] + \frac{\partial}{\partial z} [(x+az)]$$

$$= 1 + 1 + a.$$

In order that $\mathbf{V}$ be solenoidal vector, we should have $\nabla \cdot \mathbf{V} = 0$

i.e. $a+2=0$ or $a=-2$.

(c) Given $\mathbf{U} = (2x^2+8y^2zx)\mathbf{i} + (3x^3y-3xy)\mathbf{j} - (4y^2z^2+2x^3z)\mathbf{k}$

$$\therefore \nabla \cdot \mathbf{U} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot [(2x^2+8y^2zx)\mathbf{i} + (3x^3y-3xy)\mathbf{j} - (4y^2z^2+2x^3z)\mathbf{k}]$$

$$= \frac{\partial}{\partial x} (2x^2+8y^2zx) + \frac{\partial}{\partial y} (3x^3y-3xy) + \frac{\partial}{\partial z} (-4y^2z^2-2x^3z)$$

$$= 4x + 8y^2z + 3x^3 - 3x + 8y^2z + 2x^3 \neq 0.$$

Hence $\mathbf{U}$ is not a solenoidal vector.

But $xyz^2\mathbf{U} = (2x^3yz^2+8x^2y^3z^2)\mathbf{i} + (3x^4y^2z^2-3x^2y^2z^2)\mathbf{j}$

$$- (4xy^3z^4+2x^4yz^3)\mathbf{k}$$

$$\therefore \nabla \cdot (xyz^2\mathbf{U}) = \frac{\partial}{\partial x} (2x^3yz^2+8x^2y^3z^2) + \frac{\partial}{\partial y} (3x^4y^2z^2-3x^2y^2z^2)$$

$$- \frac{\partial}{\partial z} (4xy^3z^4+2x^4yz^3)\mathbf{k}$$

$$= 6x^2yz^2 + 16xy^3z^2 + 6x^4yz^2 - 6x^2yz^2$$

$$- 16xy^3z^3 - 6x^4yz^2$$

$$= 0.$$

Hence $xyz^2\mathbf{U}$ is a solenoidal vector.

1.31. THE CURL OR ROTATION OF A VECTOR POINT FUNCTION

(Agra, 1952, 61, 63)

Let $\mathbf{f}(x, y, z) = f_1\mathbf{i} + f_2\mathbf{j} + f_3\mathbf{k}$ be a continuous differentiable vector-point function; then the curl of $\mathbf{f}$ or rotation of $\mathbf{f}$ is given by

$$\left(\mathbf{i} \times \frac{\partial \mathbf{f}}{\partial x} + \mathbf{j} \times \frac{\partial \mathbf{f}}{\partial y} + \mathbf{k} \times \frac{\partial \mathbf{f}}{\partial z} \right)$$

and is written as $\text{curl } \mathbf{f}$ or $\nabla \times \mathbf{f}$ or $\text{rot } \mathbf{f}$.

$$\text{i.e., } \nabla \times \mathbf{f} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times (f_1\mathbf{i} + f_2\mathbf{j} + f_3\mathbf{k})$$

$$= \left(\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} \right) \mathbf{i} + \left(\frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} \right) \mathbf{j} + \left(\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) \mathbf{k}$$

$$= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ f_1 & f_2 & f_3 \end{vmatrix} \quad (\text{Agra, 1967})$$

It is clear that **curl f** or **rotation f** is a vector quantity and read **del cross f**.

Note. If $\text{curl } f = 0$, f is known as *Irrotational Vector*.

Interpretation of the curl f. If a rigid body is in motion, the curl of its linear velocity at any point gives twice its angular velocity.

(Agra, 1954, 72 ; Rohilkhand, 1954)

Consider the motion of a rigid body rotating with angular velocity ω about an axis OA ; O , being a fixed point of the body. Let r be the position vector of a point P of the body. Draw PQ perpendicular from P to the axis OA . Then,




Fig. 1.39

Linear velocity V of P due to circular motion $V = |\mathbf{V}| = \omega QP = \omega r \sin \theta = |\omega \times \mathbf{r}|$

i.e., $\mathbf{V} = \omega \times \mathbf{r}$
where, $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$

and $\omega = \omega_1\mathbf{i} + \omega_2\mathbf{j} + \omega_3\mathbf{k}$

But we know that $\text{curl } \mathbf{V} = \nabla \times \mathbf{V} = \nabla \times (\omega \times \mathbf{r})$

$$\begin{aligned} &= \nabla \times \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \omega_1 & \omega_2 & \omega_3 \\ x & y & z \end{vmatrix} \\ &= \nabla \times [(\omega_2 z - \omega_3 y)\mathbf{i} + (\omega_3 x - \omega_1 z)\mathbf{j} + (\omega_1 y - \omega_2 x)\mathbf{k}] \\ &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \omega_2 z - \omega_3 y & \omega_3 x - \omega_1 z & \omega_1 y - \omega_2 x \end{vmatrix} \\ &= 2[\omega_1\mathbf{i} + \omega_2\mathbf{j} + \omega_3\mathbf{k}] \\ &= 2\omega. \quad \text{Which proves the proposition.} \end{aligned}$$

1.32. CURL OF THE SUM OF TWO VECTOR-POINT FUNCTIONS

If $\mathbf{u}$ and $\mathbf{v}$ be two vector-point functions given by

$$\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}$$

$$\mathbf{v} = v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}$$

$$\nabla \times (\mathbf{u} + \mathbf{v}) = \nabla \times [(u_1 + v_1)\mathbf{i} + (u_2 + v_2)\mathbf{j} + (u_3 + v_3)\mathbf{k}]$$

$$\begin{aligned} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u_1 + v_1 & u_2 + v_2 & u_3 + v_3 \end{vmatrix} \\ &= \mathbf{i} \left[\frac{\partial}{\partial y} (u_3 + v_3) - \frac{\partial}{\partial z} (u_2 + v_2) \right] + \mathbf{j} \left[\frac{\partial}{\partial z} (u_1 + v_1) - \frac{\partial}{\partial x} (u_3 + v_3) \right] \\ &\quad + \mathbf{k} \left[\frac{\partial}{\partial x} (u_2 + v_2) - \frac{\partial}{\partial y} (u_1 + v_1) \right] \\ &= \mathbf{i} \left[\frac{\partial u_3}{\partial y} - \frac{\partial u_2}{\partial z} \right] + \mathbf{j} \left[\frac{\partial u_1}{\partial z} - \frac{\partial u_3}{\partial x} \right] + \mathbf{k} \left[\frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} \right] \\ &\quad + \mathbf{i} \left[\frac{\partial v_3}{\partial y} - \frac{\partial v_2}{\partial z} \right] + \mathbf{j} \left[\frac{\partial v_1}{\partial z} - \frac{\partial v_3}{\partial x} \right] + \mathbf{k} \left[\frac{\partial v_2}{\partial x} - \frac{\partial v_1}{\partial y} \right] \\ &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u_1 & u_2 & u_3 \end{vmatrix} + \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix} \quad (\text{by the definition}) \end{aligned}$$

$$\nabla \times (\mathbf{u} + \mathbf{v}) = \nabla \times \mathbf{u} + \nabla \times \mathbf{v}.$$

$$\text{curl } (\mathbf{u} + \mathbf{v}) = \text{curl } \mathbf{u} + \text{curl } \mathbf{v}.$$

Thus curl of sum of two vector point functions is equal to the sum of their curls.

This result may be generalized for any number of vector point functions.

Ex. If $\mathbf{r}$ is the position vector of a variable point with respect to fixed origin such that $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ then $\text{curl } \mathbf{r} = 0$. (Agra, 1967)

$$\begin{aligned} \text{curl } \mathbf{r} &= \left(\mathbf{i} \times \frac{\partial}{\partial x} + \mathbf{j} \times \frac{\partial}{\partial y} + \mathbf{k} \times \frac{\partial}{\partial z} \right) (x\mathbf{i} + y\mathbf{j} + z\mathbf{k}) \\ &= \mathbf{i} \left[\frac{\partial}{\partial y} (z) - \frac{\partial}{\partial z} (y) \right] + \mathbf{j} \left[\frac{\partial}{\partial z} (x) - \frac{\partial}{\partial x} (z) \right] \\ &\quad + \mathbf{k} \left[\frac{\partial}{\partial x} (y) - \frac{\partial}{\partial y} (x) \right] \\ &= 0. \end{aligned}$$

1.33. CURL OF THE PRODUCT OF TWO VECTOR-POINT FUNCTIONS

We have to consider the curl of the forms uv and $\mathbf{u} \times \mathbf{v}$ where u is a scalar and $\mathbf{u}, \mathbf{v}$ vector point functions.

$$\text{Suppose, } \mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}$$

$$\mathbf{v} = v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}$$

and u is a scalar point function.

$$\text{Then, } \text{curl}(uv) = \nabla \times (uv) = \left(\mathbf{i} \times \frac{\partial}{\partial x} + \mathbf{j} \times \frac{\partial}{\partial y} + \mathbf{k} \times \frac{\partial}{\partial z} \right) (uv_1\mathbf{i} + uv_2\mathbf{j} + uv_3\mathbf{k})$$

$$= \nabla \times (uv_1\mathbf{i} + uv_2\mathbf{j} + uv_3\mathbf{k})$$

$$= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ uv_1 & uv_2 & uv_3 \end{vmatrix}$$

$$= \left[\frac{\partial}{\partial y}(uv_3) - \frac{\partial}{\partial z}(uv_2) \right] \mathbf{i} + \left[\frac{\partial}{\partial z}(uv_1) - \frac{\partial}{\partial x}(uv_3) \right] \mathbf{j} + \left[\frac{\partial}{\partial x}(uv_2) - \frac{\partial}{\partial y}(uv_1) \right] \mathbf{k}$$

$$= \left[u \frac{\partial v_3}{\partial y} + v_3 \frac{\partial u}{\partial y} - u \frac{\partial v_2}{\partial z} - v_2 \frac{\partial u}{\partial z} \right] \mathbf{i} + \left[u \frac{\partial v_1}{\partial z} + v_1 \frac{\partial u}{\partial z} - u \frac{\partial v_3}{\partial x} - v_3 \frac{\partial u}{\partial x} \right] \mathbf{j} + \left[u \frac{\partial v_2}{\partial x} + v_2 \frac{\partial u}{\partial x} - u \frac{\partial v_1}{\partial y} - v_1 \frac{\partial u}{\partial y} \right] \mathbf{k}$$

$$= u \left[\left(\frac{\partial v_3}{\partial y} - \frac{\partial v_2}{\partial z} \right) \mathbf{i} + \left(\frac{\partial v_1}{\partial z} - \frac{\partial v_3}{\partial x} \right) \mathbf{j} + \left(\frac{\partial v_2}{\partial x} - \frac{\partial v_1}{\partial y} \right) \mathbf{k} \right] + \left[\left(\frac{\partial u}{\partial y} v_3 - \frac{\partial u}{\partial z} v_2 \right) \mathbf{i} + \left(\frac{\partial u}{\partial z} v_1 - \frac{\partial u}{\partial x} v_3 \right) \mathbf{j} + \left(\frac{\partial u}{\partial x} v_2 - \frac{\partial u}{\partial y} v_1 \right) \mathbf{k} \right]$$

$$= u \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix} + \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} & \frac{\partial u}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix}$$

$$= u \text{curl } \mathbf{v} + (\text{grad } u) \times \mathbf{v}$$

$$\text{i.e. } \nabla \times (uv) = u \nabla \times \mathbf{v} + (\nabla u) \times \mathbf{v}. \quad (\text{Agra, 1955, 57, 6})$$

$$\text{Again } \text{curl}(\mathbf{u} \times \mathbf{v}) = \nabla \times (\mathbf{u} \times \mathbf{v})$$

$$= \nabla \times \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

$$= \nabla \times [(v_3u_2 - v_2u_3)\mathbf{i} + (v_1u_3 - v_3u_1)\mathbf{j} + (v_2u_1 - v_1u_2)\mathbf{k}]$$

$$= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ (v_3u_2 - v_2u_3) & (v_1u_3 - v_3u_1) & (v_2u_1 - v_1u_2) \end{vmatrix}$$

$$= \mathbf{i} \left[\frac{\partial}{\partial y} (v_2u_1 - v_1u_2) - \frac{\partial}{\partial z} (v_1u_3 - v_3u_1) \right]$$

$$+ \mathbf{j} \left[\frac{\partial}{\partial z} (v_3u_2 - v_2u_3) - \frac{\partial}{\partial x} (v_2u_1 - v_1u_2) \right]$$

$$+ \mathbf{k} \left[\frac{\partial}{\partial x} (v_1u_3 - v_3u_1) - \frac{\partial}{\partial y} (v_3u_2 - v_2u_3) \right]$$

$$= \mathbf{i} \left[u_1 \frac{\partial v_3}{\partial y} + v_3 \frac{\partial u_1}{\partial y} - u_2 \frac{\partial v_1}{\partial y} - v_1 \frac{\partial u_2}{\partial y} - u_3 \frac{\partial v_1}{\partial z} - v_1 \frac{\partial u_3}{\partial z} + u_1 \frac{\partial v_3}{\partial z} + v_3 \frac{\partial u_1}{\partial z} \right]$$

$$+ \mathbf{j} \left[u_2 \frac{\partial v_3}{\partial z} + v_3 \frac{\partial u_2}{\partial z} - u_3 \frac{\partial v_2}{\partial z} - v_2 \frac{\partial u_3}{\partial z} - u_1 \frac{\partial v_2}{\partial x} - v_2 \frac{\partial u_1}{\partial x} + u_1 \frac{\partial v_2}{\partial x} + v_2 \frac{\partial u_1}{\partial x} \right]$$

$$+ \mathbf{k} \left[u_3 \frac{\partial v_1}{\partial x} + v_1 \frac{\partial u_3}{\partial x} - v_3 \frac{\partial u_1}{\partial x} - u_1 \frac{\partial v_3}{\partial x} - u_2 \frac{\partial v_3}{\partial y} - v_3 \frac{\partial u_2}{\partial y} + v_2 \frac{\partial u_3}{\partial y} + u_3 \frac{\partial v_2}{\partial y} \right]$$

$$= (u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}) \left[\left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}) \right]$$

$$- (v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}) \left[\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right] \cdot (u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k})$$

$$+ \left[(v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}) \cdot \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \right] (u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k})$$

$$- \left[(u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}) \cdot \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \right] (v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k})$$

$$= u \text{div } \mathbf{v} - \mathbf{v} \text{div } \mathbf{u} + (\mathbf{v} \cdot \nabla) \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{v}$$

$$\text{Hence } \text{Curl } \mathbf{f} = \mathbf{i} \times \frac{\partial \mathbf{f}}{\partial x} + \mathbf{j} \times \frac{\partial \mathbf{f}}{\partial y} + \mathbf{k} \times \frac{\partial \mathbf{f}}{\partial z} = \Sigma \mathbf{i} \times \frac{\partial \mathbf{f}}{\partial x}$$

$$\text{Now } \text{Curl}(\mathbf{u} \times \mathbf{v}) = \Sigma \mathbf{i} \times \frac{\partial [\mathbf{u} \times \mathbf{v}]}{\partial x}$$

$$= \Sigma \mathbf{i} \times \left[\mathbf{u} \times \frac{\partial \mathbf{v}}{\partial x} + \mathbf{v} \times \frac{\partial \mathbf{u}}{\partial x} \right]$$

$$= \Sigma \mathbf{i} \times \left(\mathbf{u} \times \frac{\partial \mathbf{v}}{\partial x} \right) + \Sigma \mathbf{i} \times \left(\mathbf{v} \times \frac{\partial \mathbf{u}}{\partial x} \right)$$

$$= \Sigma \left(\mathbf{i} \cdot \frac{\partial \mathbf{v}}{\partial x} \right) \mathbf{u} - \Sigma (\mathbf{i} \cdot \mathbf{u}) \frac{\partial \mathbf{v}}{\partial x} + \Sigma (\mathbf{i} \cdot \mathbf{v}) \frac{\partial \mathbf{u}}{\partial x} - \Sigma \left(\mathbf{i} \cdot \frac{\partial \mathbf{u}}{\partial x} \right) \mathbf{v}$$

$$= \left(\Sigma \mathbf{i} \cdot \frac{\partial \mathbf{v}}{\partial x} \right) \mathbf{u} - (\Sigma \mathbf{i} \cdot \mathbf{u}) \frac{\partial \mathbf{v}}{\partial x} + [\Sigma (\mathbf{i} \cdot \mathbf{v})] \frac{\partial \mathbf{u}}{\partial x} - \left[\Sigma \left(\mathbf{i} \cdot \frac{\partial \mathbf{u}}{\partial x} \right) \right] \mathbf{v}$$

$$= u \text{div } \mathbf{v} - \mathbf{v} \text{div } \mathbf{u} + (\mathbf{v} \cdot \nabla) \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{v}$$

1.14. TO EXPRESS GRADIENT OF SCALAR PRODUCT IN TERMS OF CURL

We have to show that

$$\text{grad}(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \times \text{curl } \mathbf{v} + \mathbf{v} \times \text{curl } \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{u}$$

We know that

$$\text{grad}(\mathbf{u} \cdot \mathbf{v}) = \Sigma \mathbf{i} \frac{\partial}{\partial x} (\mathbf{u} \cdot \mathbf{v})$$

$$\begin{aligned}
 &= \Sigma \mathbf{i} \left[\mathbf{u} \cdot \frac{\partial \mathbf{v}}{\partial x} + \mathbf{v} \cdot \frac{\partial \mathbf{u}}{\partial x} \right] \\
 &= \Sigma \mathbf{i} \left(\mathbf{u} \cdot \frac{\partial \mathbf{v}}{\partial x} \right) + \Sigma \mathbf{i} \left(\mathbf{v} \cdot \frac{\partial \mathbf{u}}{\partial x} \right) \quad \dots (1)
 \end{aligned}$$

And $\mathbf{u} \times \left(\mathbf{i} \times \frac{\partial \mathbf{v}}{\partial x} \right) = \left(\mathbf{u} \cdot \frac{\partial \mathbf{v}}{\partial x} \right) \mathbf{i} - (\mathbf{u} \cdot \mathbf{i}) \frac{\partial \mathbf{v}}{\partial x}$

or $\left(\mathbf{u} \cdot \frac{\partial \mathbf{v}}{\partial x} \right) \mathbf{i} = \left[\mathbf{u} \times \left(\mathbf{i} \times \frac{\partial \mathbf{v}}{\partial x} \right) \right] + (\mathbf{u} \cdot \mathbf{i}) \frac{\partial \mathbf{v}}{\partial x}$

$$\begin{aligned}
 \therefore \Sigma \left(\mathbf{u} \cdot \frac{\partial \mathbf{v}}{\partial x} \right) \mathbf{i} &= \Sigma \left[\mathbf{u} \times \left(\mathbf{i} \times \frac{\partial \mathbf{v}}{\partial x} \right) \right] + \Sigma (\mathbf{u} \cdot \mathbf{i}) \frac{\partial \mathbf{v}}{\partial x} \\
 &= \mathbf{u} \times \text{curl } \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{v} \quad \dots (2)
 \end{aligned}$$

Similarly $\Sigma \left(\mathbf{v} \cdot \frac{\partial \mathbf{u}}{\partial x} \right) \mathbf{i} = \mathbf{v} \times \text{curl } \mathbf{u} + (\mathbf{v} \cdot \nabla) \mathbf{u} \quad \dots (3)$

Substituting values of (2) and (3) in (1) we find,

$$\begin{aligned}
 \text{grad } (\mathbf{u} \cdot \mathbf{v}) &= \mathbf{u} \times \text{curl } \mathbf{v} + (\mathbf{u} \cdot \nabla) \mathbf{v} + \mathbf{v} \times \text{curl } \mathbf{u} + (\mathbf{v} \cdot \nabla) \mathbf{u} \\
 &= \mathbf{u} \times \text{curl } \mathbf{v} + \mathbf{v} \times \text{curl } \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{u}.
 \end{aligned}$$

1.35. TO EXPRESS DIVERGENCE OF VECTOR PRODUCT IN TERMS OF CURL

We have to show that $\text{div } (\mathbf{u} \times \mathbf{v}) = \text{curl } \mathbf{u} \cdot \mathbf{v} - \text{curl } \mathbf{v} \cdot \mathbf{u}$.

(Agra, 1971, 64, 61, 50)

We know that,

$$\begin{aligned}
 \text{div } (\mathbf{u} \times \mathbf{v}) &= \Sigma \mathbf{i} \cdot \frac{\partial}{\partial x} (\mathbf{u} \times \mathbf{v}) \\
 &= \Sigma \mathbf{i} \cdot \left[\frac{\partial \mathbf{u}}{\partial x} \times \mathbf{v} + \mathbf{u} \times \frac{\partial \mathbf{v}}{\partial x} \right] \\
 &= \left[\Sigma \mathbf{i} \cdot \left(\frac{\partial \mathbf{u}}{\partial x} \times \mathbf{v} \right) + \Sigma \mathbf{i} \cdot \left(\mathbf{u} \times \frac{\partial \mathbf{v}}{\partial x} \right) \right] \\
 &= \Sigma \mathbf{i} \times \frac{\partial \mathbf{u}}{\partial x} \cdot \mathbf{v} + \Sigma \mathbf{i} \cdot \mathbf{u} \times \frac{\partial \mathbf{v}}{\partial x} \\
 &\quad \text{(interchanging dot and cross)} \\
 &= \left(\Sigma \mathbf{i} \times \frac{\partial \mathbf{u}}{\partial x} \right) \cdot \mathbf{v} + \left(\Sigma \mathbf{i} \times \frac{\partial \mathbf{v}}{\partial x} \right) \cdot \mathbf{u} \\
 &= \text{curl } \mathbf{u} \cdot \mathbf{v} - \text{curl } \mathbf{v} \cdot \mathbf{u}.
 \end{aligned}$$

Problem 91. Prove the following :

(i) $\text{div grad } \phi = \nabla^2 \phi$.

(ii) $\text{curl grad } \phi = \nabla \times (\nabla \phi) = 0$ (Agra, 1956, 67)

(iii) $\text{div curl } \mathbf{f} = \nabla \cdot (\nabla \times \mathbf{f}) = 0$ (Agra, 1956, 67, 71)

(iv) $\text{curl curl } \mathbf{f} = \nabla \times (\nabla \times \mathbf{f}) = \text{grad div } \mathbf{f} - \nabla^2 \mathbf{f}$ (Agra, 1959, 61, 64, 69)

(v) $\text{grad div } \mathbf{V} = \nabla (\nabla \cdot \mathbf{V}) = \text{curl curl } \mathbf{V} + \nabla^2 \mathbf{V}$.

$$\begin{aligned}
 \text{(i) } \text{div grad } \phi &= \nabla \cdot (\nabla \phi) \\
 &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot \left(\mathbf{i} \frac{\partial \phi}{\partial x} + \mathbf{j} \frac{\partial \phi}{\partial y} + \mathbf{k} \frac{\partial \phi}{\partial z} \right) \\
 &= \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} + \frac{\partial^2 \phi}{\partial z^2} \\
 &= \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \phi \\
 &= \nabla^2 \phi
 \end{aligned}$$

Note. This follows that

$$\text{div grad } \equiv \left\{ \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right\} \equiv \nabla^2 \cdot \text{i.e., Laplacian operator.}$$

$$\begin{aligned}
 \text{(ii) } \text{curl grad } \phi &= \nabla \times (\nabla \phi) \\
 &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times \left(\mathbf{i} \frac{\partial \phi}{\partial x} + \mathbf{j} \frac{\partial \phi}{\partial y} + \mathbf{k} \frac{\partial \phi}{\partial z} \right) \\
 &= \left(\frac{\partial^2 \phi}{\partial y \partial z} - \frac{\partial^2 \phi}{\partial z \partial y} \right) \mathbf{i} + \left(\frac{\partial^2 \phi}{\partial z \partial x} - \frac{\partial^2 \phi}{\partial x \partial z} \right) \mathbf{j} \\
 &\quad + \left(\frac{\partial^2 \phi}{\partial x \partial y} - \frac{\partial^2 \phi}{\partial y \partial x} \right) \mathbf{k} \\
 &= 0
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) } \text{div curl } \mathbf{f} &= \nabla \cdot (\nabla \times \mathbf{f}). \\
 &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot \left(\mathbf{i} \times \frac{\partial \mathbf{f}}{\partial x} + \mathbf{j} \times \frac{\partial \mathbf{f}}{\partial y} + \mathbf{k} \times \frac{\partial \mathbf{f}}{\partial z} \right) \\
 &\quad \text{(Taking } \mathbf{f} = f_1 \mathbf{i} + f_2 \mathbf{j} + f_3 \mathbf{k}) \\
 &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot \left[\left(\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} \right) \mathbf{i} \right. \\
 &\quad \left. + \left(\frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} \right) \mathbf{j} + \left(\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) \mathbf{k} \right] \\
 &= \frac{\partial}{\partial x} \left[\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} \right] + \frac{\partial}{\partial y} \left[\frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} \right] + \frac{\partial}{\partial z} \left[\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right] \\
 &= \frac{\partial^2 f_3}{\partial x \partial y} - \frac{\partial^2 f_2}{\partial x \partial z} + \frac{\partial^2 f_1}{\partial y \partial z} - \frac{\partial^2 f_3}{\partial y \partial x} + \frac{\partial^2 f_2}{\partial z \partial x} - \frac{\partial^2 f_1}{\partial z \partial y} \\
 &= 0.
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv) } \text{curl curl } \mathbf{f} &= \nabla \times (\nabla \times \mathbf{f}) \\
 &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times \left(\mathbf{i} \times \frac{\partial \mathbf{f}}{\partial x} + \mathbf{j} \times \frac{\partial \mathbf{f}}{\partial y} + \mathbf{k} \times \frac{\partial \mathbf{f}}{\partial z} \right) \\
 &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times \left[\left(\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} \right) \mathbf{i} \right. \\
 &\quad \left. + \left(\frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} \right) \mathbf{j} + \left(\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) \mathbf{k} \right]
 \end{aligned}$$

$$\begin{aligned}
&= \left[\frac{\partial}{\partial y} \left(\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) - \frac{\partial}{\partial z} \left(\frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} \right) \right] \mathbf{i} \\
&\quad + \left[\frac{\partial}{\partial z} \left(\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} \right) - \frac{\partial}{\partial x} \left(\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) \right] \mathbf{j} \\
&\quad + \left[\frac{\partial}{\partial x} \left(\frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} \right) - \frac{\partial}{\partial y} \left(\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} \right) \right] \mathbf{k} \\
&= \left[\left(\frac{\partial^2 f_2}{\partial y \partial x} + \frac{\partial^2 f_3}{\partial z \partial x} \right) \mathbf{i} - \left(\frac{\partial^2 f_1}{\partial y^2} + \frac{\partial^2 f_1}{\partial z^2} \right) \mathbf{i} \right] \\
&\quad + \left[\left(\frac{\partial^2 f_3}{\partial z \partial y} + \frac{\partial^2 f_1}{\partial x \partial y} \right) \mathbf{j} - \left(\frac{\partial^2 f_2}{\partial x^2} + \frac{\partial^2 f_2}{\partial y^2} \right) \mathbf{j} \right] \\
&\quad + \left[\left(\frac{\partial^2 f_1}{\partial x \partial z} + \frac{\partial^2 f_2}{\partial y \partial z} \right) \mathbf{k} - \left(\frac{\partial^2 f_3}{\partial x^2} + \frac{\partial^2 f_3}{\partial y^2} \right) \mathbf{k} \right]
\end{aligned}$$

(On adding and subtracting $\frac{\partial^2 f_1}{\partial x^2}$, $\frac{\partial^2 f_2}{\partial y^2}$, $\frac{\partial^2 f_3}{\partial z^2}$ respectively in each factor of each bracket)

$$= \left[\left(\frac{\partial^2 f_1}{\partial x^2} + \frac{\partial^2 f_2}{\partial x \partial y} + \frac{\partial^2 f_3}{\partial x \partial z} \right) \mathbf{i} - \left(\frac{\partial^2 f_1}{\partial x^2} + \frac{\partial^2 f_1}{\partial y^2} + \frac{\partial^2 f_1}{\partial z^2} \right) \mathbf{i} \right] + \text{two similar terms}$$

$$= \Sigma \mathbf{i} \frac{\partial}{\partial x} \left(\frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y} + \frac{\partial f_3}{\partial z} \right) - \Sigma \left(\frac{\partial^2 f_1}{\partial x^2} + \frac{\partial^2 f_1}{\partial y^2} + \frac{\partial^2 f_1}{\partial z^2} \right) \mathbf{i} \\
= \text{grad div } \mathbf{f} - \nabla^2 \mathbf{f}$$

Note. If $\text{div } \mathbf{f} = 0$, then $\text{grad div } \mathbf{f} = 0$ and so $\text{curl curl } \mathbf{f} = -\nabla^2 \mathbf{f}$.

(Agra, 1980)

(v) $\text{grad div } \mathbf{V} = \nabla (\nabla \cdot \mathbf{V})$.

$$= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \left(\mathbf{i} \cdot \frac{\partial \mathbf{V}}{\partial x} + \mathbf{j} \cdot \frac{\partial \mathbf{V}}{\partial y} + \mathbf{k} \cdot \frac{\partial \mathbf{V}}{\partial z} \right)$$

(on taking $\mathbf{V} = V_1 \mathbf{i} + V_2 \mathbf{j} + V_3 \mathbf{k}$)

$$= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \left(\frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y} + \frac{\partial f_3}{\partial z} \right)$$

$$= \left(\frac{\partial^2 f_1}{\partial x^2} + \frac{\partial^2 f_2}{\partial x \partial y} + \frac{\partial^2 f_3}{\partial x \partial z} \right) \mathbf{i} + \left(\frac{\partial^2 f_1}{\partial y \partial x} + \frac{\partial^2 f_2}{\partial y^2} + \frac{\partial^2 f_3}{\partial y \partial z} \right) \mathbf{j} \\
+ \left(\frac{\partial^2 f_1}{\partial z \partial x} + \frac{\partial^2 f_2}{\partial z \partial y} + \frac{\partial^2 f_3}{\partial z^2} \right) \mathbf{k}$$

$$= \left[\mathbf{i} \cdot \frac{\partial^2 \mathbf{f}}{\partial x^2} + \mathbf{i} \cdot \frac{\partial^2 \mathbf{f}}{\partial x \partial y} + \mathbf{i} \cdot \frac{\partial^2 \mathbf{f}}{\partial x \partial z} \right] + \text{two similar terms}$$

$$= \Sigma \left(\mathbf{i} \cdot \frac{\partial^2 \mathbf{f}}{\partial x^2} + \mathbf{i} \cdot \frac{\partial^2 \mathbf{f}}{\partial x \partial y} + \mathbf{i} \cdot \frac{\partial^2 \mathbf{f}}{\partial x \partial z} \right)$$

$$= \Sigma \frac{\partial^2 \mathbf{f}}{\partial x^2} + \nabla \times (\mathbf{V} \times \mathbf{f}) = \nabla^2 \mathbf{f} + \text{curl curl } \mathbf{f}.$$

Problem 92. Find curl of the following functions

(a) $\mathbf{f} = \frac{x\mathbf{i} + y\mathbf{j}}{x+y}$

(b) $\mathbf{f} = x \cos z \mathbf{i} + y \log x \mathbf{j} - z^2 \mathbf{k}$

(c) $\mathbf{f} = x^2 \mathbf{i} + y^2 \mathbf{j} + z^2 \mathbf{k}$

(d) $\mathbf{f} = f_1 \mathbf{i} + f_2 \mathbf{j} + f_3 \mathbf{k}$; then

$$\begin{aligned}
\text{curl } \mathbf{f} &= \mathbf{i} \times \frac{\partial \mathbf{f}}{\partial x} + \mathbf{j} \times \frac{\partial \mathbf{f}}{\partial y} + \mathbf{k} \times \frac{\partial \mathbf{f}}{\partial z} \\
&= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times \mathbf{f} \\
&= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times (f_1 \mathbf{i} + f_2 \mathbf{j} + f_3 \mathbf{k}) \\
&= \left(\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} \right) \mathbf{i} + \left(\frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} \right) \mathbf{j} + \left(\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) \mathbf{k}
\end{aligned}$$

(a) We have $\mathbf{f} = \frac{x}{x+y} \mathbf{i} + \frac{y}{x+y} \mathbf{j} + 0 \mathbf{k}$

$$f_1 = \frac{x}{x+y}, f_2 = \frac{y}{x+y}, f_3 = 0$$

$$\frac{\partial f_3}{\partial y} = 0; \frac{\partial f_2}{\partial z} = 0; \frac{\partial f_1}{\partial z} = 0; \frac{\partial f_3}{\partial x} = 0$$

$$\frac{\partial f_2}{\partial x} = -\frac{y}{(x+y)^2}; \frac{\partial f_1}{\partial y} = -\frac{x}{(x+y)^2}$$

$$\begin{aligned}
\text{curl } \mathbf{f} &= 0 \mathbf{i} + 0 \mathbf{j} + \left[-\frac{y}{(x+y)^2} + \frac{x}{(x+y)^2} \right] \mathbf{k} \\
&= \frac{x-y}{(x+y)^2} \mathbf{k}
\end{aligned}$$

(b) We have $\mathbf{f} = x \cos z \mathbf{i} + y \log x \mathbf{j} - z^2 \mathbf{k}$

$$f_1 = x \cos z, f_2 = y \log x, f_3 = -z^2,$$

so that $\frac{\partial f_3}{\partial y} = 0; \frac{\partial f_2}{\partial z} = 0; \frac{\partial f_1}{\partial z} = -x \sin z;$

$$\frac{\partial f_3}{\partial x} = 0; \frac{\partial f_2}{\partial x} = \frac{y}{x}; \frac{\partial f_1}{\partial y} = 0$$

$$\text{curl } \mathbf{f} = 0 \mathbf{i} - x \sin z \mathbf{j} + \frac{y}{x} \mathbf{k}.$$

(c) We have $\mathbf{f} = x^2 \mathbf{i} + y^2 \mathbf{j} + z^2 \mathbf{k}$

$$f_1 = x^2, f_2 = y^2, f_3 = z^2.$$

so that $\frac{\partial f_3}{\partial y} = 0, \frac{\partial f_2}{\partial z} = 0, \frac{\partial f_1}{\partial z} = 0$

$$\frac{\partial f_3}{\partial x} = 0, \frac{\partial f_2}{\partial x} = 0, \frac{\partial f_1}{\partial y} = 0.$$

$$\therefore \text{curl } \mathbf{f} = 0\mathbf{i} + 0\mathbf{j} + 0\mathbf{k} = \mathbf{0}.$$

Problem 93. If $u = x^2yz$, $v = xy - 3z^2$ find

(a) $\nabla \cdot [(\nabla u) \cdot (\nabla v)]$

(b) $\nabla \cdot [(\nabla u) \times (\nabla v)]$

(c) $\nabla \times [(\nabla u \times \nabla v)]$.

Given $u = x^2yz$.

$$\therefore \nabla u = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (x^2yz)$$

$$= (2xyz) \mathbf{i} + (x^2z) \mathbf{j} + (x^2y) \mathbf{k}$$

and

$$v = xy - 3z^2$$

$$\therefore \nabla v = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (xy - 3z^2)$$

$$= y\mathbf{i} + x\mathbf{j} - 6z\mathbf{k}.$$

So that $(\nabla u) \cdot (\nabla v) = [(2xyz) \mathbf{i} + (x^2z) \mathbf{j} + (x^2y) \mathbf{k}] \cdot [y\mathbf{i} + x\mathbf{j} - 6z\mathbf{k}]$

$$= 2xy^2z + x^3z - 6x^2yz$$

and $[(\nabla u) \times (\nabla v)] = (2xyz\mathbf{i} + x^3z\mathbf{j} + x^2y\mathbf{k}) \times (y\mathbf{i} + x\mathbf{j} - 6z\mathbf{k})$

$$= (-6x^2z^2 - x^2y) \mathbf{i} + (x^2y^2 + 12xyz^2) \mathbf{j} + (2x^2yz - x^2yz) \mathbf{k}$$

Thus,

(a) $\nabla \cdot [\nabla u \cdot \nabla v] = \left[\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right] [2xy^2z + x^3z - 6x^2yz]$

$$= [2y^2z + 3x^2z - 12xyz] \mathbf{i} + [4xyz - 6x^2z] \mathbf{j} + [2xy^2 + x^3 - 6x^2y] \mathbf{k}$$

(b) $\nabla \cdot [\nabla u \times \nabla v]$

$$= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot [(-6x^2z^2 - x^2y) \mathbf{i} + (x^2y^2 + 12xyz^2) \mathbf{j} + (2x^2yz - x^2yz) \mathbf{k}]$$

$$= \frac{\partial}{\partial x} (-6x^2z^2 - x^2y) + \frac{\partial}{\partial y} (x^2y^2 + 12xyz^2) + \frac{\partial}{\partial z} (2x^2yz - x^2yz)$$

$$= [-12xz^2 - 3x^2y + 2x^2y + 12xz^2 + 2x^2y - x^2y]$$

$$= 0.$$

(c) $\nabla \times [\nabla u \times \nabla v]$

$$= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times [(-6x^2z^2 - x^2y) \mathbf{i} + (x^2y^2 + 12xyz^2) \mathbf{j} + (2x^2yz - x^2yz) \mathbf{k}]$$

$$= \left[\frac{\partial}{\partial y} (2x^2yz - x^2yz) - \frac{\partial}{\partial z} (x^2y^2 + 12xyz^2) \right] \mathbf{i}$$

$$+ \left[\frac{\partial}{\partial z} (6x^2z^2 - x^2y) - \frac{\partial}{\partial x} (2x^2yz - x^2yz) \right] \mathbf{j}$$

$$+ \left[\frac{\partial}{\partial x} (x^2y^2 + 12xyz^2) - \frac{\partial}{\partial y} (-6x^2z^2 - x^2y) \right] \mathbf{k}$$

$$= [2x^2y - x^2z - 24xyz] \mathbf{i} + [-12x^2z - 4xyz + 2xyz] \mathbf{j} + [2xy^2 + 12yz^2 + x^3] \mathbf{k}$$

$$= (x^2y - 24xyz) \mathbf{i} - (12x^2z + 2xyz) \mathbf{j} + (2xy^2 + x^3 + 12yz^2) \mathbf{k}.$$

Problem 94. If $\mathbf{v} = yz^2\mathbf{i} - 3xz^2\mathbf{j} + 2xyz\mathbf{k}$ and $\mathbf{v} = 3xi + 4zj - xyk$

and $\phi = xyz$ find,

(a) $\mathbf{u} \times (\nabla \phi)$

(b) $(\mathbf{u} \times \nabla) \phi$

(c) $(\nabla \times \mathbf{u}) \times \mathbf{v}$

(d) $\mathbf{v} \cdot \nabla \times \mathbf{u}$.

Given

$$\phi = xyz$$

$$\nabla \phi = \left[\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right] (xyz)$$

$$= yz\mathbf{i} + zx\mathbf{j} + xy\mathbf{k}.$$

Thus,

(a) $\mathbf{u} \times (\nabla \phi) = (yz^2\mathbf{i} - 3xz^2\mathbf{j} + 2xyz\mathbf{k}) \times (yz\mathbf{i} + zx\mathbf{j} + xy\mathbf{k})$

$$= [-3x^2yz^2 - 2x^2yz^2] \mathbf{i} + [2xy^2z^2 - xy^2z^2] \mathbf{j}$$

$$+ [xyz^3 + 3xyz^3] \mathbf{k}$$

$$= -5x^2yz^2\mathbf{i} + xy^2z^2\mathbf{j} + 4xyz^3\mathbf{k}.$$

(b) $\mathbf{u} \times \nabla = (yz^2\mathbf{i} - 3xz^2\mathbf{j} + 2xyz\mathbf{k}) \times \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right)$

$$= \left(-3xz^2 \frac{\partial}{\partial z} - 2xyz \frac{\partial}{\partial y} \right) \mathbf{i} + \left(2xyz \frac{\partial}{\partial x} - yz^2 \frac{\partial}{\partial z} \right) \mathbf{j}$$

$$+ \left(yz^2 \frac{\partial}{\partial y} + 3xz^2 \frac{\partial}{\partial x} \right) \mathbf{k}$$

So that $[(\mathbf{u} \times \nabla)] \phi = \left[\left(-3xz^2 \frac{\partial}{\partial z} - 2xyz \frac{\partial}{\partial y} \right) \mathbf{i} \right.$

$$+ \left(2xyz \frac{\partial}{\partial x} - yz^2 \frac{\partial}{\partial z} \right) \mathbf{j} + \left(yz^2 \frac{\partial}{\partial y} + 3xz^2 \frac{\partial}{\partial x} \right) \mathbf{k} \left. \right] (xyz)$$

$$= (-3x^2yz^2 - 2x^2yz^2) \mathbf{i} + (2xy^2z^2 - xy^2z^2) \mathbf{j}$$

$$+ (xyz^3 + 3xyz^3) \mathbf{k}$$

$$= -5x^2yz^2\mathbf{i} + xy^2z^2\mathbf{j} + 4xyz^3\mathbf{k}.$$

(c) $\nabla \times \mathbf{u} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times (yz^2\mathbf{i} - 3xz^2\mathbf{j} + 2xyz\mathbf{k})$

$$= \left[\frac{\partial}{\partial y} (2xyz) - \frac{\partial}{\partial z} (-3xz^2) \right] \mathbf{i}$$

$$+ \left[\frac{\partial}{\partial z} (yz^2) - \frac{\partial}{\partial x} (2xyz) \right] \mathbf{j} + \left[\frac{\partial}{\partial x} (-3xz^2) - \frac{\partial}{\partial y} (yz^2) \right] \mathbf{k}$$

$$= [2xz + 6xz] \mathbf{i} + [2yz - 2yz] \mathbf{j} + [-3z^2 - z^2] \mathbf{k}$$

$$= 8xz\mathbf{i} + 0\mathbf{j} - 4z^2\mathbf{k}.$$

(d) $(\nabla \times \mathbf{u}) \times \mathbf{v} = (8xz\mathbf{i} + 0\mathbf{j} - 4z^2\mathbf{k}) \times (3xi + 4zj - xyk)$

$$= 16xz^3\mathbf{i} + (-12xz^2 + 8x^2yz)\mathbf{j} + 32xz^2\mathbf{k}$$

$$= 16xz^3\mathbf{i} + (8x^2yz - 12xz^2)\mathbf{j} + 32xz^2\mathbf{k}.$$

$$(d) \quad \nabla \times \mathbf{u} = 8xzi + 0\mathbf{j} - 4z^2\mathbf{k} \text{ [by (c)]}$$

$$\therefore \mathbf{v} \cdot (\nabla \times \mathbf{u}) = (3xi + 4z\mathbf{j} - xy\mathbf{k}) \cdot (8xzi + 0\mathbf{j} - 4z^2\mathbf{k})$$

$$= 24x^2z + 4xyz^2.$$

Problem 95. Prove the following relations:

$$(a) \quad \nabla \cdot [\nabla u \times \nabla v] = 0.$$

$$(b) \quad \nabla \cdot [\mathbf{a} \times \mathbf{r}] = 0.$$

$$(c) \quad \nabla \times [\mathbf{a} \times \mathbf{r}] = 2\mathbf{a}$$

$$(a) \quad \text{Suppose that } \nabla u = \mathbf{U} \text{ and } \nabla v = \mathbf{V}.$$

$$\text{So that } \nabla \cdot (\nabla u \times \nabla v) = \nabla \cdot (\mathbf{U} \times \mathbf{V})$$

$$= \mathbf{V} \cdot (\nabla \times \mathbf{U}) - \mathbf{U} \cdot (\nabla \times \mathbf{V})$$

$$= \nabla v \cdot (\nabla \times \nabla u) - \nabla u \cdot (\nabla \times \nabla v)$$

$$= 0 \quad (\because \text{curl grad } u = 0 \text{ etc.})$$

$$(b) \quad \text{If } \mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k} \text{ and } \mathbf{r} = xi + y\mathbf{j} + z\mathbf{k}$$

$$\text{Then } \nabla \cdot [\mathbf{a} \times \mathbf{r}] = \left[\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right] \cdot [a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}] \times (xi + y\mathbf{j} + z\mathbf{k})$$

$$= \left[\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right] \cdot [(a_2z - a_3y)\mathbf{i} + (a_3x - a_1z)\mathbf{j} + (a_1y - a_2x)\mathbf{k}]$$

$$= \frac{\partial}{\partial x} (a_2z - a_3y) + \frac{\partial}{\partial y} (a_3x - a_1z) + \frac{\partial}{\partial z} (a_1y - a_2x)$$

$$= 0.$$

$$(c) \quad \text{We have,}$$

$$\nabla \times (\mathbf{a} \times \mathbf{r}) = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times [(a_2z - a_3y)\mathbf{i} + (a_3x - a_1z)\mathbf{j} + (a_1y - a_2x)\mathbf{k}]$$

$$= \left[\frac{\partial}{\partial y} (a_1y - a_2x) - \frac{\partial}{\partial z} (a_3x - a_1z) \right] \mathbf{i}$$

$$+ \left[\frac{\partial}{\partial z} (a_2z - a_3y) - \frac{\partial}{\partial x} (a_1y - a_2x) \right] \mathbf{j}$$

$$+ \left[\frac{\partial}{\partial x} (a_3x - a_1z) - \frac{\partial}{\partial y} (a_1y - a_2x) \right] \mathbf{k}$$

$$= 2a_1\mathbf{i} + 2a_2\mathbf{j} + 2a_3\mathbf{k} = 2(a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}) = 2\mathbf{a}.$$

Problem 96. If $\hat{\mathbf{e}}$ is a unit vector, prove that

$$(a) \quad \nabla \cdot (\hat{\mathbf{e}} \cdot \mathbf{r}) \hat{\mathbf{e}} = 1. \quad (b) \quad \nabla \times (\hat{\mathbf{e}} \cdot \mathbf{r}) \hat{\mathbf{e}} = 0$$

$$(c) \quad \nabla \cdot [\hat{\mathbf{e}} \times \mathbf{r}] \times \hat{\mathbf{e}} = 2. \quad (d) \quad \nabla \times [(\hat{\mathbf{e}} \times \mathbf{r}) \times \hat{\mathbf{e}}] = 0.$$

Taking e_1, e_2, e_3 as the components of $\hat{\mathbf{e}}$ along principal axes,

$$\text{we have } \hat{\mathbf{e}} = e_1\mathbf{i} + e_2\mathbf{j} + e_3\mathbf{k} \text{ where } |\hat{\mathbf{e}}| = \sqrt{e_1^2 + e_2^2 + e_3^2} = 1 \quad \dots (1)$$

$$\text{and let } \mathbf{r} = xi + y\mathbf{j} + z\mathbf{k} \quad \dots (2)$$

(a) We have,

$$\nabla \cdot [(\hat{\mathbf{e}} \cdot \mathbf{r}) \hat{\mathbf{e}}] = \nabla \cdot [(e_1\mathbf{i} + e_2\mathbf{j} + e_3\mathbf{k}) \cdot (xi + y\mathbf{j} + z\mathbf{k})] (e_1\mathbf{i} + e_2\mathbf{j} + e_3\mathbf{k})$$

$$= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (e_1\mathbf{i} + e_2\mathbf{j} + e_3\mathbf{k}) (e_1x + e_2y + e_3z)$$

$$= \left(e_1 \frac{\partial}{\partial x} + e_2 \frac{\partial}{\partial y} + e_3 \frac{\partial}{\partial z} \right) (e_1x + e_2y + e_3z)$$

$$= e_1^2 + e_2^2 + e_3^2 = 1 \quad \text{by (1)}$$

(b) We have,

$$\nabla \times [(\hat{\mathbf{e}} \cdot \mathbf{r}) \hat{\mathbf{e}}] = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times (e_1\mathbf{i} + e_2\mathbf{j} + e_3\mathbf{k}) (e_1x + e_2y + e_3z)$$

$$\text{as in (a)}$$

$$= \mathbf{i} [e_2e_3 - e_3e_2] + \mathbf{j} [e_1e_3 - e_3e_1] + \mathbf{k} [e_1e_2 - e_2e_1]$$

$$= 0$$

(c) We have,

$$\nabla \cdot [(\hat{\mathbf{e}} \times \mathbf{r}) \times \hat{\mathbf{e}}] = \nabla \cdot [(e_1\mathbf{i} + e_2\mathbf{j} + e_3\mathbf{k}) \times (xi + y\mathbf{j} + z\mathbf{k})] \times (e_1\mathbf{i} + e_2\mathbf{j} + e_3\mathbf{k})$$

$$= \nabla \cdot [\{ (e_2z - e_3y)\mathbf{i} + (e_3x - e_1z)\mathbf{j} + (e_1y - e_2x)\mathbf{k} \} \times (e_1\mathbf{i} + e_2\mathbf{j} + e_3\mathbf{k})]$$

$$= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot [\mathbf{i} \{ e_3(e_3x - e_1z) - e_2(e_1y - e_2x) \}$$

$$+ \mathbf{j} \{ e_1(e_1y - e_2x) - e_3(e_2z - e_3y) \} + \mathbf{k} \{ e_2(e_2z - e_3y) - e_1(e_3x - e_1z) \}]$$

$$= (e_3^2 + e_2^2) + (e_1^2 + e_3^2) + (e_2^2 + e_1^2)$$

$$= 2(e_1^2 + e_2^2 + e_3^2) = 2 \quad \text{by (1)}$$

(d) We have

$$\nabla \times [(\hat{\mathbf{e}} \times \mathbf{r}) \times \hat{\mathbf{e}}] = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times [\mathbf{i} \{ e_3(e_3x - e_1z) - e_2(e_1y - e_2x) \}$$

$$+ \mathbf{j} \{ e_1(e_1y - e_2x) - e_3(e_2z - e_3y) \}$$

$$+ \mathbf{k} \{ e_2(e_2z - e_3y) - e_1(e_3x - e_1z) \}] \text{ as in (c)}$$

$$= \mathbf{i} [-e_2e_3 + e_3e_2] + \mathbf{j} [-e_3e_1 + e_1e_3] + \mathbf{k} [-e_1e_2 + e_2e_1]$$

$$= 0$$

Problem 97. If $r = (x^2 + y^2 + z^2)^{1/2}$ and $\left(\frac{1}{r}\right)$ is a solution of Laplace's

$$\text{equation show that } \nabla^2 \left(\frac{1}{r} \right) = 0.$$

(Agra, 1957)

Hence or otherwise, evaluate $\nabla \times \left(\frac{\mathbf{r}}{r^2} \right)$.

(a) Given $r = (x^2 + y^2 + z^2)^{1/2}$

and $\nabla^2 \equiv \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right)$

$$\therefore \nabla^2 \left(\frac{1}{r} \right) = \left(\frac{\partial}{\partial x^2} + \frac{\partial}{\partial y^2} + \frac{\partial}{\partial z^2} \right) \left[\frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right]$$

But $\frac{\partial}{\partial x} \left(\frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right) = \left[-\frac{x}{(x^2 + y^2 + z^2)^{3/2}} \right]$

$$\frac{\partial^2}{\partial x^2} \left[\frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right] = \frac{\partial}{\partial x} \left[\frac{-x}{(x^2 + y^2 + z^2)^{3/2}} \right]$$

$$= \frac{3x^2}{(x^2 + y^2 + z^2)^{5/2}} - \frac{1}{(x^2 + y^2 + z^2)^{3/2}}$$

Similarly $\frac{\partial^2}{\partial y^2} \left[\frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right] = \frac{3y^2}{(x^2 + y^2 + z^2)^{5/2}} - \frac{1}{(x^2 + y^2 + z^2)^{3/2}}$

$$\frac{\partial^2}{\partial z^2} \left[\frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right] = \frac{3z^2}{(x^2 + y^2 + z^2)^{5/2}} - \frac{1}{(x^2 + y^2 + z^2)^{3/2}}$$

$$\therefore \nabla^2 \left(\frac{1}{r} \right) = \frac{3(x^2 + y^2 + z^2)}{(x^2 + y^2 + z^2)^{5/2}} - \frac{3}{(x^2 + y^2 + z^2)^{3/2}} = \frac{3}{(x^2 + y^2 + z^2)^{3/2}} - \frac{3}{(x^2 + y^2 + z^2)^{3/2}} = 0.$$

(b) Suppose that $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$

and given $r = (x^2 + y^2 + z^2)^{1/2}$

$$\therefore \nabla \times \left(\frac{\mathbf{r}}{r^2} \right) = \nabla \times [(x^2 + y^2 + z^2)^{-1} (x\mathbf{i} + y\mathbf{j} + z\mathbf{k})]$$

$$= [\nabla (x^2 + y^2 + z^2)^{-1}] \times [x\mathbf{i} + y\mathbf{j} + z\mathbf{k}]$$

$$+ (x^2 + y^2 + z^2)^{-1} [\nabla \times (x\mathbf{i} + y\mathbf{j} + z\mathbf{k})]$$

$$= \left[\nabla \left(\frac{1}{x^2 + y^2 + z^2} \right) \right] \times (x\mathbf{i} + y\mathbf{j} + z\mathbf{k})$$

$$= \left[\left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \left(\frac{1}{(x^2 + y^2 + z^2)} \right) \right] \times [x\mathbf{i} + y\mathbf{j} + z\mathbf{k}]$$

$$= \frac{-2}{(x^2 + y^2 + z^2)^2} [x\mathbf{i} + y\mathbf{j} + z\mathbf{k}] \times [x\mathbf{i} + y\mathbf{j} + z\mathbf{k}]$$

$$= 0 \text{ [Vector product of two equal vectors being zero]}$$

Problem 98. If $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$, $\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$ and $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ then prove that $\text{grad} [(\mathbf{r} \times \mathbf{a}) \cdot (\mathbf{r} \times \mathbf{b})] = (\mathbf{b} \times \mathbf{r}) \times \mathbf{a} + (\mathbf{a} \times \mathbf{r}) \times \mathbf{b}$.

$$\text{grad} [(\mathbf{r} \times \mathbf{a}) \cdot (\mathbf{r} \times \mathbf{b})] = \text{grad} [r^2 (\mathbf{a} \cdot \mathbf{b}) - (\mathbf{a} \cdot \mathbf{r}) (\mathbf{b} \cdot \mathbf{r})]$$

$$= \text{grad} [r^2 (\mathbf{a} \cdot \mathbf{b})] - \text{grad} [(\mathbf{a} \cdot \mathbf{r}) (\mathbf{b} \cdot \mathbf{r})]$$

$$= (\mathbf{a} \cdot \mathbf{b}) \text{grad } r^2 + r^2 \text{grad } (\mathbf{a} \cdot \mathbf{b})$$

$$- (\mathbf{b} \cdot \mathbf{r}) \text{grad } (\mathbf{a} \cdot \mathbf{r}) - (\mathbf{a} \cdot \mathbf{r}) \text{grad } (\mathbf{b} \cdot \mathbf{r})$$

$$= (\mathbf{a} \cdot \mathbf{b}) 2\mathbf{r} - (\mathbf{b} \cdot \mathbf{r}) \mathbf{a} - (\mathbf{a} \cdot \mathbf{r}) \mathbf{b}$$

$$= (\mathbf{a} \cdot \mathbf{b}) \mathbf{r} - (\mathbf{a} \cdot \mathbf{r}) \mathbf{b} + (\mathbf{b} \cdot \mathbf{a}) \mathbf{r} - (\mathbf{b} \cdot \mathbf{r}) \mathbf{a}$$

$$= -\mathbf{a} \times (\mathbf{b} \times \mathbf{r}) - \mathbf{b} \times (\mathbf{a} \times \mathbf{r})$$

$$= (\mathbf{b} \times \mathbf{r}) \times \mathbf{a} + (\mathbf{a} \times \mathbf{r}) \times \mathbf{b}.$$

Problem 99. If $\mathbf{r}$ and r have their usual meaning show that

(a) $\text{div } r^n \mathbf{r} = (n+3) r^n$. (b) $\text{curl } r^n \mathbf{r} = 0$.

(Agra, 1974)

Suppose that $r^n \mathbf{r} = u\mathbf{V}$ then

$$\text{div } (r^n \mathbf{r}) = \text{div } (u\mathbf{V}) = u \text{div } \mathbf{V} + (\text{grad } u) \cdot \mathbf{V}$$

$$= r^n \text{div } \mathbf{r} + (\text{grad } r^n) \cdot \mathbf{r}$$

$$= r^n (3) + (nr^{n-2} \mathbf{r}) \cdot \mathbf{r}$$

$$= 3r^n + nr^{n-2} r^2 = (n+3) r^n.$$

(b) $\text{curl } (r^n \mathbf{r}) = \text{curl } (u\mathbf{V})$

$$= (\text{grad } u) \times \mathbf{V} + u \text{curl } \mathbf{V}$$

$$= (\text{grad } r^n) \times \mathbf{r} + r^n \text{curl } \mathbf{r}$$

$$= (nr^{n-2} \mathbf{r}) \times \mathbf{r} + 0 \quad \because \text{curl } \mathbf{r} = 0$$

$$= 0 \quad \text{as } \mathbf{r} \times \mathbf{r} = 0.$$

Problem 100. If $\mathbf{a} = \alpha x\mathbf{i} + \beta y\mathbf{j} + \gamma z\mathbf{k}$, show that

(a) $\nabla \cdot (\mathbf{a} \cdot \mathbf{r}) = 2\mathbf{a}$.

(b) If $\mathbf{r}$ and r have their usual meaning prove

$$\text{curl } (\mathbf{a} \cdot \mathbf{r}) \mathbf{a} = 0.$$

(c) $\mathbf{a} \cdot \mathbf{r} = (\alpha x\mathbf{i} + \beta y\mathbf{j} + \gamma z\mathbf{k}) \cdot (x\mathbf{i} + y\mathbf{j} + z\mathbf{k})$

$$= (\alpha x^2 + \beta y^2 + \gamma z^2)$$

$$\therefore \text{grad } (\mathbf{a} \cdot \mathbf{r}) = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (\alpha x^2 + \beta y^2 + \gamma z^2)$$

$$= (2\alpha x\mathbf{i} + 2\beta y\mathbf{j} + 2\gamma z\mathbf{k}) = 2\mathbf{a}.$$

(d) If $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$ and $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$

$$\mathbf{a} \cdot \mathbf{r} = (a_1x + a_2y + a_3z)$$

so that $(\mathbf{a} \cdot \mathbf{r}) \mathbf{a} = (a_1x + a_2y + a_3z) (a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k})$

$$= (a_1x + a_2y + a_3z) a_1\mathbf{i} + (a_1x + a_2y + a_3z) a_2\mathbf{j}$$

$$+ (a_1x + a_2y + a_3z) a_3\mathbf{k}$$

$$\therefore \text{grad } (\mathbf{a} \cdot \mathbf{r}) \mathbf{a} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times [(a_1x + a_2y + a_3z) a_1\mathbf{i} + \dots + \dots]$$

$$= \left\{ \frac{\partial}{\partial y} [(a_1x + a_2y + a_3z)a_3] - \frac{\partial}{\partial z} [(a_1x + a_2y + a_3z)a_2] \right\} i + \dots + \dots$$

$$= (a_2a_3 - a_3a_2) i + \dots + \dots$$

$$= 0.$$

Problem 101. If u, V, w are point functions and $uV = \nabla w$, prove that $V \cdot \text{curl } V = 0$.

We are given that

$$uV = \nabla w.$$

Taking curl of both sides we have

$$\text{curl } (uV) = \text{curl } (\nabla w) = 0$$

$$\text{or } (\text{grad } u) \times V + u \text{ curl } V = 0.$$

Multiplying by V scalarly, we find

$$V \cdot (\text{grad } u) \times V + V \cdot u \text{ curl } V = 0$$

$$\text{or } V \cdot u \text{ curl } V = 0 \text{ as } V \cdot (\text{grad } u) \times V = 0$$

$$\text{or } V \cdot \text{curl } V = 0 \text{ as } u \neq 0.$$

Problem 102. If a is a constant unit vector show that

$$a \cdot [\nabla (V \cdot a) - \nabla \times (V \times a)] = \nabla \cdot V = \text{div } V$$

We have $\text{grad } (A \cdot B) = (A \cdot \nabla) B + (B \cdot \nabla) A + A \times \text{curl } B + B \times \text{curl } A$

$$\nabla (V \cdot a) = (V \cdot \nabla) a + (a \cdot \nabla) V + V \times \text{curl } a + a \times \text{curl } V$$

$$= (V \cdot \nabla) a + (a \cdot \nabla) V + a \times \text{curl } V \quad \dots$$

But a being a constant vector $\text{curl } a = 0$.

$$\text{Also we have } \text{curl } (A \times B) = (B \cdot \nabla) A - (A \cdot \nabla) B + A \text{ div } B - B \text{ div } A$$

$$\text{So that } \nabla \times (V \times a) = (a \cdot \nabla) V - (V \cdot \nabla) a + V \text{ div } a - a \text{ div } V$$

$$= (a \cdot \nabla) V - (V \cdot \nabla) a - a \text{ div } V \quad \dots$$

$\therefore \text{div } a = 0$, a being constant

Subtraction of (2) from (1) yields

$$\nabla (V \cdot a) - \nabla \times (V \times a) = 2 (V \cdot \nabla) a + a \times \text{curl } V + a \text{ div } V$$

$$\text{Where } (V \cdot \nabla) a = \left(v_1 \frac{\partial}{\partial x} + v_2 \frac{\partial}{\partial y} + v_3 \frac{\partial}{\partial z} \right) a; \text{ if } V = v_1 i + v_2 j + v_3 k$$

$$= 0, a \text{ being constant}$$

$$\text{Hence, } a \cdot \{ \nabla (V \cdot a) - \nabla \times (V \times a) \} = a \cdot \{ a \times \text{curl } V + a \text{ div } V \}$$

$$= a \cdot a \times \text{curl } V + a \cdot a \text{ div } V$$

$$= \text{div } V \quad \therefore a \cdot a = 1, a \text{ being unit vector}$$

$$\text{And } a \cdot a \times \text{curl } V = a \times a \cdot \text{curl } V = 0$$

Problem 103. Prove that $\text{div } (u \text{ grad } v) = u \nabla^2 v + (\text{grad } u) \cdot (\text{grad } v)$ where u and v both are scalar point functions.

$$\text{We have } \text{div } (u \text{ grad } v) = \text{div } (uV) \text{ where } V = \text{grad } v$$

$$= u \text{ div } V + V \cdot \text{grad } u$$

$$= u \text{ div } (\text{grad } v) + (\text{grad } v) \cdot (\text{grad } u)$$

$$= u \nabla \cdot (\nabla v) + (\text{grad } v) \cdot (\text{grad } u)$$

$$= u \nabla^2 v + (\text{grad } v) \cdot (\text{grad } u).$$

Problem 104. Prove that $\text{curl } (u \text{ grad } v) = \nabla u \times \nabla v = -\text{curl } (v \text{ grad } u)$ where u and v are both scalar point functions.

We have

$$\text{curl } (u \text{ grad } v) = \text{curl } (uV) \text{ where } V = \text{grad } v$$

$$= u \text{ curl } V - V \times \text{grad } u$$

$$= u \text{ curl } (\text{grad } v) - \text{grad } v \times \text{grad } u$$

$$= 0 - \text{grad } v \times \text{grad } u.$$

$$= \text{grad } u \times \text{grad } v$$

$$= \nabla u \times \nabla v = -(\nabla v \times \nabla u) = -\text{curl } (v \text{ grad } u).$$

Problem 105. Prove that $\nabla \cdot (S \nabla \times A) = (\nabla \times A) \cdot (\nabla S)$.

$$\text{We have } \nabla \cdot (S \nabla \times A) = \text{div } (S \text{ curl } A)$$

$$= \text{div } (SV) \text{ where } \text{curl } A = V$$

$$= S \text{ div } V + (\text{grad } S) \cdot V$$

$$= S \text{ div } (\text{curl } A) + (\text{grad } S) \cdot (\text{curl } A)$$

$$= 0 + (\text{grad } S) \cdot (\text{curl } A)$$

$$= (\nabla \times A) \cdot (\nabla S).$$

Problem 106. Prove that

$$\nabla \cdot (\nabla u \times \nabla v) = \nabla u \cdot (\nabla \cdot \nabla v) - \nabla v \cdot (\nabla \cdot \nabla u) + (\nabla v \cdot \nabla) \nabla u - (\nabla u \cdot \nabla) \nabla v$$

where u and v both are scalar point functions.

Suppose that $\nabla u = A$ and $\nabla v = B$ then,

$$\nabla \cdot (\nabla u \times \nabla v) = \text{curl } (A \times B)$$

$$= A \text{ div } B - B \text{ div } A + (B \cdot \text{grad}) A - (A \cdot \text{grad}) B$$

$$= \nabla u \cdot (\nabla \cdot \nabla v) - \nabla v \cdot (\nabla \cdot \nabla u) + (\nabla v \cdot \nabla) \nabla u - (\nabla u \cdot \nabla) \nabla v.$$

Problem 107. Prove that the values of $\text{div } F$ and $\text{curl } F$ are independent of the choice of rectangular axes i.e. they are invariant.

(Lucknow 1952, 59)

Taking the mutual direction cosines of two sets of rectangular axes as shown in adjoining scheme and (x, y, z) , (x', y', z') as corresponding coordinates

We have $\mathbf{i}' = l_1 \mathbf{i} + m_1 \mathbf{j} + n_1 \mathbf{k}$
 $x' = l_1 x + m_1 y + n_1 z$ etc. } ... (1)

And $\mathbf{i} = l_1 \mathbf{i}' + l_2 \mathbf{j}' + l_3 \mathbf{k}'$
 $x = l_1 x' + l_2 y' + l_3 z'$ etc. } ... (2)

	x	y
	$\mathbf{i}$	$\mathbf{j}$
x'	l_1	m_1
y'	l_2	m_2
z'	l_3	m_3

Also $\frac{\partial}{\partial x} \equiv \frac{\partial}{\partial x'} \frac{\partial x'}{\partial x} + \frac{\partial}{\partial y'} \frac{\partial y'}{\partial x} + \frac{\partial}{\partial z'} \frac{\partial z'}{\partial x} = l_1 \frac{\partial}{\partial x'} + l_2 \frac{\partial}{\partial y'} + l_3 \frac{\partial}{\partial z'}$

Similarly, $\frac{\partial}{\partial y} \equiv m_1 \frac{\partial}{\partial x'} + m_2 \frac{\partial}{\partial y'} + m_3 \frac{\partial}{\partial z'}$; $\frac{\partial}{\partial z} \equiv n_1 \frac{\partial}{\partial x'} + n_2 \frac{\partial}{\partial y'} + n_3 \frac{\partial}{\partial z'}$

Using $l_1^2 + l_2^2 + l_3^2 = 1$ etc. and $l_1 m_1 + l_2 m_2 + l_3 m_3 = 0$ etc.,

We have $\mathbf{F} = F_1 \mathbf{i} + F_2 \mathbf{j} + F_3 \mathbf{k} = F_1 (l_1 \mathbf{i}' + l_2 \mathbf{j}' + l_3 \mathbf{k}') + \dots + \dots$
 $= (l_1 F_1 + m_1 F_2 + n_1 F_3) \mathbf{i}' + (l_2 F_1 + m_2 F_2 + n_2 F_3) \mathbf{j}'$
 $+ (l_3 F_1 + m_3 F_2 + n_3 F_3) \mathbf{k}'$

So that $(\text{div } \mathbf{F})_{x', y', z'} = \frac{\partial}{\partial x'} (l_1 F_1 + m_1 F_2 + n_1 F_3) + \frac{\partial}{\partial y'} (l_2 F_1 + m_2 F_2 + n_2 F_3) + \frac{\partial}{\partial z'} (l_3 F_1 + m_3 F_2 + n_3 F_3)$

$= \left(l_1 \frac{\partial}{\partial x'} + l_2 \frac{\partial}{\partial y'} + l_3 \frac{\partial}{\partial z'} \right) F_1 + \dots + \dots$

$= \frac{\partial}{\partial x} F_1 + \frac{\partial}{\partial y} F_2 + \frac{\partial}{\partial z} F_3 = (\text{div } \mathbf{F})_{x, y, z}$

Similarly $(\text{curl } \mathbf{F})_{x', y', z'} = (\text{curl } \mathbf{F})_{x, y, z}$

Which show the required invariance.

Problem 108. Find $\text{div grad } r^m$ and verify that $\nabla \times \nabla r^m = 0$.

If $r = |\mathbf{r}|$, $\mathbf{r}$ being position vector of a point, then

$\nabla r^m = m r^{m-2} \mathbf{r}$.

Also, $\text{div grad } r^m = \nabla \cdot (\nabla r^m)$

$= \nabla \cdot (m r^{m-2} \mathbf{r})$

$= m \nabla \cdot (r^{m-2} \mathbf{r})$

$= m [\nabla r^{m-2} \cdot \mathbf{r} + r^{m-2} \text{div } \mathbf{r}]$

$= m [(m-2) r^{m-4} \mathbf{r} \cdot \mathbf{r} + r^{m-2} \cdot 3] \quad \because \text{div } \mathbf{r} = 3$

$= m (m-2) r^{m-4} r^2 + 3m r^{m-2}$

$= m (m-2) r^{m-2} + 3m r^{m-2}$

$= m (m+1) r^{m-2}$.

$\text{curl grad } r^m = \nabla \times (m r^{m-2} \mathbf{r})$

$= \nabla \times [m r^{m-2} x \mathbf{i} + m r^{m-2} y \mathbf{j} + m r^{m-2} z \mathbf{k}]$

$= \left(\mathbf{i} \times \frac{\partial}{\partial x} + \mathbf{j} \times \frac{\partial}{\partial y} + \mathbf{k} \times \frac{\partial}{\partial z} \right)$

$[m r^{m-2} x \mathbf{i} + m r^{m-2} y \mathbf{j} + m r^{m-2} z \mathbf{k}]$

$= \left[\frac{\partial}{\partial y} (m r^{m-2} z) - \frac{\partial}{\partial z} (m r^{m-2} y) \right] \mathbf{i} + \dots + \dots$

$= \left[m (m-2) r^{m-3} z \frac{\partial r}{\partial y} \right.$

$\left. - m (m-2) r^{m-3} y \frac{\partial r}{\partial z} \right] \mathbf{i} + \dots + \dots$

$= \left[m (m-2) r^{m-3} z \frac{y}{r} \right.$

$\left. - m (m-2) r^{m-3} y \frac{z}{r} \right] \mathbf{i} + \dots + \dots$

$= 0 \mathbf{i} + 0 \mathbf{j} + 0 \mathbf{k} = 0$.

Problem 109. Show that $(\mathbf{V} \cdot \nabla) \mathbf{V} = \frac{1}{2} \nabla V^2 - \mathbf{V} \times \text{curl } \mathbf{V}$.

(Allahabad, 1958)

We have $\text{grad } (\mathbf{a} \cdot \mathbf{b}) = \mathbf{a} \times \text{curl } \mathbf{b} + \mathbf{b} \times \text{curl } \mathbf{a} + (\mathbf{a} \cdot \nabla) \mathbf{b} + (\mathbf{b} \cdot \nabla) \mathbf{a}$

Putting $\mathbf{a} = \mathbf{b} = \mathbf{V}$, this becomes

$\text{grad } (V^2) = \mathbf{V} \times \text{curl } \mathbf{V} + \mathbf{V} \times \text{curl } \mathbf{V} + (\mathbf{V} \cdot \nabla) \mathbf{V} + (\mathbf{V} \cdot \nabla) \mathbf{V}$

$\nabla V^2 = 2 \mathbf{V} \times \text{curl } \mathbf{V} + 2 (\mathbf{V} \cdot \nabla) \mathbf{V}$

$(\mathbf{V} \cdot \nabla) \mathbf{V} = \frac{1}{2} \nabla V^2 - \mathbf{V} \times \text{curl } \mathbf{V}$.

Problem 110. (a) Show that $\mathbf{E} = \frac{\mathbf{r}}{r^2}$ is irrotational.

(b) If $\mathbf{A}$ and $\mathbf{B}$ are irrotational, prove that $\mathbf{A} \times \mathbf{B}$ is solenoidal.

At given, $\mathbf{E} = \frac{\mathbf{r}}{r^2} = \frac{x \mathbf{i} + y \mathbf{j} + z \mathbf{k}}{x^2 + y^2 + z^2}$.

In order to show that $\mathbf{E}$ is irrotational, we have to prove that

$\nabla \times \left(\frac{\mathbf{r}}{r^2} \right) = 0$.

$\nabla \times \left(\frac{\mathbf{r}}{r^2} \right) = \nabla \times \left(\frac{x \mathbf{i} + y \mathbf{j} + z \mathbf{k}}{x^2 + y^2 + z^2} \right)$

$= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times \left[\frac{x}{x^2 + y^2 + z^2} \mathbf{i} \right.$
 $\left. + \frac{y}{x^2 + y^2 + z^2} \mathbf{j} + \frac{z}{x^2 + y^2 + z^2} \mathbf{k} \right]$

$$\begin{aligned}
&= \left[\frac{\partial}{\partial y} \left(\frac{z}{x^2+y^2+z^2} \right) - \frac{\partial}{\partial z} \left(\frac{y}{x^2+y^2+z^2} \right) \right] \mathbf{i} \\
&\quad + \left[\frac{\partial}{\partial z} \left(\frac{x}{x^2+y^2+z^2} \right) - \frac{\partial}{\partial x} \left(\frac{z}{x^2+y^2+z^2} \right) \right] \mathbf{j} \\
&\quad + \left[\frac{\partial}{\partial x} \left(\frac{y}{x^2+y^2+z^2} \right) - \frac{\partial}{\partial y} \left(\frac{x}{x^2+y^2+z^2} \right) \right] \mathbf{k} \\
&= \left[\frac{-2zy}{x^2+y^2+z^2} + \frac{2zy}{x^2+y^2+z^2} \right] \mathbf{i} + \dots + \dots \\
&= 0.
\end{aligned}$$

showing that $\mathbf{E}$ is an irrotational vector.

(b) If $\mathbf{A}$ and $\mathbf{B}$ are irrotational, then

$$\text{curl } \mathbf{A} = 0 \text{ and } \text{curl } \mathbf{B} = 0.$$

In order to prove that $\mathbf{A} \times \mathbf{B}$ is solenoidal, we have to show that

$$\text{div } (\mathbf{A} \times \mathbf{B}) = 0.$$

$$\text{Now } \text{div } (\mathbf{A} \times \mathbf{B}) = (\text{curl } \mathbf{A}) \cdot \mathbf{B} - (\text{curl } \mathbf{B}) \cdot \mathbf{A} = 0.$$

Hence $\mathbf{A} \times \mathbf{B}$ is solenoidal.

Problem 111. Show that $r^n \cdot \mathbf{r}$ is an irrotational vector for any value of n , but is a solenoidal only if $n = -3$. (Agra, 19)

$$\text{We have } \text{curl } r^n \mathbf{r} = r^n \text{curl } \mathbf{r} - \mathbf{r} \times \text{grad } r^n$$

$$= -\mathbf{r} \times (nr^{n-2} \mathbf{r}) \quad \text{as } \text{curl } \mathbf{r} = 0$$

$$= -\mathbf{r} \times \mathbf{r} (nr^{n-2}) = 0.$$

showing that $r^n \mathbf{r}$ is an irrotational vector for any value of n .

$$\text{Again } \text{div } r^n \mathbf{r} = r^n \text{div } \mathbf{r} + \mathbf{r} \cdot \text{grad } r^n$$

$$= 3r^n + \mathbf{r} \cdot (nr^{n-2} \mathbf{r})$$

$$= 3r^n + nr^{n-2} r^2 = (n+3)r^n$$

which is zero if $n+3=0$ or $n=-3$

This shows that $r^n \mathbf{r}$ is solenoidal when $n=-3$.

Problem 112. Prove that $\nabla \times (\nabla \times \mathbf{A}) = -\nabla^2 \mathbf{A} + \nabla (\nabla \cdot \mathbf{A})$.

Take $\nabla = \mathbf{R}$

$$\text{So that } \nabla \times (\nabla \times \mathbf{A}) = \mathbf{R} \times (\mathbf{R} \times \mathbf{A})$$

$$\text{But } \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b} (\mathbf{a} \cdot \mathbf{c}) - (\mathbf{a} \cdot \mathbf{b}) \mathbf{c}$$

$$\therefore \mathbf{R} \times (\mathbf{R} \times \mathbf{A}) = \mathbf{R} (\mathbf{R} \cdot \mathbf{A}) - (\mathbf{R} \cdot \mathbf{R}) \mathbf{A}$$

$$\text{or } \nabla \times (\nabla \times \mathbf{A}) = \nabla (\nabla \cdot \mathbf{A}) - (\nabla \cdot \nabla) \mathbf{A}$$

$$= \nabla (\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A}.$$

Problem 113. If $\nabla \cdot \mathbf{E} = 0$, $\nabla \cdot \mathbf{H} = 0$, $\nabla \times \mathbf{E} = -\frac{\partial \mathbf{H}}{\partial t}$, $\nabla \times \mathbf{H} = \frac{\partial \mathbf{E}}{\partial t}$,

show that $\mathbf{E}$ and $\mathbf{H}$ satisfy $\nabla^2 u = \frac{\partial^2 u}{\partial t^2}$.

At given,

$$\nabla \times (\nabla \times \mathbf{E}) = \nabla \times \left(-\frac{\partial \mathbf{H}}{\partial t} \right)$$

$$= -\frac{\partial}{\partial t} (\nabla \times \mathbf{H})$$

$$= -\frac{\partial}{\partial t} \left(\frac{\partial \mathbf{E}}{\partial t} \right) = -\frac{\partial^2 \mathbf{E}}{\partial t^2}.$$

But by Problem 112, we have

$$\nabla \times (\nabla \times \mathbf{E}) = -\nabla^2 \mathbf{E} + \nabla (\nabla \cdot \mathbf{E}) = -\nabla^2 \mathbf{E}$$

$$\text{So that } \nabla^2 \mathbf{E} = \frac{\partial^2 \mathbf{E}}{\partial t^2}.$$

$$\text{Similarly } \nabla \times (\nabla \times \mathbf{H}) = \nabla \times \left(\frac{\partial \mathbf{E}}{\partial t} \right)$$

$$= \frac{\partial}{\partial t} (\nabla \times \mathbf{E}) = \frac{\partial}{\partial t} \left(-\frac{\partial \mathbf{H}}{\partial t} \right) = -\frac{\partial^2 \mathbf{H}}{\partial t^2}.$$

$$\text{But } \nabla \times (\nabla \times \mathbf{H}) = -\nabla^2 \mathbf{H} + \nabla (\nabla \cdot \mathbf{H}) = -\nabla^2 \mathbf{H}$$

$$\text{So that } \nabla^2 \mathbf{H} = \frac{\partial^2 \mathbf{H}}{\partial t^2}.$$

$$\therefore \mathbf{E} \text{ and } \mathbf{H} \text{ satisfy the equation } \nabla^2 u = \frac{\partial^2 u}{\partial t^2}.$$

Problem 114. Show that the solution to the Maxwell's equations

$$\nabla \times \mathbf{H} = \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t}, \quad \nabla \times \mathbf{E} = \frac{1}{c} \frac{\partial \mathbf{H}}{\partial t}, \quad \nabla \cdot \mathbf{H} = 0, \quad \nabla \cdot \mathbf{E} = 4\pi\rho$$

where ρ is the charge density, x, y, z and c is the velocity of light assumed to be constant, are given by

$$\mathbf{E} = -\nabla\phi - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t}, \quad \mathbf{H} = \nabla \times \mathbf{A}$$

where $\mathbf{A}$ and ϕ called the vector and scalar potentials respectively, satisfy the equations

$$(1) \quad \nabla \cdot \mathbf{A} + \frac{1}{c} \frac{\partial \phi}{\partial t} = 0, \quad (2) \quad \nabla^2 \phi - \frac{1}{c^2} \frac{\partial^2 \phi}{\partial t^2} = 4\pi\rho,$$

$$(3) \quad \nabla^2 \mathbf{A} - \frac{1}{c^2} \frac{\partial^2 \mathbf{A}}{\partial t^2} = \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t}.$$

Maxwell's equations are given to be

$$\nabla \times \mathbf{H} = \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t}$$

$$\nabla \times \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{H}}{\partial t}$$

$$\nabla \cdot \mathbf{H} = 0$$

$$\text{and } \nabla \cdot \mathbf{E} = 4\pi\rho$$

We have to show that the solutions of these equations are given

$$\mathbf{E} = -\nabla\phi - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t}$$

$$\text{and } \mathbf{H} = \nabla \times \mathbf{A}$$

$$\text{where } \mathbf{A} \text{ and } \phi \text{ are given by } \nabla \cdot \mathbf{A} + \frac{1}{c} \frac{\partial \phi}{\partial t} = 0,$$

$$\nabla^2 \phi - \frac{1}{c^2} \frac{\partial^2 \phi}{\partial t^2} = 4\pi\rho,$$

$$\text{and } \nabla^2 \mathbf{A} = \frac{1}{c^2} \frac{\partial^2 \mathbf{A}}{\partial t^2}.$$

Putting $\mathbf{H} = \nabla \times \mathbf{A}$ from (6), we have

$$\text{L.H.S. of (3)} = \nabla \cdot \nabla \times \mathbf{A}$$

$$= [\nabla \nabla \mathbf{A}]$$

$$= 0 \quad \text{by the property of scalar triple product.}$$

This shows that the equation (6) is a solution of (3).

Again putting $\mathbf{A} = \nabla \times \mathbf{A}$ in (2), we get

$$\nabla \times \mathbf{E} = -\frac{1}{c} \frac{\partial}{\partial t} (\nabla \times \mathbf{A})$$

or $\nabla \times \left(\mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t} \right) = 0$ which shows that the bracketed expression is the gradient of some scalar function say ϕ therefore,

$$\mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t} = -\text{grad } \phi = -\nabla \phi$$

$$\text{i.e. } \mathbf{E} = -\nabla \phi - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t}$$

which is the equation (5) showing that equation (5) is a solution of (3).

1.36. CURVILINEAR CO-ORDINATES

We know that the equations

$$u = f_1(x, y, z); v = f_2(x, y, z); w = f_3(x, y, z)$$

where u, v, w are parameters, represent three families of surfaces when expressed in the form

$$u = \text{const.}, v = \text{const.}, w = \text{const.},$$

where u, v, w are continuously differentiable functions defined in

region R of space.

Suppose that the three surfaces $u = \text{const.}$, $v = \text{const.}$, $w = \text{const.}$, intersect in a point P of the region R . The values of u, v, w for the three surfaces intersecting at P are called the *curvilinear co-ordinates* of the point P . The three surfaces are then known as *co-ordinate surfaces*. The three surfaces intersect pairwise in three curves known as *co-ordinate curves*. Only one co-ordinate is variable on each of the co-ordinate curves. The curve on which u varies known as u -curve and similarly v -curve and w -curve are those on which v and w respectively are variable. One variable is constant on each of the co-ordinate surfaces. The surface on which u is constant is known as u -surface and similarly v -surface and w -surface are those on which v and w respectively are constant.

Using the equation (1) the rectangular co-ordinates (x, y, z) and therefore the position vector $\mathbf{r}$ of any point in the region of space can be expressed in terms of curvilinear co-ordinates. Since there is a one to one correspondence between x, y, z and u, v, w the position vector $\mathbf{r}$ is a vector function of u, v, w .

Note. The loci of $u = C_1$, $v = C_2$, $w = C_3$; C_1, C_2, C_3 being constants represent the co-ordinate surfaces and the equations of the co-ordinate curves then are

$$v = C_2, w = C_3; w = C_3, u = C_1; u = C_1, v = C_2.$$

1.37. ORTHOGONAL CURVILINEAR CO-ORDINATES

A system of orthogonal curvilinear co-ordinates is one which corresponds to the points of intersection of a triply orthogonal system of three families of surfaces

$$u(x, y, z) = \text{const.}, v(x, y, z) = \text{const.}, w(x, y, z) = \text{const.}$$

such that, through each point P in any region R of space there is one and only one member of each family, each of the three surfaces cutting the other two orthogonally. In short the curvilinear co-ordinates u, v, w are said to be orthogonal if the co-ordinate curves are mutually perpendicular at every point $P(x, y, z)$ of space.

Let us suppose that $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ form a right handed system of unit vectors tangent to the co-ordinate curves u, v, w respectively at P and directed towards increasing u, v, w . Then we have

$$\left. \begin{aligned} \mathbf{e}_1 &= \mathbf{e}_2 \times \mathbf{e}_3, \mathbf{e}_2 = \mathbf{e}_3 \times \mathbf{e}_1 \text{ and } \mathbf{e}_3 = \mathbf{e}_1 \times \mathbf{e}_2 \\ \mathbf{e}_1 \cdot \mathbf{e}_2 &= \mathbf{e}_2 \cdot \mathbf{e}_3 = \mathbf{e}_3 \cdot \mathbf{e}_1 = 0 \end{aligned} \right\} \quad \dots (1)$$

Let the arc lengths measured along the co-ordinate curves in the three directions of u, v, w be respectively s_1, s_2, s_3 . Now consider a small parallelepiped whose diagonal is the element of arc ds of a curve tangent to PQ at P and faces coincide with planes u, v and length of edges are ds_1, ds_2, ds_3 . Therefore,

$$ds^2 = ds_1^2 + ds_2^2 + ds_3^2. \quad \dots (2)$$

Let us now introduce the three numbers h_1, h_2, h_3 known as *metrical coefficients* with the property

$$\frac{ds_1}{du} = h_1, \frac{ds_2}{dv} = h_2, \frac{ds_3}{dw} = h_3$$

$$\text{i.e. } ds_1 = h_1 du, ds_2 = h_2 dv, ds_3 = h_3 dw. \quad \dots (1)$$

Substituting, the values of ds_1, ds_2, ds_3 from (3) in (2) we get

$$ds^2 = h_1^2 du^2 + h_2^2 dv^2 + h_3^2 dw^2. \quad \dots (2)$$




Fig. 1.40




Fig. 1.41

Now if $\mathbf{r}$ be the position vector of P , referred to the origin of rectangular co-ordinate system, the tangents to the co-ordinate curves at P are parallel to the directions of $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ and have the magnitudes h_1, h_2, h_3 respectively. Therefore,

$$\frac{\partial \mathbf{r}}{\partial u} = h_1 \mathbf{e}_1, \frac{\partial \mathbf{r}}{\partial v} = h_2 \mathbf{e}_2, \frac{\partial \mathbf{r}}{\partial w} = h_3 \mathbf{e}_3. \quad \dots (3)$$

$$\text{These give, } \frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} = h_1 h_2 \mathbf{e}_1 \times \mathbf{e}_2 = h_1 h_2 \mathbf{e}_3 \quad \text{from (1)}$$

$$\text{i.e. } \frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} = \frac{h_1 h_2}{h_3} \frac{\partial \mathbf{r}}{\partial w} \quad \text{from (5)}$$

$$\text{Similarly } \frac{\partial \mathbf{r}}{\partial v} \times \frac{\partial \mathbf{r}}{\partial w} = \frac{h_2 h_3}{h_1} \frac{\partial \mathbf{r}}{\partial u} \quad \dots (6)$$

$$\text{and } \frac{\partial \mathbf{r}}{\partial w} \times \frac{\partial \mathbf{r}}{\partial u} = \frac{h_3 h_1}{h_2} \frac{\partial \mathbf{r}}{\partial v} \quad \dots (7)$$

$$\text{Also } \left[\frac{\partial \mathbf{r}}{\partial u} \frac{\partial \mathbf{r}}{\partial v} \frac{\partial \mathbf{r}}{\partial w} \right] = \left(\frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} \right) \cdot \frac{\partial \mathbf{r}}{\partial w} = h_1 h_2 \mathbf{e}_3 \cdot h_3 \mathbf{e}_3 = h_1 h_2 h_3 \quad \dots (7)$$

1.38. CONDITION FOR ORTHOGONALITY

We have mentioned that the curvilinear co-ordinates of a point in space are determined by three continuously differentiable scalar functions

$$u = f_1(x, y, z), v = f_2(x, y, z), w = f_3(x, y, z)$$

when these functions u, v, w are not functionally related, then these equations can be solved to give x, y, z in terms of u, v, w such that

$$x = g_1(u, v, w), y = g_2(u, v, w), z = g_3(u, v, w)$$

where x, y, z are continuously differentiable functions of u, v, w .

Now the position vector $\mathbf{r}$ of any point in space, referred to the origin of rectangular axes along which the unit vectors are $\mathbf{i}, \mathbf{j}, \mathbf{k}$, is expressed as

$$\begin{aligned} \mathbf{r} &= x\mathbf{i} + y\mathbf{j} + z\mathbf{k} \\ &= g_1(u, v, w)\mathbf{i} + g_2(u, v, w)\mathbf{j} + g_3(u, v, w)\mathbf{k} \\ &= \mathbf{F}(u, v, w) \text{ say.} \end{aligned}$$

Now if $u = C_1, v = C_2, w = C_3$ where C_1, C_2, C_3 are constants, represent the co-ordinate surfaces, then the co-ordinate curves are

$$v = C_2, w = C_3; w = C_3, u = C_1; u = C_1, v = C_2$$

The co-ordinate curve $v = C_2, w = C_3$ through (C_1, C_2, C_3) is the curve as

$$\mathbf{r} = \mathbf{F}(u, C_2, C_3), u \text{ being the parameter.}$$

Now the tangent to the curve $v = C_2, w = C_3$ is parallel to the vector $\frac{\partial \mathbf{r}}{\partial u}$.

Similarly the tangents to the curves $w = C_3, u = C_1$ and $u = C_1, v = C_2$ are respectively parallel to the vectors $\frac{\partial \mathbf{r}}{\partial v}$ and $\frac{\partial \mathbf{r}}{\partial w}$.

Since the dot product of two parallel vectors is zero, it therefore follows that the curvilinear co-ordinate system will be orthogonal if

$$\frac{\partial \mathbf{r}}{\partial u} \cdot \frac{\partial \mathbf{r}}{\partial v} = \frac{\partial \mathbf{r}}{\partial v} \cdot \frac{\partial \mathbf{r}}{\partial w} = \frac{\partial \mathbf{r}}{\partial w} \cdot \frac{\partial \mathbf{r}}{\partial u} = 0. \quad \dots (1)$$

These are the required conditions for orthogonality.

COROLLARY. The line element ds derived in relation (4) of § 1.37, may be deduced from the conditions of orthogonality.

We have $\mathbf{r} = \mathbf{F}(u, v, w)$

$$\text{so that } d\mathbf{r} = \frac{\partial \mathbf{r}}{\partial u} du + \frac{\partial \mathbf{r}}{\partial v} dv + \frac{\partial \mathbf{r}}{\partial w} dw$$

$$\begin{aligned} d\mathbf{r} \cdot d\mathbf{r} &= \left\{ \frac{\partial \mathbf{r}}{\partial u} du + \frac{\partial \mathbf{r}}{\partial v} dv + \frac{\partial \mathbf{r}}{\partial w} dw \right\} \cdot \left\{ \frac{\partial \mathbf{r}}{\partial u} du + \frac{\partial \mathbf{r}}{\partial v} dv + \frac{\partial \mathbf{r}}{\partial w} dw \right\} \\ &= \left(\frac{\partial \mathbf{r}}{\partial u} \right)^2 du^2 + \left(\frac{\partial \mathbf{r}}{\partial v} \right)^2 dv^2 + \left(\frac{\partial \mathbf{r}}{\partial w} \right)^2 dw^2 \end{aligned}$$

$$+ 2 \frac{\partial \mathbf{r}}{\partial u} \cdot \frac{\partial \mathbf{r}}{\partial v} du dv + 2 \frac{\partial \mathbf{r}}{\partial v} \cdot \frac{\partial \mathbf{r}}{\partial w} dv dw + 2 \frac{\partial \mathbf{r}}{\partial w} \cdot \frac{\partial \mathbf{r}}{\partial u} dw du \quad \dots (2)$$

Applying the conditions of orthogonality, this reduces to

$$(ds)^2 = \left(\frac{\partial \mathbf{r}}{\partial u} \right)^2 du^2 + \left(\frac{\partial \mathbf{r}}{\partial v} \right)^2 dv^2 + \left(\frac{\partial \mathbf{r}}{\partial w} \right)^2 dw^2 \quad \dots (3)$$

Putting $(d\mathbf{r})^2 = |d\mathbf{r}|^2 = ds^2$

$$\left(\frac{\partial \mathbf{r}}{\partial u}\right)^2 = \left|\frac{d\mathbf{r}}{du}\right|^2 = h_1^2, \quad \left(\frac{\partial \mathbf{r}}{\partial v}\right)^2 = h_2^2, \quad \left(\frac{\partial \mathbf{r}}{\partial w}\right)^2 = h_3^2.$$

The relation (3) yields

$$ds^2 = h_1^2 du^2 + h_2^2 dv^2 + h_3^2 dw^2.$$

1.39. RECIPROCAL SETS OF TWO TRIADS OF MUTUALLY ORTHOGONAL VECTORS

If u, v, w be a set of curvilinear co-ordinates of a point P whose position vector is $\mathbf{r}$ with respect to the origin of a rectangular co-ordinate system, then the two sets of triads of mutually orthogonal vectors, $\nabla u, \nabla v, \nabla w$ and $\frac{\partial \mathbf{r}}{\partial u}, \frac{\partial \mathbf{r}}{\partial v}, \frac{\partial \mathbf{r}}{\partial w}$ are reciprocal to each other.

We have shown in equation (7) of §1.37 that if $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ form a right handed system of unit vectors tangent to the co-ordinate curves u, v, w respectively at P and directed towards increasing u, v, w , then

$$\left[\frac{\partial \mathbf{r}}{\partial u} \frac{\partial \mathbf{r}}{\partial v} \frac{\partial \mathbf{r}}{\partial w}\right] = h_1 h_2 h_3. \quad \dots (1)$$

Now $\nabla u, \nabla v, \nabla w$ are the vectors lying along the normals to the co-ordinate surfaces which are the level surfaces of the function u, v, w . But the curvilinear co-ordinate system is orthogonal, therefore the conditions of orthogonality stated in §1.38 when applied to the orthogonal system of vectors $\nabla u, \nabla v, \nabla w$, yield

$$\nabla u \cdot \nabla v = 0, \quad \nabla v \cdot \nabla w = 0, \quad \nabla w \cdot \nabla u = 0. \quad \dots (2)$$

Let us now assume that the mutually orthogonal unit vectors are

$$\frac{\nabla u}{|\nabla u|}, \quad \frac{\nabla v}{|\nabla v|}, \quad \frac{\nabla w}{|\nabla w|}, \quad \dots (3)$$

which form a right handed system.

Here $|\nabla u|$ is the directional derivative of u along the direction of the normal to the surface $u = C_1$ i.e. along the tangent to the curve $v = C_2, w = C_3$. Hence if ds_1 represents the differential of length along this curve, then we can state

$$|\nabla u| = \frac{du}{ds_1}.$$

But from the Corollary of §1.38 the line element ds_1 along the curve $v = C_2, w = C_3$ will be obtained by putting $dv = 0, dw = 0$ in $ds^2 = h_1^2 du^2 + h_2^2 dv^2 + h_3^2 dw^2$, whence we have

$$ds_1^2 = h_1^2 du^2$$

giving

$$\frac{du}{ds_1} = \frac{1}{h_1},$$

so that

$$|\nabla u| = \frac{1}{h_1}, \quad \text{i.e.} \quad h_1 = \frac{1}{|\nabla u|}.$$

Similarly,

$$h_2 = \frac{1}{|\nabla v|} \quad \text{and} \quad h_3 = \frac{1}{|\nabla w|}.$$

Since, $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ is a system of orthogonal unit vectors therefore (3) and (4) give

$$\frac{\nabla u}{|\nabla u|} = \frac{\nabla u}{1/h_1} = \mathbf{e}_1 \quad \text{i.e.} \quad \nabla u = \frac{\mathbf{e}_1}{h_1}$$

and similarly

$$\nabla v = \frac{\mathbf{e}_2}{h_2}, \quad \nabla w = \frac{\mathbf{e}_3}{h_3}.$$

So that $[\nabla u \nabla v \nabla w] = (\nabla u \times \nabla v) \cdot \nabla w \quad \dots (5)$

$$= \left(\frac{\mathbf{e}_1}{h_1} \times \frac{\mathbf{e}_2}{h_2}\right) \cdot \frac{\mathbf{e}_3}{h_3} = \frac{\mathbf{e}_3}{h_1 h_2} \cdot \frac{\mathbf{e}_3}{h_3} \quad \text{as} \quad \mathbf{e}_1 \times \mathbf{e}_2 = \mathbf{e}_3 \quad \text{and} \quad \mathbf{e}_3 \cdot \mathbf{e}_3 = 1$$

$\dots (6)$

$$= \frac{1}{h_1 h_2 h_3}.$$

Multiplying (1) and (6), together, we get

$$[\nabla u \nabla v \nabla w] \left[\frac{\partial \mathbf{r}}{\partial u} \frac{\partial \mathbf{r}}{\partial v} \frac{\partial \mathbf{r}}{\partial w}\right] = 1$$

which follows that $\nabla u, \nabla v, \nabla w$ form a set of vectors reciprocal to $\frac{\partial \mathbf{r}}{\partial u}, \frac{\partial \mathbf{r}}{\partial v}, \frac{\partial \mathbf{r}}{\partial w}$.

1.40. GRADIENT IN TERMS OF ORTHOGONAL CURVILINEAR COORDINATES

Let $\phi(u, v, w)$ be any scalar point function given in terms of orthogonal curvilinear coordinates u, v, w .

Since u, v, w may be supposed to be the functions of rectangular Cartesian coordinates x, y, z , therefore

$$\frac{\partial \phi}{\partial x} = \frac{\partial \phi}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial \phi}{\partial v} \frac{\partial v}{\partial x} + \frac{\partial \phi}{\partial w} \frac{\partial w}{\partial x} \quad \dots (1)$$

$$\frac{\partial \phi}{\partial y} = \frac{\partial \phi}{\partial u} \frac{\partial u}{\partial y} + \frac{\partial \phi}{\partial v} \frac{\partial v}{\partial y} + \frac{\partial \phi}{\partial w} \frac{\partial w}{\partial y} \quad \dots (2)$$

$$\frac{\partial \phi}{\partial z} = \frac{\partial \phi}{\partial u} \frac{\partial u}{\partial z} + \frac{\partial \phi}{\partial v} \frac{\partial v}{\partial z} + \frac{\partial \phi}{\partial w} \frac{\partial w}{\partial z} \quad \dots (3)$$

Let us now suppose that $\mathbf{i}, \mathbf{j}, \mathbf{k}$ are the unit vectors along the rectangular axes x, y, z respectively and $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ are the mutually orthogonal unit vectors along the tangents to the coordinate curves u, v, w . Then by relation (5) of §1.39, we have

$$\nabla u = \frac{\mathbf{e}_1}{h_1}, \quad \nabla v = \frac{\mathbf{e}_2}{h_2}, \quad \nabla w = \frac{\mathbf{e}_3}{h_3} \quad \dots (4)$$

Now multiplying (1) by $\mathbf{i}$, (2) by $\mathbf{j}$ and (3) by $\mathbf{k}$ and then adding together we get

$$\left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \phi = \frac{\partial \phi}{\partial x} \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) u$$

$$\frac{\partial \phi}{\partial v} \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) v + \frac{\partial \phi}{\partial w} \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) w$$

i.e.,

$$\nabla \phi = \frac{\partial \phi}{\partial u} \nabla u + \frac{\partial \phi}{\partial v} \nabla v + \frac{\partial \phi}{\partial w} \nabla w,$$

$$= \frac{1}{h_1} \frac{\partial \phi}{\partial u} \mathbf{e}_1 + \frac{1}{h_2} \frac{\partial \phi}{\partial v} \mathbf{e}_2 + \frac{1}{h_3} \frac{\partial \phi}{\partial w} \mathbf{e}_3 \text{ by (4) } \dots$$

which is the required expression. It is obvious that the components of grad ϕ i.e. $\nabla \phi$ along the unit vectors $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ are respectively

$$\frac{1}{h_1} \frac{\partial \phi}{\partial u}, \frac{1}{h_2} \frac{\partial \phi}{\partial v}, \frac{1}{h_3} \frac{\partial \phi}{\partial w}. \quad (\text{Agra, 1971})$$

1.41. DIVERGENCE IN TERMS OF ORTHOGONAL CURVILINEAR COORDINATES

(Agra, 1971)

Let $\mathbf{F}(u, v, w)$ be a vector point function given in terms of orthogonal curvilinear coordinates u, v, w and let F_1, F_2, F_3 be the components of $\mathbf{F}$ along $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ the unit vectors along u, v, w axes. Then,

$$\mathbf{F} = F_1 \mathbf{e}_1 + F_2 \mathbf{e}_2 + F_3 \mathbf{e}_3$$

$$= F_1 \mathbf{e}_2 \times \mathbf{e}_3 + F_2 \mathbf{e}_3 \times \mathbf{e}_1 + F_3 \mathbf{e}_1 \times \mathbf{e}_2 \quad \text{since } \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3 \text{ mutually orthogonal vectors}$$

$$= F_1 h_2 h_3 \nabla v \times \nabla w + F_2 h_3 h_1 \nabla w \times \nabla u + F_3 h_1 h_2 \nabla u \times \nabla v$$

by (5) of § 1.39

$$\therefore \nabla \cdot \mathbf{F} = \nabla \cdot (F_1 h_2 h_3 \nabla v \times \nabla w + F_2 h_3 h_1 \nabla w \times \nabla u + F_3 h_1 h_2 \nabla u \times \nabla v)$$

$$= \nabla \cdot (F_1 h_2 h_3 \nabla v \times \nabla w) + \nabla \cdot (F_2 h_3 h_1 \nabla w \times \nabla u) + \nabla \cdot (F_3 h_1 h_2 \nabla u \times \nabla v) \dots$$

By the properties of divergence and curl, we have

$$\nabla \cdot (F_1 h_2 h_3 \nabla v \times \nabla w) = F_1 h_2 h_3 \nabla \cdot (\nabla v \times \nabla w) + \nabla v \times \nabla w \cdot \nabla (F_1 h_2 h_3)$$

where $\nabla \cdot (\nabla v \times \nabla w) = \nabla w \cdot \text{curl } \nabla v - \nabla v \cdot \text{curl } \nabla w$

$$= 0$$

and $\nabla (F_1 h_2 h_3) = \frac{\partial}{\partial x} (F_1 h_2 h_3) \nabla u + \frac{\partial}{\partial v} (F_1 h_2 h_3) \nabla v + \frac{\partial}{\partial w} (F_1 h_2 h_3) \nabla w$

So that $\nabla \cdot (F_1 h_2 h_3 \nabla v \times \nabla w)$

$$= \nabla v \times \nabla w \cdot \left\{ \frac{\partial}{\partial u} (F_1 h_2 h_3) \nabla u + \frac{\partial}{\partial v} (F_1 h_2 h_3) \nabla v + \frac{\partial}{\partial w} (F_1 h_2 h_3) \nabla w \right\}$$

$$= \nabla v \times \nabla w \cdot \nabla u \frac{\partial}{\partial u} (F_1 h_2 h_3) \text{ other terms vanish, by the properties of scalar triple product}$$

$$= \frac{\mathbf{e}_2}{h_2} \times \frac{\mathbf{e}_3}{h_3} \cdot \frac{\mathbf{e}_1}{h_1} \frac{\partial}{\partial u} (F_1 h_2 h_3) \text{ by (5) of § 1.39.}$$

$$= \frac{1}{h_1 h_2 h_3} [\mathbf{e}_2 \times \mathbf{e}_3 \cdot \mathbf{e}_1] \frac{\partial}{\partial u} (F_1 h_2 h_3)$$

$$= \frac{1}{h_1 h_2 h_3} \frac{\partial}{\partial u} (F_1 h_2 h_3) \text{ since } \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3 \text{ being unit vectors } [\mathbf{e}_2 \times \mathbf{e}_3 \cdot \mathbf{e}_1] = 1$$

Similarly $\nabla \cdot (F_2 h_3 h_1 \nabla w \times \nabla u) = \frac{1}{h_1 h_2 h_3} \frac{\partial}{\partial v} (F_2 h_3 h_1)$

and $\nabla \cdot (F_3 h_1 h_2 \nabla u \times \nabla v) = \frac{1}{h_1 h_2 h_3} \frac{\partial}{\partial w} (F_3 h_1 h_2)$

Substituting these values in (1) we get

$$\nabla \cdot \mathbf{F} = \frac{1}{h_1 h_2 h_3} \left[\frac{\partial}{\partial x} (F_1 h_2 h_3) + \frac{\partial}{\partial v} (F_2 h_3 h_1) + \frac{\partial}{\partial w} (F_3 h_1 h_2) \right] \dots (2)$$

which gives the required expression for div $\mathbf{F}$.

1.42. CURL IN TERMS OF ORTHOGONAL CURVILINEAR COORDINATES

(Agra 1971, 74)

Let $\mathbf{F}(u, v, w)$ be a vector function given in terms of orthogonal curvilinear coordinates u, v, w and let F_1, F_2, F_3 be the components of $\mathbf{F}$ along $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ the unit vectors along u, v, w axes. Then,

$$\mathbf{F} = F_1 \mathbf{e}_1 + F_2 \mathbf{e}_2 + F_3 \mathbf{e}_3$$

$$= h_1 F_1 \nabla u + h_2 F_2 \nabla v + h_3 F_3 \nabla w \text{ by (5) of § 1.39.}$$

$$\nabla \times \mathbf{F} = \nabla \times \{ h_1 F_1 \nabla u + h_2 F_2 \nabla v + h_3 F_3 \nabla w \}$$

$$= \nabla \times (h_1 F_1 \nabla u) + \nabla \times (h_2 F_2 \nabla v) + \nabla \times (h_3 F_3 \nabla w) \dots (1)$$

But, we have by the properties of curl

$$\nabla \times (h_1 F_1 \nabla u) = \nabla (h_1 F_1) \times \nabla u + h_1 F_1 \nabla \times \nabla u$$

$$= \nabla (h_1 F_1) \times \nabla u \text{ since } \nabla \times \nabla u = 0$$

$$= \left\{ \frac{\partial}{\partial u} (h_1 F_1) \nabla u + \frac{\partial}{\partial v} (h_1 F_1) \nabla v + \frac{\partial}{\partial w} (h_1 F_1) \nabla w \right\} \times \nabla u$$

$$= \frac{\partial}{\partial v} (h_1 F_1) \nabla v \times \nabla u + \frac{\partial}{\partial w} (h_1 F_1) \nabla w \times \nabla u,$$

the other term vanishes

$$= \frac{1}{h_1 h_2} \frac{\partial}{\partial v} (h_1 F_1) \mathbf{e}_2 \times \mathbf{e}_1 + \frac{1}{h_3 h_1} \frac{\partial}{\partial w} (h_1 F_1) \mathbf{e}_3 \times \mathbf{e}_1$$

by (5) of § 1.39

$$= -\frac{1}{h_1 h_2} \frac{\partial}{\partial v} (h_1 F_1) \mathbf{e}_3 + \frac{1}{h_3 h_1} \frac{\partial}{\partial w} (h_1 F_1) \mathbf{e}_2.$$

Similarly

$$\nabla \times (h_2 F_2 \nabla v) = -\frac{1}{h_2 h_3} \frac{\partial}{\partial w} (h_2 F_2) \mathbf{e}_1 + \frac{1}{h_2 h_1} \frac{\partial}{\partial u} (h_2 F_2) \mathbf{e}_3$$

and
$$\nabla \times (h_3 F_3 \nabla w) = -\frac{1}{h_3 h_1} \frac{\partial}{\partial w} (h_2 F_3) \mathbf{e}_2 + \frac{1}{h_3 h_2} \frac{\partial}{\partial v} (h_3 F_3) \mathbf{e}_1$$

Substituting these values in (1), we get

$$\begin{aligned} \nabla \times \mathbf{F} &= \frac{1}{h_2 h_3} \left[\frac{\partial}{\partial v} (h_3 F_3) - \frac{\partial}{\partial w} (h_2 F_2) \right] \mathbf{e}_1 \\ &\quad + \frac{1}{h_3 h_1} \left[\frac{\partial}{\partial w} (h_1 F_1) - \frac{\partial}{\partial u} (h_3 F_3) \right] \mathbf{e}_2 \\ &\quad + \frac{1}{h_1 h_2} \left[\frac{\partial}{\partial u} (h_2 F_2) - \frac{\partial}{\partial v} (h_1 F_1) \right] \mathbf{e}_3 \\ &= \frac{1}{h_1 h_2 h_3} \begin{vmatrix} h_1 \mathbf{e}_1 & h_2 \mathbf{e}_2 & h_3 \mathbf{e}_3 \\ \frac{\partial}{\partial u} & \frac{\partial}{\partial v} & \frac{\partial}{\partial w} \\ h_1 F_1 & h_2 F_2 & h_3 F_3 \end{vmatrix} \dots (2) \end{aligned}$$

which is the required expression for curl $\mathbf{F}$.

This result may also be expressed as

$$\nabla \times \mathbf{F} = \frac{1}{h_1 h_2 h_3} \begin{vmatrix} h_1 \mathbf{e}_1 & h_2 \mathbf{e}_2 & h_3 \mathbf{e}_3 \\ \frac{\partial}{\partial u} & \frac{\partial}{\partial v} & \frac{\partial}{\partial w} \\ h_1 \mathbf{e}_1 \cdot \mathbf{F} & h_2 \mathbf{e}_2 \cdot \mathbf{F} & h_3 \mathbf{e}_3 \cdot \mathbf{F} \end{vmatrix} \quad \text{since } \mathbf{e}_1 \cdot \mathbf{F} = F_1 \text{ etc.}$$

1.43. LAPLACIAN (∇^2) IN TERMS OF ORTHOGONAL CURVILINEAR CO-ORDINATES

By (5) of § 1.40, we have

$$\nabla \phi = \frac{1}{h_1} \frac{\partial \phi}{\partial u} \mathbf{e}_1 + \frac{1}{h_2} \frac{\partial \phi}{\partial v} \mathbf{e}_2 + \frac{1}{h_3} \frac{\partial \phi}{\partial w} \mathbf{e}_3.$$

$$\begin{aligned} \therefore \nabla^2 \phi &= \nabla \cdot \nabla \phi \\ &= \left(\mathbf{e}_1 \frac{\partial}{\partial u} + \mathbf{e}_2 \frac{\partial}{\partial v} + \mathbf{e}_3 \frac{\partial}{\partial w} \right) \left(\frac{1}{h_1} \frac{\partial \phi}{\partial u} \mathbf{e}_1 + \frac{1}{h_2} \frac{\partial \phi}{\partial v} \mathbf{e}_2 + \frac{1}{h_3} \frac{\partial \phi}{\partial w} \mathbf{e}_3 \right) \\ &= \frac{1}{h_1 h_2 h_3} \left[\frac{\partial}{\partial u} \left(\frac{h_2 h_3}{h_1} \frac{\partial \phi}{\partial u} \right) + \frac{\partial}{\partial v} \left(\frac{h_3 h_1}{h_2} \frac{\partial \phi}{\partial v} \right) + \frac{\partial}{\partial w} \left(\frac{h_1 h_2}{h_3} \frac{\partial \phi}{\partial w} \right) \right] \end{aligned}$$

which is the required expression.

1.44. EQUIVALENT EXPRESSIONS FOR $\nabla \phi$, $\nabla \cdot \mathbf{F}$ AND $\nabla \times \mathbf{F}$ IN RECTANGULAR CO-ORDINATES

In §§ 1.40, 1.41 and 1.42 we have derived the expressions for $\nabla \phi$, $\nabla \cdot \mathbf{F}$ and $\nabla \times \mathbf{F}$ in terms of curvilinear co-ordinate system as follows:

$$\nabla \phi = \frac{1}{h_1} \frac{\partial \phi}{\partial u} \mathbf{e}_1 + \frac{1}{h_2} \frac{\partial \phi}{\partial v} \mathbf{e}_2 + \frac{1}{h_3} \frac{\partial \phi}{\partial w} \mathbf{e}_3 \dots (1)$$

$$\nabla \cdot \mathbf{F} = \frac{1}{h_1 h_2 h_3} \left[\frac{\partial}{\partial u} (F_1 h_2 h_3) + \frac{\partial}{\partial v} (F_2 h_3 h_1) + \frac{\partial}{\partial w} (F_3 h_1 h_2) \right] \dots (2)$$

and
$$\begin{aligned} \nabla \times \mathbf{F} &= \frac{1}{h_2 h_3} \left[\frac{\partial}{\partial v} (h_3 F_3) - \frac{\partial}{\partial w} (h_2 F_2) \right] \mathbf{e}_1 \\ &\quad + \frac{1}{h_3 h_1} \left[\frac{\partial}{\partial w} (h_1 F_1) - \frac{\partial}{\partial u} (h_3 F_3) \right] \mathbf{e}_2 \\ &\quad + \frac{1}{h_1 h_2} \left[\frac{\partial}{\partial u} (h_2 F_2) - \frac{\partial}{\partial v} (h_1 F_1) \right] \mathbf{e}_3 \dots (3) \end{aligned}$$

where $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ are unit vectors along u, v, w axes

In order to get the equivalent expressions for these quantities in cartesian rectangular co-ordinates, if we use the transformations,

$$u=x, \quad v=y, \quad w=z$$

we have

$$ds^2 = dx^2 + dy^2 + dz^2$$

so that

$$h_1 = h_2 = h_3 = 1,$$

and the unit vectors $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ are taken as usual unit vectors $\mathbf{i}, \mathbf{j}, \mathbf{k}$.

With these substitutions, the relations (1), (2) and (3) become

$$\nabla \phi = \frac{\partial \phi}{\partial x} \mathbf{i} + \frac{\partial \phi}{\partial y} \mathbf{j} + \frac{\partial \phi}{\partial z} \mathbf{k}$$

$$\nabla \cdot \mathbf{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$$

and

$$\nabla \times \mathbf{F} = \left(\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z} \right) \mathbf{i} + \left(\frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x} \right) \mathbf{j} + \left(\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) \mathbf{k}.$$

1.45. CYLINDRICAL CO-ORDINATES AS A SPECIAL CURVILINEAR SYSTEM

Let P be a point in space such that its curvilinear co-ordinates are (u, v, w) and cartesian co-ordinates are (x, y, z) . Let the projection of P in the xy -plane be Q whose polar co-ordinates in the plane are (r, θ) . Then the cylindrical or cylindrical co-ordinates of the point P are specified by

$$u=r, \quad v=\theta, \quad w=z.$$

These co-ordinates are transformed to cartesian co-ordinates by the help of the relations

$$x=r \cos \theta, \quad y=r \sin \theta, \quad z=z.$$

$$r^2 = x^2 + y^2, \quad \theta = \tan^{-1} \frac{y}{x}, \quad z=z.$$

Since x is a function of r, θ, z ,

we have

$$dx = \frac{\partial x}{\partial r} dr + \frac{\partial x}{\partial \theta} d\theta + \frac{\partial x}{\partial z} dz$$

$$= \cos \theta dr - r \sin \theta d\theta.$$

Similarly

$$dy = \sin \theta dr + r \cos \theta d\theta$$




Fig. 1.42

and

$$dz = dz.$$

Therefore, the relation $(ds)^2 = (dx)^2 + (dy)^2 + (dz)^2$ gives

$$\begin{aligned} (ds)^2 &= (\cos \theta dr - r \sin \theta d\theta)^2 + (\sin \theta dr + r \cos \theta d\theta)^2 + (dz)^2 \\ &= (dr)^2 + r^2 (d\theta)^2 + (dz)^2 \end{aligned}$$

Also from relation (4) of § 1.37, we have

$$ds^2 = h_1^2 du^2 + h_2^2 dv^2 + h_3^2 dw^2$$

which when transformed by the substitutions $u=r$, $v=\theta$, $w=z$ becomes

$$ds^2 = h_1^2 dr^2 + h_2^2 d\theta^2 + h_3^2 dz^2$$

Comparing (1) and (2), we get

$$h_1=1, \quad h_2=r, \quad h_3=1.$$

With these substitutions, the functions $\nabla\phi$, $\nabla^2\phi$, $\nabla \cdot \mathbf{F}$ and $\nabla \times \mathbf{F}$ in cylindrical coordinates become

$$\nabla\phi = \frac{\partial\phi}{\partial r} \mathbf{e}_1 + \frac{1}{r} \frac{\partial\phi}{\partial\theta} \mathbf{e}_2 + \frac{\partial\phi}{\partial z} \mathbf{e}_3$$

$$\nabla^2\phi = \frac{\partial^2\phi}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2\phi}{\partial\theta^2} + \frac{\partial^2\phi}{\partial z^2} + \frac{1}{r} \frac{\partial\phi}{\partial r}$$

$$\nabla \cdot \mathbf{F} = \frac{1}{r} \frac{\partial}{\partial r}(rF_1) + \frac{1}{r} \frac{\partial F_2}{\partial\theta} + \frac{\partial F_3}{\partial z} \quad (\text{Rohilkhand, 1976})$$

$$\begin{aligned} \nabla \times \mathbf{F} &= \left(\frac{1}{r} \frac{\partial F_3}{\partial\theta} - \frac{\partial F_2}{\partial z} \right) \mathbf{e}_1 + \left(\frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial r} \right) \mathbf{e}_2 \\ &\quad + \left\{ \frac{1}{r} \frac{\partial}{\partial r}(rF_2) - \frac{1}{r} \frac{\partial F_1}{\partial\theta} \right\} \mathbf{e}_3. \end{aligned}$$

(Agra, 1974; Rohilkhand 1976)

1.46. SPHERICAL POLAR COORDINATES AS A SPECIAL CURVILINEAR SYSTEM

Let P be a point in space such that its curvilinear coordinates are (u, v, w) and cartesian coordinates are (x, y, z) . Then spherical polar coordinates of the point P are specified

$$u=r, \quad v=\theta, \quad w=\phi$$

where $r (=OP)$ is the distance of the point P from the origin O , θ is the angle between OP and the z -axis and ϕ is the angle included between the xz -plane and the plane OPZ .

These coordinates are transformed to cartesian coordinates by the help of the relations

$$\begin{aligned} x &= r \sin \theta \cos \phi, \\ y &= r \sin \theta \sin \phi, \\ z &= r \cos \theta. \end{aligned}$$




Fig. 1.43

We have,

$$\begin{aligned} dx &= \frac{\partial x}{\partial r} dr + \frac{\partial x}{\partial\theta} d\theta + \frac{\partial x}{\partial\phi} d\phi \\ &= \sin \theta \cos \phi dr - r \sin \theta \sin \phi d\theta - r \sin \theta \cos \phi d\phi. \end{aligned}$$

Similarly

$$\begin{aligned} dy &= \sin \theta \sin \phi dr + r \cos \theta \sin \phi d\theta + r \sin \theta \cos \phi d\phi \\ dz &= \cos \theta dr - r \sin \theta d\theta. \end{aligned}$$

$(ds)^2 = (dx)^2 + (dy)^2 + (dz)^2$ gives

$$(ds)^2 = (dr)^2 + r^2 (d\theta)^2 + r^2 \sin^2 \theta (d\phi)^2. \quad \dots (1)$$

Also with these transformations, the relation (4) of § 1.37 becomes

$$(ds)^2 = (dr)^2 + r^2 (d\theta)^2 + r^2 \sin^2 \theta (d\phi)^2. \quad \dots (2)$$

Comparing (1) and (2) we get

$$h_1=1, \quad h_2=r, \quad h_3=r \sin \theta.$$

With these substitutions, the functions $\nabla\psi$, $\nabla^2\psi$, $\nabla \cdot \mathbf{F}$ and $\nabla \times \mathbf{F}$ in spherical polar coordinates become

$$\nabla\psi = \frac{\partial\psi}{\partial r} \mathbf{e}_1 + \frac{1}{r} \frac{\partial\psi}{\partial\theta} \mathbf{e}_2 + \frac{1}{r \sin \theta} \frac{\partial\psi}{\partial\phi} \mathbf{e}_3$$

$$\nabla^2\psi = \frac{\partial^2\psi}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2\psi}{\partial\theta^2} + \frac{1}{\sin \theta} \frac{\partial^2\psi}{\partial\phi^2} + \frac{2}{r} \frac{\partial\psi}{\partial r} + \frac{\cot \theta}{r^2} \frac{\partial\psi}{\partial\theta}$$

$$\nabla \cdot \mathbf{F} = \frac{1}{r^2} \frac{\partial}{\partial r}(r^2 F_1) + \frac{1}{r \sin \theta} \frac{\partial}{\partial\theta}(\sin \theta F_2) + \frac{1}{r \sin \theta} \frac{\partial F_3}{\partial\phi}$$

(Rohilkhand, 1976)

$$\begin{aligned} \nabla \times \mathbf{F} &= \frac{1}{r \sin \theta} \left[\frac{\partial}{\partial\theta} (\sin \theta F_3) - \frac{\partial F_2}{\partial\phi} \right] \mathbf{e}_1 \\ &\quad + \frac{1}{r} \left[\frac{1}{\sin \theta} \frac{\partial F_1}{\partial\phi} - \frac{\partial}{\partial r}(r F_3) \right] \mathbf{e}_2 + \frac{1}{r} \left[\frac{\partial}{\partial r}(r F_2) - \frac{\partial F_1}{\partial\theta} \right] \mathbf{e}_3. \end{aligned}$$

(Rohilkhand, 1976)

Problem 115. Find an expression for ds^2 in curvilinear coordinates (u, v, w) . Then determine ds^2 for the special case of an orthogonal

system. Let the position vector of a point in space be $\mathbf{r}$, where $\mathbf{r}$ is a vector function of u, v, w . Then,

$$d\mathbf{r} = \frac{\partial \mathbf{r}}{\partial u} du + \frac{\partial \mathbf{r}}{\partial v} dv + \frac{\partial \mathbf{r}}{\partial w} dw$$

$$\begin{aligned} ds^2 &= d\mathbf{r} \cdot d\mathbf{r} \quad (\text{by the assumption in § 1.38}) \\ &= \left(\frac{\partial \mathbf{r}}{\partial u} du + \frac{\partial \mathbf{r}}{\partial v} dv + \frac{\partial \mathbf{r}}{\partial w} dw \right)^2 \\ &= \left(\frac{\partial \mathbf{r}}{\partial u} \right)^2 du^2 + \left(\frac{\partial \mathbf{r}}{\partial v} \right)^2 dv^2 + \left(\frac{\partial \mathbf{r}}{\partial w} \right)^2 dw^2 + 2 \frac{\partial \mathbf{r}}{\partial u} \cdot \frac{\partial \mathbf{r}}{\partial v} du dv \\ &\quad + 2 \frac{\partial \mathbf{r}}{\partial v} \cdot \frac{\partial \mathbf{r}}{\partial w} dv dw + 2 \frac{\partial \mathbf{r}}{\partial w} \cdot \frac{\partial \mathbf{r}}{\partial u} dw du. \end{aligned}$$

But by § 1.38 the conditions of orthogonality are

$$\frac{\partial \mathbf{r}}{\partial u} \cdot \frac{\partial \mathbf{r}}{\partial v} = \frac{\partial \mathbf{r}}{\partial v} \cdot \frac{\partial \mathbf{r}}{\partial w} = \frac{\partial \mathbf{r}}{\partial w} \cdot \frac{\partial \mathbf{r}}{\partial u} = 0.$$

So that the above relation becomes

$$ds^2 = \left(\frac{\partial \mathbf{r}}{\partial u} \right)^2 du^2 + \left(\frac{\partial \mathbf{r}}{\partial v} \right)^2 dv^2 + \left(\frac{\partial \mathbf{r}}{\partial w} \right)^2 dw^2.$$

Problem 116. If $x = uv \cos w$, $y = uv \sin w$, $z = \frac{1}{2}(u^2 - v^2)$; find h_1, h_2, h_3 and show that $ds^2 = (u^2 + v^2)(du^2 + dv^2) + uv dw^2$.

We have

$$dx = \frac{\partial x}{\partial u} du + \frac{\partial x}{\partial v} dv + \frac{\partial x}{\partial w} dw$$

$$= u \cos w du + v \cos w dv - uv \sin w dw$$

$$dy = \frac{\partial y}{\partial u} du + \frac{\partial y}{\partial v} dv + \frac{\partial y}{\partial w} dw$$

$$= u \sin w du + v \sin w dv + uv \cos w dw$$

$$dz = \frac{\partial z}{\partial u} du + \frac{\partial z}{\partial v} dv + \frac{\partial z}{\partial w} dw$$

$$= u du - v dv + 0 \cdot dw.$$

$$\begin{aligned} \therefore ds^2 &= dx^2 + dy^2 + dz^2 \\ &= (u \cos w du + v \cos w dv - uv \sin w dw)^2 \\ &\quad + (u \sin w du + v \sin w dv + uv \cos w dw)^2 \\ &\quad + (u du - v dv)^2 \\ &= (u^2 + v^2) du^2 + (u^2 + v^2) dv^2 + u^2 v^2 dw^2. \end{aligned}$$

Now comparing this relation with

$$ds^2 = h_1^2 du^2 + h_2^2 dv^2 + h_3^2 dw^2.$$

We get $h_1 = \sqrt{(u^2 + v^2)}$, $h_2 = \sqrt{(u^2 + v^2)}$, $h_3 = uv$.

Problem 117. If $u = 2x + 3$, $v = y - 4$, $w = z + 2$, show that u, v, w are orthogonal and find ds^2 and the metrical coefficients h_1, h_2, h_3 .

The given relations can be expressed as

$$x = \frac{u}{2} - \frac{3}{2}, \quad y = v + 4, \quad z = w - 2.$$

Then, if $\mathbf{r}$ be the position vector of a point in space,

$$\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$$

$$= \left(\frac{u}{2} - \frac{3}{2} \right) \mathbf{i} + (v + 4) \mathbf{j} + (w - 2) \mathbf{k}$$

$$\text{So that } \frac{\partial \mathbf{r}}{\partial u} = \frac{1}{2} \mathbf{i}, \quad \frac{\partial \mathbf{r}}{\partial v} = \mathbf{j}, \quad \frac{\partial \mathbf{r}}{\partial w} = \mathbf{k}.$$

The system of coordinates u, v, w to be orthogonal, we must have

$$\frac{\partial \mathbf{r}}{\partial u} \cdot \frac{\partial \mathbf{r}}{\partial v} = 0, \quad \frac{\partial \mathbf{r}}{\partial v} \cdot \frac{\partial \mathbf{r}}{\partial w} = 0, \quad \frac{\partial \mathbf{r}}{\partial w} \cdot \frac{\partial \mathbf{r}}{\partial u} = 0.$$

$$\text{Here } \frac{\partial \mathbf{r}}{\partial u} \cdot \frac{\partial \mathbf{r}}{\partial v} = \frac{1}{2} \mathbf{i} \cdot \mathbf{j} = 0.$$

$$\text{Similarly } \frac{\partial \mathbf{r}}{\partial v} \cdot \frac{\partial \mathbf{r}}{\partial w} = 0 \quad \text{and} \quad \frac{\partial \mathbf{r}}{\partial w} \cdot \frac{\partial \mathbf{r}}{\partial u} = 0.$$

Hence u, v, w are orthogonal.

Now, to find ds^2 , we have

$$\begin{aligned} dx &= \frac{\partial x}{\partial u} du + \frac{\partial x}{\partial v} dv + \frac{\partial x}{\partial w} dw \\ &= \frac{1}{2} du \end{aligned}$$

Similarly

$$\begin{aligned} dy &= dv, \quad \text{and} \quad dz = dw \\ ds^2 &= dx^2 + dy^2 + dz^2 \\ &= \frac{1}{4} du^2 + dv^2 + dw^2. \end{aligned}$$

Again to find h_1, h_2, h_3 let us compare the last relation with

$$ds^2 = h_1^2 du^2 + h_2^2 dv^2 + h_3^2 dw^2.$$

Whence, we get

$$h_1 = \frac{1}{2}, \quad h_2 = 1, \quad h_3 = 1.$$

Problem 118. For spherical coordinates

$$x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi, \quad z = r \cos \theta.$$

(a) Prove that the components of $\frac{\partial \mathbf{r}}{\partial r}, \frac{\partial \mathbf{r}}{\partial \theta}, \frac{\partial \mathbf{r}}{\partial \phi}$ are given by

$$\frac{\partial \mathbf{r}}{\partial r} = \sin \theta \cos \phi \mathbf{i} + \sin \theta \sin \phi \mathbf{j} + \cos \theta \mathbf{k}$$

$$\frac{\partial \mathbf{r}}{\partial \theta} = r \cos \theta \cos \phi \mathbf{i} + r \cos \theta \sin \phi \mathbf{j} - r \sin \theta \mathbf{k}$$

$$\frac{\partial \mathbf{r}}{\partial \phi} = -r \sin \theta \sin \phi \mathbf{i} + r \sin \theta \cos \phi \mathbf{j}.$$

(b) Verify the mutual orthogonality of $\frac{\partial \mathbf{r}}{\partial r}, \frac{\partial \mathbf{r}}{\partial \theta}, \frac{\partial \mathbf{r}}{\partial \phi}$.

(c) Find expressions for $\nabla r, \nabla \theta, \nabla \phi$ and then show that they constitute a set of vectors reciprocal to $\frac{\partial \mathbf{r}}{\partial r}, \frac{\partial \mathbf{r}}{\partial \theta}, \frac{\partial \mathbf{r}}{\partial \phi}$.

If $\mathbf{r}$ be the position vector of a point in space, then

$$\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}.$$

(a) We have

$$\frac{\partial \mathbf{r}}{\partial r} = \frac{\partial x}{\partial r} \mathbf{i} + \frac{\partial y}{\partial r} \mathbf{j} + \frac{\partial z}{\partial r} \mathbf{k}$$

$$= \sin \theta \cos \phi \mathbf{i} + \sin \theta \sin \phi \mathbf{j} + \cos \theta \mathbf{k}$$

$$\text{and } \frac{\partial \mathbf{r}}{\partial \theta} = \frac{\partial x}{\partial \theta} \mathbf{i} + \frac{\partial y}{\partial \theta} \mathbf{j} + \frac{\partial z}{\partial \theta} \mathbf{k}$$

$$= r \cos \theta \cos \phi \mathbf{i} + r \cos \theta \sin \phi \mathbf{j} - r \sin \theta \mathbf{k}.$$

$$\frac{\partial \mathbf{r}}{\partial \phi} = \frac{\partial x}{\partial \phi} \mathbf{i} + \frac{\partial y}{\partial \phi} \mathbf{j} + \frac{\partial z}{\partial \phi} \mathbf{k}$$

$$= -r \sin \theta \sin \phi \mathbf{i} + r \sin \theta \cos \phi \mathbf{j} + 0 \cdot \mathbf{k}.$$

(b) Taking the values of $\frac{\partial \mathbf{r}}{\partial r}, \frac{\partial \mathbf{r}}{\partial \theta}$ and $\frac{\partial \mathbf{r}}{\partial \phi}$ from (a) we have

$$\begin{aligned}\frac{\partial \mathbf{r}}{\partial r} \cdot \frac{\partial \mathbf{r}}{\partial \theta} &= (\sin \theta \cos \phi \mathbf{i} + \sin \theta \sin \phi \mathbf{j} + \cos \theta \mathbf{k}) \cdot (r \cos \theta \cos \phi \mathbf{i} \\ &\quad + r \cos \theta \sin \phi \mathbf{j} - r \sin \theta \mathbf{k}) \\ &= r \sin \theta \cos \theta \cos^2 \phi + r \sin \theta \cos \theta \sin^2 \phi - r \sin \theta \cos \theta \\ &= r \sin \theta \cos \theta - r \sin \theta \cos \theta \\ &= 0.\end{aligned}$$

$$\text{Similarly } \frac{\partial \mathbf{r}}{\partial \theta} \cdot \frac{\partial \mathbf{r}}{\partial \phi} = 0 \text{ and } \frac{\partial \mathbf{r}}{\partial \phi} \cdot \frac{\partial \mathbf{r}}{\partial r} = 0.$$

Since all the scalar products vanish, it follows that the vectors $\frac{\partial \mathbf{r}}{\partial r}$, $\frac{\partial \mathbf{r}}{\partial \theta}$, $\frac{\partial \mathbf{r}}{\partial \phi}$ are mutually orthogonal.

(c) We have

$$r^2 = x^2 + y^2 + z^2, \quad \theta = \cos^{-1} \frac{z}{\sqrt{x^2 + y^2 + z^2}} \text{ and } \phi = \tan^{-1} \frac{y}{x}.$$

$$\begin{aligned}\therefore \nabla r &= \frac{\partial r}{\partial x} \mathbf{i} + \frac{\partial r}{\partial y} \mathbf{j} + \frac{\partial r}{\partial z} \mathbf{k} \\ &= \frac{x}{r} \mathbf{i} + \frac{y}{r} \mathbf{j} + \frac{z}{r} \mathbf{k} = \sin \theta \cos \phi \mathbf{i} + \sin \theta \sin \phi \mathbf{j} + \cos \theta \mathbf{k} \\ \nabla \theta &= \frac{\partial \theta}{\partial x} \mathbf{i} + \frac{\partial \theta}{\partial y} \mathbf{j} + \frac{\partial \theta}{\partial z} \mathbf{k} = \frac{\cos \theta \cos \phi}{r} \mathbf{i} + \frac{\cos \theta \sin \phi}{r} \mathbf{j} \\ &\quad - \frac{\sin \theta}{r} \mathbf{k}\end{aligned}$$

$$\begin{aligned}\nabla \phi &= \frac{\partial \phi}{\partial x} \mathbf{i} + \frac{\partial \phi}{\partial y} \mathbf{j} + \frac{\partial \phi}{\partial z} \mathbf{k} \\ &= -\frac{\sin \phi}{r \sin \theta} \mathbf{i} + \frac{\cos \phi}{r \sin \theta} \mathbf{j} + 0 \cdot \mathbf{k}.\end{aligned}$$

$$\begin{aligned}\text{So that } [\nabla r \nabla \theta \nabla \phi] &= (\nabla r \times \nabla \theta) \cdot \nabla \phi \\ &= \left(-\frac{\sin \phi}{r} \mathbf{i} + \frac{\cos \phi}{r} \mathbf{j} + 0 \cdot \mathbf{k} \right) \cdot \left(-\frac{\sin \phi}{r \sin \theta} \mathbf{i} + \frac{\cos \phi}{r \sin \theta} \mathbf{j} + 0 \cdot \mathbf{k} \right) \\ &= \frac{\sin^2 \phi}{r^2 \sin \theta} + \frac{\cos^2 \phi}{r^2 \sin \theta} = \frac{1}{r^2 \sin \theta}.\end{aligned}$$

$$\begin{aligned}\text{Also } \left[\frac{\partial \mathbf{r}}{\partial r}, \frac{\partial \mathbf{r}}{\partial \theta}, \frac{\partial \mathbf{r}}{\partial \phi} \right] &= \left(\frac{\partial \mathbf{r}}{\partial r} \times \frac{\partial \mathbf{r}}{\partial \theta} \right) \cdot \frac{\partial \mathbf{r}}{\partial \phi} \\ &= (-r \sin \phi \mathbf{i} + r \cos \phi \mathbf{j} + 0 \cdot \mathbf{k}) \cdot (-r \sin \theta \sin \phi \mathbf{i} + r \sin \theta \cos \phi \mathbf{j}) \\ &= r^2 \sin \theta \sin^2 \phi + r^2 \sin \theta \cos^2 \phi \\ &= r^2 \sin \theta.\end{aligned}$$

$$\therefore [\nabla r \nabla \theta \nabla \phi] \left[\frac{\partial \mathbf{r}}{\partial r} \frac{\partial \mathbf{r}}{\partial \theta} \frac{\partial \mathbf{r}}{\partial \phi} \right] = 1$$

which shows that ∇r , $\nabla \theta$, $\nabla \phi$ is a set of vectors reciprocal

$$\frac{\partial \mathbf{r}}{\partial r}, \frac{\partial \mathbf{r}}{\partial \theta}, \frac{\partial \mathbf{r}}{\partial \phi}.$$

1.47. INTEGRATION OF A VECTOR

We know that integration is the reverse process to differentiation. Thus if $\mathbf{F}$ and $\mathbf{r}$ be two vector functions such that the derivative of $\mathbf{F}$ with respect to t is equal to $\mathbf{r}$,

$$\frac{d\mathbf{F}}{dt} = \mathbf{r},$$

then $\mathbf{F} + \mathbf{C}$ is called the *indefinite integral* or simply the *integral* of $\mathbf{r}$ with regard to t and is denoted by

$$\int \mathbf{r} dt = \mathbf{F} + \mathbf{C}$$

where the constant vector $\mathbf{C}$ is known as the constant of integration.

In order to apply the integration to definite problems the value of $\mathbf{F}$ may be determined from some initial or geometrical conditions.

The process of finding a vector $\mathbf{F}$ whose derivative with respect to t is equal to $\mathbf{r}$ is known as *integration* of a vector.

In general the nature of the constant of integration is the same as that of the integrand.

We thus write down the values of the following integrals:

$$\int \left(\mathbf{r} \cdot \frac{d\mathbf{s}}{dt} + \mathbf{s} \cdot \frac{d\mathbf{r}}{dt} \right) dt = \mathbf{r} \cdot \mathbf{s} + c \text{ where } c \text{ is a scalar}$$

$$\int 2\mathbf{r} \cdot \frac{d\mathbf{r}}{dt} dt = \int \frac{d\mathbf{r}^2}{dt} dt = \mathbf{r}^2 + c = \mathbf{r} \cdot \mathbf{r} + c, c \text{ being a scalar.}$$

$$\int 2 \frac{d\mathbf{r}}{dt} \cdot \frac{d^2\mathbf{r}}{dt^2} dt = \frac{d\mathbf{r}}{dt} \cdot \frac{d\mathbf{r}}{dt} + c = \left(\frac{d\mathbf{r}}{dt} \right)^2 + c, c \text{ being a scalar}$$

$$\int \mathbf{r} \times \frac{d^2\mathbf{r}}{dt^2} dt = \mathbf{r} \times \frac{d\mathbf{r}}{dt} + \mathbf{c}, \text{ where } \mathbf{c} \text{ is a vector.}$$

$$\int \left(\frac{1}{r} \frac{d\mathbf{r}}{dt} - \frac{d\mathbf{r}}{dt} \cdot \frac{\mathbf{r}}{r^2} \right) dt = \frac{\mathbf{r}}{r} + \mathbf{c} = \hat{\mathbf{r}} + \mathbf{c}, c \text{ being a vector.}$$

When $\mathbf{a}$ is a constant vector, then we have

$$\int \mathbf{a} \times \frac{d\mathbf{r}}{dt} dt = \mathbf{a} \times \mathbf{r} + \mathbf{c}, c \text{ being a vector.}$$

Problem 119. Find $\mathbf{r}$ from the equation $\frac{d^2\mathbf{r}}{dt^2} = \mathbf{a}t + \mathbf{b}$, where $\mathbf{a}$ and $\mathbf{b}$ are known constant vectors; given that both $\mathbf{r}$ and $\frac{d\mathbf{r}}{dt}$ vanish when $t=0$.

The given vector equation is

$$\frac{d^2\mathbf{r}}{dt^2} = \mathbf{a}t + \mathbf{b}.$$

Integrating with regard to t we get

$$\frac{d^2\mathbf{r}}{dt^2} = \mathbf{a} \frac{t^2}{2} + \mathbf{b}t + \mathbf{c} \quad \dots (1)$$

$$\text{Initially when } t=0, \frac{d\mathbf{r}}{dt} = 0,$$

$$\therefore c=0.$$

Thus the equation (1) becomes

$$\frac{d\mathbf{r}}{dt} = a \frac{t^2}{2} + b t.$$

Integrating it again with regard to t ,

$$\mathbf{r} = a \frac{t^3}{6} + b \frac{t^2}{2} + \mathbf{d}$$

when

$$t=0, \mathbf{r}=0,$$

$\therefore$

$$\mathbf{d}=0.$$

Hence

$$\mathbf{r} = \frac{1}{6} a t^3 + \frac{1}{2} b t^2.$$

Problem 120. Solve for $\mathbf{r}$: the equation $\mathbf{a} \times \frac{d^2 \mathbf{r}}{dt^2} = \mathbf{b}$, where $\mathbf{a}$ and $\mathbf{b}$ are constant.

Here

$$\mathbf{a} \times \frac{d^2 \mathbf{r}}{dt^2} = \mathbf{b}.$$

Integrating

$$\mathbf{a} \times \frac{d\mathbf{r}}{dt} = \mathbf{b} t + \mathbf{c}.$$

Integrating again

$$\mathbf{a} \times \mathbf{r} = \frac{1}{2} \mathbf{b} t^2 + \mathbf{c} t + \mathbf{d}$$

where $\mathbf{c}$ and $\mathbf{d}$ are constant vectors.

Problem 121. Given that $\mathbf{r}(t) = 2\mathbf{i} - \mathbf{j} + 2\mathbf{k}$ when $t=2$, and $\mathbf{r}(t) = 4\mathbf{i} - 2\mathbf{j} + 3\mathbf{k}$ when $t=3$.

Show that $\int_2^3 \mathbf{r} \cdot \frac{d\mathbf{r}}{dt} dt = 10$.

$$\begin{aligned} \text{Here } \int_2^3 \mathbf{r} \cdot \frac{d\mathbf{r}}{dt} dt &= \frac{1}{2} \int_2^3 2\mathbf{r} \cdot \frac{d\mathbf{r}}{dt} dt = \frac{1}{2} \left[\mathbf{r}^2 \right]_2^3 \\ &= \frac{1}{2} [(4\mathbf{i} - 2\mathbf{j} + 3\mathbf{k})^2 - (2\mathbf{i} - \mathbf{j} + 2\mathbf{k})^2] \\ &= \frac{1}{2} [16 + 4 + 9 - (4 + 1 + 4)] = 10. \end{aligned}$$

Problem 122. Evaluate the integral $\int_C \mathbf{F} \cdot d\mathbf{r}$

where $\mathbf{F} = c[-3a \sin^2 \theta \cos \theta \mathbf{i} + a(2 \sin \theta - 3 \sin^3 \theta) \mathbf{j} + b \sin 2\theta \mathbf{k}]$ the curve Γ is given by $\mathbf{r} = a \cos \theta \mathbf{i} + a \sin \theta \mathbf{j} + b\theta \mathbf{k}$, θ varying $\pi/4$ to $\pi/2$.

$$\text{Given } \mathbf{r} = a \cos \theta \mathbf{i} + a \sin \theta \mathbf{j} + b\theta \mathbf{k}.$$

$$\therefore \frac{d\mathbf{r}}{d\theta} = -a \sin \theta \mathbf{i} + a \cos \theta \mathbf{j} + b \mathbf{k}.$$

Now

$$\begin{aligned} \int_{\Gamma} \mathbf{F} \cdot d\mathbf{r} &= \int_{\pi/4}^{\pi/2} \mathbf{F} \cdot \frac{d\mathbf{r}}{d\theta} d\theta = \int_{\pi/4}^{\pi/2} c[-3a \sin^2 \theta \cos \theta \mathbf{i} \\ &\quad + a(2 \sin \theta - 3 \sin^3 \theta) \mathbf{j} + b \sin 2\theta \mathbf{k}] \\ &\quad \cdot (-a \sin \theta \mathbf{i} + a \cos \theta \mathbf{j} + b \mathbf{k}) d\theta \end{aligned}$$

$$\begin{aligned} &= c \int_{\pi/4}^{\pi/2} \{3a^2 \sin^3 \theta \cos \theta + a^2(2 \sin \theta - 3 \sin^3 \theta) \cos \theta \\ &\quad + b^2 \sin 2\theta\} d\theta \\ &= c \int_{\pi/4}^{\pi/2} \{a^2(3 \sin^3 \theta \cos \theta + 2 \sin \theta \cos \theta - 3 \sin^3 \theta \cos \theta) \\ &\quad + b^2 \sin 2\theta\} d\theta \\ &= c(a^2 + b^2) \int_{\pi/4}^{\pi/2} \sin 2\theta d\theta = c(a^2 + b^2) \left[-\frac{\cos 2\theta}{2} \right]_{\pi/4}^{\pi/2} \\ &= \frac{c}{2} (a^2 + b^2). \end{aligned}$$

Problem 123. Evaluate $\int_C \mathbf{F} \cdot d\mathbf{r}$ when $\mathbf{F} = xy\mathbf{i} + yz\mathbf{j} + zx\mathbf{k}$, where c is

the curve $\mathbf{r} = t\mathbf{i} + t^2\mathbf{j} + t^3\mathbf{k}$, t varying from -1 to $+1$.

Equating coefficients of like vectors in

$$\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k} = t\mathbf{i} + t^2\mathbf{j} + t^3\mathbf{k},$$

we get the parametric equations of the curve as

$$x=t, y=t^2, z=t^3.$$

Now from $\mathbf{r} = t\mathbf{i} + t^2\mathbf{j} + t^3\mathbf{k}$, we have

$$\frac{d\mathbf{r}}{dt} = \mathbf{i} + 2t\mathbf{j} + 3t^2\mathbf{k}.$$

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= \int_{-1}^1 \mathbf{F} \cdot \frac{d\mathbf{r}}{dt} dt \\ &= \int_{-1}^1 (xy\mathbf{i} + yz\mathbf{j} + zx\mathbf{k}) \cdot (\mathbf{i} + 2t\mathbf{j} + 3t^2\mathbf{k}) dt \\ &= \int_{-1}^1 (xy + 2tyz + 3t^2zx) dt \\ &= \int_{-1}^1 \{t^3 + 2t^6 + 3t^6\} dt = \int_{-1}^1 (t^3 + 5t^6) dt \\ &= \left[\frac{t^4}{4} + \frac{5t^7}{7} \right]_{-1}^1 = \frac{1}{4} - \frac{1}{4} + \frac{5}{7} - \left(-\frac{5}{7} \right) = \frac{10}{7}. \end{aligned}$$

Problem 124. Evaluate $\int_1^2 (\mathbf{A} \cdot \mathbf{B} \times \mathbf{C}) dt$ and $\int_1^2 \mathbf{A} \times (\mathbf{B} \times \mathbf{C}) dt$

$$\text{where } \mathbf{A} = t\mathbf{i} - 3\mathbf{j} + 2t\mathbf{k}, \mathbf{B} = \mathbf{i} - 2\mathbf{j} + 2\mathbf{k} \text{ and } \mathbf{C} = 3\mathbf{i} + t\mathbf{j} - \mathbf{k}.$$

$$\begin{aligned} \text{We have } \mathbf{B} \times \mathbf{C} &= (\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}) \times (3\mathbf{i} + t\mathbf{j} - \mathbf{k}) \\ &= (2 - 2t)\mathbf{i} + (6 + 1)\mathbf{j} + (t + 6)\mathbf{k} \\ &= 2(1 - t)\mathbf{i} + 7\mathbf{j} + (t + 6)\mathbf{k} \end{aligned}$$

$$\begin{aligned} \therefore \mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) &= (t\mathbf{i} - 3\mathbf{j} + 2t\mathbf{k}) \cdot [2(1 - t)\mathbf{i} + 7\mathbf{j} + (t + 6)\mathbf{k}] \\ &= (2t - 2t^2) - 21 + 2t^2 + 12t \\ &= 14t - 21. \end{aligned}$$

$$\begin{aligned}\therefore \int_1^2 [\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C})] dt &= \int_1^2 (14t - 21) dt \\ &= \left[7t^2 - 21t \right]_1^2 \\ &= 28 - 42 - 7 + 21 = 0.\end{aligned}$$

Again where $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = \mathbf{B} (\mathbf{A} \cdot \mathbf{C}) - \mathbf{C} (\mathbf{A} \cdot \mathbf{B})$
 $\mathbf{A} \cdot \mathbf{C} = (t\mathbf{i} - 3\mathbf{j} + 2t\mathbf{k}) \cdot (3\mathbf{i} + t\mathbf{j} - \mathbf{k})$
 $= 3t - 3t - 2t = -2t.$

So that $\mathbf{B} (\mathbf{A} \cdot \mathbf{C}) = (\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}) (-2t)$
 $= -2t\mathbf{i} + 4t\mathbf{j} - 4t\mathbf{k}$

and $(\mathbf{A} \cdot \mathbf{B}) = (t\mathbf{i} - 3\mathbf{j} + 2t\mathbf{k}) \cdot (\mathbf{i} - 2\mathbf{j} + 2\mathbf{k})$
 $= t + 6 + 4t$
 $= (5t + 6).$

So that $\mathbf{C} (\mathbf{A} \cdot \mathbf{B}) = (3\mathbf{i} + t\mathbf{j} - \mathbf{k}) (5t + 6)$
 $= (15t + 18)\mathbf{i} + (5t^2 + 6t)\mathbf{j} - (5t + 6)\mathbf{k}.$

$$\begin{aligned}\text{Thus, } \int_1^2 \mathbf{A} \times (\mathbf{B} \times \mathbf{C}) dt &= \int_1^2 \mathbf{B} (\mathbf{A} \cdot \mathbf{C}) dt - \int_1^2 \mathbf{C} (\mathbf{A} \cdot \mathbf{B}) dt \\ &= \int_1^2 (-2t\mathbf{i} + 4t\mathbf{j} - 4t\mathbf{k}) dt \\ &\quad - \int_1^2 [(15t + 18)\mathbf{i} + (5t^2 + 6t)\mathbf{j} - (5t + 6)\mathbf{k}] dt \\ &= \mathbf{i} \int_1^2 (-18 - 17t) dt + \mathbf{j} \int_1^2 (-2t - 5t^2) dt + \mathbf{k} \int_1^2 (t + 6) dt \\ &= \mathbf{i} \left[-18t - \frac{17}{2}t^2 \right]_1^2 + \mathbf{j} \left[-t^2 - \frac{5}{3}t^3 \right]_1^2 + \mathbf{k} \left[\frac{1}{2}t^2 + 6t \right]_1^2 \\ &= \mathbf{i} \left[-36 + 18 - 34 + \frac{17}{2} \right] + \mathbf{j} \left[-4 + 1 - \frac{40}{3} + \frac{5}{3} \right] + \mathbf{k} \left[2 - \frac{1}{2} + 12 - 6 \right] \\ &= -\frac{87}{2}\mathbf{i} - \frac{44}{3}\mathbf{j} + \frac{15}{2}\mathbf{k}.\end{aligned}$$

1.48. THE LINE INTEGRAL

The integration of a vector along a curve is known as line integral.

Suppose that $\mathbf{F}(\mathbf{r})$ is a continuous vector point function and $\mathbf{r} = \mathbf{r}(s)$ is the given curve. Take any arc C of the given curve between the points A and B , for which length s of the arc has the values a and b respectively. If $\mathbf{t}$ is a unit tangent at a point of the curve, then $\mathbf{F} \cdot \mathbf{t}$ gives the resolute of $\mathbf{F}$ in the direction of the tangent. The definite integral of $\mathbf{F} \cdot \mathbf{t}$ with respect to s , between the limits a and b , is called the line integral of the vector $\mathbf{F}$ along the curve from A to B and is written as

$$\int_a^b \mathbf{F} \cdot \mathbf{t} ds = \text{Lim} \sum_{a}^b \mathbf{F} \cdot \mathbf{t} \delta s \quad \dots$$

But we know that $\mathbf{t} = \frac{d\mathbf{r}}{ds}$

$\therefore$ (1) becomes.

$$\int_A^B \mathbf{F} \cdot d\mathbf{r} = \text{Lim} \sum_A^B \mathbf{F} \cdot \delta \mathbf{r}$$

where A and B are the end points of arc of integration, and $\delta \mathbf{r}$ is the infinitesimal vector, δs parallel to the tangent at the point considered.

The integration round a closed curve is denoted by the symbol $\oint$, while in mechanics this integral is known as the circulation of $\mathbf{F}$ about the closed curve C , being the velocity of the fluid.

Problem 125. Compute $I = \int (x dy - y dx)$ over the

- Straight line $y = x$ from $(0, 0)$ to $(1, 1)$
- Parabola $y = x^2$ from $(0, 0)$ to $(1, 1)$
- Circle $x = a \cos t$, $y = 1 + \sin t$; $-\frac{\pi}{2} \leq t \leq 0$
- Integrate also round the square $(0, 0)$; $(1, 0)$; $(1, 1)$; $(0, 1)$.
- The line integral $I = \int (x dy - y dx)$ over the line

$y = x$, $dy = dx$, is given by

$$\begin{aligned}\int (x dx - y dx) &= \int (x dx - x dx) \text{ limits of } x \text{ are from } 0 \text{ to } 1. \\ &= \int_0^1 0 dx = 0.\end{aligned}$$

- The line integral

$$I = \int (x dy - y dx) \text{ over the parabola } y = x^2, dy = 2x dx$$

- $I = \int (2x^2 dx - x^2 dx)$ limits of x being from 0 to 1

$$\begin{aligned}&= \int_0^1 x^2 dx \\ &= \left[\frac{x^3}{3} \right]_0^1 = \frac{1}{3}.\end{aligned}$$

$$\begin{aligned}(i) \quad I &= \int_{-\pi/2}^0 [\cos^2 t + (4 \sin t) \sin t] dt \text{ as } x = a \cos t \\ &\quad y = 1 + \sin t \\ &= \left[\frac{\pi}{2} - 1 \right]\end{aligned}$$

$$(d) \quad I=0+\int_0^1 dy-\int_1^0 dx+0 \text{ the figure being a square and} \\ \text{integration is along straight lines} \\ =1+1=2.$$

Problem 126. Find the condition that the line integral

$$\int_{r_1}^{r_2} \mathbf{F} \cdot d\mathbf{r} = \int_{t_1}^{t_2} \mathbf{F} \cdot \frac{d\mathbf{r}}{dt} dt$$

taken between the points A and B , is independent of the curve $\mathbf{r}=\mathbf{r}(t)$ joining the points on a smooth curve.

We observe that, if $\int \mathbf{F} \cdot d\mathbf{r}$ is independent of path in a certain region then $\oint \mathbf{F} \cdot d\mathbf{r}=0$ for all closed curves in the region.




Fig. 1.44

For if AP_1BP_2A is a closed curve, then

$$\oint \mathbf{F} \cdot d\mathbf{r} = \int_{AP_1B} \mathbf{F} \cdot d\mathbf{r} + \int_{BP_2A} \mathbf{F} \cdot d\mathbf{r} \\ = \int_{AP_1B} \mathbf{F} \cdot d\mathbf{r} - \int_{AP_2B} \mathbf{F} \cdot d\mathbf{r} \\ = 0.$$

Hence, if $\int_{r_1}^{r_2} \mathbf{F} \cdot d\mathbf{r}$ is independent of path then $\oint \mathbf{F} \cdot d\mathbf{r}=0$.

Conversely, if $\oint \mathbf{F} \cdot d\mathbf{r}=0$ over all closed curves of the region, then $\int \mathbf{F} \cdot d\mathbf{r}$ is the same over any two paths from A to B that do not cross.

Problem 127. If $d\mathbf{r}$ is the infinitesimal vector and $\mathbf{r}=\mathbf{r}(t)$ is the equation of a curve, then evaluate the integrals

$$(a) \quad \int_C \phi \, d\mathbf{r} \quad (b) \quad \int_C \mathbf{F} \cdot d\mathbf{r}$$

along the curve C where $\mathbf{F}$ is a continuous vector and ϕ is a continuous point function.

If $\mathbf{r}=xi+yj+zk$
then $d\mathbf{r}=i \, dx+j \, dy+k \, dz$.

Thus

$$(a) \quad \int_C \phi \, d\mathbf{r} = \int_A^B \phi (i \, dx + j \, dy + k \, dz) \\ = i \int_{x_1}^{x_2} \phi \, dx + j \int_{y_1}^{y_2} \phi \, dy + k \int_{z_1}^{z_2} \phi \, dz$$

where A and B are initial and final points of the curve with co-

ordinates (x_1, y_1, z_1) and (x_2, y_2, z_2) . Thus the integral $\int_C \phi \, d\mathbf{r}$ can be evaluated when y and z are known in terms of x for points on the curve C .

(b) If $\mathbf{F}(x, y, z)=f_1\mathbf{i}+f_2\mathbf{j}+f_3\mathbf{k}$ then as in (a), we find

$$\int_C \mathbf{F} \cdot d\mathbf{r} = f_1 \int_C dx + f_2 \int_C dy + f_3 \int_C dz$$

Problem 128. If $\mathbf{F}=\nabla\phi$ everywhere in a region R of space, defined by $a_1 \leq x \leq a_2, b_1 \leq y \leq b_2, c_1 \leq z \leq c_2$, where $\phi(x, y, z)$ is single valued and has continuous derivative in R , then show that

(i) $\int_A^B \mathbf{F} \cdot d\mathbf{r}$ is independent of path C in R joining A and B .

(ii) The converse of (i) is true

(iii) $\int_A^B \mathbf{F} \cdot d\mathbf{r}=0$ around any closed curve C in R .

We have

$$(i) \quad \text{grad } \phi = \nabla\phi = \left(\mathbf{i} \frac{\partial\phi}{\partial x} + \mathbf{j} \frac{\partial\phi}{\partial y} + \mathbf{k} \frac{\partial\phi}{\partial z} \right) \\ d\mathbf{r} = dx\mathbf{i} + dy\mathbf{j} + dz\mathbf{k} \\ \int_A^B \mathbf{F} \cdot d\mathbf{r} = \int_A^B \left(\mathbf{i} \frac{\partial\phi}{\partial x} + \mathbf{j} \frac{\partial\phi}{\partial y} + \mathbf{k} \frac{\partial\phi}{\partial z} \right) \cdot (dx\mathbf{i} + dy\mathbf{j} + dz\mathbf{k}) \\ = \int_A^B \left(\frac{\partial\phi}{\partial x} dx + \frac{\partial\phi}{\partial y} dy + \frac{\partial\phi}{\partial z} dz \right) \\ = \int_A^B d\phi \\ = \phi(B) - \phi(A) \\ = \phi(x_2, y_2, z_2) - \phi(x_1, y_1, z_1) \quad \dots (1)$$

where (x_1, y_1, z_1) and (x_2, y_2, z_2) being co-ordinates of A and B respectively.

It follows from (1) that the integral depends only on points A and B and not on the path joining them. This is true if only $\phi(x, y, z)$ is single valued at all points.

Note. If $\int_A^B \mathbf{F} \cdot d\mathbf{r}$ is independent of the path C joining A and B then $\mathbf{F}$ is called a conservative field. Thus if $\mathbf{F}=\nabla\phi$ then $\nabla\phi$ is conservative vector field and ϕ is its scalar potential.

(ii) Conversely, if $\int_C \mathbf{F} \cdot d\mathbf{r}$ is independent of the path C joining any two points then there exists a function ϕ such that $\mathbf{F}=\nabla\phi$.

Supposing the line integral to be independent of path, we have

$$\phi(x, y, z) = \int_{(x_1, y_1, z_1)}^{(x, y, z)} \mathbf{F} \cdot d\mathbf{r}$$

$$= \int_{(x_1, y_1, z_1)}^{(x, y, z)} \mathbf{F} \cdot \frac{d\mathbf{r}}{ds} ds.$$

Differentiating we get $\frac{d\phi}{ds} = \mathbf{F} \cdot \frac{d\mathbf{r}}{ds}$.

and also we have $\frac{d\phi}{ds} = \nabla\phi \cdot \frac{d\mathbf{r}}{ds}$.

Subtracting (2) from (3), we find

$$(\nabla\phi - \mathbf{F}) \cdot \frac{d\mathbf{r}}{ds} = 0$$

This will hold independent of $\frac{d\mathbf{r}}{ds}$, if

$$\nabla\phi - \mathbf{F} = 0$$

or $\mathbf{F} = \nabla\phi$.

(iii) We have from (1)

$$\int_A^B \mathbf{F} \cdot d\mathbf{r} = \phi(B) - \phi(A).$$

If the integration is taken round the closed curve i.e. when points B and A coincide then

$$\int_A^B \mathbf{F} \cdot d\mathbf{r} = \int_A^A \mathbf{F} \cdot d\mathbf{r} = \phi(A) - \phi(A) = 0$$

or $\oint \nabla\phi \cdot d\mathbf{r} = 0$.

Problem 129. If C is a simple closed curve in the xy plane enclosing the origin. Show that

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$$

where

$$\mathbf{F} = \frac{iy + jx}{x^2 + y^2}.$$

Given that

$$\mathbf{F} = \frac{-y}{x^2 + y^2} \mathbf{i} + \frac{x}{x^2 + y^2} \mathbf{j}$$

and

$$d\mathbf{r} = dx\mathbf{i} + dy\mathbf{j} \text{ when } \mathbf{r} = x\mathbf{i} + y\mathbf{j}$$

$$\therefore \mathbf{F} \cdot d\mathbf{r} = \frac{-y dx + x dy}{x^2 + y^2}.$$

So that $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \frac{x dy - y dx}{x^2 + y^2}$.

Let us change to polars by putting $x = r \cos \theta$, $y = r \sin \theta$ i.e.,

$$x^2 + y^2 = r^2 \text{ and } \theta = \tan^{-1} \frac{y}{x},$$

so that

$$d\theta = -\frac{1}{1 + \frac{y^2}{x^2}} \cdot \frac{x dy - y dx}{x^2} \\ = -\frac{x dy - y dx}{x^2 + y^2}$$

But the curve being closed, if there is a point P on it such that the limit of θ at P is ϕ (say), then its upper limit will also be ϕ .

$$\oint \mathbf{F} \cdot d\mathbf{r} = \int_{\phi}^{\phi} d\theta \\ = \left[\theta \right]_{\phi}^{\phi} \\ = \phi - \phi = 0.$$

Problem 130. Show that

$$\int_C \left(\frac{-y}{x^2 + y^2} \mathbf{i} + \frac{x}{x^2 + y^2} \mathbf{j} \right) \cdot d\mathbf{r} = 2\pi$$

where C is the circle $x^2 + y^2 = 1$ in the xy plane described in the counter-clockwise direction.

We have

$$\left(\frac{-y}{x^2 + y^2} \mathbf{i} + \frac{x}{x^2 + y^2} \mathbf{j} \right) \cdot [dx\mathbf{i} + dy\mathbf{j}] = \left[\frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy \right] \\ \int_C \left[\frac{-y dx + x dy}{x^2 + y^2} \right] = \int_C \left(\frac{-y dx + x dy}{x^2 + y^2} \right) \\ = \int_0^{2\pi} d\theta \text{ changing to polars} \\ = 2\pi.$$

Problem 131. (a) If $\nabla \times \mathbf{F} = 0$ (i.e. $\mathbf{F}$ is irrotational), prove that $\mathbf{F}$ is conservative.

(b) If $\mathbf{F}$ is conservative field, prove that $\text{curl } \mathbf{F} = \nabla \times \mathbf{F} = 0$ i.e. ($\mathbf{F}$ is irrotational).

We know that if $\int_A^B \mathbf{F} \cdot d\mathbf{r}$ is independent of path C joining A and B then $\mathbf{F}$ is called a conservative field.

Suppose,

$$\mathbf{F} = f_1 \mathbf{i} + f_2 \mathbf{j} + f_3 \mathbf{k}.$$

Then $\nabla \times \mathbf{F} = 0$ gives

$$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ f_1 & f_2 & f_3 \end{vmatrix} = 0$$

$$\left(\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} \right) \mathbf{i} + \left(\frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} \right) \mathbf{j} + \left(\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) \mathbf{k} = 0.$$

This will hold only if,

$$\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} = 0, \frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} = 0, \frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} = 0$$

$$\text{i.e. } \frac{\partial f_3}{\partial y} = \frac{\partial f_2}{\partial z}, \frac{\partial f_1}{\partial z} = \frac{\partial f_3}{\partial x}, \frac{\partial f_2}{\partial x} = \frac{\partial f_1}{\partial y}$$

then, we have to show that $\mathbf{F} = \nabla \phi$.

Now work done $= \int_C \mathbf{F} \cdot d\mathbf{r} = \int_C (f_1 dx + f_2 dy + f_3 dz)$ where C is the path from (x_1, y_1, z_1) to (x, y, z) . Take in particular the path joining along straight line (x_1, y_1, z_1) to (x, y_1, z_1) then to (x, y, z) and then to (x, y, z) and denote the work done along this path by $\phi(x, y, z)$. We thus have

$$\begin{aligned} \phi(x, y, z) &= \int_{x_1}^x f_1(x, y_1, z_1) dx + \int_{y_1}^y f_2(x, y, z_1) dy \\ &\quad + \int_{z_1}^z f_3(x, y, z) dz \end{aligned}$$

Which follows that

$$\frac{\partial \phi}{\partial z} = f_3(x, y, z)$$

$$\frac{\partial \phi}{\partial y} = f_2(x, y, z_1) + \int_{z_1}^z \frac{\partial f_2}{\partial y}(x, y, z) dz$$

$$= f_2(x, y, z_1) + \int_{z_1}^z \frac{\partial f_2}{\partial z}(x, y, z) dz$$

$$= f_2(x, y, z_1) + \left[f_2(x, y, z) \right]_{z_1}^z$$

$$= f_2(x, y, z)$$

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= f_1(x, y_1, z_1) + \int_{y_1}^y \frac{\partial f_1}{\partial x}(x, y, z_1) dy \\ &\quad + \int_{z_1}^z \frac{\partial f_1}{\partial x}(x, y, z) dz \end{aligned}$$

$$= f_1(x, y_1, z_1) + \int_{y_1}^y \frac{\partial f_1}{\partial y}(x, y, z_1) dy$$

$$+ \int_{z_1}^z \frac{\partial f_1}{\partial z}(x, y, z) dz$$

$$= f_1(x, y_1, z_1) + \left[f_1(x, y, z_1) \right]_{y_1}^y + \left[f_1(x, y, z) \right]_{z_1}^z$$

$$= f_1(x, y, z)$$

$$\text{Then } \mathbf{F} = f_1 \mathbf{i} + f_2 \mathbf{j} + f_3 \mathbf{k} = \frac{\partial \phi}{\partial x} \mathbf{i} + \frac{\partial \phi}{\partial y} \mathbf{j} + \frac{\partial \phi}{\partial z} \mathbf{k} = \nabla \phi.$$

(b) If $\mathbf{F}$ is a conservative field

$$\mathbf{F} = \nabla \phi$$

$$\therefore \text{curl } \mathbf{F} = \nabla \times \nabla \phi = 0.$$

Problem 132. If $\mathbf{F} = \cos y \mathbf{i} - x \sin y \mathbf{j} - \cos z \mathbf{k}$, show that the field is conservative.

$\mathbf{F}$ is conservative if

$$\nabla \times \mathbf{F} = 0$$

Here

$$\mathbf{F} = \cos y \mathbf{i} - x \sin y \mathbf{j} - \cos z \mathbf{k}$$

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \cos y & -x \sin y & \cos z \end{vmatrix}$$

$$\begin{aligned} &= \left[\frac{\partial}{\partial y} (\cos z) - \frac{\partial}{\partial z} (-x \sin y) \right] \mathbf{i} + \left[\frac{\partial}{\partial z} (\cos y) - \frac{\partial}{\partial x} (\cos z) \right] \mathbf{j} \\ &\quad + \mathbf{k} \left[\frac{\partial}{\partial x} (-x \sin y) - \frac{\partial}{\partial y} (\cos y) \right] \\ &= 0 \mathbf{i} + 0 \mathbf{j} + (-\sin y + \sin y) \mathbf{k} = 0. \end{aligned}$$

14. THE SURFACE INTEGRALS

We know that the parametric equations of a surface

$$x = x(u, v), y = y(u, v), z = z(u, v)$$

may be combined into a single vector equation

$$\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k} = \mathbf{f}(u, v).$$

A surface $\mathbf{r} = \mathbf{f}(u, v)$ is termed as **smooth** if $\mathbf{f}(u, v)$ possesses continuous first order partial derivatives. In the following discussion we shall assume that the surface under consideration is smooth.

Let a smooth surface S be given by $\mathbf{F}(\mathbf{r}) = \mathbf{r} = \mathbf{f}(u, v)$. Consider S to be the two sided surface one side of which being treated as the positive side. If S is a closed surface the outer surface may be taken as positive surface. Let $\mathbf{n}$ be an outward drawn unit normal vector at any point of the positive side.

Let us divide S into any finite number of elementary sub-surfaces and take a point (x_p, y_p, z_p) in an elementary sub-surface. Let $\mathbf{n}_p$ be the unit normal vector to this sub-surface at (x_p, y_p, z_p) drawn to the positive side of S .

If ΔS_p be the magnitude of the area of the sub-surface under consideration, then the vector area of this sub-surface S_p

$$\Delta \mathbf{S}_p = \mathbf{n}_p \Delta S_p$$

Multiplying both sides by $\mathbf{F}(x_p, y_p, z_p)$, we get




Fig. 1.45

$$\mathbf{F}(x_p, y_p, z_p) \cdot \Delta \mathbf{S}_p = \mathbf{F}(x_p, y_p, z_p) \cdot \mathbf{n}_p \Delta S_p$$

Consider the sum

$$\sum_{p=1}^{p=M} \mathbf{F}(x_p, y_p, z_p) \cdot \Delta \mathbf{S}_p = \sum_{p=1}^{p=M} \mathbf{F}(x_p, y_p, z_p) \cdot \mathbf{n}_p \Delta S_p$$

where summation extends as to include all sub-surfaces of S .

Take the limit of this sum as $M \rightarrow \infty$ in such a way that the largest dimension of each ΔS_p approaches to zero. This limit, if it exists, is termed as the **surface integral** of the normal component of $\mathbf{F}(\mathbf{r})$ over S and is denoted by

$$\int_S \mathbf{F}(\mathbf{r}) \cdot d\mathbf{S} = \int_S \mathbf{F} \cdot \mathbf{n} dS$$

or simply
$$\int_S \mathbf{F} \cdot d\mathbf{S} = \int_S \mathbf{F} \cdot \mathbf{n} dS$$

or
$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S \mathbf{F} \cdot \mathbf{n} dS$$

or
$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S [f_1(x, y, z) dy dz + f_2(x, y, z) dz dx + f_3(x, y, z) dx dy]$$

when
$$\mathbf{F} = f_1(x, y, z) \mathbf{i} + f_2(x, y, z) \mathbf{j} + f_3(x, y, z) \mathbf{k}$$

We sometimes use surface integral $\iint_S \mathbf{F} \cdot \mathbf{n} dS$ to call as the **flux**

F over S. The notation $\oint_S$ is used to indicate integration over closed surface S .

Other forms of surface integrals are:

(a) $\iint_S \phi dS$ (b) $\iint_S \phi \mathbf{n} dS$ (c) $\iint_S \mathbf{F} \times d\mathbf{S}$

where ϕ is a scalar function.

Note 1. If $\mathbf{F} = f_1(x, y, z) \mathbf{i} + f_2(x, y, z) \mathbf{j} + f_3(x, y, z) \mathbf{k}$ then it can be verified that

$$\begin{aligned} \iint_S \mathbf{F} \times d\mathbf{S} &= \mathbf{i} \iint_S (f_2 dx dy - f_3 dz dx) \\ &+ \mathbf{j} \iint_S (f_3 dy dz - f_1 dx dy) + \mathbf{k} \iint_S (f_1 dz dx + f_2 dy dz) \end{aligned}$$

and

$$\begin{aligned} \iint_S \phi \mathbf{n} dS &= \iint_S \phi d\mathbf{S} = \mathbf{i} \iint_S \phi dy dz + \mathbf{j} \iint_S \phi dz dx \\ &+ \mathbf{k} \iint_S \phi dx dy \end{aligned}$$

Note 2. Solenoidal vector function. A vector point function is called **solenoidal** in a region if its flux i.e., the surface integral $\oint \mathbf{F} \cdot d\mathbf{S}$ across

any closed surface in the region is zero.

Problem 133. Supposing that the surface S has projection R on the xy plane show that

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS = \iint_R \mathbf{F} \cdot \mathbf{n} \frac{dx dy}{|\mathbf{n} \cdot \mathbf{k}|}$$

The surface integrals, may be conveniently evaluated by expressing them as double integrals taken over the projected area of the surface on one of the coordinate planes. This is only possible if a line perpendicular to the coordinate plane (chosen) meets the surface in only one point.

Referred to the Fig. 1.45 and §1.49, we know that the surface integral is the limit of sum

$$\sum_{p=1}^{p=M} \mathbf{F}(x_p, y_p, z_p) \cdot \mathbf{n}_p \Delta S_p$$

Now the projection of ΔS_p on xy plane is

$$|\mathbf{n}_p \Delta S_p| \cdot \mathbf{k} \quad \dots (1)$$

(As projected area = original area (cosine of angle between the normal of surface and coordinate plane)]

$$= \Delta S_p |\mathbf{n}_p \cdot \mathbf{k}|$$

Also the area of projection of ΔS_p on xy plane

$$= \Delta x_p \Delta y_p \text{ (i.e. } \delta x_p \cdot \delta y_p) \quad \dots (2)$$

From (1) and (2), we get, $\Delta S_p |\mathbf{n}_p \cdot \mathbf{k}| = \Delta x_p \Delta y_p$

$$\Delta S_p = \frac{\Delta x_p \Delta y_p}{|\mathbf{n}_p \cdot \mathbf{k}|}$$

Thus the limit of sum becomes

$$= \sum_{p=1}^{p=M} \mathbf{F} \cdot \mathbf{n}_p \frac{\Delta x_p \Delta y_p}{|\mathbf{n}_p \cdot \mathbf{k}|}$$

Proceeding to the limit, when $M \rightarrow \infty$, Δx_p and Δy_p both $\rightarrow 0$

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS = \iint_R \mathbf{F} \cdot \mathbf{n} \frac{dx dy}{|\mathbf{n} \cdot \mathbf{k}|}$$

Problem 134. Evaluate

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS$$

$$\mathbf{F} = 18z\mathbf{i} - 12z\mathbf{j} + 3y\mathbf{k}$$

that part of plane $2x + 3y + 6z = 12$ which is located in the first octant.

The projection of the plane $2x + 3y + 6z = 12$

$$z = 0 \text{ plane is}$$

$$2x + 3y = 12$$




Fig. 1.46

is $\nabla(2x+3y+6z) = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (2x+3y+6z)$
 $= 2\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}.$

Unit vector along $2\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}$

$$= \frac{2\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}}{\sqrt{4+9+36}}$$

$$= \frac{2}{7}\mathbf{i} + \frac{3}{7}\mathbf{j} + \frac{6}{7}\mathbf{k}.$$

Now

$$(\mathbf{n} \cdot \mathbf{k}) = \left(\frac{2}{7}\mathbf{i} + \frac{3}{7}\mathbf{j} + \frac{6}{7}\mathbf{k} \right) \cdot \mathbf{k} = \frac{6}{7}.$$

Also

$$\mathbf{F} \cdot \mathbf{n} = (18z\mathbf{i} - 12\mathbf{j} + 3y\mathbf{k}) \cdot \left(\frac{2}{7}\mathbf{i} + \frac{3}{7}\mathbf{j} + \frac{6}{7}\mathbf{k} \right)$$

$$= \frac{36z - 36 + 18y}{7}.$$

But we have,

$$z = \frac{12-2x-3y}{6}$$

$\therefore$

$$\mathbf{F} \cdot \mathbf{n} = \frac{6(12-2x-3y) - 36 + 18y}{7} = \frac{36-12x}{7}$$

Hence

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS = \iint_R \left(\frac{36-12x}{7} \right) \frac{dx dy}{\frac{6}{7}}$$

$$= \iint_R (6-2x) dx dy$$

To integrate it consider the relation

$$2x+3y=12$$

i.e.,

$$y = \frac{12-2x}{3}$$

i.e. referred to Fig. 1.46 the projection of the plane ABC on xy plane is OAB .

By the problem, 133 we have

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS = \iint_R \mathbf{F} \cdot \mathbf{n} \frac{dx dy}{|\mathbf{n} \cdot \mathbf{k}|}$$

Also we know that $\nabla\phi$ is a vector perpendicular to the surface

$$\therefore \phi(x, y, z) = \text{constant}.$$

As such a normal vector perpendicular to the plane

$$2x+3y+6z=12$$

To cover the whole area BOA

$$y \text{ varies from } 0 \text{ to } \frac{12-2x}{3}$$

x varies from 0 to 6.

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS = \int_{x=0}^6 \int_{y=0}^{(12-2x)/3} (6-2x) dx dy$$

$$= \int_{x=0}^6 \left[6y - 2xy \right]_0^{(12-2x)/3} dx$$

$$= \int_{x=0}^6 \left[24 - 4x - 8x + \frac{4}{3}x^2 \right] dx$$

$$= \int_0^6 \left[24 - 12x + \frac{4}{3}x^2 \right] dx$$

$$= \left[24x - \frac{12x^2}{2} + \frac{4}{3} \cdot \frac{x^3}{3} \right]_0^6$$

$$= 144 - 216 + 96 = 24.$$

Problem 135. Evaluate $\int_S (x^3\mathbf{i} + y^3\mathbf{j} + z^3\mathbf{k}) \cdot d\mathbf{S}$

where S is the surface of the sphere

$$x^2 + y^2 + z^2 = 1.$$

The projection of $x^2 + y^2 + z^2 = 1$

on the plane $z=0$ is

$$x^2 + y^2 = 1$$

A normal vector to the surface (1) is

$$\nabla\phi = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (x^2 + y^2 + z^2)$$

$$= 2[\mathbf{i}x + \mathbf{j}y + \mathbf{k}z]$$

This normal vector along $2[\mathbf{i}x + \mathbf{j}y + \mathbf{k}z]$ is given by

$$\mathbf{n} = \frac{2\mathbf{i}x + 2\mathbf{j}y + 2\mathbf{k}z}{2\sqrt{(x^2 + y^2 + z^2)}} \text{ where } x^2 + y^2 + z^2 = 1$$

$$= \mathbf{i}x + \mathbf{j}y + \mathbf{k}z$$

$$\mathbf{n} \cdot \mathbf{k} = (\mathbf{i}x + \mathbf{j}y + \mathbf{k}z) \cdot \mathbf{k} = z.$$

$$\mathbf{F} \cdot \mathbf{n} = (x^3\mathbf{i} + y^3\mathbf{j} + z^3\mathbf{k}) \cdot (\mathbf{i}x + \mathbf{j}y + \mathbf{k}z)$$

$$= x^4 + y^4 + z^4.$$

$$\int_S (x^3\mathbf{i} + y^3\mathbf{j} + z^3\mathbf{k}) \cdot d\mathbf{S} = 2 \iint_R \mathbf{F} \cdot \mathbf{n} \frac{dx dy}{|\mathbf{n} \cdot \mathbf{k}|}$$

$$= 2 \iint_R (x^4 + y^4 + z^4) \frac{dx dy}{z}$$

$$= 2 \iint_R \frac{[x^4 + y^4 + (1 - x^2 - y^2)^2]}{\sqrt{(1 - x^2 - y^2)}} dx dy$$

To cover the whole area of $x^2 + y^2 = 1$

y varies from $-\sqrt{1-x^2}$ to $+\sqrt{1-x^2}$

and x varies from -1 to $+1$.

$$\begin{aligned}
 \therefore \int_S (x^3 \mathbf{i} + y^3 \mathbf{j} + z^3 \mathbf{k}) dS \\
 &= 2 \int_{x=-1}^{x=1} \int_{y=-\sqrt{1-x^2}}^{y=+\sqrt{1-x^2}} \frac{x^4 + y^4 + (1-x^2-y^2)^2}{\sqrt{1-x^2-y^2}} dx dy \\
 &= 8 \int_0^1 \int_0^{\sqrt{1-x^2}} \left[\frac{x^4}{\sqrt{1-x^2-y^2}} + (-\frac{1}{2}) \{ y^3 (-2y)(1-x^2-y^2)^{-3/2} \right. \\
 &\quad \left. + (1-x^2-y^2)^{3/2} \right] dy dx \\
 &= 8 \int_0^1 \left[x^4 \sin^{-1} \frac{y}{\sqrt{1-x^2}} - y^3 (1-x^2-y^2)^{1/2} \right. \\
 &\quad \left. + 3 \int y^2 (1-x^2-y^2)^{1/2} dy + \int (1-x^2-y^2)^{3/2} dy \right]_0^{\sqrt{1-x^2}} dx \\
 &= 8 \int_0^1 \left[\frac{\pi}{2} x^4 - 0 - \left\{ y \cdot (1-x^2-y^2)^{3/2} \right\}_0^{\sqrt{1-x^2}} \right. \\
 &\quad \left. + \int_0^{\sqrt{1-x^2}} (1-x^2-y^2)^{3/2} dy + \int_0^{\sqrt{1-x^2}} (1-x^2-y^2) dy \right] dx \\
 &= 8 \int_0^1 \left[\frac{\pi}{2} x^4 + 2 \int_0^{\sqrt{1-x^2}} (1-x^2-y^2)^{3/2} dy \right] dx \\
 &\quad \text{Put } y = \sqrt{1-x^2} \sin \theta \\
 &\quad dy = \sqrt{1-x^2} \cos \theta d\theta \\
 &= 8 \int_0^1 \left[\frac{\pi}{2} x^4 + 2 \int_0^{\pi/2} (1-x^2)^2 \cdot \cos^4 \theta d\theta \right] dx \\
 &= 8 \int_0^1 \left[\frac{\pi}{2} x^4 + 2(1-x^2)^2 \cdot \frac{\Gamma(\frac{5}{2}) \Gamma(\frac{1}{2})}{2\Gamma(3)} \right] dx \\
 &= 8 \int_0^1 \left[\frac{\pi}{2} x^4 + (1+x^4-2x^2) \cdot \frac{3}{8} \pi \right] dx \\
 &= \pi \int_0^1 (4x^4 + 3 + 3x^4 - 6x^2) dx \\
 &= \pi \int_0^1 (3 - 6x^2 + 7x) dx \\
 &= \pi \left[3x - 2x^3 + \frac{7}{2}x^2 \right]_0^1 \\
 &= \pi \left[3 - 2 + \frac{7}{2} \right] = \frac{5}{2}\pi.
 \end{aligned}$$

Problem 136. Evaluate $\iint_S [(x^3 - yz) dy dz - 2x^2y dz dx + z dx dy]$ over the surface of a cube bounded by the coordinate planes and planes $x=y=z=a$.

Here the surface is the cube in positive octant. To evaluate the integral let us project the given surface on the three coordinate planes

$$\iint_S [(x^3 - yz) dy dz - 2x^2y dz dx + z dx dy]$$

$$= \iint_S (x^3 - yz) dy dz + \iint_S -2x^2y dz dx + \iint_S z dx dy$$

For the first integral

$$\iint_S (x^3 - yz) dy dz$$

Let normal vector to the face $ABB'A'$ in the outward direction

Let unit normal vector to the face $ABB'A'$ in the outward direction is $+\mathbf{i}$.

$$\iint_S (x^3 - yz) dy dz$$

$$= \iint_S (x^3 - yz) dy dz \quad \text{for } x=0 \text{ and } x=a$$

$$= \mathbf{i} \cdot \mathbf{i} \iint_S (x^3 - yz) dy dz + \mathbf{i} \cdot \mathbf{i} \iint_S (x^3 - yz) dy dz$$

$$= - \iint_S (0 - yz) dy dz + \iint_S (a^3 - yz) dy dz$$

$$= \int_{y=0}^a \int_{z=0}^a yz dy dz + \int_{y=0}^a \int_{z=0}^a (a^3 - yz) dy dz$$

$$= \int_0^a \int_0^a a^3 dy dz = a^5.$$

Similarly other parts of integral are

$$- \iint_S 2x^2y dz dx$$

$$= - \left[\iint_{y=0} \mathbf{j} \cdot (-\mathbf{j}) 2x^2y dz dx + \iint_{y=a} \mathbf{j} \cdot \mathbf{j} 2x^2y dz dx \right]$$

$$= - \int_0^a \int_0^a 2ax^2 dx dz = -\frac{2}{3}a^5.$$

$$\iint_S z dx dy = \iint_{z=0} \mathbf{k} \cdot (-\mathbf{k}) z dx dy + \iint_{z=a} \mathbf{k} \cdot \mathbf{k} z dx dy$$

$$= \int_0^a \int_0^a a dx dy = a^3.$$




Fig. 1.47

Hence the value of the integral is $=\frac{1}{3}a^5 + a^3$.

Problem 137. If $\mathbf{F} = 4xz\mathbf{i} - y^2\mathbf{j} + yz\mathbf{k}$ evaluate

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS$$

where S is the surface of the cube bounded by $x=0$, $x=1$, $y=0$, $y=1$, $z=0$, $z=1$.

For the face $ORR'O$




Fig. 1.48

$$\mathbf{n} = -\mathbf{i}, \quad x=0$$

$$\iint_{ORR'O} \mathbf{F} \cdot \mathbf{n} dS$$

$$= \iint (-y^2\mathbf{j} + yz\mathbf{k}) \cdot (-\mathbf{i}) dy dz$$

$$= \int_0^1 \int_0^1 0 dy dz$$

$$= 0.$$

For the face $PQQ'P'$

$$\mathbf{n} = \mathbf{i}, \quad x=1$$

$$\iint_{PQQ'P'} \mathbf{F} \cdot \mathbf{n} dS = \int_0^1 \int_0^1 (4z\mathbf{i} - y^2\mathbf{j} + yz\mathbf{k}) \cdot \mathbf{i} dy dz$$

$$= 4 \int_0^1 \int_0^1 z dy dz = 2.$$

For the face $POO'P'$

$$\mathbf{n} = -\mathbf{j}, \quad y=0$$

$$\iint_{POO'P'} \mathbf{F} \cdot \mathbf{n} dS = \int_0^1 \int_0^1 (4xz\mathbf{i}) \cdot (-\mathbf{j}) dx dz$$

$$= \int_0^1 \int_0^1 0 dx dz = 0.$$

For the face $QRR'Q'$

$$\mathbf{n} = \mathbf{j}, \quad y=1$$

$$\iint_{QRR'Q'} \mathbf{F} \cdot \mathbf{n} dS = \int_0^1 \int_0^1 (4xz\mathbf{i} - \mathbf{j} + z\mathbf{k}) \cdot (\mathbf{j}) dx dz$$

$$= - \int_0^1 \int_0^1 dx dz = -1.$$

For the face $OPQR$

$$\mathbf{n} = -\mathbf{k}, \quad z=0$$

$$\iint_{OPQR} \mathbf{F} \cdot \mathbf{n} dS = \int_0^1 \int_0^1 (-y^2\mathbf{j}) \cdot (-\mathbf{k}) dx dy$$

$$= \int_0^1 \int_0^1 0 dx dy = 0.$$

For the face $O'P'Q'R'$

$$\mathbf{n} = \mathbf{k} \text{ and } z=1$$

$$\iint_{O'P'Q'R'} \mathbf{F} \cdot \mathbf{n} dS = \int_0^1 \int_0^1 (4xz\mathbf{i} - y^2\mathbf{j} + yz\mathbf{k}) \cdot \mathbf{k} dx dy$$

$$= \int_0^1 \int_0^1 y dx dy = \frac{1}{2}.$$

Adding all together,

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS = 2 + 0 - 1 + 0 + \frac{1}{2} + 0 = \frac{5}{2}.$$

Problem 138. If $\mathbf{F} = y\mathbf{i} + (x - 2xz)\mathbf{j} - xy\mathbf{k}$, evaluate

$$\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS$$

where S is the surface of the sphere $x^2 + y^2 + z^2 = a^2$.

Then

$$\mathbf{F} = y\mathbf{i} + (x - 2xz)\mathbf{j} - xy\mathbf{k}.$$

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y & (x - 2yz) & -xy \end{vmatrix}$$

$$= x\mathbf{i} + y\mathbf{j} - 2z\mathbf{k}.$$

To evaluate,

$$\iint_S (x\mathbf{i} + y\mathbf{j} - 2z\mathbf{k}) \cdot \mathbf{n} dS = \int_R \int (x\mathbf{i} + y\mathbf{j} + z\mathbf{k}) \cdot \mathbf{n} \frac{dx dy}{|\mathbf{n} \cdot \mathbf{k}|}$$

Normal vector

$$\nabla \phi = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (x^2 + y^2 + z^2)$$

$$= 2x\mathbf{i} + 2y\mathbf{j} + 2z\mathbf{k}.$$

Unit normal vector $\mathbf{n}$ along $2x\mathbf{i} + 2y\mathbf{j} + 2z\mathbf{k}$

$$= \frac{2(x\mathbf{i} + y\mathbf{j} + z\mathbf{k})}{2\sqrt{(x^2 + y^2 + z^2)}}$$

$$= \frac{x\mathbf{i} + y\mathbf{j} + z\mathbf{k}}{a} \quad \because x^2 + y^2 + z^2 = a^2$$

$$(x\mathbf{i} + y\mathbf{j} - 2z\mathbf{k}) \cdot \mathbf{n} = (x\mathbf{i} + y\mathbf{j} - 2z\mathbf{k}) \cdot \left(\frac{x\mathbf{i} + y\mathbf{j} + z\mathbf{k}}{a} \right)$$

$$= \frac{x^2 + y^2 - 2z^2}{a}$$

$$\mathbf{n} \cdot \mathbf{k} = \left(\frac{x\mathbf{i} + y\mathbf{j} + z\mathbf{k}}{a} \right) \cdot \mathbf{k} = \frac{z}{a}$$

Hence the required integrand

$$I = \int_R \int \left(\frac{x^2 + y^2 - 2z^2}{a} \right) \frac{dx dy}{z/a}$$

$$= \int_{x=-a}^a \int_{y=-\sqrt{(a^2-x^2)}}^{\sqrt{(a^2-x^2)}} \frac{x^2 + y^2 - 2(a^2 - x^2 - y^2)}{\sqrt{(a^2 - x^2 - y^2)}} dx dy$$

as $x^2 + y^2 + z^2 = a^2$

$$= \int_{x=-a}^a \int_{y=-\sqrt{(a^2-x^2)}}^{\sqrt{(a^2-x^2)}} \frac{3(x^2 + y^2) - 2a^2}{\sqrt{(a^2 - x^2 - y^2)}} dx dy$$

put $x = r \cos \theta$, $y = r \sin \theta$.

$$= \int_{\theta=0}^{2\pi} \int_{r=0}^a \frac{3r^2 - 2a^2}{\sqrt{(a^2 - r^2)}} r dr d\theta$$

$$= \int_{\theta=0}^{2\pi} \int_{r=0}^a \frac{3(r^2 - a^2) + a^2}{\sqrt{(r^2 - a^2)}} r d\theta dr$$

$$= \int_{\theta=0}^{2\pi} \int_{r=0}^a \left[-3r \sqrt{(r^2 - a^2)} + \frac{a^2 r}{\sqrt{(r^2 - a^2)}} \right] d\theta dr$$

$$= \int_{\theta=0}^{2\pi} \left[(a^2 - r^2)^{3/2} - a^2 \sqrt{(a^2 - r^2)} \right]_0^a d\theta$$

$$= \int_{\theta=0}^{2\pi} [a^3 - a^3] d\theta$$

$$= 0.$$

Problem 139. Evaluate

$$(a) \int_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS$$

$$(b) \int_S \phi \mathbf{n} dS$$

where $\mathbf{F} = (x+2y)\mathbf{i} - 3z\mathbf{j} + x\mathbf{k}$ and $\phi = 4x+3y-2z$ and S is the surface $2x+y+2z=6$ bounded by $x=0$, $x=1$, $y=0$, $y=2$.

(a) Given

$$\mathbf{F} = (x+2y)\mathbf{i} - 3z\mathbf{j} + x\mathbf{k}$$

$$\therefore \nabla \times \mathbf{F} = \mathbf{i} \left[\frac{\partial}{\partial x} (-3z) - \frac{\partial}{\partial z} (x+2y) \right] + \mathbf{j} \left[\frac{\partial}{\partial z} (x+2y) - \frac{\partial}{\partial x} (-3z) \right] + \mathbf{k} \left[\frac{\partial}{\partial x} (x+2y) - \frac{\partial}{\partial y} (-3z) \right]$$

$$= 3\mathbf{i} - \mathbf{j} - 2\mathbf{k}.$$

Normal vector to the surface $2x+y+2z=6$ is

$$\nabla(2x+y+2z) = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (2x+y+2z)$$

$$= 2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

Unit normal vector $\mathbf{n}$ along $2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$

$$= \frac{2\mathbf{i} + \mathbf{j} + 2\mathbf{k}}{\sqrt{(4+1+4)}}$$

$$= \frac{2\mathbf{i} + \mathbf{j} + 2\mathbf{k}}{3}$$

$$(\nabla \times \mathbf{F}) \cdot \mathbf{n} = (3\mathbf{i} - \mathbf{j} - 2\mathbf{k}) \cdot \left(\frac{2\mathbf{i} + \mathbf{j} + 2\mathbf{k}}{3} \right)$$

$$= \frac{1}{3} [6 - 1 - 4]$$

$$= \frac{1}{3}$$

$$\mathbf{n} \cdot \mathbf{k} = \left[\frac{2\mathbf{i} + \mathbf{j} + 2\mathbf{k}}{3} \right] \cdot \mathbf{k}$$

$$= \frac{2}{3}$$

Projecting the given surface on the plane $z=0$, we find:

$$\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS = \int_R \int (\nabla \times \mathbf{F}) \cdot \mathbf{n} \frac{dx dy}{|\mathbf{n} \cdot \mathbf{k}|}$$

$$= \int_R \int \frac{1}{3} \cdot \frac{dx dy}{\frac{2}{3}}$$

$$= \frac{1}{2} \int_{x=0}^1 \int_{y=0}^2 dx dy$$

$$= \frac{2}{2} = 1.$$

$$(b) \int_S \phi \mathbf{n} dS = \int_R \int \phi \mathbf{n} \frac{dx dy}{|\mathbf{n} \cdot \mathbf{k}|}$$

$$= \int_R \int (4x+3y-2z) \frac{(2\mathbf{i} + \mathbf{j} + 2\mathbf{k})}{3} \frac{dx dy}{\frac{2}{3}}$$

$$= \frac{1}{2} \int_R \int (4x+3y-6+2x+y) (2\mathbf{i} + \mathbf{j} + \mathbf{k}) dx dy$$

$$= \frac{1}{2} \int_{x=0}^1 \int_{y=0}^2 [6x+4y-6] [2\mathbf{i} + \mathbf{j} + \mathbf{k}] dx dy$$

$$= \frac{1}{2} \int_{x=0}^1 [12x+8-12] [2\mathbf{i} + \mathbf{j} + \mathbf{k}] dx dy$$

$$= \frac{1}{2} \int_{x=0}^1 (12x-4) (2\mathbf{i} + \mathbf{j} + \mathbf{k}) dx dy$$

$$= \frac{1}{2} [6-4] [2\mathbf{i} + \mathbf{j} + \mathbf{k}] dx dy = 2\mathbf{i} + \mathbf{j} + \mathbf{k}.$$

1.50. THE VOLUME INTEGRALS

Let $\mathbf{F}(\mathbf{r})$ be a continuous vector point function and V a region enclosed by a surface

$$\mathbf{r} = \mathbf{f}(u, v).$$




Fig. 1.49

in the region. The limit of the sum when $N \rightarrow \infty$ in such a way that the dimensions of ΔV_k tend to zero, if it exists, is denoted by the symbol $\int_V \mathbf{F}(\mathbf{r}) dV$ or simply $\int_V \mathbf{F} dV$ or $\iiint_V \mathbf{F} dV$ and termed the volume integral or space integral.

Its cartesian equivalence is

$$\int_V \mathbf{F} dV = \mathbf{i} \iiint_V F_1 dx dy dz + \mathbf{j} \iiint_V F_2 dx dy dz + \mathbf{k} \iiint_V F_3 dx dy dz$$

If ϕ is a continuous scalar-point function, in V , then,

$$\iiint_V \phi dV$$

is also known as volume integral or space integral.

Problem 140. Evaluate $\iiint_V \mathbf{F} dV$

where

$$\mathbf{F} = 2xz\mathbf{i} - x\mathbf{j} + y^2\mathbf{k}$$

and V is the region bounded by the surfaces

$$x=0, y=0, y=6, z=x^2, z=4.$$

The given solid is a parabolic cylinder with its axis parallel to the z -axis. The part of volume to be determined is shown in figure 1.50.

If we sub-divide the given volume into a large number of elementary cubes of volume δV , then the volume integral is

Let us subdivide the region V into N cubes having volumes

$$\Delta V_k = \Delta x_k \cdot \Delta y_k \cdot \Delta z_k \quad k=1, 2, 3, \dots, N$$

as shown in figure 1.49.

Take a point (x_k, y_k, z_k) within this cube and consider the sum

$$\sum_{k=1}^N \mathbf{F}(x_k, y_k, z_k) \Delta V_k$$

taken over all possible

$$\begin{aligned} & \iiint_V \mathbf{F} dV \\ &= \iiint_V (2xz\mathbf{i} - x\mathbf{j} + y^2\mathbf{k}) dx dy dz \\ &= \mathbf{i} \iiint_V 2xz dx dy dz \\ &\quad - \mathbf{j} \iiint_V x dx dy dz \\ &\quad + \mathbf{k} \iiint_V y^2 dx dy dz. \end{aligned}$$

Now to cover the whole volume,

x varies from 0 to the line in which $z=x^2$ meets the plane $z=4$ i.e., from 0 to $x^2=4$ or $x=2$.

And y varies from 0 to the plane $y=6$ i.e. limits of y are from 0 to 6. Also z varies from x^2 to 4.

Thus

$$\begin{aligned} \iiint_V \mathbf{F} dV &= 2\mathbf{i} \int_{x=0}^2 \int_{y=0}^6 \int_{z=x^2}^4 xz dx dy dz \\ &\quad - \mathbf{j} \int_{x=0}^2 \int_{y=0}^6 \int_{z=x^2}^4 x dx dy dz \\ &\quad + \mathbf{k} \int_{x=0}^2 \int_{y=0}^6 \int_{z=x^2}^4 y^2 dx dy dz \\ &= 2\mathbf{i} \int_{x=0}^2 \int_{y=0}^6 x \left[\frac{z^2}{2} \right]_{x^2}^4 dx dy \\ &\quad - \mathbf{j} \int_{x=0}^2 \int_{y=0}^6 x \left[z \right]_{x^2}^4 dx dy \\ &\quad + \mathbf{k} \int_{x=0}^2 \int_{y=0}^6 y^2 \left[z \right]_{x^2}^4 dx dy \\ &= 2\mathbf{i} \int_{x=0}^2 \int_{y=0}^6 (16x - x^5) dx dy \\ &\quad - \mathbf{j} \int_{x=0}^2 \int_{y=0}^6 (4x - x^3) dx dy \\ &\quad + \mathbf{k} \int_{x=0}^2 \int_{y=0}^6 (4 - x^2) y^2 dx dy \\ &= \mathbf{i} \int_{x=0}^2 [96x - 6x^5] dx - \mathbf{j} \int_{x=0}^2 [24x - 6x^3] dx \\ &\quad + \mathbf{k} \int_{x=0}^2 72(4 - x^2) dx \end{aligned}$$




Fig. 1.50

$$= \mathbf{i} \left[48x - x^6 \right]_0^2 - \mathbf{j} \left[12x - \frac{2}{3}x^4 \right]_0^2 + \mathbf{k} \left[4x - \frac{x^3}{3} \right]_0^2$$

$$= 120\mathbf{i} - 24\mathbf{j} + 384\mathbf{k}.$$

Problem 141. Let $\phi = 45x^2y$ and let V denote the closed region bounded by the planes $4x + 2y + z = 8$, $x = 0$, $y = 0$, $z = 0$ evaluate

$$\iiint_V \phi \, dV$$

Referred to the figure 1.49 we have,

$$\iiint_V \phi \, dV = \int_V 45x^2y \, dx \, dy \, dz$$

z varies from 0 to $8 - 4x - 2y$

y varies from 0 to $4 - 2x$

x varies from 0 to 2.

$$\therefore \iiint_V \phi \, dV = \int_{x=0}^2 \int_{y=0}^{4-2x} \int_{z=0}^{8-4x-2y} 45x^2y \, dx \, dy \, dz$$

$$= \int_{x=0}^2 \int_{y=0}^{4-2x} 45x^2y (8 - 4x - 2y) \, dx \, dy$$

$$= 45 \int_{x=0}^2 x^2 \left[4y^2 - 2xy^2 - \frac{2}{3}y^3 \right]_0^{4-2x} dx$$

$$= 45 \int_{x=0}^2 \frac{x^2}{3} [12(4-2x)^2 - 6x(4-2x)^2 - 2(4-2x)^3] dx$$

$$= 45 \int_{x=0}^2 \frac{x^2}{3} (4-2x)^2 [12 - 6x - 2(4-2x)] dx$$

$$= 45 \int_{x=0}^2 \frac{x^2}{3} (4-2x)^3 dx$$

$$= 15 \int_0^2 x^2 [64 - 32x + 16x^2 - 8x^3] dx$$

$$= 15 \int_0^2 [64x^2 - 32x^3 + 16x^4 - 8x^5] dx$$

$$= 15 \left[\frac{64x^3}{3} - \frac{32x^4}{4} + \frac{16x^5}{5} - \frac{8x^6}{6} \right]_0^2$$

$$= 15 \left[\frac{64 \times 8}{3} - 128 + \frac{16 \times 32}{5} - \frac{4 \times 64}{3} \right]$$

$$= [64 \times 8 \times 5 - 128 \times 15 + 16 \times 32 \times 3 - 4 \times 64 \times 5]$$

$$= 128.$$

Problem 142. Evaluate

$$(a) \iiint_V (\nabla \cdot \mathbf{F}) \, dV. \quad (b) \iiint_V (\nabla \times \mathbf{F}) \, dV.$$

where V is the closed region bounded by the planes $x = 0$, $y = 0$, $z = 0$ and $2x + 2y + z = 4$,

$$\mathbf{F} = (2x^2 - 3z) \mathbf{i} + 2xy \mathbf{j} - 4x \mathbf{k}.$$

$$\mathbf{F} = (2x^2 - 3z) \mathbf{i} - 2xy \mathbf{j} - 4x \mathbf{k}$$

$$\nabla \cdot \mathbf{F} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) [(2x^2 - 3z) \mathbf{i} - 2xy \mathbf{j} - 4x \mathbf{k}]$$

$$= \left[\frac{\partial}{\partial x} (2x^2 - 3z) + \frac{\partial}{\partial y} (-2xy) + \frac{\partial}{\partial z} (-4x) \right]$$

$$= 4x - 2x$$

$$= 2x.$$

$$\nabla \times \mathbf{F} = \mathbf{i} \left[\frac{\partial}{\partial y} (-4x) - \frac{\partial}{\partial z} (-2xy) \right] + \mathbf{j} \left[\frac{\partial}{\partial z} (2x^2 - 3z) \right.$$

$$\left. - \frac{\partial}{\partial x} (-4x) \right] + \mathbf{k} \left[\frac{\partial}{\partial x} (-2xy) - \frac{\partial}{\partial y} (2x^2 - 3z) \right]$$

$$= \mathbf{i} [0] + \mathbf{j} [-3 + 4] + \mathbf{k} [-2y] = \mathbf{j} - 2y \mathbf{k}.$$

z varies from 0 to $4 - 2x - 2y$

y varies from 0 to $2 - x$

x varies from 0 to 2.

$$\iiint_V (\nabla \cdot \mathbf{F}) \, dV = \iiint_V 2x \, dx \, dy \, dz$$

$$= \int_{x=0}^2 \int_{y=0}^{2-x} \int_{z=0}^{4-2x-2y} 2x \, dx \, dy \, dz$$

$$= \int_{x=0}^2 \int_{y=0}^{2-x} 2x (4 - 2x - 2y) \, dx \, dy$$

$$= \int_{x=0}^2 2x [(4-2x)(2-x) - (2-x)^2] dx$$

$$= \int_{x=0}^2 2x [8 - 4x - 4x + 2x^2 - 4 + 4x - x^2] dx$$

$$= \int_{x=0}^2 2x [4 - 4x + x^2] dx$$

$$= 2 \int_0^2 [4x - 4x^2 + x^3] dx$$

$$= 2 \left[\frac{4x^2}{2} - \frac{4x^3}{3} + \frac{x^4}{4} \right]_0^2$$

$$= 2 \left[8 - \frac{32}{3} + 4 \right] = \frac{8}{3}.$$

$$\iiint_V (\nabla \times \mathbf{F}) \, dV = \iiint_V [\mathbf{j} - 2y \mathbf{k}] \, dx \, dy \, dz$$

$$= \int_{x=0}^2 \int_{y=0}^{2-x} \int_{z=0}^{4-2x-2y} [\mathbf{j} - 2y \mathbf{k}] \, dx \, dy \, dz$$

$$\begin{aligned}
&= \int_0^2 \int_0^{2-x} (\mathbf{j} - 2y\mathbf{k}) (4 - 2x - 2y) dx dy \\
&= \int_0^2 \left[\mathbf{j} \left\{ (4-2x)y - y^2 \right\} - \mathbf{k} \left\{ (4-2x)y^2 - \frac{2}{3}y^3 \right\} \right]_0^{2-x} dx \\
&= \int_0^2 \left[\mathbf{j} \left\{ (4-2x)(2-x) - (2-x)^2 \right\} - \mathbf{k} \left\{ (4-2x)(2-x)^2 - \frac{2}{3}(2-x)^3 \right\} \right] dx \\
&= \int_0^2 \left[\mathbf{j} \left\{ 8-8x+2x^2-4-x^2+4x \right\} - \mathbf{k} \left\{ 16-24x+12x^2-2x^3+\frac{4}{3}(8-12x+6x^2-x^3) \right\} \right] dx \\
&= \int_0^2 \left[\mathbf{j} \left\{ 4-4x+x^2 \right\} - \frac{\mathbf{k}}{3} \left\{ 16-24x+12x^2-2x^3 \right\} \right] dx \\
&= \left[\mathbf{j} \left\{ 4x-2x^2+\frac{x^3}{3} \right\} - \frac{\mathbf{k}}{3} \left\{ 16x-12x^2+4x^3-\frac{1}{2}x^4 \right\} \right]_0^2 \\
&= \mathbf{i} \left\{ 8-8+\frac{8}{3} \right\} - \frac{\mathbf{k}}{3} \left\{ 32-48+32-8 \right\} \\
&= \frac{8}{3}\mathbf{j} - \frac{8}{3}\mathbf{k} = \frac{8}{3}(\mathbf{j} - \mathbf{k}).
\end{aligned}$$

1.51. GAUSS' DIVERGENCE THEOREM

(Agra, 1958, 60)

This theorem gives us a powerful device to transform the volume integral into surface integral and its statement is:

If $\mathbf{F}$ is a continuous differentiable vector-point function and S is a closed surface enclosing a volume V , then

$$\int_S \mathbf{F} \cdot \mathbf{n} dS = \int_V \text{div } \mathbf{F} dV.$$

where $\mathbf{n}$ is the unit normal drawn outward.

In other words, "The normal surface integral of a function $\mathbf{F}$ over the boundary of a closed region is equal to the space integral of divergence of $\mathbf{F}$ taken throughout the enclosed space."




Fig. 1.51

Taking $\mathbf{i}, \mathbf{j}, \mathbf{k}$ as the unit vectors along the axes of x, y and z respectively, we have

$$\mathbf{F} = F_1(x, y, z)\mathbf{i} + F_2(x, y, z)\mathbf{j} + F_3(x, y, z)\mathbf{k}$$

F_1, F_2, F_3 and their derivatives in any direction being assumed to be uniform, finite and continuous.

Suppose S is a closed surface such that any line parallel to a coordinate axis cuts it at the most in two points. Let the z coordinates of these two points be

$$z = f_1(x, y), \quad z = f_2(x, y) \text{ respectively.}$$

As such the lower and upper

surfaces S_2 and S_1 of S are given by

$$z = f_2(x, y) \text{ and } z = f_1(x, y) \text{ respectively.}$$

Now consider the integral

$$\begin{aligned}
\iiint_V \frac{\partial F_3}{\partial z} dx dy dz &= \iint_R \left[\int_{f_2}^{f_1} \frac{\partial F_3}{\partial z} dz \right] dx dy \\
&\quad (R \text{ being projection of } S \text{ on } xy \text{ plane}) \\
&= \iint_R [F_3(x, y, f_1) - F_3(x, y, f_2)] dx dy \\
&= \iint_R [F_3(x, y, f_1) - F_3(x, y, f_2)] dx dy \\
&= \iint_R F_3(x, y, f_1) dx dy - \iint_R F_3(x, y, f_2) dx dy.
\end{aligned}$$

For S_1 , we have

$$dx dy = \cos \gamma_1 dS_1 = \mathbf{k} \cdot \mathbf{n}_1 dS_1$$

$\mathbf{n}_1$ is a unit normal vector to the surface dS_1 in outward direction.

For S_2 , we have

$$dx dy = \cos \gamma_2 dS_2 = -\mathbf{k} \cdot \mathbf{n}_2 dS_2$$

$\mathbf{n}_2$ is a unit normal vector to the surface dS_2 in outward direction.

$$\begin{aligned}
\iint_R F_3(x, y, f_1) dx dy &= \iint_{S_1} F_3 \mathbf{k} \cdot \mathbf{n}_1 dS_1 \\
\iint_R F_3(x, y, f_2) dx dy &= -\iint_{S_2} F_3 \mathbf{k} \cdot \mathbf{n}_2 dS_2 \\
\text{Hence} \quad \iint_R F_3(x, y, f_1) dx dy - \iint_R F_3(x, y, f_2) dx dy \\
&= \iint_{S_1} F_3 \mathbf{k} \cdot \mathbf{n}_1 dS_1 + \iint_{S_2} F_3 \mathbf{k} \cdot \mathbf{n}_2 dS_2 \\
&= \iint_S F_3 \mathbf{k} \cdot (\mathbf{n}_1 dS_1 + \mathbf{n}_2 dS_2) \\
&= \iint_S F_3 \mathbf{k} \cdot \mathbf{n} dS; \quad [\because \mathbf{n}S = \mathbf{n}_1 S_1 + \mathbf{n}_2 S_2]
\end{aligned}$$

Consequently

$$\iiint_V \frac{\partial F_3}{\partial z} dx dy dz = \iint_S F_3 \mathbf{k} \cdot \mathbf{n} dS. \quad \dots (1)$$

Analogously projecting S on other coordinate planes, we may find

$$\iiint_V \frac{\partial F_2}{\partial y} dx dy dz = \iint_S F_2 \mathbf{j} \cdot \mathbf{n} dS. \quad \dots (2)$$

$$\iiint_V \frac{\partial F_1}{\partial x} dx dy dz = \iint_S F_1 \mathbf{i} \cdot \mathbf{n} dS. \quad \dots (3)$$

Adding (1), (2) and (3), we get

$$\begin{aligned} & \iiint_V \left(\frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z} \right) dx dy dz \\ &= \iint_S [F_1 \mathbf{i} \cdot \mathbf{n} + F_2 \mathbf{j} \cdot \mathbf{n} + F_3 \mathbf{k} \cdot \mathbf{n}] dS \\ & \iiint_V \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (F_1 \mathbf{i} + F_2 \mathbf{j} + F_3 \mathbf{k}) dx dy dz \\ &= \iint_S [F_1 \mathbf{i} + F_2 \mathbf{j} + F_3 \mathbf{k}] \cdot \mathbf{n} dS \\ & \iiint_V \operatorname{div} \mathbf{F} dV = \iint_S \mathbf{F} \cdot \mathbf{n} dS = \iint_S \mathbf{F} \cdot d\mathbf{S}. \end{aligned}$$

Note. The theorem can be extended to surfaces which are such the lines parallel to the coordinate axes meet them in more than two points. This is also true for multiply connected regions.

1.52. DEDUCTIONS FROM GAUSS' THEOREM

(1) The volume integral of the gradient of a scalar point function can be expressed in terms of the values assumed by the function at the boundary of the region i.e.

$$\iiint_V \nabla \phi dV = \iint_S \phi \mathbf{n} dS \quad (\text{Agra, 1960, p. 10})$$

Gauss' divergence theorem, is

$$\iiint_V \operatorname{div} \mathbf{F} dV = \iint_S \mathbf{F} \cdot \mathbf{n} dS$$

Assume $\mathbf{F} = \phi \mathbf{a}$

where $\mathbf{a}$ is a constant vector. Then it becomes

$$\iiint_V \nabla \cdot (\phi \mathbf{a}) dV = \iint_S \phi \mathbf{a} \cdot \mathbf{n} dS$$

Since $\nabla \cdot (\phi \mathbf{a}) = (\nabla \phi) \cdot \mathbf{a} = \mathbf{a} \cdot (\nabla \phi)$ and $\phi \mathbf{a} \cdot \mathbf{n} = \mathbf{a} \cdot (\phi \mathbf{n})$

$$\therefore \iiint_V \mathbf{a} \cdot (\nabla \phi) dV = \iint_S \mathbf{a} \cdot (\phi \mathbf{n}) dS$$

$$\text{or } \mathbf{a} \cdot \iiint_V \nabla \phi dV = \mathbf{a} \cdot \iint_S \phi \mathbf{n} dS, \mathbf{a} \text{ being constant}$$

$$\text{or } \iiint_V \nabla \phi dV = \iint_S \phi \mathbf{n} dS.$$

(2) The dot product of Gauss's divergence theorem is replaced by cross product i.e.

$$\iiint_V \nabla \times \mathbf{A} dv = \iint_S \mathbf{n} \times \mathbf{A} dS = \iint_S d\mathbf{S} \times \mathbf{A}.$$

Putting $\mathbf{F} = \mathbf{a} \times \mathbf{A}$ in Gauss' divergence theorem, we get,

$$\iiint_V \operatorname{div} \mathbf{F} dV = \iint_S \mathbf{F} \cdot \mathbf{n} dS$$

where $\mathbf{a}$ is a constant vector,

$$\iiint_V \nabla \cdot (\mathbf{a} \times \mathbf{A}) dV = \iint_S (\mathbf{a} \times \mathbf{A}) \cdot \mathbf{n} dS.$$

Since

$$\nabla \cdot (\mathbf{a} \times \mathbf{A}) = -\mathbf{a} \cdot (\nabla \times \mathbf{A})$$

$$(\mathbf{a} \times \mathbf{A}) \cdot \mathbf{n} = \mathbf{a} \cdot (\mathbf{A} \times \mathbf{n}) = (\mathbf{A} \times \mathbf{n}) \cdot \mathbf{a}.$$

$$-\iiint_V \mathbf{a} \cdot (\nabla \times \mathbf{A}) dv = \iint_S \mathbf{a} \cdot (\mathbf{A} \times \mathbf{n}) dS.$$

$$-\mathbf{a} \cdot \iiint_V \nabla \times \mathbf{A} dv = \mathbf{a} \cdot \iint_S \mathbf{A} \times \mathbf{n} dS, \mathbf{a} \text{ being constant}$$

$$-\iiint_V \nabla \times \mathbf{A} dv = \iint_S \mathbf{A} \times \mathbf{n} dS$$

$$\iiint_V \nabla \times \mathbf{A} dv = -\iint_S \mathbf{A} \times \mathbf{n} dS = \iint_S \mathbf{n} \times \mathbf{A} dS.$$

Note. Definition of $\operatorname{grad} \phi$, $\operatorname{div} \mathbf{F}$ and $\operatorname{curl} \mathbf{F}$ can be put as

$$(i) \operatorname{grad} \phi = \lim_{V \rightarrow 0} \frac{\int_S \mathbf{n} \phi ds}{V}$$

$$(ii) \operatorname{div} \mathbf{F} = \lim_{V \rightarrow 0} \frac{\int_S \mathbf{n} \cdot \mathbf{F} ds}{V}$$

$$(iii) \operatorname{curl} \mathbf{F} = \lim_{V \rightarrow 0} \frac{\int_S \mathbf{n} \times \mathbf{F} ds}{V}$$

Below prove (i), the other two can be proved on similar

Let us take a point P enclosed in a small region of volume V bounded by a surface S . Then the first deduction of §1.52 gives

$$\int_S \mathbf{n} \phi dS = \int_V \nabla \phi dV = V[(\nabla \phi)_0 + \epsilon]$$

where $(\nabla \phi)_0$ denotes the value of $\nabla \phi$ at P and $\epsilon \rightarrow 0$ as $V \rightarrow 0$

$$\lim_{V \rightarrow 0} \frac{\int_S \mathbf{n} \phi dS}{V} = (\nabla \phi)_0 = \operatorname{grad} \phi.$$

1.53. PHYSICAL INTERPRETATION OF GAUSS' DIVERGENCE THEOREM

The Gauss' Theorem can be stated as

$$\int_S \mathbf{V} \cdot \mathbf{n} dS = \int_V \text{div } \mathbf{V} dV,$$

where the vector point function $\mathbf{V}$ denotes the velocity vector of an incompressible fluid of unit density and S denotes any closed surface drawn in the space of the fluid, enclosing a volume V .

Since the scalar product $\mathbf{V} \cdot \mathbf{n}$ represents the velocity-component at a point of the surface S in the direction of the outward drawn normal, therefore $\mathbf{V} \cdot \mathbf{n} dS$ expresses the amount of fluid flowing out in unit time through the element of surface dS . As such the integral round the surface S , i.e., $\int_S \mathbf{V} \cdot \mathbf{n} dS$ gives the amount of fluid flowing out of

the surface S in unit time. But in order to maintain the continuity of the flow the total amount of fluid flowing outwards must be continually supplied so that inside the region there are sources producing fluid.

Now the $\text{div } \mathbf{V}$ at any point represents the amount of fluid passing through that point per unit time per unit volume. So $\text{div } \mathbf{V}$ may be regarded as the source-intensity of the incompressible fluid at any point. Thus the integration round the volume V , i.e., $\int_V \text{div } \mathbf{V} dV$

denotes the amount of fluid supplied by the sources inside S per unit time.

Hence the equality $\int_S \mathbf{V} \cdot \mathbf{n} dS = \int_V \text{div } \mathbf{V} dV$ is justified.

i.e., the total volume per second of a moving fluid flowing out from a closed surface S is equal to the total volume per second of fluid flowing out from all volume elements in S .

1.54. GAUSS' THEOREM

If S be a closed surface and $\mathbf{r}$ be the position vector of a point (x, y, z) with respect to an origin O , then

$$\int_S \frac{\mathbf{n} \cdot \mathbf{r}}{r^3} dS$$

is equal to 0 or 4π according as O lies outside or inside of S .

First case. When origin O lies outside the closed surface S , the divergence theorem gives

$$\int_S \mathbf{n} \cdot \left(\frac{\mathbf{r}}{r^3} \right) dS = \iiint_V \nabla \cdot \left(\frac{\mathbf{r}}{r^3} \right) dV$$

But

$$\begin{aligned} \nabla \cdot \left(\frac{\mathbf{r}}{r^3} \right) &= \nabla \cdot (r^{-3} \mathbf{1}) \\ &= (\nabla r^{-3}) \cdot \mathbf{r} + (r^{-3}) \nabla \cdot \mathbf{r} \\ &= -3r^{-5} \mathbf{r} \cdot \mathbf{r} - (r^{-3}) (3) \\ &= -3r^{-3} + 3r^{-3} = 0 \end{aligned}$$

$$\nabla \cdot \left(\frac{\mathbf{r}}{r^3} \right) \text{ is zero everywhere inside } V$$

Provided $r \neq 0$ in V i.e., O is outside the closed surface S .

$$\text{Then } \iint_S \frac{\mathbf{n} \cdot \mathbf{r}}{r^3} dS = 0.$$

Second case. When O lies inside the closed surfaces.

Here $\mathbf{F}$ is not defined at the point O within S , so we cannot apply divergence theorem. To overcome this difficulty let us surround O by a small sphere of radius a and surface S_1 , with its centre at O and lying within S . For S_1 outward drawn normal will be directed towards O and function $\mathbf{F}$ will be continuous and differentiable at all points within S and S_1 .

The divergence theorem can now be applied to the two closed surfaces S_1 and S i.e.,

$$\int_S \frac{\mathbf{n} \cdot \mathbf{r}}{r^3} dS = \iint_S \frac{\mathbf{n} \cdot \mathbf{r}}{r^3} dS + \iint_{S_1} \frac{\mathbf{n} \cdot \mathbf{r}}{r^3} dS = \iiint_V \nabla \cdot \left(\frac{\mathbf{r}}{r^3} \right) dV = 0$$

(being outside the region SS_1)

$$\iint_S \frac{\mathbf{n} \cdot \mathbf{r}}{r^3} dS = - \iint_{S_1} \frac{\mathbf{n} \cdot \mathbf{r}}{r^3} dS$$

$$\text{But on } S_1; \quad r=a \quad \text{and} \quad \mathbf{n} = -\frac{\mathbf{r}}{a}$$

$$\frac{\mathbf{n} \cdot \mathbf{r}}{r^3} = \frac{\left(-\frac{\mathbf{r}}{a} \right) \cdot \mathbf{r}}{a^3} = -\frac{\mathbf{r} \cdot \mathbf{r}}{a^4} = -\frac{a^2}{a^4} = -\frac{1}{a^2}.$$

$$\begin{aligned} \text{Hence } \iint_S \frac{\mathbf{n} \cdot \mathbf{r}}{r^3} dS &= \iint_{S_1} \frac{1}{a^2} dS \\ &= \frac{1}{a^2} \iint_{S_1} dS \\ &= \frac{1}{a^2} (4\pi a^2) \\ &= 4\pi. \end{aligned}$$




Fig. 1.52

1.55. TWO GREEN'S IDENTITIES

First Identity. If ϕ and ψ are scalar point functions having continuous derivatives of the second order at least, then

$$\iiint_V (\phi \nabla^2 \psi + \nabla \psi \cdot \nabla \phi) dV = \iint_S (\phi \nabla \psi) \cdot d\mathbf{S}.$$

The divergence theorem is

$$\iiint_V \nabla \cdot \mathbf{F} dV = \iint_S \mathbf{F} \cdot \mathbf{n} dS.$$

Taking

$$\mathbf{F} = \phi \nabla \psi, \text{ we have}$$

$$\iiint_V \nabla \cdot (\phi \nabla \psi) dV = \iint_S (\phi \nabla \psi) \cdot \mathbf{n} dS = \iint_S (\phi \nabla \psi) \cdot d\mathbf{n}.$$

But

$$\begin{aligned} \nabla \cdot (\phi \nabla \psi) &= \phi (\nabla \cdot \nabla \psi) + (\nabla \phi) \cdot (\nabla \psi) \\ &= \phi (\nabla^2 \psi) + (\nabla \phi) \cdot (\nabla \psi). \end{aligned}$$

$$\therefore \iiint_V \nabla \cdot (\phi \nabla \psi) dV = \iiint_V [\phi \nabla^2 \psi + \nabla \phi \cdot \nabla \psi] dV$$

$$\text{or } \iiint_V [\phi \nabla^2 \psi + \nabla \phi \cdot \nabla \psi] dV = \iint_S (\phi \nabla \psi) \cdot d\mathbf{S}$$

Second identity:

$$\iiint_V (\phi \nabla^2 \psi - \psi \nabla^2 \phi) dV = \iint_S (\phi \nabla \psi - \psi \nabla \phi) \cdot d\mathbf{S}$$

Putting $\mathbf{F} = \psi \nabla \phi$, in divergence theorem, we get

$$\iiint_V \nabla \cdot (\psi \nabla \phi) dV = \iint_S (\psi \nabla \phi) \cdot \mathbf{n} dS = \iint_S (\psi \nabla \phi) \cdot d\mathbf{n}$$

But

$$\begin{aligned} \nabla \cdot (\psi \nabla \phi) &= \psi (\nabla \cdot \nabla \phi) + (\nabla \psi) \cdot (\nabla \phi) \\ &= \psi \nabla^2 \phi + (\nabla \psi) \cdot (\nabla \phi) \end{aligned}$$

$$\therefore \iiint_V \nabla \cdot (\psi \nabla \phi) dV = \iiint_V [\psi \nabla^2 \phi + (\nabla \psi) \cdot (\nabla \phi)] dV$$

$$\text{And } \iiint_V [\psi \nabla^2 \phi + (\nabla \psi) \cdot (\nabla \phi)] dV = \iint_S [\psi \nabla \phi] \cdot d\mathbf{S}$$

Green's first identity, is

$$\iiint_V [\phi \nabla^2 \psi + (\nabla \phi) \cdot (\nabla \psi)] dV = \iint_S [\phi \nabla \psi] \cdot d\mathbf{S}.$$

Subtracting (1) from (2), we find

$$\iiint_V [\phi \nabla^2 \psi - \psi \nabla^2 \phi] dV = \iint_S [\phi \nabla \psi - \psi \nabla \phi] \cdot d\mathbf{S}.$$

Problem 143. Verify divergence theorem for $\mathbf{F} = x^2\mathbf{i} + y^2\mathbf{j} + z^2\mathbf{k}$ over the cube $0 \leq x, y, z \leq 1$.

The divergence theorem is

$$\iiint_V \nabla \cdot \mathbf{F} dV = \iint_S \mathbf{F} \cdot \mathbf{n} dS.$$

We have

$$\mathbf{F} = x^2\mathbf{i} + y^2\mathbf{j} + z^2\mathbf{k}$$




Fig. 1.53

$$\begin{aligned} \nabla \cdot \mathbf{F} &= \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (x^2\mathbf{i} + y^2\mathbf{j} + z^2\mathbf{k}) \\ &= 2x + 2y + 2z. \end{aligned}$$

Now the volume integral

$$\begin{aligned} \iiint_V \nabla \cdot \mathbf{F} dV &= \iiint_V 2(x+y+z) dx dy dz \\ &= \int_{x=0}^1 \int_{y=0}^1 \int_{z=0}^1 2(x+y+z) dx dy dz \\ &= 2 \int_{x=0}^1 \int_{y=0}^1 (x+y+\frac{1}{2}) dx dy \\ &= 2 \int_{x=0}^1 (x+\frac{1}{2}+\frac{1}{2}) dx \\ &= 2 [\frac{1}{2} + \frac{1}{2} + \frac{1}{2}] \\ &= 3. \end{aligned} \quad \dots (1)$$

The surface integral is contributed by six faces of the cube, i.e.

$$\begin{aligned} \iint_S \mathbf{F} \cdot \mathbf{n} dS &= \iint_{S_1} \mathbf{F} \cdot \mathbf{n} dS_1 + \iint_{S_2} \mathbf{F} \cdot \mathbf{n} dS_2 + \iint_{S_3} \mathbf{F} \cdot \mathbf{n} dS_3 \\ &\quad (\text{face } x=0) \quad (\text{face } x=1) \quad (\text{face } y=0) \\ &\quad + \iint_{S_4} \mathbf{F} \cdot \mathbf{n} dS_4 + \iint_{S_5} \mathbf{F} \cdot \mathbf{n} dS_5 + \iint_{S_6} \mathbf{F} \cdot \mathbf{n} dS_6 \\ &\quad (\text{face } y=1) \quad (\text{face } z=0) \quad (\text{face } z=1) \end{aligned}$$

For face S_1 , $x=0$, $\mathbf{n} = -\mathbf{i}$

$$\begin{aligned} \iint_{S_1} \mathbf{F} \cdot \mathbf{n} dS_1 &= \iint_{S_1} -x^2 dS_1 \\ &= \int_{y=0}^1 \int_{z=0}^1 0 dS_1 = 0 \end{aligned}$$

For face S_2 , $x=1$, $\mathbf{n} = \mathbf{i}$.

$$\iint_{S_2} \mathbf{F} \cdot \mathbf{n} dS_2 = \iint_{S_2} (x^2\mathbf{i} + y^2\mathbf{j} + z^2\mathbf{k}) \cdot (\mathbf{i}) dS_2$$

$$\begin{aligned}
 &= \iint_{S_2} x^2 dS_2 = \iint_{S_2} dS_2 \\
 &= \int_{y=0}^1 \int_{z=0}^1 dx dz = 1.
 \end{aligned}$$

For face S_3 , $y=0$, $\mathbf{n}=-\mathbf{j}$,

$$\begin{aligned}
 \iint_{S_3} \mathbf{F} \cdot \mathbf{n} dS_3 &= \iint_{S_3} 0 dS_3 \\
 &= 0.
 \end{aligned}$$

For face S_4 , $y=1$, $\mathbf{n}=\mathbf{j}$.

$$\begin{aligned}
 \iint_{S_4} \mathbf{F} \cdot \mathbf{n} dS_4 &= \iint_{S_4} 4 dS_4 \\
 &= \int_{x=0}^1 \int_{z=0}^1 dx dz = 1.
 \end{aligned}$$

For face S_5 , $z=0$, $\mathbf{n}=-\mathbf{k}$

$$\begin{aligned}
 \iint_{S_5} \mathbf{F} \cdot \mathbf{n} dS_5 &= \iint_{S_5} 0 dS_5 \\
 &= 0.
 \end{aligned}$$

For face S_6 , $z=1$, $\mathbf{n}=\mathbf{k}$

$$\begin{aligned}
 \iint_{S_6} \mathbf{F} \cdot \mathbf{n} dS_6 &= \iint_{S_6} dS_6 \\
 &= \int_{x=0}^1 \int_{y=0}^1 dx dy = 1.
 \end{aligned}$$

$$\begin{aligned}
 \text{Therefore, } \iint_S \mathbf{F} \cdot \mathbf{n} dS &= 0+1+0+1+0+1 \\
 &= 3.
 \end{aligned}$$

From (1) and (2) it follows that the volume integral = surface integral.

Problem 144. Evaluate $\iint_S \mathbf{r} \cdot \mathbf{n} dS$

where $\mathbf{r}$ is the position vector of any point on the closed surface.

The divergence theorem is

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS = \iiint_V \nabla \cdot (\mathbf{F}) dV.$$

Given

$$\mathbf{F} = (xi + yj + zk)$$

$$\therefore \iint_S \mathbf{r} \cdot \mathbf{n} dS = \iiint_V \nabla \cdot (xi + yj + zk) dV.$$

$$\begin{aligned}
 \text{Now } \nabla \cdot (xi + yj + zk) &= \left(i \frac{\partial}{\partial x} + j \frac{\partial}{\partial y} + k \frac{\partial}{\partial z} \right) \cdot (xi + yj + zk) \\
 &= 3.
 \end{aligned}$$

$$\text{Therefore } \iint_S \mathbf{r} \cdot \mathbf{n} dS = \iiint_V 3 dV = 3V$$

where V is the volume of solid bounded by closed surface S .

Problem 145. (a) State and prove Gauss' theorem.

(b) If ρ denotes the charge density and $\mathbf{j}$ the current density due to charges, show that the equation $\frac{\partial \rho}{\partial t} + \text{div } \mathbf{j} = 0$ expresses conservation of the total charge. (Agra, 1966)

(c) If N be the outward flux of electrostatic intensity $\mathbf{E}$ through a closed surface S , then

$$N = \int_S \mathbf{E} \cdot d\mathbf{S} = 4\pi Q,$$

where Q denotes the total charge enclosed by the surface S .

According to the definition the outward flux of electric intensity through any closed surface is proportional to the total charge within the surface.

We know that the flux of a vector $\mathbf{E}$ across the surface element $d\mathbf{S}$ is defined as the scalar $\mathbf{E} \cdot d\mathbf{S}$. Therefore, if S be any surface, closed or open, the flux of $\mathbf{E}$ across S is $\int_S \mathbf{E} \cdot d\mathbf{S}$.

Now the electrostatic intensity or the electrostatic field vector, i.e., at a point due to n charges $e_1, e_2, \dots, e_n$ is defined as

$$\mathbf{E} = \sum_{i=1}^n \frac{e_i}{r_i^3} \mathbf{r}_i,$$

where $\mathbf{r}_i$ denotes the position vector of the point relative to the i th charge e_i .

$$\int_S \mathbf{E} \cdot d\mathbf{S} = \int_S \left[\sum_{i=1}^n \frac{e_i \mathbf{r}_i}{r_i^3} \right] \cdot d\mathbf{S} = \sum_{i=1}^n e_i \int_S \frac{\mathbf{r}_i}{r_i^3} \cdot d\mathbf{S}$$

$$= \sum_{i=1}^n e_i \int d\omega_i = \sum_{i=1}^n e_i \omega_i,$$

(since the solid angle $d\omega$ subtended at a point by a surface element of area dS is given by $d\omega = \frac{\mathbf{r}}{r^3} \cdot d\mathbf{S}$)

where ω_i is the solid angle subtended by the closed surface at the i th charge e_i . But $\omega_i = 4\pi$ or 0 according as e_i is inside or outside the surface and $\sum e_i = Q$.

Hence
$$\int_S \mathbf{E} \cdot d\mathbf{S} = 4\pi Q.$$

(b) Suppose that a charge of density ρ is flowing with velocity $\mathbf{V}$.




Fig. 1.54

The charge crossing the surface element dS in a unit time is

$$\rho \mathbf{V} \cdot d\mathbf{S} = \mathbf{j} \cdot d\mathbf{S}$$

where $\mathbf{j}$ is given to be current density vector or conduction current vector.

Now the current say i across any surface S drawn in the medium is given by $i = \int_S \mathbf{j} \cdot d\mathbf{S}$ while the total charge inside a closed surface

S enclosing a volume v is given by $Q = \int_v \rho dv$.

Assuming that there are no sources or sinks inside the surface, the rate at which the charge is decreasing is

$$-\int_v \frac{\partial \rho}{\partial t} dv.$$

Since this is due to the outward flow of charge, we have

$$(1) \text{ and } (2), \quad -\int_v \frac{\partial \rho}{\partial t} dv = \int_S \mathbf{j} \cdot d\mathbf{S} = \int_v \text{div } \mathbf{j} dv$$

by Gauss' divergence theorem

or
$$\int_v \left(\frac{\partial \rho}{\partial t} + \text{div } \mathbf{j} \right) dv = 0.$$

The volume being arbitrary, the integrand must vanish identically and thus we have the equation of continuity or the equation of conservation of charge as

$$\frac{\partial \rho}{\partial t} + \text{div } \mathbf{j} = 0.$$

Problem 146. Prove that $\int_S \nabla \phi \times \nabla \psi \cdot d\mathbf{S} = 0$.

Put $\mathbf{F} = \nabla \phi \times \nabla \psi$ in Gauss' divergence theorem

$$\begin{aligned} \iint_S (\nabla \phi \times \nabla \psi) \cdot d\mathbf{S} &= \iint_S \mathbf{F} \cdot d\mathbf{S} = \iiint_V \nabla \cdot \mathbf{F} dV \\ &= \iiint_V \nabla \cdot (\nabla \phi \times \nabla \psi) dV. \end{aligned}$$

But $\nabla \cdot (\nabla \phi \times \nabla \psi) = \nabla \psi \cdot \text{curl } \nabla \phi - \nabla \phi \cdot \text{curl } \nabla \psi = 0$.

$$\iiint_V \nabla \cdot (\nabla \phi \times \nabla \psi) dV = \iiint_V [\nabla \psi \cdot \text{curl } \nabla \phi - \nabla \phi \cdot \text{curl } \nabla \psi] dV = 0.$$

Since
$$\int_S (\nabla \phi \times \nabla \psi) \cdot d\mathbf{S} = 0$$

Problem 147. (a) If $\mathbf{H} = \text{curl } \mathbf{A}$, prove that

$$\int_S \mathbf{H} \cdot \mathbf{n} dS = 0 \text{ for any closed surface } S.$$

(b) If $\mathbf{n}$ is unit outward normal to any closed surface of area S show

$$\iiint_V \text{div } \mathbf{n} dV = S.$$

(c) Given $\mathbf{H} = \text{curl } \mathbf{A}$. The Divergence Theorem gives

$$\iint_S \mathbf{H} \cdot \mathbf{n} dS = \iiint_V \nabla \cdot \mathbf{H} dV.$$

$$\nabla \cdot \mathbf{H} = \nabla \cdot (\text{curl } \mathbf{A}) = 0$$

$$\iint_S \mathbf{H} \cdot \mathbf{n} dS = 0.$$

(d) The Divergence Theorem gives

$$\begin{aligned} \iiint_V \text{div } \mathbf{n} dV &= \iint_S \mathbf{n} \cdot \mathbf{n} dS \\ &= \iint_S dS = S. \end{aligned}$$

Problem 148. Prove that

$$(a) \int_V \mathbf{F} \cdot \nabla \phi dV = \int_S \phi \mathbf{F} \cdot d\mathbf{S} - \int_V \phi \nabla \cdot \mathbf{F} dV.$$

$$(b) \int_S \nabla \phi \cdot \text{curl } \mathbf{F} dV = \int_S (\mathbf{F} \times \nabla \phi) \cdot d\mathbf{S}.$$

(c) The divergence theorem gives

$$\begin{aligned} \int_V \mathbf{F} \cdot \nabla \phi dV &= \int_V \nabla \cdot (\phi \mathbf{F}) dV - \int_V \phi \nabla \cdot \mathbf{F} dV \\ &= \int_V [\phi \nabla \cdot \mathbf{F} + \mathbf{F} \cdot \nabla \phi] dV - \int_V \phi \nabla \cdot \mathbf{F} dV \\ &= \int_V \mathbf{F} \cdot \nabla \phi dV + \int_V \phi \nabla \cdot \mathbf{F} dV - \int_V \phi \nabla \cdot \mathbf{F} dV \\ &= \int_V \mathbf{F} \cdot \nabla \phi dV. \end{aligned}$$

(b) Applying divergence theorem in R. H. S. of

$$\begin{aligned}\int_V \nabla \phi \operatorname{curl} \mathbf{F} dV &= \int_S (\mathbf{F} \times \nabla \phi) \cdot d\mathbf{S}, \text{ we get} \\ \int_V \nabla \phi \operatorname{curl} \mathbf{F} dV &= \int_V \nabla \cdot (\mathbf{F} \times \nabla \phi) dV \\ &= \int_V [\nabla \phi \cdot \operatorname{curl} \mathbf{F} - \mathbf{F} \cdot \operatorname{curl} \nabla \phi] dV \\ &= \int_V \nabla \phi \cdot \operatorname{curl} \mathbf{F} dV \text{ as } \operatorname{curl} \nabla \phi = 0.\end{aligned}$$

Problem 149. Show that volume enclosed by the surface S is

$$V = \frac{1}{3} \oint_S \nabla \cdot (\mathbf{r}^2) d\mathbf{S}$$

where $\mathbf{r}$ is the position vector to a point of $d\mathbf{S}$.

The divergence theorem gives

$$\begin{aligned}\int_S \nabla \cdot \mathbf{r}^2 d\mathbf{S} &= \int_V \nabla \cdot (\nabla \cdot \mathbf{r}^2) dV \\ &= \int_V \nabla \cdot [2\mathbf{r}^2] dV \\ &= 2 \int_V \nabla \cdot (\mathbf{r}) dV \\ &= 2 \int_V 3 dV \\ &= 6 \int_V dV = 6V\end{aligned}$$

$$\therefore V = \frac{1}{6} \int_S \nabla \cdot \mathbf{r}^2 d\mathbf{S}.$$

Problem 150. Prove the following:

$$(a) \iiint_V \frac{dV}{r^2} = \iint_S \frac{\mathbf{r} \cdot \mathbf{n}}{r^2} d\mathbf{S}.$$

$$(b) \iint_S r^5 \mathbf{n} d\mathbf{S} = \iiint_V 5r^3 \mathbf{r} dV.$$

(a) The divergence theorem gives

$$\begin{aligned}\iint_S \left(\frac{\mathbf{r}}{r^2} \right) \cdot \mathbf{n} d\mathbf{S} &= \iiint_V \nabla \cdot \left(\frac{\mathbf{r}}{r^2} \right) dV \\ &= \iiint_V \nabla \cdot \left(\frac{1}{r^2} \mathbf{r} \right) dV \\ &= \iiint_V \left[\frac{1}{r^2} \nabla \cdot \mathbf{r} + \nabla \left(\frac{1}{r^2} \right) \cdot \mathbf{r} \right] dV \\ &= 3 \iiint_V \frac{dV}{r^2} + \iiint_V -\frac{2}{r^4} \mathbf{r} \cdot \mathbf{r} dV\end{aligned}$$

$$= 3 \iiint_V \frac{dV}{r^2} - 2 \iiint_V \frac{dV}{r^2} = \iiint_V \frac{dV}{r^2}$$

We have

$$\begin{aligned}\iint_S r^5 \mathbf{n} d\mathbf{S} &= \iiint_V \nabla \cdot (r^5 \mathbf{r}) dV \\ &= \iiint_V [5r^4 \mathbf{r} + r^5 \nabla \cdot \mathbf{r}] dV = 5 \iiint_V r^3 \mathbf{r} dV.\end{aligned}$$

Problem 151. Show that Green's second identity can be written as

$$\iiint_V (\phi \nabla^2 \psi - \psi \nabla^2 \phi) dV = \iint_S \left[\phi \frac{\partial \psi}{\partial n} - \psi \frac{\partial \phi}{\partial n} \right] d\mathbf{S}.$$

Green's identity is

$$\begin{aligned}\iiint_V (\phi \nabla^2 \psi - \psi \nabla^2 \phi) dV &= \iint_S [\phi \nabla \psi - \psi \nabla \phi] \cdot d\mathbf{S} \\ &= \iint_S (\phi \nabla \psi - \psi \nabla \phi) \cdot \mathbf{n} d\mathbf{S}\end{aligned}$$

where $\mathbf{n}$ is unit normal vector.

$$\nabla \psi \cdot \mathbf{n} = \frac{\partial \psi}{\partial n}, \quad \nabla \phi \cdot \mathbf{n} = \frac{\partial \phi}{\partial n}.$$

$$\iiint_V (\phi \nabla^2 \psi - \psi \nabla^2 \phi) dV = \iint_S \left(\phi \frac{\partial \psi}{\partial n} - \psi \frac{\partial \phi}{\partial n} \right) d\mathbf{S}.$$

GREEN'S THEOREM IN THE PLANE

Let a closed region in the x - y plane bounded by a simple closed curve C and ϕ and ψ be two continuously differentiable functions of x and y . Then Green's theorem in the plane is stated as

$$\oint_C (\psi dx + \phi dy) \equiv \iint_R \left(\frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \right) dx dy$$

where C is traversed in positive (anti-clockwise) direction.

Let the region R be bounded by single closed curve C having the property that any straight line parallel to the axes cuts C in at most two points and let the parallels AM (say $x=a$), BN (say $x=b$), QF (say $y=c$) and DE (say $y=d$) limit the curve C in the xy plane as shown in Fig. 1.55.

Suppose the equations of the curves AQB and ADB are respectively

$$y = f_1(x) \text{ and } y = f_2(x).$$

$$\iint_R \frac{\partial \psi}{\partial y} dx dy = \int_{x=a}^b \int_{y=f_1(x)}^{f_2(x)} \frac{\partial \psi}{\partial y} dx dy$$




Fig. 1.55

Again if the equations to the curves DAQ and DBQ are

$x=F_1(y)$ and $x=F_2(y)$, respectively, then

$$\begin{aligned} \iint_R \frac{\partial \phi}{\partial x} dx dy &= \int_{y=c}^d \int_{x=F_1(y)}^{F_2(y)} \frac{\partial \phi}{\partial x} dy dx \\ &= \int_c^d \left[\phi(x, y) \right]_{F_1(y)}^{F_2(y)} dy \\ &= \int_c^d [\phi(F_2, y) - \phi(F_1, y)] dy \\ &= \int_c^a \phi(F_2, y) dy + \int_a^c \phi(F_1, y) dy \\ &= \oint_C \phi dx \end{aligned}$$

i.e., $\oint_C \phi dy = \iint_R \frac{\partial \phi}{\partial x} dx dy.$

Adding (1) and (2), we find that

$$\oint_C (\psi dx + \phi dy) = \iint_R \left(\frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \right) dx dy.$$

1.57. VECTOR FORMS OF GREEN'S THEOREM IN PLANE

Vector treatment of Green's Theorem yields two different forms

Form 1. If $\mathbf{f} = \psi\mathbf{i} + \phi\mathbf{j}$ and $\mathbf{r} = x\mathbf{i} + y\mathbf{j}$, then Green's theorem takes the form

$$\oint_C \mathbf{f} \cdot d\mathbf{r} = \iint_R (\nabla \times \mathbf{f}) \cdot \mathbf{k} dS.$$

$$\begin{aligned} &= \int_a^b \left[\psi(x, y) \right]_{f_1(x)}^{f_2(x)} dx \\ &= \int_a^b [\psi(x, f_2) - \psi(x, f_1)] dx \\ &= - \left[\int_a^b \psi(x, f_1) dx \right. \\ &\quad \left. + \int_b^a \psi(x, f_2) dx \right] \\ &= - \oint_C \psi dx \\ \text{i.e. } \oint_C \psi dx &= - \iint_R \frac{\partial \psi}{\partial y} dx dy \end{aligned}$$

Green's theorem in the plane is

$$\oint_C (\psi dx + \phi dy) = \iint_R \left(\frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \right) dx dy \quad \dots (1)$$

$$\mathbf{r} = x\mathbf{i} + y\mathbf{j}$$

$$d\mathbf{r} = dx\mathbf{i} + dy\mathbf{j},$$

$$\psi dx + \phi dy = (\psi\mathbf{i} + \phi\mathbf{j}) \cdot (dx\mathbf{i} + dy\mathbf{j}) = \mathbf{f} \cdot d\mathbf{r} \quad \dots (2)$$

$$\nabla \times \mathbf{f} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \psi & \phi & 0 \end{vmatrix} = -\frac{\partial \phi}{\partial z} \mathbf{i} + \frac{\partial \psi}{\partial z} \mathbf{j} + \left(\frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \right) \mathbf{k}$$

$$(\nabla \times \mathbf{f}) \cdot \mathbf{k} = \frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \quad \dots (3)$$

Substituting values from (2) and (3) in (1), we get

$$\oint_C \mathbf{f} \cdot d\mathbf{s} = \iint_R (\nabla \times \mathbf{f}) \cdot \mathbf{k} dS$$

$ds = dx dy$ represents the element of area.

Form 2. If $\mathbf{f} = \psi\mathbf{i} + \phi\mathbf{j}$, $dS = dx dy$, $\mathbf{g} = \mathbf{f} \times \mathbf{k}$ and $\mathbf{n}$ be the outward normal to C , then Green's theorem gives

$$\oint_C \mathbf{g} \cdot \mathbf{n} ds = \iint_R \nabla \cdot \mathbf{g} dS,$$

where ds is an element of the curve.

If $\mathbf{r}$ be the position vector of any point P and $\mathbf{T}$ be the unit tangent vector to the curve, then

$$\mathbf{r} = x\mathbf{i} + y\mathbf{j}, \text{ so that } d\mathbf{r} = dx\mathbf{i} + dy\mathbf{j} \text{ and}$$

$$\mathbf{T} = \frac{d\mathbf{r}}{ds}.$$

Since $\mathbf{n}$ and $\mathbf{k}$, both being vectors normal to the tangent at any point on the curve, the definition of cross

product yields

$$\mathbf{k} \times \mathbf{n} = \mathbf{T}.$$

Thus, $\psi dx + \phi dy$

$$= (\psi\mathbf{i} + \phi\mathbf{j}) \cdot (dx\mathbf{i} + dy\mathbf{j})$$

$$= \mathbf{f} \cdot d\mathbf{r}$$

$$= \mathbf{f} \cdot \frac{d\mathbf{r}}{ds} ds$$

$$= \mathbf{f} \cdot \mathbf{T} ds, \left(\because \mathbf{T} = \frac{d\mathbf{r}}{ds} \right)$$

$$= \mathbf{f} \cdot (\mathbf{k} \times \mathbf{n}) ds$$

$$= (\mathbf{f} \times \mathbf{k}) \cdot \mathbf{n} ds$$




Fig. 1.56

($\therefore$ in a scalar triple product dot and cross can be interchanged)
 $= \mathbf{g} \cdot \mathbf{n} \, ds.$

Also $\mathbf{g} = \mathbf{f} \times \mathbf{k} = (\psi \mathbf{i} + \phi \mathbf{j}) \times \mathbf{k} = \phi \mathbf{i} - \psi \mathbf{j}$

so that $\nabla \cdot \mathbf{g} = \left(\frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k} \right) \cdot (\phi \mathbf{i} - \psi \mathbf{j})$
 $= \frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y}.$

Substituting these values in Green's theorem i.e.

$$\oint_C (\psi \, dx + \phi \, dy) = \iint_R \left(\frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \right) dx \, dy$$

we get

$$\oint_C \mathbf{g} \cdot \mathbf{n} \, ds = \iint_R \nabla \cdot \mathbf{g} \, dS.$$

Note. Physical Interpretation of Form 1. Vector form 1 of Green's theorem is

$$\oint_C \mathbf{f} \cdot d\mathbf{r} = \iint_R (\nabla \times \mathbf{f}) \cdot \mathbf{k} \, dS.$$

Assuming that $\mathbf{f}$ represents a force field acting on a particle whose position vector is $\mathbf{r}$, the integral $\oint_C \mathbf{f} \cdot d\mathbf{r}$ may be interpreted as the net pressing work done in moving the particle around the closed path and it may be evaluated by the value of $\nabla \times \mathbf{f}$.

As a particular case if $\nabla \times \mathbf{f} = 0$ i.e. if $\mathbf{f} = \nabla \theta$, θ being scalar, then the integral around a closed path is zero. It follows that the work done by a particle in moving from one point of the plane to the other point of the plane is independent of the path traced in moving from one point to another in the plane. In other words this fact can be expressed by saying that the force field is **conservative**.

Conversely, if the integral around a closed path is independent of the path joining any two points in the plane i.e. the integral around the closed path is zero, then $\nabla \times \mathbf{f} = 0$ where $\mathbf{f} = \psi \mathbf{i} + \phi \mathbf{j}$.

i.e. $-\frac{\partial \phi}{\partial z} \mathbf{i} + \frac{\partial \psi}{\partial z} \mathbf{j} + \left(\frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \right) \mathbf{k} = 0.$

Giving $\frac{\partial \phi}{\partial x} = \frac{\partial \psi}{\partial y}.$

1.58. GREEN'S FORMULA

Suppose that $\psi = \frac{1}{r}$ is a scalar point function which has uniform

and continuous derivatives upto the second order in a region enclosed by a closed surface S .

Take a fixed point P within the region, such that r is the distance from P to a variable point of the region. Now since ψ becomes infinite at P , therefore to remove this singularity, we enclose P by small sphere of radius ϵ . The surface of this sphere as S_1 . Clearly in the region V , bounded by S and S_1 , ψ is twice continuously differentiable.




Fig. 1.57

But we know that,

$$\nabla \left(\frac{1}{r} \right) = \nabla \left[\frac{1}{(x^2 + y^2 + z^2)^{1/2}} \right] = -\frac{\mathbf{r}}{r^2}$$

$$\nabla^2 \psi = \nabla \cdot \left(-\frac{\mathbf{r}}{r^2} \right) = 0.$$

Thus, applying Green's identity to the region bounded by S and S_1 we get

$$\begin{aligned} \int_V \left[-\frac{1}{r} \nabla^2 \phi + 0 \right] dv &= \int_S \left[\phi \frac{\partial}{\partial n} \left(\frac{1}{r} \right) - \frac{1}{r} \frac{\partial \phi}{\partial n} \right] dS \\ &+ \int_{S_1} \left[\phi \frac{\partial}{\partial n} \left(\frac{1}{r} \right) - \frac{1}{r} \frac{\partial \phi}{\partial n} \right] dS_1 \quad \dots (1) \end{aligned}$$

In the surface S_1 the direction of unit normal drawn outward from the region considered will be towards P , so that

$$\begin{aligned} \frac{\partial}{\partial n} \left(\frac{1}{r} \right) \text{ on } S_1 &= \left[\nabla \left(\frac{1}{r} \right) \cdot \mathbf{n} \right]_{r=\epsilon} \\ &= \left[\left(-\frac{\mathbf{r}}{r^3} \right) \cdot \left(-\frac{\mathbf{r}}{r} \right) \right]_{r=\epsilon} \\ &= \left[\frac{(\mathbf{r})^2}{r^4} \right]_{r=\epsilon} = \left[\frac{1}{r^2} \right]_{r=\epsilon} \\ &= \frac{1}{\epsilon^2}. \end{aligned}$$

$$\int_{S_1} \phi \frac{\partial}{\partial n} \left(\frac{1}{r} \right) dS = \frac{1}{\epsilon^2} \int_{S_1} \phi \, dS_1.$$

Taking to the limit as $\epsilon \rightarrow 0$, we find

$$\lim_{\epsilon \rightarrow 0} \int_{S_1} \phi \frac{\partial}{\partial n} \left(\frac{1}{r} \right) dS_1 = \frac{4\pi\epsilon^2}{\epsilon^2} \phi(P) = 4\pi \phi(P) \quad \dots (2)$$

$$\lim_{\epsilon \rightarrow 0} \int \frac{1}{r} \frac{\partial \phi}{\partial n} dS = 0 \quad \dots (3)$$

In the limiting case when $\varepsilon \rightarrow 0$, (1) yields with the help of (2) and (3),

$$\int_V -\frac{1}{r} \nabla^2 \phi \, dv = \int_S \left[\phi \frac{\partial}{\partial n} \left(\frac{1}{r} \right) - \frac{1}{r} \frac{\partial \phi}{\partial n} \right] dS + 4\pi \phi(P)$$

Thus
$$4\pi \phi(P) = \int_S \left[\frac{1}{r} \frac{\partial \phi}{\partial n} - \phi \frac{\partial}{\partial n} \left(\frac{1}{r} \right) \right] dS - \int_V \frac{1}{r} \nabla^2 \phi \, dv$$

or
$$4\pi \phi(P) = \int_S \left[\frac{1}{r} \nabla \phi - \phi \nabla \left(\frac{1}{r} \right) \right] \cdot \mathbf{n} \, dS - \int_V \frac{1}{r} \nabla^2 \phi \, dv$$

$$4\pi \phi(P) = \int_S \left[\frac{1}{r} \nabla \phi - \phi \nabla \left(\frac{1}{r} \right) \right] \cdot d\mathbf{S} - \int_V \frac{1}{r} \nabla^2 \phi \, dv$$

which is known as Green's formula.

1.59. POISSON'S EQUATION WITH ITS SOLUTION

Let ϕ be a scalar point function vanishing outside a finite region, then the equation

$$\nabla^2 \phi = -4\pi \rho,$$

is known as Poisson's equation.

Poisson's equation is $\nabla^2 \phi = -4\pi \rho$

Green's formula is

$$4\pi \phi(P) = \int_S \left[\frac{1}{r} \nabla \phi - \phi \nabla \left(\frac{1}{r} \right) \right] \cdot d\mathbf{S} - \int_V \frac{1}{r} \nabla^2 \phi \, dv$$

(for a region bounded by a surface S)

$$= \int_S \left[\frac{1}{r} \nabla \phi - \phi \nabla \left(\frac{1}{r} \right) \right] \cdot d\mathbf{S} + 4\pi \int_V \frac{\rho}{r} \, dv \text{ by (1)}$$

In case the region V tends to infinity S also recedes to infinity.

Supposing that for large values of r , ϕ is of the form $\frac{\lambda}{r}$ which remains bounded, $|\nabla \phi|$ is the form $\frac{\lambda}{r^2}$.

So that
$$\int_S \left[\frac{1}{r} \nabla \phi - \phi \nabla \left(\frac{1}{r} \right) \right] \cdot d\mathbf{S} \rightarrow 0.$$

Then
$$4\pi \phi(P) = 4\pi \int_V \frac{\rho}{r} \, dv$$

or
$$\phi(P) = \int_V \frac{\rho}{r} \, dv.$$

The volume integral being carried over the whole space remains same as the volume integral over the region outside at which ϕ is zero.

Vector equivalent

If $\mathbf{f} = f_1 \mathbf{i} + f_2 \mathbf{j} + f_3 \mathbf{k}$.

$$\mathbf{F} = F_1 \mathbf{i} + F_2 \mathbf{j} + F_3 \mathbf{k}$$

$$\nabla^2 \mathbf{F} = -4\pi \mathbf{f}.$$

Then

$$\mathbf{F}(P) = \int_V \frac{\mathbf{f}}{r} \, dv.$$

$$\nabla^2 \mathbf{F} = -4\pi \mathbf{f} \text{ is equivalent to}$$

$$\nabla^2 F_1 = -4\pi f_1; \nabla^2 F_2 = -4\pi f_2; \nabla^2 F_3 = -4\pi f_3$$

Thus for a point P of the region

$$F_1(P) = \int_V \frac{f_1}{r} \, dv; F_2(P) = \int_V \frac{f_2}{r} \, dv; F_3(P) = \int_V \frac{f_3}{r} \, dv.$$

Imposing suitable conditions on f_1, f_2, f_3 , and multiplying these equations by $\mathbf{i}, \mathbf{j}, \mathbf{k}$ respectively and then adding, we have

$$(\mathbf{i} F_1(P) + \mathbf{j} F_2(P) + \mathbf{k} F_3(P)) = \int_V (\mathbf{i} f_1 + \mathbf{j} f_2 + \mathbf{k} f_3) \frac{1}{r} \, dv$$

$$\mathbf{F}(P) = \int_V \frac{\mathbf{f}}{r} \, dv.$$

LAPLACE'S EQUATION WITH ITS SOLUTION

If a twice differentiable scalar point function ϕ $\nabla^2 \phi = 0$ is true for every point of the region, the function ϕ is said to be harmonic in the region.

The equation $\nabla^2 \phi = 0 \dots (1)$ is called Laplace's equation. Green's formula is

$$4\pi \phi(P) = \int_S \left[\frac{1}{r} \nabla \phi - \phi \nabla \left(\frac{1}{r} \right) \right] \cdot d\mathbf{S} - \int_V \frac{1}{r} \nabla^2 \phi \, dv$$

$$= \int_S \left[\frac{1}{r} \nabla \phi - \phi \nabla \left(\frac{1}{r} \right) \right] \cdot d\mathbf{S} \text{ by (1)}$$

It follows that the harmonic function ϕ at any point within the region can be expressed in terms of the values of ϕ and $\frac{\partial \phi}{\partial n}$ at any point of the surface enclosing the region.

Problem 152. Verify Green's theorem in the plane for

$$\oint_C (xy + y^2) dx + x^2 dy$$

where C is the closed curve of the region bounded by

$$y = x^2, \text{ and } y = x.$$

The shaded region shown in Fig. 1.58, represents the positive direction traversed by the closed region C made up of a parabola and a straight line.

Given, $\psi = (xy + y^2)$ and $\phi = x^2$.

Evaluating the integral along $z = x^2$, we have,




Fig. 1.58

Evaluating the line integral along $y = x$ we have

$$\int \psi dx + \phi dy = \int [(x^2 + x^2) dx + y^2 dy] = \int_{x=1}^0 2x^2 dx + \int_{y=1}^0 y^2 dy = -1.$$

$\therefore$ the required integral $= \frac{1}{2} - 1 = -\frac{1}{2}$.

Also $\frac{\partial \phi}{\partial x} = 2x$ and $\frac{\partial \psi}{\partial y} = x + 2y$.

Thus, $\iint_R \left(\frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \right) dx dy$

$$\begin{aligned} &= \iint_R [2x - x - 2y] dx dy \\ &= \int_{x=0}^1 \int_{y=x^2}^x (x - 2y) dx dy \\ &= \int_{x=0}^1 [xy - y^2] dx \\ &= \int_{x=0}^1 [x^2 - x^2 - x^3 + x^4] dx \\ &= \int_{x=0}^1 (x^4 - x^3) dx \\ &= -\frac{1}{20}. \end{aligned}$$

$$\begin{aligned} &\int \psi dx + \phi dy \\ &= \int [(x \cdot x^2 + x^4) dx + y dy] \\ &= \int_{x=0}^1 (x^3 + x^4) dx + \int_{y=0}^1 y dy \\ &= \left[\frac{x^4}{4} + \frac{x^5}{5} \right]_0^1 + \left[\frac{y^2}{2} \right]_0^1 \\ &= \frac{1}{4} + \frac{1}{5} + \frac{1}{2} \\ &= \frac{19}{20}. \end{aligned}$$

It is evident that $\int \psi dx + \phi dy = \iint \left(\frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \right) dx dy = -\frac{1}{20}$.

Thus Green's theorem is verified.

Problem 153. Evaluate $\oint_C [(y - \sin x) dx + \cos x dy]$ where C is

a triangle whose vertices are $(0, 0)$; $\left(\frac{\pi}{2}, 0\right)$; $\left(\frac{\pi}{2}, 1\right)$

(a) directly, (b) by using Green's theorem in the plane.

(a) The line integral along OQ on which $y = 0$ and x varies from

$$\int_{x=0}^{\pi/2} \sin x dx = -1.$$

The line integral along QP on which

x and y varies from 0 to 1

$$\int_{y=0}^1 ((y-1) \cdot 0 + 0 dy) = 0.$$

The line integral along PO for which

x and y varies from $\frac{\pi}{2}$ to 0,

$$\begin{aligned} &= \int_{\pi/2}^0 \left\{ \left(\frac{2x}{\pi} - \sin x \right) dx + \frac{2}{\pi} \cos x dx \right\} \\ &= \left[\frac{x^2}{\pi} + \cos x + \frac{2}{\pi} \sin x \right]_{\pi/2}^0 = 1 - \frac{\pi}{4} - \frac{2}{\pi}. \end{aligned}$$

Hence the line integral along $C = -1 + 0 + 1 - \frac{\pi}{4} - \frac{2}{\pi} = -\frac{\pi}{4} - \frac{2}{\pi}$.

(b) In order to use Green's theorem we have

$$\psi = y - \sin x, \phi = \cos x, \frac{\partial \psi}{\partial y} = 1, \frac{\partial \phi}{\partial x} = -\sin x$$

$$\oint_C (\psi dx + \phi dy) = \iint_R \left(\frac{\partial \phi}{\partial x} - \frac{\partial \psi}{\partial y} \right) dx dy = \iint_R (-y \sin x - 1) dx dy$$

$$= \int_{x=0}^{\pi/2} \int_{y=0}^{2x/\pi} [(-\sin x - 1) dy] dx = \int_{x=0}^{\pi/2} \left(-y \sin x - y \right) \Big|_0^{2x/\pi} dx$$

$$= \int_0^{\pi/2} \left(-\frac{2x}{\pi} \sin x - \frac{2x}{\pi} \right) dx$$

$$= \left[-\frac{2}{\pi} (-x \cos x + \sin x) - \frac{x^2}{\pi} \right]_0^{\pi/2}$$




Fig. 1.59

$$= -\frac{2}{\pi} - \frac{\pi}{4}.$$

Problem 154. Compute $\oint \{(xy-x^2) dx + x^2y dy\}$ over the triangle bounded by lines $y=0$, $x=1$, $y=x$ and verify by Green's theorem.

The line integral along OP where $y=0$ and x varies from 0 to 1

$$= \int_{x=0}^1 -x^2 dx = -\frac{1}{3}.$$

The line integral along PQ for which $x=1$ and y varies from 0 to 1

$$= \int_{y=0}^1 \{(y-1) \cdot 0 + y dy\} = \frac{1}{2}.$$

The line integral along QO for which $x=y$ and y varies from 1 to 0

$$= \int_1^0 \{(y^2-y^2) dy + y^3 dy\} = -\int_1^0 y^3 dy = -\frac{1}{4}.$$




Fig. 1.60

$$= \int_{x=0}^1 [x^3 - x^2] dx = \left[\frac{x^4}{4} - \frac{x^3}{3} \right] = -\frac{1}{12},$$

which verifies the Green's theorem.

1.61. STOKES' THEOREM IN SPACE

This states that if, $\mathbf{F}$ is a vector function, which is uniform, finite and continuous along with its derivative in any direction, then the tangential line integral of $\mathbf{F}$ over any closed surface S bounded by a curve C is equal to the normal surface integral of $\text{curl } \mathbf{F}$ over S ; i.e.,

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS = \iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S}$$

where $\mathbf{n}$ is the unit normal vector at any point of S drawn in the direction in which a right handed screw would move when rotated in the sense of description of C .

Consider a surface S such that its projections on the

Thus the total integral along C
 $= -\frac{1}{3} + \frac{1}{2} - \frac{1}{4} = -\frac{1}{12}.$

Now, by Green's theorem, we have

$$\begin{aligned} & \int \{(xy-x^2) dx + x^2y dy\} \\ &= \iint \left[\frac{\partial (x^2y)}{\partial x} - \frac{\partial (xy-x^2)}{\partial y} \right] dx dy \\ &= \int_{x=0}^1 \int_{y=0}^x [2xy - x] dx dy \end{aligned}$$

planes are regions bounded by simple closed curves as shown in Fig. 1.61.

Take the equation of surface as

$$z = f_1(x, y)$$

$$y = f_2(x, z) \text{ or } x = f_3(y, z)$$

$$\mathbf{F} = F_1\mathbf{i} + F_2\mathbf{j} + F_3\mathbf{k},$$

then we have to prove that

$$\begin{aligned} \iint_S \nabla \times (F_1\mathbf{i} + F_2\mathbf{j} + F_3\mathbf{k}) \cdot \mathbf{n} dS \\ = \oint_C \mathbf{F} \cdot d\mathbf{r} \end{aligned}$$

Let us first consider,

$$\iint_S [\nabla \times (F_1\mathbf{i})] \cdot \mathbf{n} dS$$

$$= \left[\left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times F_1\mathbf{i} \right] \cdot \mathbf{n} dS$$

$$= \left[\frac{\partial F_1}{\partial z} \mathbf{j} - \frac{\partial F_1}{\partial y} \mathbf{k} \right] \cdot \mathbf{n} dS$$

$$= \left[\frac{\partial F_1}{\partial z} \mathbf{n} \cdot \mathbf{j} - \frac{\partial F_1}{\partial y} \mathbf{n} \cdot \mathbf{k} \right] dS \quad \dots (1)$$

$$\text{and } \mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k} = x\mathbf{i} + y\mathbf{j} + f_1(x, y)\mathbf{k}$$

$$\text{so that } \frac{\partial \mathbf{r}}{\partial y} = \mathbf{j} + \frac{\partial f_1}{\partial y} \mathbf{k} \text{ since } z = f_1(x, y)$$

then $\frac{\partial \mathbf{r}}{\partial y}$ is perpendicular to $\mathbf{n}$ as $\frac{\partial \mathbf{r}}{\partial y}$ is the tangent to the surface S .

$$\mathbf{n} \cdot \frac{\partial \mathbf{r}}{\partial y} = \mathbf{n} \cdot \mathbf{j} + \frac{\partial f_1}{\partial y} \mathbf{n} \cdot \mathbf{k} = 0.$$

$$\text{Hence } \mathbf{n} \cdot \mathbf{j} = -\frac{\partial f_1}{\partial y} \mathbf{n} \cdot \mathbf{k} = -\frac{\partial z}{\partial y} \mathbf{n} \cdot \mathbf{k}.$$

Thus (1) yields

$$\iint_S [\nabla \times (F_1\mathbf{i})] \cdot \mathbf{n} dS = - \left[\frac{\partial F_1}{\partial z} \cdot \frac{\partial z}{\partial y} + \frac{\partial F_1}{\partial y} \right] \mathbf{n} \cdot \mathbf{k} dS \quad \dots (2)$$

On the surface S , we have

$$F_1(x, y, z) = F_1[x, y, f_1(x, y)] = F(x, y)$$

$$\frac{\partial F_1}{\partial y} + \frac{\partial F_1}{\partial z} \cdot \frac{\partial z}{\partial y} = \frac{\partial F}{\partial y} \quad \dots (3)$$

On relation (2) with the help of (3) gives

$$\iint_S [\nabla \times (F_1\mathbf{i})] \cdot \mathbf{n} dS = - \frac{\partial F}{\partial y} \mathbf{n} \cdot \mathbf{k} dS = - \frac{\partial F}{\partial y} dx dy$$

$$\iint_S [\nabla \times (F_1\mathbf{i})] \cdot \mathbf{n} dS = \iint_R - \frac{\partial F}{\partial y} dx dy \quad \dots (4)$$




Fig. 1.61

where R is the projection of S on xy plane.

Green's theorem in plane gives

$$\oint_{C_1} F dx = - \iint_R \frac{\partial F}{\partial y} dx dy \text{ where } C_1 \text{ is the boundary of } R.$$

Now at each point (x, y) of the curve C_1 the value of F being the same as that of F_1 at each point (x, y, z) of C and dx being the same for both the curves C and C_1 , we conclude that

$$\oint_{C_1} F dx = \oint_C F_1 dx$$

$$\text{i.e.} \quad \oint_C F_1 dx = - \iint_R \frac{\partial F}{\partial y} dx dy$$

The equations (4) and (5), give

$$\iint_S [\nabla \times (F_1 \mathbf{i})] \cdot \mathbf{n} dS = \oint_C F_1 dx$$

Similar contributions are made by the projections on the other two planes, therefore

$$\iint_S [\nabla \times (F_2 \mathbf{j})] \cdot \mathbf{n} dS = \oint_C F_2 dy$$

$$\iint_S [\nabla \times (F_3 \mathbf{k})] \cdot \mathbf{n} dS = \oint_C F_3 dz$$

Adding (6), (7) and (8), we find

$$\iint_S [\nabla \times (F_1 \mathbf{i} + F_2 \mathbf{j} + F_3 \mathbf{k})] \cdot \mathbf{n} dS = \oint_C (F_1 dx + F_2 dy + F_3 dz)$$

$$\iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS = \oint_C \mathbf{F} \cdot d\mathbf{r}.$$

Problem 155. If ϕ is continuously differentiable scalar point function then show that

$$\oint_C \phi d\mathbf{r} = \iint_S \mathbf{n} \times \nabla \phi dS.$$

Put $\mathbf{F} = \mathbf{a} \phi$, where $\mathbf{a}$ is a constant vector, in the Stokes's theorem

$$\oint_C (\mathbf{a} \phi) \cdot d\mathbf{r} = \iint_S [\nabla \times (\mathbf{a} \phi)] \cdot \mathbf{n} dS$$

But we know that

$$\nabla \times (\mathbf{a} \phi) = \text{grad } \phi \times \mathbf{a} + \phi \text{ curl } \mathbf{a} = \nabla \phi \times \mathbf{a}$$

$$\begin{aligned} \therefore \mathbf{a} \cdot \oint_C \phi d\mathbf{r} &= \iint_S (\nabla \phi \times \mathbf{a}) \cdot \mathbf{n} dS \\ &= - \iint_S \mathbf{a} \times \nabla \phi \cdot \mathbf{n} dS \\ &= - \iint_S \mathbf{a} \cdot \nabla \phi \times \mathbf{n} dS = \mathbf{a} \cdot \iint_S \nabla \phi \times \mathbf{n} dS \end{aligned}$$

Since $\mathbf{a}$ is a constant vector

$$\oint_C \phi d\mathbf{r} = - \iint_S \nabla \phi \times \mathbf{n} dS = \iint_S \mathbf{n} \times \nabla \phi dS.$$

Problem 156. Show that

$$\oint_C d\mathbf{r} \times \mathbf{F} = \iint_S (\mathbf{n} \times \nabla) \times \mathbf{F} dS.$$

Putting

$$\mathbf{f} = \mathbf{a} \times \mathbf{F}, \quad \mathbf{a} \text{ being a constant vector;}$$

Stokes' theorem gives

$$\oint_C (\mathbf{a} \times \mathbf{F}) \cdot d\mathbf{r} = \iint_S [\nabla \times (\mathbf{a} \times \mathbf{F})] \cdot \mathbf{n} dS$$

But we know that

$$\begin{aligned} \nabla \times (\mathbf{a} \times \mathbf{F}) &= \mathbf{a} \nabla \cdot (\mathbf{F}) - \mathbf{F} \nabla \cdot (\mathbf{a}) + (\mathbf{F} \cdot \nabla) \mathbf{a} - (\mathbf{a} \cdot \nabla) \mathbf{F} \\ &= \mathbf{a} \nabla \cdot \mathbf{F} - (\mathbf{a} \cdot \nabla) \mathbf{F} \end{aligned}$$

$$[(\mathbf{a} \cdot \nabla) \mathbf{F}] \cdot \mathbf{n} = (\mathbf{a} \cdot \nabla) (\mathbf{F} \cdot \mathbf{n}).$$

$$\oint_C (\mathbf{a} \times \mathbf{F}) \cdot d\mathbf{r} = \iint_S [\mathbf{a} \cdot (\nabla \cdot \mathbf{F}) \mathbf{n} - \mathbf{a} \cdot \nabla (\mathbf{F} \cdot \mathbf{n})] dS$$

$$\mathbf{a} \cdot \oint_C \mathbf{F} \times d\mathbf{r} = \mathbf{a} \cdot \iint_S [(\nabla \cdot \mathbf{F}) \mathbf{n} - \nabla (\mathbf{F} \cdot \mathbf{n})] dS$$

Since $\mathbf{a}$ is a constant vector

$$\oint_C \mathbf{F} \times d\mathbf{r} = \iint_S [(\nabla \cdot \mathbf{F}) \mathbf{n} - \nabla (\mathbf{F} \cdot \mathbf{n})] dS$$

$$= \iint_S -[(\mathbf{n} \times \nabla) \times \mathbf{F}] dS$$

$$\oint_C d\mathbf{r} \times \mathbf{F} = \iint_S [(\mathbf{n} \times \nabla) \times \mathbf{F}] dS.$$

Stokes' theorem in the plane is sometimes known as Green's theorem in the plane.

Problem 157. Verify Stokes' theorem for $\mathbf{F} = (2x - y) \mathbf{i} - yz^2 \mathbf{j} - y^2 z \mathbf{k}$, where S is the upper half surface of the sphere $x^2 + y^2 + z^2 = 1$ and C is its boundary.

Stokes' theorem is

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS.$$

In the xy -plane the boundary C of the surface S is a circle $x^2 + y^2 = 1$.

Let $x = \cos t$, $y = \sin t$ and $z = 0$ for which $0 \leq t \leq 2\pi$ so that these are the parametric equations of C .

$$\begin{aligned} \oint_C \mathbf{F} \cdot d\mathbf{r} &= \int_C [(2x - y) \mathbf{i}] \cdot [dx \mathbf{i} + dy \mathbf{j} + dz \mathbf{k}] = \int_C (2x - y) dx \\ &= \int_{t=0}^{2\pi} [2 \cos t - \sin t] (-\sin t dt) \end{aligned}$$

$$= \int_{t=0}^{2\pi} (2 \sin t \cos t - \sin^2 t) dt$$

$$= \pi.$$

and

$$\nabla \times \mathbf{F} = \mathbf{i} \left[\frac{\partial}{\partial y} (-y^2 z) - \frac{\partial}{\partial z} (-y z^2) \right]$$

$$+ \mathbf{j} \left[\frac{\partial}{\partial z} (2x - y) - \frac{\partial}{\partial x} (-y^2 z) \right] + \mathbf{k} \left[\frac{\partial}{\partial x} (-y^2 z) - \frac{\partial}{\partial y} (2x - y) \right]$$

$$= 0\mathbf{i} + 0\mathbf{j} + \mathbf{k} = \mathbf{k}.$$

$$\therefore \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS = \iint_S \mathbf{k} \cdot \mathbf{n} dS$$

$$= \iint_R dx dy \text{ since } \mathbf{k} \cdot \mathbf{n} dS = dx dy$$

R being the projection of S on xy plane.

$$\text{Thus, } \iint_R dx dy = \int_{x=-1}^{x=+1} \int_{y=-\sqrt{1-x^2}}^{y=+\sqrt{1-x^2}} dx dy$$

$$= \int_0^1 \sqrt{1-x^2} \cdot 4 \frac{\pi}{4} = \pi$$

which verifies Stokes' theorem.

Problem 158. Evaluate $\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS$ for

$$\mathbf{F} = (y-z+2)\mathbf{i} + (yz+4)\mathbf{j} - xz\mathbf{k},$$

where S is the surface of the cube $x=y=z=0$; $x=y=z=2$ above xy plane.

Stokes' theorem is

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS$$

Here the boundary C of the surface S is a square bounded by the lines $x=0$, $x=2$, $y=0$, $y=2$ in the xy plane. So

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_{OP} \mathbf{F} \cdot d\mathbf{r} + \int_{PQ} \mathbf{F} \cdot d\mathbf{r} + \int_{QR} \mathbf{F} \cdot d\mathbf{r} + \int_{RO} \mathbf{F} \cdot d\mathbf{r}$$

Along OP , $y=0$ and x varies from 0 to 2.

$$\therefore \int_{OP} \mathbf{F} \cdot d\mathbf{r} = \int_0^2 [2\mathbf{i} + 4\mathbf{j}] \cdot d\mathbf{r}$$

$$dy=0 \text{ and } dz=0$$

$$= 2 \int_0^2 dx = 4.$$

Along PQ , $x=2$ and y varies from 0 to 2, so that $dx=0$, $dz=0$.




Fig. 1.62

$$\int_{RO} \mathbf{F} \cdot d\mathbf{r} = \int [(y+z)\mathbf{i} + 4\mathbf{j}] \cdot d\mathbf{r}$$

$$= 4 \int_0^2 dy = 8.$$

Along QR , $y=2$ and x varies from 2 to 0; so that $dy=0$, $dz=0$.

$$\int_{QR} \mathbf{F} \cdot d\mathbf{r} = \int [4\mathbf{j}] \cdot [dx\mathbf{i}]$$

$$= 4 \int_2^0 dx = -8.$$

Along RO , $x=0$ and y varies from 2 to 0 so that $dx=0$, $dz=0$

$$\int_{RO} \mathbf{F} \cdot d\mathbf{r} = \int [4\mathbf{j}] \cdot [dy\mathbf{j}] = 4 \int_2^0 dy = -8.$$

$$\text{Hence } \int_C \mathbf{F} \cdot d\mathbf{r} = 4 + 8 - 8 - 8 = -4.$$

Problem 159. Verify Stokes' theorem for the vector $\mathbf{F} = (z, x, y)$ over the half of the sphere $x^2 + y^2 + z^2 = a^2$ lying above xy plane.

The projection of the surface on $z=0$ plane is a circle $x^2 + y^2 = a^2$, boundary C (say),

$$\mathbf{F} = z\mathbf{i} + x\mathbf{j} + y\mathbf{k}$$

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C (0\mathbf{i} + x\mathbf{j} + y\mathbf{k}) \cdot (dx\mathbf{i} + dy\mathbf{j})$$

$$= \int_C x dy$$

$$= \int_{y=-a}^{y=a} \pm \sqrt{a^2 - y^2} dy$$

$$= 2 \int_0^a \sqrt{a^2 - y^2} dy,$$

$y = a \sin \theta$, so that $dy = a \cos \theta d\theta$

$$= 2 \int_0^{\pi/2} a^2 \cos^2 \theta d\theta = \pi a^2.$$




Fig. 1.63

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ z & x & y \end{vmatrix} = (\mathbf{i} - \mathbf{j} + \mathbf{k})$$

$$\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS = \iint_S (\mathbf{i} - \mathbf{j} + \mathbf{k}) \cdot \mathbf{n} dS$$

$$\begin{aligned}
&= \iint_S (\mathbf{i} \cdot \mathbf{n} - \mathbf{j} \cdot \mathbf{n} + \mathbf{k} \cdot \mathbf{n}) dS \\
&= \iint_S \mathbf{i} \cdot \mathbf{n} dS - \iint_S \mathbf{j} \cdot \mathbf{n} dS + \iint_S \mathbf{k} \cdot \mathbf{n} dS \\
&= \iint_{R_1} dy dz - \iint_{R_2} dx dz + \iint_{R_3} dx dy \\
&= \iint_{R_3} dx dy,
\end{aligned}$$

R_1, R_2, R_3 being projections on zy, yz, xy planes, respectively, projection on zy plane being the same as that on xz plane, the two integrals cancel out.

$$\begin{aligned}
\text{Thus } \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS &= \int_{x=-a}^a \int_{y=-\sqrt{a^2-x^2}}^{\sqrt{a^2-x^2}} dx dy \\
&= 2 \int_{-a}^a \sqrt{a^2-x^2} dx \\
&= 4 \int_0^a \sqrt{a^2-x^2} dx \\
&= 4\pi \frac{a^2}{4} = \pi a^2.
\end{aligned}$$

$$\text{Hence, } \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS = \int_C \mathbf{F} \cdot d\mathbf{r}$$

which verifies Stokes' theorem.

Problem 160. Prove that

$$\int_C \mathbf{r} \times d\mathbf{r} = 2 \iint_S d\mathbf{S}$$

where S is a diaphragm enclosing a circuit C .

Putting $\mathbf{F} = \mathbf{a} \times \mathbf{r}$, where $\mathbf{a}$ is a constant vector, in Stokes' theorem, i.e.

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS,$$

We get

$$\int_C (\mathbf{a} \times \mathbf{r}) \cdot d\mathbf{r} = \iint_S \nabla \times (\mathbf{a} \times \mathbf{r}) \cdot \mathbf{n} dS$$

$$\text{But, } \nabla \times (\mathbf{a} \times \mathbf{r}) = \mathbf{a} \nabla \cdot \mathbf{r} - (\mathbf{a} \cdot \nabla) \mathbf{r} = 3\mathbf{a} - 2\mathbf{a} = \mathbf{a}.$$

$$\therefore \int_C (\mathbf{a} \cdot \mathbf{r}) \times d\mathbf{r} = \iint_S \mathbf{a} \cdot \mathbf{n} dS$$

$$\mathbf{a} \cdot \int_C \mathbf{r} \times d\mathbf{r} = 2\mathbf{a} \cdot \iint_S d\mathbf{S}$$

Since $\mathbf{a}$ is an arbitrary constant vector

$$\int_C \mathbf{r} \times d\mathbf{r} = 2 \iint_S d\mathbf{S}.$$

Problem 161. Prove that a necessary and sufficient condition that

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$$

every closed curve C is that $\nabla \times \mathbf{F} = 0$ identically.

The condition is sufficient :

Since if $\nabla \times \mathbf{F} = 0$

then Stokes' theorem gives at once,

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS = 0.$$

the condition is necessary: Since if $\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$ round any

curve C , then taking $\nabla \times \mathbf{F} \neq 0$ at some point P , there will be a region with P as an interior point where $\nabla \times \mathbf{F} \neq 0$ provided $\mathbf{F}$ is continuous. Assuming S to be the surface contained in the region whose unit normal $\mathbf{n}$ at each point has the same direction as that of $\nabla \times \mathbf{F}$, we may express $\nabla \times \mathbf{F} = \lambda \mathbf{n}$ where λ is a positive scalar. Thus Stokes' theorem gives

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS = \lambda \iint_S \mathbf{n} \cdot \mathbf{n} dS = \lambda \iint_S dS \neq 0$$

which yields positive contribution.

This is contrary to our hypothesis and hence $\nabla \times \mathbf{F} = 0$.

The necessary and sufficient condition for $\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$ is that

$$\nabla \times \mathbf{F} = 0.$$

Problem 162. Prove the following :

$$\int_C \mathbf{r} \cdot d\mathbf{r} = 0.$$

$$\int_C [\phi \nabla \phi] \cdot d\mathbf{r} = 0.$$

$$\int_C [\phi \nabla \psi] \cdot d\mathbf{r} = - \int_C \psi \nabla \phi \cdot d\mathbf{r}.$$

By Stokes' theorem

$$\int_C \mathbf{r} \cdot d\mathbf{r} = \iint_S [\nabla \times \mathbf{r}] \cdot \mathbf{n} dS = 0 \quad \text{as } \nabla \times \mathbf{r} = 0.$$

By Stokes' theorem

$$[\phi \nabla \phi] \cdot d\mathbf{r} = \iint_S [\nabla \times (\phi \nabla \phi)] \cdot \mathbf{n} dS.$$

$$\nabla \times (\phi \nabla \phi) = \phi \nabla \times (\nabla \phi) + (\nabla \phi) \times (\nabla \phi) = 0 + 0 = 0$$

$$\therefore \int_C (\phi \nabla \phi) \cdot d\mathbf{r} = 0.$$

(c) By Stokes' theorem

$$\int_C [\phi \nabla \psi] \cdot d\mathbf{r} = \iint_S [\nabla \times (\phi \nabla \psi)] \cdot \mathbf{n} \, dS.$$

$$\text{But } \nabla \times (\phi \nabla \psi) = \phi \nabla \times (\nabla \psi) + (\nabla \phi) \times (\nabla \psi) \\ = \nabla \phi \times \nabla \psi.$$

$$\therefore \int_C [\phi \nabla \psi] \cdot d\mathbf{r} = \iint_S (\nabla \phi \times \nabla \psi) \cdot \mathbf{n} \, dS \\ = - \iint_S (\nabla \psi \times \nabla \phi) \cdot \mathbf{n} \, dS$$

$$\text{and } - \int_C \psi \nabla \phi \cdot d\mathbf{r} = - \iint_S \nabla \times (\psi \nabla \phi) \cdot \mathbf{n} \, dS$$

(by Stokes' theorem)

$$= - \iint_S \{ \psi \nabla \times (\nabla \phi) + \nabla \psi \times \nabla \phi \} \cdot \mathbf{n} \, dS$$

$$= - \iint_S (\nabla \psi \times \nabla \phi) \cdot \mathbf{n} \, dS \\ \text{as } \psi \nabla \times (\nabla \phi) = 0$$

It is evident from (1) and (2),

$$\int_C \phi \nabla \psi \cdot d\mathbf{r} = - \int_C \psi \nabla \phi \cdot d\mathbf{r}.$$

$$\text{Problem 163. If } \oint_C \mathbf{E} \cdot d\mathbf{r} = -\frac{1}{C} \frac{\partial}{\partial t} \iint_S \mathbf{H} \cdot d\mathbf{S},$$

where S is any surface bounded by curve C , show that

$$\nabla \times \mathbf{E} = -\frac{1}{C} \frac{\partial \mathbf{H}}{\partial t}.$$

$$\text{Let } \nabla \times \mathbf{E} = -\frac{1}{C} \frac{\partial \mathbf{H}}{\partial t}$$

Then Stokes' theorem yields

$$\oint_C \mathbf{E} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{E}) \cdot \mathbf{n} \, dS = \iint_S (\nabla \times \mathbf{E}) \cdot \mathbf{n} \, dS$$

$$= \iint_S \left(-\frac{1}{C} \frac{\partial \mathbf{H}}{\partial t} \right) \cdot \mathbf{n} \, dS \quad \text{by (1)}$$

$$= -\frac{1}{C} \iint_S \frac{\partial}{\partial t} (\mathbf{H}) \cdot \mathbf{n} \, dS$$

$$= -\frac{1}{C} \frac{\partial}{\partial t} \iint_S \mathbf{H} \cdot d\mathbf{S}$$

the integral being independent of t .

Problem 164. If the normal surface integral of a vector point function over every open surface is equal to the tangential line integral of the vector function $\mathbf{F}$ round its boundary, prove that

$$\mathbf{G} = \text{curl } \mathbf{F}.$$

We know that the normal surface integral of a vector point function is given by $\iint_S \mathbf{G} \cdot d\mathbf{S}$

if S is a surface.

And the tangential line integral of vector point function $\mathbf{F}$ is given by $\oint_C \mathbf{F} \cdot d\mathbf{r}$.

We are given that

$$\iint_S \mathbf{G} \cdot d\mathbf{S} = \oint_C \mathbf{F} \cdot d\mathbf{r} \quad \dots (1)$$

Stokes' theorem, yields

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, dS = \iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S}$$

$$\iint_S \mathbf{G} \cdot d\mathbf{S} = \iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S} \quad \text{by (1)}$$

It follows that

$$\mathbf{G} = \nabla \times \mathbf{F} \\ \mathbf{G} = \text{curl } \mathbf{F}.$$

SOME THEOREMS

THEOREM 1. The necessary and sufficient condition that a vector function $\mathbf{F}$ be an **Irrotational** vector function in a simply connected region is that $\text{curl } \mathbf{F} = 0$ at every point of the region.

This condition is necessary, since if $\mathbf{F}$ be irrotational then there is a scalar ϕ such that $\mathbf{F} = \nabla \phi$

$$\text{curl } \mathbf{F} = \text{curl } (\nabla \phi) = \nabla \times \nabla \phi = 0$$

This condition is also sufficient, since if $\text{curl } \mathbf{F} = 0$ then it follows from Stokes' theorem that

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S \text{curl } \mathbf{F} \cdot d\mathbf{S} = 0$$

that $\mathbf{F}$ is irrotational.

THEOREM 2. The necessary and sufficient condition that a vector function $\mathbf{F}$ be a **Solenoidal** vector function in a simply connected region is that $\text{div } \mathbf{F} = 0$ at every point of the region.

This condition is necessary, since if $\mathbf{F}$ be solenoidal then at any point we have by §1.52 (note)

$$\text{div } \mathbf{F} = \lim_{V \rightarrow 0} \iint_S \frac{\mathbf{F} \cdot d\mathbf{S}}{V} = 0$$

that $\mathbf{F}$ is solenoidal.

The condition is sufficient, since if $\text{div } \mathbf{F}=0$, then Gauss' divergence theorem yields,

$$\int_S \mathbf{F} \cdot d\mathbf{S} = \int_V \text{div } \mathbf{F} dV = 0$$

showing that the Flux $\int_S \mathbf{F} \cdot d\mathbf{S}$ across every closed surface is zero.

Note. $\text{div } \text{curl } \mathbf{F}=0 \Rightarrow$ divergence of every curl is zero i.e., curl of every function is solenoidal.

THEOREM 3. If $\mathbf{F}$ is a continuously differentiable vector point function such that $\text{div } \mathbf{F}=0$, then there exists another vector point function $\mathbf{f}$ such that $\mathbf{F}=\text{curl } \mathbf{f}$.

Firstly to show that $\mathbf{f}$ is any function whose curl is $\mathbf{F}$, take a given function $\mathbf{f}+\nabla\phi$ whose curl is $\mathbf{F}$, ϕ being continuously differentiable scalar point function.

Assuming that $\text{curl } \mathbf{f}=\mathbf{F}=\text{curl } \mathbf{g}$, we have $\text{curl } (\mathbf{g}-\mathbf{f})=0$ which follows that $\mathbf{g}-\mathbf{f}$ is the gradient of some scalar ϕ i.e.,

$$\mathbf{g}-\mathbf{f}=\nabla\phi \text{ giving } \mathbf{g}=\mathbf{f}+\nabla\phi$$

But if ϕ be any scalar point function, then

$$\begin{aligned} \text{curl } (\mathbf{f}+\nabla\phi) &= \text{curl } \mathbf{f} + \text{curl } \nabla\phi \\ &= \mathbf{F} + \nabla \times \nabla\phi = \mathbf{F} \end{aligned}$$

which proves the proposition.

Now to prove the main theorem, let us suppose that

$$\mathbf{F}=F_1\mathbf{i}+F_2\mathbf{j}+F_3\mathbf{k}$$

$$\mathbf{f}=f_1\mathbf{i}+f_2\mathbf{j}+f_3\mathbf{k}$$

$$\text{Then, } \text{curl } \mathbf{f} = \nabla \times \mathbf{f} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ f_1 & f_2 & f_3 \end{vmatrix}$$

$$= \left(\frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} \right) \mathbf{i} + \left(\frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x} \right) \mathbf{j} + \left(\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) \mathbf{k}$$

So that $\mathbf{F}=\text{curl } \mathbf{f}$ gives on comparison of the coefficients of $\mathbf{i}, \mathbf{j}, \mathbf{k}$,

$$F_1 = \frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z}, F_2 = \frac{\partial f_1}{\partial z} - \frac{\partial f_3}{\partial x}, F_3 = \frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y}$$

$$\text{Also, } \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z} = 0$$

If we suppose that $f_1=0$, then (1) gives

$$F_2 = -\frac{\partial f_3}{\partial x} \text{ giving } f_3 = -\int_{x_0}^x F_2 dx + \psi(y, z)$$

$$\text{and } F_3 = \frac{\partial f_2}{\partial x} \text{ giving } f_2 = \int_{x_0}^x F_3 dx + \psi(y, z),$$

y, z being parameters

$$\text{We get from (3) and (4), } \frac{\partial f_3}{\partial y} = -\int_{x_0}^x \frac{\partial F_2}{\partial y} dx + \frac{\partial \psi}{\partial y} \quad \dots (5)$$

$$\frac{\partial f_3}{\partial z} = \int_{x_0}^x \frac{\partial F_3}{\partial z} dx + \frac{\partial \psi}{\partial z} \quad \dots (6)$$

$$\begin{aligned} \text{So that } F_1 &= \frac{\partial f_3}{\partial y} - \frac{\partial f_2}{\partial z} = -\int_{x_0}^x \left(\frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z} \right) dx + \frac{\partial \psi}{\partial y} - \frac{\partial \psi}{\partial z} \\ &= \int_{x_0}^x \frac{\partial F_1}{\partial x} dx + \frac{\partial \psi}{\partial y} - \frac{\partial \psi}{\partial z} \text{ with the help of (2)} \\ &= F_1(x, y, z) - F_1(x_0, y, z) + \frac{\partial \psi}{\partial y} - \frac{\partial \psi}{\partial z} \end{aligned}$$

$$F_1(x_0, y, z) = \frac{\partial \psi}{\partial y} - \frac{\partial \psi}{\partial z}$$

If we now suppose that $\psi=0$, then

$$\frac{\partial \phi}{\partial y} = F_1(x_0, y, z) \text{ gives } \phi = \int_{y_0}^y F_1(x_0, y, z) dy$$

As such we find from (3) and (4) etc.

$$f_1=0,$$

$$f_2 = \int_{x_0}^x F_3 dx$$

$$f_3 = -\int_{x_0}^x F_2 dx + \phi(y, z),$$

$$\phi(y, z) = \int_{y_0}^y F_1(x_0, y, z) dy$$

It is evident that f_1, f_2, f_3 as determined here are the components of a vector $\mathbf{f}$ whose curl is $\mathbf{F}$.

$$\text{Hence } \text{div } \mathbf{F}=0 \Rightarrow \mathbf{F}=\text{curl } \mathbf{f}.$$

THE CLASSIFICATION OF VECTOR FIELDS

(Kanpur 1968, Agra 1954, 63, 65)

If $\nabla \times \mathbf{F}=0$ i.e. $\nabla \times \mathbf{F}=0$ then $\mathbf{F}=\text{grad } \phi$ or $\mathbf{F}$ is called as a

Gradient Field or a Zero Curl Field. Also if $\text{div } \mathbf{F}=0$ i.e. $\nabla \cdot \mathbf{F}=0$

then $\mathbf{F}=\text{curl } \mathbf{f}$ or $\mathbf{F}$ is called as a Solenoidal field. It is conventional

to classify the vector fields into four:

(i) When $\text{curl } \mathbf{F}=0$ and $\text{div } \mathbf{F}=0$, then the first condition shows

that the field is lamellar or irrotational

implies $\mathbf{F}=\text{grad } \phi$ and in

the second condition it gives $\text{div } \mathbf{F}=0$ i.e. $\nabla^2 \phi=0$ i.e. Laplace's

equation showing that the field is solenoidal or incompressible. On the whole

the field is termed as a type of field,

which is irrotational motion of incom-

pressible fluid as shown in Fig. 1.63.




Fig. 1.63




Fig. 1.64

$\mathbf{f} \neq 0$ or $\nabla \times (\nabla \times \mathbf{f}) \neq 0$
i.e., $\text{grad div } \mathbf{f} - \nabla^2 \mathbf{f} \neq 0$

This shows that if $\mathbf{f}$ is solenoidal then we must have $\text{div } \mathbf{f} = 0$, so that $\text{grad div } \mathbf{f} = 0$ and as such $\nabla^2 \mathbf{f} \neq 0$. Hence such a field is termed as the type of field which is *rotational motion of incompressible fluid* as shown in Fig. 1.65.

(4) When $\text{curl } \mathbf{F} \neq 0$, also $\text{div } \mathbf{F} \neq 0$. This type of vector field is most general and it is termed as the type of field which is *rotational motion of compressible fluid* as shown in Fig. 1.66.




Fig. 1.66

In fact this field is made up of two fields namely (i) Lamellar vector field (i.e. having no curl but may have div only), (ii) Solenoidal vector field (i.e. having no div but may have curl only). Mathematically,

$$\mathbf{F} = \text{grad } \phi + \text{curl } \mathbf{f}$$

$$\begin{aligned} \text{So that } \text{div } \mathbf{F} &= \text{div} (\text{grad } \phi + \text{curl } \mathbf{f}) \\ &= \text{div grad } \phi \quad \because \text{div curl } \mathbf{f} = 0 \\ &= \nabla^2 \phi \end{aligned}$$

But $\text{div } \mathbf{F} \neq 0$, therefore $\nabla^2 \phi \neq 0$ which determines ϕ .

$$\begin{aligned} \text{Again } \text{curl } \mathbf{F} &= \text{curl} (\text{grad } \phi + \text{curl } \mathbf{f}) = \text{curl curl } \mathbf{f} \\ &\quad \because \text{curl grad } \phi = 0 \\ &= -\nabla^2 \mathbf{f} \end{aligned}$$

But $\text{curl } \mathbf{F} \neq 0$, $\therefore \nabla^2 \mathbf{f} \neq 0$, where $\mathbf{f}$ is solenoidal vector field and this determines $\mathbf{f}$.

Such a decomposition of vector field comprising Lamellar and solenoidal field is known as *Helmholtz's theorem*.

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 165. Prove that if $\mathbf{A}$, $\mathbf{B}$ and $\mathbf{C}$ are three non-coplanar vectors, then vector $\mathbf{F}$ can be put in the form $\mathbf{F} = \alpha \mathbf{B} \times \mathbf{C} + \beta \mathbf{C} \times \mathbf{A} + \gamma \mathbf{A} \times \mathbf{B}$. Determine α , β and γ .

(Agra, 1968)

Given that $\mathbf{A}$, $\mathbf{B}$, $\mathbf{C}$ are non-coplanar vectors and we have to show that $\mathbf{B} \times \mathbf{C}$, $\mathbf{C} \times \mathbf{A}$ and $\mathbf{A} \times \mathbf{B}$ are also non-coplanar. They will be so if their scalar triple product is not zero i.e. if $[\mathbf{B} \times \mathbf{C}, \mathbf{C} \times \mathbf{A}, \mathbf{A} \times \mathbf{B}] \neq 0$

$$\begin{aligned} \text{Now } [\mathbf{B} \times \mathbf{C}, \mathbf{C} \times \mathbf{A}, \mathbf{A} \times \mathbf{B}] &= (\mathbf{B} \times \mathbf{C}) \cdot (\mathbf{C} \times \mathbf{A}) \times (\mathbf{A} \times \mathbf{B}) \\ &= (\mathbf{B} \times \mathbf{C}) \cdot \{[\mathbf{CAB}] \mathbf{A} - [\mathbf{CAA}] \mathbf{B}\} \\ &= [\mathbf{BCA}] [\mathbf{CAB}] \quad \because [\mathbf{CAA}] = 0 \\ &= [\mathbf{ABC}] [\mathbf{ABC}] \\ &= [\mathbf{ABC}]^2 \end{aligned} \quad \dots (1)$$

But $[\mathbf{A}, \mathbf{B}, \mathbf{C}] \neq 0$ since $\mathbf{A}$, $\mathbf{B}$, $\mathbf{C}$ are non-coplanar.

It therefore follows from (1) that $[\mathbf{B} \times \mathbf{C}, \mathbf{C} \times \mathbf{A}, \mathbf{A} \times \mathbf{B}] \neq 0$

i.e., $\mathbf{B} \times \mathbf{C}$, $\mathbf{C} \times \mathbf{A}$, $\mathbf{A} \times \mathbf{B}$ are three non-coplanar vectors and as such any vector $\mathbf{F}$ can be expressed in the form

$$\mathbf{F} = \alpha \mathbf{B} \times \mathbf{C} + \beta \mathbf{C} \times \mathbf{A} + \gamma \mathbf{A} \times \mathbf{B} \quad \dots (2)$$

Now to determine α , β , γ , multiply (2) scalarly by $\mathbf{A}$

$$\mathbf{A} \cdot \mathbf{F} = \alpha \mathbf{A} \cdot \mathbf{B} \times \mathbf{C}, \text{ other two scalar triple products vanish}$$

giving

$$\alpha = \frac{\mathbf{A} \cdot \mathbf{F}}{[\mathbf{ABC}]}$$

Similarly multiplying (2) scalarly by $\mathbf{B}$ and $\mathbf{C}$ successively we find

$$\beta = \frac{\mathbf{B} \cdot \mathbf{F}}{[\mathbf{ABC}]} \quad \text{and} \quad \gamma = \frac{\mathbf{C} \cdot \mathbf{F}}{[\mathbf{ABC}]}$$

Problem 166. If $\mathbf{A}(t)$ be a vector function of the scalar variable t and be of constant length, then show that $\frac{d}{dt} \mathbf{A}(t)$ is a vector perpendicular to $\mathbf{A}(t)$. (Agra, 1968)

Given vector $\mathbf{A}$ is of constant length i.e., $|\mathbf{A}| = \text{constant}$ and is a function of t . We know that $\mathbf{A} \cdot \mathbf{A} = A^2 = |\mathbf{A}|^2 = A^2$, A being module of $\mathbf{A}$.

$$\begin{aligned} \text{Differentiation gives, } 2\mathbf{A} \cdot \frac{d\mathbf{A}}{dt} &= 2A \frac{dA}{dt} \\ &= 0 \end{aligned}$$

$$\therefore \frac{dA}{dt} = 0 \text{ when } A = \text{constant.}$$

$$\text{Or } \mathbf{A}(t) \cdot \frac{d}{dt} \mathbf{A}(t) = 0$$

which follows that $\mathbf{A}(t)$ is a vector perpendicular to $\frac{d}{dt} \mathbf{A}(t)$.

Problem 167. Show that $\text{div} (\nabla u \times \nabla v) = 0$ (Agra, 1960)

$$\begin{aligned} \text{By 1.35, } \text{div} (\nabla u \times \nabla v) &= \nabla v \cdot \text{curl } \nabla u - \nabla u \cdot \text{curl } \nabla v \\ &= \nabla v \cdot \text{curl grad } u - \nabla u \cdot \text{curl grad } v \\ &= 0 \text{ as curl grad } u = \text{curl grad } v = 0 \text{ by Problem 91(ii)} \end{aligned}$$

Problem 168. Show that $\nabla \cdot (\mathbf{a} \times \mathbf{r}) = 0$, $\mathbf{a}$ being a constant vector. (Agra, 1958)

Take $\mathbf{a} = a_1 \mathbf{i} + a_2 \mathbf{j} + a_3 \mathbf{k}$, $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ and verify it.

Problem 169. In the gravitational field of a mass m , the potential is given by $\frac{m}{r}$, where r is the distance from the mass, given by $r^2 = x^2 + y^2 + z^2$. Obtain the components of force vector by differentiation. Find the curl of the force and show that it is zero. (Rohilkhand, 1977; Agra, 1955)

$$\text{Here } \text{if } V = -\frac{m}{r} \text{ then } F_x = -\frac{\partial V}{\partial x} \text{ etc and } \mathbf{F} = F_x \mathbf{i} + F_y \mathbf{j} + F_z \mathbf{k}.$$

It is easy to verify $\nabla \times \mathbf{F} = 0$.

Problem 170. Find the Cartesian components of vector C_i which is perpendicular to the vectors $A (2i-j-4k)$ and $B (3i-j-k)$. (Agra, 1933)

It is easy to find the components of $A \times B$ which is perpendicular to A and B both. Ans. $-3, -10, 1$

Problem 171. Prove that $(A \times B) \cdot (C \times A) = (A \cdot C)(B \cdot A) - (B \cdot C)(A \cdot A)$ (Agra, 1961)

Problem 172. The rectangular components of a vector A are

$$A_x = y \frac{\partial f}{\partial z} - z \frac{\partial f}{\partial y}, \quad A_y = z \frac{\partial f}{\partial x} - x \frac{\partial f}{\partial z}, \quad A_z = x \frac{\partial f}{\partial y} - y \frac{\partial f}{\partial x}$$

Where f is given function of (x, y, z) .

Express A as vector product of two vectors and evaluate $A \cdot r$ and $A \cdot \text{grad } f$. (Agra, 1961)

Here $A = A_x i + A_y j + A_z k = (xi + yj + zk) \times \nabla f = r \times \nabla f$ etc. and it is easy to show that $A \cdot r = 0 = A \cdot \text{grad } f$.

Problem 173. Establish Poisson's and Laplace's Equations.

Gauss's theorem, for a volume distribution of density ρ , gives

$$N = \int_S A \cdot n \, dS = 4\pi \int_V \rho \, dv,$$

where N is the flux of the electrostatic intensity A at a point of closed surface S on which the positive unit normal is n .

With the help of divergence theorem, we find

$$\int_V (\text{div } A) \, dv = 4\pi \int_V \rho \, dv$$

or

$$\int_V (\text{div } A - 4\pi\rho) \, dv = 0,$$

which is true for all volumes however small.

$\therefore \text{div } A - 4\pi\rho = 0$, i.e., $\text{div } A = 4\pi\rho$

or

$$\nabla \cdot A = 4\pi\rho.$$

$$\therefore \nabla \cdot (\nabla A) = -4\pi\rho$$

or

$$\nabla^2 A = -4\pi\rho.$$

This is known as Poisson's equation.

In free space, $\rho = 0$.

$$\therefore \nabla^2 A = 0.$$

This is known as Laplace's equation.

Problem 174. Show that $\text{curl } \frac{a \times r}{r^3} = -\frac{a}{r^3} + \frac{3r}{r^5} (a \cdot r)$, where a is a constant vector and $r = xi + yj + zk$.

$$\begin{aligned} \text{Here, } \text{curl } \frac{a \times r}{r^3} &= \nabla \times \frac{a \times r}{r^3} = \left(i \frac{\partial}{\partial x} + j \frac{\partial}{\partial y} + k \frac{\partial}{\partial z} \right) \times \frac{a \times r}{r^3} \\ &= i \times \frac{\partial}{\partial x} \left(\frac{a \times r}{r^3} \right) + j \times \frac{\partial}{\partial y} \left(\frac{a \times r}{r^3} \right) + k \times \frac{\partial}{\partial z} \left(\frac{a \times r}{r^3} \right) \end{aligned}$$

$$\text{Now } \frac{\partial}{\partial x} \left(\frac{a \times r}{r^3} \right) = -\frac{3}{r^4} \frac{\partial r}{\partial x} a \times r + \frac{1}{r^3} \left(a \times \frac{\partial r}{\partial x} \right) = -\frac{3x}{r^5} a \times r + \frac{a}{r^4} \times i$$

$$\therefore r = xi + yj + zk \text{ gives } \frac{\partial r}{\partial x} = i$$

and

$$r^2 = x^2 + y^2 + z^2 \text{ gives } \frac{\partial r}{\partial x} = \frac{x}{r}$$

$$\begin{aligned} \therefore i \times \frac{\partial}{\partial x} \left(\frac{a \times r}{r^3} \right) &= -\frac{3x}{r^5} [i \times (a \times r)] + \frac{1}{r^3} i \times (a \times i) \\ &= -\frac{3x}{r^5} [(i \cdot r) a - (i \cdot a) r] + \frac{1}{r^3} [(i \cdot i) a - (i \cdot a) i] \end{aligned}$$

$$= -\frac{3x}{r^5} [xa - (i \cdot a) r] + \frac{1}{r^3} [a - (i \cdot a) i]$$

$$= -\frac{3x^2}{r^5} a + \frac{3r}{r^5} (ix \cdot a) + \frac{a}{r^3} - \frac{(i \cdot a) i}{r^3}.$$

$$\text{Similarly, } j \times \frac{\partial}{\partial y} \left(\frac{a \times r}{r^3} \right) = -\frac{3y^2}{r^5} a + \frac{3r}{r^5} (jy \cdot a) + \frac{a}{r^3} - \frac{(j \cdot a) j}{r^3}$$

$$k \times \frac{\partial}{\partial z} \left(\frac{a \times r}{r^3} \right) = -\frac{3z^2}{r^5} a + \frac{3r}{r^5} (kz \cdot a) + \frac{a}{r^3} - \frac{(k \cdot a) k}{r^3}.$$

$$i \times \frac{\partial}{\partial x} \left(\frac{a \times r}{r^3} \right) + j \times \frac{\partial}{\partial y} \left(\frac{a \times r}{r^3} \right) + k \times \frac{\partial}{\partial z} \left(\frac{a \times r}{r^3} \right)$$

$$= -\frac{3(x^2 + y^2 + z^2)}{r^5} a + \frac{3r}{r^5} \{xi + yj + zk\} \cdot a$$

$$+ \frac{3a}{r^3} - \frac{1}{r^3} [(i \cdot a) i + (j \cdot a) j + (k \cdot a) k]$$

$$= -\frac{3a}{r^3} + \frac{3r}{r^5} r \cdot a + \frac{3a}{r^3} - \frac{a}{r^3} \text{ since if } a = a_1 i + a_2 j + a_3 k,$$

$$i \cdot a = a_1, j \cdot a = a_2, k \cdot a = a_3.$$

$$\therefore (i \cdot a) i + (j \cdot a) j + (k \cdot a) k = a_1 i + a_2 j + a_3 k = a$$

$$= -\frac{a}{r^3} + \frac{3r}{r^5} (a \cdot r). \quad \dots (2)$$

Hence from (1) and (2),

$$\text{Curl } \frac{a \times r}{r^3} = -\frac{a}{r^3} + \frac{3r}{r^5} (a \cdot r).$$

Problem 175. Give example in support that vector methods have been used to results in a simple and elegant form.

Ever since the development of Quaternion analysis by W.R. Hamilton and of 'Ausdehnungslehre' by Grassman, it has been a growing feeling that new methods and ideas be applied more simply and more directly to many of the conceptions of geometry, mechanics and mathematical physics, than those long adopted. The methods of Vector Analysis are adopted on the grounds of simplicity, simplicity and directness.

As an example, Faraday the great physicist with his mind's eye visualized the force emerging from the magnet and so he had a visual conception of the power in which the electro-magnetic waves were travelling through the ether and him and so divergence and divergence theorem to him had simple meaning.

Numerous other examples can be quoted from geometry, mechanics and mathematical physics.

Problem 176. Show that

(a) The vector product of two vectors is a vector.

(b) The gradient of a scalar function is a vector.

(c) The divergence of a vector function is a scalar. (Agra, 1965)

Problem 177. What is Green's theorem. Use it to solve the equation.

$$\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} + \frac{\partial^2 \psi}{\partial z^2} = -4\pi\rho(x, y, z). \quad (\text{Agra, 1965})$$

Problem 178. (a) Prove that $(A \times \nabla) \times R = -2A$

where A is any vector field and R is a vector drawn from the point $P: (x, y, z)$.

- (b) Find the directional derivative of the function $\phi(x, y, z) = 2xy + z^2$ in the direction of the vector $\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}$ at the point $(1, -1, 3)$.
- (c) If the divergence of a vector field $\mathbf{H}$ vanishes, show that it can be expressed as the curl of a vector field $\mathbf{A}$. (Bombay, 1965)
- (a) Let $\mathbf{A} = A_x \mathbf{i} + A_y \mathbf{j} + A_z \mathbf{k}$ and $\mathbf{R} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$

$$\text{then } \mathbf{A} \times \nabla = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ A_x & A_y & A_z \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \end{vmatrix} = \left(\frac{\partial A_y}{\partial z} - \frac{\partial A_z}{\partial y} \right) \mathbf{i} + \left(\frac{\partial A_z}{\partial x} - \frac{\partial A_x}{\partial z} \right) \mathbf{j} + \left(\frac{\partial A_x}{\partial y} - \frac{\partial A_y}{\partial x} \right) \mathbf{k}$$

$$\text{So that } (\mathbf{A} \times \nabla) \times \mathbf{R} = \left\{ \left(\frac{\partial A_y}{\partial z} - \frac{\partial A_z}{\partial y} \right) \mathbf{i} + \left(\frac{\partial A_z}{\partial x} - \frac{\partial A_x}{\partial z} \right) \mathbf{j} + \left(\frac{\partial A_x}{\partial y} - \frac{\partial A_y}{\partial x} \right) \mathbf{k} \right\} \times (x\mathbf{i} + y\mathbf{j} + z\mathbf{k})$$

$$= -2(A_x \mathbf{i} + A_y \mathbf{j} + A_z \mathbf{k}) = -2\mathbf{A}$$

It is also easy to show that $(\mathbf{A} \times \nabla) \cdot \mathbf{R} = 0$

(b) $\phi = 2xy + z^2$

$$\therefore \nabla \phi = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) (2xy + z^2) = 2y \mathbf{i} + 2x \mathbf{j} + 2z \mathbf{k}$$

$$= -2 \mathbf{i} + 2 \mathbf{j} + 6 \mathbf{k} \text{ at } x=1, y=-1, z=3$$

A unit vector in the direction of $\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}$ is $\frac{\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}}{\sqrt{1^2 + 2^2 + 2^2}} = \frac{1}{3}(\mathbf{i} + 2\mathbf{j} + 2\mathbf{k})$

Since the directional derivative along a unit vector $\mathbf{u}$ is $\mathbf{u} \cdot \text{grad } \phi$, therefore the required directional derivative in the existing case is

$$= \frac{1}{3}(\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}) \cdot (-2\mathbf{i} + 2\mathbf{j} + 6\mathbf{k})$$

$$= \frac{1}{3}[-2 + 4 + 12] = \frac{14}{3}$$

(c) It is easy to show that $\text{div } \mathbf{H} = 0 \Rightarrow \mathbf{H} = \text{curl } \mathbf{A}$ by Theorem (3) of §1.62.

Problem 179. (a) Prove the divergence theorem of Gauss.

(b) Prove that $\iint_S (\mathbf{N} \times \mathbf{F}) \cdot d\mathbf{S} = \iiint_V (\nabla \times \mathbf{F}) \cdot d\mathbf{V}$

where $\mathbf{F}$ is a vector field and $\mathbf{N}$ is the normal unit vector to surface S enclosing the volume. (Bombay, 1965)

These are well known theorems.

Problem 180. (a) Prove the following by using vector methods:

(1) The medians of a triangle meet in a point of trisection of each other.

(2) $\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$

(b) Prove that

(1) $[\mathbf{A} \times \mathbf{B}] \times [\mathbf{A} \times \mathbf{C}] = ([\mathbf{A} \times \mathbf{B}] \cdot \mathbf{C}) \mathbf{A}$

(2) $\nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2}{\partial \phi^2}$

(a) — (1) See Problem 14 (b).

(Nagpur, 1965; Agri, 1965)

(2) Take $\mathbf{i}, \mathbf{j}$ unit vectors along OX, OY two mutually perpendicular axes and consider two coplanar lines OA and OB making angles α and β with OX . If $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}}$ be unit vectors along OA and OB respectively then

$$\hat{\mathbf{a}} = \cos \alpha \mathbf{i} + \sin \alpha \mathbf{j}$$

$$\hat{\mathbf{b}} = \cos \beta \mathbf{i} + \sin \beta \mathbf{j}$$

If $\hat{\mathbf{e}}$ be a unit vector perpendicular to $\hat{\mathbf{a}}$ and $\hat{\mathbf{b}}$ both, then

$$\hat{\mathbf{b}} \times \hat{\mathbf{a}} = 1.1 \sin(\alpha + \beta) \hat{\mathbf{e}}$$

$$\text{i.e., } \sin(\alpha + \beta) \hat{\mathbf{e}} = (\cos \beta \mathbf{i} + \sin \beta \mathbf{j}) \times (\cos \alpha \mathbf{i} + \sin \alpha \mathbf{j})$$

$$= (\sin \alpha \cos \beta + \cos \alpha \sin \beta) \hat{\mathbf{e}}$$

$$\therefore \mathbf{i} \times \mathbf{j} = \hat{\mathbf{e}}$$

$$\therefore \sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta.$$

(b) (1) L.H.S. $=[\mathbf{A} \times \mathbf{B}] \times [\mathbf{A} \times \mathbf{C}]$

$$= ([\mathbf{A} \times \mathbf{B}] \cdot \mathbf{C}) \mathbf{A} - ([\mathbf{A} \times \mathbf{B}] \cdot \mathbf{A}) \mathbf{C}$$

$$= ([\mathbf{A} \times \mathbf{B}] \cdot \mathbf{C}) \mathbf{A} \quad \because [\mathbf{A} \times \mathbf{B}] \cdot \mathbf{A} = [\mathbf{A} \mathbf{B} \mathbf{A}] = 0.$$

(2) We have $\nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ in cartesian coordinates.

The transformations to polar coordinates are

$$x = r \sin \theta \cos \phi, y = r \sin \theta \sin \phi, z = r \cos \theta \quad \dots (1)$$

Let us first change x, y, z to u, ϕ, z (cylindrical coordinates) by the transformation

$$x = u \cos \phi, y = u \sin \phi, z = z \quad \dots (2)$$

So that $u = \sqrt{x^2 + y^2}$ and $\phi = \tan^{-1} \frac{y}{x}$

Also, $\frac{\partial u}{\partial x} = \cos \phi, \frac{\partial u}{\partial y} = \sin \phi, \frac{\partial \phi}{\partial x} = -\frac{\sin \phi}{r}, \frac{\partial \phi}{\partial y} = \frac{\cos \phi}{r}$

Now, $\frac{\partial v}{\partial x} = \frac{\partial v}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial v}{\partial \phi} \frac{\partial \phi}{\partial x} = \cos \phi \frac{\partial v}{\partial u} - \frac{\sin \phi}{r} \frac{\partial v}{\partial \phi}$, similarly $\frac{\partial v}{\partial y} = \sin \phi \frac{\partial v}{\partial u} + \frac{\cos \phi}{r} \frac{\partial v}{\partial \phi}$

$$\left(\frac{\partial v}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial y} \right)^2 = \left(\frac{\partial v}{\partial u} \right)^2 + \left(\frac{1}{u} \frac{\partial v}{\partial \phi} \right)^2 \quad \dots (3)$$

Again $\frac{\partial^2 v}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial v}{\partial x} \right) = \left(\cos \phi \frac{\partial}{\partial u} - \frac{\sin \phi}{u} \frac{\partial}{\partial \phi} \right) \left(\cos \phi \frac{\partial v}{\partial u} - \frac{\sin \phi}{u} \frac{\partial v}{\partial \phi} \right)$

$$= \cos \phi \left(\cos \phi \frac{\partial^2 v}{\partial u^2} - \frac{\sin \phi}{u} \frac{\partial^2 v}{\partial u \partial \phi} + \frac{\sin \phi}{u^2} \frac{\partial v}{\partial \phi} \right) - \frac{\sin \phi}{u}$$

$$\left(\cos \phi \frac{\partial^2 v}{\partial u \partial \phi} - \sin \phi \frac{\partial v}{\partial r} - \frac{\sin \phi}{u} \frac{\partial^2 v}{\partial \phi^2} - \frac{\cos \phi}{u} \frac{\partial v}{\partial \phi} \right)$$

$$= \left(\cos^2 \phi \frac{\partial^2 v}{\partial u^2} - \frac{2 \sin \phi \cos \phi}{u} \frac{\partial^2 v}{\partial u \partial \phi} + \frac{\sin^2 \phi}{u^2} \frac{\partial^2 v}{\partial \phi^2} + \frac{\sin^2 \phi}{u} \frac{\partial v}{\partial u} \right) + \frac{2 \sin \phi \cos \phi}{u^2} \frac{\partial v}{\partial \phi}$$




Fig. 1.67

Similarly $\frac{\partial^2 v}{\partial y^2} = \sin^2 \phi \frac{\partial^2 v}{\partial u^2} + \frac{2 \sin \phi \cos \phi}{u} \frac{\partial^2 v}{\partial u \partial \phi} + \frac{\cos^2 \phi}{u^2} \frac{\partial^2 v}{\partial \phi^2} + \frac{\cos^2 \phi}{u^2} \frac{\partial v}{\partial \phi}$

Adding $\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = \frac{\partial^2 v}{\partial u^2} + \frac{1}{u} \frac{\partial v}{\partial u} + \frac{1}{u^2} \frac{\partial^2 v}{\partial \phi^2}$

Adding $\frac{\partial^2 v}{\partial z^2}$ on either side of (4),

$$\nabla^2 v = \frac{\partial^2 v}{\partial u^2} + \frac{1}{u^2} \frac{\partial^2 v}{\partial \phi^2} + \frac{\partial^2 v}{\partial z^2} + \frac{1}{u} \frac{\partial v}{\partial u}$$

Now by putting $u = r \sin \theta$, $z = r \cos \theta$, the transformations (2) reduce to the form (1) and by applying (4), we get

$$\frac{\partial^2 v}{\partial u^2} + \frac{\partial^2 v}{\partial z^2} = \frac{\partial^2 v}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2 v}{\partial \theta^2} + \frac{1}{r} \frac{\partial v}{\partial r}$$

and $\frac{1}{u} \frac{\partial v}{\partial u} = \frac{1}{r \sin \theta} \left(\sin \theta \frac{\partial v}{\partial r} + \frac{\cos \theta}{r} \frac{\partial v}{\partial \theta} \right)$

Substituting values from (6) and (7) in (5) we find

$$\begin{aligned} \nabla^2 v &= \frac{\partial^2 v}{\partial r^2} + \frac{2}{r} \frac{\partial v}{\partial r} + \frac{1}{r^2} \frac{\partial^2 v}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial v}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 v}{\partial \phi^2} \\ &= \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial v}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial v}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 v}{\partial \phi^2} \end{aligned}$$

i.e. $\nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2}{\partial \phi^2}$

Problem 181. Evaluate the following :

(i) $\frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-\frac{x^2}{a^2}} e^{ikx} dx$

(ii) $\frac{1}{(2\pi)^3} \iiint \frac{e^{-r/a}}{r} e^{i\mathbf{k} \cdot \mathbf{r}} dx dy dz$ [$r = \sqrt{x^2 + y^2 + z^2}$, $\mathbf{r} = (x, y, z)$]

Problem 182. (a) Starting from the definition $\nabla^2 \psi = \frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} + \frac{\partial^2 \psi}{\partial z^2}$ express $\nabla^2 \psi$ in polar and cylindrical coordinates.

(b) Write the general solution of the equation $\nabla^2 \psi = 0$ in polar coordinates.

(See Problem 180).

Problem 183. (a) Prove the following identities :

(i) $\vec{\nabla} \cdot (\vec{\nabla} \times \mathbf{v}) = 0$, (ii) $\vec{\nabla} \times (\vec{\nabla} \phi) = 0$

(b) Show that if a scalar point function ϕ depends only on the magnitude of position vector $\mathbf{r}$, then $\vec{\nabla} \phi = \frac{d\phi}{dr} \mathbf{e}_r$, $\mathbf{e}_r$ being the unit vector in the direction of $\mathbf{r}$.

(c) Show that $\vec{\nabla} \times \mathbf{r} = 0$

(Here $\mathbf{v}$ is a vector function of $\mathbf{r}$ and $\vec{\nabla} = \mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z}$ in the usual notation.)

Problem 184. (a) Define (i) The dot product and (ii) The cross product of vectors. Show that $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = \mathbf{B} (\mathbf{A} \cdot \mathbf{C}) - \mathbf{C} (\mathbf{A} \cdot \mathbf{B})$

4. If ϕ is any scalar and $\mathbf{A}$ any vector, then show that

(i) $\text{div } \phi \mathbf{A} = \mathbf{A} \cdot \text{grad } \phi + \phi \text{ div } \mathbf{A}$

(ii) $\text{curl } \phi \mathbf{A} = \phi \text{ curl } \mathbf{A} - \mathbf{A} \times \text{grad } \phi$

[Vikram 1967, (i) 69 also]

Problem 185. Explain how the Laplacian operator ∇^2 can be expressed in any one of orthogonal curvilinear coordinates. Hence express it in cylindrical and spherical polar coordinates. (Vikram, 1967)

See §1.45 and §1.46.

Problem 186. Using the Theorem of Gauss prove the following identities:

(i) $\int_V \vec{\nabla} \cdot \phi \mathbf{d}\tau = \oint_S \phi \mathbf{n} \cdot d\mathbf{S}$, (ii) $\int_V \vec{\nabla} \times \mathbf{F} \cdot d\mathbf{\tau} = \oint_S \mathbf{n} \times \mathbf{F} \cdot d\mathbf{S}$

Here S is a closed surface enclosing the volume V and $\mathbf{n}$ is the unit normal to the surface element dS . (Agra, 1968)

Problem 187. Explain the meanings of the operations $\vec{\nabla} \cdot \vec{A}$, $\vec{\nabla} \cdot \vec{A}$ and $\vec{\nabla} \times \vec{A}$. Show that

$\vec{\nabla} \cdot (\mathbf{A} \times \mathbf{B}) = (\mathbf{B} \cdot \vec{\nabla}) \mathbf{A} - (\mathbf{A} \cdot \vec{\nabla}) \mathbf{B} - \mathbf{B} (\vec{\nabla} \cdot \mathbf{A}) + \mathbf{A} (\vec{\nabla} \cdot \mathbf{B})$ (Agra, 1972)

Meanings are clear from §1.24, §1.28 etc. For second part see §1.33.

Problem 188. (a) Prove the Green's theorem

$$\iiint_T (\phi \nabla^2 \psi - \psi \nabla^2 \phi) dx = \iint_S \left(\phi \frac{\partial \psi}{\partial n} - \psi \frac{\partial \phi}{\partial n} \right) ds$$

4. A vector field is given by $\mathbf{A} = (x^2 + xy^2) \mathbf{i} + (y^2 + x^2y) \mathbf{j}$

show that the field is irrotational and find the scalar potential. (Bombay, 1970)

First part is the well known theorem and for second show $\vec{\nabla} \times \mathbf{A} = 0$.

Problem 189. Establish an expression for the components of the curl of a vector. Express it in orthogonal curvilinear coordinates and hence obtain the radial component of $\text{curl } \mathbf{A}$ in spherical polar coordinates (r, θ, ϕ) . (Agra, 1974)

See §1.42, §1.45.

Problem 190. Find the divergence, the gradient of the divergence and the curl of

where $\mathbf{r} = r \hat{\mathbf{r}}$, where $\mathbf{r}$ is a position vector. (Agra, 1974)

See Problem 98.

Problem 191. If $\text{div } \mathbf{A} = 0$, show that $\text{curl } \text{curl } \mathbf{A} = -\nabla^2 \mathbf{A}$ (Agra, 1975)

Problem 192. (a) Clearly explain the physical significance of divergence and express them in spherical polar and circular cylindrical coordinates.

(b) If a vector function $\mathbf{F}$ depends on both space coordinate (x, y, z) and time t then

$$d\mathbf{F} = (d\mathbf{r} \cdot \vec{\nabla}) \mathbf{F} + \frac{\delta}{\delta t} \mathbf{F} dt \quad (\text{Rohilkhand, 1976})$$

Problem 193. Define Solenoidal and Irrotational vector fields. Show that a vector field is irrotational for every n , but is solenoidal only if $n = -3$, where $\mathbf{r}$ is position vector of a particle.

(b) The gravitational field of mass m , the potential at a distance r is given by

obtain the components of force and show that its curl is zero.

(c) See Problems 111 and 169.

(Rohilkhand, 1977)

CHAPTER 2

MATRICES

2.1 DEFINITIONS AND NOTATIONS

A set of mn numbers, real or complex, arranged in a rectangular array of m rows and n columns such as

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

is called a **matrix** of order $m \times n$.

In other words a scheme of detached coefficients a_{ij} arranged in m rows and n columns is called a **matrix of order m by n** or an **$m \times n$ matrix** or a **matrix of type $m \times n$** .

In case $m=n$, the rectangular array becomes a square and so a **matrix** having number of rows and number of columns equal is called a **Square Matrix** of order n .

Any matrix obtained by deleting any number of rows and any number of columns from a given matrix is said to be a **Sub-Matrix** of the given matrix.

The mn numbers a_{ij} , ($i=1, 2, \dots, m$; $j=1, 2, \dots, n$, $i \neq j$) constituting the $m \times n$ matrix are called its **elements** or **constituents**. The elements a_{ii} ($i=j$) of a square matrix A are called its **diagonal elements**.

and their sum as **trace** of A denoted by $\text{tr. } A = \sum_{i=1}^n a_{ii}$

A matrix is usually denoted by capital letters like A (in clear type) or $[a_{ij}]$, where a_{ij} represents the (i, j) th element i.e., the element in the i th row and j th column of the matrix.

Thus an $m \times n$ matrix may be expressed as

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \quad \begin{array}{l} \text{where } 1 \leq i \leq m \\ \text{and } 1 \leq j \leq n \\ \text{but } i \neq j \end{array}$$

We have so far used only a pair of brackets i.e. $[]$ to denote a matrix, but a pair of parentheses i.e. $()$ and double bars i.e. $\| \|$ are also sometimes used to indicate a matrix.

A matrix having all of its elements zero is said to be a **Null Matrix** and denoted by O e.g.

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

A square matrix of order n having all its diagonal elements unity and zero elements everywhere else is called a **unit matrix** or an **identity matrix** and denoted by I_n . Thus

$$I_n = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}$$

It is possible that a matrix may have only a single row or a single column such as

$$[a_1, a_2, \dots, a_p] \quad \text{and} \quad \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_q \end{bmatrix}$$

The first one being a matrix of order $1 \times p$ is called a **row matrix** and the second one being a matrix of order $q \times 1$ is called a **column matrix**.

A single element constitutes a matrix of order 1×1 .

In relation to matrices, the numbers are usually known as scalars; they behave as operators exactly like ordinary numbers as multipliers and hence are called scalars.

ILLUSTRATIVE EXAMPLES

$\begin{bmatrix} 2 & 3 & -1 \\ 4 & -5 & 6 \end{bmatrix}$ is a matrix of order 2×3 .

$\begin{bmatrix} 2 & -3 & 4 \\ 5 & 6 & -2 \\ 1 & 0 & 4 \end{bmatrix}$ is a square matrix of order 3.

$\begin{bmatrix} 2 & 0 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ is a sub-matrix of the matrix $\begin{bmatrix} 2 & 0 & 3 & -1 & 2 \\ 4 & 5 & 6 & 7 & 8 \\ 9 & 4 & 2 & 1 & 5 \end{bmatrix}$

4, 0, 6 are the diagonal elements of the matrix $A = \begin{bmatrix} 4 & 5 & 6 \\ 2 & 0 & 3 \\ 2 & -5 & 6 \end{bmatrix}$ whose elements are 4, 5, 6

2, 3, 2, -5, 6 i.e. if $[a_{ij}] = \begin{bmatrix} 4 & 5 & 6 \\ 2 & 0 & 3 \\ 2 & -5 & 6 \end{bmatrix}$ then $a_{11}=4$, $a_{12}=5$, $a_{13}=6$ etc.

Also trace of A i.e. $\text{tr } A = 4 + 0 + 6 = 10$.

$\begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ is a 3×2 null matrix.

$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ is a unit matrix of order 3.

7. $\begin{bmatrix} 2 & 3 & -5 \end{bmatrix}$ is a 1×3 row matrix.

8. $\begin{bmatrix} 2 \\ 1 \\ 5 \\ 7 \end{bmatrix}$ is a 4×1 column matrix.

2.2. EQUALITY OF MATRICES

Two matrices **A** and **B** defined as

$$\mathbf{A} = [a_{ij}] \text{ and } \mathbf{B} = [b_{ij}]$$

are said to be equal if both are of the same order $m \times n$ i.e. **A** has the same number of rows and columns as **B** and each element a_{ij} of **A** is equal to the corresponding element b_{ij} of **B** i.e. $a_{ij} = b_{ij}$ for each pair of subscripts i and j .

Hence for equality, the two matrices must be identical in every respect or broadly speaking, the two matrices are equal if and only if one is a duplicate of the other.

ILLUSTRATIVE EXAMPLES

1. If $\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}$

then $\mathbf{A} = \mathbf{B}$ if and only if

$$a_{11} = b_{11}, a_{12} = b_{12}, a_{21} = b_{21}, a_{22} = b_{22}, a_{31} = b_{31}, a_{32} = b_{32}.$$

2. The matrices $\begin{bmatrix} 1 & 2 \\ 3 & -4 \end{bmatrix}$ and $\begin{bmatrix} 1 & 2 & 5 \\ 3 & -4 & 0 \end{bmatrix}$ being of different

order are not comparable for equality.

3. The matrices $\begin{bmatrix} 2 & 3 \\ 0 & 1 \\ 5 & 4 \end{bmatrix}$ and $\begin{bmatrix} 2 & 3 \\ 6 & 1 \\ 5 & 4 \end{bmatrix}$ are comparable but not

equal as the element of 2nd row and 1st column of the first matrix is not equal to the corresponding element of the second matrix.

4. The matrices $\begin{bmatrix} 2 & 1 & 5 \\ 0 & 9 & 7 \end{bmatrix}$ and $\begin{bmatrix} 2 & 1 & 5 \\ 0 & 9 & 7 \end{bmatrix}$ are equal.

5. The matrices $\begin{bmatrix} 4 & 5 & 27 \\ 6 & 0 & 3 \end{bmatrix}$ and $\begin{bmatrix} 2^2 & 5 & 3^3 \\ 3.2 & 0 & \sqrt{9} \end{bmatrix}$ are equal.

COROLLARY. Equivalence Relation on Matrices

If there are three matrices **A**, **B**, **C** such that they satisfy the following three properties.

(1) Reflexivity $\mathbf{A} = \mathbf{A}$

(2) Symmetry $\mathbf{A} = \mathbf{B}$ implies that $\mathbf{B} = \mathbf{A}$

(3) Transitivity $\mathbf{A} = \mathbf{B}$ and $\mathbf{B} = \mathbf{C}$ imply that $\mathbf{A} = \mathbf{C}$

Then the equality of matrices is said to form an equivalence relation.

3. ADDITION OF MATRICES

Two matrices $\mathbf{A} = [a_{ij}]$ and $\mathbf{B} = [b_{ij}]$ are said to be conformable for addition if they are of the same order i.e. they have the same number of rows and the same number of columns. The sum of the two matrices **A** and **B** is then defined as the matrix each of whose elements is the sum of the corresponding elements of **A** and **B** i.e.

$$\mathbf{A} + \mathbf{B} = [a_{ij}] + [b_{ij}] \\ = [a_{ij} + b_{ij}]$$

For example

$$\text{If } \mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1n} \\ b_{21} & b_{22} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots \\ b_{m1} & b_{m2} & \dots & b_{mn} \end{bmatrix}$$

$$\mathbf{A} + \mathbf{B} = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \dots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \dots & a_{2n} + b_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \dots & a_{mn} + b_{mn} \end{bmatrix}$$

As an other example if

$$\mathbf{A} = \begin{bmatrix} 2 & 0 & 3 \\ -4 & 1 & 5 \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} -1 & 2 & 0 \\ 3 & -4 & -5 \end{bmatrix}$$

$$\text{then } \mathbf{A} + \mathbf{B} = \begin{bmatrix} 2-1 & 0+2 & 3+0 \\ -4+3 & 1-4 & 5-5 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ -1 & -3 & 0 \end{bmatrix}$$

COROLLARY 1. $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr} \mathbf{A} + \text{tr} \mathbf{B}$ e.g. if $\mathbf{A} = \begin{bmatrix} -1 & 2 \\ 2 & 3 \end{bmatrix}$ and

$$\mathbf{B} = \begin{bmatrix} 2 & 3 \\ 0 & 1 \end{bmatrix} \text{ then } \text{tr}(\mathbf{A} + \mathbf{B}) = 1 + 4 = 5 = 2 + 3 \\ = \text{tr} \mathbf{A} + \text{tr} \mathbf{B}$$

COROLLARY 2. Subtraction. The difference of two matrices **A** and **B** which are conformable for addition may be defined as the sum of the matrices **A** and $(-\mathbf{B})$ where $(-\mathbf{B})$ is the matrix obtained by multiplying every element of **B** by -1 . Thus

$$\mathbf{A} - \mathbf{B} = \mathbf{A} + (-\mathbf{B})$$

The elements of the difference matrix $\mathbf{A} - \mathbf{B}$ are obtained by subtracting the elements of **B** from the corresponding elements of **A**.

So if $\mathbf{A} = [a_{ij}]$ and $\mathbf{B} = [b_{ij}]$, then

$$\mathbf{A} - \mathbf{B} = [a_{ij}] - [b_{ij}] = [a_{ij} - b_{ij}]$$

As an illustrative example if

$$\mathbf{A} = \begin{bmatrix} 2 & 5 & -3 \\ 0 & 7 & -8 \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} 4 & 0 & 3 \\ -2 & 0 & 5 \end{bmatrix}$$

$$\text{then } \mathbf{A} - \mathbf{B} = \begin{bmatrix} 2-4 & 5-0 & -3-3 \\ 0-(-2) & 7-0 & -8-5 \end{bmatrix} = \begin{bmatrix} -2 & 5 & -6 \\ 2 & 7 & -13 \end{bmatrix}$$

COROLLARY 3. Multiplication of a matrix by a number (scalar)

If **A** is a matrix of any order say $m \times n$ defined as

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & \dots & a_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & \dots & a_{mn} \end{bmatrix}$$

then the addition law follows

$$A + A = \begin{bmatrix} a_{11} & a_{12} & \dots & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & \dots & a_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & \dots & a_{mn} \end{bmatrix} + \begin{bmatrix} a_{11} & a_{12} & \dots & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & \dots & a_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & \dots & a_{mn} \end{bmatrix}$$

$$\text{or } 2A = 2 \begin{bmatrix} a_{11} & a_{12} & \dots & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & \dots & a_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & \dots & a_{mn} \end{bmatrix} = \begin{bmatrix} 2a_{11} & 2a_{12} & \dots & \dots & 2a_{1n} \\ 2a_{21} & 2a_{22} & \dots & \dots & 2a_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ 2a_{m1} & 2a_{m2} & \dots & \dots & 2a_{mn} \end{bmatrix}$$

Similarly

$$2A + A = 3A = 3 \begin{bmatrix} a_{11} & a_{12} & \dots & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & \dots & a_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & \dots & a_{mn} \end{bmatrix} = \begin{bmatrix} 3a_{11} & 3a_{12} & \dots & \dots & 3a_{1n} \\ 3a_{21} & 3a_{22} & \dots & \dots & 3a_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ 3a_{m1} & 3a_{m2} & \dots & \dots & 3a_{mn} \end{bmatrix}$$

or in general if k is a number, real or complex and A is a matrix then kA the matrix obtained by multiplying every element of A by k is said to be the scalar multiple of A .

As an illustrative example if $A = \begin{bmatrix} 0 & 2 & 3 \\ -1 & 5 & 4 \end{bmatrix}$, then

$$2A = 2 \begin{bmatrix} 0 & 2 & 3 \\ -1 & 5 & 4 \end{bmatrix} = \begin{bmatrix} 0 & 4 & 6 \\ -2 & 10 & 8 \end{bmatrix}$$

$$3A = 3 \begin{bmatrix} 0 & 2 & 3 \\ -1 & 5 & 4 \end{bmatrix} = \begin{bmatrix} 0 & 6 & 9 \\ -3 & 15 & 12 \end{bmatrix}$$

It is easy to verify that

$$A + A = 2A = 5A - 3A \text{ etc.}$$

and similarly $(3+i)A = A + A(1+i) + A$ etc., where $i = \sqrt{-1}$

COROLLARY 4. Linear Combination of Matrices

The same law of addition can be applied to combine any number of matrices. Thus if $A, B, C, \dots, K$ be a finite number of matrices each of order $m \times n$ (say) and $\alpha, \beta, \gamma, \dots, k$ be scalars then

$$\alpha A + \beta B + \gamma C + \dots + kK = [\alpha a_{ij} + \beta b_{ij} + \gamma c_{ij} + \dots + k k_{ij}]$$

Where $A = [a_{ij}]$, $B = [b_{ij}]$, $C = [c_{ij}]$, $\dots$, $K = [k_{ij}]$

As an illustrative example if

$$A = \begin{bmatrix} 2 & 3 & 0 \\ 4 & -1 & 2 \end{bmatrix}, B = \begin{bmatrix} 1 & -2 & 3 \\ 0 & 4 & 5 \end{bmatrix}, C = \begin{bmatrix} 2 & -1 & 5 \\ 3 & 0 & 4 \end{bmatrix}$$

then

$$3A - 4B + 2C = \begin{bmatrix} 6 & -4 & +4 & 9 & -(-8) & +(-2) & 0 & -12 & +10 \\ 12 & -0 & +6 & -3 & -16 & +0 & 6 & -20 & +8 \end{bmatrix} = \begin{bmatrix} 6 & 15 & -2 \\ 18 & -19 & -6 \end{bmatrix}$$

COROLLARY 5. $\text{tr.}(\lambda A) = \lambda \text{tr.} A$, λ being scalar e.g. if $A = \begin{bmatrix} 2 & -2 \\ 0 & -1 \end{bmatrix}$ so

$$\lambda A = \begin{bmatrix} 2\lambda & -2\lambda \\ 0 & -\lambda \end{bmatrix} \text{ clearly } \text{tr.}(\lambda A) = 2\lambda - \lambda = \lambda \text{ and } \lambda \text{tr.} A = \lambda(2 - 1) = \lambda$$

so $\text{tr.} \lambda A = \lambda \text{tr.} A$.

1.4. PROPERTIES OF MATRIX-ADDITION

(1) The Commutative Law

If A and B are two matrices of the same order say $m \times n$, then

$$A + B = B + A$$

Let $A = [a_{ij}]$ and $B = [b_{ij}]$, $i = 1, 2, \dots, m$
 $j = 1, 2, \dots, n$

We have

$$\begin{aligned} A + B &= [a_{ij}] + [b_{ij}] = [a_{ij} + b_{ij}] \\ &= [b_{ij} + a_{ij}] \text{ since } b_{ij} \text{ and } a_{ij} \text{ are scalars} \\ &= [b_{ij}] + [a_{ij}] \\ &= B + A \end{aligned}$$

i.e. the commutative law of addition holds.

(2) The Associative Law

If A, B, C are three matrices of the same order say $m \times n$, then

$$(A + B) + C = A + (B + C)$$

Let $A = [a_{ij}]$, $B = [b_{ij}]$ and $C = [c_{ij}]$, $i = 1, 2, \dots, m$
 $j = 1, 2, \dots, n$

We have $(A + B) + C = ([a_{ij}] + [b_{ij}]) + [c_{ij}]$

$$\begin{aligned} &= ([a_{ij} + b_{ij}]) + [c_{ij}] \\ &= [(a_{ij} + b_{ij}) + c_{ij}] \\ &= [a_{ij} + (b_{ij} + c_{ij})], a_{ij}, b_{ij}, c_{ij} \text{ being scalars} \\ &= [a_{ij}] + [(b_{ij} + c_{ij})] \\ &= A + (B + C) \end{aligned}$$

i.e. the associative law of addition holds.

(3) The Distributive Law

If A and B are two matrices of the same order say $m \times n$ and k is a scalar then $k(A + B) = kA + kB$

Let $A = [a_{ij}]$ and $B = [b_{ij}]$, $i = 1, 2, \dots, m$
 $j = 1, 2, \dots, n$

We have, $k(A+B) = k[a_{ij} + b_{ij}]$
 $= [k(a_{ij} + b_{ij})]$
 $= [ka_{ij} + kb_{ij}]$
 $= [ka_{ij}] + [kb_{ij}]$
 $= k[a_{ij}] + k[b_{ij}], k \text{ being a scalar}$
 $= kA + kB$

i.e. the distributive law of addition holds.

[4] Existence of Additive Identity

If A be a matrix of any order say $m \times n$ and O a null matrix of the same order such that when it is added to A , leaves it unchanged

$$A + O = A$$

i.e. $A + O = A$
 then O is said to be the additive identity of A .

Its proof immediately follows from the fact that if $A = [a_{ij}]$ and O is a null matrix i.e. a matrix having each of its elements zero, then

$$A + O = [a_{ij} + 0]$$

$$= [a_{ij}] \text{ since a zero added to any scalar leaves it unchanged.}$$

$$= A$$

Because of this fact O is said to be an additive identity of A .

[5] Existence of Additive Inverse

If A be a matrix of any order say $m \times n$, and there exists a matrix $-A$ of the same order such that if it is added to A gives a null matrix O .

$$A + (-A) = O$$

i.e. $A + (-A) = O$
 then $(-A)$ is said to be the additive inverse of A .

Let $A = [a_{ij}]$
 Then, $-A = -[a_{ij}] = [-a_{ij}]$
 So that $A + (-A) = [a_{ij}] + [-a_{ij}]$
 $= [a_{ij} - a_{ij}]$
 $= O$

Because of this fact $(-A)$ is said to be an additive inverse of A .

[6] The Cancellation Law

If A, B, C are three matrices conformable for addition then the relation $A + B = A + C$

holds if and only if $B = C$

Let $A = [a_{ij}], B = [b_{ij}], C = [c_{ij}], i = 1, 2, \dots, m$
 $j = 1, 2, \dots, n.$

Then the relation $A + B = A + C$ follows that (i, j) th elements on either side are identically equal i.e.

$$a_{ij} + b_{ij} = a_{ij} + c_{ij}$$

which yields

$$b_{ij} = c_{ij}, \text{ since } a_{ij}, b_{ij}, c_{ij} \text{ all are scalars.}$$

(i, j)th element of $B = (i, j)$ th element C for all values of i and j .

$$\text{As such } B = C$$

Hence the relation $A + B = A + C$ holds if and only if $B = C$

Problem 1. If $A = \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \end{bmatrix}, B = \begin{bmatrix} 3 & -1 & 2 \\ 4 & 2 & 5 \end{bmatrix}$ and $C = \begin{bmatrix} 4 & 1 & 2 \\ 0 & 3 & 2 \end{bmatrix}$ then find

(i) $A + B$

(ii) $A - C$

(iii) $-2A$

(iv) $2A + 3B - 4C$

(i) We have

$$A + B = \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \end{bmatrix} + \begin{bmatrix} 3 & -1 & 2 \\ 4 & 2 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 1+3 & 2-1 & -3+2 \\ 5+4 & 0+2 & 2+5 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 1 & -1 \\ 9 & 2 & 7 \end{bmatrix}$$

(ii) We have

$$A - C = \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \end{bmatrix} - \begin{bmatrix} 4 & 1 & 2 \\ 0 & 3 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 1-4 & 2-1 & -3-2 \\ 5-0 & 0-3 & 2-2 \end{bmatrix}$$

$$= \begin{bmatrix} -3 & 1 & -5 \\ 5 & -3 & 0 \end{bmatrix}$$

(iii) We have

$$-2A = -2 \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} -2 & -4 & 6 \\ -10 & 0 & -4 \end{bmatrix}$$

(iv) We have

$$2A + 3B - 4C = 2 \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \end{bmatrix} + 3 \begin{bmatrix} 3 & -1 & 2 \\ 4 & 2 & 5 \end{bmatrix} - 4 \begin{bmatrix} 4 & 1 & 2 \\ 0 & 3 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 2+9-16 & 4-3-4 & -6+6-8 \\ 10+12-0 & 0+6-12 & 4+15-8 \end{bmatrix}$$

$$= \begin{bmatrix} -5 & -3 & -8 \\ 22 & -6 & 11 \end{bmatrix}$$

Problem 2. If $A = \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \\ 1 & -1 & 1 \end{bmatrix}, B = \begin{bmatrix} 3 & -1 & 2 \\ 4 & 2 & 5 \\ 2 & 0 & 3 \end{bmatrix}$
 and $C = \begin{bmatrix} 4 & 1 & 2 \\ 0 & 3 & 2 \\ 1 & -2 & 3 \end{bmatrix}$

Verify that $A + (B - C) = (A + B) - C$
and determine the matrix D such that $A + D = B$

We have

$$\begin{aligned} A + (B - C) &= \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \\ 1 & -1 & 1 \end{bmatrix} + \left\{ \begin{bmatrix} 3 & -1 & 2 \\ 4 & 2 & 5 \\ 2 & 0 & 3 \end{bmatrix} - \begin{bmatrix} 4 & 1 & 2 \\ 0 & 3 & 2 \\ 1 & -2 & 3 \end{bmatrix} \right\} \\ &= \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \\ 1 & -1 & 1 \end{bmatrix} + \begin{bmatrix} -1 & -2 & 0 \\ 4 & -1 & 3 \\ 1 & 2 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & -3 \\ 9 & -1 & 5 \\ 2 & 1 & 0 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} \text{and } (A + B) - C &= \left\{ \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \\ 1 & -1 & 1 \end{bmatrix} + \begin{bmatrix} 3 & -1 & 2 \\ 4 & 2 & 5 \\ 2 & 0 & 3 \end{bmatrix} \right\} - \begin{bmatrix} 4 & 1 & 2 \\ 0 & 3 & 2 \\ 1 & -2 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 1 & -1 \\ 9 & 2 & 7 \\ 3 & -1 & 4 \end{bmatrix} - \begin{bmatrix} 4 & 1 & 2 \\ 0 & 3 & 2 \\ 1 & -2 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & -3 \\ 9 & -1 & 5 \\ 2 & 1 & 0 \end{bmatrix} \end{aligned}$$

From (1) and (2) it follows that

$$A + (B - C) = (A + B) - C$$

Hence the required relation is verified.

Now given that

$$A + D = B$$

Adding $(-A)$ to both sides, we get

$$A + D - A = B - A$$

or $D + A - A = B - A$ by commutative law

or $D = B - A$ since $A - A = 0$ by existence of additive inverse

$$\begin{aligned} &= \begin{bmatrix} 3 & -1 & 2 \\ 4 & 2 & 5 \\ 2 & 0 & 3 \end{bmatrix} - \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \\ 1 & -1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 2 & -3 & 5 \\ -1 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix} \end{aligned}$$

Which is the required matrix.

2.5. MULTIPLICATION OF MATRICES

Two matrices A and B are conformable for multiplication if and only if the number of columns in A is equal to the number of rows in B . The product of the two matrices A and B denoted by AB is then defined as the matrix whose elements in the i th row and j th column is the sum of the products of the elements in the i th row of A by the corresponding elements in the j th column of B .

the sum of the products of the elements in the i th row of A by the corresponding elements in the j th column of B .

In other words the product AB of two matrices conformable for multiplication, is the matrix whose element in the i th row and j th column is the inner or scalar product of the i th row of A by the j th column of B , while the inner product or scalar product of two numbers x and y with components $x_1, x_2, \dots, x_n$ and $y_1, y_2, \dots, y_n$ is equal to

$$x_1 y_1 + x_2 y_2 + \dots + x_n y_n$$

It should be noted that inner product of two numbers with unequal numbers of components is not defined.

As an illustrative example if

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{bmatrix} \text{ and } B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{bmatrix}$$

then it is clear that the two matrices are conformable for multiplication since the number of columns in A is equal to the number of rows in B .

$$AB = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} + a_{13}b_{31} & a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} \\ a_{21}b_{11} + a_{22}b_{21} + a_{23}b_{31} & a_{21}b_{12} + a_{22}b_{22} + a_{23}b_{32} \\ a_{31}b_{11} + a_{32}b_{21} + a_{33}b_{31} & a_{31}b_{12} + a_{32}b_{22} + a_{33}b_{32} \\ a_{41}b_{11} + a_{42}b_{21} + a_{43}b_{31} & a_{41}b_{12} + a_{42}b_{22} + a_{43}b_{32} \end{bmatrix}$$

It is worth noting that the product BA is not defined, since the number of columns in B is not equal to the number of rows in A .

In the product AB , the matrix A is known as Prefactor and B as Post factor.

As an illustration in generalized form if

$$A = [a_{ij}], \text{ a matrix of order } m \times n$$

$$B = [b_{jk}], \text{ a matrix of order } n \times p$$

then $AB = C$ (say) is a matrix of order $m \times p$

$$C = [c_{ik}] \text{ is a matrix of order } m \times p \text{ such that}$$

$$c_{ik} = \sum_{j=1}^n a_{ij} b_{jk}$$

$$= a_{i1}b_{1k} + a_{i2}b_{2k} + \dots + a_{in}b_{nk}; \quad i=1, 2, \dots, m \text{ and } j=1, 2, \dots, p$$

Thus

$$C = \begin{bmatrix} c_{11} & c_{12} & \dots & c_{1p} \\ c_{21} & c_{22} & \dots & c_{2p} \\ \dots & \dots & \dots & \dots \\ c_{m1} & c_{m2} & \dots & c_{mp} \end{bmatrix}$$

COROLLARY. $tr(AB) = tr(BA)$, all matrices being square of order n .

If $A=[a_{ij}]$, $B=[b_{ij}]$ then $AB=C$ say $=[c_{ij}]$

where
$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

Let $BA=D=[d_{ij}]$ where $d_{ij} = \sum_{k=1}^n b_{ik} a_{kj}$

Thus $tr. (AB) = \sum_i c_{ii} = \sum_i \sum_k a_{ik} b_{ki}$

$$= \sum_{k=1}^n \sum_{i=1}^n a_{ik} b_{ki} \text{ (on interchanging the order of summation)}$$

$$= \sum_{k=1}^n \left(\sum_{i=1}^n b_{ki} a_{ik} \right)$$

$$= \sum_{k=1}^n d_{kk} = tr. (BA)$$

which proves the proposition.

2.6. PROPERTIES OF MATRIX-MULTIPLICATION

[1] The Commutative Law for Multiplication does not hold in general

Consider the matrices

$$A = \begin{bmatrix} 1 & -2 \\ 2 & 3 \\ -3 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 3 \end{bmatrix}$$

These are conformable for multiplication and so

$$AB = \begin{bmatrix} 1+0 & 0-2 & 2-6 \\ 2+0 & 0+3 & 4+9 \\ -3+0 & 0+1 & -6+3 \end{bmatrix} = \begin{bmatrix} 1 & -2 & -4 \\ 2 & 3 & 13 \\ -3 & 1 & -3 \end{bmatrix}$$

and $BA = \begin{bmatrix} 1+0-6 & -2+0+2 \\ 0+2-9 & 0+3+3 \end{bmatrix} = \begin{bmatrix} -5 & 0 \\ -7 & 6 \end{bmatrix}$

It is apparent that the product matrix AB is of order 3×3 while the product matrix BA is of order 2×2 and therefore the two product matrices are quite different i.e.,

$$AB \neq BA$$

This follows that the commutative law of multiplication does not hold.

Had the order of the matrices AB and BA been the same, then it would be possible that $AB \neq BA$ if every or at least one element in AB would differ from the corresponding element in BA . Though there are a few exceptions in which case the commutative law holds good. Such cases will be considered in the discussion of special matrices.

In fact for a given pair of matrices A and B it is possible that the products AB and BA may not be conformable. For example if A is of order $m \times n$ and B of order $n \times p$, then the product AB is conformable and will be of order $m \times p$ while the product BA is not conformable since number of columns in B is not equal to number of rows in A . Thus the product AB exists while BA does not.

[2] The Distributive Law for Multiplication holds Good

If A, B, C be three matrices of suitable orders such that the products $A(B+C)$ and AB, AC are conformable then

$$A(B+C) = AB + AC.$$

Suppose that

$$A = [a_{ij}] \text{ is of order } m \times n$$

$$B = [b_{jk}] \text{ is of order } n \times p$$

and

$$C = [c_{jk}] \text{ is also of order } n \times p$$

Then, $(B+C)$ is of order $n \times p$ and A is of order $m \times n$ so that $A(B+C)$ is conformable and of order $m \times p$. Also AB and AC both are of order $m \times p$ so that the sum matrix $(AB+AC)$ is of order $m \times p$. Hence the matrices $A(B+C)$ and $AB+AC$ are of the same order so that they are comparable for equality.

Now,

$$(i, k) \text{th element of } A(B+C) = \sum_{j=1}^n a_{ij} (b_{jk} + c_{jk})$$

$$= \sum_{j=1}^n a_{ij} b_{jk} + \sum_{j=1}^n a_{ij} c_{jk}$$

$$= (i, k) \text{th element of } AB + (i, k) \text{th element of } AC$$

$$= (i, k) \text{th element of } (AB+AC)$$

$$\text{for all } i=1, 2, \dots, m \text{ and } k=1, 2, \dots, p$$

$$\therefore A(B+C) = AB + AC$$

A similar procedure will show that

$$(B+C)A = BA + CA$$

where B, C, A are of orders $m \times n, m \times n, n \times p$ respectively.

Hence the matrix multiplication is distributive with respect to addition.

[3] The Associative Law for Multiplication holds Good

If A, B, C be three matrices of suitable orders such that the products $(AB)C$ and $A(BC)$ are conformable then

$$(AB)C = A(BC)$$

Suppose that

$$A = [a_{ij}] \text{ is of order } m \times n$$

$$B = [b_{jk}] \text{ is of order } n \times p$$

and

$$C = [c_{ki}] \text{ is of order } p \times q$$

Then, (AB) is of order $m \times p$ and C of order $p \times q$ so that $(AB)C$ is of order $m \times q$. Also (BC) is of order $m \times q$ and A of order $m \times n$ so that $A(BC)$ is of order $m \times q$. Hence the matrices $(AB)C$ and $A(BC)$ are of the same order so that they are comparable for equality

$$\text{Now } (i, k) \text{th element of } AB = \sum_{j=1}^n a_{ij} b_{jk}$$

$$\text{So that } (i, l) \text{th element of } (AB)C = \sum_{k=1}^p \left\{ \sum_{j=1}^n a_{ij} b_{jk} \right\} c_{kl}$$

$$= \sum_{k=1}^p \sum_{j=1}^n a_{ij} b_{jk} c_{kl}$$

$$\text{Also, } (j, l) \text{th element of } (BC) = \sum_{k=1}^p b_{jk} c_{kl}$$

$$\text{So that } (i, l) \text{th element of } A(BC) = \sum_{j=1}^n \left\{ \sum_{k=1}^p b_{jk} c_{kl} \right\} a_{ij}$$

$$= \sum_{j=1}^n \sum_{k=1}^p a_{ij} b_{jk} c_{kl}$$

$$= \sum_{k=1}^p \sum_{j=1}^n a_{ij} b_{jk} c_{kl}$$

$$= (i, l) \text{th element of } (AB)C$$

$$\text{for all } i=1, 2, \dots, m, l=1, 2, \dots, q$$

$$\therefore A(BC) = (AB)C.$$

Hence the matrix multiplication is associative.

[4] If A be a matrix of order $m \times n$ and O a null matrix of order $n \times p$ then the product AO is another null matrix of order $m \times p$ i.e.

$$AO_{n, p} = O_{m, p}$$

Also if O be an $m \times n$ null matrix and A a matrix of order $n \times p$ then their product is a null matrix of order $m \times p$ i.e.

$$O_{m, n} A = O_{m, p}$$

Conclusively if A be an n -rowed square matrix and O an n -rowed null matrix, then

$$AO = OA = O$$

If the product of two matrices A and B is a null matrix then it is essential that either of them is a null matrix i.e.

$AB = O$, it does not necessarily mean that at least one of A and B is a null-matrix. (Meerut, 1967; Gorakhpur, 1961)

An illustrative example if

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$AB = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} -1+1 & 1-1 \\ -2+2 & 2-2 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = O$$

The product of two non-zero matrices A and B is a zero matrix.

If I be an m -rowed unit matrix and A an $m \times n$ matrix then

$$I_m A = A$$

Suppose that $A = [a_{ij}]$ a matrix of order $m \times n$.

Then, $I_m A$ is a matrix of order $m \times n$ and so is comparable to A .

$$\text{Now, } (i, j) \text{th element of } I_m A = \sum_{k=1}^m (i, k) \text{th element of } I_m \cdot a_{kj}$$

In the (i, k) th element of I_m is zero except when $k=i$, therefore right hand sum of the above expression will have only one term different from zero and that is the i th term which is equal to

$$(i, i) \text{th element of } I_m \cdot a_{ij} = 1 \cdot a_{ij} \\ = a_{ij}$$

$$(i, j) \text{th element of } I_m A = a_{ij}$$

$$= (i, j) \text{th element of } A$$

It follows that

$$I_m A = A$$

Similarly it may be shown that if A be an $m \times n$ matrix and I_n an n -rowed unit matrix, then

$$A I_n = A$$

Note. This result shows the existence of a multiplicative identity.

(i) Positive Integral Powers of Square Matrices

If A is a square matrix of order n (say), then

$$A^2 = AA$$

By the associative law of multiplication leads

$$A^3 A = (AA) A = A(AA) = AA^2$$

or $A^3 = AAA = A^2A = AA^2$

In the generalized form if p, q are two positive integers, then

$$\begin{aligned} A^p A^q &= (AA \dots A, p \text{ times}) (AA \dots A, q \text{ times}) \\ &= AA \dots A, (p+q) \text{ times} \\ &= A^{p+q} \end{aligned}$$

and $(A^p)^q = (AA \dots A, p \text{ times})^q$
 $= A^q A^q \dots A^q, p \text{ times}$
 $= A^{pq}$

COROLLARY. If I is a unit matrix of any order, then

$$I^2 = I^3 = \dots = I^p = I.$$

Problem 3. Find the product of the matrices

$$A = \begin{bmatrix} 2 & 1 & 2 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 2 & -1 & 0 \\ 0 & 4 & 1 \\ -2 & 1 & 0 \\ 1 & -3 & 2 \end{bmatrix}$$

The matrix A is of the order 2×4 and B is of the order 4×3 that they are conformable for product and the product matrix will be of the order 2×3 .

Now,

$$\begin{aligned} AB &= \begin{bmatrix} 2 & 1 & 2 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & -1 & 0 \\ 0 & 4 & 1 \\ -2 & 1 & 0 \\ 1 & -3 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 2 \cdot 2 + 1 \cdot 0 + 2 \cdot (-2) + 1 \cdot 1 & 2 \cdot (-1) + 1 \cdot 4 + 2 \cdot 1 + 1 \cdot (-3) \\ 1 \cdot 2 + 1 \cdot 0 + 1 \cdot (-2) + 1 \cdot 1 & 1 \cdot (-1) + 1 \cdot 4 + 1 \cdot 1 + 1 \cdot (-3) \end{bmatrix} \\ &= \begin{bmatrix} 1 & 1 & 3 \\ 1 & 1 & 3 \end{bmatrix} \end{aligned}$$

Problem 4. If $A = \begin{bmatrix} 1 & -2 & 3 \\ -4 & 2 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 3 \\ 4 & 5 \\ 2 & 1 \end{bmatrix}$

find AB and show that $AB \neq BA$.

Since A is of the order 2×3 and B of the order 3×2 , therefore AB and BA are conformable for the products AB and BA both. AB will be of the order 2×2 while BA will be of the order 3×3 .

$$\begin{aligned} \text{Now, } AB &= \begin{bmatrix} 1 & -2 & 3 \\ -4 & 2 & 5 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 4 & 5 \\ 2 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 \cdot 2 - 2 \cdot 4 + 3 \cdot 2 & 1 \cdot 3 - 2 \cdot 5 + 3 \cdot 1 \\ -4 \cdot 2 + 2 \cdot 4 + 5 \cdot 2 & -4 \cdot 3 + 2 \cdot 5 + 5 \cdot 1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & -4 \\ 10 & 3 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} \text{and } BA &= \begin{bmatrix} 2 & 3 \\ 4 & 5 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 3 \\ -4 & 2 & 5 \end{bmatrix} \\ &= \begin{bmatrix} 2 \cdot 1 - 3 \cdot 4 & -2 \cdot 2 + 3 \cdot 2 & 2 \cdot 3 + 3 \cdot 5 \\ 4 \cdot 1 - 5 \cdot 4 & -4 \cdot 2 + 5 \cdot 2 & 4 \cdot 3 + 5 \cdot 5 \\ 2 \cdot 1 - 1 \cdot 4 & -2 \cdot 2 + 1 \cdot 2 & 2 \cdot 3 + 1 \cdot 5 \end{bmatrix} \\ &= \begin{bmatrix} -10 & 2 & 21 \\ -16 & 2 & 37 \\ -2 & -2 & 11 \end{bmatrix} \end{aligned}$$

It is clear that AB and BA being of the different order cannot be equal.

$$AB \neq BA.$$

Problem 5. If $A_\alpha = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$ and $A_\beta = \begin{bmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{bmatrix}$

show that

$$A_{\alpha+\beta} = \begin{bmatrix} \cos(\alpha+\beta) & \sin(\alpha+\beta) \\ -\sin(\alpha+\beta) & \cos(\alpha+\beta) \end{bmatrix}$$

and prove that

$$A_\alpha A_\beta = A_\beta A_\alpha = A_{\alpha+\beta}$$

We have

$$\begin{aligned} A_\alpha A_\beta &= \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{bmatrix} \\ &= \begin{bmatrix} \cos \alpha \cos \beta - \sin \alpha \sin \beta & \cos \alpha \sin \beta + \sin \alpha \cos \beta \\ -\sin \alpha \cos \beta - \cos \alpha \sin \beta & -\sin \alpha \sin \beta + \cos \alpha \cos \beta \end{bmatrix} \\ &= \begin{bmatrix} \cos(\alpha+\beta) & \sin(\alpha+\beta) \\ -\sin(\alpha+\beta) & \cos(\alpha+\beta) \end{bmatrix} \\ &= A_{\alpha+\beta}. \end{aligned}$$

Similarly $A_\beta A_\alpha = A_{\alpha+\beta}$

Hence $A_\alpha A_\beta = A_\beta A_\alpha = A_{\alpha+\beta}$.

Problem 6. If $A = \begin{bmatrix} 3 & -4 \\ 1 & 1 \end{bmatrix}$ prove that $A^n = \begin{bmatrix} 1+2n & -4n \\ n & 1-2n \end{bmatrix}$

where n is any positive integer.

We have

$$\begin{aligned} A^2 &= AA = \begin{bmatrix} 3 & -4 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 & -4 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 5 & -8 \\ 2 & -3 \end{bmatrix} = \begin{bmatrix} 1+2 \cdot 2 & -4 \cdot 2 \\ 2 & 1-2 \cdot 2 \end{bmatrix} \\ A^3 &= A^2 A = \begin{bmatrix} 5 & -8 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} 3 & -4 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 7 & -12 \\ 3 & -5 \end{bmatrix} \\ &= \begin{bmatrix} 1+2 \cdot 3 & -4 \cdot 3 \\ 3 & 1-2 \cdot 3 \end{bmatrix} \\ A^4 &= A^3 A = \begin{bmatrix} 7 & -12 \\ 3 & -5 \end{bmatrix} \begin{bmatrix} 3 & -4 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 9 & -16 \\ 4 & -7 \end{bmatrix} = \begin{bmatrix} 1+2 \cdot 4 & -4 \cdot 4 \\ 4 & 1-2 \cdot 4 \end{bmatrix} \end{aligned}$$

Assuming that the result is true for an integral power n , we have

$$A^n = \begin{bmatrix} 1+2n & -4n \\ n & 1-2n \end{bmatrix}$$

Multiplying both sides by A ,

$$A^{n+1} = \begin{bmatrix} 1+2n & -4n \\ n & 1-2n \end{bmatrix} \begin{bmatrix} 3 & -4 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1+2(n+1) & -4(n+1) \\ n+1 & 1-2(n+1) \end{bmatrix}$$

Thus if the law is true for A^n , it is also true for A^{n+1} . But it is true for $n=2, 3, 4, \dots$; hence by the method of induction the same law is true for any positive integral value n .

2.7. PARTITIONING OF MATRICES

In many cases it is found rather convenient to regard a matrix as made up of a certain number of its submatrices which may be treated as the elements of the matrix. This can be done by drawing any number of lines parallel to the rows and columns of the given matrix. The submatrices contained in rectangles so formed are then treated as the elements of the matrices. This process of dividing a given matrix into certain number of sub-matrix elements is known as the *Partitioning of matrix*.

As an illustrative example if A is a matrix defined by

$$A = \begin{bmatrix} 1 & 2 & 4 \\ -2 & 0 & -5 \\ 0 & 3 & 6 \end{bmatrix}$$

then this matrix can be partitioned in different forms by drawing lines parallel to row or column or both.

In the first instance, we may partition it as

$$A = \begin{bmatrix} 1 & 2 & 4 \\ -2 & 0 & -5 \\ \dots & \dots & \dots \\ 0 & 3 & 6 \end{bmatrix} = \begin{bmatrix} A_{11} \\ A_{21} \end{bmatrix} \text{ where } A_{11} = \begin{bmatrix} 1 & 2 & 4 \\ -2 & 0 & -5 \end{bmatrix} \text{ and } A_{21} = \begin{bmatrix} 0 & 3 & 6 \end{bmatrix}$$

In the second instance we may have

$$A = \begin{bmatrix} 1 & \vdots & 2 & 4 \\ -2 & \vdots & 0 & -5 \\ 0 & \vdots & 3 & 6 \end{bmatrix} = [A_{11} \ A_{12}] \text{ where } A_{11} = \begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix} \text{ and } A_{12} = \begin{bmatrix} 2 & 4 \\ 0 & -5 \\ 3 & 6 \end{bmatrix}$$

In the third instance we may express

$$A = \begin{bmatrix} 1 & 2 & \vdots & 4 \\ -2 & 0 & \vdots & -5 \\ \dots & \dots & \dots & \dots \\ 0 & 3 & \vdots & 6 \end{bmatrix}$$

$$= \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \text{ where } A_{11} = \begin{bmatrix} 1 & 2 \\ -2 & 0 \end{bmatrix}, A_{12} = \begin{bmatrix} 4 \\ -5 \end{bmatrix},$$

$$A_{21} = \begin{bmatrix} 0 & 3 \end{bmatrix}, A_{22} = \begin{bmatrix} 6 \end{bmatrix}$$

As another example if we have two matrices of the same order say

$$A = \begin{bmatrix} 0 & 2 & -5 \\ 1 & 3 & 4 \\ -1 & 5 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} -2 & 0 & 3 \\ 5 & 4 & 2 \\ 3 & -1 & 6 \end{bmatrix}$$

partitioned identically as below

$$A = \begin{bmatrix} 0 & 2 & -5 \\ 1 & 3 & 4 \\ \dots & \dots & \dots \\ -1 & 5 & 2 \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

$$\text{and } B = \begin{bmatrix} -2 & 0 & 3 \\ 5 & 4 & 2 \\ \dots & \dots & \dots \\ 3 & -1 & 6 \end{bmatrix} = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

then the two matrices A and B are said to be **identically partitioned**.

The two identically partitioned matrices A and B are comparable for addition if the corresponding sub-matrices in A and B are of the same order i.e. the order of A_{11} , B_{11} is the same, the order of A_{12} , B_{12} is the same, the order of A_{21} , B_{21} is the same and the order of A_{22} , B_{22} is the same in the above example and then we have

$$A+B = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} + \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} A_{11}+B_{11} & A_{12}+B_{12} \\ A_{21}+B_{21} & A_{22}+B_{22} \end{bmatrix}$$

2.8. PRODUCT OF MATRICES BY PARTITIONING

Let A be a matrix of order $m \times n$ and B be a matrix of order $n \times p$ so that the matrices A and B are conformable for multiplication. Now if the two matrices are partitioned, then the partitioned matrices will be conformable for multiplication 'when to each partition line of A parallel to the columns there corresponds a partition line of B parallel to its rows such that the number of columns of A lying between two adjacent partition lines is the same as the number of rows of B lying between corresponding adjacent partition lines.' Broadly speaking if in A there lie a partition line after third column, another line after fifth column, then in B a corresponding line should be after third row and the another line after fifth row.

It is notable that partition lines in A parallel to its columns may be drawn in any arbitrary manner. Two matrices A and B partitioned in the above described manner are said to be *conformably partitioned* for multiplication, for, with such partitions the two matrices can be multiplied as usual, with sub-matrices as the elements.

As an illustrative example if A be a matrix of order 4×5 and B a matrix of order 5×6 partitioned as below:

$$A = \begin{bmatrix} 2 & 1 & 3 & -1 & 2 \\ 3 & 2 & 2 & 5 & 0 \\ \hline 4 & 5 & 0 & 5 & 3 \\ 0 & 2 & 6 & 7 & 4 \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix} \text{ of order } 2 \times 5$$

where A_{11} is of order 2×2

A_{12} " 2×2

A_{13} " 2×1

A_{21} " 2×2

A_{22} " 2×2

A_{23} " 2×1

$$\text{and } B = \begin{bmatrix} 0 & 2 & 3 & 4 & -2 & 3 \\ 5 & 4 & -2 & 5 & 6 & 2 \\ \hline 2 & 0 & 3 & 1 & 4 & 5 \\ -2 & 3 & 4 & 5 & -6 & 0 \\ \hline 0 & 4 & -1 & 2 & 7 & 8 \end{bmatrix} = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \\ B_{31} & B_{32} \end{bmatrix} \text{ of order } 3 \times 6$$

where B_{11} is of order 2×3

B_{12} " 2×3

B_{21} " 2×3

B_{22} " 2×3

B_{31} " 1×3

B_{32} " 1×3

Then it is clear that A and B are conformable for multiplication. Also the partitioned matrices A and B are conformable for multiplication, for, the partitioned matrix A is of order 2×3 and the partitioned matrix B is of order 3×2 .

Thus,

$$\begin{aligned} AB &= \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \\ B_{31} & B_{32} \end{bmatrix} \\ &= \begin{bmatrix} A_{11} B_{11} + A_{12} B_{21} + A_{13} B_{31} & A_{11} B_{12} + A_{12} B_{22} + A_{13} B_{32} \\ A_{21} B_{11} + A_{22} B_{21} + A_{23} B_{31} & A_{21} B_{12} + A_{22} B_{22} + A_{23} B_{32} \end{bmatrix} \\ &= \begin{bmatrix} [2 \ 1] [0 \ 2 \ 3] + [3 \ -1] [2 \ 0 \ 3] + [2] [0 \ 4 \ -1] \\ [3 \ 2] [5 \ 4 \ -2] + [2 \ 5] [-2 \ 3 \ 4] + [0] [2 \ 7 \ 8] \\ [2 \ 1] [4 \ -2 \ 3] + [3 \ -1] [1 \ 4 \ 5] + [2] [2 \ 7 \ 8] \\ [3 \ 2] [5 \ 6 \ 2] + [2 \ 5] [5 \ -6 \ 0] + [0] [0 \ 4 \ -1] \\ [4 \ 5] [0 \ 2 \ 3] + [0 \ 5] [2 \ 0 \ 3] + [3] [0 \ 4 \ -1] \\ [0 \ 2] [5 \ 4 \ -2] + [6 \ 7] [-2 \ 3 \ 4] + [4] [2 \ 7 \ 8] \\ [4 \ 5] [4 \ -2 \ 3] + [0 \ 5] [1 \ 4 \ 5] + [3] [2 \ 7 \ 8] \\ [0 \ 2] [5 \ 6 \ 2] + [6 \ 7] [5 \ -6 \ 0] + [4] [0 \ 4 \ -1] \end{bmatrix} \\ &= \begin{bmatrix} 5 & 8 & 4 \\ 10 & 14 & 5 \end{bmatrix} + \begin{bmatrix} 8 & -3 & 5 \\ -6 & 15 & 26 \end{bmatrix} + \begin{bmatrix} 0 & 8 & -2 \\ 0 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 13 & 2 & 8 \\ 22 & 6 & 13 \end{bmatrix} + \begin{bmatrix} -2 & 18 & 15 \\ 27 & -22 & 10 \end{bmatrix} + \begin{bmatrix} 4 & 14 & 16 \\ 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

$$\begin{bmatrix} 25 & 28 & 2 \\ 10 & 8 & -4 \end{bmatrix} + \begin{bmatrix} -10 & 15 & 20 \\ -2 & 21 & 46 \end{bmatrix} + \begin{bmatrix} 0 & 12 & -3 \\ 0 & 16 & -4 \end{bmatrix} \\ + \begin{bmatrix} 41 & 22 & 22 \\ 10 & 12 & 4 \end{bmatrix} + \begin{bmatrix} 25 & -30 & 0 \\ 41 & -18 & 30 \end{bmatrix} + \begin{bmatrix} 6 & 21 & 24 \\ 8 & 28 & 32 \end{bmatrix}$$

$$\begin{bmatrix} 13 & 13 & 7 \\ 4 & 29 & 31 \end{bmatrix} \begin{bmatrix} 15 & 34 & 39 \\ 49 & -16 & 23 \\ 72 & 13 & 46 \end{bmatrix}$$

$$\begin{bmatrix} 13 & 13 & 7 & 15 & 34 & 39 \\ 4 & 29 & 31 & 49 & -16 & 23 \\ 15 & 55 & 19 & 72 & 13 & 46 \\ 8 & 45 & 38 & 59 & 22 & 66 \end{bmatrix}$$

SPECIAL MATRICES WITH THEIR PROPERTIES

When defining a matrix, we have already mentioned a few of the types of matrices like square matrix, null matrix, row and column matrices, unit matrix etc., but in this section we have to discuss almost all the special types of matrices with their properties in comparison to others. Some types will be discussed in more detailed manner, a few others which are rarely used in different branches of pure mathematics applied mathematics and mechanics.

Square Matrix and Special Square Matrices

A matrix A having the number of rows and columns equal is called a square matrix *e.g.*

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

A matrix A is said to be a square matrix of order n or an n -rowed square matrix or simply n -square matrix.

A matrix which is not a square matrix may be called as a rectangular matrix, *e.g.*

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{33} \end{bmatrix}$$

This is a matrix of order 2×3 .

Determinant of a square matrix. The determinant of a square matrix A is the determinant which has got for its elements all the elements of the matrix A in the same places. It is denoted by $|A|$. If $A = [a_{ij}]$ then $|A| = |a_{ij}|$.

Another example if A be a n -square matrix given by

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

$$\text{then } |A| = \begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix}$$

It is easy to show that the determinant of the product matrix is equal to the product of the determinants of the square matrices **A** and **B** of the same order i.e.,

$$|AB| = |A| |B|$$

Or, in general the determinant of the product of any number of square matrices of the same order is the product of the determinants of those matrices i.e.,

$$|ABC\dots K| = |A| |B| |C| \dots |K|$$

Special square matrices. While discussing the properties of multiplication we have mentioned the commutative property relation to square matrices **A** and **B** of the same order say n , we say that if $AB=BA$ then **A** and **B** commute and if $AB=-BA$ then **A** and **B** anti-commute. It can be easily verified that any square matrix **A** commutes with itself and also with identity matrix I_n .

$$AA=AA$$

and

$$AI_n = I_n A$$

If k is a positive integer, then the matrix **A** with the property $A^{k+1}=A$ is said to be *periodic*. In case k is the least positive integer such that $A^{k+1}=A$, then **A** is said to be of *Period k*.

In the case when $k=1$, so that $A^2=A$, then **A** is said to be *idempotent* and if p is some positive integer such that $A^p=0$ then **A** is said to be *nilpotent*. In case p is the least positive integer such that $A^p=0$, **A** is said to be nilpotent of *index p*.

ILLUSTRATIVE EXAMPLES

1. $\begin{bmatrix} 5 & 0 & 4 \\ 2 & 3 & 9 \\ -1 & 5 & 0 \end{bmatrix}$ is a square matrix of order 3

2. $\begin{bmatrix} 2 & 3 & 4 \\ 5 & -2 & 0 \end{bmatrix}$ is a rectangular matrix of order 2×3

3. The determinant of the 3-square matrix $\begin{bmatrix} 2 & 0 & 4 \\ 5 & -2 & 1 \\ 0 & 3 & 2 \end{bmatrix}$ is 0

4. The matrices $\begin{bmatrix} \alpha & \beta \\ \beta & \alpha \end{bmatrix}$ and $\begin{bmatrix} \gamma & \delta \\ \delta & \gamma \end{bmatrix}$ commute for all values of $\alpha, \beta, \gamma, \delta$

since

$$\begin{bmatrix} \alpha & \beta \\ \beta & \alpha \end{bmatrix} \begin{bmatrix} \gamma & \delta \\ \delta & \gamma \end{bmatrix} = \begin{bmatrix} \alpha\gamma + \beta\delta & \alpha\delta + \beta\gamma \\ \beta\gamma + \alpha\delta & \beta\delta + \alpha\gamma \end{bmatrix} = \begin{bmatrix} \gamma & \delta \\ \delta & \gamma \end{bmatrix} \begin{bmatrix} \alpha & \beta \\ \beta & \alpha \end{bmatrix}$$

5. The matrix $\begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$ is idempotent, since

$$\begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}^2 = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix} \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix} = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$$

6. The matrix $\begin{bmatrix} 1 & -2 & -6 \\ -3 & 2 & 9 \\ 2 & 0 & -3 \end{bmatrix}$ is periodic of period 2, since

$$\begin{bmatrix} 1 & -2 & -6 \\ -3 & 2 & 9 \\ 2 & 0 & -3 \end{bmatrix}^{2+1} = \begin{bmatrix} 1 & -2 & -6 \\ -3 & 2 & 9 \\ 2 & 0 & -3 \end{bmatrix}^2 \begin{bmatrix} 1 & -2 & -6 \\ -3 & 2 & 9 \\ 2 & 0 & -3 \end{bmatrix} = \begin{bmatrix} -5 & -6 & -6 \\ 9 & 10 & 9 \\ -4 & -4 & -3 \end{bmatrix} \begin{bmatrix} 1 & -2 & -6 \\ -3 & 2 & 9 \\ 2 & 0 & -3 \end{bmatrix} = \begin{bmatrix} 1 & -2 & -6 \\ -3 & 2 & 9 \\ 2 & 0 & -3 \end{bmatrix}$$

7. The matrix $\begin{bmatrix} 1 & 1 & 3 \\ 5 & 2 & 6 \\ -2 & -1 & -3 \end{bmatrix}$ is nilpotent of index 3, since

$$\begin{bmatrix} 1 & 1 & 3 \\ 5 & 2 & 6 \\ -2 & -1 & -3 \end{bmatrix}^3 = \begin{bmatrix} 1 & 1 & 3 \\ 5 & 2 & 6 \\ -2 & -1 & -3 \end{bmatrix}^2 \begin{bmatrix} 1 & 1 & 3 \\ 5 & 2 & 6 \\ -2 & -1 & -3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 3 & 3 & 9 \\ -1 & -1 & -3 \end{bmatrix} \begin{bmatrix} 1 & 1 & 3 \\ 5 & 2 & 6 \\ -2 & -1 & -3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \mathbf{O}$$

(2) Row Matrix and Column Matrix (Row and Column Vectors)

A matrix having only one row and any number (>1) of columns is said to be a *row matrix* or a *row vector* e.g., a row matrix of order $1 \times n$ is

$$[a_{11} \ a_{12} \ \dots \ a_{1n}] \text{ or } [a_1 \ a_2 \ \dots \ a_n]$$

A matrix having only one column and any number (>1) of rows is said to be a *column matrix* or a *column vector* e.g., a column matrix of order $m \times 1$ is

$$\begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} \text{ or } \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{bmatrix}$$

In expressing vectors as matrices, the elements of the row vector or column vector are known as the components of the vectors. Thus

we can define a n -vector as 'an ordered n -tuple of n real or complex numbers written in a horizontal or in a vertical line.'

Algebraic operations on vectors. If A and B be two vectors with components $a_1, a_2, \dots, a_n$ and $b_1, b_2, \dots, b_n$ respectively, then it is easy to show that

$$(i) \quad A+B=[a_1, a_2, \dots, a_n]+[b_1, b_2, \dots, b_n] \\ = [a_1+b_1, a_2+b_2, \dots, a_n+b_n]$$

$$(ii) \quad kA=k[a_1, a_2, \dots, a_n], \quad k \text{ being a scalar} \\ = [ka_1, ka_2, \dots, ka_n]$$

$$(iii) \quad A \cdot B = a_1 b_1 + a_2 b_2 + \dots + a_n b_n$$

$$(iv) \quad A \cdot A = |A|^2 = a_1^2 + a_2^2 + \dots + a_n^2 \\ = |a_1|^2 + |a_2|^2 + \dots + |a_n|^2$$

$$(v) \quad A \text{ vector will be a unit vector if } |A|=1$$

$$(vi) \quad \text{The vectors } A \text{ and } B \text{ will be orthogonal if } A \cdot B = 0$$

$$(vii) \quad \text{The vector } O[0, 0, \dots, 0] \text{ is said to be a null vector}$$

$$(viii) \quad A+O=O+A=A$$

Linearly dependent and Independent sets of vectors. A set of n vectors $A_1, A_2, \dots, A_n$ is said to be *linearly dependent* if there exists a set of n (of which at least one is non-zero) scalars $k_1, k_2, \dots, k_n$ such that

$$k_1 A_1 + k_2 A_2 + \dots + k_n A_n = O$$

In the case when $k_1=k_2=\dots=k_n=0$, the set of n vectors $A_1, A_2, \dots, A_n$ is said to be *linearly independent* if

$$k_1 A_1 + k_2 A_2 + \dots + k_n A_n = O.$$

A vector A is known as a *linear combination* of the set of vectors $A_1, A_2, \dots, A_n$ if it is expressible as

$$A = k_1 A_1 + k_2 A_2 + \dots + k_n A_n$$

where $k_1, k_2, \dots, k_n$ are scalars.

III. ILLUSTRATIVE EXAMPLES

1. $[1 \ 0 \ 2 \ 4]$ is a 1×4 row matrix (or row vector).

2. $\begin{bmatrix} 2 \\ 3 \\ 5 \end{bmatrix}$ is a 3×1 column matrix (or column vector).

3. If $A=[4, 6, 0, 3]$ and $B=[2, 3, 4, 0]$ are two vectors, then

$$A+B=[4, 6, 0, 3]+[2, 3, 4, 0]=[6, 9, 4, 3]$$

$$3A-2B=3[4, 6, 0, 3]-2[2, 3, 4, 0]$$

$$=[12, 18, 0, 9]-[4, 6, 8, 0]=[8, 12, -8, 9] \text{ etc.}$$

4. The set of vectors $[1, 2, 3], [2, -2, 0]$ is linearly independent since $k_1[1, 2, 3]+k_2[2, -2, 0]=0$ is equivalent to a system of equations

$$k_1+2k_2=0, \quad 2k_1-2k_2=0, \quad 3k_1=0$$

which are satisfied only if $k_1=k_2=0$.

MATRICES

5. The set of vectors $[2, 4, 10], [3, 6, 15]$ is linearly dependent, since

$$k_1[2, 4, 10]+k_2[3, 6, 15]=0 \text{ is equivalent to a system of equations}$$

$$2k_1+3k_2=0, \quad 4k_1+6k_2=0, \quad 10k_1+15k_2=0$$

which are satisfied when $k_1=3, k_2=-2$ i.e., non-zero values of k_1 and k_2 .

[3] Null Matrix

If all the elements of a matrix are zero then it is said to be a *null matrix* and denoted by O e.g.,

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ is a null matrix of order } 3 \times 4.$$

The commutative law of multiplication holds good in case of a null matrix e.g., if

$$A = \begin{bmatrix} 2 & -1 \\ 3 & 1 \end{bmatrix} \text{ and } O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\text{then } AO = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = OA = O$$

We have also proved that a null matrix acts as an additive identity of any matrix i.e., as in the above example

$$A+O = \begin{bmatrix} 2 & -1 \\ 3 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & -1 \\ 3 & 1 \end{bmatrix} = A$$

[4] Unit Matrix or Identity Matrix

A square matrix of order n having unit elements in the Principal or leading diagonal and zero elements everywhere else, is called a *unit matrix* or *identity matrix* and denoted by I_n . Thus

$$I_n = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix} \text{ and for } n=4, I_4 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \text{ etc.}$$

It is clear that a square matrix $A=[a_{ij}]$ is the unit matrix if

$$a_{ij}=1 \text{ for } i=j \\ =0 \text{ for } i \neq j$$

It is also evident that

$$I_n = I_n^2 = I_n^3 = \dots$$

The commutative law of multiplication also holds good in case of a unit matrix e.g., if

$$A = \begin{bmatrix} -1 & 2 & 0 \\ 3 & 4 & 5 \\ 2 & 0 & -3 \end{bmatrix} \text{ and } I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{then } AI = IA = A = \begin{bmatrix} -1 & 2 & 0 \\ 3 & 4 & 5 \\ 2 & 0 & -3 \end{bmatrix}$$

[5] Upper and Lower Triangular Matrices, Diagonal and Scalar Matrices

A square matrix A of any order n , defined by

$$A = [a_{ij}]$$

is said to be an *upper triangular matrix* if its elements $a_{ij} = 0$ for $i > j$ where i, j are positive integers ranging over from 1 to n . Thus an upper triangular matrix of order n is

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22} & a_{23} & \dots & a_{2n} \\ 0 & 0 & a_{33} & \dots & a_{3n} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & a_{nn} \end{bmatrix}$$

The square matrix $A = [a_{ij}]$ of order n is said to be a *lower triangular matrix* if its elements $a_{ij} = 0$ for $i < j$ where i, j are integers such that $1 \leq i \leq n, 1 \leq j \leq n$. Thus a lower triangular matrix of order n is

$$\begin{bmatrix} a_{11} & 0 & 0 & \dots & 0 \\ a_{21} & a_{22} & 0 & \dots & 0 \\ a_{31} & a_{32} & a_{33} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{bmatrix}$$

A square matrix of order n which is both upper and lower triangular is called a *diagonal matrix* e.g.,

$$\begin{bmatrix} a_{11} & 0 & 0 & \dots & 0 \\ 0 & a_{22} & 0 & \dots & 0 \\ 0 & 0 & a_{33} & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & a_{nn} \end{bmatrix}$$

This is denoted by $\text{diag}[a_{11}, a_{22}, a_{33}, \dots, a_{nn}]$ and the sum $a_{11} + a_{22} + \dots + a_{nn}$ is called its *trace* i.e., $\text{tr } A = \sum_i a_{ii}$

In a diagonal matrix if all the diagonal elements are equal to a scalar quantity say λ i.e., $a_{11} = a_{22} = a_{33} = \dots = a_{nn} = \lambda$, then the matrix is called a *scalar matrix* e.g.,

$$\begin{bmatrix} \lambda & 0 & 0 & \dots & 0 \\ 0 & \lambda & 0 & \dots & 0 \\ 0 & 0 & \lambda & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \lambda \end{bmatrix}$$

It bears the name scalar matrix due to the reason that if a square matrix A of order n be multiplied by the scalar matrix of order n , then it is equivalent to multiplying the matrix A by a scalar.

In case $\lambda = 1$, the scalar matrix reduces to a *unit matrix*.

In other words an n -rowed square matrix $[a_{ij}]$ is called a *scalar matrix* when

$$\begin{aligned} a_{ij} &= \lambda \text{ (some scalar) for } i=j, \\ &= 0 \text{ for } i \neq j. \end{aligned}$$

ILLUSTRATIVE EXAMPLES

1. $\begin{bmatrix} 1 & 2 & 3 \\ 0 & -1 & 5 \\ 0 & 0 & -2 \end{bmatrix}$ is a 3-rowed upper triangular matrix.

2. $\begin{bmatrix} 1 & 0 & 0 \\ -2 & 5 & 0 \\ 3 & 4 & -1 \end{bmatrix}$ is a 3-rowed lower triangular matrix.

3. $\begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ is a 3-rowed diagonal matrix. Its trace is $2+1+3$ i.e., 6

4. $\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$ is a 3-rowed scalar matrix. That's why? Because if

we multiply it by a 3-rowed square matrix $A = \begin{bmatrix} 1 & -1 & 2 \\ 3 & 4 & 5 \\ -2 & 5 & 4 \end{bmatrix}$

then

$$\begin{bmatrix} 1 & -1 & 2 \\ 3 & 4 & 5 \\ -2 & 5 & 4 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 2 & -2 & 4 \\ 6 & 8 & 10 \\ -4 & 10 & 8 \end{bmatrix} = 2 \begin{bmatrix} 1 & -1 & 2 \\ 3 & 4 & 5 \\ -2 & 5 & 4 \end{bmatrix} = 2A$$

i.e., A has been multiplied by the scalar 2.

Problem 7. Show that if $AB=A$ and $BA=B$ then A and B are idempotent.

We have $ABA = (AB)A$

$$= AA \text{ as } AB=A$$

$$= A^2$$

... (1)

Also $ABA = A(BA)$

$$= AB \text{ as } BA=B$$

$$= A \text{ as } AB=A$$

... (2)

From (1) and (2) it follows that

$$A^2 = A = ABA$$

Hence A is idempotent.

Again consider

$$BAB = B(AB)$$

$$= BA \text{ as } AB=A$$

$$= B \text{ as } BA=B$$

and also

$$BAB = (BA)B$$

$$= BB \text{ as } BA=B$$

$$= B^2$$

It is clear that $BAB = B^2 = B$

Hence B is also idempotent.

Problem 8. If A and B are n -square matrices then show that A and B commute if and only if $A - \lambda I$ and $B - \lambda I$ commute for every scalar λ .

If $A - \lambda I$ and $B - \lambda I$ commute, then

$$(A - \lambda I)(B - \lambda I) = (B - \lambda I)(A - \lambda I)$$

$$\text{i.e. } AB - \lambda(A + B) + \lambda^2 I = BA - \lambda(B + A) + \lambda^2 I \quad \therefore AI = A \text{ etc.}$$

The comparison gives

$$AB = BA \text{ since } A + B = B + A$$

i.e. A and B commute.

Conversely if A and B commute then $AB = BA$.

$$\begin{aligned} \text{Consider } (A - \lambda I)(B - \lambda I) &= AB - \lambda(A + B) + \lambda^2 I \\ &= BA - \lambda(B + A) + \lambda^2 I \\ &= (B - \lambda I)(A - \lambda I) \end{aligned}$$

which follows that $A - \lambda I$ and $B - \lambda I$ commute.

Problem 9. Derive a rule for forming the product BA of an $m \times n$ matrix B and $A = \text{diag}(a_{11}, a_{22}, \dots, a_{nn})$

$$\text{Given } A = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix}$$

and suppose that

$$B = \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1n} \\ b_{21} & b_{22} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots \\ b_{m1} & b_{m2} & \dots & b_{mn} \end{bmatrix}$$

then consider the product

$$\begin{aligned} BA &= \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1n} \\ b_{21} & b_{22} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots \\ b_{m1} & b_{m2} & \dots & b_{mn} \end{bmatrix} \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix} \\ &= \begin{bmatrix} b_{11}a_{11} & b_{12}a_{22} & \dots & b_{1n}a_{nn} \\ b_{21}a_{11} & b_{22}a_{22} & \dots & b_{2n}a_{nn} \\ \dots & \dots & \dots & \dots \\ b_{m1}a_{11} & b_{m2}a_{22} & \dots & b_{mn}a_{nn} \end{bmatrix} \end{aligned}$$

which follows that the product BA of an $m \times n$ matrix B by a n -square diagonal matrix A is obtained by multiplying the first column of B by a_{11} , the second column of B by a_{22} and so on. This gives the required rule.

[6] The Transpose of a Matrix

The matrix of order $n \times m$ obtained from any matrix A of order $m \times n$, by interchanging its rows and columns is called the *transpose* of A and is denoted by A' or A^T . Thus if $A = [a_{ij}]$,

then A' or $A^T = [a'_{ji}]$ where $a'_{ji} = a_{ij}$ *i.e.*, the (j, i) th element of A is the (i, j) th element of A , *e.g.*, if

$$A = \begin{bmatrix} 5 & 4 & -2 \\ 1 & 0 & 2 \\ 3 & 4 & -5 \\ 6 & -4 & 2 \end{bmatrix}, \text{ then } A' = \begin{bmatrix} 5 & 1 & 3 & 6 \\ 4 & 0 & 4 & -4 \\ -2 & 2 & -5 & 2 \end{bmatrix}$$

Properties of the transpose of a matrix

i. The properties of the transpose of a matrix coincides with itself, *i.e.*, if A be a matrix then $(A')' = A$.

Let $A = [a_{ij}]$ be the matrix of order $m \times n$.

Then $A' = [a'_{ji}]$ is the matrix of order $n \times m$.

So that $(A')'$ will be the matrix of order $m \times n$.

As such A and $(A')'$ are the matrices of the same order.

Again, (i, j) th element of $A = (j, i)$ th element of A'
 $= (i, j)$ th element of $(A')'$

Hence $(A')' = A$.

ii. The determinant of the transpose of a square matrix is the same as the determinant of the matrix.

Let $A = [a_{ij}]$ be the square matrix of order n .

Then the transpose of A *i.e.*, A' will be the square matrix of order n .

$$\begin{aligned} \text{Now, } |A| &= |a_{ij}| = \begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix} \\ \text{and } |A'| &= |a'_{ji}| = \begin{vmatrix} a_{11} & a_{21} & \dots & a_{n1} \\ a_{12} & a_{22} & \dots & a_{n2} \\ \vdots & \vdots & \dots & \vdots \\ a_{1n} & a_{2n} & \dots & a_{nn} \end{vmatrix} \end{aligned}$$

$$= \begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix} \quad \begin{array}{l} \text{since the interchange} \\ \text{of rows and columns in} \\ \text{a determinant does not} \\ \text{change the value of the} \\ \text{determinant.} \end{array}$$

$$= |A|.$$

Hence the determinant of the transpose of a square matrix is equal to the determinant of the matrix.

iii. If k be any scalar and A a matrix, then

$$(kA)' = kA'$$

Let $A = [a_{ij}]$ be the matrix of order $m \times n$. Then A' and so kA' is a matrix of order $n \times m$. Also (kA) is the matrix of order $m \times n$ and so $(kA)'$ is the matrix of order $n \times m$. Thus the matrices kA' and $(kA)'$ are of the same order.

Now,

$$\begin{aligned} (i, j) \text{th element of } (kA)' &= (j, i) \text{th element of } kA \\ &= k \text{ times the } (j, i) \text{th element of } A \\ &= k a_{ji} \end{aligned}$$

$=k$ times the (i, j) th element of A'
 $= (i, j)$ th element of kA'

Hence $(kA)' = kA'$.

IV. The transpose of the sum of two matrices A and B (conformable for addition) is the sum of their transposes i.e.,

$$(A+B)' = A' + B'$$

Let $A = [a_{ij}]$ be the matrix of order $m \times n$
 and $B = [b_{ij}]$ „ „ „ $m \times n$.

Then $(A+B)$ will be the matrix of order $m \times n$
 and $(A+B)'$ „ „ „ $n \times m$.

Also A' is the matrix of order $n \times m$
 and B' „ „ „ $n \times m$
 so that $(A' + B')$ is the matrix of order $n \times m$.

Thus, $(A+B)'$ and $(A' + B')$ will be the matrices of the same order $n \times m$.

Again (j, i) th element of $(A+B)' = (i, j)$ th element of $A+B$
 $= a_{ij} + b_{ij}$

and (j, i) th element of $(A' + B') = (j, i)$ th element of $A' + (j, i)$ th element of B'
 $= (i, j)$ th element of $A + (i, j)$ th element of B
 $= a_{ij} + b_{ij}$
 $= (j, i)$ th element of $(A+B)'$

Hence $(A+B)' = A' + B'$.

V. Reversal law for a transpose. The transpose of the product of two matrices A, B (conformable for multiplication) is the product of their transposes taken in reverse order i.e.,

$$(AB)' = B'A'$$

Let $A = [a_{ij}]$ be the matrix of order $m \times n$
 and $B = [b_{jk}]$ be the matrix of order $n \times p$.

Then AB is the matrix of order $m \times p$ and so $(AB)'$ is the matrix of order $p \times m$.

Also, A' is the matrix of order $n \times m$ and B' is the matrix of order $p \times n$ so that $B'A'$ is the matrix of order $p \times m$.

As such the order of the matrices $(AB)'$ and $B'A'$ is the same.

Again, (k, i) th element of $(AB)' = (i, k)$ th element of AB

$$= \sum_{j=1}^n a_{ij} b_{jk} \quad \dots (1)$$

Now $A' = [a'_{ji}]$ where $a'_{ji} = a_{ij}$
 and $B' = [b'_{kj}]$ where $b'_{kj} = b_{jk}$

$$(k, i) \text{th element of } B'A' = \sum_{j=1}^n b'_{kj} a'_{ji} = \sum_{j=1}^n b_{jk} a_{ij} = \sum_{j=1}^n a_{ij} b_{jk} \\ = (k, i) \text{th element of } (AB)' \text{ from (1)}$$

Hence $(AB)' = B'A'$

Note. The general reversal law for a transpose may be stated as:

If $A, B, C, \dots, K$ be any number of matrices conformable for multiplication in order, then

$$(ABC \dots JK)' = K'J' \dots C'B'A'$$

[7] Symmetric and Skew-symmetric Matrices

A square matrix A is said to be **symmetric** if its transpose coincides with itself i.e., $A' = A$.

In other words a square matrix $A = [a_{ij}]$ is said to be symmetric if $a_{ij} = a_{ji}$ for all integral values of i and j e.g. if

$$A = \begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix} \text{ then } A' = \begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix}$$

i.e., $A' = A$ and hence A is the symmetric matrix.

It is evident that the total number of independent elements in a symmetric matrix of order n is $\frac{n^2-n}{2} + n$ i.e., $\frac{1}{2}n(n+1)$ since all the diagonal elements are independent and of the remaining elements n the equidistant elements from the diagonal are the same, so that the number of independent elements other than the diagonal elements is $\frac{1}{2}(n^2-n)$.

Again, a square matrix A such that $A' = -A$ is said to be **skew-symmetric**. In other words a square matrix A is skew-symmetric if $a_{ij} = -a_{ji}$ for all integral values of i and j . It follows that $a_{ii} = -a_{ii}$ or $2a_{ii} = 0$ i.e., $a_{ii} = 0$ which shows that all the diagonal elements of a skew-symmetric matrix are zero.

As an illustrative example, the matrix

$$\begin{bmatrix} 0 & 1 & 2 & -3 \\ -1 & 0 & -4 & 5 \\ -2 & 4 & 0 & 6 \\ 3 & -5 & -6 & 0 \end{bmatrix}$$

is skew-symmetric.

The number of independent elements in a skew-symmetric matrix is clearly $\frac{1}{2}(n^2-n)$.

Problem 10. Every square matrix can be uniquely expressed as the sum of a symmetric matrix and a skew-symmetric matrix.

Let A be any square matrix. Then we have

$$A = \frac{1}{2}(A+A') + \frac{1}{2}(A-A')$$

Denoting $\frac{1}{2}(A+A')$ by P and $\frac{1}{2}(A-A')$ by Q we have

$$A = P + Q.$$

$$\begin{aligned} \text{Now } P' &= \left\{ \frac{1}{2}(A+A') \right\}' \\ &= \frac{1}{2} \{A' + (A')'\} \\ &= \frac{1}{2} \{A' + A\} \because (A')' = A \\ &= \frac{1}{2} (A+A') \\ &= P \end{aligned}$$

which follows that P is a symmetric matrix.

$$\begin{aligned} \text{Also } Q' &= \left\{ \frac{1}{2}(A-A') \right\}' \\ &= \frac{1}{2} \{A' - (A')'\} \\ &= \frac{1}{2} \{A' - A\} \\ &= -\frac{1}{2} (A-A') \\ &= -Q \end{aligned}$$

which follows that Q is a skew-symmetric matrix.

Thus the square matrix A is expressible as the sum of a symmetric matrix P and a skew-symmetric matrix Q .

Now to show that this representation is unique, let us assume that if possible A can be expressed as

$$A = R + S$$

where R is a symmetric matrix and S a skew-symmetric matrix. Consider, $A' = (R+S)' = R' + S'$

$$= R - S \text{ as } R = R' \text{ and } S' = -S$$

$$\therefore \frac{1}{2}(A+A') = P = R \text{ and } \frac{1}{2}(A-A') = Q = S$$

i.e., R is not different from P and similarly S is not different from Q , showing that this representation is unique.

[8] The Conjugate of a Matrix

If the elements of a matrix A are complex quantities, then the matrix obtained from A , on replacing its elements by the corresponding conjugate complex numbers, is said to be the **conjugate matrix** of A and is denoted by $\bar{A}$.

Thus if $A = [a_{ij}]$, then $\bar{A} = [\bar{a}_{ij}]$ where $\bar{a}_{ij}$ denotes the conjugate of a_{ij} .

As an illustrative example if

$$A = \begin{bmatrix} 1-5i & -2+7i & 5 & 2i \\ -5+6i & 3-4i & -i & -2 \\ -3-5i & 4+5i & 0 & -7i \end{bmatrix}$$

$$\text{then } \bar{A} = \begin{bmatrix} 1+5i & -2-7i & 5 & -2i \\ -5-6i & 3+4i & i & -2 \\ -3+5i & 4-5i & 0 & 7i \end{bmatrix}$$

The properties of the conjugate matrices

i. The conjugate of the conjugate of a matrix A coincides with itself i.e., $(\bar{\bar{A}}) = A$

Supposing that $A = [a_{ij}]$ is the matrix of order $m \times n$, the matrices A and $\bar{A}$ will also be the matrices of the same order $m \times n$.

Now (i, j) th element of $\bar{A}$ = conjugate of the (i, j) th element of A
 $= \bar{a}_{ij}$

and (i, j) th element of $(\bar{\bar{A}})$ = conjugate of the (i, j) th element of $\bar{A}$
 $=$ conjugate of $\bar{a}_{ij}$
 $= a_{ij}$, since if $a_{ij} = \alpha + i\beta$, then $\bar{a}_{ij} = \alpha - i\beta$
and $(\bar{\bar{a}}_{ij}) = \alpha + i\beta = a_{ij}$
 $= (i, j)$ th element of A

Hence $(\bar{\bar{A}}) = A$.

ii. The conjugate of the sum of two matrices A and B (conformable for addition) is the sum of their conjugates i.e., $(\overline{A+B}) = \bar{A} + \bar{B}$

Suppose that $A = [a_{ij}]$ and $B = [b_{ij}]$ are the matrices of the same order $m \times n$. Then the matrices $\bar{A}$ and $\bar{B}$ will also be of the same order $m \times n$ and the order of the sum matrix $(A+B)$ and so that of its conjugate $(\overline{A+B})$ will be the same i.e., $m \times n$

Now $A+B = [a_{ij}] + [b_{ij}] = [a_{ij} + b_{ij}]$

$$\begin{aligned} (\overline{A+B}) &= \text{conjugate of } [a_{ij} + b_{ij}] \\ &= \text{conjugate of } a_{ij} + \text{conjugate of } b_{ij} \\ &= \bar{a}_{ij} + \bar{b}_{ij} \\ &= \bar{A} + \bar{B} \end{aligned}$$

Hence $(\overline{A+B}) = \bar{A} + \bar{B}$.

iii. If α be a complex number and A a matrix of any order say $m \times n$,

$$(\overline{\alpha A}) = \bar{\alpha} \bar{A}$$

Let $A = [a_{ij}]$ be the matrix of order $m \times n$.

Then $\bar{A}$ will also be the matrix of the order $m \times n$.

$$\begin{aligned} \text{Now, } \alpha A &= \alpha [a_{ij}] \\ &= [\alpha a_{ij}] \end{aligned}$$

$$(\overline{\alpha A}) = \text{conjugate of } [\alpha a_{ij}]$$

$$= [\bar{\alpha} a_{ij}]$$

$$= \bar{\alpha} [\bar{a}_{ij}]$$

$$= \bar{\alpha} \bar{A}$$

IV. The conjugate of the product of two matrices A and B (conformable for multiplication) is the product of their conjugates i.e.

$$(\overline{AB}) = \bar{A} \bar{B}$$

Suppose that $A = [a_{ij}]$ is the matrix of order $m \times n$ and $B = [b_{jk}]$ is the matrix of order $n \times p$

So that AB is the matrix of order $m \times p$.

The order of $(\overline{AB})$ will also be $m \times p$.

Again the orders of $\bar{A}$ and $\bar{B}$ being $m \times n$ and $n \times p$ respectively, the order of $\bar{A} \bar{B}$ will be $m \times p$. Thus the orders of the matrices $(\overline{AB})$ and $\bar{A} \bar{B}$ are equal.

Now, (i, k) th element of $(\overline{AB}) =$ conjugate of (i, j) th element of AB

$$= \text{conjugate of } \sum_{j=1}^n a_{ij} b_{jk}$$

$$= \sum_{j=1}^n \overline{a_{ij} b_{jk}}$$

$$= \sum_{j=1}^n \bar{a}_{ij} \bar{b}_{jk}$$

$$= (i, j)\text{th element of } \bar{A} \bar{B}$$

Hence $\overline{AB} = \bar{A} \bar{B}$.

[9] The conjugate transpose of a Matrix

The matrix, which is the conjugate of the transpose of a matrix A is said to be the conjugate transpose of A and is denoted by A° or A' or $\bar{A}^T$.

As an illustrative example if

$$A = \begin{bmatrix} -2+3i & 3-4i & i \\ -5i & -5-3i & 4+i \end{bmatrix}$$

$$\text{then } A' = \begin{bmatrix} -2-3i & -5i \\ 3-4i & -5-3i \\ i & 4+i \end{bmatrix}$$

$$\text{So that } A^{\circ} \text{ or } \bar{A}' = \begin{bmatrix} -2-3i & 5i \\ 3+4i & -5+3i \\ -i & 4-i \end{bmatrix}$$

It is easy to see that $\bar{A}' = (\bar{A})'$

The properties of conjugate transposed matrices

I. The conjugate transpose of the conjugate transpose of a matrix A coincides with itself i.e., $(A^{\circ})^{\circ} = A$

we have $A^{\circ} = \bar{A}' = (\bar{A})'$

$$\therefore (A^{\circ})' = \{(\bar{A})'\}' = \bar{A} \text{ since } (B')' = B$$

so that $(A^{\circ})^{\circ} \text{ i.e., } (\bar{A}')' = (\bar{A})$

$$= A \because (\bar{A}) = A$$

II. The conjugate transpose of the sum of two matrices A and B (conformable for addition) is the sum of their conjugate transposes i.e.,

$$(A+B)^{\circ} = A^{\circ} + B^{\circ}$$

we have $(A+B)^{\circ} = (\bar{A} + \bar{B})'$

$$= (\bar{A}' + \bar{B}')$$

$$= \bar{A}' + \bar{B}'$$

$$= A^{\circ} + B^{\circ}$$

III. If α be a complex number and A a matrix of any order then

$$(\alpha A)^{\circ} = \bar{\alpha} A^{\circ}$$

we have $(\alpha A)^{\circ} = (\overline{\alpha A})'$

$$= \bar{\alpha} \bar{A}'$$

$$= \bar{\alpha} A^{\circ}$$

IV. The conjugate transpose of the product of two matrices A and B (conformable for multiplication) is the product of their conjugate transposes in reverse order i.e.

$$(AB)^{\circ} = B^{\circ} A^{\circ}$$

we have $(AB)^{\circ} = (\overline{AB})' = \bar{B}' \bar{A}'$

$$= \bar{B}' \bar{A}'$$

$$= B^{\circ} A^{\circ}$$

[10] Hermitian and Skew-Hermitian Matrices

A square matrix $A = [a_{ij}]$ is said to be Hermitian if A coincides with

its conjugate transpose i.e., $A = A^{\circ} (= \bar{A}')$ or if the transpose of A

coincides with its conjugate i.e., $\bar{A}' = A$

Thus $a_{ij} = \bar{a}_{ji}$ for all integral values of i and j .

So that $a_{ii} = \bar{a}_{ii}$ for $i=j$

Which follows that every diagonal element of a Hermitian matrix is equal to its conjugate and it is only possible if all the diagonal elements are real.

As an illustrative example, the matrix

$$\begin{bmatrix} 5 & 2+3i & -i \\ 2-3i & 3 & -3-4i \\ i & -3+4i & 0 \end{bmatrix} \text{ is Hermitian.}$$

A square matrix $A=[a_{ij}]$ is said to be *Skew-Hermitian* if

$$A^0 = -A \text{ or } A' = -\bar{A} \text{ i.e., } \bar{a}_{ij} = -a_{ji} \text{ for all integral values of } i \text{ and } j$$

$$\text{Thus } \bar{a}_{ii} = -a_{ii} \text{ for } i=j$$

$$\text{i.e., } a_{ii} + \bar{a}_{ii} = 0$$

Which follows that every diagonal element of a Skew-Hermitian matrix is either zero or a pure imaginary number.

As an illustrative example, the matrix

$$\begin{bmatrix} 3i & -3+4i & 4-5i \\ +3-4i & -4i & 5 \\ -4-5i & -5 & 0 \end{bmatrix} \text{ is Skew-Hermitian.}$$

Problem 11. Every square matrix A can be uniquely expressed as the sum of a Hermitian matrix and a Skew-Hermitian matrix.

We can express

$$A = \frac{1}{2}(A + A^0) + \frac{1}{2}(A - A^0)$$

$$= P + Q \text{ where } P = \frac{1}{2}(A + A^0), Q = \frac{1}{2}(A - A^0)$$

$$\therefore P^0 = \frac{1}{2}(A + A^0)^0$$

$$= \frac{1}{2}\{A^0 + (A^0)^0\}$$

$$= \frac{1}{2}(A^0 + A) \therefore (A^0)^0 = A$$

$$= \frac{1}{2}(A + A^0)$$

$$= P$$

which follows that P is Hermitian.

$$\text{Also } Q^0 = \frac{1}{2}(A - A^0)^0$$

$$= \frac{1}{2}\{A^0 - (A^0)^0\}$$

$$= \frac{1}{2}(A^0 - A)$$

$$= -\frac{1}{2}(A - A^0)$$

$$= -Q$$

which follows that Q is Skew-Hermitian.

Hence the square matrix A is expressible as the sum of a Hermitian and a Skew-Hermitian matrices.

In order to show that this representation is unique, let us assume if possible that there is another way of representation say

$$A = R + S$$

where R is Hermitian and S is Skew-Hermitian.

$$\text{Consider } A^0 = (R + S)^0 = R^0 + S^0$$

$$= R - S \text{ as } R \text{ is Hermitian and } S \text{ is Skew-Hermitian.}$$

$$\text{These give } R = \frac{1}{2}(A + A^0) = P$$

$$\text{and } S = \frac{1}{2}(A - A^0) = Q$$

which show that R is not different from P and similarly S is not different from Q . Hence this representation is unique.

Problem 12. If A, B, C are three matrices conformable for multiplication in the given order, then show that

$$(ABC)' = C'B'A'$$

We have

$$(ABC)' = \{(AB)C\}'$$

$$= C'(AB)' \text{ by the reversal law of transposes}$$

$$= C'B'A'$$

Problem 13. If A and B be symmetric matrices, then show that AB is symmetric if and only if A and B commute.

Since A and B are symmetric matrices, therefore

$$A' = A \text{ and } B' = B.$$

$$\text{We have } (AB)' = B'A' = BA$$

$$\dots (1)$$

$$\text{But } AB \text{ is symmetric if and only if } (AB)' = AB$$

$$\dots (2)$$

From (1) and (2) it follows that AB is symmetric if

$$AB = BA$$

i.e., if A and B commute.

Conversely if A and B commute, $AB = BA$, then

$$(AB)' = B'A' = BA = AB.$$

Thus if A and B commute AB is symmetric.

Hence AB is symmetric if and only if A and B commute.

Problem 14. If A is a m -square matrix and P is a matrix of order n then show that $B = P'AP$ will be symmetric or skew-symmetric according as A is symmetric or skew-symmetric.

Assuming that A is symmetric, we have

$$B' = (P'AP)' = \{P'(AP)\}' = (AP)'(P')' \text{ by reversal law of transposes}$$

$$= P'A'P \therefore (AP)' = P'A' \text{ and } (P')' = P$$

$$= P'AP \therefore A \text{ is symmetric so that } A' = A$$

$$\text{Thus } (P'AP)' = P'AP$$

which follows that $P'AP$ is symmetric if A is symmetric.

Again if we assume that A is skew-symmetric i.e., $A' = -A$ then

$$B' = (P'AP)' = -P'AP \text{ (proceeding as above)}$$

which follows that $P'AP$ is skew-symmetric when A is skew-symmetric.

Problem 15. If A and B are Hermitian, show that $AB + BA$ is Hermitian and $AB - BA$ is Skew-Hermitian.

A and B are Hermitian

$$A^0 = A \text{ and } B^0 = B$$

$$\text{Consider } (AB + BA)^0 = (AB)^0 + (BA)^0$$

$$= B^0A^0 + A^0B^0$$

$$= BA + AB$$

$$= AB + BA$$

which follows that $AB + BA$ is Hermitian.

Again we have

$$(AB - BA)^0 = (AB)^0 - (BA)^0$$

$$= B^0A^0 - A^0B^0$$

$$= BA - AB$$

$$= -(AB - BA)$$

which follows that $AB - BA$ is skew-Hermitian.

Problem 16. Prove that every Hermitian matrix A can be expressed as $B + iC$ where B is real and symmetric and C is real and skew-symmetric.

$\therefore B$ and C are real,

$\therefore B^0 = B'$ and $C^0 = C'$

(because the conjugate of a real number is itself)

Again since B is symmetric and C is skew-symmetric, we have

$$B' = B \text{ and } C' = -C$$

(1) and (2) yield,

$$B^0 = B \text{ and } C^0 = -C$$

Now assume that

$$A = B + iC$$

$$\therefore A^0 = (B + iC)^0$$

$$= B^0 - iC^0$$

$$= B - i(-C)$$

$$= B + iC = A$$

$\therefore$ Conjugate of i is $-i$

which follows that A is Hermitian.

Hence the result follows.

[11] Adjugate Matrix or Adjoint of a Matrix

If $A = [a_{ij}]$ be a square matrix of any order say n and A_{ij} represent the cofactor of the element a_{ij} in the determinant $|A|$ i.e., $|a_{ij}|$ the square matrix A , then the transpose of the matrix $[A_{ij}]$ is called the **adjugate** or **adjoint** of A and is denoted by $\text{adj } A$.

$$\text{Thus if } A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}, \text{ then } \text{adj } A = \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \dots & \dots & \dots & \dots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix}$$

The determinant of the $\text{adj } A$ i.e., $|\text{adj } A|$ is said to be the adjugate determinant of A or the adjugate of $|A|$.

As an illustrative example if $A = \begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix}$, then

$$\text{cofactor of } a = \begin{vmatrix} b & f \\ f & c \end{vmatrix} = bc - f^2 \text{ in } |A| = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix}$$

$$,, \quad h = - \begin{vmatrix} h & f \\ g & c \end{vmatrix} = fg - ch \quad ,,$$

$$,, \quad g = \begin{vmatrix} h & b \\ g & f \end{vmatrix} = hf - bg \quad ,,$$

etc.

$$\therefore \text{Adj } A = \begin{bmatrix} bc - f^2 & gf - ch & hf - bg \\ gf - ch & ac - g^2 & gh - af \\ hf - bg & gh - af & ab - h^2 \end{bmatrix}$$

An important relation between a matrix A and its adjugate.

If A be a square matrix of any order say n and I the unit matrix of the same order then

$$A (\text{adj } A) = |A| I = (\text{adj } A) A$$

Let $A = [a_{ij}]$ be the square matrix of order n , i.e.,

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

Then if A_{ij} is the cofactor of a_{ij} for all integral values of i and j , we have

$$\text{adj } A = \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \dots & \dots & \dots & \dots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix}$$

Since the orders of A and $\text{adj } A$ are the same, therefore they are conformable for product. Moreover both of them being square matrices, their product is commutative.

Now we know by the properties of a determinant that if the elements of a row (column) of a determinant are multiplied by their own cofactors then the sum of the products is the value of the determinant and the sum of the products of the elements of any row (column) by the cofactors of the corresponding elements of another row (column) is zero i.e., if

$$|A| = \begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix}$$

and $A_{11}, A_{12}, \dots$ are cofactors of $a_{11}, a_{12}, \dots$ etc., then

$$a_{11} A_{11} + a_{12} A_{12} + \dots + a_{1n} A_{1n} = |A| \text{ etc.}$$

$$\text{and } a_{11} A_{21} + a_{12} A_{22} + \dots + a_{1n} A_{2n} = 0 \text{ etc.}$$

Applying these results, we thus have

$$\begin{aligned} A (\text{adj } A) &= \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \dots & \dots & \dots & \dots \\ A_{1n} & A_{2n} & \dots & A_{nn} \end{bmatrix} \\ &= \begin{bmatrix} |A| & 0 & \dots & 0 \\ 0 & |A| & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & |A| \end{bmatrix} \\ &= |A| \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{bmatrix} \\ &= |A| I. \end{aligned}$$

Similarly it may be shown that $(\text{adj } A) A = |A| I$

$$\text{Hence } A (\text{adj } A) = |A| I = (\text{adj } A) A \quad \dots (1)$$

It follows that multiplication of A and $\text{adj } A$ is commutative and that their product is a scalar matrix having every diagonal element as $|A|$.

Note 1. We have shown above that

$$A(\text{adj } A) = \begin{bmatrix} |A| & 0 & \dots & 0 \\ 0 & |A| & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & |A| \end{bmatrix}$$

Taking determinants of either side we get

$$|A(\text{adj } A)| = \begin{vmatrix} |A| & 0 & \dots & 0 \\ 0 & |A| & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & |A| \end{vmatrix}$$

$$\text{or } |A| \cdot |\text{adj } A| = |A|^n \begin{vmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{vmatrix} \text{ by the properties of determinants.}$$

$$\text{i.e., } |A| \cdot |\text{adj } A| = |A|^n$$

$$\text{which gives } |\text{adj } A| = |A|^{n-1}$$

Note 2. If the square matrix A of order n is such that $|A| = 0$, then $A(\text{adj } A) = (\text{adj } A)A = 0$.

Note 3. If A and B are two n -square matrices then

$$\text{adj } AB = \text{adj } B \cdot \text{adj } A$$

which may be shown as below:

Applying the result (1) we have

$$AB(\text{adj } AB) = |AB| \cdot I = (\text{adj } AB)AB$$

$$\begin{aligned} \text{Now } AB \cdot \text{adj } B \cdot \text{adj } A &= A(B \cdot \text{adj } B) \cdot \text{adj } A \\ &= A(|B| \cdot I) \cdot \text{adj } A \text{ by (1)} \\ &= |B| \cdot (A \cdot \text{adj } A) \\ &= |B| \cdot (|A| \cdot I) \text{ by (1)} \\ &= |AB| \cdot I \end{aligned}$$

$$\begin{aligned} \text{and } (\text{adj } B \cdot \text{adj } A)AB &= \text{adj } B[(\text{adj } A)A]B \\ &= \text{adj } B \cdot |A| \cdot I \cdot B \\ &= |A| \cdot \{(\text{adj } B)B\} \\ &= |A| \cdot |B| \cdot I \\ &= |AB| \cdot I \end{aligned}$$

$$\therefore AB(\text{adj } B \cdot \text{adj } A) = (AB)I = (\text{adj } B \cdot \text{adj } A)AB$$

From (α) and (β) we conclude that

$$\text{adj } AB = \text{adj } B \cdot \text{adj } A$$

Note 4. The result of note 3 can be extended to the case of three square matrices A, B, C i.e.,

$$\text{adj } ABC = \text{adj } C \cdot \text{adj } B \cdot \text{adj } A.$$

It is easy to show it as below:

By result (3) we have

$$\begin{aligned} \text{adj } A(BC) &= \text{adj } BC \cdot \text{adj } A \\ &= \text{adj } B \cdot \text{adj } C \cdot \text{adj } A \text{ by (3)} \end{aligned}$$

[12] Singular and Non-singular Matrices

A square matrix $A = [a_{ij}]$ is known as *singular* matrix if its determinant $|A| = 0$ i.e., $|a_{ij}| = 0$. In case $|A| \neq 0$ the matrix A is known as *non-singular* matrix.

ILLUSTRATIVE EXAMPLES

1. The matrix $A = \begin{bmatrix} 21 & 17 & 7 & 10 \\ 24 & 22 & 6 & 10 \\ 6 & 8 & 2 & 3 \\ 5 & 7 & 1 & 2 \end{bmatrix}$ is singular

$$\text{since } \begin{vmatrix} 21 & 17 & 7 & 10 \\ 24 & 22 & 6 & 10 \\ 6 & 8 & 2 & 3 \\ 5 & 7 & 1 & 2 \end{vmatrix} = 0$$

2. The matrix $A = \begin{bmatrix} 12 & 3 & 7 \\ 27 & 7 & 17 \\ 26 & 9 & 22 \end{bmatrix}$ is non-singular since $|A| = 3 \neq 0$.

[13] Reciprocal Matrix or Inverse of a Matrix

[Agra 1970; Allahabad 65; Gorakhpur 1963, 65]

If there exists a square matrix B of the same order as that of A such that $AB = BA = I$, I being a unit matrix of the same order then B is said to be the inverse of A and is denoted by A^{-1} . Thus by definition,

$$AA^{-1} = A^{-1}A = I \quad \dots (1)$$

When the inverse of A exists, then A is said to be *Invertible*.

Inverse of a matrix in terms of its adjugate

According to the definition if A is an invertible matrix, then

$$AA^{-1} = A^{-1}A = I \quad \dots (\alpha)$$

Also, we have from §2.9[11],

$$A(\text{adj } A) = (\text{adj } A)A = |A| \cdot I$$

$$\text{or } A \left(\frac{1}{|A|} \text{adj } A \right) = \left(\frac{1}{|A|} \text{adj } A \right) A = I \quad \dots (\beta)$$

From (α) and (β) it is obvious that

$$AA^{-1} = A \left(\frac{1}{|A|} \text{adj } A \right) \text{ and } A^{-1}A = \left(\frac{1}{|A|} \text{adj } A \right) A$$

either of which lead to

$$A^{-1} = \frac{1}{|A|} \text{adj } A \quad \dots (2)$$

which gives the inverse of a square matrix A in terms of its adjugate.

THEOREM 1. The necessary and sufficient condition for a square matrix to be invertible is that it is non-singular.

To prove that the condition is *necessary* let us assume that A is a given square matrix which is invertible and let B be its inverse.

Then, $AB = I$ (by definition of inverse)

Taking determinant of either side we get,

$$|AB| = |I|$$

$$\text{or } |A| |B| = 1$$

which is only possible if neither of $|A|$ and $|B|$ is zero i.e., if the matrices A and B are non-singular or if $|A| \neq 0$.

To Prove that the condition is sufficient, let us assume that $|A| \neq 0$ and there exists a matrix B such that

$$B = \frac{1}{|A|} \text{adj } A$$

$$\text{Then, } AB = A \frac{1}{|A|} \text{adj } A$$

$$= \frac{1}{|A|} (A \text{adj } A)$$

$$= \frac{1}{|A|} |A| I \text{ from §2.9[11]}$$

$$= I$$

and similarly $BA = I$

$$\text{i.e., } AB = I = BA$$

which shows that A has an inverse i.e., A is invertible.

THEOREM 2. The inverse of a matrix is unique. [Agra, 1970]

Let A be an invertible (square) matrix and if possible let us assume that B and C are two inverses of A , then

$$AB = BA = I$$

$$AC = CA = I$$

$$\text{Now } CAB = C(AB) = CI = C$$

$$\text{Also, } CAB = (CA)B = IB = B$$

$$\text{so that } CAB = B = C$$

i.e., B is not different from C

Hence the inverse of a matrix is unique i.e., there exists only one inverse matrix to a given matrix.

Properties of inverse matrices

I. The Reversal law for inverses. If A and B are two n -square non-singular matrices (conformable for multiplication)

$$\text{then } (AB)^{-1} = B^{-1} A^{-1}$$

Since A and B are non-singular square matrices, they are invertible

Let their inverses be A^{-1} and B^{-1} respectively.

$$\therefore AA^{-1} = A^{-1}A = I$$

$$\text{and } BB^{-1} = B^{-1}B = I$$

Now $|A| \neq 0$, $|B| \neq 0$ imply that $|AB| = |A| |B| \neq 0$ which shows that AB is also invertible.

Let us now consider a matrix C given by

$$C = B^{-1} A^{-1}$$

$$\text{Then, } C(AB) = (B^{-1} A^{-1})(AB)$$

$$= B^{-1} (A^{-1} A) B$$

$$= B^{-1} (I) B$$

$$= B^{-1} B \quad \therefore IB = B$$

$$= I$$

$$\text{i.e., } (B^{-1} A^{-1})(AB) = I$$

Similarly it can be shown that

$$(AB)(B^{-1} A^{-1}) = I$$

These results follow that $B^{-1} A^{-1}$ is the inverse of AB i.e.,

$$(AB)^{-1} = B^{-1} A^{-1} \quad \dots (3)$$

Note. The result can be extended to any number of square matrices which are conformable for multiplication i.e.,

$$(ABC \dots JK)^{-1} = K^{-1} J^{-1} \dots C^{-1} B^{-1} A^{-1}$$

II. If A is a non-singular matrix, then the inverse of its inverse coincides with itself i.e., $(A^{-1})^{-1} = A$

$$\text{We have } AA^{-1} = A^{-1}A = I$$

Which can be interpreted as that A is the inverse of A^{-1} i.e.,

$$(A^{-1})^{-1} = A \quad \dots (4)$$

III. If A is a non-singular matrix, then the transpose (conjugate transpose) of an inverse is the inverse of the transpose (conjugate transpose) of A i.e.

$$(A^{-1})' = (A')^{-1} \text{ and } (A^{-1})^0 = (A^0)^{-1}$$

Since A is non-singular, it is invertible and therefore

$$AA^{-1} = A^{-1}A = I \quad \dots (\gamma)$$

Taking transpose of either side and using $I' = I$, we have

$$(AA^{-1})' = (A^{-1}A)' = I$$

$$\text{or } (A^{-1})'A' = A'(A^{-1}) = I \text{ by reversal law of transposes.}$$

Which follows that $(A^{-1})'$ is the inverse of A' i.e.

$$(A')^{-1} = (A^{-1})' \quad \dots (5)$$

Again taking conjugate transpose of either side of (γ) , we have

$$(AA^{-1})^0 = (A^{-1}A)^0 = I^0$$

$$\text{or } (A^{-1})^0 A^0 = A^0 (A^{-1})^0 = I \quad \therefore I^0 = I$$

which shows that $(A^{-1})^0$ is the inverse of A^0 i.e.

$$(A^0)^{-1} = (A^{-1})^0 \quad \dots (6)$$

IV. If A is invertible then

$$\text{tr. } (ACA^{-1}) = \text{tr. } C$$

[Agra, 1973]

If

$$B = CA^{-1}, \text{ then}$$

$$\text{tr. } (ACA^{-1}) = \text{tr. } (AB)$$

$$= \text{tr. } (BA) \text{ by Cor. of §2.5}$$

$$= \text{tr. } (CA^{-1}A)$$

$$= \text{tr. } CI$$

$$= \text{tr. } C$$

Problem 17. Compute the adjoint of $\begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$

Let $A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$

then $|A| = \begin{vmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{vmatrix}$

The elements of the first row of $|A|$ are 0, 1, 2 and their cofactors are

$$\begin{vmatrix} 2 & 3 \\ 1 & 1 \end{vmatrix}, -\begin{vmatrix} 1 & 3 \\ 3 & 1 \end{vmatrix}, \begin{vmatrix} 1 & 2 \\ 3 & 1 \end{vmatrix} \text{ i.e., } -1, 8, -5 \text{ respectively.}$$

The elements of the second row of $|A|$ are 1, 2, 3 and their cofactors are

$$-\begin{vmatrix} 1 & 2 \\ 1 & 1 \end{vmatrix}, \begin{vmatrix} 0 & 2 \\ 3 & 1 \end{vmatrix}, -\begin{vmatrix} 0 & 1 \\ 3 & 1 \end{vmatrix} \text{ i.e., } 1, -6, 3 \text{ respectively.}$$

The elements of the third row of $|A|$ are 3, 1, 1 and their cofactors are

$$\begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix}, -\begin{vmatrix} 0 & 2 \\ 1 & 3 \end{vmatrix}, \begin{vmatrix} 0 & 1 \\ 1 & 2 \end{vmatrix} \text{ i.e., } -1, 2, -1 \text{ respectively.}$$

$\therefore$ The matrix having its elements as cofactors of $|A|$ is

$$\begin{bmatrix} -1 & 8 & -5 \\ 1 & -6 & 3 \\ -1 & 2 & -1 \end{bmatrix}$$

and therefore $\text{Adj } A = \text{transpose of}$

$$\begin{bmatrix} -1 & 8 & -5 \\ 1 & -6 & 3 \\ -1 & 2 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix}$$

Problem 18. Verify the following:

- The adjoint of an identity matrix is the identity matrix.
- The adjoint of a scalar matrix is a scalar matrix.
- The adjoint of a diagonal matrix is a diagonal matrix.
- The adjoint of a symmetric matrix is a symmetric matrix.
- The adjoint of the transpose of a matrix is equal to the transpose of the adjoint matrix.

(i) Let $I = \begin{bmatrix} 1 & 0 & 0 \dots 0 \\ 0 & 1 & 0 \dots 0 \\ 0 & 0 & 1 \dots 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & 0 \dots 1 \end{bmatrix}$ be an identity matrix of order $n \times n$.

If A_{ij} denotes the cofactor of (i, j) th element of $|I|$ [for all integral values of i and j ranging from 1 to n , then it is clear that

$$A_{ij} = 1 \text{ for } i=j \\ = 0 \text{ for } i \neq j$$

So that $\text{Adj } I = \begin{bmatrix} 1 & 0 & 0 \dots 0 \\ 0 & 1 & 0 \dots 0 \\ 0 & 0 & 1 \dots 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & 0 \dots 1 \end{bmatrix} = I$.

(ii) If $A = \begin{bmatrix} \lambda & 0 \dots 0 \\ 0 & \lambda \dots 0 \\ \dots & \dots & \dots \\ 0 & 0 \dots \lambda \end{bmatrix}$ be a scalar matrix of order $n \times n$, then

as in (i).

$$A_{ij} = \lambda^{n-1} \text{ for } i=j \\ = 0 \text{ for } i \neq j$$

So that $\text{Adj } A = \begin{bmatrix} \lambda^{n-1} & 0 & \dots & 0 \\ 0 & \lambda^{n-1} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \lambda^{n-1} \end{bmatrix}$ which is a scalar matrix.

In other words $A = \lambda I$

$\therefore \text{Adj } A = \text{adj } \lambda I = \lambda^{n-1} I$ which is a scalar matrix.

(iii) Let $D = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$ be a diagonal matrix of order 3×3 . We

shall verify the result for it, but the result can be verified for a diagonal matrix of any order in the similar way.

Then $D_{11} = bc, D_{22} = ac, D_{33} = ab$

$$D_{12} = D_{13} = D_{21} = D_{23} = D_{31} = D_{32} = 0.$$

Thus, $\text{adj } D = \begin{bmatrix} bc & 0 & 0 \\ 0 & ca & 0 \\ 0 & 0 & ba \end{bmatrix}$

which is also a diagonal matrix.

(iv) If A is a symmetric matrix, then $A' = A$ and so $|A'| = |A|$. Thus if A_{ij} denotes the cofactor of (i, j) th element in $|A|$, then

$$A_{ij} = A_{ji} \text{ in } |A|$$

As such the adjoint matrix of the symmetric matrix is also symmetric.

(v) Let $A = \begin{bmatrix} a_{11} & a_{12} \dots a_{1n} \\ a_{21} & a_{22} \dots a_{2n} \\ \dots & \dots & \dots \\ a_{n1} & a_{n2} \dots a_{nn} \end{bmatrix}$

Then, if A_{ij} denotes the cofactor of (i, j) th element,

$$\text{adj } A = \begin{bmatrix} A_{11} & A_{21} \dots A_{n1} \\ A_{12} & A_{22} \dots A_{n2} \\ \dots & \dots & \dots \\ A_{1n} & A_{2n} \dots A_{nn} \end{bmatrix}$$

$$\therefore (\text{Adj } A) = \begin{bmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{n1} & A_{n2} & \dots & A_{nn} \end{bmatrix}$$

$$\text{Again, } A' = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

$$\text{So that } \text{adj } A' = \begin{bmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{n1} & A_{n2} & \dots & A_{nn} \end{bmatrix}$$

It is clear that $(\text{adj } A)' = \text{adj } A'$

Problem 19. If A is a square matrix of order n , then show that

$$\text{adj}(\text{adj } A) = |A|^{n-2} A$$

We have $(\text{adj } A)A = |A| I$

... (1)

We can thus express by replacing A by $\text{adj } A$, as

$$\{\text{adj}(\text{adj } A)\} \text{adj } A = |\text{adj } A| I$$

So that, $\{\text{adj}(\text{adj } A)\} (\text{adj } A) A = |\text{adj } A| A$

or $\text{adj}(\text{adj } A) |A| I = |\text{adj } A| A$ from (1) and since $IA = A$

or $|A| \{\text{adj}(\text{adj } A)\} = |\text{adj } A| A$ $\therefore |\text{adj } A| = |A|^{n-1}$

$\therefore \text{adj}(\text{adj } A) = |A|^{n-2} A$

Problem 20. Find the inverse of the matrix $\begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$

$$\text{Let } A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

$$\therefore |A| = \begin{vmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{vmatrix} = 0 - 1(-1) + 1.1 \text{ expanding about first column} \\ = 2.$$

Now the matrix having its elements as the cofactors of the corresponding elements of $|A|$

$$= \begin{bmatrix} \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} & \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} & \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix} \\ \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} & \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} & \begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} \\ \begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix} & \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix} & \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} \end{bmatrix} = \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix}$$

Its transpose gives the $\text{adj } A$ i.e.,

$$\text{adj } A = \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix}$$

Now we have $A^{-1} = \frac{1}{|A|} \text{adj } A$

$$= \frac{1}{2} \begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \end{bmatrix}$$

Problem 21. If $D = \text{diag}[d_1, d_2, \dots, d_n]$, $d_1, d_2, \dots, d_n \neq 0$, then prove that

$$D^{-1} = \text{diag}[d_1^{-1}, d_2^{-1}, \dots, d_n^{-1}] \quad [\text{Banaras, 1960}]$$

Given

$$D = \begin{bmatrix} d_1 & 0 & 0 & \dots & 0 \\ 0 & d_2 & 0 & \dots & 0 \\ 0 & 0 & d_3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & d_n \end{bmatrix}$$

So that $|D| = \begin{vmatrix} d_1 & 0 & 0 & \dots & 0 \\ 0 & d_2 & 0 & \dots & 0 \\ 0 & 0 & d_3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & d_n \end{vmatrix} = d_1 d_2 \dots d_n \neq 0$ since $d_1, d_2, \dots, d_n \neq 0$

i.e., $|D|$ is non-singular and therefore D is invertible

$$\text{Now } \text{adj } D = \begin{bmatrix} d_2 d_3 \dots d_n & 0 & \dots & 0 & 0 \\ 0 & d_1 d_3 \dots d_n & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & d_1 d_2 \dots d_{n-1} \end{bmatrix}$$

$$\therefore D^{-1} = \frac{1}{|D|} \text{adj } D = \begin{bmatrix} \frac{1}{d_1} & 0 & 0 & \dots & 0 \\ 0 & \frac{1}{d_2} & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & \frac{1}{d_n} \end{bmatrix} \\ = \text{diag} \left[\frac{1}{d_1}, \frac{1}{d_2}, \dots, \frac{1}{d_n} \right] \\ = \text{diag}[d_1^{-1}, d_2^{-1}, \dots, d_n^{-1}]$$

Problem 22. Prove that the inverse of $\begin{bmatrix} A & O \\ B & C \end{bmatrix}$ is $\begin{bmatrix} A^{-1} & O \\ -C^{-1}BA^{-1} & C^{-1} \end{bmatrix}$,

where A, C are non-singular and hence find the inverse of the matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

Suppose that the inverse of the given matrix i.e., $\begin{bmatrix} A & O \\ B & C \end{bmatrix}$ is $\begin{bmatrix} P & Q \\ R & S \end{bmatrix}$

where P, Q, R, S are the submatrices of the inverse matrix which has been partitioned so as to be conformable for premultiplication with the matrix $\begin{bmatrix} A & O \\ B & C \end{bmatrix}$ i.e.,

$$\begin{bmatrix} A & O \\ B & C \end{bmatrix} \begin{bmatrix} P & Q \\ R & S \end{bmatrix} = \begin{bmatrix} I & O \\ O & I \end{bmatrix}$$

$$\text{or } \begin{bmatrix} AP & AQ \\ BP+CR & BQ+CS \end{bmatrix} = \begin{bmatrix} I & O \\ O & I \end{bmatrix}$$

Comparison yields,

$$AP=I \text{ i.e., } P=A^{-1} \text{ as } A \text{ is non-singular} \quad \dots (1)$$

$$AQ=O \text{ i.e., } Q=O \text{ as } A \text{ is non-singular} \quad \dots (2)$$

$$BP+CR=O \text{ and } BQ+CS=I$$

$$\text{or } BA^{-1}+CR=O \text{ and } CS=I \text{ from (1) and (2) respectively}$$

$$\text{or } C^{-1}BA^{-1}+C^{-1}CR=O \text{ and } S=C^{-1} \text{ as } C \text{ is non-singular}$$

$$\text{or } R=-C^{-1}BA^{-1} \text{ as } C^{-1}CR=IR=R \quad \dots (3)$$

$$\text{and } S=C^{-1} \quad \dots (4)$$

Putting the values of P, Q, R, S , the inverse of the matrix $\begin{bmatrix} A & O \\ B & C \end{bmatrix}$ is

$$\begin{bmatrix} P & Q \\ R & S \end{bmatrix} \text{ i.e., } \begin{bmatrix} A^{-1} & O \\ -C^{-1}BA^{-1} & C^{-1} \end{bmatrix}$$

Now to find the inverse of $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix}$, let us partition it as

$$\begin{bmatrix} 1 & 0 & \vdots & 0 & 0 \\ 1 & 1 & \vdots & 0 & 0 \\ \hline 1 & 1 & \vdots & 1 & 0 \\ 1 & 1 & \vdots & 1 & 1 \end{bmatrix}$$

When it is compared with $\begin{bmatrix} A & O \\ B & C \end{bmatrix}$, we have

$$A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, C = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

$$\text{Now it is easy to compute } A^{-1} = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix}$$

$$C^{-1} = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix}$$

As above, $\begin{bmatrix} P & Q \\ R & S \end{bmatrix}$ is the inverse of $\begin{bmatrix} A & O \\ B & C \end{bmatrix}$ when

$$P=A^{-1} = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix}$$

$$Q=O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\begin{aligned} R &= C^{-1}BA^{-1} = -\begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} \\ &= -\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} \\ &= -\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \end{aligned}$$

$$= \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix}$$

and

$$S=C^{-1} = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix}$$

$$\text{Hence the inverse of } \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix} \text{ is } \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\text{i.e., } \begin{bmatrix} 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix}$$

[14] Unitary Matrix and Orthogonal Matrix

A square matrix A having its elements as complex numbers is said to be unitary if

$$A' = (\bar{A})^{-1} \text{ or } (\bar{A})' = A^{-1} \text{ or } A^{\circ}A = I$$

In case A is real i.e., the matrix A consists of real numbers as its elements, the matrix A will be unitary if $A'A=I$ for, in this case $A^{\circ}=A'$.

As an illustrative example the matrix

$$A = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \text{ is unitary, since}$$

$$A^{\circ} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$\begin{aligned} \text{So that } A^{\circ}A &= \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I. \end{aligned}$$

Hence the matrix A is unitary.

A real unitary matrix is known as orthogonal matrix i.e., a real matrix A is orthogonal if $A'A=I=AA'$ for, in this case $A'=A^{-1}$.

The determinant of an orthogonal matrix is +1 or -1.

The orthogonal matrix is said to be proper or improper according to its determinant is +1 or -1.

$$\text{Since } |AA'| = |I| = 1$$

i.e., $|A| = |A'| = 1$
 or $|A|^2 = 1$ as $|A| = |A'|$
 we have $|A| = \pm 1$

As an illustrative example the matrix

$$A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \text{ is orthogonal, since}$$

$$AA' = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

and $|A| = |A'| = \begin{vmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{vmatrix} = \cos^2 \theta + \sin^2 \theta = 1$

[15] Normal Matrix and Normal Form of the Matrix

A square matrix A is said to be **Normal** if $AA^0 = A^0A$.

The normal matrices include diagonal, real symmetric, real skew symmetric, orthogonal, Hermitian, Skew-Hermitian and unitary matrices.

As an illustrative example the unitary matrix

$$A = \begin{bmatrix} \frac{1+i}{2} & \frac{-1+i}{2} \\ \frac{1+i}{2} & \frac{1-i}{2} \end{bmatrix} \text{ is Normal, since}$$

$$AA^0 = I = A^0A$$

The normal form of a matrix is $\begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}$

[16] Elementary Matrix and Elementary Operations

A square matrix of order n is said to be **elementary matrix**, if it is obtained from a unit matrix I_n by subjecting it to any of the following elementary operations (transformations):

(i) Interchange of any two rows (columns) to be denoted by R_{ij} (C_{ij}) for the interchange of i th and j th rows (columns) and the elementary matrix obtained may be denoted by E_{ij} .

(ii) Multiplication of elements of any row (column) by any non-zero scalar, to be denoted by $R_i(\lambda)$ ($C_i(\lambda)$) for the multiplication of i th row (column) by $\lambda \neq 0$ and the elementary matrix obtained may be denoted by $E_i(\lambda)$.

(iii) Addition to the elements of any row (column) the corresponding elements of another row (column) multiplied by any non-zero scalar, to be denoted by $R_{ij}(\lambda)$ ($C_{ij}(\lambda)$) for the addition to i th row (column) of the j th row (column) multiplied by $\lambda \neq 0$ and the elementary matrix obtained may be denoted by $E_{ij}(\lambda)$ for row operation and by $E'_{ij}(\lambda)$ for column operation since $E'_{ij}(\lambda)$ is the transpose of $E_{ij}(\lambda)$.

It may be verified that

$$|E_{ij}| = -1$$

$$|E_i(\lambda)| = \lambda \neq 0$$

$$|E_{ij}(\lambda)| = |E'_{ij}(\lambda)| = 1$$

which show that no elementary matrix is singular.

As illustrative examples, the elementary matrices obtained from

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ are}$$

$$E_{12} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_2(\lambda) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } E_{23}(\lambda) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & \lambda \\ 0 & 0 & 1 \end{bmatrix}$$

An elementary operation is said to be the **Row operation** or **Column operation** according as it is applied to rows or columns.

By the elementary transformations or rather say that by a series of elementary row and column transformations, a matrix can be reduced to the normal form. This part will be discussed in the section 'Rank of a matrix' in more details.

THEOREM. Every elementary row (column) transformation of a matrix can be brought about by pre-multiplication (post-multiplication) with the corresponding elementary matrix. [Allahabad, 1966]

Consider two matrices A and B of orders $m \times n$ and $n \times p$ respectively such that

$$A = \begin{bmatrix} R_1 \\ R_2 \\ \vdots \\ R_m \end{bmatrix} \text{ and } B = [C_1, C_2, \dots, C_p]$$

where $R_1, R_2, \dots, R_m$ are the rows of A and $C_1, C_2, \dots, C_p$ are the columns of B . Then by the 'row by column' rule for multiplication we have

$$AB = \begin{bmatrix} R_1 C_1 & R_1 C_2 & \dots & R_1 C_p \\ R_2 C_1 & R_2 C_2 & \dots & R_2 C_p \\ \vdots & \vdots & \ddots & \vdots \\ R_m C_1 & R_m C_2 & \dots & R_m C_p \end{bmatrix}$$

which follows that if the rows of A are subjected to any elementary (column) transformation, then the rows (columns) of AB are also subjected to the same transformation.

Conclusively every elementary row (column) transformation of the product AB can be effected by subjecting the prefactor A (post factor B) to the same row (column) transformation.

We shall apply this result to the required proposition. Suppose A is a matrix of order $m \times n$ and I an identity matrix of order m . Then it is obvious that

$$A = IA.$$

The above result can be applied to show that every elementary row transformation of the product A can be effected by subjecting the prefactor I to the same transformation i.e., by pre-multiplying A by the corresponding elementary matrix.

Again taking $A=AI$, it can be similarly shown that every column transformation of the product A can be effected by subjecting the post factor I to the same transformation *i.e.*, by post-multiplying A by the corresponding elementary matrix.

As an illustrative example, take

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix}$$

and suppose that R_{12} operation gives

$$B = \begin{bmatrix} a_{21} & a_{22} & a_{23} & a_{24} \\ a_{11} & a_{12} & a_{13} & a_{14} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix}$$

Take the elementary matrix $E_{12} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ obtained from the

identity matrix $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ of order 3×3 .

$$\begin{aligned} \text{Consider } E_{12}A &= \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} \\ &= \begin{bmatrix} a_{21} & a_{22} & a_{23} & a_{24} \\ a_{11} & a_{12} & a_{13} & a_{14} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} \\ &= B \end{aligned}$$

which shows that the product B is effected by subjecting the prefactor E_{12} to the same row operation *i.e.*, by pre-multiplying the matrix A by the corresponding elementary matrix.

Inverses of the elementary matrices

E_{ij} being the elementary matrix obtained by interchanging i th and j th rows (columns) of an identity matrix I , may yield back the original identity matrix I on interchanging i th and j th rows. But by the preceding theorem the interchange of i th and j th rows (columns) of E_{ij} may be effected by pre-multiplying (post-multiplying) by the corresponding elementary matrix E_{ij} *i.e.*

$$\begin{aligned} E_{ij} E_{ij} &= I \\ (E_{ij})^{-1} &= E_{ij} \end{aligned}$$

or

which shows that E_{ij} is its own inverse.

Again if $E_i(\lambda)$ is a matrix obtained by multiplying the i th row (column) of an identity matrix I by λ then I may be obtained back by multiplying the i th row (column) of $E_i(\lambda)$ by λ^{-1} . But by the preceding theorem the multiplication of i th row (column) of $E_i(\lambda)$ by λ^{-1} (*i.e.*, λ^{-1}) may be obtained by pre-multiplying (post-multiplying) $E_i(\lambda)$ by the corresponding elementary matrix $E_i(\lambda^{-1})$ *i.e.*

$$\begin{aligned} E_i(\lambda^{-1}) E_i(\lambda) &= I & [E_i(\lambda) E_i(\lambda^{-1})] \\ \text{so that} & & \{E_i(\lambda)\}^{-1} = E_i(\lambda^{-1}) \end{aligned}$$

which follows that the inverse of $E_i(\lambda)$ is $E_i(\lambda^{-1})$.

Further $E_{ij}(\lambda)$ being the elementary matrix obtained by adding the elements of j th row multiplied by λ to the elements of i th row of an identity matrix I , the original identity matrix I may be obtained back by adding to the elements of i th row of $E_{ij}(\lambda)$ the corresponding elements of j th row multiplied by $-\lambda$. But the same row transformation on $E_{ij}(\lambda)$ may also be effected on pre-multiplying $E_{ij}(\lambda)$ by the corresponding elementary matrix $E_{ij}(-\lambda)$ *i.e.*,

$$E_{ij}(-\lambda) E_{ij}(\lambda) = I$$

or

$$\{E_{ij}(\lambda)\}^{-1} = E_{ij}(-\lambda)$$

which shows that the inverse of $E_{ij}(\lambda)$ is $E_{ij}(-\lambda)$.

The same result may be shown to hold in case of column operation.

Conclusively the inverse of an elementary matrix is also an elementary matrix of the same type.

[17] Equivalent Matrix

A matrix B is called the equivalent to a matrix A if B can be obtained from A by a sequence of elementary transformations and denoted by $B \sim A$.

As an illustrative example if

$$A = \begin{bmatrix} -6 & -2 & -4 & 5 \\ 3 & 4 & 5 & -1 \\ 6 & 2 & 4 & -3 \end{bmatrix}$$

$$\text{then } A \sim \begin{bmatrix} 0 & 0 & 0 & 2 \\ 3 & 4 & 5 & -1 \\ 6 & 2 & 4 & -3 \end{bmatrix} \text{ by } R_{13} (1)$$

THEOREM 1. Two matrices A and B are equivalent if and only if there exist non-singular matrices R and C such that

$$RAC = B$$

where $R = R_1 R_2 \dots R_n$ and $C = C_1 C_2 \dots C_m$; R 's being operations affecting rows and C 's those affecting columns.

If A and B are equivalent matrices, then B can be obtained from A by a series of elementary operations. But the elementary row (column) transformation can be effected by pre-multiplying (post-multiplying) A by the corresponding elementary matrix, therefore if we denote the elementary row transformations by $R_1, R_2, \dots, R_n$ and elementary column transformations by $C_1, C_2, \dots, C_m$ where R_1 represents the first elementary matrix corresponding to the first elementary row transformation, etc., and similarly C_1 represents the first elementary matrix corresponding to the first elementary column transformation etc., then

$$(R_n \dots R_2 R_1) A (C_1 C_2 \dots C_m) = B$$

$$RAC = B$$

where R and C are non-singular matrices.

COROLLARY. There are three fundamental properties of the relation $RAC = B$:

(i) *Reflexivity*. Every matrix A is equivalent to itself i.e., if we take $R=I$, $C=I$, then $IAI=A$

(ii) *Symmetry*. If a matrix B is equivalent to another matrix A , then A is also equivalent to B , for if

$$B=RAC$$

then $A=R^{-1}B C^{-1}$

where R^{-1} and C^{-1} are non-singular matrices.

(iii) *Transitivity*. If A is equivalent to B and B is equivalent to C then A is also equivalent to C , for if

$$A=PBQ \text{ and } B=RCS$$

then $A=PRCSQ=(PR)C(SQ)$

where PR and SQ being the products of non-singular matrices, are themselves non-singular.

THEOREM 2. If a square matrix A is reduced to an identity matrix I by a series of elementary row operations, then the same series of row operations applied to I yields the inverse of A i.e., A^{-1} .

Suppose that $E_1, E_2, \dots, E_r$ are the elementary matrices and I the identity matrix, such that

$$(E_r \dots E_2 E_1) A = I$$

Then, post-multiplying both sides by A^{-1} , we get

$$(E_r \dots E_2 E_1) A A^{-1} = I A^{-1}$$

$$\text{i.e., } (E_r \dots E_2 E_1) I = A^{-1}$$

which proves the proposition.

Note. In practice, we write, $A=IA$

and perform a series of elementary row operations on A and prefactor I till A is reduced to I and I is reduced to B such that

$$I=BA$$

which follows that B is inverse of A

$$\text{As an illustrative example if } A = \begin{bmatrix} 1 & 3 & -2 \\ -3 & 0 & -5 \\ 2 & 5 & 0 \end{bmatrix}$$

then take $A \quad I$

$$\begin{bmatrix} 1 & 3 & -2 \\ -3 & 0 & -5 \\ 2 & 5 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{or } \begin{bmatrix} 1 & 3 & -2 \\ 0 & 9 & -11 \\ 0 & -1 & 4 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix} \text{ by } R_{21}(3) \text{ and } R_{31}(-2)$$

$$\text{or } \begin{bmatrix} 1 & 0 & 10 \\ 0 & 1 & 21 \\ 0 & -1 & 4 \end{bmatrix} \sim \begin{bmatrix} -5 & 0 & 3 \\ -13 & 1 & 8 \\ -2 & 0 & 1 \end{bmatrix} \text{ by } R_{13}(3) \text{ and } R_{23}(8)$$

$$\text{or } \begin{bmatrix} 1 & 0 & 10 \\ 0 & 1 & 21 \\ 0 & 0 & 25 \end{bmatrix} \sim \begin{bmatrix} -5 & 0 & 3 \\ -13 & 1 & 8 \\ -15 & 1 & 9 \end{bmatrix} \text{ by } R_{32}(1)$$

$$\text{by } R_{13}(-2/5) \text{ and } R_{23}(-\frac{2}{25})$$

$$\text{by } R_3(\frac{1}{25})$$

$$\text{This gives, } A^{-1} = \begin{bmatrix} 1 & -\frac{2}{5} & -\frac{3}{5} \\ -\frac{2}{5} & \frac{4}{25} & \frac{1}{25} \\ -\frac{3}{5} & \frac{1}{25} & \frac{9}{25} \end{bmatrix}$$

In practice we may write the steps taken in this example as,

$$\text{write } \begin{bmatrix} 1 & 3 & -2 \\ -3 & 0 & -5 \\ 2 & 5 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A \text{ (the form } A=IA)$$

$$\text{by } R_{21}(3) \text{ and } R_{31}(-2)$$

$$\text{by } R_{13}(3) \text{ and } R_{23}(8)$$

$$\text{by } R_{32}(1)$$

$$\text{by } R_{13}(-\frac{2}{5}) \text{ and } R_{23}(-\frac{2}{25})$$

$$\text{by } R_3(\frac{1}{25})$$

which gives the required inverse.

(18) Canonical Matrix

This is a non-zero matrix in which

(i) the first few rows have non-zero elements while the elements of succeeding rows may be all zero,

(ii) the first non-zero row is unity, and

(iii) all the other elements of a column which contains the first non-zero element as unity, are zero.

As an illustrative example, the matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ is Canonical.}$$

Problem 23. Show that the matrix $\begin{bmatrix} \alpha+i\gamma & -\beta+i\delta \\ \beta+i\delta & \alpha-i\gamma \end{bmatrix}$ is unitary if

$$\alpha^2 + \gamma^2 + \delta^2 = I.$$

Denoting the given matrix by A , it will be unitary if

$$A^{\circ}A=I$$

$$\begin{aligned}\text{Consider, } A^*A &= \begin{bmatrix} \alpha - i\gamma & \beta - i\delta \\ -\beta - i\delta & \alpha + i\gamma \end{bmatrix} \begin{bmatrix} \alpha + i\gamma & -\beta + i\delta \\ \beta + i\delta & \alpha - i\gamma \end{bmatrix} \\ &= \begin{bmatrix} \alpha^2 + \beta^2 + \gamma^2 + \delta^2 & 0 \\ 0 & \alpha^2 + \beta^2 + \gamma^2 + \delta^2 \end{bmatrix}\end{aligned}$$

which becomes an identity matrix if

$$\alpha^2 + \beta^2 + \gamma^2 + \delta^2 = 1$$

Hence A is unitary if $\alpha^2 + \beta^2 + \gamma^2 + \delta^2 = 1$.

Problem 24. If A is a unitary matrix and $B = AP$ where $P \neq 0$, then show that PB^{-1} is unitary.

Let $C = PB^{-1}$

Then $C = P(AP)^{-1}$

$= P P^{-1} A$ by reversal law of inverses.

or $C = IA = A$

which follows that C is unitary, since A is unitary.

Problem 25. If A is real skew-symmetric matrix such that $A^2 + I = 0$, then show that A is orthogonal.

Given $A^2 + I = 0$

$\therefore A$ is given to be real skew-symmetric,

$\therefore A = -A'$

Premultiplying both sides by A , we get

$$A^2 = -AA'$$

or $-A^2 = AA'$

Adding (1) and (2) we find,

$$A A' = I$$

which follows that the matrix A is orthogonal.

Problem 26. Reduce the matrix A to its normal form, where

$$A = \begin{bmatrix} 0 & 1 & 2 & -2 \\ 4 & 0 & 2 & 6 \\ 2 & 1 & 3 & 1 \end{bmatrix}$$

We have, $A \sim \begin{bmatrix} 1 & 0 & 2 & -2 \\ 0 & 4 & 2 & 6 \\ 1 & 2 & 3 & 1 \end{bmatrix}$ by C_{12}

$$\sim \begin{bmatrix} 1 & 0 & 2 & -2 \\ 0 & 4 & 2 & 6 \\ 0 & 2 & 1 & 3 \end{bmatrix} \text{ by } R_{31}(-1)$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 4 & 2 & 6 \\ 0 & 2 & 1 & 3 \end{bmatrix} \text{ by } C_{21}(-2) \text{ and } C_{31}(2)$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & \frac{1}{2} & \frac{3}{2} \\ 0 & 2 & 1 & 3 \end{bmatrix} \text{ by } R_2(\frac{1}{4})$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & \frac{1}{2} & \frac{3}{2} \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } R_{32}(-2)$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } C_{32}(-\frac{1}{2}) \text{ and } C_{42}(-\frac{3}{2})$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by partitioning}$$

$$\sim \begin{bmatrix} I_2 & 0 \\ 0 & 0 \end{bmatrix}$$

which gives the normal form of A .

Problem 27. Applying elementary transformations, find the inverse of the matrix

$$A = \begin{bmatrix} 0 & 2 & 1 & 3 \\ 1 & 1 & -1 & -2 \\ 1 & 2 & 0 & 1 \\ -1 & 1 & 2 & 6 \end{bmatrix}$$

We may write, $A = IA$

$$\text{i.e., } \begin{bmatrix} 0 & 2 & 1 & 3 \\ 1 & 1 & -1 & -2 \\ 1 & 2 & 0 & 1 \\ -1 & 1 & 2 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} A$$

$$\text{or } \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 2 & 1 & 4 \\ 0 & 3 & 2 & 7 \\ -1 & 1 & 2 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix} A \text{ by } R_{12}(1), R_{24}(1) \text{ and } R_{34}(1)$$

$$\text{or } \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 2 & 1 & 4 \\ 0 & 3 & 2 & 7 \\ 0 & 4 & 2 & 7 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \end{bmatrix} A \text{ by } R_{41}(1)$$

$$\text{or } \begin{bmatrix} 1 & 3 & 0 & 1 \\ 0 & 2 & 1 & 4 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & -1 & 1 & 0 \\ 1 & -1 & 0 & -1 \end{bmatrix} A \text{ by } R_{32}(-1) \text{ and } R_{42}(-2)$$

$$\text{or } \begin{bmatrix} 1 & 3 & 0 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 & -1 \\ 4 & -3 & 0 & -3 \\ 3 & -4 & 1 & -3 \\ 1 & -1 & 0 & -1 \end{bmatrix} A \text{ by } R_{14}(1), R_{24}(4) \text{ and } R_{34}(3)$$

$$\text{or } \begin{bmatrix} 1 & 3 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 & -1 \\ 1 & 1 & -1 & 0 \\ 3 & -4 & 1 & -3 \\ -1 & 1 & 0 & 1 \end{bmatrix} A \text{ by } R_{23}(-1) \text{ and } R_4(-1)$$

$$\text{or } \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -1 & -3 & 3 & -1 \\ 1 & 1 & -1 & 0 \\ 2 & -5 & 2 & -3 \\ -1 & 1 & 0 & 1 \end{bmatrix} \begin{matrix} \text{A by } R_{12}(-3) \text{ and} \\ R_{32}(-1) \end{matrix}$$

$$\text{which follows that } A^{-1} = \begin{bmatrix} -1 & -3 & 3 & -1 \\ 1 & 1 & -1 & 0 \\ 2 & -5 & 2 & -3 \\ -1 & 1 & 0 & 1 \end{bmatrix}$$

[19] Derogatory and Non-Derogatory Matrices

An n -rowed matrix is said to be derogatory or non-derogatory according as the degree of its minimal equation is less than or equal to n .

Note. If $m(x)$ is a scalar polynomial of the lowest degree with leading coefficient unity, such that $m(A)=0$ then the polynomial $m(x)$ and the equation $m(x)=0$ are respectively known as the **minimal polynomial** and the **minimal equation** of the matrix A . It should be committed to memory (i) the minimal polynomial of a matrix is unique (ii) the minimal polynomial of a matrix is a divisor of every polynomial that annihilates the matrix.

For example if A be a n -square matrix with each element unity, then $A^2=nA$ and the polynomial x^2-nx annihilates A . Hence x^2-nx is the minimal polynomial of A .

THEOREM. Every unit matrix of order ≥ 2 is derogatory.

If I be a unit matrix of order $n (\geq 2)$, then the polynomial $m(x) = x-1$ annihilates I so that $x-1$ is the minimal polynomial of I , since degree of $x-1$ is 1 ($< n$) therefore I is derogatory.

As an illustration $A = \begin{bmatrix} 7 & 4 & -1 \\ 4 & 7 & -1 \\ -4 & -4 & 4 \end{bmatrix}$ is derogatory.

Note. The minimal polynomial of a matrix is a divisor of every polynomial that annihilates the matrix.

Let $m(x)$ be a minimal polynomial of A and $h(x)$ be another polynomial that annihilates A . Then Division algorithm leads

$$h(x) = m(x)q(x) + r(x) \quad \dots (1)$$

where $r(x)$ is a zero polynomial or its degree is less than that of $m(x)$.

Put $x=A$ in (1),

$$h(A) = m(A)q(A) + r(A) \quad \dots (2)$$

$\therefore m(x)$ and $h(x)$ both annihilate A , (2) gives

$$0 = 0q(A) + r(A) \text{ i.e., } r(A) = 0$$

showing that $r(x)$ also annihilates A . If $r(x) \neq 0$, then it is a non-zero polynomial of degree less than that of $m(x)$ and thereby contradicting that $m(x)$ is minimal polynomial and hence the only possibility is that $r(x)$ is zero polynomial.

Then (1) gives $h(x) = m(x)q(x)$

i.e., $m(x)$ is a divisor of $h(x)$.

2.10. RANK OF A MATRIX

While defining a matrix, we have already mentioned that the matrix constituted by the array of elements which are left after deleting any number of rows or columns or both of a matrix is said to be its **sub-matrix**. A square sub-matrix of a square matrix is known as a **principal sub-matrix** which may be obtained by deleting the corresponding rows and columns of the square matrix. If it is obtained by deleting only some of the last rows and the corresponding columns of the square matrix, then it is called as **leading sub-matrix**.

It is worth-noting that a square sub-matrix is not always obtained from a square matrix, but it may be obtained from any matrix after deleting certain numbers of row and columns such that the remaining matrix may have equal number of rows and columns.

The square sub-matrices of a matrix play an important role in deciding the rank of the matrix, for, the maximum order of the non-singular square sub-matrix of a matrix determines its **rank**.

We have also mentioned that the determinant of a square sub-matrix of a matrix A is said to be the **minor** of the matrix. The minor is principal or leading according as the corresponding sub-matrix is principal or leading.

As an illustrative example of $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$, then 1, 2, 3, 4, 5, 6,

7, 8, 9 are the minors of order 1;

$$\begin{vmatrix} 1 & 2 \\ 4 & 5 \end{vmatrix}, \begin{vmatrix} 1 & 3 \\ 4 & 6 \end{vmatrix}, \begin{vmatrix} 2 & 3 \\ 5 & 6 \end{vmatrix}, \begin{vmatrix} 1 & 2 \\ 7 & 8 \end{vmatrix}, \begin{vmatrix} 1 & 3 \\ 7 & 9 \end{vmatrix}, \begin{vmatrix} 2 & 3 \\ 8 & 9 \end{vmatrix},$$

$$\begin{vmatrix} 4 & 5 \\ 7 & 8 \end{vmatrix}, \begin{vmatrix} 4 & 6 \\ 7 & 9 \end{vmatrix}, \begin{vmatrix} 5 & 6 \\ 8 & 9 \end{vmatrix} \text{ are the minors of order 2.}$$

The sub-matrix $\begin{bmatrix} 5 & 6 \\ 8 & 9 \end{bmatrix}$ is a principal sub-matrix of A .

The sub-matrix $\begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix}$ is the leading sub-matrix of A .

It is also clear that the square sub-matrices of orders 1, 2, 3 are all non-singular, which follows that the maximum order of the non-singular square sub-matrix of A is 3 and hence the rank of A is 3.

Definition of Rank of a Matrix

Let A be a given matrix of order $m \times n$ and ρ be a natural number such that

$$\rho \leq \min(m, n)$$

where $\min(m, n) = \text{the smaller of } m \text{ and } n \text{ for } m \neq n$
 $= m = n \text{ for } m = n$

Let us now delete any $(m-\rho)$ rows and $(n-\rho)$ columns of the matrix A , so that the retained elements constitute a square sub-matrix of order ρ , whose determinant is a minor of the matrix A , of order ρ . Clearly there corresponds a system of minors of A to each admissible

value of ρ . A definite positive integral value r of ρ , with the following two properties is known as the **rank** of the matrix A :

- (i) There is at least one minor of order r , which does not vanish.
- (ii) All the minors of order $(r+1)$ vanish.

Conclusively the rank r of a matrix is the largest integer for which the statement "not all minors of order r are zero" is valid.

Thus the rank of a null-matrix is zero, the rank of a non-singular square matrix of order n is n and the rank of a singular square matrix of order n is less than n .

The rank of a matrix A is denoted by $\rho(A)$ or Rank (A) .

In determining the rank of a matrix, the following statement is often found useful :

"If in a given matrix a certain r -rowed determinant is not zero and all the $(r+1)$ -rowed determinants of which this r -rowed determinant is a first minor are zero, then all $(r+1)$ -rowed determinants of the matrix are zero".

ILLUSTRATIVE EXAMPLES

1. The rank of the matrix $\begin{bmatrix} 1 & 3 & 4 & 7 \\ 2 & 4 & 5 & 8 \\ 3 & 1 & 2 & 3 \end{bmatrix}$ is 3, for the minor of

order 4 is obviously zero and none of the minors of order 3 are zero.

2. The rank of $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 7 \\ 3 & 6 & 10 \end{bmatrix}$ is 2.

3. The rank of $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \end{bmatrix}$ is 1.

2.11. SOME THEOREMS ON RANK

THEOREM 1. Elementary operations do not change the rank of a matrix.

Let $A=[a_{ij}]$ be the matrix of order $m \times n$ and rank r and let B be the matrix obtained from A by elementary operations. Suppose that $r' (\neq r)$ is the rank of B .

Case I. Let us first assume that B is the elementary matrix obtained from A by interchange of a pair of rows.

Suppose that B_0 is any $(r+1)$ -rowed square sub-matrix of B , so that $(r+1)$ rows of the sub-matrix B_0 of B are also the rows of some uniquely determined sub-matrix A_0 of A , with the only difference that the identical rows of A_0 and B_0 may or may not occur in the same relative positions. But we know that an interchange of a pair of rows of a determinant results in the change of its sign, therefore we may have

$$|B_0| = |A_0| \text{ or } |B_0| = -|A_0|$$

Now the rank of the matrix A being r every $(r+1)$ -rowed minor

of A vanishes i.e., $|A_0| = 0$ and hence $|B_0| = 0$, showing that every $(r+1)$ -rowed minor of B also vanishes. As such r' , the rank of B can not exceed r , the rank of A i.e.,

$$r' \leq r$$

Moreover A can also be obtained from B by interchange of rows and therefore a similar procedure will yield that

$$r \leq r'$$

The results $r' \leq r$ and $r \leq r'$ imply that $r = r'$ i.e., the interchange of a pair of rows does not alter the rank of a matrix.

It can be similarly shown that the same result holds for column-operation.

Case II. Let us again assume that B is the elementary matrix obtained from A by multiplying the elements of a row by a non-zero scalar say $\lambda \neq 0$.

If $|B_0|$ is any $(r+1)$ -rowed minor of B , then $(r+1)$ rows of the sub-matrix B_0 of B are the $(r+1)$ rows of A_0 of A or one of the rows of $(r+1)$ is multiplied by λ the corresponding row of A_0 , which happens when B_0 contains the affected row. Applying the property of a determinant that the multiplication of any of its row by λ results in the multiplication of the determinant by λ , we have

$$|B_0| = |A_0| \text{ or } |B_0| = \lambda |A_0|$$

But the rank of A being r , every minor $|A_0|$ of order $r+1$ vanishes i.e., $|A_0| = 0$ and hence $|B_0| = 0$ showing that every $(r+1)$ -rowed minor of B also vanishes. As such r' , the rank of B cannot exceed r , the rank of A i.e.,

$$r' \leq r$$

Moreover A can be obtained from B by multiplying the row of B (which was assumed to be multiplied by λ) by $\frac{1}{\lambda}$ and therefore a similar procedure will yield

$$r \leq r'$$

Thus $r' \leq r$ and $r \leq r'$ imply that $r = r'$ i.e., the multiplication of the elements of a row of a matrix by a non-zero scalar λ , does not alter its rank.

It can be similarly shown that the same result holds for column-operation.

Case III. Let us further assume that B is the elementary matrix obtained by adding to the elements of i th row of the matrix A , the products with any non-zero scalar λ of the corresponding elements of another say j th row of the matrix.

If $|B_0|$ is any $(r+1)$ -rowed minor of B corresponding to the minor $|A_0|$ of A , then it is apparent that if the sub-matrix A_0 does not contain the i th row of A , $|B_0| = |A_0|$, also if i th and j th rows of A are contained in A_0 , $|B_0| = |A_0|$, because by the property of a determinant, if the elements of any row are increased by the same multiple

of the corresponding elements of another row, the determinant remains unchanged.

But if i th row of A is contained in A_0 and j th row not, then

$$|B_0| = |A_0| + \lambda |C_0|$$

where C_0 is a $(r+1)$ -rowed square sub-matrix of A such that all the rows of C_0 except the i th coincide with those of A_0 while i th row is obtained by replacing the elements of A_0 in the i th row by λ times the corresponding elements in the j th row. Clearly $|C_0|$ is a $(r+1)$ rowed minor of A .

But the rank of A being r , every minor of A of order $r+1$ vanishes i.e., $|A_0| = 0$, $|C_0| = 0$ and hence $|B_0| = 0$ showing that every $(r+1)$ rowed minor of B also vanishes. As such r' the rank of B cannot exceed r , the rank of A i.e.,

$$r' \leq r$$

Moreover A can be obtained from B by an elementary transformation of the same type and therefore a similar procedure will yield

$$r \leq r'$$

Thus $r' \leq r$ and $r \leq r'$ imply that $r = r'$ i.e., the addition to the elements of a row of a matrix, the products with any non-zero scalar λ of the corresponding elements of another row of the matrix does not alter its rank.

Conclusively the elementary operations do not alter the rank of a matrix.

COROLLARY 1. *Equivalent matrices have the same rank.*

If A and B are two equivalent matrices, then by the definition of equivalent matrices B can be obtained from A by a series of elementary operations and hence by the preceding theorem, the rank of A = the rank of B .

COROLLARY 2. *The rank of the transpose of a matrix is the same as that of the original matrix.* (Meerut, 1967)

If A is a matrix then its transpose A' is obtained from A by interchanging its rows and columns i.e., if

$$A = [a_{ij}] \text{ of order } m \times n,$$

then $A' = [a_{ji}]$ of order $n \times m$

Let r be the rank of A and r' that of A' .

If P be a square sub-matrix of A of order r such that $|P| \neq 0$, then P' will be a sub-matrix of A' of order r .

But $|P| = |P'|$ by the properties of determinants

$$|P'| = |P| \neq 0 \text{ cos } |P| \neq 0.$$

Thus by the same argument as in the above theorem

$$r' \geq r$$

Again if Q be a square sub-matrix of A of order $(r+1)$, then Q' will be a corresponding sub-matrix of A' of order $r+1$.

But the rank of A being r , $|Q| = 0$

and since $|Q| = |Q'|$

$\therefore |Q'| = |Q| = 0$ showing that every minor of A' of order $(r+1)$ is zero, therefore

$$r' \leq r$$

Hence $r' \geq r$ and $r' \leq r$ imply that $r' = r$ i.e., the rank of the transpose of a matrix is the same as that of the original matrix.

THEOREM 2. *Every non-singular matrix of order $m \times n$ and rank $r (> 0)$ can be reduced to one of the following normal forms :*

$$\begin{bmatrix} I_r & O \\ O & O \end{bmatrix}, \begin{bmatrix} I_r \\ O \end{bmatrix}, [I_r : O], [I_r]$$

by a series of elementary operations.

Suppose that $A = [a_{ij}]$ is the given non-zero matrix of order $m \times n$ and rank r . So A being non-zero will have at least one non-zero element say $a_{kl} = \lambda \neq 0$. If we interchange the k th row with the first row and the l th column with the first column, we find a matrix B with its leading element as λ i.e.,

$$B = \begin{bmatrix} \lambda & b_{12} & b_{13} & \dots & b_{1l} & \dots & b_{1n} \\ b_{21} & b_{22} & b_{23} & \dots & b_{2l} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ b_{m1} & b_{m2} & b_{m3} & \dots & b_{ml} & \dots & b_{mn} \end{bmatrix}$$

If we divide the elements of the first row of B by λ , we get another matrix C with its leading element unity, let it be given by

$$C = \begin{bmatrix} 1 & c_{12} & c_{13} & \dots & c_{1l} & \dots & c_{1n} \\ c_{21} & c_{22} & c_{23} & \dots & c_{2l} & \dots & c_{2n} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ c_{m1} & c_{m2} & c_{m3} & \dots & c_{ml} & \dots & c_{mn} \end{bmatrix}$$

If we now subtract the suitable multiples of the first column of C from the remaining columns and the suitable multiples of the first row of C from the remaining rows, then we get a matrix D with leading element unity and all the other elements of the first row and first column as zero i.e., of the form

$$D = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & d_{22} & d_{23} & \dots & d_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & d_{m2} & d_{m3} & \dots & d_{mn} \end{bmatrix}$$

which may be expressed as

$$D = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & & & & \\ \vdots & & A_1 & & \\ 0 & & & & \end{bmatrix} \text{ or } \begin{bmatrix} I & O \\ O & A_1 \end{bmatrix} \text{ by partitioning}$$

where A_1 is a matrix of the type $(m-1) \times (n-1)$.

If $A_1 \neq 0$, then proceeding as above, we can find an elementary matrix from A_1 with leading element unity and all other elements of

the first row and the first column as zero. Continuing this process t times, we shall get an elementary matrix of either of the forms.

$$(i) \begin{bmatrix} I_r & O \\ O & O \end{bmatrix} \text{ when } r < m, n$$

$$(ii) \begin{bmatrix} I_r & \\ O & \end{bmatrix} \text{ when } m=r, r < n$$

$$(iii) [I_r : O] \text{ when } r < m, n=r$$

$$(iv) [I_r] \text{ when } m=n=r.$$

THEOREM 3. If A is a matrix of rank r , then there exist non-singular matrices P and Q such that

$$PAQ = \begin{bmatrix} I_r & O \\ O & O \end{bmatrix}$$

Suppose that A is a matrix of order $m \times n$ and rank r . Then as given in the preceding theorem, if $r < m, n$, A can be reduced to the normal form $\begin{bmatrix} I_r & O \\ O & O \end{bmatrix}$ by elementary operations. But by theorem 1 of

§2.9, the elementary row (column) operation is effected by pre-multiplying (post-multiplying) the corresponding elementary matrix of suitable size, therefore corresponding to A the matrix of order $m \times m$ and rank r , there exist elementary matrices $P_1, P_2, \dots, P_s$ each of order m and $Q_1, Q_2, \dots, Q_t$ each of order n such that

$$P_s P_{s-1} \dots P_1 A Q_1 Q_2 \dots Q_t = \begin{bmatrix} I_r & O \\ O & O \end{bmatrix}$$

Here each of the elementary matrices $P_1, P_2, \dots$ etc, and $Q_1, Q_2, \dots$ etc, being non-singular, their products are also non-singular and if we take

$$P = P_s P_{s-1} \dots P_1, Q = Q_1 Q_2 \dots Q_t$$

$$\text{then } PAQ = \begin{bmatrix} I_r & O \\ O & O \end{bmatrix}$$

where P is of order $m \times m$ and Q i.e., of order $n \times n$.

THEOREM 4. If A is a matrix of order $m \times n$ and rank r , then there exists a non-singular matrix

$$(i) P \text{ such that } PA = \begin{bmatrix} G \\ O \end{bmatrix}$$

where G is a matrix of order $r \times n$ and rank r and O is the null matrix of order $(m-r) \times n$.

$$(ii) Q \text{ such that } AQ = [HO]$$

where H is a matrix of order $m \times r$ and rank r and O is the null-matrix of order $m \times (n-r)$.

By the preceding theorem there exist non-singular matrices P, Q such that $PAQ = \begin{bmatrix} I_r & O \\ O & O \end{bmatrix}$

(i) If $Q = Q_1 Q_2 \dots Q_t$, where $Q_1, Q_2, \dots, Q_t$ are elementary matrices, then

$$PA Q_1 Q_2 \dots Q_t = \begin{bmatrix} I_r & O \\ O & O \end{bmatrix} \quad \dots (1)$$

Post-multiplying the left hand side of (1) in succession by the elementary matrices $Q_t^{-1}, Q_{t-1}^{-1}, \dots, Q_2^{-1}, Q_1^{-1}$, and effecting the corresponding column operation in the right hand side of (1) while no column operation would affect the last $(m-r)$ zero rows, we shall find a relation of the form

$$PA = \begin{bmatrix} G \\ O \end{bmatrix}$$

But we know that the elementary operations, do not alter the rank and hence the rank of PA and so of G will be r , which is the rank of A .

(ii) Take $P = P_1 P_2 \dots P_s$; where $P_1, P_2, \dots, P_s$ are elementary matrices, then we have as in case (i),

$$P_1 P_2 P_3 \dots P_s A Q = \begin{bmatrix} I_r & O \\ O & O \end{bmatrix} \quad \dots (2)$$

Pre-multiplying the left hand side of (2) in succession by the elementary matrices $P_1^{-1}, P_2^{-1}, \dots, P_s^{-1}$ and effecting the corresponding row operation in the right hand side of (2) while no row operation would affect the last $(n-r)$ zero columns, we shall find a relation of the form

$$AQ = [H O]$$

where the rank of AQ and also of H will be r , since the elementary operations do not alter the rank and the rank of A is r .

THEOREM 5. The rank of a product of two matrices A and B whose ranks are r_1, r_2 respectively cannot exceed the rank of either matrix i.e., if r be the rank of product AB , then

$$r \leq r_1 \text{ and } r \leq r_2. \quad (\text{Agra, 1968})$$

Let A be the matrix of order $m \times n$ and B the matrix of order $n \times p$, so that the product AB is conformable and has order $m \times p$.

By the preceding theorem, A being the matrix of order $m \times n$ and rank r_1 , there exists a non-singular matrix P which is the product of elementary matrices, such that

$$PA = \begin{bmatrix} G \\ O \end{bmatrix} \quad \dots (1)$$

where G is a matrix of order $r_1 \times n$ and rank r_1 and O is the null matrix of order $(m-r_1) \times n$.

Post-multiplying either side of (1) by B , we get

$$PAB = \begin{bmatrix} G \\ O \end{bmatrix} B \quad \dots (2)$$

But P being the product of elementary matrices, we have

$$\text{rank of } (PAB) = \text{rank of } (AB) = r \quad \dots (3)$$

For, the elementary operations do not alter the rank.

From (2) and (3) it follows that

$$\text{rank of } \begin{bmatrix} G \\ O \end{bmatrix} B = r \quad \dots (4)$$

But $\begin{bmatrix} G \\ O \end{bmatrix} B$ has at most r_1 non-zero rows which arise by the multiplication of r_1 non-zero rows of G with the columns of B , therefore
 $\text{rank of } \begin{bmatrix} G \\ O \end{bmatrix} B \leq r_1$... (1)

Thus from (4) and (5) it is apparent that
 $r \leq r_1$ i.e., the rank of $(AB) \leq$ the rank of the prefactor A ... (2)
 Again $r = \text{rank of } (AB)$

$= \text{rank of } (AB)'$ by corollary 2 of Theorem 1.
 $= \text{rank of } (B'A') \because (AB)' = (B'A')$ by reversal law of transpose

$\leq \text{rank of the prefactor } B'$ by (6)

$\leq \text{rank of } (B)$

$\leq r_2$

i.e., the rank of $(AB) \leq$ the rank of the post-factor B .

THEOREM 6. The rank of a matrix does not alter by pre-multiplication (or post-multiplication) with any non-singular matrix.

Let A be any matrix and P a non-singular matrix such that the product PA is conformable and let

$$B = PA \quad \dots (1)$$

Then by the preceding theorem,

$$\text{rank of } (B) = \text{rank of } (PA) \leq \text{rank of } (A) \quad \dots (2)$$

(1) may also be written as

$$A = P^{-1}B \quad \dots (3)$$

$\therefore \text{rank of } (A) = \text{rank of } (P^{-1}B) \leq \text{rank of } (B)$

From (2) and (3) it follows that

$$\text{rank of } (A) = \text{rank of } (B).$$

Problem 28. Find the rank of the matrix $\begin{bmatrix} 1 & 3 & 4 & 3 \\ 3 & 9 & 12 & 3 \\ 1 & 3 & 4 & 1 \end{bmatrix}$ (Agra, 1976)

Denoting the given matrix by A we have

$$A = \begin{bmatrix} 1 & 3 & 4 & 3 \\ 3 & 9 & 12 & 3 \\ 1 & 3 & 4 & 1 \end{bmatrix}$$

The matrix A is of the order 3×4 and so it can have minors of order 1, 2 and 3. The minors of order 3 are

$$\begin{vmatrix} 1 & 3 & 4 \\ 3 & 9 & 12 \\ 1 & 3 & 4 \end{vmatrix}, \begin{vmatrix} 1 & 3 & 3 \\ 3 & 9 & 3 \\ 1 & 3 & 1 \end{vmatrix}, \begin{vmatrix} 1 & 4 & 3 \\ 3 & 12 & 3 \\ 1 & 4 & 1 \end{vmatrix}, \begin{vmatrix} 3 & 4 & 3 \\ 9 & 12 & 3 \\ 3 & 4 & 1 \end{vmatrix}$$

i.e., $12 \begin{vmatrix} 1 & 1 & 1 \\ 3 & 3 & 3 \\ 1 & 1 & 1 \end{vmatrix}, 3 \begin{vmatrix} 1 & 1 & 3 \\ 3 & 3 & 3 \\ 1 & 1 & 1 \end{vmatrix}, 4 \begin{vmatrix} 1 & 1 & 3 \\ 3 & 3 & 3 \\ 1 & 1 & 1 \end{vmatrix}, 12 \begin{vmatrix} 1 & 1 & 3 \\ 3 & 3 & 3 \\ 1 & 1 & 1 \end{vmatrix}$

all of which vanish as they consist of at least two columns identical

The minors of order 2 are

$$\begin{vmatrix} 1 & 3 \\ 3 & 9 \end{vmatrix}, \begin{vmatrix} 1 & 4 \\ 3 & 12 \end{vmatrix}, \begin{vmatrix} 1 & 3 \\ 3 & 3 \end{vmatrix}, \begin{vmatrix} 3 & 4 \\ 9 & 12 \end{vmatrix}, \begin{vmatrix} 3 & 3 \\ 9 & 3 \end{vmatrix}, \begin{vmatrix} 4 & 3 \\ 12 & 3 \end{vmatrix},$$

$$\begin{vmatrix} 1 & 3 \\ 1 & 3 \end{vmatrix}, \begin{vmatrix} 1 & 4 \\ 1 & 4 \end{vmatrix}, \begin{vmatrix} 1 & 3 \\ 1 & 1 \end{vmatrix}, \begin{vmatrix} 3 & 4 \\ 3 & 4 \end{vmatrix}, \begin{vmatrix} 3 & 3 \\ 3 & 1 \end{vmatrix}, \begin{vmatrix} 4 & 3 \\ 4 & 1 \end{vmatrix},$$

$$\begin{vmatrix} 3 & 9 \\ 1 & 3 \end{vmatrix}, \begin{vmatrix} 3 & 12 \\ 1 & 4 \end{vmatrix}, \begin{vmatrix} 3 & 3 \\ 1 & 1 \end{vmatrix}, \begin{vmatrix} 9 & 12 \\ 3 & 4 \end{vmatrix}, \begin{vmatrix} 9 & 3 \\ 3 & 1 \end{vmatrix}, \begin{vmatrix} 12 & 3 \\ 4 & 1 \end{vmatrix}$$

of which do not vanish e.g.

$$\begin{vmatrix} 1 & 3 \\ 3 & 3 \end{vmatrix} = 3 - 9 = -6 \neq 0.$$

Hence the rank of A is 2

Problem 29. Find the rank of the matrix

$$A = \begin{bmatrix} 1 & 3 & 4 & 3 \\ 3 & 9 & 12 & 9 \\ -1 & -3 & -4 & -3 \end{bmatrix}$$

(Gorakhpur, 1965)

We know that the elementary operations do not alter the rank or in other words the equivalent matrices have the same rank, thus since

$$A \sim \begin{bmatrix} 1 & 3 & 4 & 3 \\ 3 & 9 & 12 & 9 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } R_{31} (1)$$

$$\sim \begin{bmatrix} 1 & 3 & 4 & 3 \\ 1 & 3 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } R_2 \left(\frac{1}{3}\right)$$

$$\sim \begin{bmatrix} 1 & 3 & 4 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } R_{21} (-1)$$

therefore the rank of A is clearly 1, for, the minors of equivalent matrix of orders 3 and 2, vanish.

Problem 30. Reduce the matrix A to the normal form and hence find its rank, where

$$A = \begin{bmatrix} 1 & 1 & 2 & 3 \\ 1 & 3 & 0 & 3 \\ 1 & -2 & -3 & -3 \\ 1 & 1 & 2 & 3 \end{bmatrix}$$

(Delhi, 1960)

$$\text{We have } A \sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 2 & -2 & 0 \\ 1 & -3 & -5 & -6 \\ 1 & 0 & 0 & 0 \end{bmatrix} \text{ by } C_{21} (-1), C_{31} (-2) \text{ and } C_{41} (-3)$$

$$A \sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & -2 & 0 \\ 0 & -3 & -5 & -6 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } R_{21} (-1), R_{31} (-1) \text{ and } R_{41} (-1)$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 3 & 5 & 6 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } R_2 \left(\frac{1}{2}\right) \text{ and } R_3(-1)$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3 & 8 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } C_{32}(1) \text{ and } C_4\left(\frac{1}{8}\right)$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } C_{24}(-3) \text{ and } C_3\left(\frac{1}{8}\right)$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ by } C_{43}(-1)$$

$$\sim \begin{bmatrix} I_3 & O \\ O & O \end{bmatrix} \text{ which is the required normal form and its rank is 3}$$

Problem 31. Find the non-singular matrices **P** and **Q** such that **PAQ** is in the normal form, where

$$A = \begin{bmatrix} 3 & 2 & -1 & 5 \\ 5 & 1 & 4 & -2 \\ 1 & -4 & 11 & -19 \end{bmatrix}$$

The matrix **A** being of order 4×3 , we may write,

$$\begin{bmatrix} 3 & 2 & -1 & 5 \\ 5 & 1 & 4 & -2 \\ 1 & -4 & 11 & -19 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} A \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\text{or } \begin{bmatrix} 1 & -4 & 11 & -19 \\ 5 & 1 & 4 & -2 \\ 3 & 2 & -1 & 5 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} A \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \text{ by } R_{13}$$

$$\text{or } \begin{bmatrix} 1 & -4 & 11 & -19 \\ 0 & 21 & -51 & -93 \\ 0 & 14 & -34 & 62 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & -5 & -3 \\ 1 & 0 & -3 & 1 \end{bmatrix} A \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

by $R_{21}(-5)$ and $R_{31}(-3)$

$$\text{or } \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 21 & -51 & 93 \\ 0 & 14 & -34 & 62 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & -5 & -3 \\ 1 & 0 & -3 & 1 \end{bmatrix} A \begin{bmatrix} 1 & 4 & -11 & 19 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

by $C_{21}(4)$, $(3)(-11)$ and $C_{41}(19)$

$$\text{or } \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 3 & 3 & 3 \\ 0 & 2 & 2 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & -5 & -3 \\ 1 & 0 & -3 & 1 \end{bmatrix} A \begin{bmatrix} 1 & \frac{4}{3} & \frac{11}{3} & \frac{19}{3} \\ 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & -\frac{1}{3} & 0 \\ 0 & 0 & 0 & \frac{1}{3} \end{bmatrix}$$

by $C_2\left(\frac{1}{3}\right)$, $C_3\left(-\frac{1}{3}\right)$ and $C_4\left(\frac{1}{3}\right)$

$$\text{or } \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & \frac{1}{3} & -\frac{5}{3} \\ \frac{1}{3} & 0 & -\frac{2}{3} \end{bmatrix} A \begin{bmatrix} 1 & \frac{4}{3} & \frac{11}{3} & \frac{19}{3} \\ 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & -\frac{1}{3} & 0 \\ 0 & 0 & 0 & \frac{1}{3} \end{bmatrix}$$

by $R_2\left(\frac{1}{3}\right)$ and $R_3\left(\frac{1}{3}\right)$

$$\text{or } \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & \frac{1}{3} & -\frac{5}{3} \\ \frac{1}{3} & 0 & -\frac{2}{3} \end{bmatrix} A \begin{bmatrix} 1 & \frac{4}{3} & \frac{11}{3} & \frac{19}{3} \\ 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & -\frac{1}{3} & 0 \\ 0 & 0 & 0 & \frac{1}{3} \end{bmatrix}$$

by $C_{32}(-1)$ and $C_{42}(-1)$

$$\text{or } \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & \frac{1}{3} & -\frac{5}{3} \\ \frac{1}{3} & -\frac{1}{3} & \frac{1}{3} \end{bmatrix} A \begin{bmatrix} 1 & \frac{4}{3} & \frac{11}{3} & \frac{19}{3} \\ 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & -\frac{1}{3} & 0 \\ 0 & 0 & 0 & \frac{1}{3} \end{bmatrix}$$

by $R_{32}(-1)$

which gives the required normal form i.e.,

$$\begin{bmatrix} I_2 & O \\ O & O \end{bmatrix} = PAQ$$

$$\text{where } P = \begin{bmatrix} 0 & 0 & 1 \\ 0 & \frac{1}{3} & -\frac{5}{3} \\ \frac{1}{3} & -\frac{1}{3} & \frac{1}{3} \end{bmatrix} \text{ and } Q = \begin{bmatrix} 1 & \frac{4}{3} & \frac{11}{3} & \frac{19}{3} \\ 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & -\frac{1}{3} & 0 \\ 0 & 0 & 0 & \frac{1}{3} \end{bmatrix}$$

Problem 32. Find the rank of the product matrix **AB** when

$$A = \begin{bmatrix} 1 & 1 & -1 \\ 2 & -3 & 4 \\ 3 & -2 & 3 \end{bmatrix} \text{ and } B = \begin{bmatrix} -1 & -2 & -1 \\ 6 & 12 & 6 \\ 5 & 10 & 5 \end{bmatrix}$$

$$\text{We have } AB = \begin{bmatrix} 1 & 1 & -1 \\ 2 & -3 & 4 \\ 3 & -2 & 3 \end{bmatrix} \begin{bmatrix} -1 & -2 & -1 \\ 6 & 12 & 6 \\ 5 & 10 & 5 \end{bmatrix} \\ = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Hence rank of $(AB) = 0$, since the rank of a null-matrix is zero.

12. SOLUTIONS OF LINEAR EQUATIONS

In this section we shall make an attempt to apply the concepts and consequences of matrices discussed in the previous sections, to find the solution of any system of linear equations, homogeneous or non-homogeneous.

It should be clearly noted that a system of n linear equations in n unknown variables has not always a solution, e.g. the system of equations

$$(i) \quad \begin{cases} 2x+3y=7 \\ 4x-5y=3 \end{cases} \text{ has a single solution namely } x=2, y=1$$

$$(ii) \quad \begin{cases} 2x-3y=1 \\ -4x+6y=2 \end{cases} \text{ has no solution}$$

$$\text{and (iii)} \quad \begin{cases} 3x+2y=5 \\ 6x+4y=10 \end{cases} \text{ has an infinite number of solutions}$$

Homogeneous linear equations

Let there be a set of m homogeneous linear equations in n variables $x_1, x_2, x_3, \dots, x_n$,

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = 0$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = 0$$

$$\dots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = 0$$

In contracted form this set of equations may be expressed as

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

Writing,

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}, \quad X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad O = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

where A, X, O are the matrices of orders $m \times n, n \times 1$ and $m \times 1$ respectively, the given set of equations is equivalent to a single matrix equation

$$AX = O \quad \dots (1)$$

Let us assume that r is the rank of the matrix A of order $m \times n$. As regards the solution of the set of homogeneous linear equations the following results can be summarised:

I. A system of homogeneous linear equations always have one or more solutions.

Since r cannot be greater than n , therefore, either $r = n$ or $r < n$.

In the first case the equation (i) will have no linearly independent solutions, for in that case trivial (zero) solution is the only solution while in the second case there will be $(n-r)$ independent solutions and therefore the equation (1) will have more than one solution.

II. The number of linearly independent solutions of $AX = O$ is $(n-r)$ i.e. if we assign arbitrary values to $(n-r)$ of the variables, then the values of the others can be uniquely determined.

Since the rank of A is r , it has r linearly independent columns. There is no loss of generality if we suppose that the first r columns on the left are linearly independent for, it amounts only to renaming the variable components of X .

Taking $C_1, C_2, \dots, C_r, \dots, C_n$ as column vectors of A , we can write

$$A = [C_1, C_2, \dots, C_r, \dots, C_n] \text{ of order } 1 \times n$$

so that the equation $AX = O$ is equivalent to

$$x_1 C_1 + x_2 C_2 + \dots + x_r C_r + \dots + x_n C_n = O \quad \dots (2)$$

Each of the column vectors $C_{r+1}, C_{r+2}, \dots, C_n$ being linear combination of $C_1, C_2, \dots, C_r$, we have the relations of the form

$$\left. \begin{aligned} C_{r+1} &= k_{11} C_1 + k_{12} C_2 + \dots + k_{1r} C_r \\ C_{r+2} &= k_{21} C_1 + k_{22} C_2 + \dots + k_{2r} C_r \\ &\dots \\ C_n &= k_{l1} C_1 + k_{l2} C_2 + \dots + k_{lr} C_r \end{aligned} \right\} \quad \dots (3)$$

$l = n - r$

Where

Now the relations (3) can be rewritten as

$$\left. \begin{aligned} k_{11} C_1 + k_{12} C_2 + \dots + k_{1r} C_r - 1 \cdot C_{r+1} + 0 \cdot C_{r+2} + \dots + 0 \cdot C_n &= O \\ k_{21} C_1 + k_{22} C_2 + \dots + k_{2r} C_r + 0 \cdot C_{r+1} - 1 \cdot C_{r+2} + \dots + 0 \cdot C_n &= O \\ &\dots \\ k_{l1} C_1 + k_{l2} C_2 + \dots + k_{lr} C_r + 0 \cdot C_{r+1} + 0 \cdot C_{r+2} + \dots - 1 \cdot C_n &= O \end{aligned} \right\} \quad \dots (4)$$

Comparison of (2) and (4) yields

$$X_1 = \begin{bmatrix} k_{11} \\ k_{12} \\ \vdots \\ k_{1r} \\ -1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad X_2 = \begin{bmatrix} k_{21} \\ k_{22} \\ \vdots \\ k_{2r} \\ 0 \\ -1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad X_{n-r} = \begin{bmatrix} k_{l1} \\ k_{l2} \\ \vdots \\ k_{lr} \\ 0 \\ 0 \\ \vdots \\ -1 \end{bmatrix} \quad \dots (5)$$

which are $(n-r)$ solutions of $AX = O$.

These $(n-r)$ solutions constitute a linearly independent set of solutions, for if we have

$$k_1 X_1 + k_2 X_2 + \dots + k_{n-r} X_{n-r} = O \quad \dots (6)$$

Then a comparison of $(r+1)$ th, $(r+2)$ th, $\dots$, n th components on either side of (6) will give

$$k_1 = 0, k_2 = 0, \dots, k_{n-r} = 0$$

In order to show that every solution of $AX = O$ is some suitable linear combination of these $(n-r)$ solutions $X_1, X_2, \dots, X_{n-r}$, let us construct a vector

$$X + x_{r+1} X_1 + x_{r+2} X_2 + \dots + x_n X_{n-r} \quad \dots (7)$$

which is also a solution of $AX = O$, as it is the linear combination of its solutions. Obviously the last $(n-r)$ components of (7) are zero. Let its first r components be $y_1, y_2, \dots, y_r$. Then the vector

$$y_1, y_2, \dots, y_r, 0, 0, \dots, 0,$$

is a solution of $AX=O$ and hence from (2), we have

$$y_1 C_1 + y_2 C_2 + \dots + y_r C_r = O$$

But the set of vectors $C_1, C_2, \dots, C_r$ is linearly independent, therefore

$$y_1 = 0, y_2 = 0, \dots, y_r = 0$$

which shows that (7) is a zero vector and thus

$$X = -x_{r+1} X_1 - x_{r+2} X_2 - \dots - x_n X_{n-r}$$

Hence every solution X is a linear combination of the $n-r$ linearly independent solutions $X_1, X_2, \dots, X_{n-r}$

Note 1. In case $n=r$, there is only one solution $x_1 = x_2 = \dots = x_n = 0$ and when $n > r$, there is a unique solution other than $x_1 = x_2 = \dots = x_n = 0$, provided that the rank of the matrix of the coefficients of the $(n-r)$ arbitrary chosen values is r .

Note 2. If $X_1, X_2, X_3, \dots, X_{n-r}$ be a set of $(n-r)$ arbitrary independent solutions of the equation $AX=O$, then its general solution is

$$X = k_1 X_1 + k_2 X_2 + \dots + k_{n-r} X_{n-r}$$

where $k_1, k_2, \dots, k_{n-r}$ is a set of $(n-r)$ arbitrary values.

III. If the number of equations is less than the number of variables, then the solution is always other than

$$x_1 = x_2 = \dots = x_n = 0$$

IV. If the number of equations is equal to the number of variables, necessary and sufficient condition for solutions other than $x_1 = x_2 = \dots = x_n = 0$ is that the determinant of the coefficients must be zero.

Non-Homogeneous Linear Equations

Let there be m non-homogeneous linear equations in variables $x_1, x_2, \dots, x_n$, such as

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2$$

$$\dots \dots \dots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

The system of given equations is equivalent to a matrix equation

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

Denoting by

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}, \quad X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \text{ and } B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

we have the matrix equation

$$AX=B$$

This equation is said to be **consistent** i.e. the equation possesses a solution if the matrices A and $[A, B]$ are of the same rank, where $[A, B]$ denotes the matrix

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \dots & \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{bmatrix}$$

Here the matrix A is called **Coefficient Matrix** and the matrix $[A, B]$ is called **Augmented Matrix**.

THEOREM. The system of non-homogeneous equations $AX=B$ possesses a solution if

$$\text{Rank of } (A) = \text{Rank of } [A, B]$$

i.e., if r and ρ be the ranks of the matrices A and $[A, B]$ respectively, then the given equations are consistent when $r=\rho$ and inconsistent when $r < \rho$.

Supposing that $C_1, C_2, \dots, C_n$ are the column vectors of the matrix A , the equation $AX=B$

$$\dots (1)$$

is equivalent to

$$[C_1, C_2, \dots, C_n] \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = B$$

$$\text{i.e., } x_1 C_1 + x_2 C_2 + \dots + x_n C_n = B \quad \dots (2)$$

The rank of the matrix A being r , it has r linearly independent columns and so there is no loss of generality if we assume that the first r columns form a linearly independent set such that each of the remaining $n-r$ columns is a linear combination of these r columns.

Case I. If the given set of equations is consistent, then there must exist n numbers say $k_1, k_2, \dots, k_n$ such that

$$k_1 C_1 + k_2 C_2 + \dots + k_n C_n = B \quad \dots (3)$$

But each of the $(n-r)$ columns $C_{r+1}, C_{r+2}, \dots, C_n$ being a linear combination of first r columns $C_1, C_2, \dots, C_r$, it follows from (3) that B is also a linear combination of $C_1, C_2, \dots, C_r$. As such the greatest number of linearly independent columns of $[A, B]$ is also r showing that r is also the rank of $[A, B]$. Conclusively the system of equations $AX=B$ is consistent if $\text{rank of } (A) = \text{rank of } [A, B]$.

Conversely if the ranks of A and $[A, B]$ are the same say r , then the greatest number of linearly independent columns of $[A, B]$ will be r . And since the first r columns $C_1, C_2, \dots, C_r$ of $[A, B]$ form a linearly independent set, therefore the column B is expressible as a linear combination of the columns $C_1, C_2, \dots, C_r$. As such there exist r scalars $k_1, k_2, \dots, k_r$ such that

$$k_1 C_1 + k_2 C_2 + \dots + k_r C_r = B$$

$$\text{i.e., } k_1 C_1 + k_2 C_2 + \dots + k_r C_r + 0 \cdot C_{r+1} + 0 \cdot C_{r+2} + \dots + 0 \cdot C_n = B \quad \dots (4)$$

Comparison of (2) and (4) leads to

$$x_1 = k_1, x_2 = k_2, \dots, x_r = k_r, x_{r+1} = 0, x_{r+2} = 0, \dots, x_n = 0$$

which constitute a solution of $AX=B$.

Hence the given system of equations is consistent if and only if the ranks of the coefficient matrix and the augmented matrix are the same.

Case II. If rank of $(A) < \text{rank of } ([A, B])$, then the equations $AX=B$ are inconsistent as is obvious from the Case I and hence the given system of equations has no solution.

Note 3. The most general solution of $AX=B$ is

$$X_0 + k_1 X_1 + k_2 X_2 + \dots + k_{n-r} X_{n-r} = X_0 + Y$$

where $X_1, X_2, \dots, X_{n-r}$ is a set of $(n-r)$ solutions of $AX=O$ and X_0 is any fixed solution of $AX=B$, also $k_1, k_2, \dots, k_{n-r}$ is any set of $(n-r)$ constants.

Here
$$A(X_0 + Y) = AX_0 + AY \\ = B + O = B$$

So that $X_0 + Y$ is a solution of the given equation i.e., $AX=B$.

Conversely if Z be any solution of the given equations, then

$$A(Z - X_0) = AZ - AX_0 \\ = B - B = O$$

showing that $Z - X_0$ is a solution of $AX=O$.

If we write $Z = X_0 + (Z - X_0)$

then it follows that every solution Z of $AX=B$ may be expressed as the sum of a fixed solution X_0 thereof and some solution $Z - X_0$ of the auxiliary equation $AX=O$.

Note 4. If A be a n -rowed non-singular matrix, X be an $n \times 1$ matrix, and B be an $n \times 1$ matrix, then the system of equations $AX=B$ has a unique solution.

A being n -rowed non-singular matrix and B being a matrix of order $n \times 1$, the ranks of both A and $[A, B]$ will be n and so the system of equations $AX=B$ is consistent i.e., possesses a solution.

If we pre-multiply either side of $AX=B$ by A^{-1} , we have

$$A^{-1}AX = A^{-1}B$$

or
$$IX = A^{-1}B$$

i.e.
$$X = A^{-1}B$$

is a solution of
$$AX=B$$

In order to show that this solution is unique, let us assume if possible that X_1, X_2 are two solutions of $AX=B$, then we have

$$AX_1=B \text{ and } AX_2=B$$

which follow that
$$AX_1=AX_2$$

or
$$A^{-1}AX_1=A^{-1}AX_2$$

or
$$IX_1=IX_2$$

i.e.,
$$X_1=X_2$$

which shows that the solution is unique.

13. CRAMER'S RULE

If a matrix equation is of the form

$$AX=B \quad \dots (1)$$

where A is a non-singular square matrix i.e., $|A| \neq 0$, then pre-multiplying either side of (1) by A^{-1} , we have

$$A^{-1}AX = A^{-1}B$$

$$IX = A^{-1}B$$

$$X = A^{-1}B$$

A^{-1} being unique, these are the only solutions.

As a particular case if we consider a set of three non-homogeneous linear equations

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2$$

$$a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3$$

the system is equivalent to the matrix equation

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

Denoting by $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$, $X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$, $B = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$

we have

$$AX=B.$$

Pre-multiplying by A^{-1} , we get as above

$$X = A^{-1}B \quad \dots (2)$$

But we know that

$$A^{-1} = \frac{1}{|A|} \text{adj } A$$

Here

$$|A| = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

and

$$\text{adj } A = \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{bmatrix}$$

where A_{11} etc. are the cofactors of the elements a_{11} etc. in the determinant $|A|$.

Here (2) may be expressed as

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \frac{1}{|A|} \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

$$= \frac{1}{|A|} \begin{bmatrix} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} & -\begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix} & \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix} \\ -\begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} & -\begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} \\ \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} & -\begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

$$= \frac{1}{|A|} \begin{bmatrix} b_1 \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - b_2 \begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix} + b_3 \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix} \\ -b_1 \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + b_2 \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} - b_3 \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} \\ b_1 \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} - b_2 \begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix} + b_3 \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \end{bmatrix}$$

$$\text{or } \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 & a_{12} & a_{13} \\ b_2 & a_{22} & a_{23} \\ b_3 & a_{32} & a_{33} \end{bmatrix} \div \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$\begin{bmatrix} a_{11} & b_1 & a_{13} \\ a_{21} & b_2 & a_{23} \\ a_{31} & b_3 & a_{33} \end{bmatrix}$$

$$\begin{bmatrix} a_{11} & a_{12} & b_1 \\ a_{21} & a_{22} & b_2 \\ a_{31} & a_{32} & b_3 \end{bmatrix}$$

which is equivalent to

$$x_1 = \frac{\begin{vmatrix} b_1 & a_{12} & a_{13} \\ b_2 & a_{22} & a_{23} \\ b_3 & a_{32} & a_{33} \end{vmatrix}}{\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}}, x_2 = \frac{\begin{vmatrix} a_{11} & b_1 & a_{13} \\ a_{21} & b_2 & a_{23} \\ a_{31} & b_3 & a_{33} \end{vmatrix}}{\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}}, x_3 = \frac{\begin{vmatrix} a_{11} & a_{12} & b_1 \\ a_{21} & a_{22} & b_2 \\ a_{31} & a_{32} & b_3 \end{vmatrix}}{\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}}$$

This result may be extended to the general case and thus gives the rule given by Cramer.

'For a given set of n non-homogeneous linear equations in n variables $x_1, x_2, \dots, x_n$ each x_i ($i=1, 2, \dots, n$) is a quotient, the denominator of which is the determinant $|A|$ of the coefficient matrix A and the numerator of which is a determinant obtained from $|A|$ by substituting the column vector $\begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$ for the i th column of $|A|$.

Problem 33. Solve : $2x+3y-z=0$

$$x-y-2z=0$$

$$4x+y-5z=0$$

The given system of equations is equivalent to the matrix equation

$$\begin{bmatrix} 2 & 3 & -1 \\ 1 & -1 & -2 \\ 4 & 1 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \mathbf{O} \text{ where } \mathbf{O} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\text{or } \begin{bmatrix} 0 & 5 & 3 \\ 1 & -1 & -2 \\ 0 & 5 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \mathbf{O} \text{ by } R_{12}(-2) \text{ and } R_{32}(-4)$$

$$\text{or } \begin{bmatrix} 0 & 5 & 3 \\ 1 & -1 & -2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \mathbf{O} \text{ by } R_{31}(-1)$$

which is equivalent to

$$5y+3z=0$$

$$x-y-2z=0$$

giving

$$x = \frac{7}{5}z, y = -\frac{3}{5}z$$

which give the solution for arbitrary values of z .

The general solution is given by

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{7}{5}z \\ -\frac{3}{5}z \\ z \end{bmatrix} = z \begin{bmatrix} \frac{7}{5} \\ -\frac{3}{5} \\ 1 \end{bmatrix} = k \begin{bmatrix} 7 \\ -3 \\ 5 \end{bmatrix} \text{ where } z=5k \text{ is an arbitrary parameter.}$$

Problem 34. Solve $4x+2y+z+3u=0$

$$6x+3y+4z+7u=0$$

$$2x+y+u=0$$

(Meerut, 1968)

The given system of equations is equivalent to the matrix equation

$$\begin{bmatrix} 4 & 2 & 1 & 3 \\ 6 & 3 & 4 & 7 \\ 2 & 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ u \end{bmatrix} = \mathbf{O}, \text{ where } \mathbf{O} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\text{or } \begin{bmatrix} 4 & 2 & 1 & 3 \\ 0 & 0 & \frac{5}{2} & \frac{5}{2} \\ 0 & 0 & -\frac{1}{2} & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ u \end{bmatrix} = \mathbf{O} \text{ by } R_{21}(-\frac{2}{5}) \text{ and } R_{31}(-\frac{1}{2})$$

$$\text{or } \begin{bmatrix} 4 & 2 & 1 & 3 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ u \end{bmatrix} = \mathbf{O} \text{ by } R_{32}(-1) \text{ and } R_2(\frac{2}{5})$$

which is equivalent to

$$4x+2y+z+3u=0$$

$$z+u=0$$

$$\text{giving, } z = -u, x = -\frac{y+u}{2}$$

The rank of $\begin{bmatrix} 4 & 2 & 1 & 3 \\ 6 & 3 & 4 & 7 \\ 2 & 1 & 0 & 1 \end{bmatrix}$ is 2 and number of unknown variables

is 4.

$\therefore$ Number of independent solutions $= 4 - 2 = 2$

If we choose $u = k_1$ and $x = k_2$

then, $y = -2k_2 - k_1$ and $z = -k_1$

which give an infinite number of non-trivial solutions.

Problem 35. Solve $x + 2y + 3z = 0$

$$3x + 4y + 4z = 0$$

$$7x + 10y + 12z = 0$$

The given system of equations is equivalent to the matrix equation

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 4 & 4 \\ 7 & 10 & 12 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \mathbf{O}, \text{ where } \mathbf{O} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

or $\begin{bmatrix} 1 & 2 & 3 \\ 0 & -2 & -5 \\ 0 & -4 & -9 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \mathbf{O}$ by $R_{21}(-2)$ and $R_{31}(-7)$

or $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & \frac{5}{2} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \mathbf{O}$ by $R_2(-\frac{1}{2})$ and then $R_{32}(4)$

which is equivalent to

$$x + 2y + 3z = 0$$

$$y + \frac{5}{2}z = 0, z = 0$$

giving $x = 0, y = 0, z = 0$.

Moreover the rank of the coefficient matrix $\begin{bmatrix} 1 & 2 & 3 \\ 3 & 4 & 4 \\ 7 & 10 & 12 \end{bmatrix}$ is 3 which

is the number of unknown variables and hence the system has only zero solution.

Problem 36. State the conditions under which a system of non-homogeneous linear equations will have

(i) no solution, (ii) a unique solution, and (iii) an infinity of solutions. (Meerut, 1967)

Let the system of non-homogeneous linear equations be equivalent to the matrix equation $\mathbf{AX} = \mathbf{B}$

where $\mathbf{A}$ is of order $m \times n$, $\mathbf{X}$ be of order $n \times 1$ and $\mathbf{B}$ be of order $m \times 1$

(i) The equation $\mathbf{AX} = \mathbf{B}$ has no solution if

$$\text{rank of } (\mathbf{A}) \neq \text{rank of } ([\mathbf{A}, \mathbf{B}])$$

(ii) The equation $\mathbf{AX} = \mathbf{B}$ has a unique solution if

$$\text{rank of } (\mathbf{A}) = \text{rank of } ([\mathbf{A}, \mathbf{B}]) = \text{number of unknown variables}$$

In particular if $\mathbf{A}$ is a square matrix, then $\mathbf{AX} = \mathbf{B}$ will have a unique solution if $|\mathbf{A}| \neq 0$.

(iii) The equations $\mathbf{AX} = \mathbf{B}$ will have an infinite number of solutions if $\text{rank of } (\mathbf{A}) = \text{rank of } ([\mathbf{A}, \mathbf{B}]) < \text{number of unknown variables}$.

Problem 37. Solve the equations by using matrix methods :

$$x + y + z = 6$$

$$x - y + z = 2$$

$$2x + y - z = 1$$

(Agra, 1970)

The given system of equations is equivalent to the matrix equation

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \\ 2 & 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ 2 \\ 1 \end{bmatrix}$$

or $\begin{bmatrix} 1 & 1 & 1 \\ 0 & -2 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ -4 \\ 3 \end{bmatrix}$ by $R_{21}(-1)$ and $R_{32}(1)$

which is equivalent to

$$x + y + z = 6$$

$$-2y = -4$$

$$3x = 3$$

giving $x = 1, y = 2, z = 3$.

Problem 38. Show that the equations

$$x + 2y - z = 3$$

$$3x - y + 2z = 1$$

$$2x - 2y + 3z = 2$$

$$x - y + z = -1$$

are consistent and solve them.

The given system of equations is equivalent to the matrix equation

$$\begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 2 & -2 & 3 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \\ -1 \end{bmatrix} \quad \dots (1)$$

Denoting the coefficient matrix by $\mathbf{A}$ and augmented matrix by $\mathbf{C}$ we have

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -1 & 2 \\ 2 & -2 & 3 \\ 1 & -1 & 1 \end{bmatrix}, \mathbf{C} = \begin{bmatrix} 1 & 2 & -1 & 3 \\ 3 & -1 & 2 & 1 \\ 2 & -2 & 3 & 2 \\ 1 & -1 & 1 & -1 \end{bmatrix}$$

Here fourth order minor of $\mathbf{A}$ vanishes but none of the third order minor of $\mathbf{A}$ vanishes, therefore the rank of $\mathbf{A}$ is 3.

Again fourth order minor of $\mathbf{C}$ is

$$\begin{vmatrix} 1 & 2 & -1 & 3 \\ 3 & -1 & 2 & 1 \\ 2 & -2 & 3 & 2 \\ 1 & -1 & 1 & -1 \end{vmatrix} = \begin{vmatrix} 4 & -1 & 2 & 3 \\ 4 & -2 & 3 & 1 \\ 4 & -4 & 5 & 2 \\ 0 & 0 & 0 & -1 \end{vmatrix} \text{ by adding the 4th}$$

column to 1st and 3rd and subtracting 4th column from 2nd

$$\begin{aligned}
 &= \begin{vmatrix} 4 & -1 & 2 \\ 4 & -2 & 3 \\ 4 & -4 & 5 \end{vmatrix} \text{ by expanding along 4th row.} \\
 &= 4 \begin{vmatrix} 1 & 1 & 2 \\ 1 & 2 & 3 \\ 1 & 4 & 5 \end{vmatrix} \\
 &= 4 \begin{vmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & 3 & 3 \end{vmatrix} \text{ by subtracting 1st row from others.} \\
 &= 4 \begin{vmatrix} 1 & 1 \\ 3 & 3 \end{vmatrix} \text{ by expanding along 1st column.} \\
 &= 0
 \end{aligned}$$

i.e., 4th order minor of C vanishes, but none of 3rd order vanishes as may be seen easily. So the rank of C is also 3

$\therefore$ rank of $(A) = \text{rank of } (C)$

i.e., rank of coefficient matrix = rank of augmented matrix.

Hence the given system of equations is *consistent*.

Now in order to solve the given system by performing $R_{21}(-1)$, $R_{31}(-2)$ and $R_{41}(-1)$ on (1) we have

$$\begin{aligned}
 &\begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & -6 & 5 \\ 0 & -3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ -8 \\ -4 \\ -4 \end{bmatrix} \\
 \text{or } &\begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & 0 & \frac{5}{7} \\ 0 & 0 & -\frac{1}{7} \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ -8 \\ \frac{20}{7} \\ -\frac{4}{7} \end{bmatrix} \text{ by } R_{32}(-\frac{6}{7}), R_{42}(-\frac{5}{7}) \\
 \text{or } &\begin{bmatrix} 1 & 2 & -1 \\ 0 & -7 & 5 \\ 0 & 0 & \frac{5}{7} \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ -8 \\ \frac{20}{7} \\ 0 \end{bmatrix} \text{ by } R_{43}(\frac{1}{5})
 \end{aligned}$$

which is equivalent to

$$\begin{aligned}
 x + 2y - z &= 3 \\
 -7y + 5z &= -8 \\
 \frac{5}{7}z &= \frac{20}{7}
 \end{aligned}$$

giving $x = -1, y = 4, z = 4$.

Problem 39. Solve by Cramer's rule the system of equations

$$\begin{aligned}
 x + 2y + 3z &= 10 \\
 2x - 3y + z &= 1 \\
 3x + y - 2z &= 9
 \end{aligned}$$

Here $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -3 & 1 \\ 3 & 1 & -2 \end{bmatrix}$, so that $|A| = \begin{vmatrix} 1 & 2 & 3 \\ 2 & -3 & 1 \\ 3 & 1 & -2 \end{vmatrix} = 52$.

By Cramer's rule $x = \frac{\begin{vmatrix} 10 & 2 & 3 \\ 1 & -3 & 1 \\ 9 & 1 & -2 \end{vmatrix}}{|A|} = \frac{156}{52} = 3$,

$$y = \frac{\begin{vmatrix} 1 & 10 & 3 \\ 2 & 1 & 1 \\ 3 & 9 & -2 \end{vmatrix}}{|A|} = \frac{104}{52} = 2$$

$$z = \frac{\begin{vmatrix} 1 & 2 & 10 \\ 2 & -3 & 1 \\ 3 & 1 & 9 \end{vmatrix}}{|A|} = \frac{52}{52} = 1$$

Hence $x = 3, y = 2, z = 1$.

Aliter we have $|A| = \begin{vmatrix} 1 & 2 & 3 \\ 2 & -3 & 1 \\ 3 & 1 & -2 \end{vmatrix} = 52$.

$$A_{11} = \begin{vmatrix} -3 & 1 \\ 1 & -2 \end{vmatrix} = 5; \quad A_{21} = -\begin{vmatrix} 2 & 3 \\ 1 & -2 \end{vmatrix} = +7;$$

$$A_{31} = \begin{vmatrix} 2 & 3 \\ -3 & 1 \end{vmatrix} = 11;$$

$$A_{12} = -\begin{vmatrix} 2 & 1 \\ 3 & -2 \end{vmatrix} = +7; \quad A_{22} = \begin{vmatrix} 1 & 3 \\ 3 & -2 \end{vmatrix} = -11;$$

$$A_{32} = -\begin{vmatrix} 1 & 3 \\ 2 & 1 \end{vmatrix} = +5;$$

$$A_{13} = \begin{vmatrix} 2 & -3 \\ 3 & 1 \end{vmatrix} = 11; \quad A_{23} = -\begin{vmatrix} 1 & 2 \\ 3 & 1 \end{vmatrix} = +5;$$

$$A_{33} = \begin{vmatrix} 1 & 2 \\ 2 & -3 \end{vmatrix} = -7.$$

Thus

$$\text{Adj } A = \begin{bmatrix} 5 & 7 & 11 \\ 7 & -11 & 5 \\ 11 & 5 & -7 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{52} \begin{bmatrix} 5 & 7 & 11 \\ 7 & -11 & 5 \\ 11 & 5 & -7 \end{bmatrix}$$

Hence the solution of the given system of equation is given by

$$\begin{aligned}
 \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \frac{1}{52} \begin{bmatrix} 5 & 7 & 11 \\ 7 & -11 & 5 \\ 11 & 5 & -7 \end{bmatrix} \begin{bmatrix} 10 \\ 1 \\ 9 \end{bmatrix} \\
 &= \frac{1}{52} \begin{bmatrix} 50+7+99 \\ 70-11+45 \\ 110+4-63 \end{bmatrix} = \frac{1}{52} \begin{bmatrix} 156 \\ 104 \\ 52 \end{bmatrix}
 \end{aligned}$$

which gives $x = 3, y = 2, z = 1$.

2.14. CHARACTERISTIC MATRIX AND CHARACTERISTIC EQUATION OF A MATRIX

Matrix polynomials. An expression of the type

$$F(x) = A_0 + A_1x + A_2x^2 + \dots + A_mx^m,$$

where $A_0, A_1, A_2 \dots A_m$ are all square matrices of the same order, is known as *matrix polynomial* of degree m provided $A_m \neq 0$. The matrix polynomial is said to be n -rowed where $A_0, A_1, A_2, \dots A_m$ are each of order n .

The non-zero coefficient A_m of the highest power of x is called the *leading coefficient*.

The degree of the sum of two matrix polynomials cannot exceed the degree of either polynomial.

The degree of the product of two matrix polynomials is less than or equal to the sum of the degrees of the two polynomials.

Characteristic matrix. If A be any n -rowed matrix and I be an n -rowed unit matrix, then the matrix polynomial $A - xI$ of the first degree is said to be the *characteristic matrix* of A .

Characteristic polynomial. The determinant $|A - xI|$ is known as characteristic polynomial of the matrix A .

Characteristic equation. The equation $|A - xI| = 0$ is said to be the characteristic equation of the matrix A .

Cayley-Hamilton Theorem

Every square matrix satisfies its own characteristic equation,

i.e. if $|A - xI| = a_0 + a_1x + a_2x^2 + \dots + a_nx^n = 0$

be the characteristic equation of a square matrix A , then

$$a_0I + a_1A + a_2A^2 + \dots + a_nA^n = O,$$

where every x has been replaced by A and so $a_0 = a_0x^0 = a_0A^0 = a_0I$.

$\text{Adj}(A - xI)$ is a matrix polynomial, since the adjoint of $(A - xI)$ is a matrix having its elements as ordinary polynomials in x .

By the property of adjugate polynomials, we have

$$(A - xI) \text{adj}(A - xI) = |A - xI| I \quad \dots (1)$$

The relation (1) stands between three matrix polynomials namely $(A - xI)$, $\text{adj}(A - xI)$ and $|A - xI| I$ of which the last one is a scalar matrix polynomial. The relation (1) also shows that $(A - xI)$ is a left factor of the scalar polynomial $|A - xI|$, so that its left functional value is O for $x = A$. Thus

$$\begin{aligned} |A - xI| I = O &= (a_0 + a_1x + a_2x^2 + \dots + a_nx^n) I \\ &= a_0I + a_1Ix + a_2Ix^2 + \dots + a_nIx^n. \end{aligned}$$

For $x = A$, it gives $a_0I + a_1A + a_2A^2 + \dots + a_nA^n = O$.

Aliter. Since $|\text{adj } C| = |C|^{n-1}$, we may suppose that

$$\text{adj}(A - xI) = B_0 + B_1x + B_2x^2 + \dots + B_{n-1}x^{n-1},$$

then $(A - xI) \text{adj}(A - xI) = |A - xI| I$ gives

$$\begin{aligned} (A - xI)(B_0 + B_1x + B_2x^2 + \dots + B_{n-1}x^{n-1}) \\ = (a_0 + a_1x + a_2x^2 + \dots + a_nx^n) I \\ = a_0I + a_1Ix + a_2Ix^2 + \dots + a_nIx^n. \end{aligned}$$

Comparing the coefficients of like powers of x ,

$$\begin{aligned} AB_0 &= a_0I \\ AB_1 - B_0 &= a_1I, \\ AB_2 - B_1 &= a_2I, \\ &\dots \dots \dots \\ -B_{n-1} &= a_nI. \end{aligned}$$

Pre-multiplying these by $I, A, A^2, \dots A^n$ in order and adding,

$$\text{we get } a_0I + a_1A + a_2A^2 + \dots + a_nA^n = O.$$

COROLLARY. Cayley-Hamilton equation is

$$a_0I + a_1A + a_2A^2 + \dots + a_nA^n = O \quad \dots (1)$$

In characteristic polynomial $|A - xI|$ it is obvious that when $x = 0$, $a_0 = |A| \neq 0$.

Thus premultiplying (1), by A^{-1} , we get

$$A^{-1} = -\frac{a_1}{a_0}I - \frac{a_2}{a_0}A - \frac{a_3}{a_0}A^2 - \dots - \frac{a_n}{a_0}A^{n-1}.$$

Note. The right and left functional values of an n -rowed matrix polynomial $G(x)$ for any n -rowed matrix C are defined as

$$G_r(C) = A_0 + A_1C + A_2C^2 + \dots + A_kC^k,$$

$$G_l(C) = A_0 + CA_1 + C^2A_2 + \dots + C^kA_k,$$

when

$$G(x) = A_0 + A_1x + A_2x^2 + \dots + A_kx^k.$$

If $G_r(C) = O$, then $C - xI$ is called a right factor of $G(x)$ and conversely and if $G_l(C) = O$, then $C - xI$ is called left factor of $G(x)$ and conversely.

Problem 40. Find the characteristic equation of the matrix

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \\ 3 & 1 & 1 \end{bmatrix}$$

and verify Cayley-Hamilton theorem for it.

Since

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \\ 3 & 1 & 1 \end{bmatrix}$$

$$\begin{aligned} A - xI &= \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \\ 3 & 1 & 1 \end{bmatrix} - x \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1-x & 2 & 3 \\ 2 & -1-x & 4 \\ 3 & 1 & 1-x \end{bmatrix} \\ \text{so that } |A - xI| &= \begin{vmatrix} 1-x & 2 & 3 \\ 2 & -1-x & 4 \\ 3 & 1 & 1-x \end{vmatrix} = -x^3 + x^2 + 18x + 30. \end{aligned}$$

Hence the characteristic equation is

$$-x^3 + x^2 + 18x + 30 = 0.$$

Now in order to verify Cayley-Hamilton's theorem, we have to show that

$$30I + 18A + A^2 - A^3 = O. \quad \dots (1)$$

Here

$$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \\ 3 & 1 & 1 \end{bmatrix},$$

$$A^2 = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \\ 3 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \\ 3 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 14 & 3 & 14 \\ 12 & 9 & 6 \\ 8 & 6 & 14 \end{bmatrix}$$

$$\text{and } A^3 = A^2 A = \begin{bmatrix} 14 & 3 & 14 \\ 12 & 9 & 6 \\ 8 & 6 & 14 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \\ 3 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 62 & 39 & 68 \\ 48 & 21 & 78 \\ 62 & 24 & 62 \end{bmatrix},$$

so that $30I + 18A + A^2 - A^3$

$$= 30 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + 18 \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 4 \\ 3 & 1 & 1 \end{bmatrix} + \begin{bmatrix} 14 & 3 & 14 \\ 12 & 9 & 6 \\ 8 & 6 & 14 \end{bmatrix} - \begin{bmatrix} 62 & 39 & 68 \\ 48 & 21 & 78 \\ 62 & 24 & 62 \end{bmatrix}$$

$$= \begin{bmatrix} 30+18+14-62 & 0+36+3-39 & 0+54+14-68 \\ 0+36+12-48 & 30-18+9-21 & 0+72+6-78 \\ 0+54+8-62 & 0+12+6-24 & 30+18+14-62 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = O.$$

Hence Cayley-Hamilton theorem is verified.

2.15. SUB-SPACES AND NULL SPACES

Sub-space of an n -vector V_n . A set S of vectors of V_n is said to be a sub-space of V_n if it is closed with regard to the operations of addition

and scalar multiplication, i.e., if $\vec{\xi}_1, \vec{\xi}_2$ be any two members of S , then $\vec{\xi}_1 + \vec{\xi}_2$ is also a member of S and when $\vec{\xi}$ be any member of S , $k\vec{\xi}$ is also a member of S , k being a scalar.

Every sub-space of V_n , being the scalar product of any vector with the scalar zero, contains the zero vector.

Any sub-space arising as a set of all linear combinations of any given set of vectors, is said to be *spanned* by the given set of vectors.

A set of vectors is called the *Basis* of a sub-space if the sub-space is spanned by the set provided the set is linearly independent.

For example if $\vec{\xi}_1, \vec{\xi}_2, \vec{\xi}_3, \dots, \vec{\xi}_n$ be a set of vectors defined as

$$\vec{\xi}_1 = [1 \ 0 \ 0 \dots 0], \vec{\xi}_2 = [0 \ 1 \ 0 \dots 0], \dots, \vec{\xi}_n = [0 \ 0 \ 0 \dots 0 \ 1],$$

then this set is a basis of the vector space V_n if

$$k_1\vec{\xi}_1 + k_2\vec{\xi}_2 + \dots + k_n\vec{\xi}_n = \vec{0},$$

which is satisfied, when $k_1=0, k_2=0, k_3=0, \dots, k_n=0$, so that the set

linearly independent and that any vector $\vec{\xi} = [a_1, a_2, \dots, a_n]$ of V_n may be expressed as $\vec{\xi} = a_1\vec{\xi}_1 + a_2\vec{\xi}_2 + \dots + a_n\vec{\xi}_n$.

Dimension of a sub-space. The number of vectors in any basis of a sub-space is said to be the dimension of the sub-space, e.g. dimension of V_n is n .

If we have a matrix A of order $m \times n$, then each of the m rows of A consisting of n elements is an n -vector and each of the n columns, consisting of m elements is an m -vector.

The *row space* of A is one which is spanned by the m -rows which is a sub-space of V_n and the *column space* of A is the space spanned by the n -columns which is sub-space of V_m .

The dimensions of row and column spaces are respectively known as *row rank* and *column rank* of the matrix.

Premultiplication by a non-singular matrix does not alter the row rank of a matrix.

The row rank of a matrix is the same as its rank.

The column rank of a matrix is the same as its rank.

Null space and nullity of a matrix. The sub-space generated by the vectors X , such that $AX = O$ is said to be the *column null space* of matrix A of order $m \times n$ and rank r and its dimension $n-r$ is said to be the *column nullity* of matrix A , i.e.,

rank of A + column nullity of A = number of columns in the matrix A .

Similarly the sub-space of the solution of $YA = O$ is said to be the *row null space* and its dimension $m-r$ is said to be the *row nullity* of the matrix A , i.e.,

rank of A + row nullity of A = number of rows in A .

In case of a square matrix the row nullity and the column nullity are equal. Thus if $\rho(A)$ denotes the rank of A , $\nu(A)$ the nullity of square matrix A and n the number of rows of columns of A , then

$$\rho(A) + \nu(A) = n.$$

Sylvester's theorem i.e., Law of nullity.

If A and B be two n -rowed square matrices, then

$$\max. \{ \nu(A), \nu(B) \} \leq \nu(AB) \leq \nu(A) + \nu(B),$$

where $\nu(A), \nu(B), \nu(AB)$ stand for the nullities of A, B and AB .

If $\rho(A), \rho(B), \rho(AB)$ denote the ranks of the matrices A, B, AB , then we have

$$\left. \begin{aligned} \rho(A) &= n - \nu(A) \\ \rho(B) &= n - \nu(B) \\ \rho(AB) &= n - \nu(AB) \end{aligned} \right\} \dots (1)$$

Now we know that corresponding to the matrix A of rank $\rho(A)=r$ (say), there exist non-singular matrices P and Q such that

$$PAQ = \begin{bmatrix} I_r & O \\ O & O \end{bmatrix}. \quad \dots (2)$$

[Note. Here if A be an $m \times n$ matrix, then P and Q are $m \times m$ and $n \times n$ matrices respectively and then the form (2) is said to be *normal form* of A while every non-zero matrix of rank r can be reduced to the normal form by a sequence of elementary transformations.]

Premultiplying by P^{-1} and postmultiplying by Q^{-1} , (2) gives

$$A = P^{-1} \begin{bmatrix} I_r & O \\ O & O \end{bmatrix} Q^{-1}. \quad \dots (3)$$

Consider another matrix C given by

$$C = P^{-1} \begin{bmatrix} O & O \\ O & O_{n-r} \end{bmatrix} Q^{-1}, \quad \dots (4)$$

so that

$$A + C = P^{-1} \begin{bmatrix} I_r & O \\ O & O_{n-r} \end{bmatrix} Q^{-1} = P^{-1} Q^{-1}.$$

It follows that $A + C$ is a non-singular matrix.

Now

$$\rho(C) = n - \rho(A)$$

and

$$\rho(A + C) = n.$$

But A and C being non-singular, we have

$$\rho\{(A + C)B\} = \rho(B),$$

i.e.,

$$\rho(B) = \rho(AB + CB)$$

$$\leq \rho(AB) + \rho(CB).$$

Also

$$\rho(CB) \leq \rho(C) = n - \rho(A).$$

Thus (5) and (6) yield

$$\rho(B) \leq \rho(AB) + n - \rho(A)$$

or

$$\rho(AB) \geq \rho(A) + \rho(B) - n,$$

i.e., if A, B be two n -rowed square matrices, then

$$\rho(A) + \rho(B) - n \leq \rho(AB) \leq \min. \{\rho(A), \rho(B)\}$$

or $n - \nu(A) + n - \nu(B) - n \leq n - \nu(AB) \leq \min. \{n - \nu(A), n - \nu(B)\}$ from (1)

or $\nu(A) + \nu(B) \geq \nu(AB) \geq \max. \{\nu(A), \nu(B)\}.$

This proves the theorem.

Problem 41. Find the basis of the row and column null spaces of the matrix

$$\begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & 1 \\ 3 & 1 & 0 & 1 \end{bmatrix}.$$

For the column null space, let us consider the matrix equation

$$\begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \\ 3 & 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = O \quad \dots (1)$$

or $\begin{bmatrix} 1 & 1 & -1 & 1 \\ 0 & -2 & 3 & -2 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = O$ on subtracting the first row from 2nd and 3 times the first row from 3rd and then subtracting the new second row from the third new row in the first matrix.

This is equivalent to

$$x + y - z + w = 0,$$

$$-2y + 3z - 2w = 0,$$

which yield $y = \frac{3}{2}z - w$, $x = -\frac{1}{2}z$,

so that the general solution of (1) is

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} -\frac{1}{2}z \\ \frac{3}{2}z - w \\ z \\ w \end{bmatrix} = z \begin{bmatrix} -\frac{1}{2} \\ \frac{3}{2} \\ 1 \\ 0 \end{bmatrix} - w \begin{bmatrix} 0 \\ -1 \\ 0 \\ 1 \end{bmatrix} \text{ where } z \text{ and } w \text{ may be treated as two parameters.}$$

Hence the two solutions $\begin{bmatrix} -\frac{1}{2} \\ \frac{3}{2} \\ 1 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ -1 \\ 0 \\ 1 \end{bmatrix}$ constitute the basis of the

column null space of the given matrix whose nullity thus is 2 (two).

Now for the row null space, let us consider the matrix equation

$$[xyz] \begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \\ 3 & 1 & 0 & 1 \end{bmatrix} = O \quad \dots (2)$$

or $[xyz] \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & -2 & 3 & -2 \\ 3 & -2 & 3 & -2 \end{bmatrix} = O$ on subtracting the first column from 2nd and fourth and adding it to the third column in the second matrix.

or $[xyz] \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & -2 & 0 & 0 \\ 3 & -2 & 0 & 0 \end{bmatrix} = O$ on adding $\frac{3}{2}$ times the second column to the third and subtracting the second column from the fourth.

which is equivalent to

$$x + y + 3z = 0,$$

$$-2y - 2z = 0,$$

giving

$$y = -z, \quad x = -2z,$$

so that the general solution of (2) is

$$[x \ y \ z] = [-2z, -z, z] = z [-2, -1, 1],$$

where z may be treated as a parameter.

Hence the solution $[-2, -1, 1]$ constitutes the basis of the row null space of the given matrix whose row nullity is thus 1 (one).

The required basis of the row and null spaces of the given matrix are respectively

$$[-2 \ -1 \ 1] \begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \\ 3 & 1 & 0 & 1 \end{bmatrix} = O$$

and

$$\begin{bmatrix} 1 & 1 & -1 & 1 \\ 3 & -1 & 2 & -1 \\ 1 & 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 3 \\ 2 \\ 0 \\ 1 \end{bmatrix} = O$$

2.16. TRANSFORMATIONS

Linear forms. An expression of the form $\sum_{j=1}^n a_{ij}x_j$ said to be linear form of the variable x_j .

Transformation. If a_{ij} be the given constants and x_j the variables then the set of equations

$$X_i = \sum_{j=1}^n a_{ij}x_j \quad (\text{for } i=1, 2, \dots, n), \quad \dots (1)$$

is called a *linear transformation* connecting the variables x_j and the variables X_j .

The square matrix $A \equiv [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$ is said to be the *matrix of transformations*.

The determinant of the matrix

$$|A| \equiv |a_{ij}| = \begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix},$$

is said to be the *determinant* or the *modulus* of the transformation.

For the sake of convenience and brevity (1) is written as

$$X = Ax, \quad \dots (2)$$

when $|A| = 0$, the transformation is called singular and when $|A| \neq 0$, the transformation is said to be non-singular.

In case of non-singular matrix the transformation may be expressed as $x = A^{-1}X$ (on premultiplying (2) by A^{-1}).

In particular the linear transformation

$$X_1 = x_1, X_2 = x_2, \dots, X_n = x_n$$

whose matrix is $\begin{bmatrix} 0 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 \end{bmatrix} \equiv I$ is said to be the *identical transformation*

and its determinant is unity.

Resultant of two linear transformations. If the two successive transformations be $Y = PX$ and $Z = QY$, then the resultant transformation is determined by $Z = QPX$ which shows that the matrix of the resultant of two linear transformations, is the product of the matrices of the transformation, due regard being paid to the order of multiplication.

Evidently the rank of the product of two matrices cannot exceed the rank of either factor and if a matrix of rank r is multiplied (in any order) by a non-singular matrix, the rank of the product is also r .

Orthogonal transformation. Any transformation $x = AX$ that transforms $\sum x^2$ into $\sum X^2$ is said to be an *orthogonal transformation* and the matrix A is known as *orthogonal matrix*.

If

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

and its transpose

$$A' = \begin{bmatrix} a_{11} & a_{21} & \dots & a_{n1} \\ a_{12} & a_{22} & \dots & a_{n2} \\ \dots & \dots & \dots & \dots \\ a_{1n} & a_{2n} & \dots & a_{nn} \end{bmatrix}$$

then

$AA' = I$ which is the necessary and sufficient condition for a square matrix X to be orthogonal,

since if

$$x_r = a_{r1}X_1 + a_{r2}X_2 + \dots + a_{rn}X_n,$$

then $\sum_{r=1}^n X_r^2 = \sum_{r=1}^n x_r^2 = \sum_{r=1}^n (a_{r1}X_1 + a_{r2}X_2 + \dots + a_{rn}X_n)^2$ gives

$$\left. \begin{aligned} a_{11}^2 + a_{21}^2 + \dots + a_{n1}^2 &= 1 \\ a_{11}a_{1j} + a_{21}a_{2j} + \dots + a_{n1}a_{nj} &= 0 \end{aligned} \right\}$$

and

for $i=1, 2, \dots, n, j=1, 2, \dots, n$ and $i \neq j$.

Now $AA' = I$ gives

$$|A| |A'| = 1,$$

where $|A| = |A'|$ as interchange of rows and columns does not alter the value of the determinant.

$$\therefore |A|^2 = 1, \text{ i.e., } |A| = \pm 1.$$

Evidently the product of two orthogonal transformations is an orthogonal transformation. For if $x = AX$ and $X = BY$ be two orthogonal transformations then $AA' = I, BB' = I$.

$$\begin{aligned} \therefore [AB][AB]' &= ABB'A' \text{ by the law of reversal transpose} \\ &= AIA' \\ &= AA' = I. \end{aligned}$$

Orthonormal Set. A set of vectors is said to be an *orthonormal set* of vectors if

- (i) each vector of the set is a normal vector
- (ii) any two vectors of the set are orthogonal

While a complex n -vector X is said to be orthogonal to another complex n -vector Y if $(X, Y) = 0$ i.e., if $X^*Y = 0$.

The relation of orthogonality in the set of all complex n -vectors is *Symmetric*. The positive square root of X^*X is known as the *length* of X .

In other words using the Kronecker delta symbol $\delta_{ij} = 0$ for $i \neq j$, $= 1$ for $i = j$.

a set S of complex n -vectors $X_1, X_2, \dots, X_k$ is termed as an orthonormal set if $(X_i, X_j) = \delta_{ij}$, $i = 1, 2, \dots, k, j = 1, 2, \dots, k$.

THEOREM 1. Every orthonormal set of vectors is linearly independent.

Suppose that the vectors $X_1, X_2, \dots, X_k$ form an orthonormal set of vectors.

- Thus (i) X_i is a normal vector for every i i.e. $X_i^* X_i = I$
 (ii) X_i, X_j are orthogonal vectors for every i, j such that $i \neq j$
 i.e., $X_i^* X_j = 0$ for every i and $j, i \neq j$

Consider a relation $a_1 X_1 + a_2 X_2 + \dots + a_k X_k = 0$... (1)
 where $a_1, a_2, \dots, a_k$ are scalars.

Premultiplying (1) by X_i^* and applying above condition (i) and (ii), we find $a_1 I = 0$ or $a_1 = 0$ as $I \neq 0$.

Similarly premultiplying (1) by $X_2^*, X_3^*, \dots$ successively, we may get $a_2 = 0, a_3 = 0, \dots, a_k = 0$.

As such all the scalars $a_1, a_2, \dots, a_k$ being zero, the relation (1) follows that $X_1, X_2, \dots, X_k$ form an orthonormal set of linearly independent vectors.

THEOREM 2. Show that a real matrix is unitary if and only if it is orthogonal.

If A is a real matrix, then $A^* = A'$

$\therefore A$ is unitary if $A^* A = I$ or if $A' A = I$ i.e., A is orthogonal.

Conversely if A is orthogonal then $A' A = I$

i.e., $A^* A = I$, i.e. A is unitary.

2.17. HERMITIAN FORMS

Any expression of the forms $\sum_{j=1}^n \sum_{i=1}^n a_{ij} \bar{x}_i x_j$

where $a_{ij} = a_{ji}$ is said to be a Hermitian form in n -variables.

It is easy to see that $\sum_{j=1}^n \sum_{i=1}^n a_{ij} \bar{x}_i x_j = X' A X$ where $A = [a_{ij}]$
 $= X^* A X$

Where X is a Column vector with components $x_1, x_2, x_3, \dots, x_n$ and X^* is a row vector with components $\bar{x}_1, \bar{x}_2, \dots, \bar{x}_n$. Also the matrix A is Hermitian as $a_{ij} = a_{ji}$.

THEOREM 3. A Hermitian form assumes only real values.

We have $(X^* A X) = \overline{X^* A X}$,

where $\overline{X^* A X}$ being a single element matrix coincides with its transpose.

$$\begin{aligned} \therefore \overline{X^* A X} &= (\overline{X^* A X})' = (\overline{X^* A X})' \\ &= (\overline{X^* A})' \overline{X}', \therefore \overline{X^*} = X' \\ &= (\overline{X})' \overline{A}' (X')' = X^* A X, \therefore (\overline{X})' = X^*, \overline{A}' = A \text{ and } (X')' = X \end{aligned}$$

so that $(X^* A X) = X^* A X$,

i.e., $X^* A X$ coincides with its conjugate and therefore it is real.

Linear Transformation of a Hermitian form. Consider a Hermitian form $X^* A X$.

Any non-singular linear transformation say $X = P Y$, i.e., $X^* = Y^* P^*$ gives

$$X^* A X = Y^* P^* A P Y = Y^* B Y \text{ (say)}$$

where $B = P^* A P$ is the matrix of the transformation.

$$\begin{aligned} \text{Now } B^* &= (P^* A P)^* = P^* A^* (P^*)^* \\ &= P^* A P = B, \end{aligned}$$

i.e., B is also Hermitian if A is Hermitian, since for a Hermitian matrix $A^* = A$.

2.18. CHARACTERISTIC ROOTS AND VECTORS OF A MATRIX (EIGEN VALUES AND EIGEN VECTORS)

A non-zero vector X is called a *characteristic vector* of a matrix A , if there is a number λ such that $A X = \lambda X$.

Here λ is called a *characteristic root* of A corresponding to the characteristic vector X and vice versa.

Characteristic roots are often known as *Proper, Latent* or *Eigen values* and characteristic vectors are known as *Proper, Latent* or *Eigen vectors* or *in variant vectors*.

We have $A X = \lambda X = \lambda I X$, I being a unit matrix

$$(A - \lambda I) X = 0.$$

Since $X \neq 0$, the matrix $(A - \lambda I)$ is singular, so that

$$| (A - \lambda I) | = 0.$$

which follows that every characteristic root λ of a matrix A is a root of its characteristic equation $| A - \lambda I | = 0$.

Conversely if λ be a root of the characteristic equation $| A - \lambda I | = 0$, then the matrix equation $(A - \lambda I) X = 0$ possesses a non-trivial i.e., non-zero solution for X , so that there exists a vector $X \neq 0$ such that $A X = \lambda I X = \lambda X$.

It follows that every root of the characteristic equation of a matrix is a characteristic root of the matrix.

Thus if $A = [a_{ij}]$ be an n -rowed square matrix and λ an indeterminate, then the characteristic equation

$$| A - \lambda I | = 0 \text{ gives } \begin{vmatrix} a_{11} - \lambda & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} - \lambda & a_{23} & \dots & a_{2n} \\ a_{31} & a_{32} & a_{33} - \lambda & \dots & a_{3n} \\ \dots & \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} - \lambda \end{vmatrix} = 0 \quad \dots (1)$$

which is an ordinary polynomial in λ of degree n and hence will give n values of λ on simplification. These n values of λ are n eigen values

of this equation. To every eigen value there corresponds an eigen vector.

As an illustrative example if

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 5 & 2 & 6 \\ -2 & -1 & -3 \end{bmatrix}$$

then its characteristic equation $|A - \lambda I| = 0$ is

$$\begin{vmatrix} 1-\lambda & 1 & 3 \\ 5 & 2-\lambda & 6 \\ -2 & -1 & -3-\lambda \end{vmatrix} = 0$$

which yields on expansion $\lambda^3 = 0$

giving $\lambda_1 = 0 = \lambda_2 = \lambda_3$ i.e., all the three eigen roots are zero.

Nature of the eigen values and eigen vectors of special types of matrices.

THEOREM. 1 The eigen values of a Hermitian matrix are all real (Rohilkhand, 1977; Agra, 1970; Vikram, 1966)

Let λ be an eigen value of a Hermitian matrix A . Then by definition there exists a vector $X \neq 0$, such that

$$AX = \lambda X.$$

Premultiplying by X° we get

$$\begin{aligned} X^\circ AX &= X^\circ \lambda X \\ &= \lambda X^\circ X = \lambda X^\circ IX \end{aligned} \quad \dots (1)$$

so that

$$\lambda = \frac{X^\circ AX}{X^\circ IX} \quad \dots (2)$$

Also by § 2.17, we have $X^\circ AX = \overline{X^\circ AX}$, so that $X^\circ AX$ is a real number and therefore in particular $X^\circ IX$ is also a real number.

Hence from (2) it follows that λ is a real number, i.e., all the eigen values of a Hermitian matrix are real.

THEOREM 2. The eigen values of a skew-Hermitian matrix are purely imaginary or zero.

If A be a skew-Hermitian matrix and $AX = \lambda X$, then

$$(iA)X = (i\lambda)X.$$

But iA is Hermitian and therefore its eigen values $i\lambda$ are real. It follows that λ is either zero or purely imaginary number.

Note. In this case $X^\circ AX = -X^\circ AX$.

THEOREM 3. The modulus of each eigen value of a unitary matrix is unity, i.e., the eigen values of a unitary form have the absolute value 1.

Let U be the unitary matrix, λ an eigen value of U and X a corresponding eigen vector; then

$$UX = \lambda X. \quad \dots (1)$$

Taking conjugate transpose of either side, we get

$$X^\circ U^\circ = \lambda X^\circ. \quad \dots (2)$$

Post-multiplying (2) by (1), we get

$$X^\circ U^\circ UX = \lambda \bar{\lambda} X^\circ X, \quad \bar{\lambda} \text{ being complex conjugate of } \lambda$$

or $X^\circ X = \lambda \bar{\lambda} X^\circ X$ as $U^\circ U = I$, U being unitary

or $(1 - \lambda \bar{\lambda}) X^\circ X = 0.$

Since $X \neq 0$, therefore $X^\circ X \neq 0$ and hence

$$1 - \lambda \bar{\lambda} = 0, \text{ i.e., } \lambda \bar{\lambda} = |\lambda|^2 = 1,$$

i.e., the modulus of λ is unity.

Note. The above three theorems when particularised, give that

(i) the eigen values of a symmetric matrix are real,

(ii) the eigen values of a skew-symmetric matrix are zero or purely imaginary, and

(iii) the eigen values of an orthogonal matrix have the absolute unity and are real, or complex conjugate in pairs.

Thus taking λ as an eigen value of the matrix A and X the corresponding eigen vector, we have

$$AX = \lambda X,$$

$$A^2 X = \lambda AX = \lambda^2 X,$$

$$A^3 X = \lambda A^2 X = \lambda^3 X.$$

$$\dots \dots \dots$$

Therefore

$$\begin{aligned} [A^n - I_{(1)} A^{n-1} + I_{(2)} A^{n-2} - \dots + (-1)^n I_{(n)} I] X \\ = [\lambda^n - I_{(1)} \lambda^{n-1} + I_{(2)} \lambda^{n-2} - \dots + (-1)^n I_{(n)}] X, \end{aligned} \quad \dots (1)$$

since $A^0 = I$.

If the characteristic equation be taken as

$$|A - \lambda I| = (-1)^n [\lambda^n - I_{(1)} \lambda^{n-1} + I_{(2)} \lambda^{n-2} - \dots + (-1)^n I_{(n)}] = 0,$$

then (1) reduces to

$$A^n - I_{(1)} A^{n-1} + I_{(2)} A^{n-2} - \dots + (-1)^n I_{(n)} I = 0, \quad \dots (2)$$

which is Cayley-Hamilton Theorem.

THEOREM 4. Every matrix equation satisfies its own characteristic equation. (Cayley-Hamilton Theorem).

It is easy to show that the theorem that every square matrix satisfies its own characteristic equation also holds for singular

matrices as well as for matrices that have repeated eigen values. Its proof has already been given in §2.14.

If A is non-singular matrix, then multiplying the result (2) above by A^{-1} , we get

$$A^{-1} = \frac{(-1)^{n+1}}{I_{(n)}} [A^{n-1} - I_{(1)} A^{n-2} + I_{(2)} A^{n-3} - \dots + (-1)^n I_{(n-1)}],$$

which gives a method of finding the inverse of a matrix.

THEOREM 5. Any two given vectors corresponding to two distinct eigen values of a Hermitian matrix are orthogonal.

Let X_1 and X_2 be two eigen vectors corresponding to two distinct eigen values λ_1, λ_2 of a Hermitian matrix A ; then

$$AX_1 = \lambda_1 X_1, \quad \dots (1)$$

$$AX_2 = \lambda_2 X_2, \quad \dots (2)$$

where from theorem (1) the numbers λ_1, λ_2 are real.

Premultiplying (1) and (2) by X_2^0 and X_1^0 respectively

$$X_2^0 AX_1 = \lambda_1 X_2^0 X_1, \quad \dots (3)$$

$$X_1^0 AX_2 = \lambda_2 X_1^0 X_2, \quad \dots (4)$$

$$\text{But } (X_2^0 AX_1)^0 = X_1^0 AX_2,$$

$$\therefore \text{ for a Hermitian matrix } A^0 = A \quad \text{and also } (X_2^0)^0 = X_2.$$

therefore we have from (3) and (4),

$$(\lambda_1 X_2^0 X_1)^0 = \lambda_2 X_1^0 X_2$$

$$\text{or } \lambda_1 X_1^0 X_2 = \lambda_2 X_1^0 X_2$$

$$\text{or } (\lambda_1 - \lambda_2) X_1^0 X_2 = 0.$$

Since $\lambda_1 - \lambda_2 \neq 0$, otherwise the roots will not be distinct, therefore the only possibility is that $X_1^0 X_2 = 0$.

It follows that X_1, X_2 are orthogonal.

COROLLARY. Any two eigen vectors corresponding to two distinct eigen values of a real symmetric matrix are orthogonal. (Agra, 1974)

THEOREM 6. Any two eigen vectors corresponding to two distinct eigen values of a unitary matrix are orthogonal.

Let X_1, X_2 be two eigen vectors corresponding to two distinct eigen values λ_1, λ_2 of unitary matrix U ; then

$$UX_1 = \lambda_1 X_1, \quad \dots (1)$$

$$UX_2 = \lambda_2 X_2, \quad \dots (2)$$

Taking conjugate transpose of (2), we get

$$X_2^0 U^0 = \bar{\lambda}_2 X_2^0, \quad \dots (3)$$

From (1) and (3), we find

$$X_2^0 U^0 UX_1 = \bar{\lambda}_2 \lambda_1 X_2^0 X_1$$

$$\text{or } X_2^0 X_1 = \bar{\lambda}_2 \lambda_1 X_2^0 X_1 \quad \text{since } U^0 U = I$$

$$\text{or } (1 - \bar{\lambda}_2 \lambda_1) X_2^0 X_1 = 0. \quad \dots (4)$$

But U being a unitary matrix, the modulus of each of its eigen values is unity, i.e., $\lambda_2 \bar{\lambda}_2 = 1$.

$$\text{So that } (1 - \lambda_2 \bar{\lambda}_1) = \lambda_2 \bar{\lambda}_2 - \bar{\lambda}_2 \lambda_1, \bar{\lambda}_2 (\lambda_2 - \lambda_1) \neq 0 \quad \dots (5)$$

From (4) and (5) it follows that $X_2^0 X_1 = 0$ i.e., X_1 and X_2 are orthogonal.

THEOREM 7. The eigen vectors corresponding to distinct characteristic roots of a matrix are linearly independent.

Assuming that $X_1, X_2, \dots, X_m$ are eigen vectors of a matrix A corresponding to distinct eigen values $\lambda_1, \lambda_2, \dots, \lambda_n$ such that $AX_i = \lambda_i X_i, i=1, 2, \dots, m$, we have to show that $X_1, X_2, \dots, X_m$ are linearly independent.

Suppose that $X_1, X_2, \dots, X_m$ are not linearly independent but they are linearly dependent, then we can choose r such that $1 \leq r < m$ and $X_1, X_2, \dots, X_r$ are linearly independent, but $X_1, X_2, \dots, X_r, X_{r+1}$ are linearly dependent so that there exist scalars $a_1, a_2, \dots, a_{r+1}$ not all zero, satisfying

$$a_1 X_1 + a_2 X_2 + \dots + a_{r+1} X_{r+1} = 0 \quad \dots (1)$$

$$A(a_1 X_1 + a_2 X_2 + \dots + a_{r+1} X_{r+1}) = A0$$

$$a_1 AX_1 + a_2 AX_2 + \dots + a_{r+1} AX_{r+1} = 0$$

$$a_1 (\lambda_1 X_1) + a_2 (\lambda_2 X_2) + \dots + a_{r+1} (\lambda_{r+1} X_{r+1}) = 0 \quad \dots (2)$$

Multiplying (1) again by λ_{r+1} and subtracting from (2), we find,

$$a_1 (\lambda_1 - \lambda_{r+1}) X_1 + a_2 (\lambda_2 - \lambda_{r+1}) X_2 + \dots + a_r (\lambda_r - \lambda_{r+1}) X_r = 0 \quad \dots (3)$$

But $X_1, X_2, \dots, X_r$ are linearly independent by hypothesis and $\lambda_1, \lambda_2, \dots, \lambda_{r+1}$ are distinct, it therefore follows from (3) that

$$a_1 = 0 = a_2 = \dots = a_r$$

$$\text{and then (1) gives } a_{r+1} X_{r+1} = 0$$

$$\text{So that } a_{r+1} = 0 \text{ as } X_{r+1} \neq 0$$

$$\text{Hence } a_1 = 0 = a_2 = \dots = a_r = a_{r+1}$$

which contradicts the assumption that $a_1, a_2, \dots, a_{r+1}$ are not all zero.

It follows from this contradiction that our initial assumption is wrong and the only possibility is that $X_1, X_2, \dots, X_m$ are linearly independent.

THEOREM 8. The minimal polynomial of a matrix is a divisor of the characteristic polynomial of that matrix.

If $f(x)$ be the characteristic polynomial of a matrix A , then by Cayley-Hamilton theorem, $f(A) = 0$ so that $f(x)$ annihilates A . Thus if $m(x)$ is the minimal polynomial of A , then $m(x)$ is divisor of $f(x)$ by [19].

THEOREM 9. Every root of the minimal equation of a matrix is a characteristic root of that matrix.

If $f(x)$ be the characteristic polynomial of a matrix A and $m(x)$ be the minimal polynomial, then by preceding theorem $m(x)$ is a divisor of $f(x)$ and so there exists a polynomial $q(x)$ such that

$$f(x) = m(x) q(x) \quad \dots (1)$$

$$\text{If } \lambda \text{ be a root of } m(x) = 0, \text{ then } m(\lambda) = 0 \quad \dots (2)$$

Putting $x=\lambda$ in (1) we get, $f(\lambda)=m(\lambda)q(\lambda)=0$ by (2) which follows that λ is also a root of $f(x)=0$

Hence λ is also a characteristic root of A .

THEOREM 10. *The characteristic polynomial and hence the eigen values of similar matrices are the same. Also if X be an eigen vector of A corresponding to the eigen value λ , then $P^{-1}X$ is an eigen vector of B corresponding to the eigen value λ where $B=P^{-1}AP$.*

Let A and B be two similar matrices. Then there exists an invertible matrix P such that

$$B=P^{-1}AP.$$

Consider $B-\lambda I=P^{-1}AP-\lambda I$

$$=P^{-1}AP-P^{-1}(\lambda I)P \quad \because P^{-1}(\lambda I)P=\lambda P^{-1}P=\lambda I \\ =P^{-1}(A-\lambda I)P$$

$$\therefore |B-\lambda I|=|P^{-1}||A-\lambda I||P| \\ =|P^{-1}||P||A-\lambda I| \quad \because \text{scalar quantities commute} \\ =|P^{-1}P||A-\lambda I| \quad \because |CD|=|C||D| \\ =|A-\lambda I| \quad \because |P^{-1}P|=|I|=1$$

which follows that A and B have same characteristic polynomial and so they have the same eigen values.

For the second part, taking λ as one of the eigen values of A and X corresponding eigen vector, we have $AX=\lambda X$

$$\therefore B(P^{-1}X)=(P^{-1}AP)P^{-1}X=P^{-1}AX \quad \because PP^{-1}=I \text{ etc.} \\ =P^{-1}(\lambda X) \text{ by (1)} \\ =\lambda(P^{-1}X)$$

which follows that $P^{-1}X$ is an eigen vector of B corresponding to the eigen value λ .

COROLLARY. *If X is similar to a diagonal matrix D then diagonal elements of D are the eigen values of A .*

Since D has its diagonal elements as its eigen values and D is similar to A it therefore follows that diagonal element of D are eigen values of A .

Problem 42 *If X is a characteristic vector of a matrix A corresponding to eigen value λ , then show that kX , k being a non-zero scalar is also an eigen vector of A corresponding to λ .*

As given, $X \neq 0$, and $AX=\lambda X$

$$k \neq 0 \text{ and } X \neq 0 \text{ give } kX \neq 0$$

$$\text{Now, } A(kX)=k(AX)=k\lambda X \text{ by (1)} \\ =\lambda(kX)$$

which follows that kX is an eigen vector of A corresponding to the eigen value λ .

Problem 43. *If X is an eigen vector of a matrix A , then prove that it cannot correspond to more than one characteristic values of A .*

If possible let us suppose that X is an eigen vector corresponding to two eigen values λ_1, λ_2 of A . Then, we have

$$AX=\lambda_1 X \text{ and } AX=\lambda_2 X$$

These give, $\lambda_1 X=\lambda_2 X$ i.e., $(\lambda_1-\lambda_2)X=0$

$$\therefore \lambda_1=\lambda_2 \text{ as } X \neq 0.$$

i.e., λ_1 and λ_2 cannot be different.

Problem 44. *Determine the eigen values and eigen vectors of the matrix $A=\begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix}$*

The characteristic equation of A is given by $|A-\lambda I|=0$ i.e.,

$$\begin{vmatrix} 5-\lambda & 4 \\ 1 & 2-\lambda \end{vmatrix}=0 \text{ i.e., } \lambda^2-7\lambda+6=0 \text{ or } (\lambda-1)(\lambda-6)=0$$

$\therefore \lambda_1=1, \lambda_2=6$ are the eigen values of A .

Now the eigen vector X_1 of A corresponding to the eigen value $\lambda_1=1$ is given by the non-zero solutions of the equation $(A-1I)X_1=0$

$$\text{i.e., } \left(\begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \text{ or } \begin{bmatrix} 4 & 4 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The coefficient matrix here is of rank 1 and therefore these equations have 2-1 i.e., 1 linear independent solution.

These are equivalent to

$$4x_1+4x_2=0 \text{ and } x_1+x_2=0$$

which yield, $x_1=-x_2$

If we take $x_1=1$, then $x_2=-1$ so that the eigen vector of A

$$X_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ corresponding to the eigen value 1.}$$

In fact every non-zero multiple of the vector X_1 is an eigen vector of A corresponding to the eigen value 1.

Again the eigen vector X of A corresponding to the eigen value $\lambda_2=6$ is given by the non-zero solution of the equation $(A-6I)X_2=0$

$$\text{i.e., } \begin{bmatrix} -1 & 4 \\ 1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Here also the coefficient matrix being of rank 1, these equations have 2-1 i.e., 1 linear independent solution.

They are equivalent to

$$-x_1+4x_2=0 \text{ and } x_1-4x_2=0$$

which are the same and satisfied by $x_1=4, x_2=1$

$$\therefore X_2 = \begin{bmatrix} 4 \\ 1 \end{bmatrix} \text{ is an eigen vector of } A \text{ corresponding to the eigen}$$

value 6. In general it may be represented by CX_2 i.e., $C \begin{bmatrix} 4 \\ 1 \end{bmatrix}$, where C

is any non-zero scalar.

Problem 45. Determine the eigen values and eigen vectors of the matrix

$$A = \begin{bmatrix} 3 & 1 & 4 \\ 0 & 2 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

The characteristic equation of A is $|A - \lambda I| = 0$ i.e.,

$$\begin{vmatrix} 3-\lambda & 1 & 4 \\ 0 & 2-\lambda & 0 \\ 0 & 0 & 5-\lambda \end{vmatrix} = 0 \text{ i.e., } (\lambda-2)(\lambda-3)(\lambda-5) = 0$$

$$\therefore \lambda_1 = 2, \lambda_2 = 3, \lambda_3 = 5$$

which are the eigen values of A.

To determine eigen vectors let us consider the eigen values one by one.

(i) When $\lambda_1 = 2$ the eigen vector X_1 is given by $(A - 2I)X_1 = 0$ i.e.

$$\begin{bmatrix} 1 & 1 & 4 \\ 0 & 0 & 6 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The rank of coefficient matrix being 2, the equation will have only 3-2 i.e., 1 linearly independent solution.

These are equivalent to $x_1 + x_2 + 4x_3 = 0$

$$6x_3 = 0$$

$$3x_3 = 0$$

The last two give $x_3 = 0$ and then first one gives $x_1 + x_2 = 0$.

Take $x_1 = 1$, then $x_2 = -1$ and $x_3 = 0$.

Hence $X_1 = C_1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$ C_1 being a scalar

(ii) When $\lambda_2 = 3$, the eigen vector X_2 is given by $(A - 3I)X_2 = 0$ i.e.

$$\begin{bmatrix} 0 & 1 & 4 \\ 0 & -1 & 6 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

which are equivalent to $x_2 + 4x_3 = 0$

$$-x_2 + 6x_3 = 0$$

$$2x_3 = 0$$

giving $x_3 = 0$, $x_2 = 0$ and x_1 is arbitrary say $x_1 = 1$, then

$$X_2 = C_2 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad C_2 \text{ being a scalar}$$

(iii) When $\lambda_3 = 5$, the eigen vector X_3 is given by $(A - 5I)X_3 = 0$ i.e.

$$\begin{bmatrix} -2 & 1 & 4 \\ 0 & -3 & 6 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

which are equivalent to $-2x_1 + x_2 + 4x_3 = 0$
 $-3x_2 + 6x_3 = 0$

giving $x_2 = 2x_3 = \frac{2}{3}x_1$ i.e. $2x_1 = 3x_2 = 6x_3$

Take $x_3 = 1$, so that $x_2 = 2$ and $x_1 = 3$

Hence $X_3 = C_3 \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ C_3 being a scalar.

Problem 46. Show that the eigen values of a diagonal matrix are given by its diagonal elements.

Let $D = \text{diag } [a_{11}, a_{22}, \dots, a_{nn}] = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix}$

Its characteristic equation is

$$|D - \lambda I| = 0 \text{ i.e. } \begin{vmatrix} a_{11}-\lambda & 0 & \dots & 0 \\ 0 & a_{22}-\lambda & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{nn}-\lambda \end{vmatrix} = 0$$

i.e. $(a_{11}-\lambda)(a_{22}-\lambda)\dots(a_{nn}-\lambda) = 0$

giving $\lambda = a_{11}, a_{22}, \dots, a_{nn}$ which are eigen values of D and these are clearly the diagonal elements of D.

Problem 47. Prove that any square matrix A and its transpose A' have the same eigen values.

If $A = [a_{ij}]_{m \times m}$ then $A' = [a_{ji}]_{m \times m}$

The characteristic equations of A and A' are

$$|A - \lambda I| = 0 \text{ and } |A' - \lambda I| = 0$$

$$\text{i.e. } \begin{vmatrix} a_{11}-\lambda & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22}-\lambda & \dots & a_{2m} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mm} \end{vmatrix} = 0 \text{ and } \begin{vmatrix} a_{11}-\lambda & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22}-\lambda & \dots & a_{m2} \\ \dots & \dots & \dots & \dots \\ a_{1m} & a_{2m} & \dots & a_{mm} \end{vmatrix} = 0$$

both the determinants have the same values as interchange of rows and columns does not alter the value of the determinant.

$$\therefore |A - \lambda I| = |A' - \lambda I| = 0$$

which follows that characteristic equations of A and A' are the same and hence they have the same eigen values.

Problem 48. If A be a non-singular matrix then prove that the eigen values of A^{-1} are the reciprocals of the eigen values of A.

If λ be an eigen value of A and X the corresponding eigen vector, then we have

$$AX = \lambda X \quad \dots (1)$$

Pre-multiplying both sides by A^{-1} and using $A^{-1}A = I$ and $IX = X$, we get

$$X = A^{-1}(\lambda X) = \lambda(A^{-1}X) \text{ or } A^{-1}X = \frac{1}{\lambda} X \text{ which in view of (1)}$$

follows that $\frac{1}{\lambda}$ is an eigen value of A^{-1} and X is the corresponding eigen vector.

Conversely if k is an eigen value of A^{-1} , then A being non singular gives its inverse A^{-1} to be non-singular and $(A^{-1})^{-1}=A$, so that $\frac{1}{k}$ is an eigen value of A .

As such each eigen value of A^{-1} is equal to the reciprocal of some eigen value of A .

Problem 49. If A is a square matrix of order n and has $\lambda_1, \lambda_2, \dots, \lambda_n$ as its eigen values then show that eigen values of kA will be $k\lambda_1, k\lambda_2, \dots, k\lambda_n$, k being a non-zero scalar.

If $A=[a_{ij}]_{n \times n}$ then its characteristic equation is $|A-\lambda I|=0$
i.e. $(\lambda-\lambda_1)(\lambda-\lambda_2)\dots(\lambda-\lambda_n)=0 \dots (1)$

The characteristic equation of kA is $|kA-\lambda I|=0$

$$\text{i.e. } \begin{vmatrix} ka_{11}-\lambda & ka_{12} & \dots & ka_{1n} \\ ka_{21} & ka_{22}-\lambda & \dots & ka_{2n} \\ \dots & \dots & \dots & \dots \\ ka_{n1} & ka_{n2} & \dots & ka_{nn}-\lambda \end{vmatrix} = 0$$

$$\text{or } \begin{vmatrix} a_{11}-\frac{\lambda}{k} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22}-\frac{\lambda}{k} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn}-\frac{\lambda}{k} \end{vmatrix} = 0$$

$$\text{or } |A-\frac{\lambda}{k}I|=0$$

$$\text{i.e. } \left(\frac{\lambda}{k}-\lambda_1\right)\left(\frac{\lambda}{k}-\lambda_2\right)\dots\left(\frac{\lambda}{k}-\lambda_n\right)=0 \text{ by (1) on putting } \frac{\lambda}{k} \text{ for } \lambda$$

Hence the eigen values of kA are $k\lambda_1, k\lambda_2, \dots, k\lambda_n$.

Problem 50. If eigen values of A are $\lambda_1, \lambda_2, \dots, \lambda_n$ then prove that those of A^2 will be $\lambda_1^2, \lambda_2^2, \dots, \lambda_n^2$.

If X be the eigen vector of A corresponding to a eigen value λ of A then

$$AX=\lambda X \dots (1)$$

$$\text{or } A(AX)=A(\lambda X)$$

$$\text{or } A^2X=\lambda(A X)=\lambda(\lambda X) \text{ by (1)}$$

$$\text{or } A^2X=\lambda^2 X$$

which follows that λ^2 is the eigen value of A^2 corresponding to the eigen vector X .

Hence if $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigen values of A , then $\lambda_1^2, \lambda_2^2, \dots, \lambda_n^2$ are the eigen values of A^2 .

Problem 51. If A and B are two square matrices, then show that AB and BA have the same eigen values. Also show that $A^{-1}B$ and BA^{-1} have the same eigen values when $|A| \neq 0$.

$$\text{We have } BA=(A^{-1}A)BA=A^{-1}(AB)A$$

So that BA and AB are similar and hence have same eigen values since similar matrices have the same eigen values.

$$\text{Similarly } BA^{-1}=(AA^{-1})BA^{-1}=A(A^{-1}B)A^{-1}$$

BA^{-1} and $A^{-1}B$ are similar and hence etc.

Problem 52. Show that the eigen roots of A° are the conjugate of the eigen roots of A .

The characteristic equations of A and A° are as given

$$|A-\lambda I|=0 \text{ and } |A^\circ-\bar{\lambda} I|=0$$

$$\text{But } |A^\circ-\bar{\lambda} I|=|(A-\lambda I)^\circ|=|\overline{A-\lambda I}|$$

$$\therefore |A^\circ|=|\overline{A}|=|\overline{A}|$$

$$|A^\circ-\bar{\lambda} I|=0 \text{ iff } |\overline{A-\lambda I}|=0$$

or $|A^\circ-\bar{\lambda} I|=0$ or iff $|A-\lambda I|=0$ since if z be a complex number then $z=0$ iff $\bar{z}=0$.

Thus $\bar{\lambda}$ is an eigen value of A° iff λ is an eigen value of A .

Problem 53. Show that the matrix $A = \begin{vmatrix} 1 & 0 & 0 \\ 1 & -1 & 0 \\ 1 & 0 & -1 \end{vmatrix}$ is derogatory.

The characteristic equation of A is $|A-\lambda I|=0$

$$\begin{vmatrix} 1-\lambda & 0 & 0 \\ 1 & -1-\lambda & 0 \\ 1 & 0 & -1-\lambda \end{vmatrix} = 0 \text{ or } (\lambda-1)(\lambda+1)^2=0$$

giving $\lambda=1, -1, -1$, so that by theorem 9, the factors $x-1$ and $x+1$ both must be divisor of the minimal polynomial of A .

We have to see whether $h(x)=(x-1)(x+1)=x^2-1$ annihilates A or not.

$$\text{Consider } A^2=A \cdot A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

$$A^2-I=0$$

Which shows that $h(x)$ annihilates A and as such $h(x)$ is the minimal polynomial of A . Also degree of $h(x)$ is less than 3 and hence A is derogatory.

Problem 54. If X be an eigen vector of $B=P^{-1}AP$ corresponding to eigen value λ , then show that $Y=PX$ is an eigen vector of A corresponding to the same eigen value.

$$\text{We have } BX=\lambda X$$

or

$$\mathbf{PBX} = \mathbf{P}\lambda\mathbf{X} \quad \text{or} \quad \mathbf{P}(\mathbf{P}^{-1}\mathbf{A}\mathbf{P})\mathbf{X} = \mathbf{P}\lambda\mathbf{X}$$

or

$$\mathbf{A}(\mathbf{P}\mathbf{X}) = \lambda(\mathbf{P}\mathbf{X}) \quad \because \quad \mathbf{P}\mathbf{P}^{-1} = \mathbf{I} \quad \text{and} \quad \mathbf{I}\mathbf{A} = \mathbf{A} \quad \text{etc.}$$

which follows that $\mathbf{P}\mathbf{X}$ is an eigen vector of $\mathbf{A}$ corresponding to the eigen value λ .

Such eigen vectors are often called as the *invariant eigen vectors*.

Diagonalization of matrices. Let $\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_n$ be n distinct eigen values of a matrix $\mathbf{A}$ and $\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \dots, \mathbf{X}_n$ be the n corresponding eigen vectors. Also let $\mathbf{X}_i$ be the column vector given by

$$\mathbf{X}_i = \begin{bmatrix} X_{1i} \\ X_{2i} \\ \dots \\ X_{ni} \end{bmatrix}$$

Consider a matrix $\mathbf{E}$ whose column vectors are the n eigen vectors such that

$$\mathbf{E} = \begin{bmatrix} \mathbf{X}_{11} & \mathbf{X}_{12} & \dots & \mathbf{X}_{1n} \\ \mathbf{X}_{21} & \mathbf{X}_{22} & \dots & \mathbf{X}_{2n} \\ \dots & \dots & \dots & \dots \\ \mathbf{X}_{n1} & \mathbf{X}_{n2} & \dots & \mathbf{X}_{nn} \end{bmatrix} = [\mathbf{X}_{ij}] \quad (\text{say})$$

Suppose that $\mathbf{D}$ is a diagonal matrix such that

$$\mathbf{D} = \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix} = \text{diag} [\lambda_1, \lambda_2, \dots, \lambda_n]$$

$$\text{Then } \mathbf{E}\mathbf{D} = \begin{bmatrix} \lambda_1\mathbf{X}_{11} & \lambda_2\mathbf{X}_{12} & \dots & \lambda_n\mathbf{X}_{1n} \\ \lambda_1\mathbf{X}_{21} & \lambda_2\mathbf{X}_{22} & \dots & \lambda_n\mathbf{X}_{2n} \\ \dots & \dots & \dots & \dots \\ \lambda_1\mathbf{X}_{n1} & \lambda_2\mathbf{X}_{n2} & \dots & \lambda_n\mathbf{X}_{nn} \end{bmatrix} = [\lambda_i\mathbf{X}_{ij}]$$

(no summation over i)

$$= (\mathbf{A}\mathbf{X}_1, \mathbf{A}\mathbf{X}_2, \dots, \mathbf{A}\mathbf{X}_n) \quad (\text{expressing matrix as vectors})$$

$$= \mathbf{A} (\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n)$$

$$= \mathbf{A}\mathbf{E}.$$

If $\mathbf{E}$ be a non-singular matrix, then premultiplying by $\mathbf{E}^{-1}$, we get

$$\mathbf{E}^{-1}\mathbf{A}\mathbf{E} = \mathbf{D}.$$

Thus premultiplying $\mathbf{A}\mathbf{E}^{-1}$ by $\mathbf{A}$ and post-multiplying by $\mathbf{E}$, we get diagonal matrix whose diagonal elements are the eigen values. The process is called the *diagonalization of the matrix A*.

A method of diagonalization in practice. Writing the characteristic equation for given matrix, the characteristic roots can be determined. If $\mathbf{A}$ be a n -square matrix and its characteristic roots are $\lambda_1, \lambda_2, \dots, \lambda_n$, then the diagonal matrix of

$$\mathbf{A} = \begin{bmatrix} \lambda_1 & 0 & 0 & \dots & 0 \\ 0 & \lambda_2 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \lambda_n \end{bmatrix}$$

$$\text{e.g., if } \mathbf{A} = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{then } |\mathbf{A} - \lambda\mathbf{I}| = \begin{vmatrix} \cos \theta - \lambda & -\sin \theta & 0 \\ \sin \theta & \cos \theta - \lambda & 0 \\ 0 & 0 & 1 - \lambda \end{vmatrix} = 0$$

$$\text{i.e., } (1 - \lambda)(1 - 2\lambda \cos \theta + \lambda^2) = 0$$

$$\text{Characteristic roots are } 1, \frac{2 \cos \theta \pm \sqrt{4 \cos^2 \theta - 4}}{2}$$

$$\text{i.e., } 1, \cos \theta \pm i \sin \theta$$

$$\text{i.e., } 1, e^{\pm i\theta}$$

$$\therefore \lambda_1 = e^{i\theta}, \lambda_2 = e^{-i\theta}, \lambda_3 = 1 \quad (\text{say})$$

$$\text{Hence the diagonal matrix is } \begin{bmatrix} e^{i\theta} & 0 & 0 \\ 0 & e^{-i\theta} & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Note. Besides this method we can diagonalize a square matrix

(i) by orthogonally similar matrices

(ii) by unitarily similar matrices

but the above method is rather convenient in practice.

THEOREM. The necessary and sufficient condition for an n -rowed matrix $\mathbf{A}$ to be similar to a diagonal matrix is that the set of eigen vectors of $\mathbf{A}$ includes a set of n linearly independent vectors.

To prove that the condition is necessary. There exists non-singular matrix $\mathbf{E}$, such that

$$\mathbf{E}^{-1}\mathbf{A}\mathbf{E} = \mathbf{D} = \text{diag. } (\lambda_1, \lambda_2, \dots, \lambda_n).$$

Pre-multiplying by $\mathbf{E}$, we get

$$\mathbf{A}\mathbf{E} = \mathbf{E}\mathbf{D}, \text{ since } \mathbf{E}\mathbf{E}^{-1} = \mathbf{I}.$$

Suppose that $\mathbf{E} = [\mathbf{X}_1 \mathbf{X}_2 \dots \mathbf{X}_n]$.

$$\therefore \mathbf{A} [\mathbf{X}_1 \mathbf{X}_2 \dots \mathbf{X}_n] = [\mathbf{X}_1 \mathbf{X}_2 \dots \mathbf{X}_n] \text{diag. } (\lambda_1, \lambda_2, \dots, \lambda_n)$$

$$[\mathbf{A}\mathbf{X}_1 \mathbf{A}\mathbf{X}_2 \dots \mathbf{A}\mathbf{X}_n] = [\lambda_1\mathbf{X}_1 \lambda_2\mathbf{X}_2 \dots \lambda_n\mathbf{X}_n].$$

This is equivalent to

$$\mathbf{A}\mathbf{X}_1 = \lambda_1\mathbf{X}_1, \mathbf{A}\mathbf{X}_2 = \lambda_2\mathbf{X}_2, \dots, \mathbf{A}\mathbf{X}_n = \lambda_n\mathbf{X}_n.$$

that $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n$ are n eigen vectors of $\mathbf{A}$ whose corresponding eigen values are $\lambda_1, \lambda_2, \dots, \lambda_n$. As these vectors constitute the column of a non-singular matrix, there exists a linearly independent set of n eigen vectors.

To prove that the condition is sufficient. Let $\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \dots, \mathbf{X}_n$ be a linearly independent set of n eigen vectors of the matrix $\mathbf{A}$ and let $\lambda_1, \lambda_2, \dots, \lambda_n$ be the corresponding eigen values. Then

$$\mathbf{A}\mathbf{X}_1 = \lambda_1\mathbf{X}_1, \mathbf{A}\mathbf{X}_2 = \lambda_2\mathbf{X}_2, \dots, \mathbf{A}\mathbf{X}_n = \lambda_n\mathbf{X}_n. \quad \dots (1)$$

If we write $\mathbf{E} = [\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n]$,

Also it is apparent that the pre-factor row matrix is the product of the pre-factor row matrix with components $x_1, x_2, \dots, x_n$ and the post-factor matrix A . Hence

$$\sum_{j=1}^n \sum_{i=1}^n a_{ij} x_i x_j = \mathbf{X}' \mathbf{A} \mathbf{X}.$$

For example,

$$ax^2 + 2hxy + by^2 = [xy] \begin{bmatrix} a & h \\ h & b \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}.$$

Congruence of quadratic forms and matrices. An n -rowed square matrix B is said to be congruent to another n -rowed square matrix A , if there exists a non-singular matrix P , such that

$$B = P'AP.$$

Properties of congruence of matrices

(i) **Reflectivity.** Every matrix A is congruent to itself as

$$A = \mathbf{I} \mathbf{A} \mathbf{I} = \mathbf{I}' \mathbf{A} \mathbf{I}.$$

(ii) **Symmetry.** If A is congruent to B , then B is also congruent to A .

If $A = P'BP$, then

$$B = (P')^{-1} A P^{-1} = (P^{-1})' A P^{-1}.$$

(iii) **Transitivity.** If A is congruent to B and B to C then A is also congruent to C .

If $A = P'BP$, $B = Q'CQ$, then

$$A = P'Q'CQP = (QP)' C (QP).$$

(iv) **The ranges of values of two congruent quadratic forms are the same.**

If the two congruent quadratic forms be $\mathbf{X}' \mathbf{A} \mathbf{X}$ and $\mathbf{Y}' \mathbf{B} \mathbf{Y}$, then there exists a non-singular matrix P , such that

$$B = P'AP.$$

To show that the sets of values assumed by the two forms are the same when the vectors X and Y range over a field, let us write $X = PY$ so that $Y = P^{-1}X$.

Suppose that $X_1' \mathbf{A} X_1$ is a value of the first form for any vector X_1 and that $P^{-1}X_1 = Y_1$; then

$$\begin{aligned} Y_1' \mathbf{B} Y_1 &= (P^{-1} X_1)' P' A P (P^{-1} X_1) \\ &= X_1' (P')^{-1} P' A P P^{-1} X_1 \\ &= X_1' \mathbf{A} X_1, \text{ since } PP^{-1} = \mathbf{I} \text{ etc.} \end{aligned}$$

Congruent transformations. A pair of elementary transformations, one row and the other column, such that each of the corresponding elementary matrices is the transpose of the other, may be called an elementary congruent transformation. It may be of the following types:

(i) Interchange of the i th and j th rows as well as of the i th and j th columns as $E_{ij} \mathbf{A} E_{ij}$ for $E_{ij}' = E_{ij}$.

(ii) Product of the i th row as well as the i th column by a non-zero scalar c as $E_i(c) \mathbf{A} E_i(c)$ for $E_i'(c) = E_i(c)$.

(iii) Addition of k times the j th row to the i th row and also the addition of k times of the j th column to the i th column as $F_{ij}(k) \mathbf{A} F_{ij}(k)$ for $E_{ij}(k) = E_{ij}(k)$.

EXAMPLE. Reduce the following symmetric matrix to a diagonal form and interpret the result in terms of quadratic forms

$$A = \begin{bmatrix} 3 & 2 & -1 \\ 2 & 2 & 3 \\ -1 & 3 & 1 \end{bmatrix}.$$

Let us write $A = \mathbf{I}' \mathbf{A} \mathbf{I}$.

$$\text{i.e., } \begin{bmatrix} 3 & 2 & -1 \\ 2 & 2 & 3 \\ -1 & 3 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mathbf{A} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

In order to perform the elementary congruent transformations the prefactor and the postfactors of A by the row and column parts can be subjected in two steps:

(i) adding $\frac{1}{3}$ times the first row to the third and $(-\frac{2}{3})$ times the first row to the second row in L.H.S. matrix as well as in the prefactor of A ; and then

(ii) adding $\frac{1}{3}$ times of the first column to the third and $(-\frac{2}{3})$ times the first column to the second column in L.H.S. matrix as well as in the postfactor of A .

We thus have

$$\begin{bmatrix} 3 & 0 & 0 \\ 0 & \frac{2}{3} & -\frac{1}{3} \\ 0 & -\frac{1}{3} & \frac{2}{3} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -\frac{2}{3} & 1 & 0 \\ \frac{1}{3} & 0 & 1 \end{bmatrix} \mathbf{A} \begin{bmatrix} 1 & -\frac{2}{3} & \frac{1}{3} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Again performing the elementary congruent transformations by adding $(-\frac{1}{2})$ times the second row to the third row in L.H.S. matrix as well as in the pre-factor of A and then adding $(-\frac{1}{2})$ times the second column to the third column in L.H.S. matrix as well as in post-factor of A , we have

$$\begin{bmatrix} 3 & 0 & 0 \\ 0 & \frac{2}{3} & 0 \\ 0 & 0 & -\frac{1}{6} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -\frac{2}{3} & 1 & 0 \\ 4 & -\frac{1}{2} & 1 \end{bmatrix} \mathbf{A} \begin{bmatrix} 1 & -\frac{2}{3} & 4 \\ 0 & 1 & -\frac{1}{2} \\ 0 & 0 & 1 \end{bmatrix}. \quad \dots (1)$$

Thus we get the diagonal matrix

$$B = \text{diag} \left(3, \frac{2}{3}, -\frac{1}{6} \right), \quad \dots (2)$$

which is congruent to the given symmetric matrix.

From (1), we have

$$P = \begin{bmatrix} 1 & -\frac{2}{3} & 4 \\ 0 & 1 & -\frac{1}{2} \\ 0 & 0 & 1 \end{bmatrix}$$

so that $X = PY$ gives

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 & -\frac{2}{3} & 4 \\ 0 & 1 & -\frac{1}{2} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

$$= \begin{bmatrix} y_1 & -\frac{2}{3}y_2 & 4y_3 \\ 0 & y_2 & -\frac{1}{2}y_3 \\ 0 & 0 & y_3 \end{bmatrix}$$

which is equivalent to

$$\begin{aligned} x_1 &= y_1 - \frac{2}{3}y_2 + 4y_3 \\ x_2 &= y_2 - \frac{1}{2}y_3 \\ x_3 &= y_3 \end{aligned} \quad \dots (3)$$

This follows that the linear transformation (3) transforms the quadratic form

$$\begin{aligned} \mathbf{X}'\mathbf{A}\mathbf{X} &= [x_1 x_2 x_3] \begin{bmatrix} 3 & 2 & -1 \\ 2 & 2 & 3 \\ -1 & 3 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \\ &= [x_1 x_2 x_3] \begin{bmatrix} 3x_1 + 2x_2 - x_3 \\ 2x_1 + 2x_2 + 3x_3 \\ -x_1 + 3x_2 + x_3 \end{bmatrix} \\ &= x_1(3x_1 + 2x_2 - x_3) + x_2(2x_1 + 2x_2 + 3x_3) \\ &\quad + x_3(-x_1 + 3x_2 + x_3) \\ &= 3x_1^2 + 2x_2^2 + x_3^2 + 4x_1x_2 + 6x_2x_3 - 2x_3x_1 \end{aligned}$$

to the diagonal form

$$\mathbf{Y}'\mathbf{P}'\mathbf{A}\mathbf{P}\mathbf{Y} = 3y_1^2 + \frac{2}{3}y_2^2 - \frac{1}{6}y_3^2 \text{ by (2).}$$

Note. Corresponding to every quadratic form $\mathbf{X}'\mathbf{A}\mathbf{X}$ there exists a non-singular linear transformation $\mathbf{X} = \mathbf{P}\mathbf{Y}$, such that the form transforms to a sum of r square terms

$$d_1y_1^2 + d_2y_2^2 + \dots + d_ry_r^2$$

where $d_1, d_2, \dots, d_r$ are scalars in the diagonal matrix and r is the rank of the matrix $\mathbf{A}$.

Index and signature. $\mathbf{X}'\mathbf{A}\mathbf{X}$ be any real quadratic form of rank r , then there exists a real non-singular linear transformation say $\mathbf{X} = \mathbf{P}\mathbf{Y}$ which transforms $\mathbf{X}'\mathbf{A}\mathbf{X}$ to

$\mathbf{Y}'\mathbf{P}'\mathbf{A}\mathbf{P}\mathbf{Y} = y_1^2 + y_2^2 + \dots + y_s^2 - y_{s+1}^2 - y_{s+2}^2 - \dots - y_r^2$, i.e., a real quadratic form is expressed as a sum and difference of the squares of the new variables, then the number of positive squares i.e., s is called the *index* of the form and the difference $s - (r - s) = 2s - r$ is known as the *signature* of the form (canonical).

Definite, Semi-definite and Indefinite-real quadratic forms

If $\mathbf{X}'\mathbf{A}\mathbf{X}$ be a real quadratic form in n variables of rank r and index s , then

If $r = n$, $s = n$, the form is said to be *positive definite* as $y_1^2 + \dots + y_n^2$

if $r = n$, $s = 0$, *negative definite* as $-y_1^2 - \dots - y_n^2$

if $r < n$, $s = r$, *positive semi-definite* as $y_1^2 + \dots + y_r^2$;
and if $r < n$, $s = 0$, *negative semi-definite* as $-y_1^2 - \dots - y_r^2$

In any other case the form is called *indefinite*.

A real symmetric matrix $\mathbf{A}$ is said to be definite, semi-definite or indefinite according as the corresponding real quadratic form $\mathbf{X}'\mathbf{A}\mathbf{X}$ is definite, semi-definite or indefinite.

For a definite n -rowed matrix $\mathbf{A}$, there exists a real non-singular matrix $\mathbf{P}$ such that $\mathbf{P}'\mathbf{A}\mathbf{P} = \mathbf{I}_n$ or $-\mathbf{I}_n$ according as $\mathbf{A}$ is positive or negative definite.

For a semi-definite n -rowed matrix $\mathbf{A}$, there exists a real non-singular matrix $\mathbf{P}$ such that $\mathbf{P}'\mathbf{A}\mathbf{P} = \begin{bmatrix} \mathbf{I}_r & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{bmatrix}$ or $\begin{bmatrix} -\mathbf{I}_r & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{bmatrix}$ according as $\mathbf{A}$ is positive or negative definite.

EXAMPLE. Determine the following form as definite, semi-definite or indefinite.

$$2x_1^2 + 2x_2^2 + 3x_3^2 - 4x_2x_3 - 4x_3x_1 + 2x_1x_2.$$

Let $\mathbf{A}$ be the real symmetric matrix corresponding to the given real quadratic form, i.e.,

$$\begin{aligned} \mathbf{X}'\mathbf{A}\mathbf{X} &= 2x_1^2 + 2x_2^2 + 3x_3^2 - 4x_2x_3 - 4x_3x_1 + 2x_1x_2 \\ &= x_1(2x_1 + x_2 - 2x_3) + x_2(x_1 + 2x_2 - 2x_3) \\ &\quad + x_3(-2x_1 - 2x_2 + 3x_3) \end{aligned}$$

$$= [x_1 x_2 x_3] \begin{bmatrix} 2x_1 & 2x_2 & -2x_3 \\ x_1 & 2x_2 & -2x_3 \\ -2x_1 & -2x_2 & 3x_3 \end{bmatrix}$$

$$= [x_1 x_2 x_3] \begin{bmatrix} 2 & 1 & -2 \\ 1 & 2 & -2 \\ -2 & -2 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\text{It is clear that } \mathbf{A} = \begin{bmatrix} 2 & 1 & -2 \\ 1 & 2 & -2 \\ -2 & -2 & 3 \end{bmatrix},$$

We write

$$\mathbf{A} = \mathbf{I}'\mathbf{A}\mathbf{I},$$

$$\begin{bmatrix} 2 & 1 & -2 \\ 1 & 2 & -2 \\ -2 & -2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mathbf{A} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Performing the elementary congruent transformations $R_1(=\frac{1}{2}) + R_2$; $R_1 + R_3$; $C_1(-\frac{1}{2}) + C_2$, $C_1 + C_3$, we get

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & \frac{3}{2} & -1 \\ 0 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -\frac{1}{2} & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \mathbf{A} \begin{bmatrix} 1 & -\frac{1}{2} & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Again performing the elementary congruent transformations, $-\frac{2}{3}R_2 + R_3$; $-\frac{2}{3}C_2 + C_3$,

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & \frac{2}{3} & 0 \\ 0 & 0 & \frac{4}{3} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -\frac{1}{3} & 1 & 0 \\ \frac{2}{3} & -\frac{2}{3} & 1 \end{bmatrix} A \begin{bmatrix} 1 & -\frac{1}{3} & \frac{4}{3} \\ 0 & 1 & -\frac{2}{3} \\ 0 & 0 & 1 \end{bmatrix}$$

so that the linear transformation $X=PY$, i.e.,

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 & -\frac{1}{3} & \frac{4}{3} \\ 0 & 1 & -\frac{2}{3} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

i.e.,

$$\begin{cases} x_1 = y_1 - \frac{1}{3}y_2 + \frac{4}{3}y_3 \\ x_2 = y_2 - \frac{2}{3}y_3 \\ x_3 = y_3 \end{cases}$$

transforms the given form to the diagonal form

$$2y_1^2 + \frac{2}{3}y_2^2 + \frac{4}{3}y_3^2,$$

so that the rank is 3 and also the index is 3.

Hence the form is positive definite.

2.20. DIFFERENTIATION AND INTEGRATION OF MATRICES

Limit of a matrix. The limit of a matrix $A(t)$ (each element of matrix being taken as function of t) as $t \rightarrow t_0$ is a matrix $A(t_0)$ i.e.,

$$\lim_{t \rightarrow t_0} A(t) = A(t_0).$$

The sequence $\{A(t)\}$ converges to A if each element of $A(t)$ converges as $t \rightarrow \infty$ to the corresponding element of A .

Derivative of a matrix. The derivative of a square matrix $A(t)$ of order n with respect to an independent variable t is denoted by

$$\frac{d}{dt} A(t) \text{ or by } A' \text{ and is defined by}$$

$$A' = \frac{d}{dt} A(t) = \begin{bmatrix} \frac{d}{dt} a_{11} & \frac{d}{dt} a_{12} & \dots & \frac{d}{dt} a_{1n} \\ \frac{d}{dt} a_{21} & \frac{d}{dt} a_{22} & \dots & \frac{d}{dt} a_{2n} \\ \dots & \dots & \dots & \dots \\ \frac{d}{dt} a_{n1} & \frac{d}{dt} a_{n2} & \dots & \frac{d}{dt} a_{nn} \end{bmatrix} = \left[\frac{d}{dt} a_{ij}(t) \right]$$

provided all the elements $a_{ij}(t)$ are differentiable.

In general the n th derivative of $A(t)$ is defined by

$$\frac{d^n A(t)}{dt^n} = \left[\frac{d^n a_{ij}(t)}{dt^n} \right]$$

It is easy to show that

$$\frac{d}{dt} [A(t) B(t)] = A(t) \frac{dB(t)}{dt} + \frac{dA(t)}{dt} B(t) = A(t) B'(t) + A'(t) B(t)$$

and $\frac{d}{dt} |A| = \sum_{i=1}^n |A_i(t)|$ where $A_i(t)$ is obtained from $A(t)$

by differentiating i th row of $|A(t)|$ and n is the order of A .

Integral of a matrix. If $A(t) = [a_{ij}(t)]$, then

$$\int_b^t A(t) dt = \left[\int_b^t a_{ij}(t) dt \right]$$

Power series. If there is a square matrix with all eigen values less than 1 in absolute value, then the series $a_0 I + a_1 A + a_2 A^2 + \dots$ is convergent. The sum of the series is denoted by $f(A)$. Following are few convergent series for every A .

$$e^A = I + A + \frac{1}{2!} A^2 + \dots$$

$$\cos A = I - \frac{1}{2!} A^2 + \frac{1}{4!} A^4 - \dots$$

$$\sin A = A - \frac{1}{3!} A^3 + \frac{1}{5!} A^5 - \dots$$

and $(I - A)^{-1} = I + A + A^2 + \dots$ converges only for all eigen values of A which are less than 1 in absolute value.

Problem 58. For a matrix A prove that if

$$e^{tA} = I + tA + \frac{t^2}{2!} A^2 + \frac{t^3}{3!} A^3 + \dots, \text{ then}$$

$$\frac{d}{dt} e^{tA} = A e^{tA}.$$

$$\begin{aligned} \text{We have } \left[\frac{d}{dt} e^{tA} \right]_{ij} &= \frac{d}{dt} \left[(I)_{ij} + t(A)_{ij} + \frac{t^2}{2!} (A^2)_{ij} + \dots \right] \\ &= [(A)_{ij} + t(A^2)_{ij} + \dots] = [A + tA^2 + \dots]_{ij} \\ &= \left[A(I + tA + \frac{t^2}{2!} A^2 + \dots) \right]_{ij} = (A e^{tA})_{ij} \end{aligned}$$

Problem 59. Show that $e^A = e^x \begin{bmatrix} \cos hx & \sin hx \\ \sin hx & \cos hx \end{bmatrix}$ when $A = \begin{bmatrix} x & x \\ x & x \end{bmatrix}$

$$\text{we have } e^A \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = e^x \begin{bmatrix} \cos hx & \sin hx \\ \sin hx & \cos hx \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = e^x \begin{bmatrix} e^x & e^x \\ e^x & e^x \end{bmatrix} = e^{2x} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$\text{Again } e^A = I + A + \frac{1}{2!} A^2 + \frac{1}{3!} A^3 + \dots = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + x \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$+ \frac{1}{2!} \cdot 2x^2 \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} + \dots$$

$$\therefore e^A \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} + x \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix} + \frac{1}{2!} 2x^2 \begin{bmatrix} 4 & 4 \\ 4 & 4 \end{bmatrix} + \dots$$

$$= \left[1 + 2x + 4 \frac{x^2}{2!} + \dots \right] \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = e^{2x} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

Hence the proposition.

2.21. SOLUTION OF A SYSTEM OF LINEAR DIFFERENTIAL EQUATIONS USING DIFFERENTIATION OF MATRICES

[A] Linear differential equations of first order

Let the system be

$$\left. \begin{aligned} \frac{dx_1}{dt} &= a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n + b_1(t) \\ \frac{dx_2}{dt} &= a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n + b_2(t) \\ &\dots\dots\dots \\ \frac{dx_n}{dt} &= a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n + b_n(t) \end{aligned} \right\} \dots (1)$$

which are equivalent to $\frac{d\mathbf{X}}{dt} = \mathbf{A}\mathbf{X} + \mathbf{B}$ $\dots (2)$

Making substitution $\mathbf{X}(t) = \mathbf{P}\mathbf{Y}(t)$, (2) yields $\dots (3)$

$$\frac{d}{dt}[\mathbf{P}\mathbf{Y}(t)] = \mathbf{P} \frac{d\mathbf{Y}}{dt} = \mathbf{A}\mathbf{P}\mathbf{Y} + \mathbf{B}(t) \quad \therefore \frac{d\mathbf{P}}{dt} = 0$$

But $\mathbf{A}$ is similar to $\mathbf{P}$ if $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D}$, $\mathbf{D}$ being Diagonal matrix

$$\therefore \frac{d\mathbf{Y}}{dt} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}\mathbf{Y} + \mathbf{P}^{-1}\mathbf{B}(t) = \mathbf{D}\mathbf{Y} + \mathbf{P}^{-1}\mathbf{B}(t) \dots (4)$$

which is equivalent to

$$\frac{dy_1}{dt} = \lambda_1 y_1 + c_1(t), \frac{dy_2}{dt} = \lambda_2 y_2 + c_2(t), \dots, \frac{dy_n}{dt} = \lambda_n y_n + c_n(t) \dots (5)$$

Their solution being given by

$$\left[y_j(t) e^{-\lambda_j t} \right]_0^t = \int_0^t c_j(t) e^{-\lambda_j t} dt \dots (6)$$

or $y_j(t) = e^{\lambda_j t} \left[y_0(t) + \int_0^t c_j(t) e^{-\lambda_j t} dt \right]$ for $j=1, 2, 3, \dots, n$.

$\therefore \mathbf{Y}(t)$ and hence $\mathbf{X}(t)$ can be determined.

[B] Linear differential equations of n th order

Regarding a linear differential equation of n th order as equivalent to n linear differential equations each of order one, and writing

$$\frac{d^n}{dt^n} x(t) + a_1 \frac{d^{n-1}}{dt^{n-1}} x(t) + \dots + a_n x(t) = b(t) \dots (7)$$

we have on making substitutions $x_1(t) = x(t)$, $x_2(t) = \frac{d}{dt} x(t)$, ...,

$$x_n(t) = \frac{d^{n-1}}{dt^{n-1}} x(t)$$

$$Dx_1(t) = x_2(t), Dx_2(t) = x_3(t), \dots, Dx_{n-1}(t) = x_n(t) \text{ and}$$

$$Dx_n(t) = -a_1 x_n(t) - a_2 x_{n-1}(t) \dots - a_n x_1(t) + b(t) \text{ where } D \equiv \frac{d}{dt}$$

the system is equivalent to Matrix-equation

$$D\mathbf{X}(t) = \begin{bmatrix} Dx_1(t) \\ Dx_2(t) \\ \dots \\ Dx_n(t) \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ -a_n & -a_{n-1} & -a_{n-2} & \dots & -a_1 \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \\ \dots \\ x_n(t) \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \dots \\ b(t) \end{bmatrix}$$

$$= \mathbf{A}\mathbf{X}(t) + \mathbf{B}(t).$$

where we have used the concept of diagonalization.

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 60. Write short note on Hermitian Matrices.

(Vikram 1967)

See §2.9, [10].

Problem 61. What is a reciprocal matrix? Show that a non-singular matrix $\mathbf{A} = [a_{ij}]$ always possesses a reciprocal.

Solve by the method of matrix theory, the linear equations.

$$x_1 + 2x_3 = 1$$

$$3x_1 - x_2 + x_3 = 2$$

$$4x_2 + 5x_3 = -1$$

(Vikram 1967)

See 2.9 [13] Theorem 1 on it for the first part.

Now, Matrix form of the given system is $\mathbf{A}\mathbf{X} = \mathbf{B}$ i.e., where

$$\begin{bmatrix} 1 & 0 & 2 \\ 3 & -1 & 1 \\ 0 & 4 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} \text{ where } \mathbf{A} = \begin{bmatrix} 1 & 0 & 2 \\ 3 & -1 & 1 \\ 0 & 4 & 5 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} \text{ and } \mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\text{By } R_{21}(-3), \begin{bmatrix} 1 & 0 & 2 \\ 0 & -1 & -5 \\ 0 & 4 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}$$

$$\text{By } R_{32}(1), \begin{bmatrix} 1 & 0 & 2 \\ 0 & -1 & -5 \\ 0 & 3 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix}$$

which are equivalent to $x_1 + 2x_3 = 1$

$$-x_2 - 5x_3 = -1$$

$$3x_2 = -2$$

Giving $x_2 = -\frac{2}{3}$, $x_3 = \frac{1}{3}$ and $x_1 = \frac{1}{3}$.

Problem 62. (a) Explain with examples what you understand by a unitary matrix and a Hermitian matrix.

(b) Find the eigen values and the eigen vectors of the matrix

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 3 & 6 \end{bmatrix}$$

(Agra, 1969)

For (a) see §2.9 [14] and §2.9 [10]

(b) Denoting the given matrix by $\mathbf{A}$, its characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \text{ i.e., } \begin{vmatrix} 1-\lambda & 1 & 1 \\ 1 & 2-\lambda & 3 \\ 1 & 3 & 6-\lambda \end{vmatrix} = 0 \text{ or } \lambda^3 - 9\lambda^2 + 9\lambda - 1 = 0$$

$$\text{or } (\lambda-1)(\lambda^2 - 8\lambda + 1) = 0$$

Giving $\lambda = 1, 4 \pm \sqrt{15}$.

New proceeding just like in problem 45, eigen vectors can be determined.

$$\text{Ans. } \begin{bmatrix} k \\ -k \\ -k/2 \end{bmatrix} \begin{bmatrix} 5 + \sqrt{15} \\ 10 + 3\sqrt{15} \\ k' \end{bmatrix} k' \text{ etc.}$$

$$\begin{bmatrix} 5 + \sqrt{15} \\ \sqrt{15} + 1 \end{bmatrix} k' \text{ etc.}$$

Problem 63. Find the characteristic equation of the matrix $\begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 1 \\ 3 & 1 & 1 \end{bmatrix}$

(Vikram, 1988)

and verify Cayley Hamilton equation for it.

Denoting the given matrix by A , its characteristic equation is

$$|A - \lambda I| = 0 \text{ i.e., } \begin{vmatrix} 1-\lambda & 2 & 3 \\ 2 & -1-\lambda & 1 \\ 3 & 1 & 1-\lambda \end{vmatrix} = 0 \text{ i.e., } \lambda^3 - \lambda^2 - 15\lambda - 15 = 0$$

and verify that $A^3 - A^2 - 15A - 15I = 0$.

Problem 64. (a) Define Hermitian and orthogonal matrices. Give one example of each type.

(b) Find the matrices C and C^{-1} required to reduce the matrix

$$A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} \text{ to the diagonal form by the transformation } C^{-1}AC$$

(Agra, 1971)

Problem 65. Show that the six matrices

$$E = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, B = \frac{1}{2} \begin{bmatrix} -1 & \sqrt{3} \\ \sqrt{3} & 1 \end{bmatrix}, C = \frac{1}{2} \begin{bmatrix} -1 & -\sqrt{3} \\ \sqrt{3} & 1 \end{bmatrix}$$

$$D = \frac{1}{2} \begin{bmatrix} -1 & -\sqrt{3} \\ \sqrt{3} & -1 \end{bmatrix}, F = \frac{1}{2} \begin{bmatrix} -1 & \sqrt{3} \\ \sqrt{3} & -1 \end{bmatrix}$$

satisfy the relations

$$A^2 = B^2 = C^2 = E$$

$$AB = D \text{ and } AC = BA = E.$$

Problem 66. Define eigen values and eigen vectors of a matrix. Prove that the eigen values of an orthogonal matrix are unimodular.

Obtain the eigen values and eigen vectors of the matrix $A = \begin{bmatrix} 8 & -12 & 3 \\ 15 & -25 & 11 \\ 24 & -42 & 19 \end{bmatrix}$

(Agra, 1971)

Problem 67. If the Hermitian matrices $\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, $\sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$, $\sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$

satisfy the equations

$$\sigma_1^2 = \sigma_2^2 = \sigma_3^2 = I, \sigma_1\sigma_2 = i\sigma_3, \sigma_2\sigma_3 = i\sigma_1, \sigma_3\sigma_1 = i\sigma_2$$

$$\sigma_1\sigma_2 - \sigma_2\sigma_1 = 2i\sigma_3, \sigma_1\sigma_3 + \sigma_3\sigma_1 = 0$$

then find the eigen values of the matrices $\sigma_1, \sigma_2, \sigma_3$.

Problem 68. Define the trace of a matrix. Prove that two matrices A and B have the same trace if $B = T^{-1}AT$ where T is a non-singular matrix.

The eigen values of a 3×3 matrix H are equal to the three cube roots of unity. Prove that $H^3 = I$, where I is a unit matrix of order 3. (Agra, 1971)

Problem 69. Show that all the matrices of Problem 65 are unitary.

Problem 70. Find the reciprocal of each of the matrices of Problem 65 and verify that

$$D^{-1} = B^{-1}A^{-1}.$$

Problem 71. Find the eigen values and normalised eigen vectors of the matrix

$$\begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

(Agra, 1971)

Problem 72. [A Significant property of diagonal Matrices]

Prove that in case of diagonal Matrices, the commutative law for multiplication holds good.

If A and B be two diagonal (square) matrices of the same order, then we have to prove that

$$AB = BA$$

Since a diagonal matrix is such that its all the elements except those in the principal diagonal are zero, we can take

$$A = [a_{ij} \delta_{ij}] \text{ and } B = [b_{ij} \delta_{ij}]$$

where $\delta_{ij} = 0$ for $i \neq j$

$$= 1 \text{ for } i = j$$

$$\therefore [AB]_{ik} = \sum_j [A]_{ij} [B]_{jk} = \sum_j a_{ij} \delta_{ij} b_{jk} \delta_{jk}$$

$$= a_{ii} b_{ii} \delta_{ii} \text{ for } j = i \text{ and no summation over } i$$

$$= b_{ii} a_{ii} \delta_{ii}$$

$$= [BA]_{ii} \text{ and zero for } k \neq i$$

which follows that AB is also a diagonal matrix and $AB = BA$.

Problem 73. If $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, find M^{-1} , transpose of M , Hermitian Conjugate of M and the eigen values of M . (Agra, 1975)

Problem 74. (a) Find eigen value and corresponding orthonormal vector of the following matrix :

$$\begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

(b) Diagonalize the following matrix :

$$A = \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix}$$

(Rohilkhand, 1976)

Problem 75. Show that eigen values of a Hermitian matrix are all real and its eigen vectors corresponding to two distinct eigen values are orthogonal.

What do you mean by diagonalization of a matrix? Show that the necessary and sufficient condition for the reduction of two matrices to the diagonal form by the same transformation is that they commute. (Rohilkhand, 1977)

See theorems 1 and 5 of § 2.18.

See § 2.18 after Problem 54 and also see Problem 72.

CHAPTER 3

TENSORS

3.1. INTRODUCTION

It is an established fact that a natural law gives a relation between different physical entities and its mathematical formulation is nothing but a relation between the sets of numbers representing those entities. Tensor Analysis forms that part of study which is rather suitable for the mathematical formulation of natural laws in forms which are invariant with respect to underlying frames of reference. In brief tensors are quantities obeying certain transformation laws. In wider sense a tensor formulation is very compact and good deal of clarity in its use. The tensor formulation was originated by G. Ricci and it became rather popular when Albert Einstein used it as a natural tool for the description of his general theory of relativity.

Actually tensor analysis is the generalization of vector analysis as is evident by considering a vector function $f(\mathbf{r})$ of a vector $\mathbf{r}$. This vector function is continuous at $\mathbf{r}=\mathbf{r}_0$ if,

$$\lim_{\mathbf{r} \rightarrow \mathbf{r}_0} f(\mathbf{r}) = f(\mathbf{r}_0)$$

and it is linear, if

$$f(\mathbf{r} + \mathbf{s}) = f(\mathbf{r}) + f(\mathbf{s}) \quad \dots (1)$$

and

$$f(\lambda \mathbf{r}) = \lambda f(\mathbf{r}) \quad \dots (2)$$

for arbitrary values of $\mathbf{r}, \mathbf{s}, \lambda$.

Now we know that as linear vector function $f(\mathbf{r})$ is completely defined only if $f(\mathbf{a}_1), f(\mathbf{a}_2)$ and $f(\mathbf{a}_3)$ are given for any three non-coplanar vectors $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$. In terms of $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$ as basis if we assume that

$$\mathbf{r} = x_1 \mathbf{a}_1 + x_2 \mathbf{a}_2 + x_3 \mathbf{a}_3 \quad \dots (3)$$

then we have from (1) and (2),

$$f(\mathbf{r}) = x_1 f(\mathbf{a}_1) + x_2 f(\mathbf{a}_2) + x_3 f(\mathbf{a}_3) \quad \dots (4)$$

As such equation (3) yields

$$x_\alpha = \mathbf{r} \cdot \mathbf{a}_\alpha, \quad \alpha = 1, 2, 3 \quad \dots (5)$$

Let us put

$$f(\mathbf{a}_\alpha) = \mathbf{b}_\alpha.$$

So that

$$f(\mathbf{r}) = (\mathbf{b}_1 \mathbf{a}_1 + \mathbf{b}_2 \mathbf{a}_2 + \mathbf{b}_3 \mathbf{a}_3) \cdot \mathbf{r} \\ = \phi \cdot \mathbf{r} \text{ (say)} \quad \dots (6)$$

where the operator $\phi = \mathbf{a}_1 \mathbf{b}_1 + \mathbf{a}_2 \mathbf{b}_2 + \mathbf{a}_3 \mathbf{b}_3$ consists of nine components in three dimensional coordinate geometry and hence it is neither scalar nor vector quantity but is a new mathematical symbol called as the *dyadic*.

Suppose that there are two vectors $\mathbf{u}$ and $\mathbf{v}$ such that components of vector $\mathbf{v}$ are linear functions of the components of vector $\mathbf{u}$ defined as

$$\left. \begin{aligned} v_x &= a_{xx} u_x + a_{xy} u_y + a_{xz} u_z \\ v_y &= a_{yx} u_x + a_{yy} u_y + a_{yz} u_z \\ v_z &= a_{zx} u_x + a_{zy} u_y + a_{zz} u_z \end{aligned} \right\} \quad \dots (7)$$

In this way the vector $\mathbf{v}$ is placed in one-to-one correspondence with the vector $\mathbf{u}$. The scheme of coefficient $a_{\alpha\beta}$ has thus an independent meaning if the correspondence is such that the passage from $\mathbf{u}$ to $\mathbf{v}$ is independent of the particular coordinate system in which the vectors are resolved into components. We call the coefficients $a_{\alpha\beta}$ in this case as the coefficients of a tensor.

It is observed that the nine components as mentioned above characterise the transformation of the components of one vector into those of other. The coefficients $a_{\alpha\beta}$ in general transform u_β into one of three parts of v_α .

The equations (7) are equivalent to a single vector equation,

$$\mathbf{v} = \phi \mathbf{u} \quad \dots (8)$$

where the operator ϕ turns $\mathbf{u}$ into $\mathbf{v}$. It is rather graphically known as *Tensor*.

The essential part of a tensor operation is the array of coefficients like $a_{\alpha\beta}$, written in the form of a matrix such as

$$\phi = \begin{bmatrix} a_{xx} & a_{xy} & a_{xz} \\ a_{yx} & a_{yy} & a_{yz} \\ a_{zx} & a_{zy} & a_{zz} \end{bmatrix} \quad \dots (9)$$

As such the *dyadic* operator turns a vector $\mathbf{r}$ into the vector function $f(\mathbf{r})$ and is expressed as the sum of dyads *ab i.e.*

$$\phi = \sum_{\alpha} \mathbf{a}_{\alpha} \mathbf{b}_{\alpha}$$

In a similar way a *triadic* is expressed as the sum of the triads $\mathbf{a}_{\alpha} \mathbf{b}_{\alpha} \mathbf{c}_{\alpha}$.

Considering it as an operator that converts vector $\mathbf{r}$ into the dyadic ϕ , we may write $\phi \cdot \mathbf{r} = \sum (\mathbf{a}_{\alpha} \mathbf{b}_{\alpha} \mathbf{c}_{\alpha}) \cdot \mathbf{r}$

Similarly a *tetradic* is the sum of tetrads, $\sum \mathbf{a}_{\alpha} \mathbf{b}_{\alpha} \mathbf{c}_{\alpha} \mathbf{d}_{\alpha}$ and etc.

All such physical quantities as scalars, vectors, dyadics, triadics, tetradics etc. are collectively known as tensors of rank 1, 2, 3, 4 etc. and as such the tensors can be regarded as generalized extended form of vectors.

The examples of dyadic *i.e.* tensor of rank two are : an operator relating dielectric displacement vector with the electric vector of an electro-magnetic wave in anisotropic medium; a stress tensor relating stress and strain in anisotropic medium in which case a component of stress $\mathbf{T}$ is a function of every component of strain $\mathbf{S}$

$$\text{i.e.} \quad T_{\alpha} = \sum_{\beta=1}^3 a_{\alpha\beta} S_{\beta} \text{ or } \mathbf{T} = \psi \mathbf{S} \quad \dots (10)$$

where ψ is a nine coefficient operator in three dimensional space.

Note. The dyadic or tensor of rank two is also known as *Stress Tensor*.

An Explanatory Note on Dyads and Dyadics

We know that the gradient of a vector $\mathbf{f}$ such as

$$\nabla \mathbf{f} = i\mathbf{f}_x + j\mathbf{f}_y + k\mathbf{f}_z \quad \dots (11)$$

is meaningless as it consists of sum of three ordered pairs of vectors, but we sometimes take it to define as *dyadic* and the ordered vector pairs as *dyads*. In (11) we regard $\nabla \mathbf{f}$ as an operator setting up a one-one correspondence between directions $\mathbf{e}$ at a point and its directional

$$\text{derivative } \frac{d\mathbf{f}}{ds} \text{ i.e. } \frac{d\mathbf{f}}{ds} = \mathbf{e} \cdot \nabla \mathbf{f} \quad \dots (12)$$

Actually the dyadic $\nabla \mathbf{f}$ replaces an infinite number of vectors $\frac{d\mathbf{f}}{ds}$, so that any sum of dyads is called as dyadic e.g. the dyadic

$$\mathbf{D} = \mathbf{a}_1 \mathbf{b}_1 + \mathbf{a}_2 \mathbf{b}_2 + \dots + \mathbf{a}_n \mathbf{b}_n \quad \dots (13)$$

is a general dyadic in which the vectors $\mathbf{a}_i$ are known as *antecedents* and $\mathbf{b}_i$ as *Consequents*, while the dyadic

$$\mathbf{D}_c = \mathbf{b}_1 \mathbf{a}_1 + \mathbf{b}_2 \mathbf{a}_2 + \dots + \mathbf{b}_n \mathbf{a}_n \quad \dots (14)$$

is said to be the conjugate of $\mathbf{D}$, such that $\mathbf{D}$ is *symmetric* if $\mathbf{D} = \mathbf{D}_c$, and *skew* if $\mathbf{D} = -\mathbf{D}_c$.

In (11) if $\mathbf{f}$ is replaced by $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ such that

$$\mathbf{r}_x = \mathbf{i}, \mathbf{r}_y = \mathbf{j}, \mathbf{r}_z = \mathbf{k}, \text{ then } \nabla \mathbf{r} = \mathbf{i}\mathbf{i} + \mathbf{j}\mathbf{j} + \mathbf{k}\mathbf{k} = \mathbf{I} \quad \dots (15)$$

where the dyadic $\mathbf{I}$ is termed as *Idemfactor* as it transforms any vector $\mathbf{V}$ into itself i.e.,

$$\mathbf{V} \cdot \mathbf{I} = \mathbf{I} \cdot \mathbf{V} = \mathbf{V} \text{ for every } \mathbf{V}.$$

3.2. TRANSFORMATION OF COORDINATES

If we focus our attention on some point of Minkowski's four dimensional world and consider the transformation from one system of coordinates (x_1, x_2, x_3, x_4) to another system (x'_1, x'_2, x'_3, x'_4) , such that

$$x'_1 = f_1(x_1, x_2, x_3, x_4) \text{ etc.}$$

then we can solve x_1, x_2, x_3, x_4 in terms of x'_1, x'_2, x'_3, x'_4 such that

$$x_1 = \phi_1(x'_1, x'_2, x'_3, x'_4) \text{ etc.}$$

and the differentials dx_1, dx_2, dx_3, dx_4 are then transformed as

$$dx'_1 = \frac{\partial x'_1}{\partial x_1} dx_1 + \frac{\partial x'_1}{\partial x_2} dx_2 + \frac{\partial x'_1}{\partial x_3} dx_3 + \frac{\partial x'_1}{\partial x_4} dx_4 \text{ etc.}$$

or symbolically

$$dx'_\mu = \sum_{\alpha=1}^4 \frac{\partial x'_\mu}{\partial x_\alpha} dx_\alpha; (\mu=1, 2, 3, 4) \text{ etc.} \quad \dots (1)$$

3.3. THE SUMMATION CONVENTION AND KRONECKER DELTA SYMBOL

Let (x_1, x_2, x_3, x_4) and $(x_1+dx_1, x_2+dx_2, x_3+dx_3, x_4+dx_4)$ be the coordinates of two neighbouring events considered in Minkowski's four dimensional space. Then the interval ds between these two neighbouring events in any coordinate system, is given by

$$ds^2 = g_{11}dx_1^2 + g_{22}dx_2^2 + g_{33}dx_3^2 + g_{44}dx_4^2 + 2g_{12}dx_1dx_2 + 2g_{13}dx_1dx_3 + 2g_{14}dx_1dx_4 + 2g_{23}dx_2dx_3 + 2g_{24}dx_2dx_4 + 2g_{34}dx_3dx_4 \quad \dots (1)$$

where the coefficients $g_{\mu\nu}$ ($\mu, \nu=1, 2, 3, 4$) are functions of x_1, x_2, x_3, x_4 . This follows that ds^2 is some quadratic function of the difference of coordinates.

Adopting the convention that whenever a literal suffix appears twice in a term that term is to be summed for values of the suffix 1, 2, 3, 4; the equation (1) can be written as

$$ds^2 = g_{\mu\nu} dx_\mu dx_\nu \quad (\mu, \nu=1, 2, 3, 4 \text{ and } g_{\mu\nu} = g_{\nu\mu}) \quad \dots (2)$$

Since μ and ν each appear twice, the right hand side of (2) indicates

$$\text{the summation } \sum_{\mu=1}^4 \sum_{\nu=1}^4$$

Any literal suffix appearing twice in a term is said to be a *dummy suffix* and it may be changed freely to any other letter not already used in that term. Also two or more dummy suffixes can be interchanged e.g.

$$g_{\alpha\beta} \frac{\partial^2 x_\alpha}{\partial x'_\mu \partial x'_\nu} \frac{\partial x_\beta}{\partial x'_\lambda} = 2g_{\alpha\beta} \frac{\partial^2 x_\beta}{\partial x'_\mu \partial x'_\nu} \frac{\partial x_\alpha}{\partial x'_\lambda}$$

(by interchanging the dummy suffixes α and β and using $g_{\beta\alpha} = g_{\alpha\beta}$)

Illustration. To prove that

$$\frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x'_\alpha}{\partial x_\nu} = \frac{\partial x_\mu}{\partial x_\nu} = 0 \text{ if } \mu \neq \nu$$

$$= 1 \text{ if } \mu = \nu \text{ where } \alpha = 1, 2, 3, 4.$$

$$\text{Here, R.H.S.} = \frac{\partial x_\mu}{\partial x'_1} \frac{\partial x'_1}{\partial x_\nu} + \frac{\partial x_\mu}{\partial x'_2} \frac{\partial x'_2}{\partial x_\nu} + \frac{\partial x_\mu}{\partial x'_3} \frac{\partial x'_3}{\partial x_\nu} + \frac{\partial x_\mu}{\partial x'_4} \frac{\partial x'_4}{\partial x_\nu}$$

$$= \frac{\partial x_\mu}{\partial x_\nu}.$$

x_μ and x_ν being the coordinates of the same system, their variations are independent and so

$$dx_\mu = 0 \text{ when } \mu \neq \nu$$

and

$$dx_\mu = dx_\nu \text{ when } \mu = \nu$$

$$\therefore \frac{dx_\mu}{dx'_\nu} = \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x'_\alpha}{\partial x_\nu} = 0 \text{ when } \mu \neq \nu$$

$$= 1 \text{ when } \mu = \nu$$

Here the multiplier $\frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x'_\alpha}{\partial x_\mu}$ acts as a *substitution operator*.

If is rather convenient to write

$$\frac{\partial x_\mu}{\partial x_\nu} = \delta_{\mu\nu} \text{ or } \delta_\nu^\mu \text{ which is known as Kronecker delta.}$$

As such the above results can be expressed as

$$\left. \begin{aligned} \delta_\nu^\mu &= 0 & \text{if } \mu \neq \nu \\ &= 1 & \text{if } \mu = \nu \end{aligned} \right\} \dots (3)$$

Thus if $A(\mu)$ be an expression involving the suffix μ , then

$$\frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x'_\alpha}{\partial x_\nu} A(\mu) = A(\nu) \dots (4)$$

for; the summation on the left, with respect to μ gives four terms corresponding to $\mu=1, 2, 3, 4$; one of which will agree with ν . Denoting the other three values by σ, τ, ρ , the left hand side of (4) is

$$= 1 \cdot A(\nu) + 0 \cdot A(\sigma) + 0 \cdot A(\tau) + 0 \cdot A(\rho) \text{ by (3)}$$

$$= A(\nu)$$

$$\text{i.e. } \delta_\nu^\mu A(\mu) = A(\nu) \dots (5)$$

$$\text{Evidently } \delta_\rho^\mu \delta_\nu^\rho = \delta_\nu^\mu \dots (6)$$

$$\text{and } \delta_\mu^\mu = 4 \dots (7)$$

$$\text{or, in the latter case, } \delta_\mu^\mu = \delta_1^1 + \delta_2^2 + \delta_3^3 + \delta_4^4 \\ = 1 + 1 + 1 + 1 = 4. \text{ by (3)}$$

3.4. TENSORS AS CLASSIFICATION OF TRANSFORMATION LAWS

We have already mentioned in §3.2 that if we consider the transformation from one system of coordinates (x_1, x_2, x_3, x_4) to another system (x'_1, x'_2, x'_3, x'_4) , then the differentials dx_1, dx_2, dx_3, dx_4 are transformed as

$$dx'_1 = \frac{\partial x'_1}{\partial x_1} dx_1 + \frac{\partial x'_1}{\partial x_2} dx_2 + \frac{\partial x'_1}{\partial x_3} dx_3 + \frac{\partial x'_1}{\partial x_4} dx_4 \text{ etc.}$$

or in short as

$$dx'_\mu = \sum_{\alpha=1}^4 \frac{\partial x'_\mu}{\partial x_\alpha} dx_\alpha \quad \mu=1, 2, 3, 4$$

Any set of four quantities transformed in accordance with this law is said to be a **Contravariant Vector**. Thus if a coordinate system (A^1, A^2, A^3, A^4) transforms to the new coordinate system (A'^1, A'^2, A'^3, A'^4) where

$$A'^\mu = \sum_{\alpha=1}^4 \frac{\partial x'_\mu}{\partial x_\alpha} A^\alpha \dots (1)$$

Then (A^1, A^2, A^3, A^4) or briefly A^μ is a contravariant vector. Hence the upper position of the suffix (which is definitely, not an exponent) is reserved to indicate contravariant vectors.

Now, if we consider an operator ϕ such that it is an invariant function of position i.e., it has a fixed value at each point independent of the coordinate system used, then the four quantities

$$\frac{\partial \phi}{\partial x_1}, \frac{\partial \phi}{\partial x_2}, \frac{\partial \phi}{\partial x_3}, \frac{\partial \phi}{\partial x_4} \text{ are transformed as}$$

$$\frac{\partial \phi}{\partial x'_1} = \frac{\partial x_1}{\partial x'_1} \frac{\partial \phi}{\partial x_1} + \frac{\partial x_2}{\partial x'_1} \frac{\partial \phi}{\partial x_2} + \frac{\partial x_3}{\partial x'_1} \frac{\partial \phi}{\partial x_3} + \frac{\partial x_4}{\partial x'_1} \frac{\partial \phi}{\partial x_4}, \text{ etc.}$$

$$\text{or in short } \frac{\partial \phi}{\partial x'_\mu} = \sum_{\alpha=1}^4 \frac{\partial x_\alpha}{\partial x'_\mu} \frac{\partial \phi}{\partial x_\alpha} \quad (\mu=1, 2, 3, 4).$$

Any set of four quantities transformed in accordance with this law is said to be a **Covariant Vector**.

Thus if A_μ be a covariant vector, its transformation law is

$$A'_\mu = \sum_{\alpha=1}^4 \frac{\partial x_\alpha}{\partial x'_\mu} A_\alpha \dots (2)$$

where the lower position of the suffix indicates covariance.

Hence the relations (1) and (2) give the laws of transformation of vectors. If we denote by $A_{\mu\nu}$ a quantity with 16 components by assigning μ and ν the values 1, 2, 3, 4 independently, then a generalization of these laws yields quantities classified as

$$\text{Contravariant Tensors } A'^{\mu\nu} = \frac{\partial x'_\mu}{\partial x_\alpha} \frac{\partial x'_\nu}{\partial x_\beta} A^{\alpha\beta} \dots (3)$$

$$\text{Covariant Tensors } A'_{\mu\nu} = \frac{\partial x_\alpha}{\partial x'_\mu} \frac{\partial x_\beta}{\partial x'_\nu} A_{\alpha\beta} \dots (4)$$

$$\text{Mixed Tensors } A'^{\mu\nu} = \frac{\partial x_\alpha}{\partial x'_\mu} \frac{\partial x'_\nu}{\partial x_\beta} A^{\alpha\beta} \dots (5)$$

There are the tensors of *second rank*.

Similarly $A_{\mu\nu\sigma}$ has 64 components and $A_{\mu\nu\sigma\tau}$ has 256 components.

Thus a tensor of higher rank is of the type

$$A'^{\tau}_{\mu\nu\sigma} = \frac{\partial x_{\alpha}}{\partial x'_{\mu}} \frac{\partial x_{\beta}}{\partial x'_{\nu}} \frac{\partial x_{\gamma}}{\partial x'_{\sigma}} \frac{\partial x'_{\tau}}{\partial x_{\delta}} A_{\alpha\beta\gamma}^{\delta} \dots (6)$$

Note. A vector is a tensor of first rank and an invariant or scalar is a tensor of zero rank.

Rank of a Tensor. The rank of a tensor is determined by the number of suffixes or indices attached to it. As a matter of fact the rank of a tensor when raised as power to the number of dimensions, yields the number of components of the tensor and hence the components of the matrix that represents the tensor. As such a tensor of rank n in four dimensional space has 4^n component. Consequently the rank of a tensor gives the number of the mode of changes of a physical quantity when passing from one system to another system which is in rotation relative to the first. It is clear from this discussion that a quantity that does not change when the axes are rotated is a tensor of zero rank, since the number of mode of changes is then zero. These quantities named as tensor of zero rank are scalars while the tensors of rank one are vectors.

Problem 1. Explain what is meant by the rank of a tensor.

(Agra, 1972)

See the note on §3.4.

Problem 2. What is a tensor?

(Vikram, 1967)

Define as discussed in above articles or see Problem 6 discussed after.

Problem 3. Define contravariant and covariant tensors. (Agra, 1967)

Contravariant tensor. If n quantities A^{α} ($\alpha=1, 2, \dots, n$) in a coordinate system $(x_1, x_2, \dots, x_n)$ are related to n other quantities A'^{α} ($\alpha=1, 2, \dots, n$) in another coordinate system $(x'_1, x'_2, \dots, x'_n)$ by the transformation laws

$$A'^{\mu} = \frac{\partial x'_{\mu}}{\partial x_{\alpha}} A^{\alpha} \text{ (Contravariant law)}$$

on change of the coordinate x_{β} to x'_{β} according to summation convention, then A^{α} are termed as the components of a contravariant vector or a contravariant tensor of the first rank.

Covariant tensor. If n quantities A_{α} ($\alpha=1, 2, \dots, n$) in a coordinate system $(x_1, x_2, \dots, x_n)$ are related to n other quantities A'_{α} ($\alpha=1, 2, \dots, n$) in another coordinate system $(x'_1, x'_2, \dots, x'_n)$ by the transformation laws

$$A'_{\mu} = \frac{\partial x_{\alpha}}{\partial x'_{\mu}} A_{\alpha} \text{ (Covariant law)}$$

according to summation convention, then A_{α} are termed as the components of a covariant vector or a covariant tensor of the first rank.

Problem 4. Show that the velocity of a fluid at any point is a contravariant vector of rank one.

Assuming that $x_{\alpha}(t)$ is the coordinate of a moving particle with the time t , we have

$$v^{\alpha} = \frac{dx_{\alpha}}{dt}$$

as the velocity of the particle.

In transformed coordinates the components of velocity are

$$V'^{\alpha} = \frac{d}{dt} x'_{\alpha}$$

$$\text{But } v^{\alpha} = \frac{d}{dt} x_{\alpha} = \frac{\partial x'_{\alpha}}{\partial x_{\beta}} \frac{dx_{\beta}}{dt} = \frac{\partial x'_{\alpha}}{\partial x_{\beta}} v_{\beta}$$

which follows that velocity is a contravariant vector of rank one.

Problem 5. Show that the law of transformation for a contravariant vector is transitive.

$$\text{We have } A'^{\mu} = \frac{\partial x'_{\mu}}{\partial x_{\alpha}} A^{\alpha}$$

$$\text{Let } A''^{\mu} = \frac{\partial x''_{\mu}}{\partial x'_{\alpha}} A'^{\alpha}$$

$$\therefore A''^{\mu} = \frac{\partial x''_{\mu}}{\partial x'_{\beta}} A^{\beta} = \frac{\partial x''_{\mu}}{\partial x'_{\beta}} \frac{\partial x'_{\beta}}{\partial x_{\alpha}} A^{\alpha} = \frac{\partial x''_{\mu}}{\partial x_{\alpha}} A^{\alpha}$$

which shows that contravariant law is transitive.

Problem 6. Find the components of a vector in polar coordinates whose components in cartesian co-ordinates are $\dot{x}$, $\dot{y}$ and $\ddot{x}$, $\ddot{y}$.

As given, suppose that

$$x_1 = x, x_2 = y,$$

$$x'_1 = r, x'_2 = \theta$$

and

$$(i) A^1 = \dot{x}, A^2 = \dot{y},$$

$$(ii) A^1 = \ddot{x}, A^2 = \ddot{y}.$$

Then, we have to find C'^1, A'^2 .

We have the transformations

$$x = r \cos \theta, y = r \sin \theta,$$

giving

$$r^2 = x^2 + y^2 \text{ and } \theta = \tan^{-1} \frac{y}{x}$$

so that

$$\frac{\partial r}{\partial x} = \frac{x}{r}, \frac{\partial r}{\partial y} = \frac{y}{r}, \frac{\partial \theta}{\partial x} = -\frac{y}{r^2} \text{ and } \frac{\partial \theta}{\partial y} = \frac{x}{r^2}.$$

Also

$$r\dot{r} = x\dot{x} + y\dot{y} \text{ and } \dot{\theta} = \frac{x\dot{y} - y\dot{x}}{r^2}.$$

(i) Transformation law as already defined gives

$$\begin{aligned} A'^{\mu} &= \frac{\partial x'_{\mu}}{\partial x_{\alpha}} A^{\alpha} \quad (\alpha=1, 2) \\ &= \frac{\partial x'_{\mu}}{\partial x_1} A^1 + \frac{\partial x'_{\mu}}{\partial x_2} A^2 \\ &= \frac{\partial x'_{\mu}}{\partial x_1} \dot{x} + \frac{\partial x'_{\mu}}{\partial x_2} \dot{y} \text{ as } A^1 = \dot{x} \text{ and } A^2 = \dot{y} \\ \therefore A'^1 &= \frac{\partial x'_1}{\partial x_1} \dot{x} + \frac{\partial x'_1}{\partial x_2} \dot{y} \\ &= \frac{\partial r}{\partial x} \dot{x} + \frac{\partial r}{\partial y} \dot{y} \text{ as } x_1 = x, x_2 = y \text{ and } x'_1 = r \\ &= \frac{x}{r} \dot{x} + \frac{y}{r} \dot{y} \\ &= \frac{x\dot{x} + y\dot{y}}{r} = \frac{r\dot{r}}{r} = \dot{r} \end{aligned}$$

and

$$\begin{aligned} A'^2 &= \frac{\partial x'_2}{\partial x_1} \dot{x} + \frac{\partial x'_2}{\partial x_2} \dot{y} \\ &= \frac{\partial \theta}{\partial x} \dot{x} + \frac{\partial \theta}{\partial y} \dot{y} \text{ as } x'_2 = \theta, x_1 = x \text{ and } x_2 = y \\ &= -\frac{y}{r^2} \dot{x} + \frac{x}{r^2} \dot{y} \\ &= \frac{x\dot{y} - y\dot{x}}{r^2} = \dot{\theta} \end{aligned}$$

$$\begin{aligned} \text{(ii) } A'^1 &= \frac{\partial r}{\partial x} \dot{x} + \frac{\partial r}{\partial y} \dot{y} \text{ as in part (i)} \\ &= \frac{x\dot{x} + y\dot{y}}{r} \end{aligned}$$

But $x\dot{x} + y\dot{y} = r\dot{r}$ gives on differentiation,

$$\begin{aligned} \text{i.e., } x\ddot{x} + \dot{x}^2 + y\ddot{y} + \dot{y}^2 &= r\ddot{r} + r\dot{r}^2, \\ x\ddot{x} + y\ddot{y} &= r\ddot{r} + \dot{r}^2 - \dot{x}^2 - \dot{y}^2 \\ &= r\ddot{r} + \dot{r}^2 - (\dot{r} \cos \theta - r \sin \theta \dot{\theta})^2 \\ &\quad - (\dot{r} \sin \theta + r \cos \theta \dot{\theta})^2 \\ &= r\ddot{r} + \dot{r}^2 - \dot{r}^2 - r^2 \dot{\theta}^2 \\ &= r\ddot{r} - r^2 \dot{\theta}^2 \end{aligned}$$

Thus

$$\begin{aligned} A'^1 &= \frac{r\ddot{r} - r^2 \dot{\theta}^2}{r} \\ &= \ddot{r} - r\dot{\theta}^2 \end{aligned}$$

and

$$\begin{aligned} A'^2 &= \frac{\partial \theta}{\partial x} \dot{x} + \frac{\partial \theta}{\partial y} \dot{y} \\ &= \frac{x\ddot{y} - y\ddot{x}}{r^2} \end{aligned}$$

But

$x\ddot{y} - y\ddot{x} = r\dot{\theta}^2$ gives on differentiation

$$\begin{aligned} \dot{x}\ddot{y} + x\dddot{y} - \dot{y}\ddot{x} - y\ddot{x} &= r^2\ddot{\theta} + 2r\dot{r}\dot{\theta} \\ \text{or } x\ddot{y} - y\ddot{x} &= r^2\ddot{\theta} + 2r\dot{r}\dot{\theta} \end{aligned}$$

$\therefore$

$$\begin{aligned} A'^2 &= \frac{r^2\ddot{\theta} + 2r\dot{r}\dot{\theta}}{r^2} \\ &= \ddot{\theta} + \frac{2\dot{r}\dot{\theta}}{r} \end{aligned}$$

3.5. SYMMETRIC AND ANTI-SYMMETRIC TENSORS

Let a tensor be such that two contravariant or covariant indices of it can be interchanged without altering the value of the tensor, then the tensor is termed as symmetrical or symmetric in these indices.

If $A^{\mu\nu}$ and $A^{\nu\mu}$ be two contravariant tensors in a certain system of co-ordinates such that

$$A^{\mu\nu} = A^{\nu\mu}$$

Then if $A^{\mu\nu}$ and $A^{\nu\mu}$ become $A'^{\mu\nu}$ and $A'^{\nu\mu}$ in another system of coordinates, the symmetry will be maintained in this system also if $A'^{\mu\nu} = A'^{\nu\mu}$.

To show it, let us consider

$$\begin{aligned} A'^{\mu\nu} &= \frac{\partial x'_{\mu}}{\partial x_{\alpha}} \frac{\partial x'_{\nu}}{\partial x_{\beta}} A^{\alpha\beta} \\ &= \frac{\partial x'_{\nu}}{\partial x_{\beta}} \frac{\partial x'_{\mu}}{\partial x_{\alpha}} A^{\beta\alpha} \text{ (on interchanging the indices)} \\ &= A'^{\nu\mu} \text{ as } A^{\beta\alpha} = A^{\alpha\beta} \end{aligned}$$

which shows the symmetry in the other system also.

Similarly if we consider two covariant tensors, $A_{\mu\nu}$ and $A_{\nu\mu}$ such that $A_{\mu\nu} = A_{\nu\mu}$ in one system

Then if they become $A'_{\mu\nu}$ and $A'_{\nu\mu}$ in another system, we have

$$\begin{aligned} A'_{\mu\nu} &= \frac{\partial x_{\alpha}}{\partial x'_{\mu}} \frac{\partial x_{\beta}}{\partial x'_{\nu}} A_{\alpha\beta} \\ &= \frac{\partial x_{\beta}}{\partial x'_{\nu}} \frac{\partial x_{\alpha}}{\partial x'_{\mu}} A_{\beta\alpha} \text{ (on interchanging the indices)} \\ &= A'_{\nu\mu} \text{ as } A_{\alpha\beta} = A_{\beta\alpha} \end{aligned}$$

In case one index is contravariant and other covariant, the symmetry cannot be easily defined. But it is notable that Kronecker delta which is a mixed tensor is symmetrical with respect to its indices.

When $A^{\mu\nu}$ is symmetrical, we have

$$A^{11}=A^{11}, A^{22}=A^{22} \text{ etc.}$$

In all, there are ${}^4C_2+4=5C_2$ components.

As an example the components of the angular momentum of a rigid body B_μ are connected with the components of its angular velocity A_α by the relations $B_\mu = \sum_{\alpha=1}^3 T_{\mu\alpha} A_\alpha$, where $T_{\mu\alpha}$ is the inertia tensor.

This tensor is symmetric, because

$$T_{\mu\alpha} = T_{\alpha\mu}. \quad \dots (1)$$

Now it has been already mentioned that a tensor can be expressed as a matrix and the columns and rows of a matrix when interchanged, the resulting tensor is the conjugate tensor. As such the conjugate of the tensor defined in §3.1 (a) may be written as

$$\phi_c = \begin{bmatrix} a_{xx} & a_{yx} & a_{zx} \\ a_{xy} & a_{yy} & a_{zy} \\ a_{xz} & a_{yz} & a_{zz} \end{bmatrix}$$

Thus the tensor ϕ_c is the conjugate of ϕ , when

$$a_{xy} = a_{yx}, a_{xz} = a_{zx} \text{ and } a_{yz} = a_{zy} \quad \dots (2)$$

If a single tensor satisfies this condition, it is called a **symmetric tensor**. It means that condition (1) is essential for a tensor to be symmetric. It has only six independent elements in three dimensional space and may be written as

$$\phi_{ym} = \begin{bmatrix} a_{xx} & a_{xy} & a_{xz} \\ a_{yx} & a_{yy} & a_{yz} \\ a_{zx} & a_{zy} & a_{zz} \end{bmatrix}$$

The relations (2) between the components of tensors follow that any symmetrical tensor corresponds with a transformation from the principal axes to another rectangular system of axes. To express the symmetrical linear functions by graphical function, let us suppose that we have two vectors $\mathbf{u}$ and $\mathbf{v}$ such that

$$\mathbf{v} = \phi_{ym} \mathbf{u} \quad \dots (3)$$

Project $\mathbf{u}$ upon $\mathbf{v}$ so that we calculate

$$\mathbf{u} \cdot \mathbf{v} = u_x v_x + u_y v_y + u_z v_z, \quad \dots (4)$$

where u_x, u_y, u_z and v_x, v_y, v_z are the components of $\mathbf{u}$ and $\mathbf{v}$ respectively along the principal axes.

Multiplying both sides of (3) scalarly by $\mathbf{u}$ we get

$$\begin{aligned} \mathbf{u} \cdot \mathbf{v} &= \mathbf{u} \cdot \phi_{ym} \mathbf{u} \\ &= u_x v_x + u_y v_y + u_z v_z \text{ from (4)} \\ &= u_x (a_{xx} u_x + a_{xy} u_y + a_{xz} u_z) + (a_{yx} u_x + a_{yy} u_y + a_{yz} u_z) u_y \\ &\quad + u_z (a_{zx} u_x + a_{zy} u_y + a_{zz} u_z) \end{aligned}$$

Say $S = a_{xx} u_x^2 + a_{yy} u_y^2 + a_{zz} u_z^2 + 2(a_{xy} u_x u_y + a_{yz} u_y u_z + a_{zx} u_z u_x)$ where S is a scalar point function.

From (5), we get

$$\frac{\partial S}{\partial u_x} = 2[a_{xx} u_x + a_{xy} u_y + a_{xz} u_z] = 2 v_x \text{ by §3.1 (7)} \quad \dots (5)$$

or

$$u_x = \frac{1}{2} \frac{\partial S}{\partial u_x}$$

Similarly

$$v_y = \frac{1}{2} \frac{\partial S}{\partial u_y} \text{ and } v_z = \frac{1}{2} \frac{\partial S}{\partial u_z}$$

So that $\mathbf{v} = v_x \mathbf{i} + v_y \mathbf{j} + v_z \mathbf{k}$; $\mathbf{i}, \mathbf{j}, \mathbf{k}$ being unit vectors along principal axes

$$\begin{aligned} &= \frac{1}{2} \left[\frac{\partial S}{\partial u_x} \mathbf{i} + \frac{\partial S}{\partial u_y} \mathbf{j} + \frac{\partial S}{\partial u_z} \mathbf{k} \right] \\ &= \frac{1}{2} \text{ grad } S. \end{aligned}$$

Which shows that $\mathbf{v}$ is a vector perpendicular to the surface $S = \text{const.}$ in the direction of the outward normal. But $S = \text{const.}$ is an equation of the second degree in the rectangular components of $\mathbf{u}$ regarding these as coordinates defining the extremity P of the vector $\mathbf{u}$ the locus of P is a conicoid.




Fig. 3.1

As a particular case if $S=1$, the surface defined under certain conditions is the **tensor ellipsoid** as shown in Fig. 3.1.

Also $\mathbf{u} \cdot \mathbf{v} = S = 1 = \text{resolute of } \mathbf{u} \text{ in the direction of } \mathbf{v}$.

In the direction of $\text{grad } S$, this resolute becomes $\frac{1}{|\mathbf{v}|}$.

Few other examples of symmetric tensor may be given as below

$$A_{\mu\alpha\beta} = A_{\alpha\mu\beta}$$

$$\text{and } A_{\mu\alpha\beta\gamma} = A_{\mu\beta\alpha\gamma} = A_{\alpha\mu\beta\gamma} = A_{\beta\alpha\mu\gamma} = A_{\alpha\beta\mu\gamma} = A_{\beta\mu\alpha\gamma}$$

Here the first tensor is symmetric in its first two indices and the second one is symmetric in first three indices.

If a tensor is such that two contravariant or covariant indices of it when interchanged, the components of the tensor alter in sign but not in magnitude, the tensor is said to be **anti-symmetric or skew-symmetric**.

Hence if $A^{\mu\nu} = -A^{\nu\mu}$

$$\begin{aligned} \text{then, } A^{\mu\nu} &= \frac{\partial x'^\mu}{\partial x^\alpha} \frac{\partial x'^\nu}{\partial x^\beta} A^{\alpha\beta} \\ &= -\frac{\partial x'^\nu}{\partial x^\beta} \frac{\partial x'^\mu}{\partial x^\alpha} A^{\beta\alpha} \\ &= -A^{\nu\mu} \text{ as } A^{\alpha\beta} = -A^{\beta\alpha} \end{aligned}$$

Similarly if $A_{\mu\nu} = -A_{\nu\mu}$, then $A'_{\mu\nu} = -A'_{\nu\mu}$

Here $A^{12} = -A^{21}$ etc. and $A^{11} = 0 = A^{22} = A^{33}$

As such number of arithmetic components is 4C_2 only.

Evidently the components of an antisymmetric tensor satisfy the relations

$$A_{\mu\nu} + A_{\nu\mu} = 0 \text{ or } A_{\mu\nu} = -A_{\nu\mu} \quad \dots (6)$$

which follows that the tensor changes its sign when indices are interchanged. If $\mu = \nu$, then (6) yields.

$$A_{\mu\mu} + A_{\mu\mu} = 0 \text{ or } 2 A_{\mu\mu} = 0 \text{ or } A_{\mu\mu} = 0 \quad \dots (7)$$

So in terms of coefficients, $a_{xx} = a_{yy} = a_{zz} = 0$

and $a_{xy} = -a_{yx}$, $a_{xz} = -a_{zx}$, $a_{yz} = -a_{zy}$

give the conditions for a tensor to be anti or skew-symmetric.

Thus ϕ expressed by §3.1 (9) will be antisymmetric if

$$\phi_{sk} = \begin{bmatrix} 0 & a_{xy} & a_{xz} \\ -a_{xy} & 0 & a_{yz} \\ -a_{xz} & -a_{yz} & 0 \end{bmatrix}$$

The matrix has only three components. The property of having only three components is possessed by vectors. This leads to the conclusion that an operation of ϕ_{sk} on the vector u , is exactly equivalent to the vector product of two vectors, since the final result is itself a vector v . For example consider the product of the coordinates of two points

$$A_{\mu\nu} = x_\mu \xi_\nu - \xi_\mu x_\nu \quad \dots (8)$$

In 3-dimensional space, suffixes can have only 3 values (1, 2, 3) and so any pair of suffixes can be replaced by a single one which is not present in the pair i.e.,

$$\left. \begin{aligned} A_{23} &= -A_{32} = A_1 \\ A_{12} &= -A_{21} = A_2 \\ A_{31} &= -A_{13} = A_3 \end{aligned} \right\} \quad \dots (9)$$

Thus the anti-symmetric system may be replaced by

$$\begin{aligned} \xi_{\mu\nu\sigma} &= +1 \text{ if } (\mu, \nu, \sigma) \text{ is an even permutation of } (1, 2, 3) \\ &= -1 \text{ if } \text{,,} \text{,, odd } \text{,,} \text{,,} \\ &= 0 \text{ if any of the suffixes are equal.} \end{aligned}$$

Now equations (9) yield,

$$A_1 = \frac{1}{2} [2 A_{33}], A_2 = \frac{1}{2} [2 A_{13}], A_3 = \frac{1}{2} [2 A_{12}]$$

$$\text{i.e., } A_1 = \frac{1}{2} \sum_{\mu, \nu=1}^3 \xi_{\mu\nu\sigma} A_{\mu\nu} \quad \dots (10)$$

There being only two terms in sum (10).

Few other examples of antisymmetric tensors are

$$A_{\mu\nu\sigma} = -A_{\nu\mu\sigma}$$

$$\text{and } A_{\mu\nu\sigma\rho} = -A_{\sigma\rho\mu\nu} = A_{\sigma\mu\nu\rho} = -A_{\mu\sigma\nu\rho} = -A_{\nu\mu\sigma\rho} = -A_{\sigma\nu\mu\rho}.$$

3.6. INVARIANT TENSORS

It is not known about any vector which has the same components in different systems of co-ordinates, but there exist tensors of higher ranks which have the same components in all the frames of reference. These tensors are called to have the *invariant components* or *invariant tensors* in general. One of the examples of such a tensor is Kronecker symbol defined as follows :

With respect to the old frame of reference (i.e., before rotation)

$$\frac{\partial x_\mu}{\partial x_\nu} = \begin{cases} 0 & \text{if } \mu \neq \nu \\ 1 & \text{if } \mu = \nu \end{cases} \text{ since } x_\mu \text{ is independent of } x_\nu.$$

$$\text{But } \frac{\partial x_\mu}{\partial x_\nu'} = \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial x_\alpha'}{\partial x_\nu}.$$

$$\text{Hence } \frac{\partial x_\mu}{\partial x_\nu} = \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial x_\alpha'}{\partial x_\nu} = \delta^\mu_\nu \quad \dots (1)$$

$$\text{where } \delta^\mu_\nu = \begin{cases} 1 & \text{if } \mu = \nu \\ 0 & \text{if } \mu \neq \nu \end{cases}$$

The symbol $\delta_{\nu}^\mu = \delta_{\mu\nu} = \delta^{\mu\nu} = \begin{cases} 1 & \text{if } \mu = \nu \\ 0 & \text{if } \mu \neq \nu \end{cases}$ is called as *Kronecker delta symbol*. In terms of new frame of reference (i.e., after the rotation), we may write

$$\begin{aligned} \delta'_{\nu}{}^\mu &= \frac{\partial x_\mu'}{\partial x_\nu'} = \frac{\partial x_\mu'}{\partial x_\alpha} \frac{\partial x_\alpha}{\partial x_\nu'} = \begin{cases} 1 & \text{if } \mu = \nu \\ 0 & \text{if } \mu \neq \nu \end{cases} \\ &= \frac{\partial x_\mu'}{\partial x_\beta} \cdot \frac{\partial x_\alpha}{\partial x_\nu'} \cdot \frac{\partial x_\beta}{\partial x_\alpha} \\ &= \frac{\partial x_\mu'}{\partial x_\beta} \frac{\partial x_\alpha}{\partial x_\nu'} \delta_\alpha^\beta \text{ since } \delta_\alpha^\beta = \frac{\partial x_\beta}{\partial x_\alpha} \end{aligned} \quad \dots (2)$$

(In summation convention).

Hence δ_{ν}^μ is invariant and transforms as mixed tensor of rank two. Similarly $\delta_{\mu\nu}$ transforms as the components of covariant tensor of rank two while $\delta^{\mu\nu}$ transforms as a contravariant tensor of rank two.

Kronecker symbol can be used as *substitution multiplier*.

$$\begin{aligned} \text{Since } A^\mu &= \frac{\partial x_\nu}{\partial x_\alpha'} \frac{\partial x_\alpha'}{\partial x_\mu} A^\nu \text{ (in summation convention)} \\ &= \delta_{\nu}^\mu A^\nu \text{ (by substitution of index).} \end{aligned}$$

$$\begin{aligned} \text{Similarly, } A_\nu &= \frac{\partial x_\alpha'}{\partial x_\nu} \frac{\partial x_\mu}{\partial x_\alpha'} A_\mu \\ &= \delta_{\nu}^\mu A_\mu \end{aligned}$$

$$\text{and } \delta_{\mu\nu} \delta_{\nu\beta} = \frac{\partial x_\mu}{\partial x_\nu} \frac{\partial x_\nu}{\partial x_\beta} = \frac{\partial x_\mu}{\partial x_\beta} = \delta_{\mu\beta}$$

$$\begin{aligned} \text{while } \delta^\mu_\mu &= \delta_1^1 + \delta_2^2 + \delta_3^3 \text{ (summation convention)} \\ &= 3 \text{ (in three dimensional geometry)} \\ &= 4 \text{ (in four dimensional geometry).} \end{aligned}$$

Secondly, we define the generalised Kronecker delta symbol, $\delta_{\beta\gamma}^{\mu\nu}$ by

$$\delta_{\beta\gamma}^{\mu\nu} = \begin{cases} 1 & \text{if } \mu=\beta, \nu=\gamma \text{ and } \mu \neq \nu \text{ (as } \delta_{12}^{12}=1), \\ -1 & \text{if } \mu=\gamma, \nu=\beta \text{ and } \mu \neq \nu \text{ (as } \delta_{21}^{12}=-1), \\ 0 & \text{for all other combinations of indices.} \end{cases}$$

Similarly we can define $\delta_{\beta\gamma\xi}^{\mu\nu\sigma}$ as an absolute (invariant) tensor of rank six.

Conclusively if both upper and lower indices of a generalised delta consist of the same distinct numbers chosen from 1, 2, 3, the delta is 1 or -1 according as the upper indices form an even or odd permutation of the lower, in all other permutations the delta is zero. We have for example

$$\delta_{12}^{12}=1, \quad \delta_{32}^{32}=-1, \quad \delta_{11}^{23}=\delta_{21}^{13}=0;$$

$$\delta_{123}^{123}=\delta_{231}^{231}=1, \quad \delta_{123}^{213}=\delta_{123}^{321}=-1, \quad \delta_{123}^{312}=0.$$

Evaluation of the various possible combinations of indices shows that

$$\left. \begin{aligned} \delta_{\beta\nu}^{\mu\beta} &= -\delta_{\nu\beta}^{\mu\nu} = 2 \delta_{\beta}^{\mu} \\ \delta_{\beta\xi}^{\mu\nu} &= \delta_{\beta}^{\mu} \delta_{\xi}^{\nu} - \delta_{\xi}^{\mu} \delta_{\beta}^{\nu} \end{aligned} \right\} \dots (3)$$

Alternating or Permutation epsilon tensor. This tensor is also invariant component tensor of third rank and anti-symmetric in every pair of indices. Let $\epsilon_{\mu\nu\sigma}$ be such a tensor; then

$$\epsilon_{\mu\nu\sigma} = -\epsilon_{\nu\mu\sigma} = \epsilon_{\sigma\mu\nu} = -\epsilon_{\mu\sigma\nu} = \epsilon_{\nu\sigma\mu} = -\epsilon_{\sigma\nu\mu} \dots (6)$$

But if $\mu=\nu$ then $\epsilon_{\mu\nu\sigma} = -\epsilon_{\nu\mu\sigma}$ gives $\epsilon_{\mu\mu\sigma} = -\epsilon_{\mu\mu\sigma}$ or

$$\epsilon_{\mu\mu\sigma} = 0 \dots (7)$$

It is clear that whenever two indices are equal, the component is zero. Moreover a tensor of third rank in three dimensional geometry has 27 components. But in case of the tensor $\epsilon_{\mu\nu\sigma}$ only 6 components are non-vanishing. All of them have the same absolute value, 3 being positive and the rest three are negative.

So $\epsilon_{123} = \epsilon_{231} = \epsilon_{312} = 1$ and $\epsilon_{213} = \epsilon_{321} = \epsilon_{132} = -1$ while all other components are zero.

Thus all the even permutations of 1, 2 and 3 correspond to the components with value +1, while all the odd permutations correspond to -1.

The transformation law for this type of tensor in three dimensional space is given by

$$\epsilon'_{\mu\nu\sigma} = \sum_{\alpha, \beta, \gamma=1}^3 a_{\mu\alpha} a_{\nu\beta} a_{\sigma\gamma} \epsilon_{\alpha\beta\gamma} = \epsilon_{\mu\nu\sigma}$$

i.e., $\epsilon_{\mu\nu\sigma}$ is invariant.

Consequently

$$\epsilon'_{\mu\nu\sigma} = \epsilon_{\mu\nu\sigma} = \begin{cases} 1 & \text{when } \mu, \nu, \sigma \text{ are an even permutation} \\ -1 & \text{when } \mu, \nu, \sigma \text{ are an odd permutation,} \\ 0 & \text{when } \mu, \nu, \sigma \text{ contain two or more repeated indices.} \end{cases} \dots (9)$$

Similarly the contravariant components can also be discussed,

$$\left. \begin{aligned} \epsilon^{123} = \epsilon^{312} = \epsilon^{231} = 1, \quad \epsilon^{132} = \epsilon^{213} = \epsilon^{321} = -1, \\ \epsilon^{112} = \epsilon^{233} = \epsilon^{111} = \dots = 0. \end{aligned} \right\} \dots (10)$$

Pseudo tensor. Let there be a tensor $\epsilon_{\mu\sigma\tau\rho}$ of rank 4, defined such that

$$\epsilon_{\mu\sigma\tau\rho} = \begin{cases} +1 & \text{if } \mu\sigma\tau\rho \text{ is an even permutation of } 0, 1, 2, 3 \\ -1 & \text{if } \mu\sigma\tau\rho \text{ is an odd permutation of } 0, 1, 2, 3 \\ 0 & \text{if two or more indices are equal.} \end{cases} \dots (11)$$

These are termed as components of **pseudo tensor** of rank four.

In case ϕ is a scalar, the quantities $\phi\epsilon_{\mu\sigma\tau\rho}$ are called as **pseudo scalars** since they have only one component.

From every antisymmetric tensor $A_{\mu\sigma}$ of the second rank a pseudo tensor $A^*_{\mu\sigma}$ of the same rank can be obtained by multiplying the former with a pseudo-tensor of rank 4,

$$A^*_{\mu\sigma} = \frac{1}{2} \sum_{\alpha, \beta=0}^3 \epsilon^{\mu\sigma\tau\rho} A_{\alpha\beta} \dots (12)$$

Thus the product of a tensor with a pseudo-tensor is a pseudo tensor. It is called **dual of a given tensor**.

A useful property of ϵ tensor.

ϵ -tensor can be used to write the cross product of two vectors **A** and **B**.

Let

$$\mathbf{D} = \mathbf{A} \times \mathbf{B},$$

$$\begin{aligned} D_1 &= A_2 B_3 - A_3 B_2 = \epsilon_{123} A_2 B_3 + \epsilon_{132} A_3 B_2; \quad A_1, A_2, A_3, B_1, \\ &\quad B_2, B_3 \text{ being components of } \mathbf{A} \text{ and } \mathbf{B}. \\ &= \epsilon_{1\nu\sigma} A_{\nu} B_{\sigma} \end{aligned}$$

Similarly $D_2 = \epsilon_{2\nu\sigma} A_{\nu} B_{\sigma}$ (in summation convention)

$$D_3 = \epsilon_{3\nu\sigma} A_{\nu} B_{\sigma}$$

$$D_{\mu} = \epsilon_{\mu\nu\sigma} A_{\nu} B_{\sigma} \dots (13)$$

Evaluating various possible combinations we may have

$$\epsilon^{\mu\nu\sigma} \epsilon_{\sigma\alpha\beta} = \delta_{\alpha\beta}^{\mu\nu} = \delta_{\nu}^{\mu} \delta_{\beta}^{\sigma} - \delta_{\beta}^{\mu} \delta_{\nu}^{\sigma}$$

Thus, if $\mathbf{r} = \mathbf{C} \cdot (\mathbf{A} \times \mathbf{B}) = \mathbf{C} \cdot \mathbf{D}$ for $\mathbf{D} = \mathbf{A} \times \mathbf{B}$.

$$\mathbf{r} = C_{\mu} D^{\mu} \text{ (in summation convention)}$$

or

$$r = C^{\mu\nu\sigma} A_\nu A_\sigma$$

$$= \epsilon^{\mu\nu\sigma} C_\mu A_\nu B_\sigma$$

Similarly vector triple product of three vectors can be given as

$$\mathbf{E} = \mathbf{C} \times (\mathbf{A} \times \mathbf{B}) = \mathbf{C} \times \mathbf{D}$$

i.e.,

$$E^\mu = \epsilon^{\mu\nu\sigma} C_\nu D_\sigma$$

$$= \epsilon^{\mu\nu\sigma} C_\nu (\epsilon_{\sigma\alpha\beta} A^\alpha B^\beta)$$

$$= \epsilon^{\mu\nu\sigma} \epsilon_{\sigma\alpha\beta} C_\nu A^\alpha B^\beta$$

$$= \delta_{\alpha\beta}^{\mu\nu} C_\nu A^\alpha B^\beta = \left(\delta_\alpha^\mu \delta_\beta^\nu - \delta_\beta^\mu \delta_\alpha^\nu \right) C_\nu A^\alpha B^\beta$$

Since $\delta_\alpha^\mu A^\alpha = A^\mu$ etc., we get

$$E_\mu = \delta_\beta^\nu A^\mu C_\nu B^\beta - \delta_\alpha^\nu B^\mu C_\nu A^\alpha$$

$$= A^\mu (C_\beta B^\beta) - B^\mu (C_\alpha A^\alpha)$$

As such $\mathbf{E} = \mathbf{A} (\mathbf{C} \cdot \mathbf{B}) - \mathbf{B} (\mathbf{C} \cdot \mathbf{A})$.

Evaluation of $\nabla \times (\mathbf{V} \times \mathbf{W})$ using ϵ -tensor.

Suppose, $\nabla \times (\mathbf{V} \times \mathbf{W}) = \nabla \times \mathbf{Z}$,
then $(\nabla \times \mathbf{Z})_\mu = \epsilon^{\mu\nu\sigma} \nabla_\nu Z_\sigma = \epsilon^{\mu\nu\sigma} \nabla_\mu (\epsilon_{\sigma\alpha\beta} V^\alpha W^\beta)$

$$\epsilon^{\mu\nu\sigma} \epsilon_{\sigma\alpha\beta} \nabla_\nu V^\alpha W^\beta = \delta_{\alpha\beta}^{\mu\nu} (V^\alpha \nabla_\nu W^\beta + W^\beta \nabla_\nu V^\alpha)$$

$$= \left(\delta_\nu^\mu \delta_\beta^\nu - \delta_\beta^\mu \delta_\alpha^\nu \right) (V^\alpha \nabla_\nu W^\beta + W^\beta \nabla_\nu V^\alpha)$$

$$= V^\mu \nabla_\beta W^\beta - V^\nu \nabla_\nu W^\mu + W^\nu \nabla_\nu V^\mu - W^\mu \nabla_\alpha V^\alpha$$

So $\nabla \times \mathbf{Z} = \nabla \times (\mathbf{V} \times \mathbf{W})$

$$= \mathbf{V} (\nabla \cdot \mathbf{W}) - (\mathbf{V} \cdot \nabla) \mathbf{W} + (\mathbf{W} \cdot \nabla) \mathbf{V} - \mathbf{W} (\nabla \cdot \mathbf{V})$$

Similarly all the vector relationships can be derived by using ϵ -tensor.

Krutkov's tensor. Let us consider a tensor $A^{\mu\nu\gamma\beta\sigma}$ of fourth rank having following properties:

- (1) Antisymmetric with respect to the first pair of indices
 $A^{\mu\nu\gamma\beta\sigma} = -A^{\nu\mu\gamma\beta\sigma}$
- (2) Antisymmetric in second pair of indices
 $A^{\mu\nu\gamma\beta\sigma} = -A^{\mu\nu\sigma\beta\gamma}$
- (3) Symmetric in cyclic order
 $A^{\mu\nu\gamma\beta\sigma} + A^{\mu\beta\sigma\gamma\nu} + A^{\mu\sigma\gamma\nu\beta} = 0$

Then in terms of second derivatives of $A^{\mu\nu\gamma\beta\sigma}$ we can form a new tensor given by

$$B^{\mu\sigma} = \sum_{\gamma, \beta=0}^3 \frac{\partial^2 A^{\mu\nu\gamma\beta\sigma}}{\partial x_\gamma \partial x_\beta} \quad \dots (14)$$

This tensor is called as *Krutkov's tensor*. If we differentiate equation (14) with respect to x , we have

$$\sum_{\sigma=0}^3 \frac{\partial B^{\mu\sigma}}{\partial x_\sigma} = 0. \quad \dots (15)$$

This is an important property of Krutkov's tensor.

Problem 6. Define a tensor. Prove that the Kronecker symbol δ_i^k is a tensor where components are the same in every coordinate system. (Agra 1966, 70)

We know that the tensors are quantities obeying certain transformation laws so that tensor analysis may be regarded as an indispensable part of study which is rather suitable for the mathematical formulation of natural laws which remain invariant when the coordinate system is changed to another. The rank of a tensor measures the number of the mode of changes of a physical quantity when passing from one system to another which is in rotation relative to the first. As such tensor of zero rank is a scalar quantity and the tensor of rank one is a vector quantity.

The laws of transformation of vector being defined by

$$A'^\mu = \sum_{\alpha=1}^4 \frac{\partial x_\mu'}{\partial x_\alpha} A^\alpha \text{ (contravariant vector)}$$

$$\text{and } A'_\mu = \sum_{\alpha=1}^4 \frac{\partial x_\alpha}{\partial x_\mu'} A_\alpha \text{ (covariant vector)}$$

In Minkowski's four dimensional space, we define the tensors of rank n as follows:

$$\text{Contravariant tensor : } A'^{\mu\nu} = \frac{\partial x_\mu'}{\partial x_\alpha} \frac{\partial x_\nu'}{\partial x_\beta} A^{\alpha\beta} \quad \dots (1)$$

$$\text{Covariant tensor : } A'_{\mu\nu} = \frac{\partial x_\alpha}{\partial x_\mu'} \frac{\partial x_\beta}{\partial x_\nu'} A_{\alpha\beta} \quad \dots (2)$$

$$\text{Mixed tensor : } A'^\mu{}_\nu = \frac{\partial x_\alpha}{\partial x_\mu'} \frac{\partial x_\nu'}{\partial x_\beta} A^\alpha{}_\beta \quad \dots (3)$$

Each one having 4^2 i.e., 16 components.

Similarly we can define the tensors of higher ranks.

Now the Kronecker delta symbol δ_i^k is defined as

$$\delta_i^k = \frac{\partial x_k}{\partial x_i} = \frac{\partial x_k}{\partial x_j'} \frac{\partial x_j'}{\partial x_i}$$

which is easily deduced from (3) by choosing A_α^β to be the Kronecker delta δ_α^β so that

$$A'^\mu{}_\nu = \frac{\partial x_\alpha}{\partial x_\mu'} \frac{\partial x_\nu'}{\partial x_\beta} \delta_\alpha^\beta = \frac{\partial x_\alpha}{\partial x_\mu'} \frac{\partial x_\nu'}{\partial x_\beta} \cdot \frac{\partial x_\beta}{\partial x_\alpha}$$

$$= \frac{\partial x_\nu'}{\partial x_\mu'} = \delta_\mu^\nu$$

and now replacing μ by i , ν by j this gives

$$\delta_i^k = A'^k{}_i = \frac{\partial x_\alpha}{\partial x_i'} \frac{\partial x_k'}{\partial x_\beta} A_\alpha^\beta = \frac{\partial x_\alpha}{\partial x_i'} \frac{\partial x_k'}{\partial x_\beta} \delta_\alpha^\beta$$

From the definition of mixed tensor, it follows that δ_i^k is a mixed tensor of order two with 16 components in 4-dimensional space.

In order to show that the components of the tensor δ_i^k are the same in every coordinate system, let us define the symbol δ_i^k as

$$\delta_i^k = 1 \text{ if } i=k \\ = 0 \text{ if } i \neq k$$

which is evident from $\delta_i^k = \frac{\partial x_k}{\partial x_i} = 0$ when $i \neq k$
 $\delta_i^k = 1$ when $i=k$

In terms of new frame of reference (or new coordinate system), we may have

$$\delta_i'^k = \frac{\partial x_k'}{\partial x_i'} = \frac{\partial x_k'}{\partial x_j} \frac{\partial x_j}{\partial x_i} = 1 \text{ if } i=k \\ = 0 \text{ if } i \neq k$$

or $\delta_i'^k = \frac{\partial x_k'}{\partial x_i} \frac{\partial x_j}{\partial x_i'} \frac{\partial x_i}{\partial x_j} = \frac{\partial x_k'}{\partial x_i} \frac{\partial x_j}{\partial x_i'} \delta_j^i$ since $\delta_j^i = \frac{\partial x_i}{\partial x_j}$

From which it is clear that δ_i^k is invariant and transforms as a mixed tensor of rank two.

Problem 7. Prove that Kronecker delta is a mixed tensor of rank two. Its solution has been given in Problem 6. (Agra 1988)

Problem 8. Show that symmetry properties of a tensor are invariant. If $A_{\lambda\mu\nu} = A_{\mu\lambda\nu}$ then we have to show that $A'_{\lambda\mu\nu} = A'_{\mu\lambda\nu}$

The definition follows:

$$A'_{\lambda\mu\nu} = \sum_{\alpha, \beta, \gamma=1}^3 \frac{\partial x_\alpha}{\partial x'_\lambda} \frac{\partial x_\beta}{\partial x'_\mu} \frac{\partial x_\gamma}{\partial x'_\nu} A_{\alpha\beta\gamma}$$

$$\text{and } A'_{\mu\lambda\nu} = \sum_{\beta, \alpha, \gamma=1}^3 \frac{\partial x_\beta}{\partial x'_\mu} \frac{\partial x_\alpha}{\partial x'_\lambda} \frac{\partial x_\gamma}{\partial x'_\nu} A_{\beta\alpha\gamma}$$

The given tensor having symmetrical in first two indices, we have

$$A_{\lambda\mu\nu} = A_{\mu\lambda\nu} \text{ and } A_{\alpha\beta\gamma} = A_{\beta\alpha\gamma}$$

Using this relation and comparing the two equations for $A'_{\lambda\mu\nu}$ and $A'_{\mu\lambda\nu}$ we find that both the equations are identical i.e., $A'_{\lambda\mu\nu} = A'_{\mu\lambda\nu}$

Which follows that the tensor in other system is also symmetrical in first two indices. Hence the properties of symmetric tensors are invariant.

3.7. RULES WHICH GOVERN TENSOR ANALYSIS

Rule I. The sum and difference of two tensors of the same rank result in a third tensor of the same rank. Moreover, if $F_{\lambda\mu\dots}$ and $G_{\lambda\mu\dots}$ are the tensors of the same rank, then $pF_{\lambda\mu\dots} + qG_{\lambda\mu\dots}$ is also a tensor of the same rank (p, q being numbers).

Suppose there are two tensors $A_{\lambda\mu}$ and $B_{\lambda\mu}$, then it will be shown that

$$A_{\lambda\mu} + B_{\lambda\mu} = C_{\lambda\mu} \text{ where } C_{\lambda\mu} \text{ is another tensor of the same rank}$$

Expressing the tensor $A_{\lambda\mu}$ in the form of a matrix, we have

$$A_{\lambda\mu} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}, B_{\lambda\mu} = \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix}$$

and

$$C_{\lambda\mu} = \begin{bmatrix} c_{11} & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{bmatrix}$$

$$\text{so that } A_{\lambda\mu} + B_{\lambda\mu} = \begin{bmatrix} a_{11}+b_{11} & a_{12}+b_{12} & a_{13}+b_{13} \\ a_{21}+b_{21} & a_{22}+b_{22} & a_{23}+b_{23} \\ a_{31}+b_{31} & a_{32}+b_{32} & a_{33}+b_{33} \end{bmatrix}$$

If the relations between the coefficients a 's and b 's be such that

$$a_{\lambda\mu} + b_{\lambda\mu} = c_{\lambda\mu}$$

$$\text{then } A_{\lambda\mu} + B_{\lambda\mu} = \begin{bmatrix} c_{11} & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{bmatrix} = C_{\lambda\mu}$$

which is a tensor of the same rank.

$$\text{Similarly } A_{\lambda\mu} - B_{\lambda\mu} = \begin{bmatrix} d_{11} & d_{12} & d_{13} \\ d_{21} & d_{22} & d_{23} \\ d_{31} & d_{32} & d_{33} \end{bmatrix} = D_{\lambda\mu}$$

where $a_{\lambda\mu} - b_{\lambda\mu} = d_{\lambda\mu}$

Here $D_{\lambda\mu}$ is again a tensor of the same rank.

Further,

$$pA_{\lambda\mu} + qB_{\lambda\mu} = \begin{bmatrix} e_{11} & e_{12} & e_{13} \\ e_{21} & e_{22} & e_{23} \\ e_{31} & e_{32} & e_{33} \end{bmatrix} = E_{\lambda\mu}$$

where $pa_{\lambda\mu} + qb_{\lambda\mu} = e_{\lambda\mu}$

Showing that $E_{\lambda\mu}$ is also a tensor of the same rank.

The rule of addition may be generalized for any number of tensors of any rank.

Suppose there are two mixed tensors T and S of rank N , having their r indices (from λ_1 to λ_r) contravariant and s indices (from μ_1 to μ_s) covariant, then laws of their transformation may be written as

$$T'^{\alpha_1 \dots \alpha_r}_{\beta_1 \dots \beta_s} = \left| \frac{\partial x}{\partial x'} \right| N \left[\frac{\partial x_{\mu_1}}{\partial x'_{\beta_1}} \dots \frac{\partial x_{\mu_s}}{\partial x'_{\beta_s}} \right] \left[\frac{\partial x'^{\alpha_1}}{\partial x_{\lambda_1}} \dots \frac{\partial x'^{\alpha_r}}{\partial x_{\lambda_r}} \right] T^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} \dots (1)$$

$$S'^{\alpha_1 \dots \alpha_r}_{\beta_1 \dots \beta_s} = \left| \frac{\partial x}{\partial x'} \right| N \left[\frac{\partial x_{\mu_1}}{\partial x'_{\beta_1}} \dots \frac{\partial x_{\mu_s}}{\partial x'_{\beta_s}} \right] \left[\frac{\partial x'^{\alpha_1}}{\partial x_{\lambda_1}} \dots \frac{\partial x'^{\alpha_r}}{\partial x_{\lambda_r}} \right] S^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} \dots (2)$$

If the sum of two tensors $T^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s}$ and $S^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s}$ be a third tensor

$$U^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s}, \text{ i.e., } U^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} = T^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} + S^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} \dots (3)$$

So that

$$U'^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} = T'^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} + S'^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s}$$

$$= \left| \frac{\partial x}{\partial x'} \right| N \left[\frac{\partial x_{\mu_1}}{\partial x'_{\beta_1}} \dots \frac{\partial x_{\mu_s}}{\partial x'_{\beta_s}} \right] \left[\frac{\partial x'_{\alpha_1}}{\partial x_{\lambda_1}} \dots \frac{\partial x'_{\alpha_r}}{\partial x_{\lambda_r}} \right] \left[T^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} + S^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} \right]$$

$$= \left| \frac{\partial x}{\partial x'} \right| N \frac{\partial x_{\mu_1}}{\partial x'_{\beta_1}} \dots \frac{\partial x_{\mu_s}}{\partial x'_{\beta_s}}, \frac{\partial x'_{\alpha_1}}{\partial x_{\lambda_1}} \dots \frac{\partial x'_{\alpha_r}}{\partial x_{\lambda_r}} U^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} \text{ by (3)} \dots (4)$$

Which is transformation equation for a tensor of rank N having r contravariant and s covariant indices and follows that the sum of two tensors of the same rank is a new tensor of the same rank.

Rule II. The direct product of two tensors gives a new tensor of rank equal to the sum of ranks of these tensors.

Consider two tensors $T^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s}$ of rank N and $S^{\sigma_1 \dots \sigma_p}_{\rho_1 \dots \rho_q}$ of rank N' .

Their transformation may be given as,

$$T'^{\alpha_1 \dots \alpha_r}_{\beta_1 \dots \beta_s} = \left| \frac{\partial x}{\partial x'} \right| N \frac{\partial x_{\mu_1}}{\partial x'_{\beta_1}} \dots \frac{\partial x_{\mu_s}}{\partial x'_{\beta_s}} \frac{\partial x'_{\alpha_1}}{\partial x_{\lambda_1}} \dots \frac{\partial x'_{\alpha_r}}{\partial x_{\lambda_r}} T^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s}$$

$$S'^{\eta_1 \dots \eta_p}_{\xi_1 \dots \xi_q} = \left| \frac{\partial x}{\partial x'} \right| N' \frac{\partial x_{\rho_1}}{\partial x'_{\xi_1}} \dots \frac{\partial x_{\rho_q}}{\partial x'_{\xi_q}} \frac{\partial x'_{\eta_1}}{\partial x_{\sigma_1}} \dots \frac{\partial x'_{\eta_p}}{\partial x_{\sigma_p}} S^{\sigma_1 \dots \sigma_p}_{\rho_1 \dots \rho_q}$$

Then $T'^{\alpha_1 \dots \alpha_r}_{\beta_1 \dots \beta_s} S'^{\eta_1 \dots \eta_p}_{\xi_1 \dots \xi_q}$

$$= \left| \frac{\partial x}{\partial x'} \right| N + N' \frac{\partial x_{\mu_1}}{\partial x'_{\beta_1}} \dots \frac{\partial x_{\mu_s}}{\partial x'_{\beta_s}} \frac{\partial x'_{\alpha_1}}{\partial x_{\lambda_1}} \dots \frac{\partial x'_{\alpha_r}}{\partial x_{\lambda_r}} \frac{\partial x_{\rho_1}}{\partial x'_{\xi_1}} \dots \frac{\partial x_{\rho_q}}{\partial x'_{\xi_q}}$$

$$\frac{\partial x'_{\eta_1}}{\partial x_{\sigma_1}} \dots \frac{\partial x'_{\eta_p}}{\partial x_{\sigma_p}} \times \left[T^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} \right] \left[S^{\sigma_1 \dots \sigma_p}_{\rho_1 \dots \rho_q} \right]$$

or $U^{\alpha_1 \dots \alpha_r, \eta_1 \dots \eta_p}_{\beta_1 \dots \beta_s, \xi_1 \dots \xi_q} = \left| \frac{\partial x}{\partial x'} \right| N + N' \frac{\partial x_{\mu_1}}{\partial x'_{\beta_1}} \dots \frac{\partial x_{\mu_s}}{\partial x'_{\beta_s}} \frac{\partial x'_{\alpha_1}}{\partial x_{\lambda_1}} \dots \frac{\partial x'_{\alpha_r}}{\partial x_{\lambda_r}}$

$$\times \frac{\partial x_{\rho_1}}{\partial x'_{\xi_1}} \dots \frac{\partial x_{\rho_q}}{\partial x'_{\xi_q}} \frac{\partial x'_{\eta_1}}{\partial x_{\sigma_1}} \dots \frac{\partial x'_{\eta_p}}{\partial x_{\sigma_p}} \times U^{\lambda_1 \dots \lambda_r, \sigma_1 \dots \sigma_p}_{\mu_1 \dots \mu_s, \rho_1 \dots \rho_q} \dots (5)$$

where $U^{\lambda_1 \dots \lambda_r, \sigma_1 \dots \sigma_p}_{\mu_1 \dots \mu_s, \rho_1 \dots \rho_q} = T^{\lambda_1 \dots \lambda_r}_{\mu_1 \dots \mu_s} S^{\sigma_1 \dots \sigma_p}_{\rho_1 \dots \rho_q}$

the equation (5) transforms a tensor of rank $N + N'$.

Rule III Contraction. The algebraic operation by which the rank of a tensor may be lowered by 2 (or by any even number) is known as contraction. (Agra 1969)

The contraction of a tensor may be affected by adding up all the components which have equal indices in a given pair. Any two indices are converted into a pair of dummy indices.

Consider a tensor of rank 3 with one contravariant index α and two covariant indices β and γ . Then we have

$$A'^{\lambda}_{\mu\nu} = \sum_{\alpha, \beta, \gamma=0}^3 A^{\alpha}_{\beta\lambda} \frac{\partial x_{\beta}}{\partial x'_{\mu}} \frac{\partial x_{\gamma}}{\partial x'_{\nu}} \frac{\partial x'_{\lambda}}{\partial x_{\alpha}}$$

Replacing ν by λ , we have

$$A'^{\lambda}_{\mu\nu} = \sum_{\lambda, \mu, \nu=0}^3 A^{\alpha}_{\beta\gamma} \frac{\partial x_{\beta}}{\partial x'_{\mu}} \frac{\partial x_{\gamma}}{\partial x'_{\nu}} \frac{\partial x'_{\lambda}}{\partial x_{\alpha}}$$

$$= \sum_{\alpha, \beta, \gamma=0}^3 A^{\alpha}_{\beta\gamma} \frac{\partial x_{\beta}}{\partial x'_{\mu}} \frac{\partial x_{\gamma}}{\partial x_{\alpha}}$$

But $\frac{\partial x_{\gamma}}{\partial x_{\alpha}} = \begin{cases} 0 & \text{if } \gamma \neq \alpha \\ 1 & \text{if } \gamma = \alpha \end{cases}$

Choosing the second condition i.e., if $\gamma = \alpha$, $\frac{\partial x_{\gamma}}{\partial x_{\alpha}} = 1$, above relation becomes

$$A'^{\lambda}_{\mu\lambda} = \sum_{\alpha, \beta=0}^3 A^{\alpha}_{\beta\alpha} \frac{\partial x_{\beta}}{\partial x'_{\mu}}$$

$$A'_{\mu} = \sum_{\beta=0}^3 \frac{\partial x_{\beta}}{\partial x'_{\mu}} A_{\beta}$$

Which denotes the law of transformation of tensors of rank one, i.e., vectors. As a general rule we equate a certain covariant index to a contravariant index, sum on repeated indices, and obtain a new tensor of lower rank. This process is termed as **contraction**. The contraction of a tensor of rank 2 yields a scalar i.e., tensor of rank zero.

Illustration. We know that the scalar product of two vectors is a scalar quantity. It follows that the scalar product of two tensors of rank one is a tensor of rank zero. As such the rank is lowered by two.

Rule IV. Extension of the rank. The differentiation of each components of a tensor of rank n with respect to x, y, z , gives a new tensor of rank $(n+1)$, e.g.,

$$\frac{\partial A_{\lambda\mu}}{\partial x_{\nu}} = B_{\lambda\mu\nu} \dots (6)$$

This rule may be proved for a simple case, where the original tensor is of rank zero, i.e., a scalar say $S(x_1, x_2, x_3, t)$ where derivatives relative to the axes K are $\frac{\partial S}{\partial x}, \frac{\partial S}{\partial y}, \frac{\partial S}{\partial z}$.

In system K' , the scalar is $S'(x'_1, x'_2, x'_3, t')$, such that

$$S(x_1, x_2, x_3, t) = S'(x'_1, x'_2, x'_3, t')$$

So

$$\left. \begin{aligned} \frac{\partial S'}{\partial x'_1} &= \frac{\partial S}{\partial x_1} \frac{\partial x_1}{\partial x'_1} + \frac{\partial S}{\partial x_2} \frac{\partial x_2}{\partial x'_1} + \frac{\partial S}{\partial x_3} \frac{\partial x_3}{\partial x'_1} \\ \frac{\partial S'}{\partial x'_2} &= \frac{\partial S}{\partial x_1} \frac{\partial x_1}{\partial x'_2} + \frac{\partial S}{\partial x_2} \frac{\partial x_2}{\partial x'_2} + \frac{\partial S}{\partial x_3} \frac{\partial x_3}{\partial x'_2} \\ \frac{\partial S'}{\partial x'_3} &= \frac{\partial S}{\partial x_1} \frac{\partial x_1}{\partial x'_3} + \frac{\partial S}{\partial x_2} \frac{\partial x_2}{\partial x'_3} + \frac{\partial S}{\partial x_3} \frac{\partial x_3}{\partial x'_3} \end{aligned} \right\}$$

which may be written as a single equation

$$\frac{\partial S'}{\partial x'_i} = \frac{\partial x_m}{\partial x'_i} \frac{\partial S}{\partial x_m} \quad (i, m=1, 2, 3)$$

This shows that $\frac{\partial S}{\partial x_m}$ transforms like the components of a tensor of rank one, i.e. vector. Thus the differentiation of the tensor of rank zero, yields a tensor of rank one. The rank of a tensor can also be extended when a tensor depends on another tensor and a differentiation is performed. For example, consider a scalar, i.e., tensor of rank zero, say S depending on tensor $A_{\lambda\mu}$, so that

$$\frac{\partial S}{\partial A_{\lambda\mu}} = B_{\mu\nu}$$

is also a tensor of rank two. Thus the rank of the tensor is extended by two.

Rule V. The Quotient Law. If $A^\lambda B_{\mu\nu}$ is a tensor for all contravariant tensors A_λ then $B_{\mu\nu}$ is also a tensor.

$$\text{We have } A'^\lambda B'_{\mu\nu} = A^\alpha B_{\beta\gamma} \frac{\partial \alpha_\beta}{\partial x'^\mu} \frac{\partial x_\gamma}{\partial x'^\nu} \frac{\partial x'_\lambda}{\partial x_\alpha}$$

$$= A'^\lambda B_{\beta\gamma} \frac{\partial x_\beta}{\partial x'^\mu} \frac{\partial x_\gamma}{\partial x'^\nu}$$

$$\text{as } A'^\lambda = A^\alpha \frac{\partial x'_\lambda}{\partial x_\alpha}$$

$$\text{or } A'^\lambda \left[B'_{\mu\nu} - B_{\beta\gamma} \frac{\partial x_\beta}{\partial x'^\mu} \frac{\partial x_\gamma}{\partial x'^\nu} \right] = 0$$

But A'^λ being arbitrary, $A'^\lambda \neq 0$ so that

$$B'_{\mu\nu} = B_{\beta\gamma} \frac{\partial x_\beta}{\partial x'^\mu} \frac{\partial x_\gamma}{\partial x'^\nu}$$

which shows that $B_{\mu\nu}$ is a covariant tensor.

Problem 9. Show that there exists no distinction between contravariant and covariant vectors if we restrict ourselves to transformations of the type

$$x'_\alpha = a^\alpha_\lambda x_\lambda + b^\alpha$$

where b^α are n constants which do not necessarily form the components of a contravariant vector and a^α_λ are constant (not necessarily forming a tensor) such that $a^\alpha_\mu a^\mu_\lambda = \delta^\alpha_\lambda$

$$\text{Given } x'_\alpha = a^\alpha_\lambda x_\lambda + b^\alpha$$

$$\text{i.e., } a^\alpha_\lambda x_\lambda = x'_\alpha - b^\alpha$$

Multiplying (2) throughout by a^α_μ and summing over the index from 1 to n , we find

$$x_\mu = a^\alpha_\mu x'_\alpha - a^\alpha_\mu b^\alpha$$

Now (1) and (3) yield,

$$\frac{\partial x'_\alpha}{\partial x_\beta} = a^\alpha_\beta \text{ and } \frac{\partial x_\beta}{\partial x'_\alpha} = a^\alpha_\beta$$

$$\text{that } \frac{\partial x'_\alpha}{\partial x_\beta} = \frac{\partial x_\beta}{\partial x'_\alpha} = a^\alpha_\beta$$

This follows that the transformation laws

$$A'^\mu = \frac{\partial x'^\mu}{\partial x_\beta} A^\beta \text{ and } A'_\mu = \frac{\partial x^\alpha}{\partial x'^\mu} A_\alpha$$

define the same type of entity without any distinction between contravariant and covariant vectors.

Problem 10. Show that $\frac{\partial A_\lambda}{\partial x_\mu}$ is not a tensor although A_λ is a covariant tensor of rank one.

$$\text{We have } A'_\alpha = \frac{\partial x_\lambda}{\partial x'_\alpha} A_\lambda \text{ (by covariant law)}$$

Differentiating w.r.t. x'_β we get

$$\begin{aligned} \frac{\partial A'_\alpha}{\partial x'_\beta} &= \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial A_\lambda}{\partial x'_\beta} + \frac{\partial^2 x_\lambda}{\partial x'_\alpha \partial x'_\beta} A_\lambda \\ &= \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial A_\lambda}{\partial x_\mu} \frac{\partial x_\mu}{\partial x'_\beta} + \frac{\partial^2 x_\lambda}{\partial x'_\alpha \partial x'_\beta} A_\lambda \\ &= \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial A_\lambda}{\partial x_\mu} + \frac{\partial^2 x_\lambda}{\partial x'_\alpha \partial x'_\beta} A_\lambda \end{aligned}$$

The presence of the second term on the right shows that $\frac{\partial A_\lambda}{\partial x_\mu}$ does

not transform as a tensor.

Problem 11. If xy , $2y-z^2$ and xz are the components of a covariant tensor in rectangular coordinates, then find its covariant components in spherical coordinates.

Suppose that $x=x_1$, $y=x_2$, $z=x_3$

Then the covariant components of a tensor A_λ are

$$A_1 = xy = x_1 x_2$$

$$A_2 = 2y - z^2 = 2x_2 - x_3^2$$

$$A_3 = xz = x_1 x_3$$

Let A'_μ be the covariant components in spherical coordinates. Then

$$x'_1 = r, x'_2 = \theta, x'_3 = \phi$$

$$A'_\mu = \frac{\partial x_\lambda}{\partial x'_\mu} A_\lambda \quad \dots (1)$$

Now the transformation equations are

$$x = r \sin \theta \cos \phi, y = r \sin \theta \sin \phi, z = r \cos \theta$$

which in existing case become

$$x_1 = x'_1 \sin x'_2 \cos x'_3, x_2 = x'_1 \sin x'_2 \sin x'_3, x_3 = x'_1 \cos x'_2$$

from (1) we have

$$\begin{aligned}
 A_1' &= \frac{\partial x_1}{\partial x_1'} A_1 + \frac{\partial x_2}{\partial x_1'} A_2 + \frac{\partial x_3}{\partial x_1'} A_3 \\
 &= (\sin x_2' \cos x_3') x_1 x_2 + \sin x_2' \sin x_3' (2x_2 - x_3^2) + \cos x_3' x_1^2 \\
 &= (\sin \theta \cos \phi) (r^2 \sin^2 \theta \sin \phi \cos \phi) + \sin \theta \sin \phi (2r \sin \theta \sin \phi \\
 &\quad - r^2 \cos^2 \theta) + \cos \theta (r^2 \sin \theta \cos \theta \cos \phi)
 \end{aligned}$$

Similarly A_2' and A_3' can be determined.

Problem 12. If $A_{\lambda\mu}$ is a skew-symmetric tensor, show that

$$B_v^\mu B_r^\sigma + B_r^\mu B_v^\sigma) A_{\mu\sigma} = 0$$

$\therefore A_{\mu\nu}$ is skew-symmetric

$\therefore A_{11} = A_{22} = \dots = 0$ and $A_{12} = -A_{21}$ etc.

Now left hand expression = $B_v^\mu B_r^\sigma + B_r^\mu B_v^\sigma$

$$\begin{aligned}
 &= (B_v^1 B_r^2 + B_r^1 B_v^2) A_{12} + (B_v^2 B_r^1 + B_r^2 B_v^1) A_{21} + \dots \\
 &= (B_v^1 B_r^2 + B_r^1 B_v^2) A_{12} + (B_v^2 B_r^1 + B_r^2 B_v^1) A_{21} + \dots \\
 &+ (B_v^1 B_r^2 + B_r^1 B_v^2) A_{12} + (B_v^2 B_r^1 + B_r^2 B_v^1) A_{21} + \dots \\
 &= 0 + 0 + \dots + (B_v^1 B_r^2 + B_r^1 B_v^2) \{A_{12} + A_{21}\} + \dots \text{ by (1)} \\
 &= (B_v^1 B_r^2 + B_r^1 B_v^2) \{0\} \text{ by (1)} \\
 &= 0.
 \end{aligned}$$

Problem 13. Show that every tensor can be expressed as the sum of two tensors, one of which is symmetric and other skew-symmetric in a pair of covariant or contravariant indices.

$$\begin{aligned}
 \text{Since } A_{\mu\nu} &= \frac{1}{2} [A_{\mu\nu} + A_{\nu\mu}] + \frac{1}{2} [A_{\mu\nu} - A_{\nu\mu}] \\
 &= B_{\mu\nu} + C_{\mu\nu} \text{ (say)}
 \end{aligned}$$

where $B_{\mu\nu} = \frac{1}{2} [A_{\mu\nu} + A_{\nu\mu}] = B_{\nu\mu}$ is symmetric

and $C_{\mu\nu} = \frac{1}{2} [A_{\mu\nu} - A_{\nu\mu}] = -C_{\nu\mu}$ is anti-symmetric

Hence the proposition.

Problem 14. Show that $A^{\mu\sigma\nu}$ is a tensor

(Agra, 1961)

$$\text{We have } A_{\alpha\beta\gamma}^{\delta} = \frac{\partial x_{\delta}}{\partial x_{\alpha}} \frac{\partial x_{\mu}}{\partial x_{\beta}} \frac{\partial x_{\sigma}}{\partial x_{\gamma}} \frac{\partial x_{\nu}}{\partial x_{\gamma}} A_{\mu\sigma\nu}^{\tau}$$

which follows that $A^{\mu\sigma\nu}$ according to transformation law, is a mixed tensor of rank 4, contravariant of order 1 and covariant of order 3.

Problem 15. If A^λ and B^μ are the components of a contravariant and covariant tensors of rank one, then show that $C_\mu^\lambda = A^\lambda B_\mu$ are the components of a mixed tensor of rank two.

$$\text{We have } A'^\lambda = A^\alpha \frac{\partial x'^\lambda}{\partial x^\alpha} \text{ and } B'_\mu = B_\beta \frac{\partial x_\beta}{\partial x'_\mu}$$

$$\text{So that } C'^\lambda_\mu = A'^\lambda B'_\mu = A^\alpha B_\beta \frac{\partial x'^\lambda}{\partial x^\alpha} \frac{\partial x_\beta}{\partial x'_\mu} = A^\alpha B_\beta \frac{\partial x'^\lambda}{\partial x^\alpha} \frac{\partial x_\beta}{\partial x'_\mu}$$

which shows that C_μ^λ transforms as a mixed tensor of rank two.

Problem 16. Show that the contracted tensor $A^{\nu\sigma\rho}_{\lambda\mu\nu}$ is a mixed tensor.

we have

$$A_{\alpha\beta\gamma}^{\delta\xi\eta} = \frac{\partial x'^\delta}{\partial x_\alpha} \frac{\partial x'^\xi}{\partial x_\beta} \frac{\partial x'^\eta}{\partial x_\gamma} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x_\nu}{\partial x'_\gamma} A_{\lambda\mu\nu}^{\tau\sigma\rho}$$

Setting $\delta = \gamma$ and summing we get

$$\begin{aligned}
 A_{\alpha\beta\gamma}^{\gamma\xi\eta} &= \frac{\partial x'^\xi}{\partial x_\alpha} \frac{\partial x'^\eta}{\partial x_\beta} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x_\nu}{\partial x'_\gamma} A_{\lambda\mu\nu}^{\tau\sigma\rho} \\
 &= \frac{\partial x'^\xi}{\partial x_\alpha} \frac{\partial x'^\eta}{\partial x_\beta} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \delta^\nu_\gamma A_{\lambda\mu\nu}^{\tau\sigma\rho} \\
 &= \frac{\partial x'^\xi}{\partial x_\alpha} \frac{\partial x'^\eta}{\partial x_\beta} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} A_{\lambda\mu\nu}^{\nu\sigma\rho}
 \end{aligned}$$

If we compare it with the transformation law

$$A_{\alpha\beta}^{\xi\eta} = \frac{\partial x'^\xi}{\partial x_\alpha} \frac{\partial x'^\eta}{\partial x_\beta} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} A_{\lambda\mu}^{\sigma\rho}$$

then we conclude that $A_{\lambda\mu\nu}^{\nu\sigma\rho}$ is a tensor of rank four, contravariant of rank 2 and covariant of rank 2.

Problem 17. If the components of a tensor are zero in one co-ordinate system, then prove that the components are zero in all co-ordinate systems.

Consider the components of a tensor in the form

$$T_{\mu_1, \mu_2, \dots, \mu_s}^{\lambda_1, \lambda_2, \dots, \lambda_r}$$

where the indices $\lambda_1, \lambda_2, \dots, \lambda_r, \mu_1, \mu_2, \dots, \mu_s$ run through the integers 1, 2, ..., n.

The components then transform according to the rule

$$T_{\mu_1, \mu_2, \dots, \mu_s}^{\lambda_1, \lambda_2, \dots, \lambda_r} = \left| \frac{\partial x}{\partial x'} \right|^N \frac{\partial x_{\lambda_1}}{\partial x'_{\alpha_1}} \frac{\partial x_{\lambda_2}}{\partial x'_{\alpha_2}} \dots \frac{\partial x_{\lambda_r}}{\partial x'_{\alpha_r}} \frac{\partial x_{\beta_1}}{\partial x'_{\mu_1}} \frac{\partial x_{\beta_2}}{\partial x'_{\mu_2}} \dots \frac{\partial x_{\beta_s}}{\partial x'_{\mu_s}} T_{\beta_1, \beta_2, \dots, \beta_s}^{\alpha_1, \alpha_2, \dots, \alpha_r}$$

where $\left| \frac{\partial x}{\partial x'} \right|$ represents a Jacobian and N is the weight of a tensor field.

In this equation $T_{\mu_1, \mu_2, \dots, \mu_s}^{\lambda_1, \lambda_2, \dots, \lambda_r}$ are the components of a tensor in K'

system of reference while $T_{\beta_1, \beta_2, \dots, \beta_s}^{\alpha_1, \alpha_2, \dots, \alpha_r}$ are those in the system of

reference K . Hence if $T_{\beta_1, \beta_2, \dots, \beta_s}^{\alpha_1, \alpha_2, \dots, \alpha_r}$ are zero then $T_{\mu_1, \mu_2, \dots, \mu_s}^{\lambda_1, \lambda_2, \dots, \lambda_r}$ are also zero.

Similarly its components are zero in all the other coordinate systems.

Problem 18. A quantity $A(\mu, \nu, \sigma, \tau)$ function of coordinates x_i transforms to another system of coordinates as

$$A'(\alpha, \beta, \gamma, \xi) = \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} \frac{\partial x_\sigma}{\partial x'_\gamma} \frac{\partial x_\tau}{\partial x'_\xi} A(\mu, \nu, \sigma, \tau)$$

Is it a tensor? If so give its order and rank.

Yes it is a tensor as it transforms according to the transformation law of mixed tensors of rank 4 having a contravariant order 3 and covariant order 1.

3.8. THE FUNDAMENTAL TENSORS

In §3.3, we have already shown that the interval ds between the two neighbouring events (x_1, x_2, x_3, x_4) and $(x_1+dx_1, x_2+dx_2, x_3+dx_3, x_4+dx_4)$ considered in Minkowski's four dimensional space, is given by

$$ds^2 = g_{\mu\nu} dx_\mu dx_\nu \quad (\mu, \nu = 1, 2, 3, 4)$$

(in summation convention by §3.3 (2))

and

$$g_{\mu\nu} = g_{\nu\mu}$$

A convenient way of writing it, is adopted as

$$ds^2 = g_{\mu\nu} (dx)^\mu (dx)^\nu. \quad \dots (1)$$

Now ds^2 being independent of the coordinate system, is an invariant or tensor of zero rank and therefore the equation (1) shows that $g_{\mu\nu} (dx)^\mu$ multiplied by an arbitrary contravariant vector $(dx)^\nu$ always gives a tensor of zero rank; hence $g_{\mu\nu} (dx)^\mu$ is a vector since the dot product of two vectors gives a scalar *i.e.* a tensor of zero rank. Again $g_{\mu\nu}$ multiplied by an arbitrary contravariant vector $(dx)^\mu$ always gives a vector and hence $g_{\mu\nu}$ is covariant tensor (by quotient law).

To show that $g_{\mu\nu}$ is a covariant tensor, let us assume that $g_{\mu\nu}$ becomes $g'_{\mu\nu}$ when the coordinate system becomes (x') from (x) . Then, ds^2 being invariant, we have

$$ds^2 = g'_{\mu\nu} (dx')^\mu (dx')^\nu \quad \dots (2)$$

From (1) and (2), it follows that

$$g'_{\mu\nu} (dx')^\mu (dx')^\nu = g_{\mu\nu} (dx)^\mu (dx)^\nu$$

$$\therefore \text{i.e.,} \quad g'_{\mu\nu} \frac{\partial x_\mu'}{\partial x_\alpha} \frac{\partial x_\nu'}{\partial x_\beta} dx_\alpha dx_\beta = g_{\mu\nu} (dx)^\mu (dx)^\nu$$

$$= g_{\alpha\beta} dx_\alpha dx_\beta \text{ since } dx_\mu' = \frac{\partial x_\mu'}{\partial x_\alpha} dx_\alpha$$

and μ, ν are dummy suffixes on the R.H.S.

The equation is true if

$$g'_{\mu\nu} \frac{\partial x_\mu'}{\partial x_\alpha} \frac{\partial x_\nu'}{\partial x_\beta} = g_{\alpha\beta} \quad \dots (3)$$

Interchanging the primed and unprimed letters we get

$$g'_{\alpha\beta} = \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial x_\nu}{\partial x_\beta'} g_{\mu\nu} \quad \dots (4)$$

Let us further assume that g stands for the determinant

$$g = \begin{vmatrix} g_{11} & g_{12} & g_{13} & g_{14} \\ g_{21} & g_{22} & g_{23} & g_{24} \\ g_{31} & g_{32} & g_{33} & g_{34} \\ g_{41} & g_{42} & g_{43} & g_{44} \end{vmatrix} \quad \dots (5)$$

and $g^{\mu\nu}$ is defined as the cofactor of $g_{\mu\nu}$ in determinant (5), divided by *i.e.*,

$$g^{\mu\nu} = \frac{\text{cofactor of } g_{\mu\nu} \text{ in } g}{g}, \quad \dots (6)$$

Considering the inner product $g_{\mu\sigma} g^{\nu\sigma}$ we have,

$$g_{\mu\sigma} g^{\nu\sigma} = g_{\mu 1} g^{\nu 1} + g_{\mu 2} g^{\nu 2} + g_{\mu 3} g^{\nu 3} + g_{\mu 4} g^{\nu 4}$$

which with the help of assumption (6), becomes

$$\begin{aligned} g_{\mu\sigma} g^{\nu\sigma} &= \frac{1}{g} [g_{\mu 1} \cdot \text{cofactor of } g_{\nu 1} \text{ in } g + g_{\mu 2} \cdot \text{cofactor of } g_{\nu 2} \text{ in } g \\ &\quad + g_{\mu 3} \cdot \text{cofactor of } g_{\nu 3} \text{ in } g + g_{\mu 4} \cdot \text{cofactor of } g_{\nu 4} \text{ in } g]. \\ &= \frac{1}{g} \times 0 \text{ i.e. } 0 \text{ when } \mu \neq \nu \\ &= \frac{1}{g} \times g \text{ i.e., } 1 \text{ when } \mu = \nu. \end{aligned}$$

For, when $\mu \neq \nu$, we get a determinant with two rows identical and when $\mu = \nu$, we reproduce the determinant g divided by itself.

We write

$$\left. \begin{aligned} g_{\mu\sigma} g^{\nu\sigma} &= g_{\mu\sigma} g^{\nu\sigma} \\ &= 0 \text{ if } \mu \neq \nu \\ &= 1 \text{ if } \mu = \nu \end{aligned} \right\} \quad \dots (7)$$

It is therefore clear in view of §3.3, that $g_{\mu\nu}$ has not the same property as the substitution operator or the Kronecker delta $\delta_{\mu\nu}$. Thus,

$$g_{\mu\nu} A^\mu = A^\nu + 0 + 0 + 0 \text{ by (7)} \\ = A^\nu. \quad \dots (8)$$

The equation (8) shows that $g_{\mu\nu}$ multiplied by a contravariant vector yields a vector and hence by quotient law, $g_{\mu\nu}$ is a tensor such that its components are the same in all coordinate system, because $g_{\mu\nu}$ becomes $g'_{\mu\nu}$ in another system of coordinates then

$$\begin{aligned} g'_{\mu\nu} &= \frac{\partial x_\alpha}{\partial x_\mu'} \frac{\partial x_\beta}{\partial x_\nu'} g_{\alpha\beta} \\ &= \frac{\partial x_\alpha}{\partial x_\mu'} \frac{\partial x_\beta}{\partial x_\nu'} \text{ since } \frac{\partial x_\alpha}{\partial x_\alpha} = g_{\alpha\beta} \frac{\partial x_\alpha}{\partial x_\beta} \\ &= 0 \text{ if } \mu \neq \nu \\ &= 1 \text{ if } \mu = \nu \end{aligned} \quad \text{by (7)}$$

Hence $g'_{\mu\nu} = g_{\mu\nu}$ ($\mu, \nu = 1, 2, 3, 4$)

Again $g^{\mu\nu}$ has the same meaning as $g_{\mu\nu}$ with $\mu = \nu$ and we have

$$\begin{aligned} g^{\nu\nu} &= g_1^1 + g_2^2 + g_3^3 + g_4^4 \\ &= 1 + 1 + 1 + 1 \text{ by (7)} \\ &= 4. \end{aligned} \quad \dots (10)$$

Further, we shall show that $g^{\nu\sigma}$ is a tensor when $g_{\mu\sigma} g^{\nu\sigma}$ is a tensor. Multiplying the covariant vector $g_{\mu\sigma} A^\mu$ by the vector $g^{\nu\sigma}$, we have

$$\begin{aligned} g_{\mu\sigma} g^{\nu\sigma} A^\mu &= g_{\mu\nu} A^\mu \text{ by (7)} \\ &= A^\nu \text{ by (8)} \end{aligned}$$

which shows that the product is always a vector.

The tensor character of $g^{\mu\nu}$ may also be shown by denoting the covariant vector $g_{\mu\nu} A^\nu$ by B_μ . Then we have

$$g_{11}A^1 + g_{12}A^2 + g_{13}A^3 + g_{14}A^4 = B_1$$

with similar three equations for B_2, B_3 and B_4 .

Solving these four equations by the method of determinants we can easily find

$$A^1 = g^{11}B_1 + g^{12}B_2 + g^{13}B_3 + g^{14}B_4$$

with similar expressions for A^2, A^3 and A^4 .

$$\text{So that } A^\mu + g^{\mu\nu} B_\nu$$

which follows by quotient law that $g^{\mu\nu}$ is a tensor.

Hence we have defined in this article, the three fundamental tensors

$$g_{\mu\nu}, g_\mu^\nu, g^{\mu\nu}$$

of covariant, mixed and contravariant characters, respectively.

Problem 19. Transform $ds^2 = dx^2 + dy^2 + dz^2$ into polar and cylindrical coordinates.

The transformation equation from cartesian to polar coordinates are

$$x = r \sin \theta \cos \phi, y = r \cos \theta \sin \phi, z = r \cos \theta.$$

If a point (x_1, x_2, x_3) in cartesian becomes (x_1', x_2', x_3') in polar system of coordinates, then

$$x_1 = x, x_2 = y, x_3 = z$$

$$\text{and } x_1' = r, x_2' = \theta, x_3' = \phi.$$

$$\text{So that } g_{11} = g_{22} = g_{33} = 1$$

$$\text{and } g_{12} = g_{13} = g_{23} = 0.$$

Now, from equation (4) of §3.8, we have

$$g'_{\alpha\beta} = \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} g_{\mu\nu}$$

$$\begin{aligned} \therefore g'_{11} &= \frac{\partial x_\mu}{\partial x'_1} \frac{\partial x_\nu}{\partial x'_1} g_{\mu\nu} \\ &= \frac{\partial x_1}{\partial x'_1} \frac{\partial x_1}{\partial x'_1} g_{11} + \frac{\partial x_2}{\partial x'_1} \frac{\partial x_2}{\partial x'_1} g_{22} + \frac{\partial x_3}{\partial x'_1} \frac{\partial x_3}{\partial x'_1} g_{33} \\ &= \left(\frac{\partial x}{\partial r} \right)^2 + \left(\frac{\partial y}{\partial r} \right)^2 + \left(\frac{\partial z}{\partial r} \right)^2 \end{aligned}$$

$$\begin{aligned} &\text{as } x_1 = x \text{ etc. and } g_{11} = g_{22} = g_{33} = 1 \\ &= \sin^2 \theta \cos^2 \phi + \sin^2 \theta \sin^2 \phi + \cos^2 \theta \\ &= \sin^2 \theta + \cos^2 \theta = 1. \end{aligned}$$

$$\text{Similarly } g'_{22} = \left(\frac{\partial x}{\partial \theta} \right)^2 + \left(\frac{\partial y}{\partial \theta} \right)^2 + \left(\frac{\partial z}{\partial \theta} \right)^2 = r^2$$

$$\text{and } g'_{33} = \left(\frac{\partial x}{\partial \phi} \right)^2 + \left(\frac{\partial y}{\partial \phi} \right)^2 + \left(\frac{\partial z}{\partial \phi} \right)^2 = r^2 \sin^2 \theta$$

Also

$$\begin{aligned} g'_{12} &= \frac{\partial x_\mu}{\partial x'_1} \frac{\partial x_\nu}{\partial x'_2} g_{\mu\nu} \\ &= \frac{\partial x_1}{\partial x'_1} \frac{\partial x_1}{\partial x'_2} g_{11} + \frac{\partial x_2}{\partial x'_1} \frac{\partial x_2}{\partial x'_2} g_{22} + \frac{\partial x_3}{\partial x'_1} \frac{\partial x_3}{\partial x'_2} g_{33} \\ &= \frac{\partial x}{\partial r} \frac{\partial x}{\partial \theta} + \frac{\partial y}{\partial r} \frac{\partial y}{\partial \theta} + \frac{\partial z}{\partial r} \frac{\partial z}{\partial \theta} \\ &= \sin \theta \cos \phi (r \cos \theta \cos \phi) + \sin \theta \sin \phi (r \cos \theta \sin \phi) + \cos \theta (-r \sin \theta) \\ &= r \sin \theta \cos \theta (\cos^2 \phi + \sin^2 \phi) - r \sin \theta \cos \theta \\ &= r \sin \theta \cos \theta - r \sin \theta \cos \theta = 0. \end{aligned}$$

Similarly, $g'_{13} = 0 = g'_{23}$.

Hence $ds'^2 = g'_{\mu\nu} (dx')^\mu (dx')^\nu$ gives

$$\begin{aligned} ds'^2 &= g'_{11} (dx'_1)^2 + g'_{22} (dx'_2)^2 + g'_{33} (dx'_3)^2 \\ &= dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2. \end{aligned}$$

Again the cylindrical transformations are

$$x = r \cos \theta, y = r \sin \theta, z = z,$$

so that

$$x_1 = x, x_2 = y, x_3 = z$$

and

$$x_1' = r, x_2' = \theta, x_3' = z.$$

Giving, $g_{11} = g_{22} = g_{33} = 1$.

Also as above and since $g_{12} = g_{13} = g_{23} = 0$, we have

$$g'_{11} = 1, g'_{22} = r^2, g'_{33} = 1$$

$$g'_{12} = 0 = g'_{13} = g'_{23}.$$

$$\begin{aligned} \text{Hence } ds'^2 &= g'_{11} (dx'_1)^2 + g'_{22} (dx'_2)^2 + g'_{33} (dx'_3)^2 \\ &= dr^2 + r^2 d\theta^2 + dz^2. \end{aligned}$$

Problem 20. Show that $g_{\mu\nu}$ is a covariant tensor of the second order.

Let ds be the distance between two neighbouring points $P(y_\alpha)$ and $Q(y_\alpha + dy_\alpha)$, then

$$ds^2 = \sum_{\beta=1}^n dy_\beta dy_\beta$$

But

$$dy_\beta = \frac{\partial y_\beta}{\partial x_\alpha} dx_\alpha$$

$\therefore$

$$\begin{aligned} ds^2 &= \sum_{\beta=1}^n \left(\frac{\partial y_\beta}{\partial x_\alpha} dx_\alpha \right) \left(\frac{\partial y_\beta}{\partial x_\gamma} dx_\gamma \right) \\ &= g_{\alpha\gamma} dx_\alpha dx_\gamma \end{aligned}$$

where

$$g_{\alpha\gamma} = \sum_{\beta=1}^n \frac{\partial y_\beta}{\partial x_\alpha} \frac{\partial y_\beta}{\partial x_\gamma} = \frac{\partial y_\beta}{\partial x_\alpha} \frac{\partial y_\beta}{\partial x_\gamma}$$

Hence in new coordinates,

$$g'_{\xi\eta} = \frac{\partial y_\beta}{\partial x'_\xi} \frac{\partial y_\beta}{\partial x'_\eta}$$

$$\begin{aligned}
&= \frac{\partial y_\beta}{\partial x_\mu} \frac{\partial x_\mu}{\partial x'_\xi} \frac{\partial y_\beta}{\partial x_\nu} \frac{\partial x_\nu}{\partial x'_\eta} \\
&= \frac{\partial x_\mu}{\partial x'_\xi} \frac{\partial x_\nu}{\partial x'_\eta} \frac{\partial y_\beta}{\partial x_\mu} \frac{\partial y_\beta}{\partial x_\nu} \\
&= \frac{\partial x_\mu}{\partial x'_\xi} \frac{\partial x_\nu}{\partial x'_\eta} g_{\mu\nu}
\end{aligned}$$

which follows that $g_{\mu\nu}$ is covariant tensor of second order.

Problem 21. Show that $g_{\alpha\beta} dx_\alpha dx_\beta$ is an invariant

$$g'_{\alpha\beta} dx'_\alpha dx'_\beta = g_{\alpha\beta} dx_\alpha dx_\beta.$$

We have $g_{\mu\nu} = g'_{\alpha\beta} \frac{\partial x'_\alpha}{\partial x_\mu} \frac{\partial x'_\beta}{\partial x_\nu}$

or $\left(g'_{\alpha\beta} \frac{\partial x'_\alpha}{\partial x_\mu} \frac{\partial x'_\beta}{\partial x_\nu} - g_{\mu\nu} \right) = 0$

or $\left(g'_{\alpha\beta} \frac{\partial x'_\alpha}{\partial x_\mu} \frac{\partial x'_\beta}{\partial x_\nu} - g_{\mu\nu} \right) dx_\mu dx_\nu = 0,$

since dx_i is arbitrary

or $g'_{\alpha\beta} \frac{\partial x'_\alpha}{\partial x_\mu} \frac{\partial x'_\beta}{\partial x_\nu} dx_\mu dx_\nu = g_{\mu\nu} dx_\mu dx_\nu$

or $g'_{\alpha\beta} dx'_\alpha dx'_\beta = g_{\mu\nu} dx_\mu dx_\nu$ by (1) of §3.2

which shows that $g_{\alpha\beta} dx_\alpha dx_\beta$ is an invariant.

3.9. ASSOCIATED TENSORS: RAISING AND LOWERING OF SUFFIXES

The process of raising and lowering of the suffices of a tensor is employed to obtain the new tensors associated with the given tensors. In fact this process of raising and lowering the suffices of a tensor changes a covariant suffix into a contravariant suffix and vice versa. These operations merely depend on the inner product of the given tensor with a fundamental tensor.

We define the raising of the suffix of a vector (i.e., a tensor of rank one) by

$$A^\mu = g^{\mu\nu} A_\nu \quad \dots (1)$$

and the lowering of the suffix of a vector, by

$$A_\mu = g_{\mu\nu} A^\nu. \quad \dots (2)$$

In general for a tensor like $A^{\gamma\delta}_{\alpha\beta\mu}$, the operation of raising μ is defined as $A^{\gamma\delta\mu}_{\alpha\beta} = g^{\mu\nu} A^{\gamma\delta}_{\alpha\beta\nu}$ $\dots (3)$

and that of lowering μ as

$$A^{\gamma\delta}_{\alpha\beta\mu} = g_{\mu\nu} A^{\gamma\delta\nu}_{\alpha\beta} \quad \dots (4)$$

The definitions are quite consistent; for, if we first raise a suffix and then lower it, the original tensor is reproduced, e.g., multiplication of (3) by $g_{\mu\sigma}$ in order to lower the suffix on the left, yields

$$\begin{aligned}
g_{\mu\sigma} A^{\alpha\delta\mu}_{\alpha\beta} &= g_{\mu\sigma} g^{\mu\nu} A^{\gamma\delta}_{\alpha\beta\nu} \\
&= g_{\sigma\nu} A^{\gamma\delta}_{\alpha\beta\nu} \text{ by (7) of §3.8} \\
&= A^{\gamma\delta}_{\alpha\beta\sigma} \text{ by (8) §3.8}
\end{aligned}$$

which verifies (4).

It is however worth notable that the raising of a suffix ν by means of $g^{\mu\nu}$ is accompanied by the substitution of μ for ν and the operation of plain substitution of μ for ν is carried by $g_{\mu}{}^\nu$; conclusively,

multiplication by $g^{\mu\nu}$ yields substitution with raising,
multiplication by $g_{\mu}{}^\nu$ yields plain substitution, and
multiplication by $g_{\mu\nu}$ yields substitution with lowering.

The operation of raising and lowering of suffices is applicable to a tensor of any type. For example if $A^{\alpha\beta}$ is a contravariant tensor of rank, two, then

$$A_{\mu}{}^{\alpha} = g_{\mu\nu} B^{\alpha\nu} \text{ so that second has been lowered}$$

$$A_{\nu\beta} = g_{\mu\nu} A^{\mu\beta} \text{ so that first has been lowered}$$

$$A_{\mu\nu} = g_{\mu\alpha} g_{\nu\beta} A^{\alpha\beta} \text{ so that both have been lowered.}$$

Similarly if $B_{\alpha\beta}$ is a covariant tensor of rank two, then

$$B_{\alpha}{}^{\mu} = g^{\mu\nu} B_{\alpha\nu} \text{ so that second has been raised,}$$

$$B_{\beta}{}^{\nu} = g^{\mu\nu} B_{\alpha\beta} \text{ so that first has been raised,}$$

$$B^{\mu\nu} = g^{\mu\alpha} g^{\nu\beta} B_{\alpha\beta} \text{ so that both have been raised.}$$

If two tensors are such that either of them can be obtained from the other by any combination of the operation of raising and lowering the suffices, then these tensors are known as **associated tensors**.

Problem 22. Show that for a rectangular system of co-ordinates the raising and lowering of a suffix leaves the components unaltered in three-dimensional space.

We have in ordinary three-dimensional space.

$$ds^2 = dx_1^2 + dx_2^2 + dx_3^2$$

where

$$x_1 = x, x_2 = y, x_3 = z$$

so that

$$g_{11} = g_{22} = g_{33} = 1$$

and

$$g_{12} = g_{13} = g_{23} = 0.$$

$$\therefore g = \begin{vmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{vmatrix} = \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 1.$$

We thus have

$$g^{11} = \frac{\text{cofactor of } g_{11} \text{ in } g}{g} = \frac{\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix}}{1} = 1.$$

Similarly
which follows that

$$g^{22}=g^{33}=1,$$

$$g_{\mu\nu}=g^{\mu\nu}=g_{\mu}{}^{\nu}$$

so that all three tensors are merely substitution operators. Hence the proposition.

Problem 23. To show that dummy suffixes have a certain freedom of movement between the tensor-factors of an expression, prove that

$$(i) A_{\alpha\beta} B^{\alpha\beta} = A^{\alpha\beta} B_{\alpha\beta},$$

$$(ii) A_{\mu\alpha} B^{\nu\alpha} = A_{\mu}{}^{\alpha} B_{\alpha}{}^{\nu}.$$

(f) Since we have

$$A_{\mu\nu} = g_{\mu\alpha} g_{\nu\beta} A^{\alpha\beta}$$

$$B^{\mu\nu} = g^{\mu\alpha} g^{\nu\beta} B_{\alpha\beta}.$$

and

Therefore

$$A_{\mu\nu} B^{\mu\nu} = g_{\mu\alpha} g_{\nu\beta} A^{\alpha\beta} g^{\mu\alpha} g^{\nu\beta} B_{\alpha\beta}$$

or

$$A_{\mu\nu} B^{\mu\nu} = g^{\mu\alpha} g_{\nu\beta} g^{\nu\beta} g_{\mu\alpha} A^{\alpha\beta} B_{\alpha\beta} = A^{\alpha\beta} B_{\alpha\beta}$$

So that

$$A_{\alpha\beta} B^{\alpha\beta} = A^{\alpha\beta} B_{\alpha\beta}.$$

(ii) Since $g^{\mu\alpha} B_{\alpha}{}^{\nu} = B^{\nu\mu}$.

$$\therefore g_{\mu}{}^{\alpha} g^{\mu\alpha} B_{\alpha}{}^{\nu} = g_{\mu}{}^{\alpha} B^{\nu\mu} = B^{\nu\alpha}$$

and similarly $g_{\nu\alpha} A_{\mu}{}^{\alpha} = A_{\mu\nu}$ gives

$$g_{\alpha}{}^{\nu} g_{\nu\alpha} A_{\mu}{}^{\alpha} = g_{\alpha}{}^{\nu} A_{\mu\nu} = A_{\mu\alpha}$$

$$A_{\mu\alpha} B^{\nu\alpha} = g_{\alpha}{}^{\nu} g_{\nu\alpha} A_{\mu}{}^{\alpha} g_{\mu}{}^{\alpha} g^{\mu\alpha} B_{\alpha}{}^{\nu}$$

$$= A_{\mu}{}^{\alpha} B_{\alpha}{}^{\nu}.$$

3.10. LENGTH OF A VECTOR (i.e. A TENSOR OF RANK ONE), ANGLE BETWEEN TWO VECTORS AND ORTHOGONALITY OF VECTORS

We deal in elementary vector algebra that the square of the length of the vector is its scalar-product into itself and the two vectors are mutually perpendicular if their scalar product is zero. Thus if A_{μ} , B_{ν} be two vectors, then condition of their orthogonality is

$$A_{\mu} B^{\mu} = 0. \quad \dots (1)$$

Also if l is the length of the vector A_{μ} (or A^{μ}), then

$$A_{\mu} A^{\mu} = l^2. \quad \dots (2)$$

In case a vector is self-perpendicular, its length will be zero, i.e.

$$A_{\mu} A^{\mu} = 0. \quad \dots (3)$$

Now we have

$$\begin{aligned} ds^2 &= g_{\mu\nu} (dx)^{\mu} (dx)^{\nu} \\ &= (g_{\mu\nu} dx^{\mu}) (dx)^{\nu} \\ &= dx_{\nu} dx^{\nu}. \end{aligned} \quad \dots (4)$$

$\therefore$ the displacement is self perpendicular if

$$\begin{aligned} dx_{\nu} dx^{\nu} &= 0, \\ ds &= 0. \end{aligned}$$

i.e.

It shows that a displacement is self-perpendicular when it is along a light track $ds=0$.

Now, if θ is the angle between the two vectors A_{μ} and B_{μ} , then

$$\begin{aligned} \cos \theta &= \frac{\text{scalar product of the two vectors}}{\sqrt{(\text{length of } A_{\mu})} \sqrt{(\text{length of } B_{\mu})}} \\ &= \frac{A_{\mu} B^{\mu}}{\sqrt{(A_{\alpha} A^{\alpha})} \sqrt{(B_{\beta} B^{\beta})}} = \frac{A_{\mu} B^{\mu}}{\sqrt{(A_{\alpha} A^{\alpha}) (B_{\beta} B^{\beta})}} \quad \dots (5) \end{aligned}$$

11. METRIC TENSOR, RIEMANNIAN SPACES

We have

$$ds^2 = dx_{\alpha} dx^{\alpha}, \text{ by } \S 3.10(4) \quad \dots (1)$$

where dx^{α} with the help of transformation (1) of § 3.2 can be expressed as

$$dx^{\alpha} = \frac{\partial x^{\alpha}}{\partial x'_{\mu}} dx'^{\mu}$$

Similarly

$$dx_{\alpha} = \frac{\partial x_{\alpha}}{\partial x'_{\mu}} dx'^{\mu} \text{ as } dx_{\nu} \equiv (dx)^{\nu}.$$

With the help of these transformations (1) can be written as

$$ds^2 = \frac{\partial x_{\alpha}}{\partial x'_{\mu}} \frac{\partial x_{\alpha}}{\partial x'_{\nu}} dx'^{\mu} dx'^{\nu} \quad \dots (2)$$

But

$$ds^2 = g_{\mu\nu} dx'^{\mu} dx'^{\nu}. \quad \dots (3)$$

Comparison of (2) and (3) yields

$$g_{\mu\nu} = \frac{\partial x_{\alpha}}{\partial x'_{\mu}} \frac{\partial x_{\alpha}}{\partial x'_{\nu}} \quad \dots (4)$$

which transforms as a tensor when a coordinate system changes from x to x' . Therefore

$$\left. \begin{aligned} \frac{\partial x_{\alpha}}{\partial x'_{\mu}} \frac{\partial x_{\alpha}}{\partial x'_{\nu}} &= \frac{\partial x_{\alpha}}{\partial x'_{\mu}} \frac{\partial x'_{\mu}}{\partial x''_{\beta}} \frac{\partial x_{\alpha}}{\partial x'_{\nu}} \frac{\partial x'_{\nu}}{\partial x''_{\gamma}} \\ g'_{\beta\gamma} &= \frac{\partial x'_{\mu}}{\partial x''_{\beta}} \frac{\partial x'_{\nu}}{\partial x''_{\gamma}} g_{\mu\nu} \end{aligned} \right\} \quad \dots (5)$$

Here $g_{\mu\nu}$ is a covariant symmetric tensor of rank two. It is called the metric tensor.

There are 'spaces' where we cannot introduce a cartesian coordinate system, e.g. two dimensional 'spaces' of that kind which include the surface of a sphere. Introduction of a coordinate system with latitude ϕ and longitude θ , makes it possible to express the distance between two neighbouring points on the spherical surface, in terms of their coordinate differentials, like

$$ds^2 = R^2 (d\phi^2 + \cos^2 \phi d\theta^2).$$

(On assuming equator as $\theta=0$, north pole as $\theta=\frac{\pi}{2}$, south pole as

$$\theta=\frac{\pi}{2})$$

Whenever we define a 'squared infinitesimal distance' which is a homogenous quadratic function of the coordinate differentials, the manifold is called a **metric space** or a **Riemannian space**. In case it is possible to introduce in a Riemannian space a coordinate system with respect to which the components of the metric tensor take the values $\delta_{\mu\alpha}$ at every point, the coordinate system is a cartesian one and the space is called a **Euclidean Space**. In fact Euclidean spaces are the particular cases of Riemannian Spaces.

In all cases where the infinitesimal distance is expressed by equation (3), ds^2 being an invariant, $g_{\mu\nu}$ is always a covariant tensor.

Problem 24. A hyper surface in a Riemannian space is given by

$$x_i = x_i(u_1, u_2):$$

then prove that the coordinate curve of the surface is given by

$$ds^2 = h_{\mu\nu} du^\mu du^\nu$$

$$\text{where } h_{\mu\nu} = g_{\alpha\beta} \frac{\partial x_\alpha}{\partial u_\mu} \frac{\partial x_\beta}{\partial u_\nu}.$$

We have

$$\begin{aligned} ds^2 &= g_{\alpha\beta} dx^\alpha dx^\beta \\ &= g_{\alpha\beta} \frac{\partial x_\alpha}{\partial u^\mu} du^\mu \frac{\partial x_\beta}{\partial u^\nu} du^\nu \\ &= g_{\alpha\beta} \frac{\partial x_\alpha}{\partial u^\mu} \frac{\partial x_\beta}{\partial u^\nu} du^\mu du^\nu \\ &= h_{\mu\nu} du^\mu du^\nu. \end{aligned}$$

Problem 25. Surface of a sphere is a two-dimensional Riemannian space. Find its fundamental metric tensor. (Agra, 1974)

If a be the fixed radius of a sphere, then its surface is given by

$$ds^2 = a^2 d\theta^2 + a^2 \sin^2 \theta d\phi^2, \text{ with equator as } \theta = \frac{\pi}{2}, \text{ north pole as } \theta = 0 \text{ and south pole as } \theta = \pi$$

so that

$$g_{11} = a^2, g_{22} = a^2 \sin^2 \theta$$

$$g_{12} = 0 = g_{21}.$$

$$\therefore g = \begin{vmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{vmatrix} = \begin{vmatrix} a^2 & 0 \\ 0 & a^2 \sin^2 \theta \end{vmatrix} = a^4 \sin^2 \theta$$

$$g^{11} = \frac{\text{cofactor of } g_{11}}{g} = \frac{a^2 \sin^2 \theta}{a^4 \sin^2 \theta} = \frac{1}{a^2}$$

$$g^{22} = \frac{\text{cofactor of } g_{22}}{g} = \frac{a^2}{a^4 \sin^2 \theta} = \frac{1}{a^2 \sin^2 \theta}$$

$$g^{12} = 0 = g^{21}.$$

3.12. CHRISTOFFEL'S 3-INDEX SYMBOLS

Here we introduce two expressions (not tensors) known as Christoffel's symbols of the first and second kind. These will be found of great utility throughout our subsequent work.

Christoffel symbol of the first kind

$$\Gamma_{\mu\nu, \sigma} \equiv [\mu\nu, \sigma] = \frac{1}{2} \left(\frac{\partial g_{\mu\sigma}}{\partial x_\nu} + \frac{\partial g_{\nu\sigma}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \right) \quad \dots (1)$$

Christoffel symbol of the second kind

$$\Gamma^\sigma_{\mu\nu} \equiv \{\mu\nu, \sigma\} = \frac{1}{2} g^{\sigma\lambda} \left(\frac{\partial g_{\mu\lambda}}{\partial x_\nu} + \frac{\partial g_{\nu\lambda}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\lambda} \right) \quad \dots (2)$$

It is obvious from these expressions that

$$[\mu\nu, \sigma] = [\nu\mu, \sigma] \quad \text{i.e.} \quad \Gamma_{\mu\nu, \sigma} = \Gamma_{\nu\mu, \sigma} \quad \dots (3)$$

$$\text{and } \{\mu\nu, \sigma\} = \{\nu\mu, \sigma\} \quad \text{i.e.} \quad \Gamma^\sigma_{\mu\nu} = \Gamma^\sigma_{\nu\mu} \quad \dots (4)$$

showing that they are symmetrical with respect to μ and ν .

We also observe that

$$\{\mu\nu, \sigma\} = g^{\sigma\lambda} [\mu\nu, \lambda] \quad \text{i.e.} \quad \Gamma^\sigma_{\mu\nu} = g^{\sigma\lambda} \Gamma_{\mu\nu, \lambda} \quad \dots (5)$$

$$\text{and } [\mu\nu, \sigma] = g_{\sigma\lambda} \{\mu\nu, \lambda\} \quad \text{i.e.} \quad \Gamma_{\mu\nu, \sigma} = g_{\sigma\lambda} \Gamma^\lambda_{\mu\nu} \quad \dots (6)$$

To prove the result (5), we have from (1), on replacing σ by λ ,

$$[\mu\nu, \lambda] = \frac{1}{2} \left(\frac{\partial g_{\mu\lambda}}{\partial x_\nu} + \frac{\partial g_{\nu\lambda}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\lambda} \right).$$

Multiplying both sides by $g^{\sigma\lambda}$, this becomes

$$\begin{aligned} g^{\sigma\lambda} [\mu\nu, \lambda] &= \frac{1}{2} g^{\sigma\lambda} \left(\frac{\partial g_{\mu\lambda}}{\partial x_\nu} + \frac{\partial g_{\nu\lambda}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\lambda} \right) \\ &= \{\mu\nu, \sigma\} \text{ from (2).} \end{aligned}$$

This proves the result (5).

Again to prove the result (6), interchanging λ and σ in (2), we have

$$\begin{aligned} \{\mu\nu, \lambda\} &= \frac{1}{2} g^{\lambda\sigma} \left(\frac{\partial g_{\mu\sigma}}{\partial x_\nu} + \frac{\partial g_{\nu\sigma}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \right) \\ &= g^{\lambda\sigma} [\mu\nu, \sigma], \text{ from (1).} \end{aligned}$$

Multiplying both sides by $g_{\sigma\lambda}$, this becomes

$$\begin{aligned} g_{\sigma\lambda} \{\mu\nu, \lambda\} &= g_{\sigma\lambda} g^{\lambda\sigma} [\mu\nu, \sigma] \\ &= [\mu\nu, \sigma], \end{aligned}$$

which proves the result (6).

Now we have from (1),

$$\begin{aligned} [\mu\nu, \sigma] + [\sigma\nu, \mu] &= \frac{1}{2} \left(\frac{\partial g_{\mu\sigma}}{\partial x_\nu} + \frac{\partial g_{\nu\sigma}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \right) \\ &\quad + \frac{1}{2} \left(\frac{\partial g_{\mu\sigma}}{\partial x_\nu} + \frac{\partial g_{\nu\sigma}}{\partial x_\mu} - \frac{\partial g_{\mu\sigma}}{\partial x_\nu} \right) \\ &= \frac{1}{2} \left(\frac{\partial g_{\mu\sigma}}{\partial x_\nu} + \frac{\partial g_{\nu\sigma}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\sigma} + \frac{\partial g_{\mu\sigma}}{\partial x_\nu} + \frac{\partial g_{\nu\sigma}}{\partial x_\mu} - \frac{\partial g_{\mu\sigma}}{\partial x_\nu} \right) \\ &= \frac{\partial g_{\mu\sigma}}{\partial x_\nu}, \quad \dots (7) \end{aligned}$$

i.e. $\Gamma_{\mu\nu, \sigma} + \Gamma_{\sigma\nu, \mu} = \frac{\partial g_{\mu\sigma}}{\partial x_\nu}$.

Problem 26. Find the Christoffel's symbols corresponding to (Agra, 1974)

(a) $ds^2 = a^2 d\theta^2 + a^2 \sin^2 \theta d\phi^2$.

(b) $ds^2 = dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2$.

(a) We have $ds^2 = a^2 d\theta^2 + a^2 \sin^2 \theta d\phi^2$,

so that $g_{11} = a^2, g_{22} = a^2 \sin^2 \theta, g_{12} = 0 = g_{21}$

$\therefore g = \begin{vmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{vmatrix} = \begin{vmatrix} a^2 & 0 \\ 0 & a^2 \sin^2 \theta \end{vmatrix} = a^4 \sin^2 \theta$,

giving $g^{11} = \frac{\text{cofactor of } g_{11}}{g} = \frac{a^2 \sin^2 \theta}{a^4 \sin^2 \theta} = \frac{1}{a^2}$.

Similarly, $g^{22} = \frac{1}{a^2 \sin^2 \theta}$ and $g^{12} = 0 = g^{21}$.

Thus Christoffel symbols of first kind are

$[22, 1] = \frac{1}{2} \left(\frac{\partial g_{21}}{\partial x_2} + \frac{\partial g_{21}}{\partial x_2} - \frac{\partial g_{22}}{\partial x_1} \right)$, where $x_1 = \theta, x_2 = \phi$ etc.

$= \frac{1}{2} \left\{ 0 + 0 - \frac{\partial}{\partial \theta} (a^2 \sin^2 \theta) \right\}$

$= -a^2 \sin \theta \cos \theta$.

Similarly, $[12, 2] = \frac{1}{2} \frac{\partial g_{22}}{\partial x_1} = \frac{1}{2} \frac{\partial}{\partial \theta} (a^2 \sin^2 \theta) = a^2 \sin \theta \cos \theta$.

The rest of all are zero, and the Christoffel symbols of second kind are

$\{22, 1\} = g^{1\lambda} [22, \lambda]$

$= g^{11} [22, 1] + g^{12} [22, 2]$

$= \frac{1}{a^2} (-a^2 \sin \theta \cos \theta) + 0 = -\sin \theta \cos \theta$.

Similarly, $\{12, 2\} = g^{2\lambda} [12, \lambda]$

$= g^{21} [12, 2] + g^{22} [12, 2]$

$= 0 + \frac{1}{a^2 \sin^2 \theta} (a^2 \sin \theta \cos \theta) = \cot \theta$

and the rest of all are zero.

(b) We have $ds^2 = dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2$,

where

$x_1 = r, x_2 = \theta, x_3 = \phi$

and

$g_{11} = 1, g_{22} = r^2, g_{33} = r^2 \sin^2 \theta$,

$g_{12} = 0 = g_{13} = \dots$ etc.

$\therefore$

$g = \begin{vmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{vmatrix}$

$= \begin{vmatrix} 1 & 0 & 0 \\ 0 & r^2 & 0 \\ 0 & 0 & r^2 \sin^2 \theta \end{vmatrix} = r^4 \sin^2 \theta$,

giving

$g^{11} = \frac{\text{cofactor of } g_{11}}{g} = \frac{\begin{vmatrix} r^2 & 0 \\ 3 & r^2 \sin^2 \theta \end{vmatrix}}{r^4 \sin^2 \theta} = \frac{r^4 \sin^2 \theta}{r^4 \sin^2 \theta} = 1$.

Similarly $g^{22} = \frac{\text{cofactor of } g_{22}}{g} = \frac{r^2 \sin^2 \theta}{r^4 \sin^2 \theta} = \frac{1}{r^2}$,

$g^{33} = \frac{\text{cofactor of } g_{33}}{g} = \frac{r^2}{r^4 \sin^2 \theta} = \frac{1}{r^2 \sin^2 \theta}$

and the rest all are zero.

Now the Christoffel's symbols of the first kind are

$[11, 1] = \frac{1}{2} \left(\frac{\partial g_{11}}{\partial x_1} + \frac{\partial g_{11}}{\partial x_1} - \frac{\partial g_{11}}{\partial x_1} \right) = 0$.

Similarly

$[11, 2] = 0 = [11, 3]$.

In a similar manner, it is easy to show that

$[22, 1] = -r, [22, 2] = 0 = [22, 3]$

$[33, 1] = -r \sin^2 \theta, [33, 2] = -r^2 \sin \theta \cos \theta, [33, 3] = 0$

$[12, 1] = 0 = [21, 1], [13, 1] = 0 = [31, 1]$

$[12, 2] = 0 = [21, 2], [12, 3] = 0 = [21, 3]$

$[13, 1] = 0 = [31, 1], [13, 2] = 0 = [31, 2]$

$[13, 3] = r \sin^2 \theta = [31, 3], [23, 1] = 0 = [32, 1]$

$[23, 2] = 0 = [32, 2], [23, 3] = r^2 \sin \theta \cos \theta = [32, 3]$

and the Christoffel symbols of the second kind are

$\{22, 1\} = g^{1\lambda} [22, \lambda]$

$= g^{11} [22, 1] + g^{12} [22, 2] + g^{13} [22, 3]$

$= -r + 0 + 0 = -r$.

Similarly $\{3, 3, 1\} = -r \sin^2 \theta, \{11, 1\} = 0$,

$\{33, 2\} = -\sin \theta \cos \theta, \{13, 3\} = \frac{1}{2}, \{23, 3\} = \cot \theta$

and the rest all are zero.

13. EQUATIONS OF A GEODESIC

(Agra, 1963, 65, 66)

We sometimes define a geodesic as the path of shortest distance on the surface between two given points on it. Here our aim is to determine the equations of a geodesic or path between two points for which

$\int ds$ is stationary.

Assuming that the initial and terminal points of the path are fixed, let the path be deformed by giving every intermediate point an arbitrary infinitesimal displacement δx_σ , so that the expression

$ds^2 = g_{\mu\nu} dx_\mu dx_\nu$ yields,

$2ds \delta(ds) = \delta(g_{\mu\nu}) dx_\mu dx_\nu + g_{\mu\nu} \delta(dx_\mu) dx_\nu + g_{\mu\nu} dx_\mu \delta(dx_\nu)$

$= dx_\mu dx_\nu \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \delta x_\sigma + g_{\mu\nu} dx_\nu \delta(dx_\mu) + g_{\mu\nu} dx_\mu \delta(dx_\nu)$

The stationary condition is

$$\int \delta(ds) = 0. \quad \dots (2)$$

Substituting the value of $\delta(ds)$ from (1) in (2), we get

$$\frac{1}{2} \int \left\{ \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \delta x_\sigma + g_{\mu\nu} \frac{dx_\nu}{ds} \frac{d}{ds} (\delta x_\mu) + g_{\mu\nu} \frac{dx_\mu}{ds} \frac{d}{ds} (\delta x_\nu) \right\} ds = 0$$

On changing the dummy suffixes in the last two terms, this becomes

$$\frac{1}{2} \int \left\{ \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \delta x_\sigma + \left(g_{\sigma\nu} \frac{dx_\nu}{ds} + g_{\mu\sigma} \frac{dx_\mu}{ds} \right) \frac{d}{ds} (\delta x_\sigma) \right\} ds = 0$$

or $\frac{1}{2} \int \left(\frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \delta x_\sigma + \frac{d}{ds} \left(g_{\sigma\nu} \frac{dx_\nu}{ds} + g_{\mu\sigma} \frac{dx_\mu}{ds} \right) \delta x_\sigma \right) ds = 0$

Integrating the second terms by parts, we have

$$\frac{1}{2} \int \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \delta x_\sigma ds + \frac{1}{2} \left[\left(g_{\sigma\nu} \frac{dx_\nu}{ds} + g_{\mu\sigma} \frac{dx_\mu}{ds} \right) \delta x_\sigma \right]_0^1 - \frac{1}{2} \int \frac{d}{ds} \left(g_{\sigma\nu} \frac{dx_\nu}{ds} + g_{\mu\sigma} \frac{dx_\mu}{ds} \right) (\delta x_\sigma) ds = 0,$$

where the integrated part vanishes as δx_σ vanishes at both the limits.

We are, therefore left with

$$\frac{1}{2} \int \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \delta x_\sigma ds - \frac{1}{2} \int \frac{d}{ds} \left(g_{\sigma\nu} \frac{dx_\nu}{ds} + g_{\mu\sigma} \frac{dx_\mu}{ds} \right) (\delta x_\sigma) ds = 0$$

or $\frac{1}{2} \int \left[\frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} - \frac{d}{ds} \left(g_{\sigma\nu} \frac{dx_\nu}{ds} + g_{\mu\sigma} \frac{dx_\mu}{ds} \right) \right] \delta(x_\sigma) ds = 0.$

This will hold for all values of the arbitrary displacement (δx_σ) at all points if

$$\frac{1}{2} \left[\frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} - \frac{d}{ds} \left(g_{\sigma\nu} \frac{dx_\nu}{ds} + g_{\mu\sigma} \frac{dx_\mu}{ds} \right) \right] = 0$$

or if $\frac{1}{2} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} - \frac{1}{2} \frac{dg_{\sigma\nu}}{ds} \frac{dx_\nu}{ds} - \frac{1}{2} g_{\sigma\nu} \frac{d^2 x_\nu}{ds^2} - \frac{1}{2} \frac{dg_{\mu\sigma}}{ds} \frac{dx_\mu}{ds} - \frac{1}{2} g_{\mu\sigma} \frac{d^2 x_\mu}{ds^2} = 0.$

But we have

$$\frac{dg_{\sigma\nu}}{ds} = \frac{\partial g_{\sigma\nu}}{\partial x_\mu} \frac{dx_\mu}{ds} \text{ and } \frac{dg_{\mu\sigma}}{ds} = \frac{\partial g_{\mu\sigma}}{\partial x_\nu} \frac{dx_\nu}{ds}$$

so that the last relation becomes

$$\frac{1}{2} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} - \frac{1}{2} \frac{\partial g_{\sigma\nu}}{\partial x_\mu} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} - \frac{1}{2} \frac{\partial g_{\mu\sigma}}{\partial x_\nu} \frac{dx_\nu}{ds} \frac{dx_\mu}{ds} - \frac{1}{2} g_{\sigma\nu} \frac{d^2 x_\nu}{ds^2} - \frac{1}{2} g_{\mu\sigma} \frac{d^2 x_\mu}{ds^2} = 0$$

$$\text{or } \frac{1}{2} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \left(\frac{\partial g_{\mu\nu}}{\partial x_\sigma} - \frac{\partial g_{\sigma\nu}}{\partial x_\mu} - \frac{\partial g_{\mu\sigma}}{\partial x_\nu} \right) - \frac{1}{2} \left(g_{\sigma\nu} \frac{d^2 x_\nu}{ds^2} + g_{\mu\sigma} \frac{d^2 x_\mu}{ds^2} \right) = 0$$

Replacing the dummy suffixes μ and ν by ε in the last term, we get,

$$\frac{1}{2} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \left(\frac{\partial g_{\mu\nu}}{\partial x_\sigma} - \frac{\partial g_{\sigma\nu}}{\partial x_\mu} - \frac{\partial g_{\mu\sigma}}{\partial x_\nu} \right) - \frac{1}{2} \left(g_{\sigma\varepsilon} \frac{d^2 x_\varepsilon}{ds^2} + g_{\varepsilon\sigma} \frac{d^2 x_\varepsilon}{ds^2} \right) = 0.$$

or $\frac{1}{2} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \left(\frac{\partial g_{\mu\nu}}{\partial x_\sigma} - \frac{\partial g_{\nu\sigma}}{\partial x_\mu} - \frac{\partial g_{\mu\sigma}}{\partial x_\nu} \right) + g_{\varepsilon\sigma} \frac{d^2 x_\varepsilon}{ds^2} = 0.$

Multiplying throughout by $-g^{\sigma\alpha}$, this becomes

$$\frac{1}{2} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} g^{\sigma\alpha} \left(\frac{\partial g_{\mu\sigma}}{\partial x_\nu} + \frac{\partial g_{\nu\sigma}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \right) + g^{\sigma\alpha} g_{\varepsilon\sigma} \frac{d^2 x_\varepsilon}{ds^2} = 0$$

or $g_{\varepsilon\sigma} \frac{d^2 x_\varepsilon}{ds^2} + \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \{ \mu\nu, \alpha \} = 0, \quad \dots (3)$

as $g^{\sigma\alpha} g_{\varepsilon\sigma} = g_{\varepsilon\alpha}$

$$\text{or } \left. \begin{aligned} \frac{d^2 x_\alpha}{ds^2} + \{ \mu\nu, \alpha \} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} &= 0 \\ \frac{d^2 x_\alpha}{ds^2} + \Gamma_{\mu\nu}^\alpha \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} &= 0 \end{aligned} \right\} \text{ since } \frac{d^2 x_\alpha}{ds^2} = g_{\varepsilon\sigma} \frac{d^2 x_\varepsilon}{ds^2} \quad \dots (4)$$

which corresponding to $\alpha=1, 2, 3, 4$ gives four equations determining a geodesic.

3.14. LAW OF TRANSFORMATION FOR CHRISTOFFEL'S SYMBOLS

(Agra, 1966, 68)

Considering the two coordinate systems x_α and x'_α in a Riemannian space the equation of geodesic line can be written as

$$\frac{d^2 x_\alpha}{ds^2} + \Gamma_{\beta\gamma}^\alpha \frac{dx_\beta}{ds} \frac{dx_\gamma}{ds} = 0 \quad \dots (1)$$

where

$$\Gamma_{\beta\gamma}^\alpha \equiv \{ \beta\gamma, \alpha \}$$

and

$$\frac{d^2 x'_\alpha}{ds^2} + \Gamma'_{\beta\gamma}^\alpha \frac{dx'_\beta}{ds} \frac{dx'_\gamma}{ds} = 0 \quad \dots (2)$$

where

$$\Gamma'_{\beta\gamma}^\alpha = \{ \beta\gamma, \alpha \}'.$$

Now we have

$$\frac{dx'_\alpha}{ds} = \frac{\partial x'_\alpha}{\partial x_\lambda} \frac{dx_\lambda}{ds} \text{ etc.}$$

and

$$\begin{aligned} \frac{d^2 x'_\alpha}{ds^2} &= \frac{d}{ds} \left(\frac{dx'_\alpha}{ds} \right) = \frac{d}{ds} \left(\frac{\partial x'_\alpha}{\partial x_\lambda} \frac{dx_\lambda}{ds} \right) \\ &= \frac{d}{ds} \left(\frac{\partial x'_\alpha}{\partial x_\lambda} \right) \frac{dx_\lambda}{ds} + \frac{\partial x'_\alpha}{\partial x_\lambda} \frac{d^2 x_\lambda}{ds^2} \\ &= \frac{\partial^2 x'_\alpha}{\partial x_\mu \partial x_\lambda} \frac{dx_\mu}{ds} \frac{dx_\lambda}{ds} + \frac{\partial x'_\alpha}{\partial x_\lambda} \frac{d^2 x_\lambda}{ds^2} \end{aligned}$$

$$= \frac{\partial^2 x'_\alpha}{\partial x_\mu \partial x_\nu} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} + \frac{\partial x'_\alpha}{\partial x_\lambda} \frac{d^2 x_\lambda}{ds^2}$$

(replacing λ by ν in the first term).

Substituting these values in (2), we get

$$\frac{\partial x'_\alpha}{\partial x_\lambda} \frac{d^2 x_\lambda}{ds^2} + \frac{\partial^2 x'_\alpha}{\partial x_\mu \partial x_\nu} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} + \Gamma'_{\beta\gamma} \frac{\partial x_\beta}{\partial x_\mu} \frac{\partial x_\gamma}{\partial x_\nu} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} = 0.$$

Multiplying by $\frac{\partial x_\lambda}{\partial x'_\alpha}$, this equation becomes

$$\frac{d^2 x_\lambda}{ds^2} + \left(\Gamma'_{\beta\gamma} \frac{\partial x'_\beta}{\partial x_\mu} \frac{\partial x'_\gamma}{\partial x_\nu} \frac{\partial x_\lambda}{\partial x'_\alpha} + \frac{\partial^2 x'_\alpha}{\partial x_\mu \partial x_\nu} \frac{\partial x_\lambda}{\partial x'_\alpha} \right) \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} = 0.$$

Interchanging α, β, γ , with λ, μ, ν respectively, this becomes

$$\frac{d^2 x_\alpha}{ds^2} + \left(\Gamma'_{\mu\nu} \frac{\partial x'_\mu}{\partial x_\beta} \frac{\partial x'_\nu}{\partial x_\gamma} \frac{\partial x_\alpha}{\partial x'_\lambda} + \frac{\partial^2 x'_\lambda}{\partial x_\beta \partial x_\gamma} \frac{\partial x_\alpha}{\partial x'_\lambda} \right) \frac{dx_\beta}{ds} \frac{dx_\gamma}{ds} = 0 \quad \dots (3)$$

Comparison of (1) and (3) yields

$$\left. \begin{aligned} \Gamma'_{\beta\gamma} &= \Gamma'_{\mu\nu} \frac{\partial x'_\mu}{\partial x_\beta} \frac{\partial x'_\nu}{\partial x_\gamma} \frac{\partial x_\alpha}{\partial x'_\lambda} + \frac{\partial^2 x'_\lambda}{\partial x_\beta \partial x_\gamma} \frac{\partial x_\alpha}{\partial x'_\lambda} \\ \{\beta\gamma, \alpha\} &= \{\mu\nu, \lambda\} \frac{\partial x'_\mu}{\partial x_\beta} \frac{\partial x'_\nu}{\partial x_\gamma} \frac{\partial x_\alpha}{\partial x'_\lambda} + \frac{\partial^2 x'_\lambda}{\partial x_\beta \partial x_\gamma} \frac{\partial x_\alpha}{\partial x'_\lambda} \end{aligned} \right\} \quad \dots (4)$$

(Agra, 1968)

This gives the law of transformation for $\Gamma'_{\beta\gamma}$, i.e. $\{\beta\gamma, \alpha\}$. It is clear from this transformation that $\Gamma'_{\beta\gamma}$ are not the components of a tensor, so that $\Gamma'_{\beta\gamma}$ may be zero in one coordinate system, but not in all coordinate systems.

We may also arrive at the Christoffel symbols and their law of transformation by another method. We have by §3.8 (4)

$$g'_{\alpha\beta} = g_{\nu\mu} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta}.$$

Differentiating it with respect to x'_γ , we have

$$\frac{\partial g'_{\alpha\beta}}{\partial x'_\gamma} = \frac{\partial g_{\lambda\mu}}{\partial x_\nu} \frac{\partial x_\nu}{\partial x'_\gamma} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} + g_{\lambda\mu} \left(\frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial^2 x_\mu}{\partial x'_\gamma \partial x'_\beta} + \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial^2 x_\lambda}{\partial x'_\gamma \partial x'_\alpha} \right) \quad \dots (5)$$

The other two equations may be obtained from it by cyclic permutations of the indices α, β, γ , and these are

$$\frac{\partial g'_{\beta\gamma}}{\partial x'_\alpha} = \frac{\partial g_{\lambda\mu}}{\partial x_\nu} \frac{\partial x_\nu}{\partial x'_\alpha} \frac{\partial x_\lambda}{\partial x'_\beta} \frac{\partial x_\mu}{\partial x'_\gamma} + g_{\lambda\mu} \left(\frac{\partial x_\lambda}{\partial x'_\beta} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\gamma} + \frac{\partial x_\mu}{\partial x'_\gamma} \frac{\partial^2 x_\lambda}{\partial x'_\alpha \partial x'_\beta} \right) \quad \dots (6)$$

$$\text{and } \frac{\partial g'_{\gamma\alpha}}{\partial x'_\beta} = \frac{\partial g_{\lambda\mu}}{\partial x_\nu} \frac{\partial x_\nu}{\partial x'_\beta} \frac{\partial x_\lambda}{\partial x'_\gamma} \frac{\partial x_\mu}{\partial x'_\alpha} + g_{\lambda\mu} \left(\frac{\partial x_\lambda}{\partial x'_\gamma} \frac{\partial^2 x_\mu}{\partial x'_\beta \partial x'_\alpha} + \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial^2 x_\lambda}{\partial x'_\beta \partial x'_\gamma} \right) \quad \dots (7)$$

Subtracting (5) from the sum of (6) and (7) after changing dummy suffixes in their first terms and then dividing by 2, we get

$$\begin{aligned} \frac{1}{2} \left(\frac{\partial g'_{\beta\gamma}}{\partial x'_\alpha} + \frac{\partial g'_{\gamma\alpha}}{\partial x'_\beta} - \frac{\partial g'_{\alpha\beta}}{\partial x'_\gamma} \right) &= \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x_\nu}{\partial x'_\gamma} \cdot \frac{1}{2} \left(\frac{\partial g_{\mu\nu}}{\partial x_\lambda} + \frac{\partial g_{\gamma\lambda}}{\partial x_\mu} - \frac{\partial g_{\lambda\mu}}{\partial x_\nu} \right) \\ &+ \frac{1}{2} g_{\lambda\mu} \left(\frac{\partial x_\lambda}{\partial x'_\beta} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\gamma} + \frac{\partial x_\lambda}{\partial x'_\gamma} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\beta} + \frac{\partial x_\mu}{\partial x'_\gamma} \frac{\partial^2 x_\lambda}{\partial x'_\alpha \partial x'_\beta} \right. \\ &\left. + \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial^2 x_\lambda}{\partial x'_\beta \partial x'_\gamma} - \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial^2 x_\mu}{\partial x'_\beta \partial x'_\gamma} - \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial^2 x_\lambda}{\partial x'_\alpha \partial x'_\gamma} \right). \end{aligned}$$

In the last term of right hand side, interchanging the dummy suffixes λ, μ wherever necessary in order to make all the second differentials of x_μ only, we get

$$\begin{aligned} \Gamma'_{\alpha\beta, \gamma} &= \Gamma_{\lambda\mu, \nu} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x_\nu}{\partial x'_\gamma} + \frac{1}{2} g_{\lambda\mu} \left(\frac{\partial x_\lambda}{\partial x'_\beta} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\gamma} + \frac{\partial x_\lambda}{\partial x'_\gamma} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\beta} \right. \\ &\left. + \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\beta} \frac{\partial x_\lambda}{\partial x'_\gamma} + \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial^2 x_\mu}{\partial x'_\beta \partial x'_\gamma} + \frac{\partial x_\lambda}{\partial x'_\beta} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\gamma} - \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial^2 x_\mu}{\partial x'_\beta \partial x'_\gamma} \right. \\ &\left. - \frac{\partial x_\lambda}{\partial x'_\beta} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\gamma} \right) \\ &= \Gamma_{\lambda\mu, \nu} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x_\nu}{\partial x'_\gamma} + g_{\lambda\mu} \frac{\partial x_\lambda}{\partial x'_\gamma} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\beta} \quad \dots (8) \end{aligned}$$

Also we have the transformation law for the contravariant fundamental tensor as

$$g'^{\gamma\epsilon} = g^{\mu\sigma} \frac{\partial x'_\gamma}{\partial x_\mu} \frac{\partial x'_\epsilon}{\partial x_\sigma} \quad \dots (9)$$

Multiplying together the corresponding sides of (8) and (9), we get

$$\begin{aligned} g'^{\gamma\epsilon} \Gamma'_{\alpha\beta, \gamma} &= g^{\mu\sigma} \Gamma_{\lambda\mu, \nu} \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x_\nu}{\partial x'_\gamma} \frac{\partial x'_\epsilon}{\partial x_\sigma} \\ &+ g^{\mu\sigma} g_{\lambda\mu} \frac{\partial x_\lambda}{\partial x'_\gamma} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\beta} \frac{\partial x'_\epsilon}{\partial x_\sigma} \\ &= \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x_\epsilon}{\partial x'_\gamma} g^{\mu\sigma} \frac{\partial x_\lambda}{\partial x'_\sigma} \frac{\partial x'_\epsilon}{\partial x_\sigma} \Gamma_{\lambda\mu, \nu} \\ &+ g^{\mu\sigma} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\beta} \frac{\partial x'_\epsilon}{\partial x_\sigma} \frac{\partial x_\lambda}{\partial x'_\gamma} \frac{\partial x'_\epsilon}{\partial x_\sigma} g_{\lambda\mu} \\ &= \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x'_\epsilon}{\partial x_\sigma} g^{\mu\sigma} \Gamma_{\lambda\mu, \epsilon} + \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\beta} g^{\mu\sigma} \frac{\partial x'_\epsilon}{\partial x_\sigma} g_{\lambda\mu} \\ &= \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x'_\epsilon}{\partial x_\sigma} \Gamma_{\lambda\mu}^\sigma + \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\beta} g^{\mu\sigma} \frac{\partial x'_\epsilon}{\partial x_\sigma} \end{aligned}$$

or

$$\Gamma'_{\alpha\beta}^{\gamma\epsilon} = \Gamma_{\lambda\mu}^\sigma \frac{\partial x_\lambda}{\partial x'_\alpha} \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x'_\epsilon}{\partial x_\sigma} + \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\beta} \frac{\partial x'_\epsilon}{\partial x_\sigma} g_{\lambda\mu}.$$

Replacing α by β , β by γ , ϵ by α , λ by μ , μ by ν and σ by λ , this becomes

$$\Gamma'_{\beta\gamma}^{\alpha} = \Gamma_{\mu\nu}^\lambda \frac{\partial x_\mu}{\partial x'_\beta} \frac{\partial x_\nu}{\partial x'_\gamma} \frac{\partial x'_\alpha}{\partial x_\lambda} + \frac{\partial^2 x_\nu}{\partial x'_\beta \partial x'_\gamma} \frac{\partial x'_\alpha}{\partial x_\lambda} g_{\mu\nu}.$$

Again replacing ν by λ in the last term,

$$\left. \begin{aligned} \Gamma_{\beta\gamma}^{\alpha} &= \Gamma_{\mu\nu}^{\lambda} \frac{\partial x_{\mu}}{\partial x_{\beta}} \frac{\partial x_{\nu}}{\partial x_{\gamma}} \frac{\partial x_{\alpha}'}{\partial x_{\lambda}} + \frac{\partial^2 x_{\lambda}}{\partial x_{\beta} \partial x_{\gamma}} \frac{\partial x_{\alpha}'}{\partial x_{\lambda}}, \\ \text{i.e. } \{\beta\gamma, \alpha\}' &= \{\mu\nu, \lambda\} \frac{\partial x_{\mu}}{\partial x_{\beta}} \frac{\partial x_{\nu}}{\partial x_{\gamma}} \frac{\partial x_{\alpha}'}{\partial x_{\lambda}} + \frac{\partial^2 x_{\lambda}}{\partial x_{\beta} \partial x_{\gamma}} \frac{\partial x_{\alpha}'}{\partial x_{\lambda}}, \end{aligned} \right\} \dots (10)$$

Interchanging primed and unprimed letters, this yields

$$\{\beta\gamma, \alpha\} = \{\mu\nu, \lambda\} \frac{\partial x_{\mu}'}{\partial x_{\beta}} \frac{\partial x_{\nu}'}{\partial x_{\gamma}} \frac{\partial x_{\alpha}}{\partial x_{\lambda}'} + \frac{\partial^2 x_{\lambda}'}{\partial x_{\beta} \partial x_{\gamma}} \frac{\partial x_{\alpha}}{\partial x_{\lambda}'},$$

which is the same transformation as obtained in (4).

In the given Euclidean space such as

$$ds^2 = (dx_1)^2 + (dx_2)^2 + \dots + (dx_n)^2,$$

$$\Gamma_{\mu\nu}^{\lambda} = 0.$$

The equation (10), yields

$$\Gamma_{\beta\gamma}^{\alpha} = \frac{\partial^2 x_{\lambda}}{\partial x_{\beta} \partial x_{\gamma}} \frac{\partial x_{\alpha}'}{\partial x_{\lambda}}.$$

If this new system is also cartesian, then $g_{\alpha\beta} = \text{constant}$

$$\text{or } \Gamma_{\beta\gamma}^{\alpha} = 0$$

$$\text{or } \frac{\partial^2 x_{\lambda}}{\partial x_{\beta} \partial x_{\gamma}} = 0 \quad \dots (11)$$

and then the symbols transform like tensor. (Agra, 1965, 66, 67)

Integrating (11) twice, we get

$$x_{\lambda} = a_{\alpha}^{\lambda} x_{\alpha}' + b^{\lambda},$$

where a_{α}^{λ} and b^{λ} are the constants of integration.

This follows that the co-ordinate transformation between two cartesian co-ordinate systems is linear.

Now inner multiplication of (10) by $\frac{\partial x_{\beta}}{\partial x_{\alpha}'}$, gives

$$\left. \begin{aligned} \Gamma_{\beta\lambda}^{\alpha} \frac{\partial x_{\beta}}{\partial x_{\alpha}'} &= \frac{\partial x_{\beta}}{\partial x_{\alpha}'} \Gamma_{\mu\nu}^{\lambda} \frac{\partial x_{\mu}}{\partial x_{\beta}} \frac{\partial x_{\nu}}{\partial x_{\gamma}} \frac{\partial x_{\alpha}'}{\partial x_{\lambda}} + \frac{\partial x_{\beta}}{\partial x_{\alpha}'} \frac{\partial^2 x_{\lambda}}{\partial x_{\beta} \partial x_{\gamma}} \frac{\partial x_{\alpha}'}{\partial x_{\lambda}} \\ &= \frac{\partial x_{\mu}}{\partial x_{\beta}'} \frac{\partial x_{\gamma}}{\partial x_{\alpha}'} \frac{\partial x_{\beta}}{\partial x_{\mu}'} \frac{\partial x_{\alpha}'}{\partial x_{\lambda}'} \Gamma_{\mu\nu}^{\lambda} + \frac{\partial x_{\beta}}{\partial x_{\alpha}'} \frac{\partial x_{\alpha}'}{\partial x_{\lambda}'} \frac{\partial^2 x_{\lambda}}{\partial x_{\beta} \partial x_{\gamma}} \\ &= \frac{\partial x_{\mu}}{\partial x_{\beta}'} \frac{\partial x_{\gamma}}{\partial x_{\alpha}'} \Gamma_{\mu\nu}^{\lambda} + \frac{\partial^2 x_{\beta}}{\partial x_{\beta} \partial x_{\gamma}}, \end{aligned} \right\} \dots (12)$$

$$\text{i.e. } \frac{\partial^2 x_{\beta}}{\partial x_{\beta} \partial x_{\gamma}'} = \Gamma_{\beta\gamma}^{\alpha} \frac{\partial x_{\beta}}{\partial x_{\alpha}'} - \Gamma_{\mu\nu}^{\lambda} \frac{\partial x_{\mu}}{\partial x_{\beta}'} \frac{\partial x_{\gamma}}{\partial x_{\alpha}'}.$$

3.15. PARALLEL DISPLACEMENT OF VECTORS

Consider an absolute contravariant vector A^{μ} ($x_1, x_2, x_3, \dots, x_n$) in a cartesian co-ordinate system. Then, we have

$$A'^{\mu} = A^{\alpha} \frac{\partial x_{\mu}'}{\partial x_{\alpha}} \text{ and } A^{\mu} = A'^{\alpha} \frac{\partial x_{\mu}}{\partial x_{\alpha}'}.$$

$$\therefore dA'^{\mu} = A^{\alpha} \frac{\partial^2 x_{\mu}'}{\partial x_{\beta} \partial x_{\alpha}} \frac{\partial x_{\beta}}{\partial x_{\gamma}'} dx_{\gamma}' + \frac{\partial x_{\mu}'}{\partial x_{\alpha}} dA^{\alpha}.$$

If we now assume that the components of A^{μ} are constants, i.e. $dA^{\mu} = 0$, then the above relation becomes

$$\begin{aligned} dA'^{\mu} &= A^{\alpha} \frac{\partial^2 x_{\mu}'}{\partial x_{\beta} \partial x_{\alpha}} \frac{\partial x_{\beta}}{\partial x_{\gamma}'} dx_{\gamma}' \\ &= A'^{\sigma} \frac{\partial x_{\alpha}}{\partial x_{\sigma}'} \frac{\partial^2 x_{\mu}'}{\partial x_{\beta} \partial x_{\alpha}} \frac{\partial x_{\beta}}{\partial x_{\gamma}'} dx_{\gamma}' \text{ as } A^{\sigma} = A'^{\sigma} \frac{\partial x_{\sigma}}{\partial x_{\sigma}'} \\ &= A'^{\sigma} \frac{\partial^2 x_{\mu}'}{\partial x_{\beta} \partial x_{\alpha}} \frac{\partial x_{\beta}}{\partial x_{\gamma}'} \frac{\partial x_{\alpha}}{\partial x_{\sigma}'}, \quad \dots (1) \end{aligned}$$

From the preceding section we have the law of transformation

$$\Gamma_{\gamma\sigma}^{\mu} = \frac{\partial^2 x_{\alpha}'}{\partial x_{\gamma}' \partial x_{\sigma}'} \frac{\partial x^{\mu}}{\partial x_{\alpha}} + \Gamma_{\beta\gamma}^{\sigma} \frac{\partial x_{\beta}}{\partial x_{\gamma}'} \frac{\partial x_{\gamma}}{\partial x_{\sigma}'} \frac{\partial x^{\mu}}{\partial x_{\alpha}}.$$

But in the cartesian co-ordinate system, we have $\Gamma_{\beta\gamma}^{\alpha} = 0$.

$$\therefore \Gamma_{\gamma\sigma}^{\mu} = \frac{\partial^2 x_{\alpha}'}{\partial x_{\gamma}' \partial x_{\sigma}'} \frac{\partial x^{\mu}}{\partial x_{\alpha}}$$

$$\text{Now } \frac{\partial x_{\mu}'}{\partial x_{\alpha}} \frac{\partial x_{\alpha}}{\partial x_{\sigma}'} = \delta_{\sigma}^{\mu}.$$

Differentiating this equation partially with respect to x_{γ}' , we obtain

$$\begin{aligned} \frac{\partial^2 x_{\mu}'}{\partial x_{\beta} \partial x_{\alpha}} \frac{\partial x_{\beta}}{\partial x_{\gamma}'} \frac{\partial x_{\alpha}}{\partial x_{\sigma}'} &= - \frac{\partial x_{\mu}'}{\partial x_{\alpha}} \frac{\partial^2 x_{\alpha}'}{\partial x_{\gamma}' \partial x_{\sigma}'} \\ \therefore \Gamma_{\gamma\sigma}^{\mu} &= - \frac{\partial^2 x_{\mu}'}{\partial x_{\beta} \partial x_{\alpha}} \frac{\partial x_{\beta}}{\partial x_{\gamma}'} \frac{\partial x_{\alpha}}{\partial x_{\sigma}'} \end{aligned}$$

With the substitution of these values in (1), we have

$$dA'^{\mu} = -A'^{\sigma} \Gamma_{\gamma\sigma}^{\mu} dx_{\gamma}' \quad \dots (2)$$

or interchanging the primed and unprimed letters,

$$dA^{\mu} = -A^{\sigma} \Gamma_{\gamma\sigma}^{\mu} dx_{\gamma}.$$

Thus if A^{μ} is parallelly displaced with respect to any Riemannian V_n along any curve, we have

$$\frac{dA^{\mu}}{ds} = -A^{\sigma} \Gamma_{\gamma\sigma}^{\mu} \frac{dx_{\gamma}}{ds}.$$

Now we write

$$\begin{aligned} \frac{\delta A^{\mu}}{\delta s} &= \frac{dA^{\mu}}{ds} + A^{\sigma} \Gamma_{\gamma\sigma}^{\mu} \frac{dx_{\gamma}}{ds} \\ &= -A^{\sigma} \Gamma_{\gamma\sigma}^{\mu} \frac{dx_{\gamma}}{ds} + A^{\sigma} \Gamma_{\gamma\sigma}^{\mu} \frac{dx_{\gamma}}{ds} \\ &= 0 \end{aligned} \quad \dots (3)$$

which represents a parallel displacement of a vector A^μ along the curve. Here $\frac{\delta A^\mu}{\delta s}$ is known as the **intrinsic derivative** of A^μ w.r.t. s .

But the equation of geodesic is

$$\frac{d^2 x_\mu}{ds^2} + \Gamma_{\alpha\beta}^\mu \frac{dx_\alpha}{ds} \frac{dx_\beta}{ds} = 0$$

or

$$\frac{d}{ds} \left(\frac{dx_\mu}{ds} \right) + \Gamma_{\alpha\beta}^\mu \left(\frac{dx_\alpha}{ds} \right) \frac{dx_\beta}{ds} = 0 \quad \dots (4)$$

So the equation (4) represents the parallel displacement of unit tangent vector $\frac{dx_\mu}{ds}$ along the geodesic.

3.16. COVARIANT DERIVATIVE OF A VECTOR (Agra, 1959, 70)

We have already introduced the transformation law of covariant vector as

$$\frac{\partial \phi}{\partial x_\mu} = \frac{\partial x_\alpha}{\partial x_\mu} \frac{\partial \phi}{\partial x_\alpha}, \text{ (by §3.4)}$$

where ϕ is an invariant function of position.

This relation follows that the derivative of an invariant is a covariant vector, but we should carefully note that the derivative of a vector is not a tensor. We shall now find certain tensors to be employed in place of ordinary derivatives of vectors.

Now consider $\frac{dx_\mu}{ds}$ when dx_μ is contravariant and ds invariant.

It is a contravariant vector and represents velocity. Thus if A_μ is a covariant vector, then the inner product $A_\mu \frac{dx_\mu}{ds}$ is invariant. As such, the rate of change of $A_\mu \frac{dx_\mu}{ds}$ per unit interval along any assigned curve must be independent of the co-ordinate system

i.e. $\frac{d}{ds} \left(A_\mu \frac{dx_\mu}{ds} \right)$ is invariant. $\dots (1)$

This leads to an assumption that we keep to the same absolute curve however the co-ordinate system is varied. This result being of no practical importance is now applied to a geodesic.

The expression (1) on performing differentiation yields

$$\frac{\partial A_\mu}{\partial x_\nu} \frac{dx_\nu}{ds} \frac{dx_\mu}{ds} + A_\mu \frac{d^2 x_\mu}{ds^2} \text{ is invariant along a geodesic} \quad \dots (2)$$

The equation of a geodesic is

$$\frac{d^2 x_\alpha}{ds^2} + \Gamma_{\mu\nu}^\alpha \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} = 0$$

$$\text{i.e. } A_\mu \frac{d^2 x_\mu}{ds^2} + A_\alpha \frac{d^2 x_\alpha}{ds^2} = -A_\alpha \Gamma_{\mu\nu}^\alpha \frac{dx_\mu}{ds} \frac{dx_\nu}{ds}.$$

Its substitution in (2) gives

$$\frac{dx_\mu}{ds} \frac{dx_\nu}{ds} \left(\frac{\partial A_\mu}{\partial x_\nu} - A_\alpha \Gamma_{\mu\nu}^\alpha \right) \text{ is invariant.}$$

Since $\frac{dx_\mu}{ds}$ and $\frac{dx_\nu}{ds}$ both are contravariant vectors, their cofactor

i.e., $\frac{\partial A_\mu}{\partial x_\nu} - \Gamma_{\mu\nu}^\alpha A_\alpha$ is a covariant tensor of rank two. (Agra, 1964)

We write

$$A_{\mu, \nu} \text{ (or simply } A_{\mu\nu}) = \frac{\partial A_\mu}{\partial x_\nu} - \Gamma_{\mu\nu}^\alpha A_\alpha \quad \dots (3)$$

and call the tensor $A_{\mu, \nu}$ the **contravariant derivative** of A_μ w.r.t. x_ν .

The result (3) may be arrived at by another method as follows:

We have

$$\Gamma_{\mu\nu}^\rho \frac{\partial x_\rho}{\partial x'_\mu} = \frac{\partial^2 x_\rho}{\partial x'_\mu \partial x'_\nu} + \frac{\partial x_\alpha}{\partial x'_\mu} \frac{\partial x_\beta}{\partial x'_\nu} \Gamma_{\alpha\beta}^\rho \quad \dots (4)$$

and the transformation law of covariant vectors gives

$$A'_\mu = \frac{\partial x_\alpha}{\partial x'_\mu} A_\alpha$$

which on differentiation with respect to x'_ν yields

$$\frac{\partial A'_\mu}{\partial x'_\nu} = \frac{\partial^2 x_\alpha}{\partial x'_\mu \partial x'_\nu} A_\alpha + \frac{\partial x_\alpha}{\partial x'_\mu} \frac{\partial x_\beta}{\partial x'_\nu} \frac{\partial A_\alpha}{\partial x_\beta}$$

Applying (4) and changing the dummy suffixes in the last term, we get

$$\begin{aligned} \frac{\partial A'_\mu}{\partial x'_\nu} = & \left(\Gamma_{\mu\nu}^\rho \frac{\partial x_\rho}{\partial x'_\mu} - \frac{\partial x_\alpha}{\partial x'_\mu} \frac{\partial x_\beta}{\partial x'_\nu} \Gamma_{\alpha\beta}^\rho \right) A_\alpha \\ & + \frac{\partial x_\alpha}{\partial x'_\mu} \frac{\partial x_\beta}{\partial x'_\nu} \frac{\partial A_\alpha}{\partial x_\beta}. \end{aligned} \quad \dots (5)$$

Again we have from law of transformation of covariant vectors

$$A_\alpha \frac{\partial x_\alpha}{\partial x'_\mu} = A'_\mu$$

so that (5) becomes

$$\frac{\partial A'_\mu}{\partial x'_\nu} - \Gamma_{\mu\nu}^\rho A'_\rho = \frac{\partial x_\alpha}{\partial x'_\mu} \frac{\partial x_\beta}{\partial x'_\nu} \left(\frac{\partial A_\alpha}{\partial x_\beta} - \Gamma_{\alpha\beta}^\rho A_\rho \right) \quad \dots (6)$$

Showing that $\frac{\partial A_\mu}{\partial x_\nu} - \Gamma_{\mu\nu}^\rho A_\rho$ obeys the law of transformation of a covariant tensor. We thus approach the result (3).

From (3), by raising a suffix we get an important associated tensor A^μ_{ν} as follows:

We have

$$A_\alpha = g_{\alpha\mu} A^\mu,$$

so that from (3) we have

$$\begin{aligned} A_{\sigma, \nu} &= \frac{\partial A_{\sigma}}{\partial x_{\nu}} - \Gamma_{\sigma \nu}^{\alpha} A_{\alpha} \\ &= \frac{\partial}{\partial x_{\nu}} (g_{\sigma \epsilon} A^{\epsilon}) - \Gamma_{\sigma \nu}^{\alpha} (g_{\alpha \epsilon} A^{\epsilon}) \\ &= g_{\sigma \epsilon} \frac{\partial A^{\epsilon}}{\partial x_{\nu}} + A^{\epsilon} \frac{\partial g_{\sigma \epsilon}}{\partial x_{\nu}} - \Gamma_{\sigma \nu}^{\alpha} A_{\alpha} \quad \text{by §3.11(6)} \\ &= g_{\sigma \epsilon} \frac{\partial A^{\epsilon}}{\partial x_{\nu}} + \Gamma_{\epsilon \nu, \sigma} A^{\epsilon} \quad \text{by §3.11(7)} \end{aligned}$$

Multiplying throughout by $g^{\mu \sigma}$, this becomes

$$g^{\mu \sigma} A_{\sigma, \nu} = g^{\mu \sigma} \frac{\partial A^{\epsilon}}{\partial x_{\nu}} + g^{\mu \sigma} \Gamma_{\epsilon \nu, \sigma} A^{\epsilon}$$

or

$$\begin{aligned} A^{\mu}{}_{\nu} &= g^{\mu \sigma} \frac{\partial A^{\epsilon}}{\partial x_{\nu}} + \Gamma_{\epsilon \nu}^{\mu} A^{\epsilon} \\ &= \frac{\partial A^{\mu}}{\partial x_{\nu}} + \Gamma_{\epsilon \nu}^{\mu} A^{\epsilon} \end{aligned} \quad \dots (7)$$

(Agra, 1970)

which is called the covariant derivative of A^{μ} .

Similarly the tensors $A^{\mu}{}_{\nu}$ and $A^{\mu \nu}$ obtained from (3) and (7) by raising the suffixes are known as contravariant derivatives of A_{μ} and A^{μ} , but they are of no practical importance.

3.17. COVARIANT DERIVATIVE OF A TENSOR (Agra, 1963)

The covariant derivatives of a tensor of rank two are formed as

$$A_{\mu \nu, \sigma} = \frac{\partial A_{\mu \nu}}{\partial x_{\sigma}} - \Gamma_{\mu \sigma}^{\alpha} A_{\alpha \nu} - \Gamma_{\nu \sigma}^{\alpha} A_{\mu \alpha} \quad \dots (1)$$

$$A^{\nu}{}_{\mu \sigma} \text{ (or } A^{\nu}{}_{\mu, \sigma}) = \frac{\partial A^{\nu}{}_{\mu}}{\partial x_{\sigma}} - \Gamma_{\mu \sigma}^{\alpha} A^{\nu}{}_{\alpha} + \Gamma_{\alpha \sigma}^{\nu} A^{\alpha}{}_{\mu} \quad \dots (2)$$

$$A^{\mu \nu}{}_{, \sigma} \text{ (or } A^{\mu \nu, \sigma}) = \frac{\partial A^{\mu \nu}}{\partial x_{\sigma}} + \Gamma_{\alpha \sigma}^{\mu} A^{\alpha \nu} + \Gamma_{\alpha \sigma}^{\nu} A^{\mu \alpha} \quad \dots (3)$$

and a general rule for giving covariant differentiation with respect to x_{σ} may be illustrated as follows:

$$\begin{aligned} A^{\rho}{}_{\lambda \mu \nu, \sigma} &= \frac{\partial A^{\rho}{}_{\lambda \mu \nu}}{\partial x_{\sigma}} - \Gamma_{\lambda \sigma}^{\alpha} A^{\rho}{}_{\alpha \mu \nu} - \Gamma_{\mu \sigma}^{\alpha} A^{\rho}{}_{\lambda \alpha \nu} - \Gamma_{\nu \sigma}^{\alpha} A^{\rho}{}_{\lambda \mu \alpha} \\ &\quad + \Gamma_{\alpha \sigma}^{\rho} A^{\alpha}{}_{\lambda \mu \nu} \end{aligned} \quad \dots (4)$$

In order to show that the quantities on the right are actually tensors, we proceed as follows:

In place of (1) of § 3.16 we may use to say that

$$\frac{d}{ds} \left(A_{\mu \nu} \frac{dx_{\mu}}{ds} \frac{dx_{\nu}}{ds} \right) \text{ is invariant}$$

$$\text{i.e.,} \quad \frac{dA_{\mu \nu}}{dx_{\sigma}} \frac{dx_{\sigma}}{ds} \frac{dx_{\mu}}{ds} \frac{dx_{\nu}}{ds} + A_{\mu \nu} \frac{dx_{\sigma}}{ds} \frac{d^2 x_{\mu}}{ds^2} + A_{\mu \nu} \frac{dx_{\sigma}}{ds} \frac{d^2 x_{\nu}}{ds^2}$$

is invariant along a geodesic.

But along a geodesic, we have

$$A_{\mu \nu} \frac{d^2 x_{\mu}}{ds^2} = A_{\alpha \nu} \frac{d^2 x_{\alpha}}{ds^2} = -A_{\alpha \nu} \Gamma_{\mu \sigma}^{\alpha} \frac{dx_{\mu}}{ds} \frac{dx_{\sigma}}{ds}$$

and

$$A_{\mu \nu} \frac{d^2 x_{\nu}}{ds^2} = A_{\mu \alpha} \frac{d^2 x_{\alpha}}{ds^2} = -A_{\mu \alpha} \Gamma_{\nu \sigma}^{\alpha} \frac{dx_{\nu}}{ds} \frac{dx_{\sigma}}{ds}$$

$$\therefore \left(\frac{\partial A_{\mu \nu}}{\partial x_{\sigma}} - \Gamma_{\mu \sigma}^{\alpha} A_{\alpha \nu} - \Gamma_{\nu \sigma}^{\alpha} A_{\mu \alpha} \right) \frac{dx_{\mu}}{ds} \frac{dx_{\nu}}{ds} \frac{dx_{\sigma}}{ds} \text{ is invariant,}$$

i.e.,

$$A_{\mu \nu, \sigma} \frac{dx_{\mu}}{ds} \frac{dx_{\nu}}{ds} \frac{dx_{\sigma}}{ds} \text{ is invariant}$$

which shows that $A_{\mu \nu \sigma}$ is a covariant tensor of rank three.

The results (2) and (3) may be obtained by raising the suffixes ν and μ as follows:

Since $A_{\mu \gamma} = g_{\gamma \epsilon} A^{\epsilon}{}_{\mu}$, therefore,

$$\begin{aligned} A_{\mu \gamma, \sigma} &= \frac{\partial A_{\mu \gamma}}{\partial x_{\sigma}} - \Gamma_{\mu \sigma}^{\alpha} A_{\alpha \gamma} - \Gamma_{\gamma \sigma}^{\alpha} A_{\mu \alpha} \\ &= \frac{\partial}{\partial x_{\sigma}} (g_{\gamma \epsilon} A^{\epsilon}{}_{\mu}) - \Gamma_{\mu \sigma}^{\alpha} g_{\alpha \gamma} A^{\epsilon}{}_{\mu} - \Gamma_{\gamma \sigma}^{\alpha} g_{\alpha \epsilon} A^{\epsilon}{}_{\mu} \\ &= g_{\gamma \epsilon} \frac{\partial A^{\epsilon}{}_{\mu}}{\partial x_{\sigma}} + A^{\epsilon}{}_{\mu} \frac{\partial g_{\gamma \epsilon}}{\partial x_{\sigma}} - \Gamma_{\mu \sigma}^{\alpha} g_{\alpha \gamma} A^{\epsilon}{}_{\mu} - \Gamma_{\gamma \sigma}^{\alpha} g_{\alpha \epsilon} A^{\epsilon}{}_{\mu} \\ &= g_{\gamma \epsilon} \frac{\partial A^{\epsilon}{}_{\mu}}{\partial x_{\sigma}} - \Gamma_{\mu \sigma}^{\alpha} g_{\alpha \gamma} A^{\epsilon}{}_{\mu} + \left(\frac{\partial g_{\gamma \epsilon}}{\partial x_{\sigma}} - \Gamma_{\gamma \sigma}^{\alpha} g_{\alpha \epsilon} \right) A^{\epsilon}{}_{\mu} \\ &= g_{\gamma \epsilon} \frac{\partial A^{\epsilon}{}_{\mu}}{\partial x_{\sigma}} - \Gamma_{\mu \sigma}^{\alpha} g_{\alpha \gamma} A^{\epsilon}{}_{\mu} + \Gamma_{\epsilon \sigma, \gamma} A^{\epsilon}{}_{\mu} \text{ by (7) of §3.12. (Agra, 1959)} \end{aligned}$$

Multiplying throughout by $g^{\nu \gamma}$, this becomes

$$g^{\nu \gamma} A_{\mu \gamma, \sigma} = g^{\nu \gamma} g_{\gamma \epsilon} \frac{\partial A^{\epsilon}{}_{\mu}}{\partial x_{\sigma}} - g^{\nu \gamma} \Gamma_{\mu \sigma}^{\alpha} g_{\alpha \gamma} A^{\epsilon}{}_{\mu} + g^{\nu \gamma} \Gamma_{\epsilon \sigma, \gamma} A^{\epsilon}{}_{\mu}$$

$$\text{i.e.,} \quad A^{\nu}{}_{\mu, \sigma} = \frac{\partial A^{\nu}{}_{\mu}}{\partial x_{\sigma}} - \Gamma_{\mu \sigma}^{\alpha} g^{\nu \gamma} A_{\alpha \gamma} + \Gamma_{\epsilon \sigma}^{\nu} A^{\epsilon}{}_{\mu} \text{ by (5) of §3.12.}$$

$$= \frac{\partial A^{\nu}{}_{\mu}}{\partial x_{\sigma}} - \Gamma_{\mu \sigma}^{\alpha} A^{\nu}{}_{\alpha} + \Gamma_{\epsilon \sigma}^{\nu} A^{\epsilon}{}_{\mu}$$

which is the result (2).

Again, since $A^{\nu}{}_{\gamma} = g_{\gamma \epsilon} A^{\epsilon \nu}$, we have

$$A^{\nu}{}_{\gamma, \sigma} = \frac{\partial A^{\nu}{}_{\gamma}}{\partial x_{\sigma}} - \Gamma_{\gamma \sigma}^{\alpha} A^{\nu}{}_{\alpha} + \Gamma_{\alpha \sigma}^{\nu} A^{\alpha}{}_{\gamma}$$

$$\begin{aligned}
&= \frac{\partial}{\partial x_\sigma} (g_{\gamma\epsilon} A^{\epsilon\nu}) - \Gamma_{\gamma\sigma}^\alpha g_{\alpha\epsilon} A^{\epsilon\nu} + \Gamma_{\alpha\sigma}^\nu A_\gamma^\alpha \\
&= g_{\gamma\epsilon} \frac{\partial A^{\epsilon\nu}}{\partial x_\sigma} + A^{\epsilon\nu} \frac{\partial g_{\gamma\epsilon}}{\partial x_\sigma} - \Gamma_{\gamma\sigma,\epsilon} A^{\epsilon\nu} + \Gamma_{\alpha\sigma}^\nu A_\gamma^\alpha \\
&\quad \text{by (6) of §3.12.}
\end{aligned}$$

$$\begin{aligned}
\text{or } A_{\gamma,\sigma}^\nu &= g_{\gamma\epsilon} \frac{\partial A^{\epsilon\nu}}{\partial x_\sigma} + A^{\epsilon\nu} \left(\frac{\partial g_{\gamma\epsilon}}{\partial x_\sigma} - \Gamma_{\gamma\sigma,\epsilon} \right) + \Gamma_{\alpha\sigma}^\nu A_\gamma^\alpha \\
&= g_{\gamma\epsilon} \frac{\partial A^{\epsilon\nu}}{\partial x_\sigma} + A^{\epsilon\nu} \Gamma_{\epsilon\sigma,\gamma} + \Gamma_{\alpha\sigma}^\nu A_\gamma^\alpha \quad \text{by (7) of §3.12.}
\end{aligned}$$

Multiplying throughout by $g^{\mu\nu}$, this becomes

$$g^{\mu\gamma} A_{\gamma,\sigma}^\nu = g^{\mu\gamma} g_{\gamma\epsilon} \frac{\partial A^{\epsilon\nu}}{\partial x_\sigma} + g^{\mu\gamma} A^{\epsilon\nu} \Gamma_{\epsilon\sigma,\gamma} + g^{\mu\gamma} \Gamma_{\alpha\sigma}^\nu A_\gamma^\alpha$$

$$\begin{aligned}
\text{i.e., } A_{\sigma}^{\mu\nu} &= \frac{\partial A^{\mu\nu}}{\partial x_\sigma} + \Gamma_{\beta\sigma}^\mu A^{\beta\nu} + \Gamma_{\alpha\sigma}^\nu A^{\mu\alpha} \\
&= \frac{\partial A^{\mu\nu}}{\partial x_\sigma} + \Gamma_{\beta\sigma}^\mu A^{\beta\nu} + \Gamma_{\alpha\sigma}^\nu A^{\mu\alpha}
\end{aligned}$$

which is the result (3).

Problem 27. Prove that $g_{\alpha\beta,\gamma} = 0$ i.e., covariant derivative of fundamental tensor vanishes identically. (Agra, 1963, 65, 68)

We have by §3.17(1).

$$g_{\alpha\beta,\gamma} = \frac{\partial g_{\alpha\beta}}{\partial x_\gamma} - \Gamma_{\alpha\gamma}^\mu g_{\mu\beta} - \Gamma_{\beta\gamma}^\mu g_{\mu\alpha} \quad \dots (1)$$

$$\text{But } \Gamma_{\alpha\gamma}^\mu = \frac{1}{2} g^{\mu\beta} \left(\frac{\partial g_{\alpha\beta}}{\partial x_\gamma} + \frac{\partial g_{\beta\gamma}}{\partial x_\alpha} - \frac{\partial g_{\alpha\gamma}}{\partial x_\beta} \right)$$

$$\begin{aligned}
\text{so that } g_{\mu\beta} \Gamma_{\alpha\gamma}^\mu &= \frac{1}{2} g_{\mu\beta} g^{\mu\beta} \left(\frac{\partial g_{\alpha\beta}}{\partial x_\gamma} + \frac{\partial g_{\beta\gamma}}{\partial x_\alpha} - \frac{\partial g_{\alpha\gamma}}{\partial x_\beta} \right) \\
&= \frac{1}{2} \left(\frac{\partial g_{\alpha\beta}}{\partial x_\gamma} + \frac{\partial g_{\beta\gamma}}{\partial x_\alpha} - \frac{\partial g_{\alpha\gamma}}{\partial x_\beta} \right) \text{ since } g_{\mu\beta} g^{\mu\beta} = 1
\end{aligned}$$

$$\text{Similarly } g_{\mu\alpha} \Gamma_{\beta\gamma}^\mu = \frac{1}{2} \left(\frac{\partial g_{\beta\alpha}}{\partial x_\gamma} + \frac{\partial g_{\alpha\gamma}}{\partial x_\beta} - \frac{\partial g_{\beta\gamma}}{\partial x_\alpha} \right).$$

Adding the last two relations, we get

$$g_{\mu\beta} \Gamma_{\alpha\gamma}^\mu + g_{\mu\alpha} \Gamma_{\beta\gamma}^\mu = \frac{\partial g_{\alpha\beta}}{\partial x_\gamma}$$

Substituting this value in (1), we get

$$\begin{aligned}
g_{\alpha\beta,\gamma} &= g_{\mu\beta} \Gamma_{\alpha\gamma}^\mu + g_{\mu\alpha} \Gamma_{\beta\gamma}^\mu - \Gamma_{\alpha\gamma}^\mu g_{\mu\beta} - \Gamma_{\beta\gamma}^\mu g_{\mu\alpha} \\
&= 0.
\end{aligned}$$

Problem 28. By starting with $A'_{\alpha\beta} = A_{\mu\nu} \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta}$, verify the first result of §3.17.

Given that

$$A'_{\alpha\beta} = A_{\mu\nu} \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} \quad \dots (1)$$

Differentiating with regard to x'_γ , we have

$$\begin{aligned}
\frac{\partial A'_{\alpha\beta}}{\partial x'_\gamma} &= \frac{\partial A_{\mu\nu}}{\partial x_\sigma} \frac{\partial x_\sigma}{\partial x'_\gamma} \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} + A_{\mu\nu} \frac{\partial^2 x_\mu}{\partial x'_\alpha \partial x'_\gamma} \frac{\partial x_\nu}{\partial x'_\beta} + A_{\mu\nu} \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial^2 x_\nu}{\partial x'_\beta \partial x'_\gamma} \\
&= \frac{\partial A_{\mu\nu}}{\partial x_\sigma} \frac{\partial x_\sigma}{\partial x'_\gamma} \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} + A_{\mu\nu} \left[\Gamma_{\alpha\gamma}^\epsilon \frac{\partial x_\mu}{\partial x'_\epsilon} - \Gamma_{\epsilon\gamma}^\mu \frac{\partial x_\epsilon}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} \right] \frac{\partial x_\nu}{\partial x'_\beta} \\
&\quad + A_{\mu\nu} \left[\Gamma_{\beta\gamma}^\epsilon \frac{\partial x_\nu}{\partial x'_\epsilon} - \Gamma_{\epsilon\gamma}^\nu \frac{\partial x_\epsilon}{\partial x'_\beta} \frac{\partial x_\mu}{\partial x'_\alpha} \right] \frac{\partial x_\mu}{\partial x'_\alpha}, \text{ by (12) §3.14.}
\end{aligned}$$

$$\begin{aligned}
\text{or } \frac{\partial A'_{\alpha\beta}}{\partial x'_\gamma} &= A_{\mu\nu} \Gamma_{\alpha\gamma}^\epsilon \frac{\partial x_\mu}{\partial x'_\epsilon} \frac{\partial x_\nu}{\partial x'_\beta} - A_{\mu\nu} \Gamma_{\beta\gamma}^\epsilon \frac{\partial x_\nu}{\partial x'_\epsilon} \frac{\partial x_\mu}{\partial x'_\alpha} \\
&= \frac{\partial A_{\mu\nu}}{\partial x_\sigma} \frac{\partial x_\sigma}{\partial x'_\gamma} \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} - A_{\mu\nu} \Gamma_{\epsilon\gamma}^\mu \frac{\partial x_\epsilon}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} \\
&\quad - A_{\mu\nu} \Gamma_{\epsilon\gamma}^\nu \frac{\partial x_\epsilon}{\partial x'_\beta} \frac{\partial x_\mu}{\partial x'_\alpha}
\end{aligned}$$

or using (1), this becomes

$$\frac{\partial A'_{\alpha\beta}}{\partial x'_\gamma} = A'_{\epsilon\nu} \Gamma_{\alpha\gamma}^\epsilon - A'_{\mu\tau} \Gamma_{\beta\gamma}^\mu \left(\frac{\partial A_{\mu\nu}}{\partial x_\sigma} - \Gamma_{\mu\sigma}^\tau A_{\tau\nu} - \Gamma_{\nu\sigma}^\tau A_{\mu\tau} \right) \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} \frac{\partial x_\sigma}{\partial x'_\gamma}$$

$$\text{i.e. } A'_{\alpha\beta,\gamma} = A_{\mu\nu,\sigma} \frac{\partial x_\mu}{\partial x'_\alpha} \frac{\partial x_\nu}{\partial x'_\beta} \frac{\partial x_\sigma}{\partial x'_\gamma}$$

showing that $A_{\mu\nu,\sigma}$ is a tensor of rank three and is the covariant derivative of $A_{\mu\nu}$ with respect to x_σ .

This verifies the first result of §3.17.

Problem 29. Show that the distributive law of ordinary differentiation holds good in case of covariant differentiation of a product, i.e. to show that (Agra, 1959, 63)

$$(i) (B_\mu C_\nu)_{,\sigma} = B_{\mu,\sigma} C_\nu + B_\mu C_{\nu,\sigma},$$

$$(ii) (B^\mu C^\nu)_{,\sigma} = B^{\mu}_{,\sigma} C^\nu + B^\mu C^{\nu}_{,\sigma}.$$

$$\begin{aligned}
(i) \quad \text{R.H.S.} &= B_{\mu,\sigma} C_\nu + B_\mu C_{\nu,\sigma} \\
&= \left(\frac{\partial B_\mu}{\partial x_\sigma} - \Gamma_{\mu\sigma}^\alpha B_\alpha \right) C_\nu + B_\mu \left(\frac{\partial C_\nu}{\partial x_\sigma} - \Gamma_{\nu\sigma}^\alpha C_\alpha \right) \\
&= \frac{\partial}{\partial x_\sigma} (B_\mu C_\nu) - \Gamma_{\mu\sigma}^\alpha (B_\alpha C_\nu) - \Gamma_{\nu\sigma}^\alpha (B_\mu C_\alpha) \\
&= (B_\mu C_\nu)_{,\sigma}.
\end{aligned}$$

$$\begin{aligned}
(ii) \quad \text{R.H.S.} &= B^{\mu}_{,\sigma} C^\nu + B^\mu C^{\nu}_{,\sigma} \\
&= \left(\frac{\partial B^\mu}{\partial x_\sigma} + \Gamma_{\alpha\sigma}^\mu B^\alpha \right) C^\nu + B^\mu \left(\frac{\partial C^\nu}{\partial x_\sigma} + \Gamma_{\alpha\sigma}^\nu C^\alpha \right) \\
&= \frac{\partial}{\partial x_\sigma} (B^\mu C^\nu) + \Gamma_{\alpha\sigma}^\mu (B^\alpha C^\nu) + \Gamma_{\alpha\sigma}^\nu (B^\mu C^\alpha) \\
&= (B^\mu C^\nu)_{,\sigma}.
\end{aligned}$$

Problem 30. Prove that

$$(a) \quad g^{\mu\nu}_{;\sigma} = 0. \quad (\text{Agra, 1959, 65})$$

$$(b) \quad \delta^{\nu}_{\mu;\sigma} = 0. \quad (\text{Agra, 1965})$$

(a) We have

$$g^{\xi\eta} g^{\xi\xi} = \delta^{\xi}_{\eta} \\ = 0 \text{ or } 1.$$

Differentiating with respect to x_{λ} , this gives

$$g^{\xi\xi} \frac{\partial g^{\xi\eta}}{\partial x_{\lambda}} + g^{\xi\eta} \frac{\partial g^{\xi\xi}}{\partial x_{\lambda}} = 0$$

Multiplying throughout by $g^{\eta\rho}$, we get

$$g^{\eta\rho} g^{\xi\xi} \frac{\partial g^{\xi\eta}}{\partial x_{\lambda}} + g^{\eta\rho} g^{\xi\eta} \frac{\partial g^{\xi\xi}}{\partial x_{\lambda}} = 0$$

$$\text{or} \quad g^{\eta\rho} g^{\xi\xi} (\Gamma^{\xi}_{\lambda\eta} - \Gamma^{\eta}_{\lambda\xi}) + g^{\xi\rho} \frac{\partial g^{\xi\xi}}{\partial x_{\lambda}} = 0 \text{ by (7) of §3.11}$$

$$\text{or} \quad g^{\xi\xi} \Gamma^{\rho}_{\xi\lambda} + g^{\eta\rho} \Gamma^{\xi}_{\lambda\eta} + \frac{\partial g^{\xi\rho}}{\partial x_{\lambda}} = 0 \text{ by (5) of §3.11} \quad \dots (1)$$

$$\text{Now } g^{\mu\nu}_{;\sigma} = \frac{\partial g^{\mu\nu}}{\partial x_{\sigma}} + \Gamma^{\mu}_{\alpha\sigma} g^{\alpha\nu} + \Gamma^{\nu}_{\alpha\sigma} g^{\alpha\mu} \\ = 0 \text{ by (1).}$$

$$(b) \quad \delta^{\nu}_{\mu;\sigma} = \frac{\partial \delta^{\nu}_{\mu}}{\partial x_{\sigma}} - \Gamma^{\alpha}_{\mu\sigma} g^{\nu}_{\alpha} + \Gamma^{\nu}_{\alpha\sigma} \delta^{\alpha}_{\mu} \\ = 0 - \Gamma^{\nu}_{\mu\sigma} + \Gamma^{\nu}_{\mu\sigma} \\ = 0.$$

Problem 31. Show that the covariant derivative of an invariant is the same as its ordinary derivative.

Let I be an invariant, so that IA_{μ} is a covariant vector.

Now the covariant derivative of IA_{μ} is

$$(IA_{\mu})_{;\nu} = \frac{\partial (IA_{\mu})}{\partial x_{\nu}} - \Gamma^{\alpha}_{\mu\nu} (IA_{\alpha}) \\ = A_{\mu} \frac{\partial I}{\partial x_{\nu}} - I \frac{\partial A_{\mu}}{\partial x_{\nu}} - \Gamma^{\alpha}_{\mu\nu} IA_{\alpha} \\ = A_{\mu} \frac{\partial I}{\partial x_{\nu}} + I \left(\frac{\partial A_{\mu}}{\partial x_{\nu}} - \Gamma^{\alpha}_{\mu\nu} A_{\alpha} \right) \\ = A_{\mu} \frac{\partial I}{\partial x_{\nu}} + IA_{\mu;\nu}.$$

But from Problem 29.

$$(IA_{\mu})_{;\nu} = I_{;\nu} A_{\mu} + IA_{\mu;\nu}$$

$$\therefore \quad I_{;\nu} A_{\mu} + IA_{\mu;\nu} = A_{\mu} \frac{\partial I}{\partial x_{\nu}} + IA_{\mu;\nu}$$

$$\text{i.e.} \quad I_{;\nu} A_{\mu} = A_{\mu} \frac{\partial I}{\partial x_{\nu}}$$

$$\text{or} \quad I_{;\nu} = \frac{\partial I}{\partial x_{\nu}}$$

which shows that the covariant derivative of an invariant is the same as its ordinary derivative.

3.18. THE CURVATURE TENSOR (RIEMANN CHRISTOFFEL TENSOR)

Consider an absolute contravariant vector A^{ϵ} . Its covariant derivative gives the mixed tensor.

$$A^{\epsilon}_{;\rho} = \frac{\partial A^{\epsilon}}{\partial x_{\rho}} + A^{\mu} \Gamma^{\epsilon}_{\mu\rho} \text{ by (7) of §3.16.}$$

If this is again differentiated covariantly, then we have

$$A^{\epsilon}_{;\rho\sigma} = \frac{\partial A^{\epsilon}_{;\rho}}{\partial x_{\sigma}} + A^{\mu}_{;\rho} \Gamma^{\epsilon}_{\mu\sigma} - A^{\epsilon}_{\mu} \Gamma^{\mu}_{\rho\sigma} \text{ by (2) of §3.17} \\ = \frac{\partial}{\partial x_{\sigma}} \left(\frac{\partial A^{\epsilon}}{\partial x_{\rho}} + A^{\mu} \Gamma^{\epsilon}_{\mu\rho} \right) + \left(\frac{\partial A^{\mu}}{\partial x_{\rho}} + A^{\beta} \Gamma^{\mu}_{\beta\rho} \right) \Gamma^{\epsilon}_{\mu\sigma} \\ - \left(\frac{\partial A^{\epsilon}}{\partial x_{\mu}} + A^{\beta} \Gamma^{\epsilon}_{\beta\mu} \right) \Gamma^{\mu}_{\rho\sigma} \\ = \frac{\partial^2 A^{\epsilon}}{\partial x_{\sigma} \partial x_{\rho}} + \frac{\partial A^{\mu}}{\partial x_{\sigma}} \Gamma^{\epsilon}_{\mu\rho} + A^{\mu} \frac{\partial \Gamma^{\epsilon}_{\mu\rho}}{\partial x_{\sigma}} + \left(\frac{\partial A^{\mu}}{\partial x_{\rho}} + A^{\beta} \Gamma^{\mu}_{\beta\rho} \right) \Gamma^{\epsilon}_{\mu\sigma} \\ - \left(\frac{\partial A^{\epsilon}}{\partial x_{\mu}} + A^{\beta} \Gamma^{\epsilon}_{\beta\mu} \right) \Gamma^{\mu}_{\rho\sigma} \quad \dots (1)$$

Similarly,

$$A^{\epsilon}_{;\sigma\rho} = \frac{\partial^2 A^{\epsilon}}{\partial x_{\rho} \partial x_{\sigma}} + \frac{\partial A^{\mu}}{\partial x_{\rho}} \Gamma^{\epsilon}_{\mu\sigma} + A^{\mu} \frac{\partial \Gamma^{\epsilon}_{\mu\sigma}}{\partial x_{\rho}} + \left(\frac{\partial A^{\mu}}{\partial x_{\sigma}} + A^{\beta} \Gamma^{\mu}_{\beta\sigma} \right) \Gamma^{\epsilon}_{\mu\rho} \\ - \left(\frac{\partial A^{\epsilon}}{\partial x_{\mu}} + A^{\beta} \Gamma^{\epsilon}_{\beta\mu} \right) \Gamma^{\mu}_{\sigma\rho} \quad \dots (2)$$

Subtracting (2) from (1), after changing the dummy suffixes wherever needed, we get

$$A^{\epsilon}_{;\rho\sigma} - A^{\epsilon}_{;\sigma\rho} = A^{\mu} \left(\frac{\partial \Gamma^{\epsilon}_{\mu\rho}}{\partial x_{\sigma}} - \frac{\partial \Gamma^{\epsilon}_{\mu\sigma}}{\partial x_{\rho}} + \Gamma^{\beta}_{\mu\rho} \Gamma^{\epsilon}_{\beta\sigma} - \Gamma^{\beta}_{\mu\sigma} \Gamma^{\epsilon}_{\beta\rho} \right).$$

(Agra, 1958)

The quotient rule follows that the cofactor of A^{μ} must be a tensor of rank four, so that we may write

$$A^{\epsilon}_{;\rho\sigma} - A^{\epsilon}_{;\sigma\rho} = A^{\mu} B^{\epsilon}_{\mu\rho\sigma}$$

where

$$B_{\mu\rho\sigma}^{\epsilon} = \frac{\partial \Gamma_{\mu\rho}^{\epsilon}}{\partial x_{\sigma}} - \frac{\partial \Gamma_{\mu\sigma}^{\epsilon}}{\partial x_{\rho}} + \Gamma_{\mu\rho}^{\epsilon} \Gamma_{\beta\sigma}^{\epsilon} - \Gamma_{\mu\sigma}^{\epsilon} \Gamma_{\beta\rho}^{\epsilon}.$$

This tensor is known as **curvature tensor**. On contraction this tensor yields two new tensors as follows:

$$(i) \quad B_{\mu\sigma\nu}^{\sigma} = \frac{\partial \Gamma_{\mu\sigma}^{\sigma}}{\partial x_{\nu}} - \frac{\partial \Gamma_{\mu\nu}^{\sigma}}{\partial x_{\sigma}} + \Gamma_{\mu\sigma}^{\beta} \Gamma_{\beta\nu}^{\sigma} - \Gamma_{\mu\nu}^{\beta} \Gamma_{\beta\sigma}^{\sigma} \\ = R_{\mu\nu} \text{ (say).} \quad \dots (4)$$

This tensor is known as **Ricci tensor**.

$$(ii) \quad B_{\sigma\mu\nu}^{\sigma} = \frac{\partial \Gamma_{\sigma\mu}^{\sigma}}{\partial x_{\nu}} - \frac{\partial \Gamma_{\sigma\nu}^{\sigma}}{\partial x_{\mu}} + \Gamma_{\sigma\mu}^{\beta} \Gamma_{\beta\nu}^{\sigma} - \Gamma_{\sigma\nu}^{\beta} \Gamma_{\beta\mu}^{\sigma} \\ = \frac{\partial \Gamma_{\sigma\mu}^{\sigma}}{\partial x_{\nu}} - \frac{\partial \Gamma_{\sigma\nu}^{\sigma}}{\partial x_{\mu}} \\ \text{(the last two terms cancel when } \mu \text{ and } \nu \text{ are interchanged in either of them)} \\ = S_{\mu\nu} \text{ (say).} \quad \dots (5)$$

$$\text{So } S_{\nu\mu} = B_{\sigma\nu\mu}^{\sigma} = \frac{\partial \Gamma_{\sigma\nu}^{\sigma}}{\partial x_{\mu}} - \frac{\partial \Gamma_{\sigma\mu}^{\sigma}}{\partial x_{\nu}} \\ = - \left(\frac{\partial \Gamma_{\sigma\mu}^{\sigma}}{\partial x_{\nu}} - \frac{\partial \Gamma_{\sigma\nu}^{\sigma}}{\partial x_{\mu}} \right) \\ = -S_{\mu\nu} \quad \dots (6)$$

This shows that the tensor $S_{\mu\nu}$ is anti-symmetric.

Again, we have from (4),

$$R_{\mu\nu} = \frac{\partial \Gamma_{\mu\sigma}^{\sigma}}{\partial x_{\nu}} - \frac{\partial \Gamma_{\mu\nu}^{\sigma}}{\partial x_{\sigma}} + \Gamma_{\mu\sigma}^{\beta} \Gamma_{\beta\nu}^{\sigma} - \Gamma_{\mu\nu}^{\beta} \Gamma_{\beta\sigma}^{\sigma} \\ = \frac{\partial \Gamma_{\mu\sigma}^{\sigma}}{\partial x_{\nu}} - \frac{\partial \Gamma_{\sigma\nu}^{\sigma}}{\partial x_{\mu}} + \Gamma_{\mu\sigma}^{\beta} \Gamma_{\beta\nu}^{\sigma} - \Gamma_{\mu\nu}^{\beta} \Gamma_{\beta\sigma}^{\sigma} \\ \text{(interchanging } \sigma \text{ and } \mu \text{ in second term)} \\ = S_{\mu\nu} + \Gamma_{\mu\sigma}^{\beta} \Gamma_{\beta\nu}^{\sigma} - \Gamma_{\mu\nu}^{\beta} \Gamma_{\beta\sigma}^{\sigma}.$$

Similarly,

$$R_{\nu\mu} = -S_{\nu\mu} + \Gamma_{\nu\sigma}^{\beta} \Gamma_{\beta\mu}^{\sigma} - \Gamma_{\nu\mu}^{\beta} \Gamma_{\beta\sigma}^{\sigma} \\ = -S_{\nu\mu} + \Gamma_{\mu\sigma}^{\beta} \Gamma_{\beta\nu}^{\sigma} - \Gamma_{\mu\nu}^{\beta} \Gamma_{\beta\sigma}^{\sigma} \\ \text{(interchanging } \mu \text{ and } \nu \text{ in second term)}$$

$$\therefore R_{\mu\nu} - R_{\nu\mu} = S_{\mu\nu} + S_{\nu\mu} \\ = S_{\mu\nu} - S_{\mu\nu} \text{ from (6)} \\ = 0$$

$$\text{i.e. } R_{\mu\nu} = R_{\nu\mu}.$$

... (7)

This shows that $R_{\mu\nu}$ is symmetric.

3.19. RIEMANN-CHRISTOFFEL TENSOR OR COVARIANT CURVATURE TENSOR

The tensor $B_{\mu\nu\sigma\rho} = g_{\rho\epsilon} B_{\mu\nu\sigma}^{\epsilon}$ is known as *Riemann Christoffel* or *covariant curvature tensor*. (Agra, 1963, 64, 65)

We have

$$B_{\mu\nu\sigma\rho} = g_{\rho\epsilon} B_{\mu\nu\sigma}^{\epsilon} \\ = g_{\rho\epsilon} \left(\Gamma_{\mu\sigma}^{\alpha} \Gamma_{\alpha\nu}^{\epsilon} - \Gamma_{\mu\nu}^{\alpha} \Gamma_{\alpha\sigma}^{\epsilon} + \frac{\partial \Gamma_{\mu\sigma}^{\epsilon}}{\partial x_{\nu}} - \frac{\partial \Gamma_{\mu\nu}^{\epsilon}}{\partial x_{\sigma}} \right) \quad \text{from (3) of § 3.18} \\ = \Gamma_{\mu\sigma}^{\alpha} (g_{\rho\epsilon} \Gamma_{\alpha\nu}^{\epsilon}) - \Gamma_{\mu\nu}^{\alpha} (g_{\rho\epsilon} \Gamma_{\alpha\sigma}^{\epsilon}) + \frac{\partial}{\partial x_{\nu}} (g_{\rho\epsilon} \Gamma_{\mu\sigma}^{\epsilon}) \\ - \Gamma_{\mu\sigma}^{\epsilon} \frac{\partial g_{\rho\epsilon}}{\partial x_{\nu}} - \frac{\partial}{\partial x_{\sigma}} (g_{\rho\epsilon} \Gamma_{\mu\nu}^{\epsilon}) + \Gamma_{\mu\nu}^{\epsilon} \frac{\partial g_{\rho\epsilon}}{\partial x_{\sigma}} \\ = \Gamma_{\mu\sigma}^{\alpha} \Gamma_{\alpha\nu, \rho} - \Gamma_{\mu\nu}^{\alpha} \Gamma_{\alpha\sigma, \rho} + \frac{\partial \Gamma_{\mu\sigma, \rho}}{\partial x_{\nu}} - \Gamma_{\mu\nu}^{\epsilon} \frac{\partial g_{\rho\epsilon}}{\partial x_{\sigma}} \\ - \frac{\partial \Gamma_{\mu\alpha, \rho}}{\partial x_{\sigma}} + \Gamma_{\mu\nu}^{\epsilon} \frac{\partial g_{\rho\epsilon}}{\partial x_{\sigma}} \\ = \Gamma_{\mu\sigma}^{\alpha} \Gamma_{\alpha\nu, \rho} - \Gamma_{\mu\nu}^{\alpha} \Gamma_{\alpha\sigma, \rho} + \frac{\partial \Gamma_{\mu\sigma, \rho}}{\partial x_{\nu}} - \frac{\partial \Gamma_{\mu\nu, \rho}}{\partial x_{\sigma}} \\ - \Gamma_{\mu\sigma}^{\alpha} \frac{\partial g_{\rho\alpha}}{\partial x_{\nu}} + \Gamma_{\mu\nu}^{\alpha} \frac{\partial g_{\rho\alpha}}{\partial x_{\sigma}} \\ \text{(replacing } \epsilon \text{ by } \alpha \text{ in the last two terms)} \\ = \Gamma_{\mu\sigma}^{\alpha} \left(\Gamma_{\alpha\nu, \rho} - \frac{\partial g_{\rho\alpha}}{\partial x_{\nu}} \right) - \Gamma_{\mu\nu}^{\alpha} \left(\Gamma_{\alpha\sigma, \rho} - \frac{\partial g_{\rho\alpha}}{\partial x_{\sigma}} \right) \\ + \frac{\partial}{\partial x_{\nu}} \cdot \frac{1}{2} \left(\frac{\partial g_{\mu\rho}}{\partial x_{\sigma}} + \frac{\partial g_{\nu\rho}}{\partial x_{\mu}} - \frac{\partial g_{\mu\sigma}}{\partial x_{\rho}} \right) \\ - \frac{\partial}{\partial x_{\nu}} \cdot \frac{1}{2} \left(\frac{\partial g_{\mu\rho}}{\partial x_{\nu}} + \frac{\partial g_{\nu\rho}}{\partial x_{\mu}} - \frac{\partial g_{\mu\nu}}{\partial x_{\rho}} \right) \\ = -\Gamma_{\mu\sigma}^{\alpha} \Gamma_{\rho\nu, \alpha} + \Gamma_{\mu\nu}^{\alpha} \Gamma_{\rho\sigma, \alpha} + \frac{1}{2} \left(\frac{\partial^2 g_{\rho\sigma}}{\partial x_{\mu} \partial x_{\beta}} - \frac{\partial^2 g_{\mu\sigma}}{\partial x_{\nu} \partial x_{\rho}} \right. \\ \left. - \frac{\partial^2 g_{\nu\rho}}{\partial x_{\mu} \partial x_{\sigma}} + \frac{\partial^2 g_{\mu\nu}}{\partial x_{\sigma} \partial x_{\rho}} \right) \\ \text{(since } \Gamma_{\alpha\nu, \rho} = -\Gamma_{\rho\nu, \alpha} \text{ etc.)} \\ = -\Gamma_{\mu\sigma}^{\alpha} \Gamma_{\rho\nu, \alpha} + \Gamma_{\mu\nu}^{\alpha} \Gamma_{\rho\sigma, \alpha} + \frac{1}{2} \left(\frac{\partial^2 g_{\rho\sigma}}{\partial x_{\mu} \partial x_{\nu}} + \frac{\partial^2 g_{\mu\nu}}{\partial x_{\rho} \partial x_{\sigma}} \right. \\ \left. - \frac{\partial^2 g_{\rho\sigma}}{\partial x_{\rho} \partial x_{\nu}} - \frac{\partial^2 g_{\mu\nu}}{\partial x_{\mu} \partial x_{\sigma}} \right) \quad \dots (1)$$

Interchanging μ and ρ , this gives

$$B_{\rho\nu\sigma\mu} = -\Gamma_{\rho\sigma}^{\alpha} \Gamma_{\mu\nu, \alpha} + \Gamma_{\rho\nu}^{\alpha} \Gamma_{\mu\sigma, \alpha} + \frac{1}{2} \left(\frac{\partial^2 g_{\mu\sigma}}{\partial x_{\rho} \partial x_{\nu}} - \frac{\partial^2 g_{\rho\nu}}{\partial x_{\mu} \partial x_{\sigma}} - \frac{\partial^2 g_{\rho\sigma}}{\partial x_{\mu} \partial x_{\nu}} - \frac{\partial^2 g_{\mu\nu}}{\partial x_{\rho} \partial x_{\sigma}} \right)$$

so that

$$\begin{aligned} B_{\mu\nu\sigma\rho} + B_{\rho\nu\sigma\mu} &= -\Gamma_{\mu\sigma}^{\alpha} \Gamma_{\rho\nu, \alpha} + \Gamma_{\mu\nu}^{\alpha} \Gamma_{\rho\sigma, \alpha} - \Gamma_{\rho\sigma}^{\alpha} \Gamma_{\mu\nu, \alpha} + \Gamma_{\rho\nu}^{\alpha} \Gamma_{\mu\sigma, \alpha} \\ &= -g^{\alpha\lambda} \Gamma_{\mu\sigma, \lambda} g_{\alpha\lambda} \Gamma_{\rho\nu}^{\lambda} + g^{\alpha\lambda} \Gamma_{\mu\nu, \lambda} g_{\alpha\lambda} \Gamma_{\rho\sigma}^{\lambda} \\ &\quad - \Gamma_{\rho\sigma}^{\lambda} \Gamma_{\mu\nu, \lambda} + \Gamma_{\rho\nu}^{\lambda} \Gamma_{\mu\sigma, \lambda} \\ &= -\Gamma_{\mu\sigma, \lambda} \Gamma_{\rho\nu}^{\lambda} + \Gamma_{\mu\nu, \lambda} \Gamma_{\rho\sigma}^{\lambda} - \Gamma_{\rho\sigma}^{\lambda} \Gamma_{\mu\nu, \lambda} + \Gamma_{\rho\nu}^{\lambda} \Gamma_{\mu\sigma, \lambda} \\ &= 0, \end{aligned}$$

$$\text{i.e. } B_{\mu\nu\sigma\rho} = -B_{\rho\nu\sigma\mu} \quad \dots (2)$$

This shows that $B_{\mu\nu\sigma\rho}$ is anti-symmetric in indices μ and ρ .

$$\text{Similarly, } B_{\mu\nu\sigma\rho} = -B_{\mu\sigma\nu\rho} \quad \dots (3)$$

i.e. $B_{\mu\nu\sigma\rho}$ is anti-symmetric in indices ν and σ .

Also, it is easy to show that

$$B_{\mu\nu\sigma\rho} = B_{\rho\sigma\nu\mu} \quad \dots (4)$$

i.e. $B_{\mu\nu\sigma\rho}$ is symmetric in index pair (μ, ν) and (ρ, σ) .

because

$$\begin{aligned} B_{\mu\nu\sigma\rho} &= -B_{\rho\nu\sigma\mu} \text{ by (2)} \\ &= -(-B_{\rho\sigma\nu\mu}) \text{ by (3)} \\ &= B_{\rho\sigma\nu\mu} \end{aligned}$$

Thus $B_{\mu\nu\sigma\rho}$ is skew-symmetric in the first indices, skew-symmetric in the last two indices and symmetric in two pairs of indices.

(Agra, 1969)

Further, it has the cyclic property i.e.

$$B_{\mu\nu\sigma\rho} + B_{\mu\sigma\rho\nu} + B_{\mu\rho\nu\sigma} = 0. \quad \dots (5)$$

(Agra, 1968)

Writing the values of all three tensors by (1), and adding them altogether, the result follows.

We have already mentioned that a general tensor of rank four has $(4)^4$ i.e. 256 components. In the existing case the double anti-symmetry reduces the number to ${}^4C_2 \times {}^4C_2$ i.e. 36 of which 30 are paired because μ, ρ can be interchanged with ν, σ , but the remaining 6 having μ, ρ as the same pair of numbers as ν, σ are without partners. As such we have 21 different components (30 paired components are equivalent to 15 different components + 6) with one further relation given by (5). Conclusively the Riemann-Christoffel tensor has 20 independent components. The scheme may be shown as:

Writing the suffixes in the order $\mu\rho\sigma\nu$, the different 21 components are

$$\begin{aligned} &B_{1212} \ B_{1223} \ B_{1313} \ B_{1324} \ B_{1423} \ B_{2323} \ B_{2424} \\ &B_{1213} \ B_{1224} \ B_{1314} \ B_{1334} \ B_{1424} \ B_{2324} \ B_{2434} \\ &B_{1214} \ B_{1234} \ B_{1323} \ B_{1414} \ B_{1434} \ B_{2334} \ B_{2434} \end{aligned}$$

With the relation $B_{1234} + B_{1342} + B_{1423} = 0$, which reduces the number by 1.

Independent number = $21 - 1 = 20$.

(Agra, 1963, 65, 76)

Problem 32. Prove that if $B_{\mu\nu\sigma}^{\epsilon} = 0$, then space is Euclidean.

We have

$$B_{\mu\nu\sigma}^{\epsilon} = \frac{\partial \Gamma_{\mu\nu}^{\epsilon}}{\partial x_{\sigma}} - \frac{\partial \Gamma_{\mu\sigma}^{\epsilon}}{\partial x_{\nu}} + \Gamma_{\mu\nu}^{\alpha} \Gamma_{\alpha\sigma}^{\epsilon} - \Gamma_{\mu\sigma}^{\alpha} \Gamma_{\alpha\nu}^{\epsilon}$$

But in Euclidean space with cartesian coordinates, we have

$$\Gamma_{\nu\sigma}^{\epsilon}(x) = 0.$$

So,

$$B_{\mu\nu\sigma}^{\epsilon} = 0 \text{ in this coordinate system.}$$

But if $B_{\mu\nu\sigma}^{\epsilon} = 0$ in one coordinate system, the components are zero in all coordinate systems.

Hence if $B_{\mu\nu\sigma}^{\epsilon} = 0$, the space is Euclidean.

Problem 33. If $R_{\mu\nu} = Kg_{\mu\nu}$, then show that $R = K$ where R is called the scalar curvature.

Given that $R_{\mu\nu} = Kg_{\mu\nu}$.

We have

$$\begin{aligned} R &= g^{\mu\nu} R_{\mu\nu} \\ &= g^{\mu\nu} Kg_{\mu\nu} \end{aligned}$$

Hence

$$R = K \text{ as } g^{\mu\nu} g_{\mu\nu} = 1.$$

Problem 34. Prove the Bianchi's Identity

$$B_{\mu\nu\sigma, \rho}^{\epsilon} + B_{\mu\sigma\rho, \nu}^{\epsilon} + B_{\mu\rho\nu, \sigma}^{\epsilon} = 0.$$

(Agra, 1964, 70)

We know that

$$B_{\mu\nu\sigma}^{\epsilon} = \frac{\partial \Gamma_{\mu\nu}^{\epsilon}}{\partial x_{\sigma}} - \frac{\partial \Gamma_{\mu\sigma}^{\epsilon}}{\partial x_{\nu}} + \Gamma_{\mu\nu}^{\alpha} \Gamma_{\alpha\sigma}^{\epsilon} - \Gamma_{\mu\sigma}^{\alpha} \Gamma_{\alpha\nu}^{\epsilon}$$

If we differentiate this equation and evaluate at the origin of a geodesic co-ordinate system, then we have

$$B_{\mu\nu\sigma, \rho}^{\epsilon} = \frac{\partial^2 \Gamma_{\mu\nu}^{\epsilon}}{\partial x_{\rho} \partial x_{\sigma}} - \frac{\partial^2 \Gamma_{\mu\sigma}^{\epsilon}}{\partial x_{\rho} \partial x_{\nu}}$$

(since the Christoffel's symbols referred to the geodesic co-ordinate system vanish at O , the origin, by §3.3.)

Similarly,

$$B_{\mu\sigma\rho,\nu}^{\epsilon} = \frac{\partial^2 \Gamma_{\mu\sigma}^{\epsilon}}{\partial x_{\nu} \partial x_{\rho}} - \frac{\partial^2 \Gamma_{\mu\rho}^{\epsilon}}{\partial x_{\nu} \partial x_{\sigma}}$$

and

$$B_{\mu\rho\nu,\sigma}^{\epsilon} = \frac{\partial \Gamma_{\mu\rho}^{\epsilon}}{\partial x_{\sigma} \partial x_{\nu}} - \frac{\partial^2 \Gamma_{\mu\nu}^{\epsilon}}{\partial x_{\sigma} \partial x_{\rho}}$$

Adding all these three equations, we get

$$B_{\mu\nu\sigma,\rho}^{\epsilon} + B_{\mu\sigma\rho,\nu}^{\epsilon} + B_{\mu\rho\nu,\sigma}^{\epsilon} = 0.$$

Problem 35. Show that the construction of a uniform vector field is only possible when the curvature tensor vanishes.

Or

Prove that when the Riemann-Christoffel tensor vanishes, the differential-equations $A_{\mu\nu} = \frac{\partial A_{\mu}}{\partial x_{\nu}} - \{\mu\nu, \alpha\} A_{\alpha} = 0$ are integrable.

(Agra, 1966)

Consider the differential equation

$$A_{\mu,\nu} = \frac{\partial A_{\mu}}{\partial x_{\nu}} - \Gamma_{\mu\nu}^{\alpha} A_{\alpha} = 0. \quad \dots (1)$$

This gives

$$\frac{\partial A_{\mu}}{\partial x_{\nu}} = \Gamma_{\mu\nu}^{\alpha} A_{\alpha}. \quad \dots (2)$$

Consider the integral of L.H.S. of (2).

$$\int \frac{\partial A_{\mu}}{\partial x_{\nu}} dx_{\nu} = \int \left[\frac{\partial A_{\mu}}{\partial x_1} dx_1 + \dots \right] \\ = \int dA_{\mu} = A_{\mu}.$$

This follows that the left hand side of (2) multiplied by x_{ν} is a perfect differential and as such the right hand side of (2) multiplied by dx_{ν} must be a perfect differential. i.e.,

$\Gamma_{\mu\nu}^{\alpha} A_{\alpha} dx_{\nu}$ is a perfect differential,

$$i.e. \quad \Gamma_{\mu\nu}^1 A_1 dx_{\nu} + \Gamma_{\mu\nu}^2 A_2 dx_{\nu} + \dots + \Gamma_{\mu 1}^{\alpha} A_{\alpha} dx_1 + \Gamma_{\mu 2}^{\alpha} A_{\alpha} dx_2 + \dots$$

is a perfect differential

$$i.e. \quad \Gamma_{\mu\nu}^{\alpha} A_{\alpha} dx_{\nu} + \Gamma_{\mu\sigma}^{\alpha} A_{\alpha} dx_{\sigma} \quad \text{is a perfect differential}$$

which is of the form $M dx + N dy = 0$ and this is perfect differential if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} \quad i.e. \quad \frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} = 0.$$

As such $\Gamma_{\mu\nu}^{\alpha} A_{\alpha} dx_{\nu} + \Gamma_{\mu\sigma}^{\alpha} A_{\alpha} dx_{\sigma}$ (i.e. $\Gamma_{\mu\nu}^{\alpha} A_{\alpha} dx_{\nu}$) is perfect differential if

$$\frac{\partial}{\partial x_{\sigma}} (\Gamma_{\mu\nu}^{\alpha} A_{\alpha}) - \frac{\partial}{\partial x_{\nu}} (\Gamma_{\mu\sigma}^{\alpha} A_{\alpha}) = 0.$$

or if

$$A_{\alpha} \left(\frac{\partial \Gamma_{\mu\nu}^{\alpha}}{\partial x_{\sigma}} - \frac{\partial \Gamma_{\mu\sigma}^{\alpha}}{\partial x_{\nu}} + \Gamma_{\mu\nu}^{\alpha} \frac{\partial A_{\alpha}}{\partial x_{\nu}} - \Gamma_{\mu\sigma}^{\alpha} \frac{\partial A_{\alpha}}{\partial x_{\nu}} \right) = 0.$$

With the help of (2), this yields

$$A_{\alpha} \left(\frac{\partial \Gamma_{\mu\nu}^{\alpha}}{\partial x_{\sigma}} - \frac{\partial \Gamma_{\mu\sigma}^{\alpha}}{\partial x_{\nu}} \right) + (\Gamma_{\mu\nu}^{\alpha} \Gamma_{\alpha\sigma}^{\epsilon} - \Gamma_{\mu\sigma}^{\alpha} \Gamma_{\alpha\nu}^{\epsilon}) A_{\epsilon} = 0.$$

Changing the dummy suffix α to ϵ in the first term, we get

$$A_{\epsilon} \left[\frac{\partial \Gamma_{\mu\nu}^{\epsilon}}{\partial x_{\sigma}} - \frac{\partial \Gamma_{\mu\sigma}^{\epsilon}}{\partial x_{\nu}} + \Gamma_{\mu\nu}^{\alpha} \Gamma_{\alpha\sigma}^{\epsilon} - \Gamma_{\mu\sigma}^{\alpha} \Gamma_{\alpha\nu}^{\epsilon} \right] = 0$$

i.e.

$$A_{\epsilon} B_{\mu\sigma\nu}^{\epsilon} = 0.$$

Since A_{ϵ} is arbitrary, therefore $B_{\mu\sigma\nu}^{\epsilon} = 0$.

Hence when $B_{\mu\sigma\nu}^{\epsilon}$ vanishes the differential dA_{μ} determined by (1)

will be a perfect differential and as such $\int dA_{\mu}$ between any two points will be independent of the path of integration. Then the vector A_{μ} can be parallelly displaced to any point thereby giving a unique result independent of the path of transfer. Displacement of such a vector gives a uniform vector field. Conclusively the construction of a uniform vector field is only possible where the curvature tensor vanishes.

3.20. SOME IMPORTANT RESULTS

We have $g_{\mu\nu} g^{\mu\alpha} = g_{\nu}^{\alpha} = 0$ or 1.

Its differentiation gives

$$g^{\mu\alpha} dg_{\mu\nu} + g_{\mu\nu} dg^{\mu\alpha} = 0$$

or

$$g^{\mu\alpha} dg_{\mu\nu} = -g_{\mu\nu} dg^{\mu\alpha}.$$

Multiplying throughout by $g^{\nu\beta}$, we get

$$g^{\mu\alpha} g^{\nu\beta} dg_{\mu\nu} = -g_{\mu\nu} g^{\nu\beta} dg^{\mu\alpha} \\ = -g_{\mu}^{\beta} dg^{\mu\alpha} \\ = -dg^{\alpha\beta}. \quad \dots (1)$$

Similarly,

$$dg_{\alpha\beta} = -g_{\mu\alpha} g_{\nu\beta} dg^{\mu\nu}. \quad \dots (2)$$

(Agra, 1963)

Multiplying (2) throughout by $A^{\alpha\beta}$, we get

$$A^{\alpha\beta} dg_{\alpha\beta} = -(g_{\mu\alpha} g_{\nu\beta} A^{\alpha\beta}) dg^{\mu\nu} \\ = -(g_{\mu\alpha} A^{\alpha\nu}) dg^{\mu\nu}$$

$$\begin{aligned}
 &= -A_{\mu\nu} dg^{\mu\nu} \\
 &= -A_{\alpha\beta} dg^{\alpha\beta} \quad (\text{replacing } \mu, \nu \text{ by } \alpha, \beta \text{ respectively}), \\
 \text{i.e. } A^{\alpha\beta} dg_{\alpha\beta} &= -A_{\alpha\beta} dg^{\alpha\beta}. \quad \dots (3) \\
 &\quad \quad \quad (\text{Agra, 1967})
 \end{aligned}$$

But for any other tensor $B_{\alpha\beta}$, we have

$$A^{\alpha\beta} dB_{\alpha\beta} = A_{\alpha\beta} dB^{\alpha\beta}.$$

Now consider dg formed by taking the differential of each of $g_{\mu\nu}$ and multiplying by its cofactor $g^{\mu\nu}$ in the determinant g .

$$\begin{aligned}
 \frac{dg}{g} &= g^{\mu\nu} dg_{\mu\nu} \\
 &= -g_{\mu\nu} dg^{\mu\nu} \text{ by (3)} \quad \dots (4)
 \end{aligned}$$

But we have

$$\begin{aligned}
 \Gamma_{\mu\sigma}^{\sigma} &= \frac{1}{2} g^{\sigma\lambda} \left\{ \frac{\partial g_{\mu\lambda}}{\partial x_{\sigma}} + \frac{\partial g_{\sigma\lambda}}{\partial x_{\mu}} - \frac{\partial g_{\mu\sigma}}{\partial x_{\lambda}} \right\} \\
 &= \frac{1}{2} g^{\sigma\lambda} \frac{\partial g_{\sigma\lambda}}{\partial x_{\mu}}, \text{ other two terms cancel, by} \\
 &\quad \text{interchange of } \sigma \text{ and } \lambda \\
 &= \frac{1}{2g} \frac{dg}{dx_{\mu}} \text{ by (4)} \quad \dots (5) \\
 &= \frac{\partial}{\partial x_{\mu}} \log \sqrt{(-g)}. \quad \dots (6) \\
 &\quad \quad \quad (\text{Agra, 1959, 64, 68})
 \end{aligned}$$

Since g is always negative for real coordinates, we therefore use $\sqrt{(-g)}$.

Again,

$$\begin{aligned}
 A_{,\nu}^{\nu} &= \frac{\partial A^{\nu}}{\partial x_{\nu}} + \Gamma_{\mu\nu}^{\nu} A^{\mu} \\
 &= \frac{\partial A^{\nu}}{\partial x_{\nu}} + \frac{1}{2g} \frac{\partial g}{\partial x_{\nu}} A^{\nu} \text{ by (5)} \\
 &= \frac{\partial A^{\nu}}{\partial x_{\nu}} + \frac{1}{2g} \frac{\partial g}{\partial x_{\nu}} A^{\nu} \text{ on replacing } \mu \text{ by } \nu \\
 \text{or } A_{,\nu}^{\nu} &= \frac{1}{\sqrt{g}} \frac{\partial}{\partial x_{\nu}} (\sqrt{g} A^{\nu}) \quad \dots (7) \\
 &\quad \quad \quad (\text{taking the absolute value of } g).
 \end{aligned}$$

Further, let

$$h_{\nu\tau} = a_{\mu\nu} a_{\sigma\tau} x_{\mu} x_{\sigma},$$

where $a_{\mu\nu}$ denotes constant coefficients.

Then

$$\begin{aligned}
 \frac{\partial h_{\nu\tau}}{\partial x_{\sigma}} &= a_{\mu\nu} a_{\sigma\tau} \left(\frac{\partial x_{\mu}}{\partial x_{\sigma}} x_{\sigma} + \frac{\partial x_{\sigma}}{\partial x_{\mu}} x_{\mu} \right) \\
 &= a_{\mu\nu} a_{\sigma\tau} (g_{\alpha}^{\mu} x_{\sigma} + g_{\alpha}^{\sigma} x_{\mu}) \text{ by §3.3.}
 \end{aligned}$$

Repeating the process,

$$\begin{aligned}
 \frac{\partial^2 h_{\nu\tau}}{\partial x_{\alpha} \partial x_{\beta}} &= a_{\mu\nu} a_{\sigma\tau} (g_{\alpha}^{\mu} g_{\beta}^{\sigma} + g_{\alpha}^{\sigma} g_{\beta}^{\mu}) \\
 &= a_{\alpha\nu} a_{\beta\tau} + a_{\beta\nu} a_{\alpha\tau}.
 \end{aligned}$$

Here changing dummy suffixes, we have

$$\frac{\partial^2}{\partial x_{\mu} \partial x_{\nu}} (a_{\mu\nu} a_{\sigma\mu} x_{\mu} x_{\sigma}) = a_{\mu\nu} a_{\sigma\tau} + a_{\sigma\nu} a_{\mu\tau}. \quad \dots (8)$$

Similarly if $a_{\mu\nu\sigma}$ is symmetrical in its suffixes, we have

$$\frac{\partial^3}{\partial x_{\mu} \partial x_{\nu} \partial x_{\sigma}} (a_{\mu\nu\sigma} x_{\mu} x_{\nu} x_{\sigma}) = 6a_{\mu\nu\sigma}. \quad \dots (9)$$

3.21. TENSOR FORMS OF OPERATORS

[A] Gradient. If ϕ be an invariant quantity then

$$\nabla\phi = \text{grad } \phi = \phi_{,\nu} = \frac{\partial\phi}{\partial x_{\nu}}$$

where $\phi_{,\nu}$ is the covariant derivative of ϕ w.r.t. x_{ν} .

[B] Divergence. Divergence of A^{μ} is defined as the contraction of its covariant derivative w.r.t. x_{μ} i.e. the contraction of $A^{\mu}_{,\mu}$, so that by (7) of § 3.20 we have

$$\text{div } A^{\mu}_{,\mu} = \sqrt{\frac{1}{g}} \frac{\partial}{\partial x_{\mu}} (\sqrt{g} A^{\mu})$$

[C] Curl. The curl of A_{μ} is defined as

$$A_{\mu,\nu} - A_{\nu,\mu} = \frac{\partial A_{\mu}}{\partial x_{\nu}} - \frac{\partial A_{\nu}}{\partial x_{\mu}}$$

Also curl $A_{\mu,\nu} = -\epsilon^{ijk} A_{\mu,j} k$

[D] Laplacian. The Laplacian of ϕ is divergence of gradient ϕ i.e. $\nabla^2\phi = \text{div } \phi_{,\nu}$.

But $\nabla\phi = \frac{\partial\phi}{\partial x_{\nu}}$ a covariant tensor of rank one defined as covariant derivative of ϕ , written as $\phi_{,\mu}$. The contravariant tensor of rank one associated with $\phi_{,\mu}$ is

$$A^{\alpha} = g^{\alpha\mu} \frac{\partial\phi}{\partial x^{\mu}}$$

Hence

$$\begin{aligned}
 \nabla^2\phi &= \text{div} \left(g^{\alpha\mu} \frac{\partial\phi}{\partial x^{\mu}} \right) \\
 &= \frac{1}{\sqrt{g}} \frac{\partial}{\partial x_{\alpha}} \left(\sqrt{g} g^{\alpha\mu} \frac{\partial\phi}{\partial x^{\mu}} \right)
 \end{aligned}$$

when g^{20} , $\sqrt{g}$ must be replaced by $\sqrt{-g}$ while the cases $g > 0$ and $g < 0$ are included in $\sqrt{|g|}$ instead of $\sqrt{g}$.

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 36. Write the law of transformation for the tensors

$$(i) A_{\mu\nu}^{\sigma}, (ii) A_{\sigma}^{\mu\nu}, (iii) B_{\mu\nu\sigma}^{\alpha\beta}, (iv) B_{\alpha\beta}^{\mu\nu\sigma}, (v) C_{\mu\nu}, (vi) C^{\mu\nu}, (vii) D_{\mu}, (viii) D^{\mu}.$$

We have,

$$(i) A_{\xi\eta}^{\zeta} = \frac{\partial x^{\zeta}}{\partial x_{\sigma}} \frac{\partial x_{\mu}}{\partial x^{\xi}} \frac{\partial x_{\nu}}{\partial x^{\eta}} A_{\mu\nu}^{\sigma}$$

$$\begin{aligned}
 \text{(ii)} \quad A'_{\zeta}^{\xi\eta} &= \frac{\partial x_{\xi}'}{\partial x_{\mu}} \frac{\partial x_{\eta}'}{\partial x_{\nu}} \frac{\partial x_{\sigma}}{\partial x_{\delta}} A_{\sigma}^{\mu\nu} \\
 \text{(iii)} \quad B'_{\xi\eta\zeta}^{\gamma\delta} &= \frac{\partial x'_{\gamma}}{\partial x_{\alpha}} \frac{\partial x'_{\delta}}{\partial x_{\beta}} \frac{\partial x_{\mu}}{\partial x'_{\xi}} \frac{\partial x_{\nu}}{\partial x'_{\eta}} \frac{\partial x_{\sigma}}{\partial x'_{\zeta}} B_{\mu\nu\sigma}^{\alpha\beta} \\
 \text{(iv)} \quad B'_{\gamma\delta}^{\xi\eta\zeta} &= \frac{\partial x'_{\xi}}{\partial x_{\mu}} \frac{\partial x'_{\eta}}{\partial x_{\nu}} \frac{\partial x'_{\zeta}}{\partial x_{\sigma}} \frac{\partial x_{\alpha}}{\partial x'_{\gamma}} \frac{\partial x_{\beta}}{\partial x'_{\delta}} B_{\alpha\beta}^{\mu\nu\sigma} \\
 \text{(v)} \quad C'_{\alpha\beta} &= \frac{\partial x_{\mu}}{\partial x_{\alpha}'} \frac{\partial x_{\nu}}{\partial x_{\beta}'} C_{\mu\nu} \\
 \text{(vi)} \quad C'_{\alpha\beta} &= \frac{\partial x_{\alpha}'}{\partial x_{\mu}} \frac{\partial x_{\beta}'}{\partial x_{\nu}} C^{\mu\nu} \\
 \text{(vii)} \quad D'_{\alpha} &= \frac{\partial x_{\mu}}{\partial x'_{\alpha}} D_{\mu} \text{ and (viii) } D'^{\alpha} = \frac{\partial x'_{\alpha}}{\partial x_{\mu}} D^{\mu}
 \end{aligned}$$

Problem 37. If the tensors a_{ik} , a_{ikj} , a_{ijkl} are antisymmetric in every pair of indices, find their independent components in four dimensions. (Agra, 1966)

If a_{ik} is a skew-symmetric set, then $a_{ik} = -a_{ki}$ for every pair of values i, k . In particular

$$a_{11} = -a_{11}, a_{22} = -a_{22}, a_{33} = -a_{33}, a_{44} = -a_{44} \text{ giving } a_{11} = a_{22} = a_{33} = a_{44} = 0.$$

i.e., the components arising for equal values of the suffixes are all separately zero. Their number is four.

In all there are 4^2 i.e., 16 components of which 4 are zero. The remaining (16-4) i.e., 12 components divide themselves into pairs such that the two members of any pair are equal and opposite as $a_{12} = -a_{21}$ etc. Thus the total number of independent components of an anti-symmetric tensor a_{ik} of order 2 and dimension 4 is 6.

Similar arguments will give the independent component of a_{ikj} and a_{ijkl} .

Problem 38. Show that the number of distinct non-vanishing components of the covariant Riemann Christoffel curvature tensor does not exceed $\frac{1}{2}n^2(n^2-1)$.

Since $B_{\mu\nu\sigma\rho}$ is skew-symmetric in μ, ρ and ν, σ therefore the number of non-vanishing components contributed by μ, ρ is nC_2 i.e., $\frac{n(n-1)}{2}$ also the contributed by ν, σ is $\frac{1}{2}n(n-1)$. Thus the number of non-vanishing components of Riemann tensor does not exceed $\frac{n(n-1)}{2} \times \frac{n(n-1)}{2}$ i.e., $\frac{n^2}{4}(n-1)^2$

But the number of relations in Bianchi's identity i.e.,

$$B_{\mu\nu\sigma\rho} + B_{\mu\sigma\rho\nu} + B_{\mu\rho\nu\sigma} = 0 \text{ is } {}^nC_3 \text{ i.e., } \frac{n^2(n-1)(n-2)}{6}$$

$\therefore$ The number of distinct non-vanishing components does not exceed

$$\begin{aligned}
 \frac{n^2}{4}(n-1)^2 - \frac{n^2(n-1)(n-2)}{6} &= \frac{n^2}{12}(n-1)[3(n-1)-2(n-2)] \\
 &= \frac{n^2}{12}(n-1)(n+1) = \frac{n^2}{12}(n^2-1)
 \end{aligned}$$

Problem 39. Establish the result

$$(A_{\mu})_{\nu\sigma} - (A_{\mu})_{\sigma\nu} = A^{\epsilon} B_{\epsilon\nu\sigma}^{\mu}$$

and prove that $B_{\epsilon\nu\sigma}^{\mu}$ has only twenty algebraically independent components.

(Agra, 1958, 65, 67)

For first part see (3) of §3.18.

For second part see §3.19.

Problem 40. Explain what is meant by covariant, contravariant and mixed tensors. If $B_{\nu\sigma}$ is an arbitrary covariant tensor and $A(\mu, \nu) B_{\nu\sigma} = C_{\mu\sigma}$ where $C_{\mu\sigma}$ is a tensor, then prove that $A(\mu, \nu)$ is a mixed tensor.

Define the Christoffel three-index symbol of the second kind. Prove that it is not a tensor. Verify that $g_{\mu\nu}$ behaves as a constant in covariant differentiation.

(Agra, 1958, 63, 65)

For definition see §3.4.

Suppose that $A(\mu, \nu) B_{\nu\sigma} = C_{\mu\sigma}$ in the coordinate system x_{α} .

Then in the transformed coordinate x_{α}' , we have

$$A'(\beta, \gamma) B'_{\gamma\epsilon} = C'_{\beta\epsilon}$$

$$\begin{aligned}
 \text{Consider, } A'(\beta, \gamma) \frac{\partial x_{\nu}}{\partial x_{\gamma}'} \frac{\partial x_{\sigma}}{\partial x_{\epsilon}'} B_{\nu\sigma} &= \frac{\partial x_{\sigma}}{\partial x_{\epsilon}'} \frac{\partial x_{\mu}}{\partial x_{\beta}'} C_{\mu\sigma} \\
 &= \frac{\partial x_{\sigma}}{\partial x_{\epsilon}'} \frac{\partial x_{\mu}}{\partial x_{\beta}'} A(\mu, \nu) B_{\nu\sigma}
 \end{aligned}$$

$$\text{or } \frac{\partial x_{\sigma}}{\partial x_{\epsilon}'} \left[\frac{\partial x_{\nu}}{\partial x_{\gamma}'} \frac{\partial x_{\mu}}{\partial x_{\beta}'} A'(\beta, \gamma) - \frac{\partial x_{\sigma}}{\partial x_{\beta}'} A(\mu, \nu) \right] B_{\nu\sigma} = 0 \quad \dots (1)$$

Multiplying by $\frac{\partial x_{\epsilon}'}{\partial x_{\eta}}$ we have, $\frac{\partial x_{\sigma}}{\partial x_{\epsilon}'} \frac{\partial x_{\epsilon}'}{\partial x_{\eta}} = \frac{\partial x_{\sigma}}{\partial x_{\eta}} = \delta_{\eta}^{\sigma}$ so that $\delta_{\eta}^{\sigma} B_{\nu\sigma} = B_{\nu\eta}$

$\therefore$ (1) becomes

$$\left[\frac{\partial x_{\nu}}{\partial x_{\gamma}'} A'(\beta, \gamma) - \frac{\partial x_{\mu}}{\partial x_{\beta}'} A(\mu, \nu) \right] B_{\nu\eta} = 0$$

$$\text{or } \frac{\partial x_{\nu}}{\partial x_{\gamma}'} A'(\beta, \gamma) - \frac{\partial x_{\mu}}{\partial x_{\beta}'} A(\mu, \nu) = 0, B_{\nu\eta} \text{ being arbitrary.}$$

$$\text{Multiplying by } \frac{\partial x_{\epsilon}'}{\partial x_{\nu}} \text{ we get } \frac{\partial x_{\nu}}{\partial x_{\gamma}'} \frac{\partial x_{\epsilon}'}{\partial x_{\nu}} A'(\beta, \gamma) = \frac{\partial x_{\mu}}{\partial x_{\beta}'} \frac{\partial x_{\epsilon}'}{\partial x_{\nu}} A(\mu, \nu)$$

$$\text{i.e., } \delta_{\gamma}^{\epsilon} A'(\beta, \gamma) = \frac{\partial x_{\mu}}{\partial x_{\beta}'} \frac{\partial x_{\epsilon}'}{\partial x_{\nu}} A(\mu, \nu)$$

$$\text{or } A'(\beta, \eta) = \frac{\partial x_{\mu}}{\partial x_{\beta}'} \frac{\partial x_{\eta}'}{\partial x_{\nu}} A(\mu, \nu)$$

which follows that $A(\mu, \nu)$ is a mixed tensor.

Now in §3.12, the Christoffel's 3-index symbol of the second kind has been defined as

$$\Gamma_{\mu\nu}^{\sigma} = \{\mu\nu, \sigma\} = \frac{1}{2} g^{\sigma\lambda} \left(\frac{\partial g_{\mu\lambda}}{\partial x_{\nu}} + \frac{\partial g_{\nu\lambda}}{\partial x_{\mu}} - \frac{\partial g_{\mu\nu}}{\partial x_{\lambda}} \right)$$

and in §3.14, we have derived the transformation law for it in equation (4) as

$$\{\beta\gamma, \alpha\} = \{\mu\nu, \lambda\} \frac{\partial x_{\mu}}{\partial x_{\beta}'} \frac{\partial x_{\nu}}{\partial x_{\gamma}'} \frac{\partial x_{\alpha}}{\partial x_{\lambda}'} + \frac{\partial^2 x_{\lambda}'}{\partial x_{\beta} \partial x_{\gamma}} \frac{\partial x_{\alpha}}{\partial x_{\lambda}'}$$

which clearly shows that $\{\beta\gamma, \alpha\}$ or $\Gamma_{\beta\gamma}^{\alpha}$ are not the components of a tensor unless the second terms on the right are zero.

Since the covariant derivative of $g_{\mu\nu}$ is zero as is shown below, the tensor $g_{\mu\nu}$ may be treated as constant in covariant differentiation.

Covariant derivative of $g_{\mu\nu}$ is given by (1) of §3.17,

$$\partial_{\mu\nu, \sigma} = \frac{\partial g_{\mu\nu}}{\partial x_{\sigma}} - \Gamma_{\mu\sigma}^{\alpha} g_{\alpha\nu} - \Gamma_{\nu\sigma}^{\alpha} g_{\mu\alpha}$$

$$\text{But } \Gamma_{\mu\sigma}^{\alpha} = \frac{1}{2} g^{\alpha\nu} \left(\frac{\partial g_{\mu\nu}}{\partial x_{\sigma}} + \frac{\partial g_{\sigma\nu}}{\partial x_{\mu}} - \frac{\partial g_{\mu\sigma}}{\partial x_{\nu}} \right)$$

$$\begin{aligned}
 \text{So that } g_{\alpha\nu} \Gamma_{\mu\sigma}^{\alpha} &= \frac{1}{2} g_{\alpha\nu} g^{\alpha\nu} \left(\frac{\partial g_{\mu\nu}}{\partial x_{\sigma}} + \frac{\partial g_{\sigma\nu}}{\partial x_{\mu}} - \frac{\partial g_{\mu\sigma}}{\partial x_{\nu}} \right) \\
 &= \frac{1}{2} \left(\frac{\partial g_{\mu\nu}}{\partial x_{\sigma}} + \frac{\partial g_{\sigma\nu}}{\partial x_{\mu}} - \frac{\partial g_{\mu\sigma}}{\partial x_{\nu}} \right) \text{ as } g_{\alpha\nu} g^{\alpha\nu} = 1
 \end{aligned}$$

Similarly, $g_{\mu\alpha} \Gamma_{\nu\alpha}^{\alpha} = \frac{1}{2} \left(\frac{\partial g_{\mu\nu}}{\partial x_{\sigma}} + \frac{\partial g_{\mu\sigma}}{\partial x_{\nu}} - \frac{\partial g_{\sigma\nu}}{\partial x_{\mu}} \right)$

$$\therefore - \left[\Gamma_{\mu\sigma}^{\alpha} g_{\alpha\nu} + \Gamma_{\nu\alpha}^{\alpha} g_{\mu\alpha} \right] = - \frac{\partial g_{\mu\nu}}{\partial x_{\sigma}}$$

$$\text{Hence } \partial_{\mu\nu, \sigma} = \frac{\partial g_{\mu\nu}}{\partial x_{\sigma}} - \frac{\partial g_{\mu\sigma}}{\partial x_{\nu}} = 0.$$

Problem 41. State the law of transformation of a mixed tensor. Show how the transformation is affected when the tensor is subjected to a Contraction.

Calculate the Christoffel 3-index symbol of the second kind $\{\mu, \sigma\}$

Verify that the covariant derivative of the tensor $g^{\nu\alpha}$ vanishes. (Agra, 1959, 65)

For first part see (5) and (6) of §3.4 and III of §3.7.

Now we have

$$\{\mu\nu, \sigma\} = \frac{1}{2} g^{\alpha\lambda} \left(\frac{\partial g_{\mu\lambda}}{\partial x_{\nu}} + \frac{\partial g_{\nu\lambda}}{\partial x_{\mu}} - \frac{\partial g_{\mu\nu}}{\partial x_{\lambda}} \right)$$

But $\nu = \sigma$, so that

$$\begin{aligned} \{\mu\sigma, \sigma\} &= \frac{1}{2} g^{\sigma\lambda} \left(\frac{\partial g_{\mu\lambda}}{\partial x_{\sigma}} + \frac{\partial g_{\sigma\lambda}}{\partial x_{\mu}} - \frac{\partial g_{\mu\sigma}}{\partial x_{\lambda}} \right) \\ &= \frac{\partial}{\partial x_{\mu}} \log \sqrt{-g} \text{ by (8) of §3.20.} \end{aligned}$$

For the last part to show that $g^{\mu\alpha}_{;\sigma} = 0$ see Problem 30 (a).

Problem 42. Define geodesics and obtain their equations with the help of a variational principle. (Agra, 1963, 65, 68)

See §3.13.

Problem 43. Prove that

$$\frac{dg_{\alpha\beta}}{ds} = -g_{\mu\alpha} g_{\nu\beta} \frac{dg^{\mu\nu}}{ds} \quad (\text{Agra, 1963})$$

We have

$$g_{\mu\alpha} g^{\mu\nu} = g_{\alpha}^{\nu} = 0 \text{ or } 1.$$

Its differentiation gives

$$\begin{aligned} g_{\mu\alpha} \frac{dg^{\mu\nu}}{ds} + g^{\mu\nu} \frac{dg_{\mu\alpha}}{ds} &= 0 \\ \text{or } g^{\mu\nu} \frac{dg_{\mu\alpha}}{ds} &= -g_{\mu\alpha} \frac{dg^{\mu\nu}}{ds} \end{aligned}$$

Multiplying throughout by $g_{\nu\beta}$, we get

$$g^{\mu\nu} g_{\nu\beta} \frac{dg_{\mu\alpha}}{ds} = -g_{\mu\alpha} g_{\nu\beta} \frac{dg^{\mu\nu}}{ds}$$

or

$$g_{\beta}^{\nu} \frac{dg_{\mu\alpha}}{ds} = -g_{\mu\alpha} g_{\nu\beta} \frac{dg^{\mu\nu}}{ds}$$

or

$$\frac{dg_{\alpha\beta}}{ds} = -g_{\mu\alpha} g_{\nu\beta} \frac{dg^{\mu\nu}}{ds}$$

Problem 44. Show with the usual notation

$$(i) \quad \frac{\partial g_{ij}}{\partial x_k} = [j, i k] + [i, j k]$$

$$(ii) \quad \left\{ \frac{i}{ik} \right\} = \frac{\partial}{\partial x_k} \log \sqrt{g}$$

(i) Using (i) of §3.12, we have

$$[j, i k] = \frac{1}{2} \left(\frac{\partial g_{ij}}{\partial x_k} + \frac{\partial g_{kj}}{\partial x_i} - \frac{\partial g_{ik}}{\partial x_j} \right) \quad \dots (1)$$

$$\text{and } [i, j k] = \frac{1}{2} \left(\frac{\partial g_{ji}}{\partial x_k} + \frac{\partial g_{ki}}{\partial x_j} - \frac{\partial g_{kj}}{\partial x_i} \right) = \frac{1}{2} \left(\frac{\partial g_{ij}}{\partial x_k} + \frac{\partial g_{ik}}{\partial x_j} - \frac{\partial g_{kj}}{\partial x_i} \right) \quad \dots (2)$$

Adding (1) and (2) we get $[j, i k] + [i, j k] = \frac{\partial g_{ij}}{\partial x_k}$

$$(ii) \text{ From (4) of §3.20 we have } \frac{dg}{g} = -g_{\mu\nu} \frac{dg^{\mu\nu}}{ds} \quad \dots (1)$$

$$\begin{aligned} \text{So that } \left\{ \frac{i}{ik} \right\} &= \{i, i k\} = \frac{1}{2} g^{i\lambda} \left(\frac{\partial g_{k\lambda}}{\partial x_i} + \frac{\partial g_{i\lambda}}{\partial x_k} - \frac{\partial g_{ki}}{\partial x_{\lambda}} \right) \\ &= \frac{1}{2} g^{i\lambda} \frac{\partial g_{i\lambda}}{\partial x_k} \text{ other two terms cancel by interchange of } i \text{ and } \lambda \\ &= \frac{1}{2g} \frac{dg}{dx_k} \text{ by (1)} \\ &= \frac{\partial}{\partial x_k} \log \sqrt{g} \text{ if } g \text{ is positive.} \end{aligned}$$

Problem 45 (a) Define the Kronecker and alternating tensors.

(b) Prove the following identities

$$(i) \quad \delta_{ii} = 3$$

$$(ii) \quad \delta_{ik} \epsilon_{ikm} = 0$$

$$(iii) \quad \epsilon_{ikl} \epsilon_{mnp} = \delta_{im} \delta_{kp} - \delta_{ip} \delta_{km} = 0 \quad (\text{Agra, 1969})$$

(a) **Kronecker delta** is a notation defined by

$$\begin{aligned} \delta_{ij} \text{ or } \delta_i^j &= 0 \text{ if } i \neq j \\ &= 1 \text{ if } i = j \end{aligned}$$

It is a mixed tensor of rank 2 as it transforms like a mixed tensor of rank two.

Alternating Tensor (or permutation tensor or Epsilon tensor) is an abstract entity of order (rank) 3 and dimension 3 such that its components are invariant for any coordinate system. It is denoted by ϵ_{ijk} and defined as

$$\epsilon_{ijk} = \begin{cases} 0, & \text{if any two of } i, j, k \text{ are equal} \\ 1, & \text{if } i, j, k \text{ is a cyclic (i.e., even) permutation of } 1, 2, 3 \\ -1, & \text{if } i, j, k \text{ is an anticyclic (i.e., odd) permutation of } 1, 2, 3 \end{cases}$$

i.e., for unequal values of the suffixes, we have

$$\epsilon_{123} = \epsilon_{231} = \epsilon_{312} = 1$$

$$\epsilon_{132} = \epsilon_{213} = \epsilon_{321} = -1$$

(b) (i) In three dimensions, we have

$$\begin{aligned} \delta_{ii} &= \delta_{11} + \delta_{22} + \delta_{33} \\ &= 1 + 1 + 1 \text{ as } \delta_{ij} = 1 \text{ for } i = j \\ &= 3 \end{aligned}$$

(ii) We have

$$\begin{aligned} \delta_{ik} &= 0 \text{ for } i \neq k \\ &= 1 \text{ for } i = k \end{aligned}$$

and

$$\begin{aligned} \epsilon_{ikm} &= \pm 1 \text{ for } i \neq k \neq m \\ &= 0 \text{ for } i = k \neq m \text{ or } i \neq k = m \text{ or } k \neq i = m \end{aligned}$$

Hence in either case

$$\delta_{ik} \epsilon_{ikm} = 0$$

(iii) We have to show that

$$\epsilon_{iks} \epsilon_{mps} = \delta_{im} \delta_{kp} - \delta_{ip} \delta_{km} = 0 \quad \dots (1)$$

Consider, $\epsilon_{iks} \epsilon_{mps} = \delta_{im} \delta_{kp} - \delta_{ip} \delta_{km}$

Either side of (1) is a tensor of order 4. In 3-dimensional space this result can be written as

$$\epsilon_{ik1} \epsilon_{mp1} + \epsilon_{ik2} \epsilon_{mp2} + \epsilon_{ik3} \epsilon_{mp3} = \delta_{im} \delta_{kp} - \delta_{ip} \delta_{km}$$

when $i = k$, we have

$$\epsilon_{2k1} = \epsilon_{ik2} = \epsilon_{ik3} = 0 \text{ so that L.H.S. of (1) = 0}$$

and then $\delta_{im} = \delta_{kp} = \delta_{ip} = \delta_{km} = 0$ so that R.H.S. of (1) = 0

The same result holds when instead of $i = k$, $m = p$.

Again when $i \neq k$ and when $m \neq p$ and if the pair of unequal values of i, k is different from the pair of unequal values of m, p then also

$$\text{L.H.S.} = 0 = \text{R.H.S.}$$

As such we are left to consider the possibilities when i, k and m, p have the pairs of values 1, 2; 1, 3; 2, 3; 2, 1; 3, 1; 3, 2

Taking the first case we find

$$i=1, k=2, m=1, p=2, i=1, k=2, m=2, p=1$$

$$i=2, k=1, m=1, p=2, i=2, k=1, m=2, p=1$$

which give L.H.S. = 1 = R.H.S.; L.H.S. = -1 = R.H.S.

$$\text{L.H.S.} = -1 = \text{R.H.S.}; \text{L.H.S.} = 1 = \text{R.H.S.} \text{ respectively.}$$

In other cases also we can show similarly that

$$\text{L.H.S. of (1)} = \text{R.H.S. of (1)}$$

Problem 46. Show $\frac{\partial A_\mu}{\partial x_\nu}$ is not a tensor even though A_μ is a covariant tensor of rank one, but the addition of a suitable quantity to $\frac{\partial A_\mu}{\partial x_\nu}$ causes the result to be a tensor. or.

We have $A_\alpha' = \frac{\partial x_\mu}{\partial x_\alpha'} A_\mu$ by §3.4 (2).

Differentiating both sides w.r.t. x_β' we get

$$\begin{aligned} \frac{\partial A_\alpha'}{\partial x_\beta'} &= \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial A_\mu}{\partial x_\beta'} + \frac{\partial^2 x_\mu}{\partial x_\alpha' \partial x_\beta'} A_\mu \\ &= \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial A_\mu}{\partial x_\nu} \frac{\partial x_\nu}{\partial x_\beta'} + \frac{\partial^2 x_\mu}{\partial x_\alpha' \partial x_\beta'} A_\mu \\ &= \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial x_\nu}{\partial x_\beta'} \frac{\partial A_\mu}{\partial x_\nu} + \frac{\partial^2 x_\mu}{\partial x_\alpha' \partial x_\beta'} A_\mu \end{aligned} \quad \dots (1)$$

Unless the second term on the right is zero, $\frac{\partial A_\mu}{\partial x_\nu}$ does not transform as a tensor.

Using (12) of §3.14, we have

$$\frac{\partial^2 x_\mu}{\partial x_\alpha' \partial x_\beta'} = \Gamma_{\beta\alpha}^\gamma \frac{\partial x_\mu}{\partial x_\gamma'} - \Gamma_{\sigma\tau}^\mu \frac{\partial x_\sigma}{\partial x_\alpha'} \frac{\partial x_\tau}{\partial x_\beta'}$$

Substituting in (1), we find

$$\begin{aligned} \frac{\partial A_\alpha'}{\partial x_\beta'} &= \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial A_\mu}{\partial x_\beta'} + \Gamma_{\beta\alpha}^\gamma \frac{\partial x_\mu}{\partial x_\gamma'} A_\mu - \Gamma_{\sigma\tau}^\mu \frac{\partial x_\sigma}{\partial x_\alpha'} \frac{\partial x_\tau}{\partial x_\beta'} A_\mu \\ &= \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial x_\nu}{\partial x_\beta'} \frac{\partial A_\mu}{\partial x_\nu} + \Gamma_{\beta\alpha}^\gamma A_\gamma' - \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial x_\nu}{\partial x_\beta'} \Gamma_{\mu\nu}^\rho A_\rho \end{aligned}$$

(by interchanging the suffixes τ, σ by μ, ν and μ by ρ in the last term)

$$\text{or } \frac{\partial A_\alpha'}{\partial x_\beta'} - \Gamma_{\beta\alpha}^\gamma A_\gamma' = \frac{\partial x_\mu}{\partial x_\alpha'} \frac{\partial x_\nu}{\partial x_\beta'} \left(\frac{\partial A_\mu}{\partial x_\nu} - \Gamma_{\mu\nu}^\rho A_\rho \right) \quad \dots (2)$$

Where $\frac{\partial A_\mu}{\partial x_\nu} - \Gamma_{\mu\nu}^\rho A_\rho$ being a covariant tensor of second rank is known to be the covariant derivative of A_μ w.r.t. x_ν and is written as

$$A_{\mu;\nu} \text{ i.e., } A_{\mu;\nu} = \frac{\partial A_\mu}{\partial x_\nu} - \Gamma_{\mu\nu}^\rho A_\rho.$$

The result (2) follows that addition of a quantity in $\frac{\partial A_\mu}{\partial x_\nu}$ renders it to be a tensor.

Problem 47. If $A_\sigma^{\mu\nu}$ and $B_\sigma^{\mu\nu}$ are tensors, then prove that their sum and difference are also tensors.

$$\text{We have } A_\gamma^{\alpha\beta} = \frac{\partial x_\alpha'}{\partial x_\mu} \frac{\partial x_\beta'}{\partial x_\nu} \frac{\partial x_\sigma}{\partial x_\gamma'} A_\sigma^{\mu\nu} \quad \dots (1)$$

$$\text{and } B_\gamma^{\alpha\beta} = \frac{\partial x_\alpha'}{\partial x_\mu} \frac{\partial x_\beta'}{\partial x_\nu} \frac{\partial x_\sigma}{\partial x_\gamma'} B_\sigma^{\mu\nu} \quad \dots (2)$$

Adding (1) and (2), we get

$$A_\gamma^{\alpha\beta} + B_\gamma^{\alpha\beta} = \frac{\partial x_\alpha'}{\partial x_\mu} \frac{\partial x_\beta'}{\partial x_\nu} \frac{\partial x_\sigma}{\partial x_\gamma'} (A_\sigma^{\mu\nu} + B_\sigma^{\mu\nu})$$

which follows that $A_\sigma^{\mu\nu} + B_\sigma^{\mu\nu}$ is a tensor of the same rank and type as $A_\sigma^{\mu\nu}$ and

$B_\sigma^{\mu\nu}$.

Subtracting (2) from (1) the second result follows.

Problem 48. What is a tensor? Distinguish between a symmetrical and an anti-symmetrical tensor.

If S_{ij} is symmetric and A_{ij} anti-symmetric in the indices, evaluate $A_{ij} S_{ji}$ see §3.5 for the first part. (Vikram, 1967; Rohilkhand, 1976)

Problem 49. Distinguish between symmetric and anti-symmetric tensors. Show that the symmetry properties of a tensor are invariant.

If A^{ij} and A_{ij} are reciprocal symmetric tensors and if u_j are components of a covariant tensor of rank one, then show that

$$A_{ij} u^i u^j = A^{ij} u_i u_j \text{ where } u^i = A^{i\alpha} u_\alpha$$

Problem 50. Discuss the application of tensor analysis to the dynamics of a particle. (Agra, 1971)

Problem 51. Explain what is meant by the rank of a tensor. Show that multiplication of tensors results in addition of their ranks and contraction reduces the rank by two.

$$\text{Given } T = \frac{1}{2} M g_{mn} \dot{x}^m \dot{x}^n \text{ prove that}$$

$$\dot{x}^i = \frac{1}{M} g^{in} \left(\frac{\partial T}{\partial \dot{x}^n} \right), g_{mn} \text{ being the metric tensor. (Agra, 1972)}$$

Problem 52. Define the intrinsic and covariant derivatives of a contravariant vector. Use the expression for the intrinsic derivative to obtain the components of the acceleration vector for the metric $ds^2 = dr^2 + r^2 ds$. (Agra, 1973)

Problem 53. The length ds of a line element in a two dimensional surface $\phi = \phi$ is given by $ds^2 = R^2 d\theta^2 + R^2 \sin^2 \theta d\phi^2$, where R is a constant. Find all components of the metric tensor g_{ab} and the Christoffel symbols of first kind for this surface. (Agra, 1974)

See Problem 25 and 26 (a).

Problem 54. Show why the introduction of an affine connection is necessary in curved space. Find Expression for the affine connection in terms of the metric tensor? (Agra, 1975)

Hint. We should note that in case of cartesian coordinates, both kinds of Christoffel 3-index symbols vanish. Also the concept of parallel displacement being independent of the existence of a metric tensor, a space with a law of parallel displacement is known as an *affinely connected space* and $\Gamma_{\mu\nu}^\lambda$ the components of an affine connection.

Problem 55. If A_{ij} is an anti-symmetric tensor and S_{ij} is symmetric, find whether or not any one of the following tensors is anti-symmetric, or symmetric:

- (i) $A_{ij} A_{ik}$ (ii) $A_{ij} S_{ik}$ (iii) $S_{ij} A_{ik}$ (iv) $S_{ij} S_{ik}$ (v) $A_{ij} S_{ik} + S_{ij} A_{ik}$
 (vi) $A_{ij} S_{ik} - S_{ij} A_{ik}$ (vii) $A_{im} A_{mn} A_{nk}$ (viii) $A_{im} S_{mn} A_{nk}$ (ix) $S_{im} A_{mn} S_{nk}$
 (x) $S_{im} S_{mn} S_{nk}$. (Agra, 1976)

CHAPTER 4

GROUP THEORY

4.1. INTRODUCTION TO SETS, MAPPINGS AND BINARY OPERATIONS

Set. A set is a collection of objects of any sort, having some properties in common, e.g. the set of all natural numbers.

The objects comprising the set are known as its *elements* or *members*.

A set is *finite* or *infinite* according as the number of its elements is finite or infinite.

The elements of a set must be *distinct* and *distinguishable*. Here *distinct* means that no element of the set is repeated and *distinguishable* means that given any object whatsoever, it is either in the set or not in the set.

The sets are denoted by braces like $\{ \}$, e.g. $\{1, 2\}$ and $\{1, 1, 2\}$ which represent the same set.

Defining property of a set. Using any of the notation $:$, 1 , $\exists$, $\forall$, for *such that*, the defining property of a set is $\{x : P(x)\}$, e.g. a set of even numbers from 2 to 20 may be expressed as

$$\{x : x=2n, n=1, 2, \dots, 10\}$$

Singleton set. A set having a single element is called as singleton set e.g. $\{a\} = \{x : x=a\}$. As another example $\{0\}$ is a singleton set having 0 as the single element.

Null set or void set or empty set. A set having no element is called an empty set and denoted by ϕ , such as $\phi = \{x : x \neq x\}$

Subset. Using the notation $\in$ for 'belong to' and $\Rightarrow$ for 'implies', if there are two sets A and B such that every element of A belongs to B , i.e.

$$a \in A \Rightarrow a \in B$$

then A is called a subset of B or said to be contained in B and denoted by $A \subset B$ or $B \supset A$ (i.e. B contains A).

Here A is subset of B and B is superset of A . e.g. $\{1, 3\}$ is subset of $\{1, 2, 3\}$ but $\{1, 2, 3\}$ is superset of $\{1, 3\}$.

Equal sets. Two sets A and B are said to be equal if

$$A \subset B \text{ and } B \subset A$$

e.g. $A = \{a, b, c, d\}$ and $B = \{b, c, a, d\}$ are equal.

The Negations $a \in A$, $A \subset B$ and $A = B$ are $a \notin A$, $a \notin B$ and $A \neq B$ respectively.

Axiom of extension. Two sets A and B are equal if and only if they have the same number of elements.

Axiom of specification. If A is a set and $S(x)$ is a condition or statement, then there corresponds a set B whose elements are exactly those elements x of A for which $S(x)$ is true i.e.

$$B = \{x \in A : S(x)\}$$

Axiom of pairing. If a and b are two objects (sets), then there exists a set A such that $a \in A$ and $b \in A$ i.e. $\{x \in A : x=a \text{ or } x=b\}$

Normal and abnormal sets. If a set contains itself as one of its elements, it is said to be an abnormal set, otherwise it is a normal set.

Equivalent set. Two sets A and B are said to be equivalent and denoted by $A \sim B$ if the number of elements of A is equal to the number of elements of B . Evidently two equal sets are equivalent but the converse is not true.

Proper subset. A set A is said to be the proper subset of B and denoted by $A \subset B$ (some authors use $\subseteq$ for a subset and $\subset$ for a proper subset), if every element of A is an element of B and there is atleast one element of B which is not the element of A so that $A \neq B$ e.g. $\{1, 3, 5\}$ is a subset of $\{1, 3, 5\}$ but it is not its proper subset while $\{1, 3, 5\}$ is a proper subset of $\{1, 3, 5, 7\}$.

Axiom of power set. The power set of a set A is the family or class of all the subsets of A and denoted by $P(A)$ e.g. if $A = \{1, 2, 3\}$ then

$$P(A) = \{\phi, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$$

Obviously when A consists of 3 elements, $P(A)$ consists of 2^3 elements. In general if A consists of m elements, $P(A)$ will consist of 2^m elements. The power set of A is also denoted by 2^A .

Operations on Sets.


Union. The union of two sets A and B denoted by $A \cup B$ and read as ' A union B ' is the set of all objects which are members of A or B (or both)

$$\text{i.e. } A \cup B = \{x \mid x \in A \text{ or } x \in B\}$$

The word 'or' used here gives the inclusive sense and/or.

e.g. if $A = \{1, 2, 3\}$, $B = \{4, 5, 6\}$, then $A \cup B = \{1, 2, 3, 4, 5, 6\}$

The union of n sets $A_1, A_2, \dots, A_n$ is defined as



Venn diagram
Fig. 4.1




Fig. 4.2

$$A_1 \cup A_2 \cup \dots \cup A_n = \bigcup_{i=1}^n A_i = \{x : x \in A_i \text{ for some } i \text{ in the range } i=1 \text{ to } i=n\}$$

Intersection. The intersection of two sets A and B denoted by $A \cap B$ and read as ' A intersection B ' is the set of all objects which are members of both A and of B i.e.

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\}$$

e.g. if $A = \{1, 2, 3, 4\}$, $B = \{2, 4, 6, 8\}$, then $A \cap B = \{2, 4\}$

The intersection of n sets $A_1, A_2, \dots, A_n$ is defined as

$$A_1 \cap A_2 \cap \dots \cap A_n = \bigcap_{i=1}^n A_i = \{x : x \in A_i \text{ for } 1 \leq i \leq n\}$$

Family or class or collection of sets. If a set consists of elements which are sets themselves, then such a set is called as 'family or class' of sets e.g. a set $A^* = \{\{1\}, \{1, 2\}, \{1, 2, 3\}\}$.

Indexed family of sets. For a given index set Λ , a collection of sets such that to each member of Λ there corresponds a member of the collection of the sets, is known as indexed family of sets and written as

$$A^* = \{A_\alpha : \alpha \in \Lambda\}$$

where $\alpha \in \Lambda$ is an index and A_α denote the indexed sets.

The arbitrary union of sets $\{A_\alpha : \alpha \in \Lambda\}$ is given by $\bigcup \{A_\alpha : \alpha \in \Lambda\} = \{x : x \in A_\alpha \text{ for at least one } \alpha \in \Lambda\}$

If $\Lambda = \phi$, then $\bigcup \{A_\alpha : \alpha \in \phi\} = \phi$.

The arbitrary Intersection of sets $\{A_\alpha : \alpha \in \Lambda\}$ is given by

$\bigcap \{A_\alpha : \alpha \in \Lambda\} = \{x : x \in A_\alpha \forall \alpha \in \Lambda\}$, $\forall$ is the notation for 'for every'.

If $\Lambda = \phi$ then $\bigcap \{A_\alpha : \alpha \in \phi\} = U$.

Mutually exclusive or disjoint sets. If there are two sets A and B such that $A \cap B = \phi$, then A and B are said to be disjoint e.g., if $A = \{1, 2\}$, $B = \{5, 6\}$ then $A \cap B = \phi$.

Universal set. All the sets under consideration are assumed to be the subsets of some fixed set called as the universal set and denoted by U .

Complementary set. The complement of a set A is denoted by A' and is defined by the set of all members of the universal set U , which are not members of A i.e. $A' = \{x : x \in U \text{ and } x \notin A\}$

e.g. if $A = \{x : x < 3\}$ then $A' = \{x : x > 3\}$

It is notable that $A \cup A' = U$; $U' = \phi$; $\phi' = U$; $A \cap A' = \phi$ and $(A')' = A$.

Important properties of operations on sets

(1) Difference operation. If A and B are two sets, then the set consisting of elements which belong to A but not to B is said to be the difference of sets A and B and is denoted by $A - B$, i.e.

$$A - B = \{x \mid x \in A \text{ and } x \notin B\}.$$

For example, if $A = \{1, 2, 3, 4\}$ and $B = \{1, 4, 7\}$, then

$$A - B = \{2, 3\} \text{ and } B - A = \{7\}.$$

It is to be noted that $A - B \neq B - A$; $A - A = \phi$, $A - \phi = A$ and $A - B \subset A$.

It may also be shown that $A - B = A \cap B'$; $(A - B) \cap B = \phi$ and $(A - B) \cup A = A$.

The symmetric difference of A and B denoted by $A \Delta B$ is defined as $A \Delta B = (A - B) \cup (B - A)$.

(2) Commutative laws

$$(i) A \cup B = B \cup A,$$

$$(ii) A \cap B = B \cap A.$$

These can be proved as follows :

We have $x \in A \cup B \Rightarrow x \in A$ or $x \in B$

$$\Rightarrow x \in B \text{ or } x \in A$$

$$\Rightarrow x \in B \cup A.$$

Consequently $A \cup B \subset B \cup A$ and also $B \cup A \subset A \cup B$

(in a similar manner).

$$\therefore A \cup B = B \cup A.$$

Similarly, if $x \in A \cap B \Rightarrow x \in A$ and $x \in B$

$$\Rightarrow x \in B \text{ and } x \in A$$

$$\Rightarrow x \in B \cap A,$$

so that $A \cap B \subset B \cap A$ and similarly $B \cap A \subset A \cap B$.

$$\therefore A \cap B = B \cap A.$$

It is easy to verify that $A \cap B \subset A$ and $A \cap B \subset B$.

(3) Associative laws

$$(i) A \cup (B \cap C) = (A \cup B) \cap C,$$

$$(ii) A \cap (B \cup C) = (A \cap B) \cup C.$$

We have $x \in (A \cup B) \cap C \Rightarrow x \in (A \cup B)$ or $x \in C$

$$\Rightarrow x \in A \text{ or } x \in B \text{ or } x \in C$$

$$\Rightarrow x \in A \text{ or } x \in (B \cup C)$$

$$\Rightarrow x \in A \cup (B \cup C),$$

so that $(A \cup B) \cap C \subset A \cup (B \cap C)$

and similarly $A \cup (B \cap C) \subset (A \cup B) \cap C$

$$\therefore A \cup (B \cap C) = (A \cup B) \cap C.$$

Again $x \in (A \cap B) \cup C \Rightarrow x \in A \cap B$ and $x \in C$

$$\Rightarrow (x \in A \text{ and } x \in B) \text{ and } x \in C$$

$$\Rightarrow x \in A \text{ and } (x \in B \text{ and } x \in C)$$

$$\Rightarrow x \in A \text{ and } x \in B \cap C$$

$$\Rightarrow x \in A \cap (B \cap C),$$

so that $(A \cap B) \cup C \subset A \cap (B \cup C)$

and similarly $A \cap (B \cup C) \subset (A \cap B) \cup C$.

$$\therefore A \cap (B \cup C) = (A \cap B) \cup C.$$

(4) Idempotent laws

$$(i) A \cup A = A,$$

$$(ii) A \cap A = A.$$

Here $x \in A \cup A \Rightarrow x \in A$ or $x \in A$.

$$\Rightarrow x \in A.$$

$\therefore A \cup A \subset A$ and similarly $A \subset A \cup A$, so that

$$A \cup A = A.$$

In a similar way it may be shown that $A \cap A = A$.

It is easy to verify that $A \cap \phi = \phi$ and $A \cup U = A$

(5) Distributive laws

$$(i) A \cap (B \cup C) = (A \cap B) \cup (A \cap C),$$

$$(ii) A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

We have, $x \in A \cap (B \cup C) \Rightarrow x \in A$ and $x \in (B \cup C)$
 $\Rightarrow x \in A$ and $(x \in B \text{ or } x \in C)$
 $\Rightarrow (x \in A \text{ and } x \in B)$
 and $(x \in A \text{ and } x \in C)$
 $\Rightarrow x \in A \cap B$ or $x \in A \cap C$
 $\Rightarrow x \in (A \cap B) \cup (A \cap C).$

Consequently $A \cap (B \cup C) \subset (A \cap B) \cup (A \cap C)$
 and similarly $(A \cap B) \cup (A \cap C) \subset A \cap (B \cup C),$
 so that $A \cap (B \cup C) = (A \cap B) \cup (A \cap C).$

Again, $x \in A \cup (B \cap C) \Rightarrow x \in A$ or $x \in B \cap C$
 $\Rightarrow x \in A$ or $(x \in B \text{ and } x \in C)$
 $\Rightarrow (x \in A \text{ or } x \in B)$
 and $(x \in A \text{ or } x \in C)$
 $\Rightarrow x \in A \cup B$ and $x \in A \cup C$
 $\Rightarrow x \in (A \cup B) \cap (A \cup C).$

Consequently $A \cup (B \cap C) \subset (A \cup B) \cap (A \cup C)$
 and similarly $(A \cup B) \cap (A \cup C) \subset A \cup (B \cap C),$
 so that $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

(6) De Morgan laws

$$(i) A - (B \cup C) = (A - B) \cap (A - C),$$

$$(ii) A - (B \cap C) = (A - B) \cup (A - C).$$

We have $A - (B \cup C) = \{x \mid x \in A \text{ and } x \notin B \cup C\}$
 $= \{x \mid x \in A \text{ and } (x \notin B \text{ and } x \notin C)\}$
 $= \{x \mid (x \in A \text{ and } x \notin B)$
 and $(x \in A \text{ and } x \notin C)\}$
 $= \{x \mid x \in (A - B) \text{ and } x \in (A - C)\}$
 $= \{x \mid x \in (A - B) \cap (A - C)\}$

$$\therefore A - (B \cup C) \subset (A - B) \cap (A - C).$$

Similarly it may be shown that $(A - B) \cap (A - C) \subset A - (B \cup C)$

Thus, $A - B \cap C = (A - B) \cup (A - C).$

De Morgan Laws are sometimes expressed as

$$(A \cup B)' = A' \cap B' \text{ and } (A \cap B)' = A' \cup B'.$$

These may be shown as follows :

$$\begin{aligned} (A \cup B)' &= \{x : x \notin A \cup B\} \\ &= \{x : x \notin A \text{ and } x \notin B\} \\ &= \{x : x \in A' \text{ and } x \in B'\} \end{aligned}$$

$$= \{x : x \in A' \cap B'\}.$$

$$(A \cup B)' \subset A' \cap B'$$

$$\text{Similarly } A' \cap B' = \{x : x \in A' \text{ and } x \in B'\}$$

$$= \{x : x \notin A \text{ and } x \notin B\}$$

$$= \{x : x \in (A \cup B)'\}$$

$$\therefore A' \cap B' \subset (A \cup B)'$$

$$\text{Hence } (A \cup B)' = A' \cap B'.$$

Similar procedure will show that $(A \cap B)' = A' \cup B'$

Cartesian product of two sets. The product of two sets A and B is the set of all distinct ordered pairs (a, b) where $a \in A$, and $b \in B$ and denoted by $A \times B$ (read as A cross B) i.e.

$$A \times B = \{(a, b) : a \in A, b \in B\}$$

e.g. if $A = \{1, 2\}$ and $B = \{p, q, r\}$, then

$$A \times B = \{(1, p), (1, q), (1, r), (2, p), (2, q), (2, r)\}$$

and

$$B \times A = \{(p, 1), (p, 2), (q, 1), (q, 2), (r, 1), (r, 2)\}$$

It is clear that $A \times B \neq B \times A$

If either A or B is a null set, then $A \times B = \phi$.

If the set A has m elements and B has n elements, then $A \times B$ or $B \times A$ has mn elements (ordered pairs).

The product of n sets $A_1, A_2, \dots, A_n$ is the set of all distinct ordered n -tuples $(a_1, a_2, \dots, a_n)$ where $a_1 \in A_1, a_2 \in A_2, \dots, a_n \in A_n$ and is defined as

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_1 \in A_1, a_2 \in A_2, \dots, a_n \in A_n\}$$

Function or mapping

Let there be two non-empty sets X and Y and there is some rule or correspondence which assigns to each element $x \in X$, a unique element $y \in Y$, then this rule or correspondence is said to be a *mapping* or a *function* and denoted by f i.e. $f : X \rightarrow Y$

and read as ' f is a function of X to Y ' or f is a mapping of X to Y '.

The set X is called the *domain* of the given function f and the set Y of all the values assumed by it is called its *Range* or *Image set*. Also Y is called the *co-domain* of f .

y is sometimes known as *image* of x and written as $y = f(x)$. Here $f(x)$ is read as '*image of x under the rule f* ' or simply ' *f of x* '. The rule f is also known as *mapping* or *transformation* or *operator* and x is also known as *preimage* of y .




Fig. 4.3

A function whose range has a single element is said to be *constant function*.

Diagrammatical representation of $y=f(x)$ with a rule f defined by $x \rightarrow f(x)$ is shown in Fig. 4.3.

If $y=x^2$, then the rule f is $x \rightarrow x^2$

which is shown in Fig. 4.4 for positive integral values of x .




Fig. 4.4

Functions defined as sets of ordered pairs. Given two non-empty sets X and Y , a function f from X to Y is a subset of $X \times Y$ provided

(i) $\forall x \in X, (x, y) \in f$ for some $y \in Y$ i.e. $\exists$ (there exists) a rule f so that every element of X has image.

(ii) $(x, y) \in f$ and $(x, y') \in f \Rightarrow y = y'$ i.e. the image is unique.

The *graph* of f is defined as the subset of $X \times Y$ given by $\{(x, f(x)) : x \in X\}$ and the *range* of f as the set of all images under f given by $f[X] = \{y \in Y : y = f(x) \text{ for some } x \in X\} = \{f(x) : x \in X\}$.

In case $A \subset X$ then the set $\{f(x) : x \in A\}$ is known as the *image* of A under f and denoted by $f[A]$. Also if $B \subset Y$, then the set $\{x \in X : f(x) \in B\}$ is known as the *inverse image* of B under f and denoted by $f^{-1}[B]$.

Extension and restriction of a function. Given two functions f and g such that f contains the domain of g and $f(x) = g(x) \forall x$ in the domain of g , the function f is said to be the *extension* of g and g is said to be the *restriction* of f .

Real and Complex functions. If range of f consists of real numbers, f is said to be a *real function* and if its range consists of complex numbers, f is said to be a *complex function*.

Onto and Into Mappings. If the range is completely filled up, the mapping is said to be *onto* and if the range is not completely filled up then it is *into*. In other words if $\exists$ at least one $y \in Y$ which is not an $f(x)$ for any $x \in X$, then the mapping f is said to be *into* otherwise it is said to be *onto* or *surjective*. The surjective function is also known as a *Surjection* or an *epimorphism*.

One-one and Many-one Mappings. Given two non-empty sets X and Y , if two different elements in X always have different images under the rule f , then f is said to be a *one-one mapping* or an *injection* or *monomorphism* of X into (onto) Y and if the two or more different elements of X have the same image under f , then f is said to be a *many-one mapping* of X into (onto) Y .

Diagrammatical representation of such functions are shown in Figures 4.5, 4.6, 4.7, 4.8.




Fig. 4.5

A function which is both surjective and injective is known as *bijective* i.e. a one-one onto mapping is also known as a *bijection* and a bijection of a set X onto itself is known as *Permutation* of X .




Fig. 4.6

If $f: X \rightarrow Y$, f is one-one if $x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2) \forall x_1, x_2 \in X$. In case f is *into*, the range of f is a proper subset of Y i.e. $f[X] \subset Y$ and $f[X] \neq Y$.




Fig. 4.7




Fig. 4.8

In case f is *onto*, the range of f is equal to Y i.e. $f[X] = Y$.

Inverse mapping. Let f represent a function (mapping) which is both *onto* and *one-one* defined as $f: X \rightarrow Y$, then its inverse mapping $f^{-1}: Y \rightarrow X$ is defined as below:

$\forall y \in Y$, if we find the unique element $x \in X$ s.t. $f(x) = y$ then x is defined to be $f^{-1}(y)$ i.e. $f^{-1}(y) = \{x : x \in X, f(x) = y\}$ which follows that $f^{-1}(y)$ is always a subset of X .

Diagrammatical representation of an inverse mapping is shown in Fig. 4.9.

One-one onto mapping is often called as *one to one correspondence*. Thus if f is a one to one correspondence between X and Y , then f^{-1} is a one to one correspondence between Y and X .




Fig. 4.9

Identity mapping. Given a non-empty set X , the identity mapping I_X on X is the mapping of X onto itself i.e. $I_X : X \rightarrow X$ and defined as $I_X(x) = x \quad \forall x \in X$

Equal mappings. Let there be two functions f and g defined as $f : X \rightarrow Y$ and $g : X \rightarrow Y$

then the functions f and g are equal iff $f(x) = g(x) \quad \forall x \in X$.

Product or Composite mapping of two mappings $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ is defined to be a mapping, $g \circ f : X \rightarrow Z$ and given by $(g \circ f)(x) = g[f(x)] \quad \forall x \in X$




Fig. 4.10

Diagrammatical representation is shown in Fig. 4.10.

Binary operation. Binary operation over a set X is a mapping from $X \times X \rightarrow X$, i.e. if f denotes a binary operation over a set X , then $f : X \times X \rightarrow X$.

Binary relation or simply relation

A binary relation in a set S is a mathematical symbol denoted by R s.t. for each ordered pair $(x, y) \in S$, the statement xRy is true or false in the sense that xRy asserts that x is related by R to y and

$\neg xRy$, the negation of it i.e. x is not related by R to y , e.g. if S be the set of all integers and R be a relation $<$ (less than) defined on S , then $5 < 8$ is true for $(5, 8) \in S$ i.e. $5R_8$ is true while $8R_5$ is false as $8 \nless 5$ which is denoted by $\neg 8R_5$.

In other words if S is a set s.t. the Cartesian product $S \times S$ involves the same set, then it is called a relation R on S . In fact the relation R on the set S is a subset of $S \times S$ i.e. $R \subset S \times S$.

Thus if $(x, y) \in R$, $(x, y) \in S$ then we have xRy .

In general a relation R from A to B between two sets A and B is a subset of $A \times B$ i.e. $R \subset A \times B$.

If there are m elements in A , n in B then there will be mn element in $A \times B$ and so there will be different mn relations from A to B .

Domain and range of a relation. If A and B are two sets and R is a relation from A to B , then the *domain* of R denoted by $\text{Dom}(R)$ is the set of first coordinates of all the ordered pairs in R and the *range* of R denoted $\text{Ran}(R)$ is the set of second coordinates of all the ordered pairs in R . Thus

$$\text{Dom}(R) = \{x : (x, y) \in R \text{ for some } y \in B\}$$

$$\text{Ran}(R) = \{y : (x, y) \in R \text{ for some } x \in A\}$$

e.g. if R be a relation in $\mathbb{Z}$ the set of natural numbers s.t.

$$R = \{(x, y) \in \mathbb{Z} \times \mathbb{Z} : x + 2y = 10\}$$

then $\text{Dom}(R) = \{2, 4, 6, 8\} \because 2 + 2 \cdot 4 = 10 = 4 + 2 \cdot 3 = 6 + 2 \cdot 2 = 8 + 2 \cdot 1$

$$\text{Ran}(R) = \{4, 3, 2, 1\}$$

Identity or diagonal relation. A relation R on a set A is known as identity relation or diagonal relation iff $xRy \Rightarrow x = y \quad \forall x, y \in A$.

Composite relation. If R be a relation from A to B and S a relation from B to C , then composite relation from A to C is denoted by SoR and is defined as the set of all ordered pairs $(x, z) \in SoR$ iff $\exists y \in B$ s.t. xRy and ySz i.e. $(x, y) \in R$ and $(y, z) \in S$.

In other words $SoR \subset \text{Dom}(R) \times \text{Ran}(R)$

Universal relation in a set. If A is any set and R is the set $A \times A$ then R is said to be the universal relation on A .

Empty or void relation. If A is a set, then every subset of $A \times A$ is a relation on A . Since the null set ϕ is the subset of all the sets and so is of the set $A \times A$, therefore ϕ is also a relation on A . Such a relation is said to be the Empty or Void relation on A .

Inverse relation. If R be a relation from a set A to another set B , then the inverse relation of R denoted by R^{-1} is defined as the inverse relation i.e. R^{-1} from B to A iff

$$R^{-1} = \{y, x) : (x, y) \in R\}$$

clearly $\text{Dom}(R^{-1}) = \text{Ran}(R)$

$$\text{Ran}(R^{-1}) = \text{Dom}(R)$$

So that $xRy \leftrightarrow yR_x^{-1}$

If $A = B$, R and R^{-1} both are the relations on A .

It is easy to verify that $(R^{-1})^{-1} = R$.

Types of Relations

I. Reflexive. A relation R on a set A is reflexive iff each member of A is R -related to itself i.e. $\forall x \in A, xRx$ or in other-words $(x, x) \in R \forall x \in A$. Evidently a relation R on A is reflexive iff $\Delta_A \subset R$, Δ_A being identity relation. e.g. if A be the set of lines in a plane and R a relation 'parallel to' then any line $x \in A$ is parallel to itself i.e. $\forall x \in A, xRx$ and so R is reflexive.

II. Symmetric. A relation R on a set A is symmetric iff $xRy \Rightarrow yRx$; $x, y \in A$

$$\text{i.e. } (x, y) \in R \Rightarrow (y, x) \in R$$

Evidently a relation R on A is symmetric iff $R^{-1} = R$.

e.g. $\forall x, y, xRy = '(x-y) \text{ is even numbers}'$ is symmetric since $y-x$ is also even, when $(x-y)$ is even.

III. Transitive. A relation R on a set A is transitive iff

$$xRy \text{ and } yRz \Rightarrow xRz; x, y, z \in A$$

$$\text{i.e. } (x, z) \in R \text{ and } (y, z) \in R \Rightarrow (x, y) \in R$$

Evidently a relation R on A is transitive iff $R \circ R \subset R$

e.g. the relation $x < y$ is transitive since if $x < y$ and $y < z$ then $x < z$.

IV. Anti-symmetric. A relation R on a set A is anti-symmetric iff we never have xRy and yRx both; $x, y \in A$ except when $x=y$

$$\text{i.e. } xRy \text{ and } yRx \Rightarrow x=y; x, y \in A$$

$$\text{or } (x, y) \in R \text{ and } (y, x) \in R \Rightarrow x=y; x, y \in A$$

Evidently a relation R on A is anti-symmetric iff $R \cap R^{-1} \subset \Delta_A$, Δ_A being identity relation or in other words $R \cap R^{-1} = \phi$.

e.g. $\forall x, y, xRy = 'x \text{ divides } y'$ in $\mathbb{Z}$ (set of natural numbers) is anti-symmetric since ' x divides y ' and ' y divides x ' $\Rightarrow x=y$; $x, y \in \mathbb{Z}$

$$\text{i.e. } xRy \text{ and } yRx \Rightarrow x=y; x, y \in \mathbb{Z}$$

Equivalence relation. A relation R over a set S is said to be an equivalence relation if it satisfies the following properties:

(i) Reflexivity, i.e. $\forall x \in S, xRx$ or $x \sim x \forall x$ ($\sim$ called wobble)

(ii) Symmetry, i.e. $xRy \Rightarrow yRx$ or $x \sim y \Rightarrow y \sim x$

(iii) Transitivity, i.e. xRy and $yRz \Rightarrow xRz$ or $x \sim y$ and $y \sim z \Rightarrow x \sim z$.

Equivalence set (or class). Let R be an equivalence relation in a non-empty set S and let x be an element of S i.e. $x \in S$; then the elements $y \in S$ satisfying xRy constitute a subset of S , known as equivalence set of x w.r.t. R , i.e.

$$S_x \text{ or } \bar{x} \text{ or } [x] = \{y : y \in S \text{ and } xRy\}$$

The equivalence set has the following properties.

(i) If $z \in [x]$ then $[z] = [x]$

(ii) $[x] = [z]$ iff xRz

(iii) If $[x] \cap [z] \neq \phi$, then $[x] = [z]$

Partition set. Given a non-empty set S a set $P = \{X, Y, Z, \dots\}$ of non-empty subsets of S is called a partition of S , provided

$$(i) X \cup Y \cup Z \cup \dots = S$$

(ii) The intersection of every pair of distinct subsets of $S \in P$ is the null set e.g. if $X, Y \in P$ then either $X=Y$ or $X \cap Y = \phi$.

e.g. if $S = \{1, 2, 3, 4, 5\}$ then $\{1, 3, 5\}$, $\{2, 4\}$, and $\{1, 2, 3\}$, $\{4, 5\}$ are two different partitions of S .

Quotient set. If R be an equivalence relation defined on a non-empty set S , then the set of mutually exclusive sets in which S is partitioned w.r.t. the equivalence relation R , is called as quotient set of S for the equivalence relation R and is denoted by $\bar{S}$ or S/R .

e.g. the set $\mathbb{I}$ of all integers for the equivalence relation modulo 5 is the set $\mathbb{I}/R = \{[0], [1], [2], [3], [4]\}$.

Note. Two integers p and q are said to be congruent modulo m denoted by $p \equiv q \pmod{m}$ if $p-q$ is exactly divisible by m i.e. $(p-q)$ is an integral multiple of m .

A note on binary operations. A binary operation (or simply composition) usually denoted by $\circ$ (or sometimes by $\cdot$, $*$, $\oplus$ etc.) is used to combine two elements of a set in order to produce another element of the set. In other words a binary operation in a set S is a function $f: S \times S \rightarrow S$. e.g. the binary operation 'addition' on any two elements of $\mathbb{I}$ (set of integers) gives another integer belonging to $\mathbb{I}$.

(i) Such an operation is commutative if $xoy = yox \forall x, y \in S$

(ii) It is associative if $xo(yoz) = (xoy)oz \forall x, y, z \in S$

(iii) It is distributive w.r.t. another binary operation $\oplus$ if $xo(y \oplus z) = (xoy) \oplus (xoz)$ (left distributive)

and $(y \oplus z)ox = (yox) \oplus (zox)$ (right distributive) $\forall x, y, z \in S$.

(iv) It is said to have identity element if $\exists e \in S$ s.t. $xoe = eox = x \forall x \in S$

(v) It is invertible if corresponding to an $x \in S \exists a y \in S$ s.t. $xoy = e = yox$.

i.e. y is the inverse of x w.r.t. ' $\circ$ ' and vice versa.

(vi) It satisfies cancellation law if $\nexists x, y, z \in S$

$$xoy = xoz \Rightarrow y = z \quad (\text{left cancellation law})$$

$$yox = zox \Rightarrow y = z \quad (\text{right cancellation law})$$

e.g. consider two binary operations $\circ$ and $\cdot$ defined on $\mathbb{I}$ (set of all integers) such that

$$xoy = x + 2y \text{ and } x \cdot y = 2xy \quad \forall x, y \in \mathbb{I}$$

Then we can verify the above laws on operations ' $\circ$ ' and ' $\cdot$ ' as follows:

$$\begin{aligned} (i) \text{ We have } xoy &= x + 2y \quad \forall x, y \in \mathbb{I} \\ &= 2y + x \\ &= yox \end{aligned}$$

$$\begin{aligned} \text{and } x \cdot y &= 2xy = 2yx \quad \forall x, y \in \mathbb{I} \\ &= y \cdot x \end{aligned}$$

i.e. commutative law holds for the operation $\circ$ and

$$(ii) \quad \begin{aligned} xo(yoz) &= xo(y+2z) \quad \forall x, y, z \in I \\ &= x+2(y+2z) = x+2y+4z \end{aligned}$$

$$\text{and} \quad \begin{aligned} (xoy) oz &= (x+2y) oz \quad \forall x, y, z \in I \\ &= x+2y+2z \end{aligned}$$

So that $xo(yoz) \neq (xoy) oz$ i.e. associative law does not hold for 'o'.

$$\text{Also} \quad \begin{aligned} x \cdot (y \cdot z) &= x \cdot (2yz) \quad \forall x, y, z \in I \\ &= 2x \cdot 2yz = 4xyz \end{aligned}$$

$$\text{and} \quad (x \cdot y) \cdot z = (2xy) \cdot z = 4xyz$$

So that $x \cdot (y \cdot z) = (x \cdot y) \cdot z$ i.e. '·' is associative.

$$(iii) \quad \begin{aligned} xo(y \cdot z) &= xo(2yz) \quad \forall x, y, z \in I \\ &= x+4yz \end{aligned}$$

$$\begin{aligned} (xoy) \cdot (xoz) &= (x+2y) \cdot (x+2z) = 2(x+2y)(x+2z) \\ &= 2x^2 + 4xz + 4xy + 8yz. \end{aligned}$$

So that $xo(y \cdot z) \neq (xoy) \cdot (xoz)$ i.e. 'o' is not left distributive w.r.t. '·'.

$$\text{Also} \quad \begin{aligned} x \cdot (yoz) &= x \cdot (y+2z) \quad \forall x, y, z \in I \\ &= 2x(y+2z) \\ &= 2xy + 4xz \end{aligned}$$

$$\text{and} \quad \begin{aligned} (x \cdot y)o(x \cdot z) &= (2xy)o(2xz) \\ &= 2xy + 4xz \end{aligned}$$

So that $x \cdot (yoz) = (x \cdot y)o(xoz)$ i.e. '·' is left distributive w.r.t. 'o'.

(iv) Assuming that $\exists$ an identity e , we have

$$xoe = eox = x \quad \forall x \in I$$

$$\text{So} \quad \begin{aligned} xoe &= x \Rightarrow x+2e = x \\ &\Rightarrow e = 0 \end{aligned}$$

$$\text{and} \quad \begin{aligned} eox &= x \Rightarrow e+2x = x \\ &\Rightarrow e = -x \end{aligned}$$

These imply that e is not unique and hence 'o' has no identity element.

Again if $x \cdot e = e \cdot x = x \quad \forall x \in I$, then

$$\begin{aligned} x \cdot e &= x \Rightarrow 2xe = x \\ &\Rightarrow e = \frac{1}{2} \end{aligned}$$

$$\text{and} \quad \begin{aligned} e \cdot x &= x \Rightarrow 2ex = x \\ &\Rightarrow e = \frac{1}{2} \end{aligned}$$

i.e. e is unique and hence '·' has an identity element which is $\frac{1}{2}$.

(v) Since 'o' has no identity element, it is not invertible.

Assuming that '·' has an inverse p , we must have

$$x \cdot p = e = p \cdot x \quad \forall x \in I$$

$$\begin{aligned} \text{Now} \quad x \cdot p &= e \Rightarrow 2xp = e = \frac{1}{2} \\ &\Rightarrow p = \frac{1}{4x} \end{aligned}$$

$$\text{and} \quad p \cdot x = e \Rightarrow 2px = \frac{1}{2} \Rightarrow p = \frac{1}{4x}$$

Hence '·' is invertible and inverse of x is $\frac{1}{4x}$.

(vi) We have $xoy = x+2y$ and $xoz = x+2z \quad \forall x, y, z \in I$

So that $xoy = xoz \Rightarrow x+2y = x+2z \Rightarrow y = z$ i.e. left cancellation law holds for 'o'.

and $x \cdot y = 2xy, x \cdot z = 2xz \quad \forall x, y, z \in I$

So that $x \cdot y = x \cdot z \Rightarrow 2xy = 2xz \Rightarrow y = z$ i.e. left cancellation law holds for '·'.

Also $yox = y+2x$ and $zox = z+2x \quad \forall x, y, z \in I$

So $yox = zox \Rightarrow y+2x = z+2x \Rightarrow y = z$ i.e. right cancellation law holds for 'o'.

and $y \cdot x = 2yz, z \cdot x = 2zx \quad \forall x, y, z \in I$

So $y \cdot x = z \cdot x \Rightarrow 2yz = 2zx$

$\Rightarrow y = z$ i.e. right cancellation law holds for '·'.

Hence cancellation law holds for o and both the binary operations.

4.2. GROUPS

A group is the simplest algebraic structure found in nature wherever symmetry exists.

A group (G, o) is a system consisting of a non-empty set G such as $G = \{a, b, c, \dots\}$ and a binary operation 'o' satisfying the following axioms:

G_1 —(Closure). If $a \in G, b \in G$ then $aob \in G$ or in other words if $a \in G, b \in G$ then $aob = c$ (closure) where $c \in G$.

G_2 —(Associativity). If $a, b, c \in G$ then $ao(boc) = (aob)oc$.

G_3 —(Existence of an identity) If $a \in G$, then $\exists$ an identity element $e \in G$ s.t. $ea = a \quad \forall a \in G$

G_4 —(Existence of an inverse) If $a \in G, \exists$ an inverse $a^{-1} \in G$ s.t. $a^{-1}oa = e$

where $e \in G$, being an identity element.

In addition to these four axioms if a fifth axiom of commutativity namely G_5 is also satisfied, i.e.

G_5 —(Commutation). If $a, b \in G$ then $aob = boa$ so that G_3 and G_4 take the forms $G_3 : eoa = aoe = a$

$$G_4 : eoa^{-1} = a^{-1}oa = e$$

then the group is said to be an Abelian group.

e.g. the set I (of all integers) with the binary operation 'o' taken as additive (+) is a group, for it satisfies all the four axioms

G_1 —if $a, b \in I$, then $a+b \in I$

G_2 —if $a, b, c \in I$ then $a+(b+c) = (a+b)+c$

G_3 —if $a \in I$ then $\exists$ an integer zero (0) such that $0+a = a$

G_4 —if $a \in I$, then $\exists$ an inverse $(-a)$ e.t. $-a+a = 0$ (identity element).

This group is also abelian or commutative as G_5 is also satisfied i.e. $G_5 - a, b \in I, a + b = b + a$.

Finite and Infinite groups. A group (G, o) consisting of a finite number of elements is said to be a *finite group*, e.g. the set $S = \{1, \omega, \omega^2\}$ where $\omega^3 = 1$, is a finite group under multiplication composition.

A group (G, o) consisting of an infinite number of elements is said to be an *infinite group* e.g. the set I (of all integers) is an infinite group under the addition composition.

Order of a group. The number of elements in a finite group is known as the order of the group. The infinite set is said to be of infinite order. As an example the set $\{1, -1\}$ under multiplication composition is a group of order 2.

4.3. ELEMENTARY PROPERTIES OF A GROUP

I. Uniqueness of identity i.e. the identity element of a group (G, o) is unique.

If possible let us assume that e and e' are two identity elements of the group (G, o) , then

$$eoa = aoe = a \quad \forall a \in G \quad \dots (1)$$

$$\text{and} \quad e'oa = aoe' = a \quad \forall a \in G \quad \dots (2)$$

$$\text{Putting } a = e' \text{ in (1), we have} \quad \dots (3)$$

$$eoe' = e'oe = e'$$

$$\text{Also putting } a = e \text{ in (2), we have} \quad \dots (4)$$

$$e'oe = eoe' = e$$

From (3) and (4) it follows that $e' = e$ i.e. there cannot be two identity elements for (G, o) and hence the identity element of a group is unique.

II. Uniqueness of Inverse i.e. in a group (G, o) every element possesses a unique inverse.

If possible let us assume that a' and a^{-1} are two inverses of a . Also let $a \in G$ and e be the identity element of G . Then we have

$$a^{-1}oa = a oa^{-1} = e \quad \dots (1)$$

$$a'oa = a oa' = e \quad \dots (2)$$

$$\text{Post-multiplying (1) by } a' \text{ we get} \quad \dots (3)$$

$$(a^{-1}oa) oa' = (a oa^{-1}) oa' = eoa' = a'$$

$$\text{and pre-multiplying (2) by } a^{-1} \text{ we get} \quad \dots (4)$$

$$a^{-1}o(a'oa) = a^{-1}o(a oa') = a^{-1}oe = a^{-1}$$

$$\text{But group-postulate } G_2 \text{ gives} \quad \dots (4)$$

$$(a^{-1}oa) oa' = a^{-1}o(a oa')$$

$\therefore$ (3) and (4) follow that $a' = a^{-1}$ i.e. the inverse of an element in a group is unique.

Aliter. Taking a^{-1} and a' two inverses of $a \in G$, we have

$$a oa' = a'oa = e \text{ and } a^{-1}oa = a oa^{-1} = e$$

$$\therefore a^{-1} = a^{-1}oe = a^{-1}o(a oa') = (a^{-1}oa) oa' = eoa' = a'$$

which follows the uniqueness of inverse of $a \in G$.

III. Cancellation laws i.e. for any group (G, o) , and $a, b, c \in G$, the following laws hold

$$(i) aob = aoc \Rightarrow b = c \text{ (left cancellation law)}$$

$$(ii) boa = coa \Rightarrow b = c \text{ (right cancellation law)}$$

(i) Taking $a^{-1} \in G$ as the inverse of $a \in G$, we have

$$aob = aoc \Rightarrow a^{-1}o(aob) = a^{-1}o(aoc) \text{ on pre-multiplying by } a^{-1}$$

$$\Rightarrow (a^{-1}oa) ob = (a^{-1}oa) oc, \text{ the composition being associative by } G_2$$

$$\Rightarrow eob = eoc, \text{ since } \exists \text{ an identity element } e \in G \text{ for 'o'}$$

$$\Rightarrow b = c.$$

(ii) Again taking $a^{-1} \in G$ as the inverse of $a \in G$, we have

$$boa = coa \Rightarrow (boa)oa^{-1} = (coa)oa^{-1} \text{ on post-multiplying by } a^{-1}$$

$$\Rightarrow bo(aoa^{-1}) = co(aoa^{-1}) \text{ by } G_2$$

$$\Rightarrow boe = coe \quad \because \exists \text{ an identity element } e \in G \text{ for 'o'}$$

$$\Rightarrow b = c.$$

Note. $aoc = cob \nRightarrow a = b$ unless the group is abelian.

IV. Uniqueness of solutions, i.e. if $a, b \in G$, then the equations $aox = b$ and $yoa = b$ have unique solutions in G .

If a^{-1} be the inverse of $a \in G$, then $a^{-1} \in G$ and $aoa^{-1} = e$ (identity element).

Now $a^{-1} \in G$ and $b \in G \Rightarrow a^{-1}ob \in G$

Putting $x = a^{-1}ob$ in the equation $aox = b$ we get

$$ao(a^{-1}ob) = b$$

$$\text{or} \quad (aoa^{-1})ob = b \text{ by } G_2$$

$$\text{or} \quad eob = b \text{ i.e. } b = b$$

which follows that $x = a^{-1}ob$ is a solution of $aox = b$.

To show that this solution is unique, let us assume that y is an element different from $a^{-1}ob$ in G s.t. it satisfies the equation $aox = b$. Then,

$$aoy = b = eob = (aoa^{-1})ob = ao(a^{-1}ob) \text{ by } G_2.$$

So that left cancellation law yields $y = a^{-1}ob$.

As such $x = y$ i.e. the solution is unique.

Again, $b \in G$ and $a^{-1} \in G \Rightarrow boa^{-1} \in G$

Putting $y = boa^{-1}$ in $yoa = b$, we get

$$(boa^{-1})oa = b \text{ or } bo(a^{-1}oa) = b \text{ by } G_2$$

$$\text{or} \quad boe = b \text{ i.e. } b = b$$

which follows that $y = boa^{-1}$ is a solution of $yoa = b$.

To show that this solution is unique, let us assume that z is an element different from boa^{-1} in G s.t. it satisfies the equation $yoa = b$. Then $zoa = b = boe = bo(a^{-1}oa) = (boa^{-1})oa$ by G_2 .

The right cancellation law gives $z = boa^{-1}$

So that $y = z$ and hence the solution is unique.

Note. The unique solution of $xox=x$ is $x=e$ in group (G, o)

V. Inverse of the inverse is itself i.e. if $a \in G$ then $(a^{-1})^{-1}=a$.

Inverse law gives $(a^{-1})^{-1}oa^{-1}=e$, e being identity element in G .

Postmultiplying by a , we get

$$[(a^{-1})^{-1}oa^{-1}]oa=eo a$$

$$(a^{-1})^{-1}o(a^{-1}oa)=a \text{ by } G_2$$

$$(a^{-1})^{-1}oe=a \text{ by } G_4$$

$$(a^{-1})^{-1}=a \text{ by } G_3$$

which proves the proposition.

VI. Reversal law i.e. if $a, b \in G$ then $(aob)^{-1}=b^{-1}oa^{-1}$.

Let e be the identity element in G .

Now $a \in G \Rightarrow a^{-1} \in G$ and $b \in G \Rightarrow b^{-1} \in G$

$$\begin{aligned} \therefore (b^{-1}oa^{-1})o(aob) &= b^{-1}o[a^{-1}o(aob)] \text{ by } G_2 \\ &= b^{-1}o[(a^{-1}oa)ob] \text{ by } G_2 \\ &= b^{-1}o[ob] \text{ by } G_4 \\ &= b^{-1}ob \text{ by } G_3 \\ &= e. \end{aligned}$$

Hence by definition of inverse element of a group $b^{-1}oa^{-1}$ is the inverse of aob i.e. $(aob)^{-1}=b^{-1}oa^{-1}$.

Note. The result may be generalized for any number of elements

$a_1, a_2, a_3, \dots, a_n \in G$, where we have

$$(a_1oa_2o\dots oa_n)^{-1}=a_n^{-1}oa_{n-1}^{-1}o\dots oa_2^{-1}oa_1^{-1}.$$

4.4. SOME DEFINITIONS

Semi-group. A set S with a binary operation ' o ' is said to be a semi-group if it satisfies the following two axioms.

S_{G_1} —(Closure). $a \in S, b \in S \Rightarrow aob \in S$.

S_{G_2} —(Associativity). If $a, b, c \in S$ then $(aob)oc=ao(boc)$.

THEOREM. A semi-group (G, o) satisfying the following postulates is a group.

(1) G has a left identity e s.t. $eo a=a \forall a \in G$.

(2) Every element a in G has a left inverse a^{-1} in G s.t. $a^{-1}oa=e$.

Since (G, o) is a semi-group, therefore by definition it follows that

(i) (G, o) satisfies the closure law.

(ii) (G, o) satisfies the associative law.

(iii) a^{-1} being the left inverse of a and e the left identity we have

$$\begin{aligned} a^{-1}o(aoe) &= (a^{-1}oa)oe \text{ by (ii)} \\ &= eoe \text{ by postulate (2)} \\ &= e \text{ by postulate (1)} \\ &= a^{-1}oa \text{ by postulate (2)} \end{aligned}$$

So that by left cancellation law, $aoe=a$ which follows that e is also a right identity.

Hence the identity element exists for the composition.

(iv) We have $a^{-1}o(aoa^{-1})=(a^{-1}oa)oa^{-1}$ by (ii)

$$\begin{aligned} \text{or } a^{-1}o(aoa^{-1}) &= eoa^{-1} \text{ by (2)} \\ &= a^{-1} \text{ by (1)} \end{aligned}$$

$$= a^{-1}oe, \text{ since identity element exists for 'o',}$$

$$\therefore aoa^{-1}=e \text{ by left cancellation law.}$$

Which shows that a^{-1} is also a right inverse of a .

Hence every element of G has an inverse.

Since all the four group axioms are satisfied, therefore a semi-group with the given two postulates is a group.

Sub-group. A sub-group of a group (G, o) is any collection of elements of G satisfying the axioms of G . In other words a non-empty subset say H of a group G is said to be the sub-group of G , if the binary operation ' o ' in G induces a binary operation in H and the elements of H obey the group axioms.

In other words a non-empty subset H of a group G is said to be a sub-group of G if it satisfies the following two axioms:

(i) $a, b \in H \Rightarrow aob \in H$,

(ii) $a \in H \Rightarrow a^{-1} \in H$

e.g., the set of even integers is a sub-group of the additive group of integers.

Proper sub-group. A subgroup of a group (G, o) other than G itself and the group consisting of the identity element alone is termed as a proper sub-group of G . e.g. the additive group of integers is a proper sub-group of the additive group of rational numbers.

Improper or trivial subgroups. The group (G, o) itself and the group consisting of identity alone i.e. $(\{e\}, o)$ are known as trivial or improper subgroups of (G, o) .

Order of an element of a group. Let a be an element of a group (G, o) i.e. $a \in G$. Then the order of a is the least positive integer n s.t. $a^n=e$.

In case $\nexists$ such integer, the order of a is said to be zero or infinite. e.g. the order of the element -1 in the multiplicative group $\{1, -1, i, -i\}$ is 2 since $(-1)^2=1$, the identity element. The order of i is 4 since $i^4=1$.

Addition modulo m . (m being an integer). If a and b are two integers and m a positive integer, then 'addition modulo m ' is denoted by $a+_mb$ and defined as $a+_mb=r$, $0 \leq r < m$ where r is the least positive remainder obtained on dividing the sum of a and b by m .

$$\begin{aligned} \text{e.g. } 12+_35 &= 2 \text{ since } 12+5=3(5)+2 \\ \text{and } -5+_410 &= 1 \text{ since } -5+10=4(1)+1 \end{aligned}$$

Multiplication modulo p . (p being a prime). If a and b are two integers and p , a positive integer, then 'multiplication modulo p ' is denoted by $a \times_p b$ and defined as

$a \times_p b = r$, $0 \leq r < p$ where r is the remainder obtained on dividing the ordinary product ab by p .

e.g. $9 \times_6 7 = 3$ since $9 \times 7 = (6) 10 + 3$

and $-7 \times_8 8 = 4$ since $-7 \times 8 = (5) (-12) + 4$

Group table or composition table. It is commonly observed that a 'table' is a convenient way of either defining a binary operation in a finite set S or tabulating the effect of a binary operation in a set S . In forming a table or say a group table we arrange the elements of a group in rows and columns of a square array such that each element of the group occurs once and only once in each row or column. The composition element aob occurs at the intersection of row and column of the elements a and b of the group after the binary operation has been performed. For example consider a set

$S = \{1, 2, 3\}$ and let $\cdot$ be the binary operation in S defined by

$\cdot: (1, 1) \rightarrow 1, (1, 2) \rightarrow 2, (1, 3) \rightarrow 3, (2, 1) \rightarrow 2, (2, 2) \rightarrow 1, (2, 3) \rightarrow 2,$

$(3, 1) \rightarrow 3, (3, 2) \rightarrow 2, (3, 3) \rightarrow 1$

then these operations can be arranged in a table as follows:

It is clear that (i, j) th square ($i, j = 1, 2, 3$) is the intersection of i th row (i.e. row labelled or faced by i) and j th column (i.e. column labelled or faced by j) and in this square we have put the element obtained by the binary operation $\cdot$ on (i, j) such as, $\cdot: (1, 2) \rightarrow 2$

$\cdot$	1	2	3
1	1	2	3
2	2	1	2
3	3	2	1

As an other example if $a, b \in G$ and e be an identity element of the group (G, o) , such that $aoa = b$ and $aob = e$ etc., then the group table is as shown here.

o	e	a	b
e	e	a	b
a	a	b	e
b	b	e	a

Problem 1. Show that three cube roots of unity form an abelian finite group under multiplication.

We have the set $G = \{1, \omega, \omega^2\}$, where $\omega^3 = 1$.

$\cdot$	1	ω	ω^2
1	1	ω	ω^2
ω	ω	ω^2	1
ω^2	ω^2	1	ω

$\therefore \omega^3 = 1$

and $\omega^4 = \omega^3 \cdot \omega = \omega$

The composition table under multiplication is as shown here.

The set G forms an abelian finite group, since it satisfies all the five axioms:

G_1 —since all the elements in group table belong to G , hence closure axiom is true.

G_2 —since multiplication of complex numbers is associative, therefore G_2 is satisfied.

G_3 —since $\exists$ an identity element $1 \in G$, G_3 is satisfied.

G_4 —since the inverses of $1, \omega, \omega^2$ are respectively $1, \omega^2, \omega \in G$, G_4 is satisfied.

G_5 —commutative property is apparently satisfied since $1 \cdot \omega = \omega \cdot 1 = \omega$ etc.

Moreover the set consists of finite number of elements and hence G_1 is an abelian finite group.

Problem 2. Show that the set of all n th roots of unity form a finite abelian group G of order n under ordinary multiplication as composition.

By De Moivre's theorem, n th roots of unity are given by

$$(1)^{1/n} = (\cos 2r\pi + i \sin 2r\pi)^{1/n} \\ = \cos \frac{2r\pi}{n} + i \sin \frac{2r\pi}{n}, \text{ where } r = 0, 1, 2, \dots, n-1$$

So n , n th roots of unity are

$$1, \cos \frac{2\pi}{n} + i \sin \frac{2\pi}{n}, \cos 2 \frac{2\pi}{n} + i \sin \frac{2 \cdot 2\pi}{n}, \dots, \cos \frac{(n-1) \cdot 2\pi}{n} \\ + i \sin \frac{(n-1) \cdot 2\pi}{n}$$

i.e. $1, e^{1 \cdot 2\pi i/n}, e^{2 \cdot 2\pi i/n}, e^{3 \cdot 2\pi i/n}, \dots, e^{(n-1) \cdot 2\pi i/n}$

Now, G_1 is satisfied since the product of any two elements of the set is the element of the set such as if $a = e^{p \cdot 2\pi i/n}, b = e^{q \cdot 2\pi i/n} \in G$,

where $0 \leq p \leq n-1, 0 \leq q \leq n-1$, then $a \cdot b = e^{2\pi i/n(p+q)}$ will belong to G if $p+q \leq n-1$. Let us assume the contrary i.e. $p+q > n-1$ so that $p+q = n+m$ where $m \leq n-2$ since the maximum value of $p+q$ can be $2(n-1)$ i.e. $2n-2$.

$$\therefore a \cdot b = e^{2\pi i/n(n+m)} = e^{2\pi i} e^{2\pi im/n} = e^{2\pi im/n}$$

$$\therefore e^{2\pi i} = \cos 2\pi + i \sin 2\pi = 1$$

which follows that $a \cdot b \in G$ since $m \leq n-2$.

G_2 is satisfied since multiplication of complex numbers is associative.

G_3 is satisfied since there exists an identity element $e^{2\pi i \cdot 0/n} = 1$

G_4 is satisfied since $\exists$ inverse of $e^{2\pi ir/n}$ as $e^{2\pi i(n-r)/n}$ since

$$e^{2\pi ir/n} e^{2\pi i(n-r)/n} = e^{2\pi in/n} = e^{2\pi i} = 1$$

G_5 is also satisfied since the multiplication of complex numbers is commutative.

Moreover the set consists of finite number of elements. Hence (G, o) is a finite abelian group.

Problem 3. Show that the set of matrices $A_\alpha = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$, where α is real, forms a group under multiplication.

Let G be the set of matrices given by $A_\alpha = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$, α being real and $A_\alpha, A_\beta, A_\gamma \in G$. Then,

G_1 is satisfied since the product of any two matrices of the set belongs to the set, as

$$A_\alpha \cdot A_\beta = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \beta & -\sin \beta \\ \sin \beta & \cos \beta \end{bmatrix} = \begin{bmatrix} \cos(\alpha+\beta) & -\sin(\alpha+\beta) \\ \sin(\alpha+\beta) & \cos(\alpha+\beta) \end{bmatrix} \\ = A_{\alpha+\beta}, \alpha+\beta \text{ being real.}$$

$\therefore \alpha, \beta$ are real and so is $\alpha+\beta$, $\therefore A_{\alpha+\beta} \in G$

G_2 is satisfied since $A_\alpha \cdot (A_\beta \cdot A_\gamma) = A_\alpha \cdot A_{\beta+\gamma} = A_{\alpha+\beta+\gamma} = A_{\alpha+\beta} \cdot A_\gamma = (A_\alpha \cdot A_\beta) \cdot A_\gamma$ i.e. the operation is associative.

G_3 is satisfied since $\exists$ an identity element $A_o \in G$, o being real such that

$$A_o \cdot A_\alpha = A_{o+\alpha} = A_\alpha \quad \forall A_\alpha \in G.$$

G_4 is satisfied since $\exists$ an inverse $A_{-\alpha} \in G \quad \forall A_\alpha \in G$, $-\alpha$ being real as α is real, such that

$$A_{-\alpha} \cdot A_\alpha = A_{-\alpha+\alpha} = A_o \text{ (the identity element)}$$

G_5 is also satisfied since $A_\alpha \cdot A_\beta = A_{\alpha+\beta} = A_{\beta+\alpha} = A_\beta \cdot A_\alpha$; α, β being real i.e. the operation is commutative.

Hence the given set of matrices forms an abelian group.

Problem 4. If OX, OY be the two rectangular axes in the cartesian plane and T_α denotes the rotation of the axes through an angle α s.t.

$$T_\alpha : (x, y) \rightarrow (x \cos \alpha + y \sin \alpha, -x \sin \alpha + y \cos \alpha)$$

then show that the set of these rotations w.r.t. the operation 'o' s.t. $T_\beta o T_\alpha$ is the resultant of two such operations, forms a group.

Let $G = \{T_\alpha : (x, y) \rightarrow x \cos \alpha + y \sin \alpha, -x \sin \alpha + y \cos \alpha\}$

G_1 is satisfied since if $T_\alpha \in G, T_\beta \in G$ then $T_\beta o T_\alpha \in G$ as

$$T_\beta o T_\alpha(x, y) = T_\beta [T_\alpha(x, y)] \\ = x \cos(\alpha+\beta) + y \sin(\alpha+\beta), -x \sin(\alpha+\beta) + y \cos(\alpha+\beta) \dots (1) \\ = T_{\alpha+\beta}(x, y) \text{ for any } (x, y) \text{ in the plane}$$

G_2 is satisfied since if $T_\alpha, T_\beta, T_\gamma \in G$, then

$$T_\gamma o (T_\beta o T_\alpha) = T_\gamma [x \cos(\alpha+\beta) + y \sin(\alpha+\beta), -x \sin(\alpha+\beta) + y \cos(\alpha+\beta)] \text{ by (1)} \\ = x \cos(\alpha+\beta+\gamma) + y \sin(\alpha+\beta+\gamma), -x \sin(\alpha+\beta+\gamma) + y \cos(\alpha+\beta+\gamma) \\ = T_{\alpha+\beta+\gamma}(x, y) \text{ etc.}$$

G_3 is satisfied since $\exists$ an identity $T_o \in G$ s.t. $T_o o T_\alpha = T_\alpha \quad \forall T_\alpha \in G$

G_4 is satisfied since $\exists$ an inverse $T_{-\alpha} \in G$ s.t. $T_{-\alpha} o T_\alpha = T_o \quad \forall T_\alpha \in G$

G_5 is satisfied by (1)

Hence the set of rotations form an abelian group.

Problem 5. Prove that residue classes modulo m form a group w.r.t. addition of residue classes.

If S be the set of residue classes modulo m , then we have

$$S = \{[0], [1], [2], \dots, [m-1]\}$$

G_1 is satisfied since if $[r_1], [r_2] \in S$ and $r_1 + r_2 = mk + r$ where r is the least positive remainder when $(r_1 + r_2)$ is divided by m and $0 \leq r < m$, then

$$[r_1] + [r_2] = [r_1 + r_2] = [r] \in S. \quad \therefore r_1 + r_2 \equiv r \pmod{m}$$

i.e. closure axiom is satisfied.

$$\text{or } r_1 + m r_2 = r$$

G_2 is satisfied since if $[r_1], [r_2], [r_3] \in S$ and $r_2 + r_3 = mk + r$ so that $r_1 + r_2 + r_3 = r_1 + mk + r = mk + (r_1 + r) = mk + r'$ (say), r' being least positive remainder when $(mk + r_1 + r)$ i.e. $r_1 + r_2 + r_3$ i.e. $r_1 + r$ is divided by m ,

$$\text{then } [r_1] + ([r_2] + [r_3]) = [r_1] + [r_2 + r_3]$$

$$= [r_1] + [r] \quad \therefore r_1 + r_2 \equiv r \pmod{m}$$

$$= [r_1 + r] = [r'] \quad \therefore r_1 + r \equiv r' \pmod{m}$$

$$= [r_1 + r_2 + r_3]$$

$$= [r_1 + r_2] + [r_3]$$

$$= ([r_1] + [r_2]) + [r_3]$$

i.e. addition of residue classes is associative.

G_3 is satisfied since $\exists$ an additive identity $[o] \in S$ s.t. $[o] + [r] = [r] \quad \forall [r] \in S$

G_4 is satisfied since $\exists$ an additive inverse $[m-r] \in S$ s.t.

$$[r] + [m-r] = [m] = [o] \quad \forall [r] \in S \text{ as } m \equiv o \pmod{m}.$$

Hence the set of residue classes modulo m form a group, w.r.t. addition.

Problem 6. Prove that the non-zero residue classes modulo m (a prime integer) w.r.t. multiplication form a group.

If S be the set of non-zero residue classes modulo m , then we have

$$S = \{[1], [2], [3], \dots, [r], \dots, [m-1]\}, 0 < r \leq m-1.$$

Defining the multiplication of classes $[r_1], [r_2], [r_3] \in S$ by $[r_1] [r_2] = [r_3]$ where $0 < r_1, r_2, r_3 \leq m-1$, we observe that

G_1 is satisfied since r_1, r_2 are prime to m and division of $r_1 r_2$ by m renders a non-zero remainder and so if $[r_1], [r_2] \in S$ then

$$[r_1] [r_2] \text{ i.e. } [r_3] \in S.$$

G_2 is satisfied since if $[r_1], [r_2], [r_3] \in S$ then taking $r_1 r_2 = mk + r$ we have $(r_1 r_2) r_3 = (mk + r) r_3 = mkr_3 + rr_3 = p + r'$ (say) where r' is the least positive remainder when $(r_1 r_2) r_3$ or $r r_3$ is divided by m .

So that

$$([r_1] \cdot [r_2]) [r_3] = [r_1 r_2] [r_3] = [r] [r_3] \quad \therefore r_1 r_2 \equiv r \pmod{m}$$

$$= [r r_3] = [r'] \quad \therefore r r_3 \equiv r' \pmod{m}$$

But $r_1 r_2 r_3 \equiv r' \pmod{m} \therefore [r'] = [r_1 r_2 r_3] = [r_1] [r_2 r_3] = [r_1] ([r_2] [r_3])$ i.e. $([r_1] \cdot [r_2]) \cdot [r_3] = [r_1] ([r_2] \cdot [r_3])$.

thereby showing that the associative law is satisfied.

G_3 is satisfied since $\exists$ an identity $[1] \in S$ s.t. $[1][r]=[r] \forall [r] \in S$

G_4 is satisfied since $\exists$ inverse of each element, as is shown below:

Multiplying each element of S by an element $[r]$, we find

$[1][r], [2][r], \dots, [m-1][r] \dots (1)$

By G_1 (closure axiom) all these $(m-1)$ elements must belong to S . Also all of them should be distinct otherwise if

$[r_1][r]=[r_2][r], [r_1], [r_2] \in S$

then left cancellation law gives $[r_1]=[r_2]$ which contradicts the hypothesis that $[r_1], [r_2]$ are distinct. Hence all the $(m-1)$ elements of (1) must be distinct and they must also be the same elements of S as already defined except that their order may be different. Conclusively in (1) there is one element which is the identity $[1]$. Suppose this identity element is $[r_1][r]=[1]$.

Which shows that $[r_1]$ is the inverse of $[r]$.

But r_1 being arbitrary, the inverse of each element exists. Hence the non-zero residue classes modulo m w.r.t. multiplication form a group.

Problem 7. *If every element of a group (G, o) is its own inverse then show that (G, o) is abelian.*

Given that (G, o) is a group.

$\therefore$ if $a, b \in G$ then $a^{-1}, b^{-1} \in G$

also if $aob \in G$ then $(aob)^{-1} \in G$

But we have $a=a^{-1}$ and $b=b^{-1}$

As such $(aob)=(aob)^{-1}=b^{-1}oa^{-1}=(boa)$

i.e. (G, o) is commutative. Hence (G, o) is abelian.

Problem. 8. *If (G, o) be a group and $a^2=e$ (identity) $\forall a \in G$, then show that the group must be commutative.*

Given that (G, o) is a group and $a^2=aoa=e$ also $aoa^{-1}=e$

$\therefore aoa=e=aoa^{-1}$.

So that left cancellation law gives $a=a^{-1}$

i.e., every element of the group is its own inverse and hence by Problem 7 it follows that the group (G, o) is commutative.

Problem 9. *Show that if a group has 3, 4 or 5 elements, then it is abelian.*

We prove the proposition for 4 elements, similar procedure can be adopted for other two.

Suppose that $G=\{e, a, b, c\}$ is a set forming the group $\{G, o\}$ where e is the identity element.

In case every element of G is its own inverse, the problem reduces

to the problem 7 which has been already discussed. Consider the other case.

Let $a^{-1}=b$. Then the only alternative is that $c^{-1}=c$, so that

$aob=boa=e$ and $coc=e$

Now

$aoc \neq e$ as $c^{-1} \neq a$

$aoc \neq a$ as $c \neq e$

$aoc \neq c$ as $a \neq e$

So the only alternative is that $aoc=b$.

Similar argument will give that $coa=b$

$\therefore aoc=coa$

Also

$boc \neq e$ as $b^{-1} \neq c$

$boc \neq b$ as $c \neq e$

$boc \neq c$ as $b \neq e$

leading that $boc=a$ and similarly $cob=a$

$\therefore boc=cob$

(1), (2) and (3) follow that the group (G, o) is commutative and hence it is abelian. The group table is as shown here.

Problem 10. *Show that any non-commutative group has at least six elements.*

Let (G, o) be a non-commutative group. It will be so if it has at least one pair of non-commuting elements a and b (say).

We shall first show that a set $\{e, a, b, aob, boa\}$ having a, b non-commuting elements i.e. $aob \neq boa$, consists of distinct elements. Taking two at a time, there are ten possibilities leading to a contradiction of $aob \neq boa$:

(i) $e=a \Rightarrow aob=eob=b=boe=boa$.

(ii) $e=b \Rightarrow aob=aob=a=eo a=boa$

(iii) $e=aob \Rightarrow aoe=eo a=(aob)oa=ao(boa)$

i.e. $e=boa$ or $aob=boa$

(iv) $e=boa \Rightarrow eoa=aoe=ao(boa)=(aob)oa$ i.e. $e=aob$ or $boa=aob$

(v) $a=b \Rightarrow aob=aoa=boa$

(vi) $a=aob \Rightarrow e=b$ thereby reducing to (ii)

(vii) $a=boa \Rightarrow e=b$ thereby reducing to (ii)

(viii) $b=aob \Rightarrow e=a$ thereby reducing to (i)

(ix) $b=boa \Rightarrow e=a$ thereby reducing to (i)

(x) $aob=boa$

Hence the elements of the set $\{e, a, b, aob, boa\}$ having (a, b) non-commuting, are all distinct.

We shall now show that at least one of the group elements aoa or $ao\ b oa$ is distinct from these five namely e, a, b, aob, boa .

To show that aoa is different from each element a, b, aob, boa , we see that

(xi) $aoa = a \Rightarrow a = e$ thereby reducing to (i)

(xii) $aoa = b \Rightarrow aob = ao\ (aoa) = (aoa)\ oa = boa$

(xiii) $aoa = aob \Rightarrow a = b$ thereby reducing to (v)

(xiv) $aoa = boa \Rightarrow a = b$ thereby reducing to (v)

These possibilities lead that either $aoa \neq e$ in which case aoa is the sixth distinct element of G or else $aoa = e$

Again we shall show that $ao\ b oa$ is different from each element e, a, b, aob, boa so that it will be the sixth distinct element of G .

Obviously $ao\ (ao\ b oa) = (aoa)\ o\ (boa) = eo\ (boa) = boa$.

Now consider the case

(xv) $ao\ b oa = e \Rightarrow boa = ao\ (ao\ b oa) = aoe = o$ thereby reducing to (vii)

(xvi) $ao\ b oa = a \Rightarrow aob = e$ thereby reducing to (iii)

(xvii) $ao\ b oa = b \Rightarrow aob = ao\ (ao\ b oa) = boa$ when $aoe = e$

(xviii) $ao\ b oa = aob \Rightarrow a = e$ thereby reducing to (i)

(xix) $ao\ b oa = boa \Rightarrow a = e$ thereby reducing to (i)

Conclusively a group with upto 5 elements is essentially abelian but for it to be non-abelian there should be at least six elements.

Problem 11. Show that non-empty semi-group (G, o) forms a group if the equations $ax=b$ and $ya=b$ have unique solutions in $G \forall$ pair of elements $a, b \in G$.

Since $ya=b$ is solvable for any $b \in G$, therefore by taking $b=a$, we find that $ya=a$ has a solution in G . Call this solution as e_1 so that $e_1 a = a$ where a is a fixed element of G .

Let $c \in G$, then $ax=c$ has a solution in G .

Thus $e_1 c = e_1 (ax) = (e_1 a) x = ax = c$

which follows that $e_1 c = c \forall c \in G$ i.e. e_1 is the left identity in G .

As e_1 exists in G , so $ya=e_1$ has a solution in G . Call this solution as a^{-1} . This follows that every element in G has a left inverse relative to the left identity. Hence by the theorem on §4.4, it follows that (G, o) is a group.

Problem 12. Show that a finite-non-empty semi-group (G, o) forms a group if $ab=ac \Rightarrow b=c$ and $ba=ca \Rightarrow b=c \forall a, b, c \in G$.

Consider a set $G = \{a_1, a_2, \dots, a_r, \dots, a_p\}$ consisting of p distinct elements. Take an element a_m and multiply it to all the elements of this group.

$$a_m a_1, a_m a_2, \dots, a_m a_r, \dots, a_m a_p.$$

All these elements will be distinct save possibly arranged in different order. If possible let us assume that

$$a_m a_r = a_m a_p \Rightarrow a_r = a_p$$

which contradicts the hypothesis that a_r and a_p are distinct elements of G . Thus

$G = \{a_m a_1, a_m a_2, \dots, a_m a_r, \dots, a_m a_p\}$ consists of p distinct elements and $a_m a_1$ will be some element say a_r of G i.e.

$$a_m a_1 = a_r \Rightarrow ax = b \text{ has a unique solution in } G$$

Similarly we can show that

$G = \{a_1 a_m, a_2 a_m, \dots, a_r a_m, \dots, a_p a_m\} \Rightarrow ya = b$ has a unique solution in G .

Hence by Problem 11, the semi-group (G, o) under given conditions forms a group.

Problem 13. Show that the set of subsets of a set with the union composition is a semi-group.

If $S_1 = \{A, B, C, \dots\}$ be the set of subsets of a set S , then

S_{G_1} is satisfied since, $A, B \in S$, and $A \subset S, B \subset S \Rightarrow A \cup B \subset S$ and $A, B \in S$, i.e. the closure law is satisfied. $\Rightarrow A \cup B \in S_1$

S_{G_2} is satisfied since if $A, B, C \in S$, then associative property of union yields, $(A \cup B) \cup C = A \cup (B \cup C)$

Hence S_1 is a semi-group (by def. § in 4.4).

Problem 14. Show that the identity of a subgroup of a group is the same as that of the group.

Let (H, o) be a subgroup of the group (G, o) and let e, e' be the identities of (G, o) and (H, o) respectively. Then

$$aoe' = a \forall a \in H$$

This equality will also hold in (G, o) as $a \in H \Rightarrow a \in G$.

Now if b be the inverse of $a \in G$, then we have

$$\begin{aligned} aoe' &= a \Rightarrow bo(aoe') \\ &\Rightarrow (boa) oe' = boa \text{ by } G_2 \text{ for } G \\ &\Rightarrow eoe' = e \quad \quad \quad boa = e \\ &\Rightarrow e' = e. \end{aligned}$$

Problem 15. Show that the inverse of an element of a subgroup of a group is the same as the inverse of the same element regarded as an element of the group.

Let (H, o) be a subgroup of the group (G, o) and let b_1 and b_2 be the inverses of an element a as member of H and G respectively. Also let e and e' be the identities of G and H respectively. Then by Problem 14, $e = e'$,

$$\begin{aligned} \text{Now } aob_1 &= e' = e \Rightarrow b_2 o (aob_1) = b_2 oe \\ &\Rightarrow (b_2 oa) ob_1 = b_2 \text{ by } G_2, G_3 \text{ for } G. \\ &\Rightarrow eob_1 = b_2 \quad \quad \quad \therefore b_2 oa = e \\ &\Rightarrow b_1 = b_2. \end{aligned}$$

Problem 16. Show that the necessary and sufficient conditions for a complex H to be a subgroup (H, o) of a group (G, o) are

(i) $a_1 b \in H \Rightarrow aob \in H \forall a, b$; and (ii) $a \in H \Rightarrow a^{-1} \in H \forall a$
(Rohilkhand, 1976)

The conditions are necessary, since (H, o) being a subgroup of (G, o) the composition in H (being also the composition in G) satisfies the closure law

$$\text{i.e. } a, b \in H \Rightarrow aob \in H \quad \forall a, b$$

which proves the first condition.

Also by Problem 14, the identity of H being the same as that of G and by Problem 15, the inverse of any element of H being the same as its inverse in G , we have

$$a \in H \Rightarrow a^{-1} \in H \quad \forall a$$

which proves the second condition.

The conditions are also sufficient, since if the conditions (i) and (ii) hold then

$$G_1 \text{ is satisfied, for } a, b \in H \Rightarrow aob \in H \text{ by condition (i)}$$

$$G_2 \text{ is satisfied, for } a, b \in H \Rightarrow aob \in H \text{ by (i) leads to}$$

$$aob, c \in H \text{ and } a, boc \in H \quad \forall a, b, c \in H$$

$$\Rightarrow \text{the same element } aoboc \in H \text{ i.e. associative law is satisfied.}$$

$$G_3 \text{ is satisfied since } a \in H \Rightarrow a^{-1} \in H \text{ by (ii) leads to}$$

$$a \in H \text{ and } a^{-1} \in H \Rightarrow a a^{-1} \in H \text{ by (i)}$$

$$\text{But } a a^{-1} = e, \text{ (identity of } G)$$

$\therefore e \in H$ is an identity in H , which is also identity in G , thereby showing the existence of an identity element in H .

G_4 is satisfied since from G_3 and condition (ii), every element of H has an inverse.

Hence (H, o) which is a subgroup of the group (G, o) satisfies all the four axioms of group.

Problem 17. Show that a necessary and sufficient condition for a complex H to be a subgroup (H, o) of a group (G, o) is that $a \in H, b \in H \Rightarrow aob^{-1} \in H$.

The condition is necessary, since when (H, o) is a subgroup of (G, o) then by condition (ii) of Problem 16, we have $b \in H \Rightarrow b^{-1} \in H$

$$\text{Also by condition (i) of the Problem 16, } a, b^{-1} \in H \Rightarrow aob^{-1} \in H.$$

$$\text{Combining these two conditions we have } a \in H, b \in H \Rightarrow aob^{-1} \in H.$$

The condition is sufficient, since if $a, b \in H \Rightarrow aob^{-1} \in H$, then we can show as below that (H, o) is a subgroup of (G, o) .

The given condition yields,

$$a \in H, a \in H \Rightarrow a o a^{-1} = e \in H, e \text{ being identity of } G.$$

$$\text{This follows that } G_3 \text{ is satisfied i.e. } \exists \text{ an identity } e \in H.$$

$$\text{Also } e \in H, a \in H \Rightarrow e o a^{-1} = a^{-1} \in H$$

i.e. G_4 is satisfied or in other words every element in H is invertible.

$$\text{As such any } b \in H \Rightarrow b^{-1} \in H$$

$$\text{So that } a \in H, b^{-1} \in H \Rightarrow a o (b^{-1})^{-1} = aob \in H$$

which follows that H satisfies closure law under 'o' i.e. G_1 is satisfied.

Now associativity of G w.r.t 'o' immediately follows the associativity of H w.r.t 'o' i.e. G_2 is satisfied.

Hence (H, o) is a group.

But (H, o) is a subset of (G, o) .

Therefore (H, o) is a subgroup of (G, o) .

Problem 18. Show that the intersection of two subgroups of a group (G, o) is a subgroup of (G, o) .

Let (H_1, o) and (H_2, o) be the two subgroups of (G, o) . Then

$$H_1 \cap H_2 \subset G.$$

$$\text{Now } a, b \in H_1 \cap H_2 \Rightarrow a, b \in H_1, a, b \in H_2$$

$$\Rightarrow aob \in H_1, aob \in H_2 \text{ since } H_1, H_2 \text{ being subgroups, satisfy group axioms.}$$

$$\Rightarrow aob \in H_1 \cap H_2 \quad \forall a, b \in H_1 \cap H_2$$

$$\text{Also } a \in H_1 \cap H_2 \Rightarrow a \in H_1 \text{ and } a \in H_2$$

$$\Rightarrow a^{-1} \in H_1 \text{ and } a^{-1} \in H_2 \text{ since } H_1, H_2 \text{ being subgroups, satisfy group axioms.}$$

$$\Rightarrow a^{-1} \in H_1 \cap H_2.$$

Hence by Problem 16, $H_1 \cap H_2$ is a subgroup of G .

Problem 19. Show that the union of two subgroups of a group (G, o) may not be subgroup of G .

Let (H_1, o) and (H_2, o) be the two subgroups of (G, o) and let

$$a \in H_1, b \in H_2, \text{ so that } a, b \in H_1 \cup H_2.$$

Now $a, b \in H_1 \cup H_2 \Rightarrow a \in H_1, b \in H_2 \nRightarrow aob \in H_1 \cup H_2$ for b may not belong to H_1 .

Hence the union of two subgroups of a group may not be subgroup of the group.

Problem 20. Show that the set $S = \{1, i, -1, -i\}$ is a subgroup of a multiplicative group of non-zero complex numbers.

Let $(G, \cdot)$ be a multiplicative group of non-zero complex numbers. Then $(S, \cdot)$ will be a subgroup of $(G, \cdot)$ if it satisfies both the conditions for a subgroup.

$$\text{The condition (i) is satisfied since } 1 \cdot i = i \in S, 1 \cdot (-1) = -1 \in S, 1 \cdot (-i) = -i \in S, i \cdot (-1) = -i \in S, i \cdot (-i) = 1 \in S, (-1) \cdot (-i) = i \in S.$$

The condition (ii) is satisfied since the inverse of 1 is $1 \in S$, the inverse of i is $-i \in S$, the inverse of -1 is $-1 \in S$ and the inverse of $-i$ is $i \in S$.

$$\text{Hence } (S, \cdot) \text{ is a subgroup of } (G, \cdot).$$

Problem 21. Show that the order of every element of a group (G, o) of finite order is finite.

If a be an element of (G, o) of finite order, then the positive integral powers of a viz. $a, a^2, a^3, a^4 \dots$ will all be the members of G .

But the order of G is finite, therefore all these elements of G can not be different.

$$\text{Suppose that } a^r = a^s, r > s.$$

Then, $a^{r-s} = a^r o a^{-s} = a^r o a^{-r} = a^{r-r} = a^0 = e$, e being the identity in G .

If $r-s=m$, then $a^{r-s} = e \Rightarrow a^m = e$, m being a positive integer as $r > s$.

This follows that $\exists$ a positive integer m s.t. $a^m = e$.

As every set of positive integers essentially possesses a least member so the set of all those positive integers s.t. $a^m = e$ has a least member known as the order of a . But a is arbitrary and hence the order of every element of G is finite.

Problem 22. Show that the order of any power of any element a of a group is atmost equal to the order of the element.

Assuming that order of $a = m$ and order of $(a^p) = n$, $p \in \mathbb{I}$ (set of integers), we have order of $a = m \Rightarrow a^m = e$, e being identity element.

$$\begin{aligned} \Rightarrow (a^m)^p &= e^p \\ \Rightarrow a^{mp} &= e \\ \Rightarrow (a^p)^m &= e \\ \Rightarrow \text{order of } (a^p) &\leq m \end{aligned}$$

which proves the proposition.

Problem 23. Show that the order of any element of a group is always equal to the order of its inverse.

Taking the orders of a and a^{-1} as m and n respectively, we have

$$a^m = e \text{ and } (a^{-1})^n = e$$

Now a^{-1} being an exponent power of a , the Problem 22 leads to order of $(a^{-1}) \leq$ order of a i.e. $n \leq m$.

Also since $a = (a^{-1})^{-1}$ i.e. a is an exponent power of a^{-1} , so by Problem 22, we have order of $a \leq$ order of (a^{-1}) i.e. $m \leq n$.

Hence $m \leq n$ and $n \leq m \Rightarrow m = n$.

Problem 24. If a, b be two elements of a group (G, o) and $ba = a^m b^n \neq a, b \in G$ then prove that the elements $a^m b^{n-2}$, $a^{m-2} b^n$ and ab^{-1} have the same order.

$$\text{We have } (a^{-1}b)^{-1} = b^{-1}(a^{-1})^{-1} = b^{-1}a$$

Since $b^{-1}a$ is the inverse of $a^{-1}b$, therefore by Problem 23, the order of $b^{-1}a$ and $a^{-1}b$ is the same.

$$\begin{aligned} \text{Now } a^m b^{n-2} &= a^m b^n b^{-2} = (ba) b^{-2} \quad \because ba = a^m b^n \\ &= b (ab^{-1}) b^{-1} \quad \because b^{-2} = b^{-1} b^{-1} \end{aligned}$$

But $b (ab^{-1}) b^{-1}$ has the same order as that of ab^{-1} since

$$\begin{aligned} [b (ab^{-1}) b^{-1}]^2 &= [b (ab^{-1}) b^{-1}] [b (ab^{-1}) b^{-1}] \\ &= [b (ab^{-1})] (b^{-1}b) [(ab^{-1}) b^{-1}] \\ &= b (ab^{-1}) (e) (ab^{-1}) b^{-1} \\ &= b (ab^{-1})^2 b^{-1} \end{aligned}$$

$$\text{or in general } [b (ab^{-1}) b^{-1}]^n = b (ab^{-1})^n b^{-1} = b e b^{-1} \text{ if order of } ab^{-1} \text{ be } n \\ = b b^{-1} = e.$$

These results follows that order of ab^{-1} is the same as that of $a^m b^{n-2}$

Again $a^{m-2} b^n = a^{-2} (a^m b^n) = a^{-2} (ba) = a^{-1} (a^{-1}b) a$
i.e. as above, the order of $a^{-1}b$ is the same as that of $a^{m-2} b^n$.

Problem 25. If the elements a, b and aob of a group (G, o) are each of order 2, then show that the group is abelian.

The order of aob being 2, we have $(aob)^2 = e$, e is the identity in G .

$$\begin{aligned} \therefore (aob) o (aob) &= e \Rightarrow ao (aob) o (aob) = aoe \\ &\Rightarrow ao (aob) o (aob) ob = ao eob \\ &\Rightarrow (aoa) o (boa) o (bob) = ao eob \text{ by associative law.} \\ &\Rightarrow a^2 o (boa) ob^2 = ao eob \\ &\Rightarrow eo (boa) oe = ao eob \text{ since the order of } a \text{ and } b \text{ is 2.} \end{aligned}$$

$$\Rightarrow boa = aob$$

which proves that a and b commute and hence the group is abelian.

4.5. THE CENTRE OF A GROUP

If (G, o) be a group and H be the set of those elements $x \in G$, which commute with each element in G i.e. the set

$$H = \{x : x \in G \text{ and } aox = xoa \quad \forall a \in G\}$$

then the set H is known as the **centre** of G .

THEOREM. The centre of G is a subgroup of (G, o) .

If H be the centre of G , then we have by definition

$$H = \{x : x \in G \text{ and } aox = xoa \quad \forall a \in G\}$$

$$\therefore x_1, x_2 \in H \Rightarrow aox_1 = x_1 o a \text{ and } aox_2 = x_2 o a \quad \forall a \in G.$$

But $aox_1 = x_1 o a = x_1 o (x_2^{-1} o x_2) o a$, since $x_2^{-1} o x_2 = e$, the identity in H and $x_1 o e o a = x_1 o a$

$$\begin{aligned} &= (x_1 o x_2^{-1}) o (x_2 o a) \\ &= (x_1 o x_2^{-1}) o (aox_2) \quad \because aox_2 = x_2 o a. \end{aligned}$$

$$\begin{aligned} \therefore aox_1 &= (x_1 o x_2^{-1}) o (aox_2) \Rightarrow (aox_1) o x_2^{-1} = (x_1 o x_2^{-1}) o a \\ &\Rightarrow ao (x_1 o x_2^{-1}) = (x_1 o x_2^{-1}) o a \\ &\Rightarrow x_1 o x_2^{-1} \text{ commutes with } a \in G \\ &\Rightarrow x_1 o x_2^{-1} \in H \end{aligned}$$

$$\text{Conclusively } x_1 \in H, x_2 \in H \Rightarrow x_1 o x_2^{-1} \in H$$

Which follows by the definition of a subgroup that (H, o) is a subgroup of (G, o) .

4.6. COSETS OR COSETS OF A SUBGROUP

Let (G, o) be a group, (H, o) be a subgroup of (G, o) and ' a ' be an element in G i.e. $a \in G$. Then the set

$$aH = \{ah : h \in H\} \text{ (not using the binary operation)}$$

i.e. the collection,

$$\begin{aligned} aoH &= \{ao h_1, ao h_2, \dots, ao h_i, \dots\}, h_i \in H \\ &= \{aox : x \in H \text{ and } a \in G\} \end{aligned}$$

is said to be the **Left Coset** of H in G ;

and the set $aH = \{ah : h \in H\}$

i.e. the collection, $aH = \{h_1 a, h_2 a, \dots, h_i a, \dots\} \quad h_i \in H$
 $= \{x a : x \in H \text{ and } a \in G\}$

is said to be the **Right Coset** of H in G .

Since $eH = H = He$, therefore H is itself a coset.

If the cosets aH and bH are such that $aH \cap bH \neq \emptyset$, then $aH = bH$, hence the cosets have no element in common with H i.e. two cosets contain either the same elements or have no elements in common. Also the cosets do not form a group.

The number of left (or right) cosets of H in G is said to be the **Index** of H in G and denoted by $(G : H)$.

THEOREM 1. If (H, o) or simply H be a subgroup of (G, o) or simply G , then H is both a left coset and a right coset.

If e be the identity in G , then $He = eH = H$, which follows that H is both a left coset as well as a right coset of H in G .

THEOREM 2. If H be a subgroup of G , then $aH = H \Leftrightarrow a \in H$.

If e be the identity in G and so is in H , then

$$aH = H \Rightarrow ae \in H$$

$$\text{i.e. } aH = H \Rightarrow a \in H \quad \dots (1)$$

Again, if $a \in H$ and $h \in H$ then

$$a \in H \Rightarrow ah \in H \quad \forall h \in H$$

$$\therefore aH \subset H$$

Also $a \in H \Rightarrow a^{-1} \in H$, H being a subgroup of the group G , satisfies group axioms.

$$\Rightarrow a^{-1}h \in H \quad \forall h \in H \text{ by closure law in } H$$

$$\Rightarrow a(a^{-1}h) \in H \quad \forall h \in H \text{ by closure law in } H$$

$$\Rightarrow h \in aH \quad \forall h \in H$$

$$\therefore H \subset aH$$

So $aH \subset H$ and $H \subset aH \Rightarrow aH = H$

$$\text{Ultimately } a \in H \Rightarrow aH = H \quad \dots (2)$$

Hence $aH = H \Leftrightarrow a \in H$ by (1) and (2).

THEOREM 3. If $a, b \in G$ and $a \neq b$ then $aH = bH \Leftrightarrow a^{-1}b \in H$ where H is a subgroup of the group G .

We have,

$$aH = bH \Rightarrow a^{-1}aH = a^{-1}bH$$

$$\Rightarrow (a^{-1}a)H = (a^{-1}b)H$$

$$\Rightarrow eH = (a^{-1}b)H, \quad e \text{ being the identity in } G \text{ and so in } H.$$

$$\Rightarrow H = (a^{-1}b)H$$

$$\therefore aH = bH \Rightarrow a^{-1}b \in H \text{ by theorem 1.} \quad \dots (1)$$

Also, if $a^{-1}b \in H$, then

$$bH = e(bH) = (aa^{-1})(bH) = a(a^{-1}b)H = aH \text{ by theorem 1.} \dots (2)$$

(1) and (2) follow that $aH = bH \Leftrightarrow a^{-1}b \in H$.

THEOREM 4. The two left cosets aH and bH of a subgroup H of a group G are either identical or disjoint.

There arise two cases :

Case I. If $aH \neq bH$, then we have to show that aH and bH are disjoint.

Let us assume if possible that $x \in aH$ and $x \in bH$.

Then $x = ay, y \in H$ and $x = bz, z \in H$

$$\therefore ay = bz \Rightarrow ayz^{-1} = bzz^{-1} \Rightarrow a(yz^{-1}) = b(zz^{-1}) = be = b$$

$$\Rightarrow (a^{-1}a)(yz^{-1}) = a^{-1}b$$

$$\Rightarrow e(yz^{-1}) = a^{-1}b$$

$$\Rightarrow yz^{-1} = a^{-1}b$$

Thus, $yz^{-1} \in H \Rightarrow a^{-1}b \in H$.

So that by theorem 3, it follows that $aH = bH$, which contradicts the hypothesis and hence two unequal cosets cannot have any element in common i.e. aH and bH are disjoint.

Case II. If aH and bH are not disjoint, then we have to show that $aH = bH$.

aH and bH are not disjoint $\Rightarrow \exists$ an element common to aH and bH

$$\Rightarrow \exists h_i, h_j \text{ s.t. } ah_i = bh_j$$

$$\Rightarrow a(h_i h_i^{-1}) = b h_j h_i^{-1}$$

$$\Rightarrow a = bh_j h_i^{-1}$$

$$\Rightarrow ah = b(h_j h_i^{-1} h) \quad \forall h \in H$$

$$\Rightarrow ah \in bH \quad \forall h \in H$$

$$\Rightarrow aH \subset bH \quad \dots (1)$$

Similarly it can be shown that

$$ah_i = bh_j \Rightarrow bH \subset aH \quad \dots (2)$$

(1) and (2) follow that $aH = bH$ i.e. aH and bH are identical.

THEOREM 5. If H be a subgroup of the group G and $a \in G$ but $a \notin H$ then $\exists$ one-one mapping of H onto aH .

Taking $f : H \rightarrow aH$ defined by $f(h) = ah, h \in H$, we have to show that the map f is onto.

Every element of the left coset aH being of the form $ah, h \in H$, and so being the f -image of h in H , the mapping f is onto.

Again to show that f is one-one, let $h_i, h_j \in H$ s.t. $ah_i = ah_j$.

Then $ah_i = ah_j \Rightarrow h_i = h_j$ by left cancellation law.

So f is one-one.

Conclusively f is a one-one mapping of H onto aH .

Note. This theorem follows that if H be a finite subgroup, the number of elements in each of its left cosets is the same as the number of elements in H i.e. equal to the order of H .

THEOREM 6. (Lagrange's theorem). *The order of every subgroup of a finite group is a divisor of the order of the group.*

Let H be a subgroup of a finite group G . So G being finite, H is also finite.

Let m and n be the order of H and G respectively.

Since the order of H is m , therefore H consists of exactly m elements or in other words every coset aH has exactly m elements, for if $h_1, h_2 \in H$, $ah_1 = ah_2$ iff $h_1 = h_2$, hence aH has the same number of elements as H .

Now if $m = n$, the theorem is self-evident.

But if $n > m$, then G being of finite order, there are only a finite number say k , of different cosets of H in G .

Taking $H = \{h_1, h_2, \dots, h_m\}$, if $a \in G$ but $a \notin H$ and binary operation of G being denoted multiplicatively, the distinct m elements

$ah_1, ah_2, \dots, ah_m \in H$ but belong to G by closure axiom.

Denoting the set formed by these m elements by H' i.e.

$$H' = \{ah_1, ah_2, \dots, ah_m\}$$

We observe that if $H \cup H'$ is a proper subset of G then there is an element say $b \in G$ s.t. $b \notin H \cup H'$. We thus have again a set of m distinct elements

$bh_1, bh_2, \dots, bh_m$ which belong to G but not to $H \cup H'$

Denoting the set of these m elements by H'' i.e.

$$H'' = \{bh_1, bh_2, \dots, bh_m\}$$

and continuing this process, we see that G can be divided into k subsets each consisting of m elements.

$\therefore$ Order of G = number of elements in G

i.e. $n = k \times \text{order of } H$

$$= km$$

which follows that the order of H is a divisor of the order of G .

COROLLARY 1. *The order of an element of a group G of finite order is a divisor of the order of the group.*

Let m be the order of the group G and $a \in G$. Then by definition, $a^m = e$, m being least positive integer and e being identity in G .

Evidently the elements $a, a^2, a^3, \dots, a^{m-1}, a^m \in G$, are all distinct and form a subgroup of order m . Also by definition, m is the order of a .

$\therefore$ The order m of a is a divisor of the order of the group.

COROLLARY 2. *A finite group of prime order has no proper subgroup.*

Let G be a finite group of order p , where p is a prime. Then by Lagrange's theorem, the order of any subgroup of G is divisor of p . But p being prime has no divisor and hence there is no proper subgroup of G .

COROLLARY 3. Fermat's theorem. *If p be a prime and ' a ' a natural number not divisible by p , then*

$$a^{p-1} \equiv 1 \pmod{p}$$

Taking the multiplicative group of non-zero residue classes modulo p and a not divisible by p , we have the equivalence class $\bar{a} \neq 0$ i.e. $[a] \neq 0$.

But the order of the group being $p-1$, it follows from Cor. 1, that

$$[a^{p-1}] = [1]$$

which yields $a^{p-1} \equiv 1 \pmod{p}$.

Problem 26. *If H be a subgroup of a group G and m, n are the orders of m and n respectively then prove that $a^n = e$, e being identity in G .*

Lagrange's theorem gives $n = km$, k being some positive integer

$$\therefore a^n = a^{km} = (a^m)^k = e^k = e.$$

Problem 27. *Find the cosets of the additive subgroup $(2\mathbb{I}, +)$ of the additive group $(\mathbb{I}, +)$, $\mathbb{I}$ being set of all integers.*

We have

$$\mathbb{I} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

and say $H = (2\mathbb{I}, +) = \{\dots, -6, -4, -2, 0, 2, 4, 6, \dots\}$

If $a \in \mathbb{I}$ then the coset of H in $\mathbb{I}$ corresponding to a is $2\mathbb{I} + a$ since the group being abelian, $\mathbb{I} + a = a + \mathbb{I}$

$$\therefore 2\mathbb{I} + 0 = \{\dots, -6, -4, -2, 0, 2, 4, 6, \dots\}$$

$$2\mathbb{I} + 1 = \{\dots, -5, -3, -1, 1, 3, 5, 7, \dots\}$$

$$2\mathbb{I} + 2 = \{\dots, -4, -2, 0, 2, 4, 6, 8, \dots\} = 2\mathbb{I}$$

$$2\mathbb{I} + 3 = \{\dots, -3, -1, 1, 3, 5, 7, 9, \dots\} = 2\mathbb{I} + 1$$

$$2\mathbb{I} + 4 = \{\dots, -2, 0, 2, 4, 6, 8, 10, \dots\} = 2\mathbb{I}$$

$$2\mathbb{I} + 5 = \{\dots, -1, 1, 3, 5, 7, 9, 11, \dots\} = 2\mathbb{I} + 1 \text{ and so on.}$$

Thus the distinct cosets of H in $\mathbb{I}$ are $2\mathbb{I}$ and $2\mathbb{I} + 1$.

Clearly $2\mathbb{I} \cup (2\mathbb{I} + 1) = \mathbb{I}$.

4.7. CYCLIC GROUPS

If a group H contains an element a s.t. it is capable of being generated by the single element a i.e. every element of H is of the form a^n for some integer n , then H is said to be a cyclic group and a is known as the generator of H . We also denote $H = \langle a \rangle$.

Hence if H is a cyclic group, then $\exists a \in H, b \in H$ s.t. $a^n = b$ (in multiplicative form) or $b = na$ (in additive form) for some integer n .

Thus $H = \{a^n : n \in \mathbb{I}\}$, $a \in H$, $\mathbb{I}$ being set of integers.

e.g. the unit circle $\{z : |z| = 1\}$ in the complex plane is a cyclic group.

Characteristics of Cyclic Group

(i) Every cyclic group is abelian.

If H be a cyclic group and a is its generator, then

$$a^m, a^n \in H \quad \forall m, n \in \mathbb{I}$$

$$\therefore a^m \circ a^n = a^{m+n} = a^{n+m} = a^n \circ a^m$$

which proves the commutative property and hence every cyclic group is abelian.

(ii) The order of a cyclic is the same as that of its generator.

Let H be a cyclic group, a its generator and e the identity element in H . Also let n be the order of a , so that $a^n = e$

Evidently, $m \in \mathbb{I}$ and $m < n \Rightarrow a^m \neq e$.

In case $m > n$, then if q be the quotient and r the least positive remainder when m is divided by n ,

$$m = nq + r$$

$$\text{So that } a^m = a^{nq+r} = a^{nq} \circ a^r = (a^n)^q \circ a^r = e^q \circ a^r = a^r$$

where $r = 0, 1, 2, \dots, (n-1)$.

By closure axiom since $a^m \in H$, therefore n distinct elements belonging to H are $a^0, a^1, a^2, a^3, \dots, a^{n-1}$ where $a^0 = e = a^n$.

As such there are only n elements in H and hence the order of the cyclic group H is also n which is the order of its generator.

(iii) The generators of a cyclic group of order n are the generators a^p where p is prime to n and $0 < p < n$.

$$\therefore a^n = e, \therefore (a^p)^n = (a^n)^p = e^p = e$$

which shows that order of $a^p \leq n$.

Taking $s \in \mathbb{I}$ s.t. $0 < s < n$, we have ps prime to n since n is neither a factor of p , nor of s .

Let $ps = nq + r$, q being quotient and r the least positive remainder when ps is divided by n and $0 \leq r < n-1$.

$$\text{Thus } (a^p)^s = a^{ps} = a^{nq+r} = a^{nq} \circ a^r = (a^n)^q \circ a^r = e^q \circ a^r = e \circ a^r = a^r$$

where $r = 0, 1, 2, \dots, n-1$.

It is clear that $a^r \neq e$

Hence the order of a^p is n and a^p is the generator of the group.

(iv) A subgroup H' of a cyclic group H is also cyclic.

Let a be the generator of H . Given that H' is a subgroup of H . Therefore every element of H and so of H' will be of the form a^n , n being an integer.

Let m be the least positive integer s.t. $a^m \in H'$.

If m does not divide n then $\exists$ integers q (quotient) and r (remainder) s.t. $n = mq + r$, $0 \leq r < m$

$$\therefore a^n = a^{mq+r} = a^{mq} \circ a^r \text{ giving } a^r = (a^{mq})^{-1} \circ a^n \quad \dots (1)$$

But $a^m \in H' \therefore$ by closure law $a^{mq} \in H'$ and so $(a^{mq})^{-1} \in H'$ since H' satisfies group axioms.

Now $a^n \in H'$ (by hypothesis)

$\therefore$ (1) yields, $a^r \in H'$ which contradicts the assumption that m is the least positive integer s.t. $a^m \in H'$

Thus the only possibility is that $r=0$ and then $n=mq$ so that $a^n = a^{mq} = (a^m)^q$.

Which follows that every element a^n of H' is of the form $(a^m)^q$ showing that a^m is the generator of H' and hence H' is cyclic.

Finite cyclic groups. If H is a cyclic group generated by a s.t. all the powers of a are not different then $H = \{a\}$ is a finite cyclic group.

If $n (>0)$ be the order of a , then $a^n = e$

Given any integer $s \exists$ two integers q and r s.t. $s = nq + r$, $0 \leq r < n$.

$$\therefore a^s = a^{nq+r} = a^{nq} \circ a^r = (a^n)^q \circ a^r = e^q \circ a^r = e \circ a^r = a^r$$

Which follows that there are at most n distinct elements $a^1, a^2, a^3, \dots, a^{n-1}, a^n = e$

To show that no two of these n elements are equal, let us assume if possible that $a^x = a^y$, $0 < y < x < n$

$$\therefore a^{x-y} = a^y \circ a^{-y} = a^0 = e$$

But $0 < n - y < n$ and order of a being n , $a^{n-y} \neq e$ i.e. $a^x \neq a^y$.

Thus H contains exactly n (finite) distinct elements

$$a^1, a^2, \dots, a^{n-1}, a^n.$$

Hence H is a finite cyclic group of order n .

Infinite cyclic groups. If H be a cyclic group generated by a s.t. all the powers of a are distinct, then $H = \{a\}$ is an infinite cyclic group.

Let a be the generator of H . Then all the powers of a being different the order of a is zero.

Let us assume, if possible that $a^s = a^r$ where $s > r$.

Then $a^{s-r} = a^r \circ a^{-r} = a^0 = e$ which contradicts the assumption that the order of a is zero.

$$\therefore a^s \neq a^r$$

i.e. H contains an infinite number of elements and hence H is an infinite cyclic group.

THEOREM 1. In an infinite cyclic group, there are exactly two distinct generators namely one generator and the other its inverse.

Let H be an infinite cyclic group and a , one of its generator. Then since $a_n = (a^{-1})^{-n}$, therefore a^{-1} is the other generator.

Also $a \neq a^{-1}$, otherwise $a = a^{-1} \Rightarrow aa^{-1} = a^2 = e = a$ finite cyclic group of order 2 which contradicts the hypothesis that the cyclic group is infinite.

To show that $\exists$ third generator, if possible suppose that b is the third generator of H , so that a and b being both generators of H , $a = b^m$ and $b = a^l$,

$$\therefore a = (a^l)^m = a^{ml} \quad \dots (1)$$

But H being infinite cyclic group, $r \neq n \Rightarrow a^r \neq a^n$,

$\therefore$ the relation (1) is satisfied if $ml=1$, m, l being both integers.

It follows that either $m=+1$ or $m=-1$

i.e. either $b=a$ or $b=a^{-1}$.

So that $\exists$ third generator of H other than a and a^{-1} .

THEOREM 2. Every subgroup of an infinite cyclic group is infinite.

Let H' be a subgroup of an infinite cyclic group H whose generator is a . Then by characteristic (iv) of groups, we have $H'=\{a^m\}$, m being least positive integer s.t. $a^m \in H'$.

Assume, if possible that H' is finite, then $(a^m)^n=e$ for some $n>0$ which follows that a is of finite order and so H is finite which contradicts the hypothesis.

Hence H' must be an infinite cyclic subgroup of H .

Problem 28. Show that the group formed by the set $\{1, \omega, \omega^2\}$, ω being cube root of unity i.e. $\omega^3=1$, is a cyclic group of order 3 with respect to multiplication.

Here $\omega^3=1$ is the identity and ω is the generator as its powers generate the elements $1, \omega, \omega^2$ as tabulated:

The group axioms are satisfied, since if $G=\{1, \omega, \omega^2\}$ w.r.t. ' $\cdot$ ' then

$G_1-1, \omega, \omega^2 \in G, 1 \cdot \omega, 1 \cdot \omega^2, \omega \cdot \omega^2 \in G$ as $\omega^3=1$

$G_2-(1 \cdot \omega) \cdot \omega^2 = 1 \cdot (\omega \cdot \omega^2) = \omega \cdot \omega^2 = \omega^3 = 1$

G_3-1 is the identity elements as $\omega \cdot 1 = \omega$ etc.

G_4 -Inverses of $1, \omega, \omega^2$ are respectively $1, \omega^2, \omega$ as

$$1 \cdot 1 = \omega \cdot \omega^2 = \omega^2 \cdot \omega = 1 \text{ (the identity element)}$$

Hence $\{1, \omega, \omega^2\}$ is a cyclic group of order 3 with generator ω .

Problem 29. Find all the generators of the cyclic group $\{a, a^2, a^3, a^4, a^5, a^6, a^7, a^8=e\}$ of order 8.

Let $H=\{a, a^2, a^3, a^4, a^5, a^6, a^7, a^8=e\}$

Since it contains all powers of a , so a is a generator.

Now $(a^3)^1=a^3, (a^3)^2=a^6, (a^3)^3=a^9=a^8oa^1=eo a=a,$

$(a^3)^4=a^{12}=a^8oa^4=eo a^4=a^4, (a^3)^5=a^{15}=a^8oa^7=eo a^7=a^7$

$(a^3)^6=a^{18}=(a^3)^3oa^2=e^2oa^2=eo a^2=a^2$

$(a^3)^7=a^{21}=(a^3)^2oa^5=e^2oa^5=a^5$

$(a^3)^8=a^{24}=(a^3)^3=e^3e$

Since powers of a^3 are the elements of H so a^3 is a generator of H . Similarly a^5 and a^7 are also the generators of H .

4.8. PERMUTATION OR TRANSFORMATION

If G be a set then a one-one onto mapping $f: G \rightarrow G$ is said to be a transformation or in case G is finite, f is said to be a permutation.

In fact the permutation is a rearrangement of the elements of the set and the permutation groups are associated with symmetry groups introduced a bit later.

Consider a set $\{1, 2, 3\}$ with three elements. Its symmetry group or permutations may be written as

$$P_1=I \text{ (identity)} = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}, P_2 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}, P_3 = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}$$

$$P_4 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}, P_5 = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}, P_6 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}$$

If we multiply P_5 and P_6 , we have

$$P_6 P_5 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix} \text{ as } \begin{matrix} 1 \rightarrow 2 \rightarrow 2 \\ 2 \rightarrow 1 \rightarrow 3 \\ 3 \rightarrow 3 \rightarrow 1 \end{matrix}$$

$$= P_2$$

$$\text{and } P_5 H_6 = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} = P_3$$

It is clear that $P_6 P_5 \neq P_5 P_6$

i.e. the permutation multiplication is not commutative, but it can be shown that permutation multiplication is associative, since

$$P_1 (P_2 P_3) = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} \left[\begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \right]$$

$$= \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}$$

$$\text{and } (P_1 P_2) P_3 = \left[\begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix} \right] \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}$$

$$\therefore P_1 (P_2 P_3) = (P_1 P_2) P_3$$

To find the inverse of $\begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}$ (say) let us assume that its inverse

is $\begin{pmatrix} 1 & 2 & 3 \\ x & y & z \end{pmatrix}$. Each of the two permutations whose product is the

Identity Permutation. $I = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}$ i.e. Pre-image of every element is

the same element), being called Inverse to each other, we have

$$\begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ x & y & z \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}$$

$$\text{i.e. } \begin{pmatrix} 1 & 2 & 3 \\ z & x & y \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}$$

Comparison gives $z=1, x=2, y=3$

Hence the inverse of $\begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}$ is $\begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}$

Thus if P^{-1} be the inverse of a permutation P , then we can easily show that

$$(i) PP^{-1} = I = P^{-1}P$$

$$(ii) P^{-1} \text{ is unique}$$

$$(iii) (P^{-1})^{-1} = P$$

(iv) $(PQ)^{-1} = Q^{-1}P^{-1}$ (Reversal law)

The set of all permutations of n elements.

The $|n|$ permutations on n elements (objects) form a group with respect to permutation multiplication. Such a group of all permutations of n elements is denoted by S_n and called the Symmetric group of degree n as it satisfies all the four group axioms.

Note 1. The number of elements in the finite set permuted is known as the degree of permutation.

Note 2. The number of elements in a permutation on n elements is $|n|$.

A permutation which replaces n elements cyclically is said to be a cyclic permutation of degree n e.g., $(1\ 2\ 3\ \dots\ n-1\ n)$ is cyclic and may be denoted by $(1\ 2\ 3\ \dots\ n)$

The number of elements permuted by a cycle is said to be its length and the disjoint cycles are those which have no common elements.

Every cycle can be uniquely expressed as product of disjoint cycles

$$\text{e.g. } \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 4 & 3 & 2 & 5 & 7 & 6 & 1 \end{pmatrix} = (1\ 4\ 5\ 7)(2\ 3)(6) \\ = (2\ 3)(1\ 4\ 5\ 7)(6)$$

$$\text{where } (1\ 4\ 5\ 7) \Rightarrow \begin{pmatrix} 1 & 4 & 5 & 7 & 2 & 3 & 6 \\ 4 & 5 & 7 & 1 & 2 & 3 & 6 \end{pmatrix} \text{ etc.}$$

A cyclic permutation such as $(a\ b)$ which interchanges the symbols leaving all other unchanged is called a Transposition.

In other words Transposition is cycle of length two of the form $(a\ b)$ i.e. it is a mapping which maps each object onto itself excepting two, each of which is mapped on the other. e.g. $(1\ 2)$ is a transposition.

Note 3. A cycle of length one is invariant.

Note 4. Transposition is its own inverse, since, if $(a\ b)$ be the transposition $\begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}$ then $(a\ b)(a\ b) = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} \\ = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} = I.$

Note 5. Any permutation can be resolved as the product of transpositions in infinitely many ways since every permutation can be expressed as a product of disjoint cycles and every cycle can be expressed as the product of transpositions in an infinite way, therefore the proposition follows.

Note 6. The order of transposition cannot be changed, since they may not be disjoint.

Even and odd permutations. A permutation is said to be even or odd

according as it is expressed as a product of an even or odd number of transpositions e.g. the permutation

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 3 & 4 & 8 & 5 & 2 & 7 & 9 & 6 & 1 \end{pmatrix}$$

is an odd permutation as it can be expressed as the product of seven transpositions such as

$$(1\ 3)(1\ 8)(1\ 6)(1\ 7)(1\ 9)(2\ 4)(2\ 5)$$

and the permutation $\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 6 & 4 & 5 & 2 & 1 & 3 \end{pmatrix}$ is even since it can be expressed

as the product of four transpositions such as $(1\ 6)(1\ 3)(1\ 5)(2\ 4)$

Note 7. For any manner of expressing a given permutation as a product of transpositions, the number of transpositions is either necessarily odd or even.

Note 8. The product of two even or two odd permutations is even while the product of an even and an odd permutation is odd.

Note 9. Of the $|n|$ permutations on n elements, $\frac{|n|}{2}$ are even and $\frac{|n|}{2}$ are odd

Alternating set or group. The set of all even permutations of degree n is known as an alternating set or group and is denoted by A_n and symbolized as

$A_n = \{\alpha : \alpha \text{ is an even permutation on a set containing } n \text{ elements}\}.$

e.g. if $E_1, E_2, E_3, \dots, E_s$ be the even permutations of n symbols then G_1 is satisfied, since the product of two even permutations is even and so A_n is closed.

G_2 is satisfied, since permutation composition is associative.

G_3 is satisfied, since the identity permutation considered as even permutation is also identity for even permutations.

G_4 is satisfied, since an $f \in A_n \exists f^{-1} \in A_n$ for, $f^{-1}of = fof^{-1} = I$ and f and I are even permutations so that f^{-1} is also an even permutation.

But the composition in A_n is not commutative since permutation composition is not commutative.

Hence A_n is a group which is non-abelian.

Also the set A_n contains $\frac{|n|}{2}$ elements, hence (A_n, o) is a non abelian group of order $\frac{|n|}{2}$.

Permutation group. Any group whose elements are permutations is said to be a permutation group.

Any subgroup of S_n (symmetric group of degree n) is essentially a permutation group.

$$\text{e.g. if } P_1 = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}, P_2 = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}, P_3 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}, P_4 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix},$$

$$P_5 = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}, P_6 = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} \text{ be the six permutations on the}$$

	P_1	P_2	P_3	P_4	P_5	P_6
P_1	P_1	P_2	P_3	P_4	P_5	P_6
P_2	P_2	P_1	P_6	P_3	P_4	P_5
P_3	P_3	P_6	P_1	P_5	P_4	P_2
P_4	P_4	P_3	P_5	P_1	P_2	P_6
P_5	P_5	P_4	P_2	P_6	P_3	P_1
P_6	P_6	P_5	P_4	P_2	P_1	P_3

set $\{1, 2, 3\}$, then the elements $P_1, P_2, P_3, P_4, P_5, P_6$ constitute all the elements of the symmetric group S_3 of degree 3, whose multiplication table is as shown have.

The elements P_1, P_4, P_6 constitute one of the several subgroups of S_3 and this is a permutation group on three symbols.

Problem 30. Show that the cycle $(1\ 2\ 3\ 3\ 5)$ may be expressed as a product of 3 cycles.

We have

$$(1\ 2\ 3\ 4\ 5) = (1\ 5)(1\ 4)(1\ 3)(1\ 2) \\ = (1\ 4\ 5)(1\ 3)(1\ 2)$$

Problem 31. Express $(1\ 2\ 3\ 4\ 5\ 6)$ as the product of disjoint cycles.

Denoting the given permutation by f , which is the permutation of 6 positive numbers, let us determine the set of images of 1 under the successive powers of f i.e.

$$f(1) = 3 \text{ as } 1 \rightarrow 3$$

$$f^2 = f[f(1)] = f(3) = 4 \text{ as } 3 \rightarrow 4$$

$$f^3 = f[f^2(1)] = f(4) = 1 \text{ as } 4 \rightarrow 1$$

which follows that the first cycle is $(1\ 3\ 4)$.

It is clear that 2 does not belong to this cycle.

We have,

$$f(2) = 6 \text{ as } 2 \rightarrow 6$$

$$f(6) = 5 \text{ and } f(5) = 2$$

i.e. another cycle is $(2\ 6\ 5)$.

Since all the six elements are exhausted in two cycles, we have

$$(1\ 2\ 3\ 4\ 5\ 6) = (1\ 3\ 4)(2\ 6\ 5)$$

Problem 32. Find all the permutations of four letters a, b, c, d which leave the expression $ab+cd$ invariant.

If a, b, c, d are distinct letters then the three functions

$$y_1 = ab+cd, \quad y_2 = ac+bd, \quad y_3 = ad+bc$$

are distinct and there are the only functions of the given four. This follows that each of the 24 permutations on a, b, c, d replaces y_1 by y_1, y_2 or y_3 so that $\frac{1}{3} \times 24$ i.e. 8 of them leave y_1 invariant which may be verified by showing that $I, (a\ b), (cd), (ab)(cd), (ac)(bd), (ad)(bc), (a\ d\ b\ c), (a\ c\ b\ d)$ leave $ab+cd$ invariant.

We have $I = \begin{pmatrix} a & b & c & d \\ a & b & c & d \end{pmatrix}$ where

$$\begin{aligned} a &\rightarrow a \\ b &\rightarrow b \\ c &\rightarrow c \\ d &\rightarrow d \end{aligned}$$

so that $ab+cd=ab+cd$

$(a\ b) = \begin{pmatrix} a & b & c & d \\ b & a & c & d \end{pmatrix}$ where

$$\begin{aligned} a &\rightarrow b \\ b &\rightarrow a \\ c &\rightarrow c \\ d &\rightarrow d \end{aligned}$$

so that $ab+cd=ba+cd=ab+cd$

$(c\ d) = \begin{pmatrix} a & b & c & d \\ a & b & d & c \end{pmatrix}$ where

$$\begin{aligned} a &\rightarrow a \\ b &\rightarrow b \\ c &\rightarrow d \\ d &\rightarrow c \end{aligned}$$

so that $ab+cd=ab+dc=ab+cd$

$(a\ b)(c\ d) = \begin{pmatrix} a & b & c & d \\ b & a & d & c \end{pmatrix}$ where

$$\begin{aligned} a &\rightarrow b \\ b &\rightarrow a \\ c &\rightarrow d \\ d &\rightarrow c \end{aligned}$$

so that $ab+cd=ba+dc=ab+cd$

$(a\ c)(b\ d) = \begin{pmatrix} a & c & b & d \\ c & a & d & b \end{pmatrix}$ where

$$\begin{aligned} a &\rightarrow c \\ c &\rightarrow a \\ b &\rightarrow d \\ d &\rightarrow b \end{aligned}$$

so that $ab+cd=cd+ab=ab+cd$

$(a\ d)(b\ c) = \begin{pmatrix} a & d & b & c \\ d & a & c & b \end{pmatrix}$ where

$$\begin{aligned} a &\rightarrow d \\ d &\rightarrow a \\ b &\rightarrow c \\ c &\rightarrow b \end{aligned}$$

so that $ab+cd=dc+ba=ab+cd$

$(a\ d\ b\ c) = \begin{pmatrix} a & d & b & c \\ d & b & c & a \end{pmatrix}$ where

$$\begin{aligned} a &\rightarrow d \\ d &\rightarrow b \\ b &\rightarrow c \\ c &\rightarrow a \end{aligned}$$

so that $ab+cd=dc+ab=ab+cd$

and $(a\ c\ b\ d) = \begin{pmatrix} a & c & b & d \\ c & b & d & a \end{pmatrix}$ where

$$\begin{aligned} a &\rightarrow c \\ c &\rightarrow b \\ b &\rightarrow d \\ d &\rightarrow a \end{aligned}$$

so that $ab+cd=cd+ba=ab+cd$

Conclusively y_1 remains invariant by the 8 permutations mentioned above.

4.9. HOMOMORPHISM AND ISOMORPHISM OF GROUPS

Homomorphism of groups. If (G, o) and (G', o') be two groups, then a mapping $f: G \rightarrow G'$ which retains the structure and is many one is called **Homomorphism of the group G with the group G'** s.t.

$$f(aob) = f(a) \cdot o' f(b), \quad \forall a, b \in G.$$

We sometimes use to say that G is homomorphic to G' and denote it by $G \cong G'$ if $\exists$ a mapping $f: G \rightarrow G'$ s.t. $f(aob) = f(a) \cdot o' f(b)$
 $\forall a, b \in G.$

Properties of homomorphism

(1) The group (G', o') is a homomorphic image of the group (G, o)

(2) The relation of homomorphism is not symmetric i.e.

$$G \cong G' \nRightarrow G' \cong G$$

(3) The homomorphic image of the identity of the group (G, o) is the identity of the group (G', o') i.e. if e, e' be the identities in G, G' respectively then $f(e) = e'$.

We have $a \in G \Rightarrow f(a) \in G'$

and $f(aoe) = f(a) \cdot o' f(e) \quad \forall a \in G$ by definition of homomorphism.

$\therefore f(a) \cdot o' e' = f(a) = f(aoe) = f(a) \cdot o' f(e)$ since $aoe = a$

and $f(a) \cdot o' e' = f(a)$

So left cancellation law gives $e' = f(e)$

(4) The homomorphic image of the inverse of any element a of a group (G, o) is the inverse of the image of a i.e. $f(a^{-1}) = [f(a)]^{-1}$
 $\forall a \in G$

We have $a^{-1}, a \in G \Rightarrow f(a^{-1}), f(a) \in G'$

$\therefore f(a^{-1}) \cdot o' f(a) = f(a^{-1}oa)$, by definition of homomorphism
 $= f(e) = e'$ by property (3)

But $f(a^{-1}) \cdot o' f(a) = e' \Rightarrow f(a^{-1}) = [f(a)]^{-1}$
 $\therefore f(a), f(a^{-1}) \in G'$

Isomorphism of groups. If (G, o) and (G', o') are two groups and mapped that

$\exists$ a one-one onto mapping $f: G \rightarrow G'$ s.t. $aob \xrightarrow{\text{mapped that}} a'ob'$ where $a \rightarrow a', b \rightarrow b', \forall ab \in G$ and $a', b' \in G'$, then the mapping f is called as **Isomorphism** and we say that G is **isomorphic** to G' and write $G \cong G'$.

e.g. if G is an additive group of all integers i.e.

$$G = \{\dots, -4, -3, -2, -1, 0, 1, 2, 3, 4, \dots\}$$

and G' is a multiplicative group of all positive and negative powers of an integer 2 i.e.

$$G' = \{2^m : m = 0, \pm 1, \pm 2, \dots\}$$

$$= \{\dots, \frac{1}{8}, \frac{1}{4}, \frac{1}{2}, 1, 2, 4, 8, \dots\}$$

Then we have $f(m) = 2^m$, m being an integer

and $f(m+n) = 2^{m+n} = 2^m \cdot 2^n = f(m) \cdot f(n)$, m, n being integers.

This shows that f is one-one onto and retains the group structure and hence $G \cong G'$

Properties of isomorphism

(i) The order of G = the order of G'

(ii) For isomorphic groups (G, o) and (G', o') the identity e' of G' is the image of identity e of G i.e. $f(e) = e'$.

If $a \in G$ and $a' \in G'$ then $a' = f(a)$

$\therefore f: G \rightarrow G'$ is one-one onto $\Rightarrow f(a) \in G' \quad \forall a \in G.$

$$\Rightarrow f(e) \in G' \quad \because e \in G$$

Now

$$aoe = a \Rightarrow f(aoe) = f(a)$$

$$\Rightarrow f(a) \cdot o' f(e) = f(a) \cdot o' e' \quad \text{by definition of isomorphism}$$

$$\Rightarrow a' \cdot o' f(e) = a' \cdot o' e'$$

$$\Rightarrow f(e) = e' \quad \text{by left cancellation law.}$$

(iii) For isomorphic groups (G, o) and (G', o') the image of inverse of any element a of G is the inverse of the image of a , i.e.

$$f(a^{-1}) = [f(a)]^{-1}$$

If e, e' are identities of G, G' respectively then by property

(ii) $f(e) = e'$

Also we have $a^{-1}oa = e = aoa^{-1} \quad \forall a \in G$

But $a^{-1}oa = e \Rightarrow f(a^{-1}oa) = f(e) \quad \forall a \in G$

$$\Rightarrow f(a^{-1}) \cdot o' f(a) = e' \quad \text{by definition of isomorphism}$$

$$\Rightarrow f(a^{-1}) = [f(a)]^{-1} \quad \text{by definition of inverse}$$

of an element in G'

(iv) For isomorphic groups (G, o) and (G', o') , the order of an element $a \in G$ is the same as the order of its image $a' \in G'$.

$f: G \rightarrow G'$ is one-one and onto.

If e, e' be identities in G, G' respectively, then

$$f(e) = e' \quad \text{and} \quad f(aob) = f(a) \cdot o' f(b) \quad \forall a, b \in G.$$

If n be the order of an element $a \in G$ then $a^n = e$

Also if m be the order of $f(a)$ then $[f(a)]^m = e'$

But

$$a^n = e \Rightarrow f(a^n) = f(e)$$

$$\Rightarrow f(aoa\ldots n \text{ times}) = e'$$

$$\Rightarrow f(a) \cdot o' f(a) \cdot o' \ldots n \text{ times} = e' \quad \text{by definition of isomorphism}$$

$$\Rightarrow [f(a)]^n = e'$$

$$\Rightarrow \text{order of } f(a) \leq n$$

$$\Rightarrow m \leq n.$$

Also,

$$[f(a)]^m = e' \Rightarrow f(a) \cdot o' f(a) \cdot o' \ldots m \text{ times} = f(e)$$

$$\Rightarrow f(aoa\ldots m \text{ times}) = f(e) \quad \text{by definition of isomorphism}$$

$$\Rightarrow f(a^m) = f(e)$$

$$\Rightarrow a^m = e \quad \because f \text{ is one-one}$$

$$\Rightarrow \text{order of } a \leq m$$

$$\Rightarrow n \leq m$$

So that $m \leq n$ and $n \leq m \Rightarrow n = m$

$\Rightarrow$ order of a = order of a' .

(v) If f is isomorphic mapping of $G \rightarrow G'$, then f^{-1} is also isomorphic.

If f is one-one and onto then f^{-1} exists and is also one-one onto.

Also if $x = f(a)$, $y = f(b)$ for $a, b \in G$ and $x, y \in G'$, then

$$a = f^{-1}(x), b = f^{-1}(y)$$

$$\text{But } f^{-1}(x \circ' y) = f^{-1}[f(a) \circ' f(b)]$$

$$= f^{-1}[f(aob)] \quad \because f \text{ is isomorphic mapping}$$

$$\Rightarrow aob \quad \because f^{-1}f(p) = p.$$

$$\Rightarrow f^{-1}(x) \circ f^{-1}(y)$$

which follows that f^{-1} retains the group structure and hence f^{-1} is isomorphic.

Automorphism of groups. An isomorphism of a group onto itself is said to be an automorphism of the group e.g. $f: G \rightarrow G'$ given by $f(a) = a^{-1}$, $a \in G$ is an automorphism iff G is an abelian group.

As another example the identity mapping $i: G \rightarrow G$ is an automorphism of group G .

In other words an automorphism f of G is a one-one transformation of G onto itself s.t. $(xy)f = (xf)(yf) \quad \forall x, y \in G$

$$\text{i.e. } f(xy) = f(x)f(y)$$

Product of Automorphisms. If $x \mapsto xf = x'$ be an automorphism of A where x' is the element of A in some order, then the mapping is automorphism and so $(xy)f = (xf)(yf) = x'y'$.

Take $x \mapsto y$ another automorphism and denote z' by $z\phi$,

$$\text{so that } (xy)f\phi = [(xy)f]\phi = [(xf)(yf)]\phi = [(xf)\phi][(yf)\phi]$$

$$= [(x)f\phi][(y)f\phi] \quad \forall x, y \in A$$

which shows that $f\phi$ is an automorphism of A and the mapping $f\phi$ is termed as product of automorphisms of f and ϕ .

The automorphism of a mathematical system forms a group.

The mapping $x \mapsto x$ is said to be the Identity automorphism in the identity element of the automorphism group. So axiom G_3 is satisfied.

G_1 is satisfied since product of two automorphisms is an automorphism.

G_2 is satisfied since if we arrange the mappings

$$f: x \mapsto x', \phi: x' \mapsto x'', \psi: x'' \mapsto x'''$$

$$\text{as } f: x \mapsto x', \phi: x' \mapsto x'', \psi: x'' \mapsto x'''$$

then $x''' = (x)f\phi\psi$ corresponding to x under the automorphism $f\phi\psi$ is uniquely determined whether it is obtained as $[(x)f]\phi\psi$ from the automorphism $f(\phi\psi)$ or as $[(x)\phi]\psi$ from $(f\phi)\psi$. Ultimately $x \mapsto x'$ is an automorphism and so $(xy)f^{-1} = [(xf^{-1}f)(yf^{-1}f)]f^{-1}$

$$= [(xf^{-1})(yf^{-1})f]f^{-1} = (xf^{-1})(yf^{-1})$$

showing that f^{-1} is an automorphism and hence G_4 is satisfied.

Conclusively the automorphisms of a mathematical system form a group.

Endomorphism of groups. A homomorphism of a group onto itself is said to be an endomorphism of the group.

Regular permutation group. A permutation group to which a group G is isomorphic is said to be a regular permutation group.

THEOREM 1. Transference of group structures.

If $(G, \circ)$ is a group and G' is a set with the multiplicative composition ' $\circ'$ ' and if $\exists$ a one-one onto mapping $f: G \rightarrow G'$ s.t. $f(aob) = f(a) \circ' f(b) \quad \forall a, b \in G$ then G' is also a group isomorphic to G for the given composition.

We have to show that G' is a group and $G' \cong G$.

Let $a' = f(a)$, $b' = f(b)$, $c' = f(c)$; $a, b, c \in G$ and $a', b', c' \in G'$ then $a' \circ' b' = f(a) \circ' f(b) = f(aob)$ is given.

G_1 is satisfied since $a, b' \in G' \Rightarrow f(a), f(b) \in G$

$$\Rightarrow a, b \in G$$

$$\Rightarrow aob \in G$$

$$\Rightarrow f(aob) \in G'$$

$$\Rightarrow f(a) \circ' f(b) \in G'$$

$$\Rightarrow a' \circ' b' \in G' \quad \forall a' b' \in G'$$

G_2 is satisfied, since $(a' \circ' b') \circ' c' = [f(a) \circ' f(b)] \circ' f(c)$

$$= f[aob(c)] \quad \because 'o' \text{ is associative}$$

$$= f(a) \circ' f(boc)$$

$$= f(a) \circ' [f(b) \circ' f(c)]$$

$$= a' \circ' (b' \circ' c')$$

G_3 is satisfied, since if e be the identity in G then

$$f(e) \circ' a' = [f(e) \circ' f(a)]$$

$$= f(eoa) = f(a)$$

$$= a'$$

$$\therefore eoa = a$$

$$\text{and } a' \circ' f(e) = f(a) \circ' f(e) = f(aoe) = f(a) = a'$$

$$\therefore f(e) \circ' a' = a' \circ' f(e) = a'$$

G_4 is satisfied, since if $a \in G$ then $a^{-1} \in G$ so that $aoa^{-1} = e = a^{-1}oa$

$$\text{and } aoe^{-1} = e \Rightarrow f(aoa^{-1}) = f(e)$$

$$\Rightarrow f(a) \circ' f(a^{-1}) = f(e)$$

$$\Rightarrow a' \circ' f(a^{-1}) = f(e)$$

$$\text{also } f(a^{-1}oa) = f(e) \Rightarrow f(a^{-1}) \circ' f(a) = f(e)$$

$$\Rightarrow f(a^{-1}) \circ' a' = f(e)$$

$$\therefore a' \circ' f(a^{-1}) = f(a^{-1}) \circ' a' = f(e).$$

Thus $f(a^{-1})$ is the inverse of $a' \in G'$ i.e. $f(a^{-1}) = (a')^{-1} = [f(a)]^{-1}$

These axioms show that G' is a group.

Again $G \cong G'$ and the relation of isomorphism is symmetric

$$\therefore G' \cong G$$

THEOREM 2. The relation of isomorphism in the set of all groups is an equivalence relation.

If G be a group belonging to the set of all groups and $x \in G$, then consider a one-one onto mapping $f: G \rightarrow G$ defined by $f(x) = x \quad \forall x \in G$.

The relation $\cong$ is reflective, since $f(x) = f(y) \Rightarrow x = y$ i.e. f is one-one and $f(xy) = xy = f(x) \cdot f(y)$, operation being multiplicative.

$\therefore$ The group structure is retained and so $G \cong G \quad \forall G \in S$, S being the set of all groups.

The relation $\cong$ is symmetric, since if f is isomorphism of G to G' , then f is one-one onto and so f^{-1} exists s.t. $f^{-1}: G' \rightarrow G$.

$\therefore$ By property (v) of isomorphism, f^{-1} is isomorphic

Thus, $G \cong G' \Rightarrow G' \cong G$

The relation $\cong$ is transitive, since if $f: G \rightarrow G'$ and $g: G' \rightarrow G''$ be two isomorphic mappings, then composite mapping $g \circ f$ is also one-one onto when $g \circ f: G \rightarrow G''$

Now $x, y \in G \Rightarrow f(x), f(y) \in G' \Rightarrow g[f(x)] \in G''$

So that $(g \circ f)(xy) = g[f(xy)]$

$= g[f(x)f(y)], f$ being isomorphic.

$\Rightarrow g f(x) g f(y), g$ being isomorphic.

i.e. $g \circ f$ retains the group compositions and also it is one-one onto, so $g \circ f$ is isomorphism and maps $G \rightarrow G''$ i.e.

$$G \cong G', G' \cong G'' \Rightarrow G \cong G''$$

Hence $\cong$ is an equivalence relation.

THEOREM 3. Cayley's Theorem.

Every finite group G of order n (say) is isomorphic with a sub-group of symmetric group S_n

or

Every finite group G of order n is isomorphic to a permutation group (or transformation group).

Let $G = \{a_1, a_2, \dots, a_n\}$ be a finite group of order n , with multiplicative composition and $a \in G$. Then n products

$a a_1, a a_2, \dots, a a_n$ are all distinct elements of G , for if possible let us assume that $a a_i = a a_j$

Let cancellation law give, $a_i = a_j$

But $a_i \neq a_j \therefore a a_i \neq a a_j$ so that $a a_1, a a_2, \dots, a a_n$ are all distinct elements of G in some order.

$\therefore$ The mapping $f_a: G \rightarrow G$ s.t. $f_a(x) = ax, a \in G, x \in G$ is one-one and onto.

Thus $f_a = \begin{pmatrix} a_1 & a_2 & \dots & a_n \\ aa_1 & aa_2 & \dots & aa_n \end{pmatrix}$ is a permutation on n symbols.

Replacing a by $a_1, a_2, \dots, a_n$ in succession, we shall have x permutations $f_{a_1}, f_{a_2}, \dots, f_{a_n}$ of which no two can be equal since if $a_1, a_2 \in G$, then $f_{a_1} = f_{a_2} \Rightarrow f_{a_1}(x) = f_{a_2}(x) \quad \forall x \in G$

$$\Rightarrow a_1 x = a_2 x \quad \forall x \in G$$

$$\Rightarrow a_1 = a_2$$

Denoting the n permutations by G' i.e.

$$G' = \{f_a : a \in G\}$$

We have

$$\begin{aligned} f_a f_b &= \begin{pmatrix} a_1 & a_2 & \dots & a_n \\ aa_1 & aa_2 & \dots & aa_n \end{pmatrix} \begin{pmatrix} a_1 & a_2 & \dots & a_n \\ ba_1 & ba_2 & \dots & ba_n \end{pmatrix} \\ &= \begin{pmatrix} a_1 & a_2 & \dots & a_n \\ aa_1 & aa_2 & \dots & aa_n \end{pmatrix} \begin{pmatrix} aa_1 & aa_2 & \dots & aa_n \\ aba_1 & aba_2 & \dots & aba_n \end{pmatrix} \\ &= \begin{pmatrix} a_1 & a_2 & \dots & a_n \\ aba_1 & aba_2 & \dots & aba_n \end{pmatrix} = f_{ab} \end{aligned}$$

But $a, b \in G \Rightarrow ab \in G$ and so $f_a, f_b \in G' \Rightarrow f_{ab} \in G'$

$\therefore$ Closure axiom is satisfied.

Now to show that the set G' with the composite composition is a group isomorphic to the given group G , let us consider a mapping

$$g: G \rightarrow G' \text{ s.t. } g(a) = f_a \quad \forall a \in G$$

Then, $g(a) = g(b) \Rightarrow f_a = f_b$

$$\Rightarrow ax = bx \quad \forall x \in G$$

$$\Rightarrow a = b \text{ by right cancellation law.}$$

So that g is one-one and therefore G, G' consist of the same number of elements. But g being one-one mapping of G to G' , g is also onto. Moreover $g(ab) = f_{ab} = f_a f_b = g(a) g(b)$

i.e. the group-composition is retained (preserved) by g .

Hence $G \cong G'$.

THEOREM 4. Every cyclic group of infinite order is isomorphic to the additive group of integers.

If G be an infinite cyclic group generated by a , then $G = \{a^i\}$ and all the powers of a are distinct.

Consider the mapping $f: G \rightarrow \mathbb{I}$ given by $f(a^i) = i$

This mapping is onto and also one-one since $i \neq j \Rightarrow a^i \neq a^j$

$$\therefore f(a^i \cdot a^j) = f(a^{i+j}) = i+j = f(a^i) + f(a^j)$$

So that f preserves the operation and hence f is an isomorphism i.e.

$$(G, \cdot) \cong (\mathbb{I}, +)$$

Problem 33. Show that the multiplicative group $G = \{I, -I, i, -i\}$ is isomorphic to the permutation group $G' = \{I, (abcd), (ac)(bd), (adcb)\}$ on four symbols.

Isomorphism of G and G' will be established if we define mapping of $G \rightarrow G'$ s.t. identity element if G is mapped to identity element of G' and inverses are mapped to inverses since then the elements of same order are mapped to elements of the same order.

In G' , the order of $(ac)(bd)$ is 2 and the order of each of $(abcd)$ and $(adcb)$ is 4.

Now

$$[(ac)(bd)]^2 = (ac)(bd)(ac)(bd) = (ac)(ac)(bd)(bd), \text{ product of disjoint cycles being abelian}$$

$$= (ac)^2 (bd)^2 = II = I \text{ as } (ac)^2 = I, (bd)^2 = I.$$

$$\text{and } (abcd)^2 = \begin{pmatrix} a & b & c & d \\ b & c & d & a \end{pmatrix} \begin{pmatrix} a & b & c & d \\ b & c & d & a \end{pmatrix} = \begin{pmatrix} a & b & c & d \\ c & d & a & b \end{pmatrix} = (ac) (bd)$$

so that $(abcd)^4 = [(ac) (bd)]^2 = I$ as above

$$\text{Similarly } (adcb)^4 = I$$

We can thus define a mapping $f: G \rightarrow G'$ given by

$$f(1) = I, f(-1) = (ac) (bd) = A \text{ (say)}, f(i) = (abcd) = B \text{ (say)}, f(-i) = (adcb) = C \text{ (say)}$$

The mapping is evidently one-one and onto. The composition tables for G and G' are as shown here.

Clearly in the Table of G if 1, -1, i , $-i$ are replaced by I, A, B, C respectively then it transforms to the Table for G' .

.	1	-1	i	-i
1	1	-1	i	-i
-1	-1	1	-i	i
i	i	-i	1	-1
-i	-i	i	-1	1

Table for G

<u>.</u>	<u>I</u>	<u>A</u>	<u>B</u>	<u>C</u>
<u>I</u>	<u>I</u>	<u>A</u>	<u>B</u>	<u>C</u>
<u>A</u>	<u>A</u>	<u>I</u>	<u>C</u>	<u>B</u>
<u>B</u>	<u>B</u>	<u>C</u>	<u>A</u>	<u>I</u>
<u>C</u>	<u>C</u>	<u>B</u>	<u>I</u>	<u>A</u>

Table for G'

Hence

$$G \cong G'.$$

Problem 34. Find the regular permutation group isomorphic to the group $G = \{a, b, c, d\}$ with the composition table.

Let G' be the required regular permutation group. Then by Cayley's theorem G' will consist of four permutations p_1, p_2, p_3, p_4 given by

.	a	b	c	d
a	a	b	c	d
b	b	a	d	c
c	c	d	a	b
d	d	c	b	a

$$p_1 = \begin{pmatrix} a & b & c & d \\ aa & ab & ac & ad \end{pmatrix} = \begin{pmatrix} a & b & c & d \\ a & b & c & d \end{pmatrix} = I \text{ (by given composition table)}$$

$$p_2 = \begin{pmatrix} a & b & c & d \\ ba & bb & bc & bd \end{pmatrix} = \begin{pmatrix} a & b & c & d \\ b & a & d & c \end{pmatrix} = \begin{pmatrix} a & b \\ b & a \end{pmatrix} \begin{pmatrix} c & d \\ d & c \end{pmatrix} = (ab) (cd)$$

$$p_3 = \begin{pmatrix} a & b & c & d \\ ca & cb & cc & cd \end{pmatrix} = \begin{pmatrix} a & b & c & d \\ c & d & a & b \end{pmatrix} = \begin{pmatrix} a & c \\ c & a \end{pmatrix} \begin{pmatrix} b & d \\ d & b \end{pmatrix} = (a c) (b d)$$

$$p_4 = \begin{pmatrix} a & b & c & d \\ da & db & dc & dd \end{pmatrix} = \begin{pmatrix} a & b & c & d \\ d & c & b & a \end{pmatrix} = \begin{pmatrix} a & d \\ d & a \end{pmatrix} \begin{pmatrix} b & c \\ c & b \end{pmatrix} = (a d) (b c)$$

$$\text{Hence } G' = \{p_1, p_2, p_3, p_4, o\} \\ = \{I, (ab) (cd), (ac) (bd), (ad) (bc)\}, o\}$$

4.10. NORMAL AND CONJUGATE SUBGROUPS

Conjugate elements. Given a group G , an element $a \in G$ is known as the conjugate to another element $b \in G$ if $\exists$ an element $x \in G$, s.t. $a = x^{-1}bx$.

If $a = x^{-1}bx$ then a is sometimes known as the transform of b by x .

Inner and Outer Automorphisms. If a be a fixed element of a group G then the conjugation $C_a: x \rightarrow a^{-1}xa$ is said to be the **Inner automorphism** and all other automorphisms are **outer**. Clearly $(a^{-1}xa)(a^{-1}ya) = a^{-1}(xy)a$ for all x, y .

e.g., the cyclic group of order 3 has no inner automorphism except the identity while it has outer automorphism $x \leftrightarrow x^2$

The inner automorphisms of any group G form an automorphism of G .

If C_a and C_b be two inner automorphism, then

$$C_a \cdot C_b = C_{ab} \text{ since } b^{-1}(a^{-1}xa)b = (ab)^{-1}x(ab)$$

where

$$C_a: x \rightarrow a^{-1}xa$$

$$C_b: x \rightarrow b^{-1}xb$$

Similarly since $(a^{-1})^{-1}(a^{-1}xa)(a^{-1}) = x$

$\therefore$ the inverse of the conjugation C_a is C_a^{-1}

Properties of conjugate elements

(i) **Conjugacy is reflexive** i.e. every element is conjugate with itself.

We have $a = x^{-1}bx$; $a, b, x \in G$

... (1)

If e be the identity element in G , then $e \in G$ and $e^{-1} = e$.

Thus replacing x by e in (1) we get $e^{-1}be = e^{-1}b = b$

$\therefore b = e^{-1}be \Rightarrow b$ is conjugate with itself.

(ii) **Conjugacy is symmetric** i.e. if a is conjugate with b , then b is conjugate with a .

We have $a = x^{-1}bx$; $a, b, x \in G$

$\therefore a = x^{-1}bx \Rightarrow xa = x(x^{-1}bx)$

$$\Rightarrow xa = ebx$$

$$\therefore x(x^{-1}bx) = (xx^{-1})bx = ebx$$

$$\Rightarrow xax^{-1} = (bx)x^{-1}$$

$$\therefore ebx = b$$

$$\Rightarrow xax^{-1} = be$$

$$\therefore (bx)x^{-1} = b(xx^{-1}) = be = b$$

$$\Rightarrow xax^{-1} = b$$

So that

$$a = x^{-1}bx \Rightarrow b = (x^{-1})^{-1}a(x^{-1}), x^{-1} \in G$$

Which shows that b is conjugate with a .

(iii) **Conjugacy is transitive** i.e. if a is conjugate with b , b is conjugate with c , then a is conjugate with c .

We have

$$a = x^{-1}bx \text{ and } b = y^{-1}cy, a, b, c, x, y \in G$$

$$\therefore a = x^{-1}bx, b = y^{-1}cy \Rightarrow a = x^{-1}(y^{-1}cy)x$$

$$\Rightarrow a = (x^{-1}y^{-1})c(yx)$$

$$\Rightarrow a = (yx)^{-1}c(yx) \text{ by reversal law of inverse}$$

$$\text{as } x, y \in G \Rightarrow xy \in G$$

$$\Rightarrow a \text{ is conjugate with } c.$$

Note 1. The above properties (i), (ii), (iii) when combined, show that conjugacy is an equivalence relation on G .

(iv) If a is conjugate with b and c both, then, b and c are conjugate with each other.

We have

$$\begin{aligned} a &= x^{-1}bx \text{ and } a = y^{-1}cy \\ \therefore a &= x^{-1}bx, a = y^{-1}cy \Rightarrow x^{-1}bx = y^{-1}cy \\ &\Rightarrow x(x^{-1}bx) = x(y^{-1}cy) \\ &\Rightarrow bx = (xy^{-1})(cy) \\ &\quad \therefore x(x^{-1}bx) = (xx^{-1})bx = ebx = bx \\ &\Rightarrow (bx)x^{-1} = (xy^{-1})(cy)x^{-1} \\ &\Rightarrow b = (xy^{-1})c(yx^{-1}) \\ &\quad \therefore (bx)x^{-1} = b(xx^{-1}) = be = b \end{aligned}$$

Now since $x, y \in G \Rightarrow x^{-1}, y^{-1} \in G$ and so $xy^{-1}, yx^{-1} \in G$.

Hence b is conjugate with c .

Class of the group. It is observed that there may be more than two elements of a group, which are conjugate with one another. The entire set of elements $a_1, a_2, a_3, \dots, a_h$ which are conjugate with one another is termed as a *class of the group*.

If a single element of a class of a group is given then the whole class may be determined. e.g. if the elements of a group G are $a_1 (= e), a_2, a_3, \dots, a_h$ then the class of A may be determined by forming the sequence

$$e^{-1}ae = a_2^{-1}aa_2, a_3^{-1}aa_3, \dots, a_h^{-1}aa_h.$$

Evidently all the elements of this sequence being conjugate to one another form a class.

Properties of classes

- (1) Every element conjugate can be divided in to classes.
- (2) Every element will appear in one and only class.
- (3) The identity element of a group being not conjugate to any other element, forms a group by itself since $x^{-1}ex = e \forall x \in G$.
- (4) No class can be a subgroup unless it contains only the identity element e .
- (5) Every element of an abelian (commutative) group being conjugate with itself since $ax = xa \Rightarrow x^{-1}ax = x^{-1}xa$

$$\Rightarrow x^{-1}ax = ea = a \forall x \in G$$

the class of an abelian group consists of a single element.

- (6) All the elements of a class have the same order.

If a be an element of a class then $x^{-1}ax$ will also be one element of it. Taking n as order of a , we have $a^n = e$, e being identity element.

$$\begin{aligned} \therefore (x^{-1}ax)^n &= (x^{-1}ax)(x^{-1}ax)^{n-1} \\ &= x^{-1}ax \cdot x^{-1}ax \cdot (x^{-1}ax)^{n-2} \\ &= a^2(x^{-1}ax)^{n-2} \dots \end{aligned}$$

$$= a^4(x^{-1}ax)^{n-4}$$

$$= \dots$$

$$= a^n = e$$

which follows that order of $x^{-1}ax$ is also n .

Hence the order of all the elements of a class is the same.

As an example consider the group G of matrices and divide the elements of this group into classes s.t. the matrices A and B belong to the same class.

We have

$$A = C^{-1}BC^{-1}, A, B, C \in G.$$

$$\text{Trace } A \text{ or } tr(A) = tr[C^{-1}BC^{-1}]$$

$$= \sum_p [C^{-1}BC^{-1}]_{pp}$$

$$= \sum_p [\sum_{qr} (C^{-1})_{pq} (B)_{qr} (C)_{rp}]$$

$$= \sum_p [(\sum_q (C^{-1})_{pq}) (\sum_r (B)_{qr}) (C)_{rp}]$$

$$= \sum_q \sum_r [(\sum_p (C)_{rp}) (C^{-1})_{pq} (B)_{qr}]$$

$$= \sum_q \sum_r [CC^{-1}]_{rq} [B]_{qr}$$

$$= \sum_q \sum_r (I)_{rq} [B]_{qr}, I \text{ being unite matrix s.t.}$$

$$\begin{aligned} |I|_{rq} &= 0 \text{ for } r \neq q \\ &= 1 \text{ for } r = q \end{aligned}$$

$$\text{Hence } r = q, tr(A) = \sum_q (B)_{qq} = tr(B)$$

which shows that all the matrices forming a class have the same trace.

Note 2. This is a result analogous to the property (6).

Note 3. If H be a subgroup of a group G and $x \in G$, then $K = x^{-1}Hx$ is a subgroup of G .

Let $H = \{h_1, h_2, \dots, h_i, \dots\}$ and $x^{-1}h_i x$ and $x^{-1}h_j x$ be elements of the set $K \subset G$. Then to show that K is a subgroup of G , it is sufficient to show that $(x^{-1}h_i x)(x^{-1}h_j x) \in K$.

We have

$$(x^{-1}h_i x)(x^{-1}h_j x)^{-1} = (x^{-1}h_i x)(x^{-1}h_j^{-1}x)$$

by reversal law of inverses and since $(x^{-1})^{-1} = x$

$$= x^{-1}h_i (xx^{-1})h_j^{-1}x$$

$$= x^{-1}h_i e h_j^{-1}x$$

$$= x^{-1}h_i h_j^{-1}x$$

$$= x^{-1}hx \quad \therefore h_i, h_j^{-1} \in H \Rightarrow h_i h_j^{-1} \in H \text{ and put } h = h_i h_j^{-1}$$

Hence $x^{-1}hx \in K$.

So K is a subgroup of G .

Conjugate subgroups. If $x, y, z, \dots$ be the elements of a group G i.e. $x, y, z, \dots \in G$, then the subgroups $H, x^{-1}Hx, y^{-1}Hy, z^{-1}Hz, \dots$, are known as the conjugate subgroups of G .

Normal Subgroups (or Normal divisor or Invariant subgroup or Self-conjugate subgroup). A subgroup H of a group G is said to be a normal subgroup of G if $\forall x \in G, x^{-1}Hx = H$ or equivalently, if $Hx = xH \forall x \in G$.

Properties of normal subgroups

(a) If e be the identity in G , then the whole group G and $\{e\}$ are normal subgroups of G

(b) Every subgroup H of a commutative group G is normal since a left coset xH is the same as the right coset Hx since

$$x \in H \Rightarrow x^{-1}Hx = H \quad \forall x \in G.$$

Every subgroup of an abelian group is a normal subgroup, since

$$a^{-1}xa = a^{-1}ax = x \quad \forall a, x \in G.$$

(c) The alternating group A_n is an invariant subgroup of the symmetric group S_n .

Since if E be an element of A_n i.e. $E(X) = X$, then we have to show that

$$P \in S_n \Rightarrow P^{-1}EP \in A_n$$

If P is even, then $P^{-1}EP$ is even and hence is an element of A_n .

If P is odd, then $P(X) = -X$ or $P^{-1}(-X) = X$

$$\text{and } (P^{-1}EP)(X) = P^{-1}E[P(X)] = P^{-1}E(-X) = P^{-1}(-X) = X$$

which follows that $P^{-1}EP \in A_n$ and hence A_n is a normal subgroup.

(d) The intersection of any two normal subgroups of a group is a normal subgroup.

If H_1, H_2 , be two normal subgroups of G and $a \in H_1 \cap H_2$, then

$$a \in H_1 \cap H_2 \Rightarrow a \in H_1, a \in H_2$$

But H_1, H_2 being normal subgroups,

$$x^{-1}ax \in H_1 \text{ and } x^{-1}ax \in H_2 \quad \forall x \in G$$

These imply that $x^{-1}ax \in H_1 \cap H_2$

$$\text{Hence } a \in H_1 \cap H_2 \Rightarrow x^{-1}ax \in H_1 \cap H_2 \quad \forall x \in G$$

and also $H_1 \cap H_2$ is a subgroup of G

$\therefore H_1 \cap H_2$ is a normal subgroup.

Factor group or quotient group. If H be a normal subgroup of a group G , then the group of all cosets of H in G is known as Factor group or quotient group of G by H and denoted by G/H .

Properties of factor group

(a) The order of a factor group G/H is equal to the index of H in G .

(b) Each quotient group of an abelian group is abelian but its converse is not true, since

$$(Hx)(Hy) = H(xy) = H(yx) = H(y)H(x)$$

and if S_3 be a symmetric group and A_3 an alternating group each of degree 3 then S_3/A_3 is an abelian group of degree 3 whereas S_3 is not abelian. The group S_3/A_3 is of order 2 and so it is abelian as every group of order 2 is abelian.

Problem 35. If H be a subgroup of the group $(\mathbb{I}, +)$, $\mathbb{I}$ being set of integers, s.t. $H = \{mx : x \in \mathbb{I}\}$ where m is a fixed integer, then find the elements of the quotient group $\mathbb{I}/H$ and mention the composition table for $\mathbb{I}/H$ for $m=5$.

Clearly $(\mathbb{I}, +)$ is an abelian group, therefore by the properties of normal subgroups, H is a normal subgroup. The elements of $\mathbb{I}/H$, which are cosets of H in $\mathbb{I}$ may be given as follows:

$$H+0 = H = \{\dots, -3m, -2m, -m, 0, m, 2m, \dots\}$$

$$H+1 = \{\dots, -3m+1, -2m+1, -m+1, 1, m+1, 2m+1, \dots\}$$

$$H+(m-1) = \{\dots, -2m-1, -m-1, -1, m-1, 2m-1, 3m-1, \dots\}$$

Then $\mathbb{I}/H$ has n distinct cosets as its elements.

When $m=5$, the cosets are

$$H, H+1, H+2, H+3, H+4.$$

The composition table is as shown here.

	H	$H+1$	$H+2$	$H+3$	$H+4$
H	H	$H+1$	$H+2$	$H+3$	$H+4$
$H+1$	$H+1$	$H+2$	$H+3$	$H+4$	H
$H+2$	$H+2$	$H+3$	$H+4$	H	$H+1$
$H+3$	$H+3$	$H+4$	H	$H+1$	$H+2$
$H+4$	$H+4$	H	$H+1$	$H+2$	$H+3$

Problem 36. If $G = \{e, a, a^2, a^3, a^4, a^5\}$ be a cyclic group of order 6 such that every subgroup of a cyclic group is normal, then if $H = \{e, a^3\}$ be a subgroup of G , find the elements of G/H and show that it is a group.

Elements of G/H are $He = \{e, a^3\}$

$$Ha = \{e, a^3\} a = \{a, a^4\}$$

$$Ha^2 = \{e, a^3\} a^2 = \{a^2, a^5\}$$

This is easy to show that G/H is a group and its composition table is as shown here.

Evidently G/H is a cyclic group generated by $\{a, a^4\}$.

	$\{e, a^3\}$	$\{a, a^4\}$	$\{a^2, a^5\}$
$\{e, a^3\}$	$\{e, a^3\}$	$\{a, a^4\}$	$\{a^2, a^5\}$
$\{a, a^4\}$	$\{a, a^4\}$	$\{a^2, a^5\}$	$\{e, a^3\}$
$\{a^2, a^5\}$	$\{a^2, a^5\}$	$\{e, a^3\}$	$\{a, a^4\}$

4.11. COMPLEXES AND KERNEL

Complex of a group. A non-empty subset H of a group G is called as a complex of the group G .

Properties of complexes. (i) If Z be a complex containing the elements a, b, c of a group G then $Z = \{a, b, c\}$

(ii) If $Z = \{a, b, c\}$ be a complex then $aZ = \{a^2, ab, ac\}$ etc.

(iii) If Z_1 and Z_2 be two complexes of a group G , then the product of Z_1, Z_2 is defined as

$$Z_1 Z_2 = \{x : x = z_1 z_2, z_1 \in Z_1, z_2 \in Z_2\}$$

Now since $z_1 \in Z_1$, $z_2 \in Z_2$ and $Z_1, Z_2 \subset G$

$\therefore z_1 z_2 = x \in G$ by closure axiom.

As such $Z_1 Z_2 \subset G$.

Which follows that $Z_1 Z_2$ is also a complex of G , obtained by multiplying every element in Z_1 with every element in Z_2 .

(iv) The subgroup H of a group G also gives a complex s.t. $HH = H^2 = H$.

(v) A group can be expressed as a sum of complexes.

If $x \in G$ and $x \notin H$, H being a subgroup of G , then the complex Hx is a right coset and xH is a left coset of H in G . But cosets are not groups and they are complexes, therefore if the group G as a whole is capable of forming a complex Z which consists of all the elements of the group, then we have

$$Z = H + Hx + Hy + \dots$$

(vi) The number of complexes in a group is equal to the index of a subgroup H in G and in fact it is the order of the group divided by the order of the subgroup H .

(vii) The product of complexes is associative.

Let Z_1, Z_2 and Z_3 be three complexes of a group G and let

$z_1 \in Z_1, z_2 \in Z_2, z_3 \in Z_3$, then

$z_1 \in Z_1, z_2 \in Z_2 \Rightarrow z_1 z_2 \in Z_1 Z_2$

$\therefore z_1 z_2 \in Z_1 Z_2, z_3 \in Z_3 \Rightarrow (z_1 z_2) z_3 \in (Z_1 Z_2) Z_3$

$$\Rightarrow z_1 z_2 z_3 \in (Z_1 Z_2) Z_3$$

But $z_1 z_2 z_3 = z_1 (z_2 z_3) \therefore z_1 z_2 z_3 \in Z_1 (Z_2 Z_3)$

Thus $z_1 z_2 z_3 \in (Z_1 Z_2) Z_3 \Rightarrow z_1 z_2 z_3 \in Z_1 (Z_2 Z_3)$

$\therefore (Z_1 Z_2) Z_3 \subset Z_1 (Z_2 Z_3)$

Similarly $Z_1 (Z_2 Z_3) \subset (Z_1 Z_2) Z_3$

So that $(Z_1 Z_2) Z_3 = Z_1 (Z_2 Z_3)$

Inverse of complex. If Z be a complex of a group G , then its inverse is given by $Z^{-1} = \{z^{-1} : z \in Z\}$

In other words the inverse of a complex Z is the set of inverses of all elements of Z .

Properties of inverse of a complex

(1) If $Z_1 Z_2$ be two complexes of a group G , then $(Z_1 Z_2)^{-1} = Z_2^{-1} Z_1^{-1}$

And $x \in (Z_1 Z_2)^{-1} \Rightarrow x = (z_1 z_2)^{-1}$ for $z_1 \in Z_1, z_2 \in Z_2$

$\Rightarrow x = z_2^{-1} z_1^{-1}$ by reversal law of inverses.

$\Rightarrow x \in Z_2^{-1} Z_1^{-1}$ by definition

$\therefore (Z_1 Z_2)^{-1} \subset Z_2^{-1} Z_1^{-1} \dots (A)$

Similarly if $y \in Z_2^{-1} Z_1^{-1} \Rightarrow y = z_2^{-1} z_1^{-1}$ when $z_2 \in Z_2, z_1 \in Z_1$

$\Rightarrow y = (z_1 z_2)^{-1}$ where $z_1 \in Z_1, z_2 \in Z_2$

$\Rightarrow y \in (Z_1 Z_2)^{-1}$ by definition

$\therefore Z_2^{-1} Z_1^{-1} \subset (Z_1 Z_2)^{-1} \dots (B)$

(A) and (B) follow that $(Z_1 Z_2)^{-1} = Z_2^{-1} Z_1^{-1}$

(2) If H be a subgroup of a group G , then $H^{-1} = H$.

And $h^{-1} \in H^{-1} \Rightarrow h \in H$

$\Rightarrow h^{-1} \in H$, H being a group

So $H^{-1} \subset H$

Similarly an $h \in H \Rightarrow h^{-1} \in H$, H being a group

$\Rightarrow h = (h^{-1})^{-1} \in H$ by definition of inverse of a complex.

So $H \subset H^{-1}$

$\therefore H^{-1} \subset H, H \subset H^{-1} \Rightarrow H^{-1} = H$.

(3) If H, K be two subgroups of a group G , then HK is also a subgroup of G iff $HK = KH$.

Taking $HK = KH$, we have $(HK)^{-1} = (KH)^{-1}$
 $= K^{-1} H^{-1}$ by Property (1)

$= KH$ by Property (2)

$= HK \therefore HK = KH$

Which shows that HK is a subgroup of G .

Again taking HK as subgroup of G , we have

$(HK)^{-1} = HK$ by Property (2)

$\therefore K^{-1} H^{-1} = HK$ by Property (1)

i.e. $KH = HK$ by Property (2)

Hence the proposition.

(4) A necessary and sufficient condition for a complex H of a group G to be a subgroup is that $HH^{-1} = H$.

The condition is necessary since if H is a subgroup of G and $ab^{-1} \in HH^{-1}$ then

$$a \in H, b \in H \Rightarrow a \in H, b^{-1} \in H \\ \Rightarrow ab^{-1} \in H$$

So $ab^{-1} \in HH^{-1} \Rightarrow ab^{-1} \in H, b \in H, b^{-1} \in H^{-1}$

i.e. $HH^{-1} \subset H$

Also H is a subgroup of $G \Rightarrow$ identity $e \in H$

If $h \in H$, then $h = he = he^{-1} \in HH^{-1}, h \in H, e^{-1} \in H^{-1}$

$\therefore H \subset HH^{-1}$

Thus $HH^{-1} \subset H, H \subset HH^{-1} \Rightarrow HH^{-1} = H$

The condition is sufficient since if $HH^{-1} = H$, then we have

$$HH^{-1} \subset H$$

Now suppose that $a, b \in H$ so that $ab^{-1} \in HH^{-1}$

$\therefore HH^{-1} \subset H$ and $ab^{-1} \in HH^{-1} \Rightarrow ab^{-1} \in H$

Ultimately $a \in H, b \in H \Rightarrow ab^{-1} \in H$

Which follows that H is a subgroup as is evident from the following discussion: Taking H a subgroup of G with the same composition as in G , the identity in H and G is the same. Also $a \in H$ and $b \in H$ give $b^{-1} \in H, H$ being a group.

$\therefore a \in H, b^{-1} \in H \Rightarrow ab^{-1} \in H \dots (a)$

Further taking H to be a non-empty subset of G s.t. $a \in H, b \in H$, and assuming that $a \in H, b \in H \Rightarrow ab^{-1} \in H$, we observe that H is non-empty and $\exists$ an $a \in H$ so that by setting $b=a$ in (α), we find

$$a \in H, a \in H \Rightarrow aa^{-1} \in H$$

$$\Rightarrow e \in H, e \text{ also being identity in } G.$$

$$\text{Now } e \in H, b \in H \Rightarrow eb^{-1} \in H \text{ by } (\alpha) \quad \dots (\beta)$$

$$\Rightarrow b^{-1} \in H$$

$$\therefore a \in H, b^{-1} \in H \Rightarrow a(b^{-1})^{-1} \in H \Rightarrow ab \in H$$

$$\text{i.e. } a \in H, b \in H \Rightarrow ab \in H$$

Here (α) and (β) fulfil the requirements for H which is a complex of G , to be its subgroup

Image of a group G under a mapping f . If $f: G \rightarrow G'$ be a homomorphism of a group G into a group G' , then $f(G) = \{f(x), x \in G\}$ is a subset of G' and is termed as the **Image of G under f** and denoted by $Im(f)$.

Kernel of f . If $f: G \rightarrow G'$ be a homomorphism of G into G' , then the subset of those elements of G which are mapped onto the identity of G' under f is said to be the **Kernel of f** and denoted by $ker(f)$ or $f^{-1}(e')$.

$$\text{i.e. } ker(f) = \{x \in G : f(x) = e'\}$$

Propositions relating to Kernel

I. A homomorphism $f: G \rightarrow G'$ is an isomorphism iff $ker f = \{e\}$.

Assuming that $f: G \rightarrow G'$ is an isomorphism, if $a \in ker f$ then

$$f(a) = e' = f(e), e' \text{ being identity in } G'.$$

Now f being one-one and $a=e$, kernel of f consists of e only. Conversely if $ker f = \{e\}$ for f to be homomorphism, and if $a, b \in G$ s.t. $f(a) = f(b)$, then $f(ab^{-1})$

$$= f(a)f(b^{-1})$$

$$= f(a)[f(b)]^{-1}$$

$$= e'$$

$$\therefore f(a) = f(b)$$

$$\therefore ab^{-1} \in ker f$$

$$\text{or } ab^{-1} = e$$

$$\text{or } a = b$$

So f is one-one and hence f is an isomorphism.

II. If f be homomorphism of G' then $ker(f)$ is an invariant subgroup of G .

If $a, b \in ker(f)$, then $f(a) = e' = f(b)$, e' being identity of G' .

$$\therefore f(ab) = f(a)f(b) = e'e' = e'$$

which implies that $ab \in ker(f)$ i.e. closure axiom is satisfied.

Now $ker(f)$ being a subset of G , associativity axiom is self-evident.

Again $f(e) = e' \Rightarrow e \in ker(f)$, e being identity in G .

$\therefore$ There exists an identity in G .

Further if $a \in ker(f)$ then $f(a^{-1}) = [f(a)]^{-1} = (e')^{-1} = e'$

which shows that $a^{-1} \in ker(f)$ when $a \in ker(f)$

This follows the existence of an inverse in G .

As such $ker(f)$ is a subgroup of G , as $ker(f)$ satisfies all the four group axioms.

Moreover $ker(f)$ is an invariant subgroup of G as is shown below:

If $g \in G$ and $h \in ker(f)$, then

$$f(g^{-1}hg) = f(g^{-1})f(h)f(g)$$

$$= [f(g)]^{-1}e'f(g)$$

$$\therefore h \in ker(f) \Rightarrow f(h) = e'$$

$$= [f(g)]^{-1}f(g)$$

$$= e'$$

$$\therefore g^{-1}hg \in ker(f).$$

Hence $ker(f)$ is an invariant subgroup of G .

Note 1. It is easy to show that $Im(f)$ is a subgroup of G' .

III. If H be a normal subgroup of a group G , then there is a homomorphism of G onto G/H .

Let $f: G \rightarrow G/H$ be given by $f(x) = Hx \forall x \in G$

$\therefore \forall x \in G, \exists$ a unique coset Hx , f is a mapping.

Also the binary operation in G/H being defined by

$$(Hx)(Hy) = H(xy)$$

We have

$$f(xy) = H(xy) = (Hx)(Hy) = f(x)f(y)$$

Which follows that f is a homomorphism and it is onto since every coset $Hx \in G/H$ has x as its preimage in G .

Note 2. Natural Homomorphism. The homomorphism $f: G \rightarrow G/H$ given by $f(x) = Hx$ is known as **Natural Homomorphism or Canonical Homomorphism** of G onto G/H .

IV. If f be a homomorphism of a group G onto a group G' with kernel K , then

$$G/K \cong G'$$

Consider the mapping $\phi: G/K \rightarrow G'$ defined by $\phi(Kx) = f(x)$

Taking $Kx = Ky$, we have $xy^{-1} \in K$ and $f(xy^{-1}) = e'$, e' being identity in G' i.e. $f(x)f(y)^{-1} = e'$

$$\text{or } f(x)[f(y)]^{-1} = e'$$

$$\text{or } f(x) = f(y).$$

This follows that ϕ is uniquely defined.

Now if $f(y) \in G'$ then Ky is the preimage of $f(y)$ in G/K under ϕ .

This follows that ϕ is onto.

Again ϕ will be one-one if $Kx = Ky$ provided $f(x) = f(y)$.

Take an element $z = xy^{-1} \in G$ i.e. $zy = x$

$$\begin{aligned}\therefore f(z) &= f(xy^{-1}) = f(x)f(y^{-1}) \\ &= f(x)[f(y)]^{-1} \\ &= e'\end{aligned}$$

$$\therefore f(x) = f(y)$$

So that $z \in K$ and $Kx = K(z y) = (Kz) y = Ky$

$\therefore \phi$ is one-one.

Further to show that ϕ preserves the structure, we have

$$\phi(Kx) \phi(Ky) = f(x) f(y) = f(xy) = \phi[K(xy)] = \phi[(Kx)(Ky)]$$

Hence ϕ is isomorphism and thus $G/K \cong G'$.

V. If f is a homomorphism from the group (G, o) into the group (G', o') then the pair $(\ker f, o)$ is a normal subgroup of (G, o) .

Evidently $\ker f \neq \phi$ (non empty) since $e \in \ker f$ and $\ker f \subset G$

Now $a, b \in \ker f \Rightarrow f(a) = e', f(b) = e'$

But $f(b^{-1}) = [f(b)]^{-1} = [e']^{-1} = e'$

$$\therefore f(aob^{-1}) = f(a)o[f(b)]^{-1} = e'oe' = e'$$

$$\therefore a, b \in \ker f \Rightarrow aob^{-1} \in \ker f$$

Hence $(\ker f, o)$ is a subgroup.

Again $\forall a \in G$ and $h \in \ker f$, we have

$$\begin{aligned}f(aoh a^{-1}) &= f(a) o f(h) o f(a^{-1}) \\ &= f(a) o f(h) o [f(a)]^{-1} \\ &= f(a) oe' o [f(a)]^{-1} \\ &= f(a) o [f(a)]^{-1} \\ &= e'\end{aligned}$$

$$\therefore \forall a \in G \text{ and } h \in \ker f \Rightarrow aoh a^{-1} \in \ker f$$

Hence $(\ker f, o)$ is a normal subgroup.

Note 3. Similarly it can be shown that image of $[Im(f), o]$ is a subgroup of (G', o') when f is a homomorphism of (G, o) into (G', o') .

Problem 37. If $GL(n, R)$ is the multiplication group of all $n \times n$ singular matrices with elements as real numbers and that G' is the multiplicative group of all non-zero real numbers, then show that the mapping $f: G \rightarrow G'$ s.t. $f(A) = |A|$ $\forall A \in G$ is a homomorphism of G onto G' and also show that

$$\ker f = \{A \in GL(n, R) : |A| = e', \text{ the identity in } G'\}.$$

Let $f: (C, +) \rightarrow (R^+, +)$ s.t. $f(x+iy) = x$.

We have to show that f is a homomorphism of $(C, \cdot)$ onto $(R, \cdot)$ and $\ker f = \{z \in C : x=0\}$ i.e. $\ker f$ is the imaginary y-axis.

If $z_1 = x_1 + iz_1 \in C, z_2 = x_2 + iy_2 \in C$, then

$$f(x_1 + iy_1) + f(x_2 + iy_2)$$

$$= f(x_1 + x_2) + if(y_1 + y_2)$$

$$= x_1 + x_2 \text{ by hypothesis}$$

$$= f(x_1 + iy_1) + f(x_2 + iy_2)$$

This shows that f is onto since $f(x+io) = x$ if $x \in R$.




Fig. 4.11

As such f is a homomorphism of $(C^*, +)$ onto $(R^+, +)$ and 1 is the identity element in $(R^+, \cdot)$

Also $\ker f$ is given by $f(x+iy) = 0 = x \quad \forall x \in R$.
i.e. $\ker f = \{z \in C : x=0=e', \text{ the identity in } R\}$

which follows that $\ker f$ is the imaginary axis.

Problem 38. If $(R, \cdot)$ be a multiplicative group and $x \in R$, then find homomorphisms and their kernels in the following mappings

$$(i) x \rightarrow |x|$$

$$(ii) x \rightarrow \frac{1}{x^2}$$

$$(i) x \rightarrow |x| \Rightarrow x \rightarrow |x| \text{ and } -x \rightarrow |x|$$

$$\therefore x \rightarrow |x| \text{ and } y \rightarrow |y| \Rightarrow xy \rightarrow |xy| \rightarrow |x| |y|$$

Thus $x \rightarrow |x|$ is homomorphism.

$$\begin{aligned}\text{Now } f: x \rightarrow |x| \Rightarrow f(x) &= |x| \Rightarrow f(xy) = |xy| \Rightarrow f(xy) = |x| |y| \\ &= |x| |y| = f(x) f(y)\end{aligned}$$

i.e. f is two-one mapping since $|f(x)| \Rightarrow f(x) = -1, 1$.

Its kernel is $|f(x)| = 1 \Rightarrow -1, +1$, i.e. $\{-1, 1\}$

$$(ii) \text{ Say } g: x \rightarrow \frac{1}{x^2} \Rightarrow g(x) = \frac{1}{x^2} \Rightarrow g(xy) = \frac{1}{(xy)^2} = \frac{1}{x^2 y^2} = \frac{1}{x^2} \cdot \frac{1}{y^2} = g(x) \cdot g(y)$$

So g is a homomorphism and it is two-one mapping since

$$g(x) = \frac{1}{x^2} \Rightarrow g(-x) = \frac{1}{x^2}$$

$$\text{Now } g(1) = e' \Rightarrow \frac{1}{x^2} = 1 \Rightarrow x = -1, 1$$

$$\therefore \ker g = \{-1, 1\}$$

4.12. GROUPS OF ISOMETRIES

Let R be the set of real numbers and S_R be the symmetric group on R . Then $I(R)$ the group of isometries of R is a subgroup of S_R and defined to be the set of all elements of S_R which preserve distance (distance between two points $a, b \in R$ is the absolute value $|a-b|$ of $a-b$ and denoted by $d(a, b)$ and the elements of such a set are known as isometries of R .

An element $\sigma \in S_R$ is called an *Isometry* iff $d(a, b) = d(a\sigma, b\sigma) \quad \forall (a, b) \in R$ semi the identity mapping $i \in I(R), I(R) \neq \phi$.

To show that $I(R)$ the group of isometries of R is a subgroup of R , suppose that $\sigma \in I(R)$ so that $\sigma^{-1} \in S_R$ as S_R is a group and $\sigma \in S_R$. Then we have to show that $\sigma^{-1} \in I(R)$.

If $a, b \in R$ then σ being an isometry, $d(a\sigma^{-1}, b\sigma^{-1}) = d[(a\sigma^{-1})\sigma, (b\sigma^{-1})\sigma] = d(a, b)$

$$\text{Thus } d(a, b) = d(a\sigma^{-1}, b\sigma^{-1}) \Rightarrow \sigma^{-1} \in I(R)$$

If $\sigma, \tau \in I(R)$ then $\tau^{-1} \in I(R)$ and

$$d[a(\sigma\tau^{-1}), b(\sigma\tau^{-1})] = d[(a\sigma)\tau^{-1}, (b\sigma)\tau^{-1}] = d(a\sigma, b\sigma) = d(a, b)$$

So that $\sigma\tau^{-1} \in I(R)$ and hence $I(R)$ is a subgroup of S_R .

If $\sigma, \tau \in I(R)$ have the same effect on two distinct real numbers a and b i.e. $a\sigma = a\tau$ and $b\sigma = b\tau$ then $\sigma = \tau$

This version is used to describe the elements of $I(R)$.


Geometrically interpreted, if $r\sigma = r + a$ where $\sigma \in I(R)$, $r \in R$ and $a\sigma = a$, then it moves the real line a units to the right and if $r\sigma = -r + a$, the real line is inverted about the origin and then moved a units to the right.

If E be the set $R^2 = K \times R$ and $(x_A, y_A) = A$, $(x_B, y_B) = B$ are two elements of E then distance $d(A, B) = \sqrt{(x_A - y_B)^2 + (y_A - y_B)^2}$


$\sigma \in S_E$ the symmetric group on E is known as an isometry if for $A, B \in E$, $d(A, B) = d(A\sigma, B\sigma)$.

The set I of all isometries of E forms a subgroup of S_E .

Assuming the Euclidean plane E covered by an infinite rigid metal



Initially
Fig. 4.12



After a movement
Fig. 4.13


lamina S let $P, Q, R, \dots$ be the points of E and $A, B, C, \dots$ be the points of S initially as shown in Fig. 4.12 and after a movement as shown in Fig. 4.13.

Define $\theta : E \rightarrow E$ as an isometry given by


$$P\theta = P_1, Q\theta = Q_1, R\theta = R_1 \text{ where } d(A, B) = d(P_1, Q_1) = d(P, Q).$$

As such we have three particular isometries.

(i) **Rotation about a point.** Take a point O of S and rotate S about



A on the top of P
Fig. 4.14



A on the top of P1
Fig. 4.15

O through an angle ψ . Such an isometry induced by the movement of S is the rotation about O through an angle ψ .

(ii) **Reflection in a line.** Choosing a line in E , turn S over this line and back to E . Such an isometry is the reflection in XY .




Fig. 4.16




Fig. 4.17

(iii) **Translation.** Choosing a line XY and an isometry corresponding to a movement of S s.t. the line $X\theta Y\theta$ is parallel to XY , θ is the translation.

In terms of mapping. Translation is the mapping

$$\tau_{a,b} : (xy) \tau_{a,b} = (x+a, y+b)$$

which is isometry for each a, b , $\tau_{a,b}$ and $(\tau_{a,b})^{-1} = \tau_{-a,-b}$

Counter clockwise rotation about the origin through an angle θ is the mapping $\rho_\theta : (x, y) \rho_\theta = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta)$ which is isometry for each θ , ρ_θ and $(\rho_\theta)^{-1} = \rho_{-\theta}$.

and reflection in OX is the mapping $\sigma_y : (x, y) \sigma_y = (x, -y)$ which is isometry and $(\sigma_y)^{-1} = \sigma_y$.

Note 1. Isometries are product of reflections, translations and rotations. In other words every isometry E is expressible as the product of a reflection, a translation and a rotation.

Symmetry groups. If S be a subset of the Euclidean plane then the set I_s of all $\sigma \in I$ s.t. $s \in S \Rightarrow s\sigma \in S$ and $t \in S \Rightarrow t\sigma \in S$, forms a subgroup of I , known as the symmetry group of S .

Algebra of symmetries of an equilateral triangle

Case I. Counter clockwise rotation of an equilateral triangle in its own plane about an axis through geometric centre O and perpendicular to the plane of the triangle ABC .

Let us define the rotations as follows:

$$R_0 : \triangle ABC \rightarrow \triangle ABC$$

$$R_1 : \triangle ABC \rightarrow \triangle CAB$$

$$R_2 : \triangle ABC \rightarrow \triangle BCA$$

where R_0 is the same position of the triangle, R_1 is the counter clockwise rotation through 120° which carries A to B , B to C , C to A and R_2 is the counter clockwise rotation through 240° which carries A to C , B to A , C to B . Evidently a counter clockwise rotation through 120° is identical with a clockwise rotation through 240° and similarly a counter clockwise rotation is identical with a clockwise one through 120° . As such R_0 ,




Fig. 4.18

R_1, R_2 are the only three distinct rotations, forming a finite abelian group.




Fig. 4.19




Fig. 4.20

Let $G = \{R_0, R_1, R_2\}$ with the binary operation $R_j \circ R_k$ being the rotation obtained by j -successive application of R_j and R_k for any two rotations R_j and R_k .

e.g. $R_1 \circ R_2$ represents the rotation through 120° followed by a rotation through 240° i.e. $R_1 \circ R_2$ is the rotation through 360° and hence

$$R_1 \circ R_2 = R_0$$

Conclusively

$$\begin{aligned} \triangle ABC &\xrightarrow{R_0} \triangle ABC \xrightarrow{R_1} \triangle CAB \xrightarrow{R_1} \triangle BCA \Rightarrow \\ &\xrightarrow{R_0 \circ R_1} \triangle ARC \xrightarrow{R_2} \triangle BCA \end{aligned}$$

	R_0	R_1	R_2
R_0	R_0	R_1	R_2
R_1	R_1	R_2	R_0
R_2	R_2	R_0	R_1

which is equivalent to $\triangle ABC \xrightarrow{R_2} \triangle BCA$

The composition table is as shown here.

This group of rotations G is abelian since, G_1 is satisfied, since every element of the table belong to S_3 .

G_2 is satisfied, since $R_0 \circ (R_1 \circ R_2) = R_0 \circ (R_0) = R_0$

and $(R_0 \circ R_1) \circ R_2 = R_1 \circ R_2 = R_0 = R_0 \circ (R_1 \circ R_2)$

G_3 is satisfied, since R_0 is an identity in S_3 .

G_4 is satisfied, since inverses of R_0, R_1, R_2 are R_0, R_2, R_1 respectively as $R_1 \circ R_2 = R_0$ etc.

Commutative property is also satisfied, since

$$R_0 \circ R_1 = R_1 = R_1 \circ R_0 \text{ etc.}$$

Case II. Rotation of an equilateral triangle ABC about the medians AD, BE, CF .

Let R_3, R_4, R_5 be the rotations about the medians AD, BE and CF respectively of equilateral triangle ABC , each through π . Clearly there are six coincident rotations of the triangle. There is correspondence between the group of rotations.

$$G' = \{R_0, R_1, R_2, R_3, R_4, R_5\}$$




Fig. 4.21




Fig. 4.22




Fig. 4.23

and the symmetric group S_3 whose elements permute A, B, C .

We have,

$$\begin{aligned} R_1 \circ R_3 = R_4 & \because \begin{array}{c} A \ R_1 \ C \ R_3 \ C \\ B \ C \ A \ B \ B \ A \end{array} \quad \text{by Fig. 4.22.} \quad \begin{array}{c} o \\ R_0 \end{array} \begin{array}{c} R_0 \\ R_0 \end{array} \begin{array}{c} R_1 \\ R_1 \end{array} \begin{array}{c} R_2 \\ R_2 \end{array} \begin{array}{c} R_3 \\ R_3 \end{array} \begin{array}{c} R_4 \\ R_4 \end{array} \begin{array}{c} R_5 \\ R_5 \end{array} \\ R_1 \circ R_4 = R_5 & \because \begin{array}{c} A \ R_1 \ C \ R_4 \ B \\ B \ C \ A \ B \ A \ C \end{array} \quad \text{by Fig. 4.23.} \quad \begin{array}{c} R_1 \\ R_2 \end{array} \begin{array}{c} R_1 \\ R_2 \end{array} \begin{array}{c} R_2 \\ R_0 \end{array} \begin{array}{c} R_4 \\ R_1 \end{array} \begin{array}{c} R_5 \\ R_5 \end{array} \begin{array}{c} R_5 \\ R_4 \end{array} \\ R_1 \circ R_5 = R_3 & \because \begin{array}{c} A \ R_1 \ C \ R_5 \ A \\ B \ C \ A \ B \ C \ B \end{array} \quad \text{by Fig. 4.21.} \quad \begin{array}{c} R_3 \\ R_4 \end{array} \begin{array}{c} R_3 \\ R_4 \end{array} \begin{array}{c} R_5 \\ R_2 \end{array} \begin{array}{c} R_4 \\ R_5 \end{array} \begin{array}{c} R_0 \\ R_1 \end{array} \begin{array}{c} R_2 \\ R_0 \end{array} \\ R_3 \circ R_5 = R_5 & \because \begin{array}{c} A \ R_2 \ B \ R_5 \ B \\ B \ C \ C \ A \ A \ C \end{array} \quad \text{by Fig. 4.23.} \quad \begin{array}{c} R_5 \\ R_5 \end{array} \begin{array}{c} R_5 \\ R_4 \end{array} \begin{array}{c} R_4 \\ R_3 \end{array} \begin{array}{c} R_5 \\ R_2 \end{array} \begin{array}{c} R_1 \\ R_1 \end{array} \begin{array}{c} R_0 \\ R_0 \end{array} \end{aligned}$$

Similarly

$$R_2 \circ R_4 = R_3, R_2 \circ R_5 = R_4 \text{ etc.}$$

In general $R_j \circ R_k$ for $j=0, 1, 2, \dots, 5$ and $k=0, 1, \dots, 5$ gives the adjoining composition table.

Clearly the set of six rotations of the equilateral triangle forms a non-abelian group.

The di-hedral group

If S be a regular polygon of n (>2) sides, then in any isometry of S , vertices are taken to vertices. Then the order of the symmetry group of a regular n -gon (polygon of sides $n>2$) can be easily determined.

In this connection the following axioms are to be noted:

(1) Every regular n -gon can be circumscribed by one and only one circle.

(2) The centre of a regular n -gon S is considered onto itself by any element of I_s .

(3) If S be a regular n -gon and $\sigma \in I_s$, then vertices of S are taken onto vertices of S by σ .

The symmetry group of the regular n -gon is said to be the di-hedral group of degree n .

Determination of the orders of the di-hedral groups

Suppose the vertices of a regular n -gon S with centre O are $A_1, A_2, \dots, A_n$ in a clock wise direction. Also suppose that σ_j , $1 \leq j \leq n$ rotates S




Fig. 4.24




Fig. 4.25

about O in clockwise direction through an angle $\frac{2\pi(j-1)}{n}$ radians (i.e., $\frac{360}{n}(j-1)$ degrees, So that $A_1 \sigma_j = A_j$ etc.

For the sake convenience we have shown here the effect of σ_3 on the regular pentagon (5-gon) in Fig 4.24.

Taking τ as the reflection about the line through A_1 and O , s.t.

$$A_1 \tau = A_1, A_1 \tau = A_n$$

The effect of τ on the regular pentagon is shown in Fig. 4.25.

The Fig. 4.26 shows the effect on a regular pentagon of the reflection τ followed by the rotation.

We observe that the elements $\sigma_1, \sigma_2, \dots, \sigma_n, \tau \sigma_1, \dots, \tau \sigma_n$ are all distinct since, of course $\sigma_j \neq \sigma_k, j \neq k$ as $A_1 \sigma_j \neq A_1 \sigma_k, j \neq k$.

In case $\tau \sigma_j = \sigma_k$, then $A_1 \tau \sigma_j = A_1 \sigma_j = A_1 \sigma_k$

$$\therefore \tau \sigma_j = \sigma_k \Rightarrow j = k$$

But $\tau \sigma_j = \sigma_j \Rightarrow \tau = \sigma_1$, the identity which contradicts the hypothesis. Finally $\tau \sigma_j = \tau \sigma_k \Rightarrow \sigma_j = \sigma_k$.

Consequently there are at least $2n$ possible elements of the dihedral group of degree n , but there are no more than $2n$ since if $\sigma \in I_s$, S being regular n -gon, then there are n possibilities for $A_1 \sigma$.

Since the vertices are taken to vertices, therefore $A_1 \sigma$ is one of A_1 ,




Fig. 4.26

..., A_n and $A_2\sigma$ has only two possibilities once $A_1\sigma$ has been determined as $d(A_1\sigma, A_2\sigma) = d(A_1, A_2)$ and $A_2\sigma$ must also be a vertex.

So $A_1\sigma$ and $A_2\sigma$ being once determined, $A_i\sigma, i=3, 4, \dots, n$ may also be determined. Thus $\exists$ at most two elements $\sigma \in I_s$ which map $A_1\sigma$ to A_j . Hence there are at most $2n$ elements of I_s and so

$$|I_s| = 2n.$$

Note. 2. The dihedral group of degree n is denoted by D_n .

As an example the elements of D_3 are $\sigma_j, \tau\sigma_j$ and $\sigma_j\sigma_2 = \sigma_j + 1, 1 \leq j \leq 2$ and $\sigma_3\sigma_2 = \sigma_1$

Also we have $\tau^{-1} = \tau$ and $\sigma_i\tau = \tau^2\sigma_i\tau = \tau\sigma_i\tau$

Now $\tau\sigma_1\tau = \sigma_1$, σ_1 being identity

$\tau\sigma_2\tau = \sigma_3$ as $A_1\tau\sigma_2\tau = A_1\sigma_2\tau = A_2\tau = A_3$

and $A_2\tau\sigma_2\tau = A_3\sigma_2\tau = A_1\tau = A_1$

So $\sigma_3\tau = \tau\sigma_2$ and $\tau\sigma_3 = \sigma_2\tau$

The multiplication table is as shown here.

	σ_1	σ_2	σ_3	τ	$\tau\sigma_2$	$\tau\sigma_3$
σ_1	σ_1	σ_1	σ_2	τ	$\tau\sigma_2$	$\tau\sigma_3$
σ_2	σ_2	σ_2	σ_3	$\tau\sigma_3$	τ	$\tau\sigma_2$
σ_3	σ_3	σ_3	σ_1	$\tau\sigma_2$	$\tau\sigma_3$	τ
τ	τ	τ	$\tau\sigma_2$	$\tau\sigma_3$	σ_1	σ_2
$\tau\sigma_2$	$\tau\sigma_2$	$\tau\sigma_3$	τ	σ_3	σ_1	σ_2
$\tau\sigma_3$	$\tau\sigma_3$	τ	$\tau\sigma_2$	σ_2	σ_3	σ_1

4.13. SOME SPECIAL GROUPS WITH LINEAR OPERATORS

Vector Space or Linear Space is an additive Abelian group L (Elements of L being called vectors) with the property that any scalar α (real or complex) and any vector $\mathbf{x}$ can be combined by the operation of scalar multiplication to yield a vector $\alpha\mathbf{x}$ s.t.

(i) $\alpha(\mathbf{x} + \mathbf{y}) = \alpha\mathbf{x} + \alpha\mathbf{y}, \mathbf{x} \in L, \mathbf{y} \in L \Rightarrow \alpha\mathbf{x} \in L$ and $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x} \in L$

(ii) $(\alpha + \beta)\mathbf{x} = \alpha\mathbf{x} + \beta\mathbf{x}$

(iii) $(\alpha\beta)\mathbf{x} = \alpha(\beta\mathbf{x})$

(iv) $1 \cdot \mathbf{x} = \mathbf{x}$

e.g. the set of $n \times n$ matrices forms a linear space.

A linear space is real or complex linear space according as the scalars are real or complex numbers.

If the linear space consists solely of the vector $\mathbf{O}$ with scalar multiplication defined by $\alpha \cdot \mathbf{O} = \mathbf{O}$ for all α , then we call it as *zero space* and denote it by $\{\mathbf{O}\}$,

A non-empty subset M of L is said to be a *subspace* or a *linear subspace* of L if M is a linear space in its own right w.r.t. linear operations in L . In case M is a proper subset of L , then we call it a *proper subspace* of L .

Basis for Linear Space. If S be a linear independent set of vectors in a linear space L , then $\exists$ a basis B for L s.t. $S \subseteq B$ and the basis for L is a linearly independent set which span the whole space L . Moreover

$S (\neq \emptyset) \subset L$ is linear independent iff $\Leftrightarrow$ each vector in the subspace $[S]$ spanned by S is uniquely expressible as a linear combination of the vectors in S e.g. the vector $\mathbf{O}$ in $[S]$ is uniquely expressible in the form

$$\mathbf{O} = \mathbf{O} \cdot \mathbf{x}_1 + \mathbf{O} \cdot \mathbf{x}_2 + \dots + \mathbf{O} \cdot \mathbf{x}_n \text{ where } S = \{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\}$$

While the vectors in the sub space $[S]$ spanned by S are the linear

combinations of the type $\mathbf{x} = \alpha_1\mathbf{x}_1 + \alpha_2\mathbf{x}_2 + \dots + \alpha_n\mathbf{x}_n$; $\alpha_1, \alpha_2, \dots, \alpha_n$ are scalars.

Dimension of a linear space. The dimension of a linear space is the number of elements in its basis, e.g. $L = \{\mathbf{O}\}$ is 0-dimensional and $L \neq \{\mathbf{O}\}$ has dimension equal to the number of elements in any basis.

A linear space is *finite-dimensional* if its dimension is 0 or a positive integer and it is *infinite-dimensional* if its dimension is not zero or a finite positive integer.

Linear transformations. If L, L' be two linear spaces with the same system of scalars, then a mapping T of L into L' is said to be a *linear transformation*

$$\text{if } T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y}) \text{ and } T(\alpha\mathbf{x}) = \alpha T(\mathbf{x})$$

or equivalently if $T(\alpha\mathbf{x} + \beta\mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$.

A set of operator $T_1, T_2, \dots$, in a linear space L forms a group if it satisfies all the four group axioms.

A linear transformation of one linear space into another is a *homomorphism* of the first space into the second, because it is a mapping which preserves the linear operations. Also it is observed that T preserves the origin ($\mathbf{O}$) and negatives for

$$T(\mathbf{O}) = T(\mathbf{O} \cdot \mathbf{O}) = \mathbf{O} \cdot T(\mathbf{O}) = \mathbf{O}$$

$$\text{and } T(-\mathbf{x}) = T((-1) \cdot \mathbf{x}) = (-1) T(\mathbf{x}) = -T(\mathbf{x}).$$

Examples: (i) $T_1((\mathbf{x}_1, \mathbf{x}_2)) = (\alpha\mathbf{x}_1, \alpha\mathbf{x}_2)$, α being real, multiplies each vector in $\mathbb{R}^2$ by into self where $T_1: \mathbb{R}^2 \rightarrow \mathbb{R}^2$

(ii) The mapping $D: D(p) = \frac{dp}{dn}$ is the linear transformation of P into itself where the linear space P of all polynomials $p(n)$ with real coefficients is defined on $[0, 1]$

(iii) The mapping $I: I(f) = \int_0^1 f(x) dx$ is the linear transformation of $C[0, 1]$ in the real linear space $\mathbb{R}$.

Characteristics of Linear Transformations

(i) If T and U be two operators transforming the linear space L into L , then $(T + U)(\mathbf{x}) = T(\mathbf{x}) + U(\mathbf{x})$... (1)

Similarly $(\alpha T)(\mathbf{x}) = \alpha T(\mathbf{x})$... (2)

is not shell if L and L' be two linear spaces with the same system of scalars, then the set of all linear transformations of L into L' is itself a linear space w.r.t. linear operations (1) and (2).

(ii) If T and U be two linear transformations on L , then their product TU is defined by $(TU)(\mathbf{x}) = T(U\mathbf{x})$

(iii) If T, U, V be three linear transformations on L , then the operation is associative

$$\text{i.e. } T(UV) = (TU)V$$

(iv) If T, U, V be three linear transformations on L , then distributive law holds

$$\text{i.e. } T(U + V) = TU + TV$$

and $(T+U)V=TV+UV$
 since, $((T+U)V)(\mathbf{x})=(T+U)(V(\mathbf{x}))=T(V(\mathbf{x}))+U(V(\mathbf{x}))$
 $= (TV)(\mathbf{x})+(UV)(\mathbf{x})$
 $= (TV+UV)(\mathbf{x}).$

(v) If T, U be two linear transformations on L and α is a scalar, then

$$\alpha(TU)=(\alpha T)U=T(\alpha U).$$

(vi) Identity transformation I is defined by $I(\mathbf{x})=\mathbf{x}$ and it is observed that

$$I \neq 0 \Leftrightarrow L \neq \{0\} \text{ and } TI=IT=I$$

Also if α is a scalar, then αI is known as scalar multiplication for $(\alpha I)(\mathbf{x})=\alpha I(\mathbf{x})=\alpha \mathbf{x}.$

(vii) A linear transformation T on L is *non-singular* if it is one-one and onto and *singular* otherwise.

(viii) If the linear transformation T on L is non-singular and one-one onto, then $\exists$ its *inverse* T^{-1} s.t. $TT^{-1}=T^{-1}T=I$.

It is easy to show that if T is non-singular then the mapping T^{-1} is also a linear transformation on L .

(ix) If L be the direct sum of the subspaces M and N s.t. $L=M \oplus N$, then each vector $z \in L$ can be uniquely expressed as $z=\mathbf{x}+\mathbf{y}$, $\mathbf{x} \in M$ and $\mathbf{y} \in N$, $\mathbf{x}$ being uniquely determined by z , we define a mapping E of L into itself s.t. $E(z)=\mathbf{x}$ and call it *projection on M along N* .

Clearly E is *idempotent* since $E^2=E$.

$$\text{Also } E^2(z)=(EE)(z)=E(E(z))=E(\mathbf{x})=\mathbf{x}=E(z)$$

(x) *Reversal law*. If T, U be two non-singular linear transformations on L , then

$$(TU)^{-1}=U^{-1}T^{-1}$$

But $(\alpha T)^{-1}=\alpha^{-1}T^{-1}$, α being a scalar.

(xi) If T be a linear transformation of L to L' , then we get an *isomorphic group* of operators in L' which transform $A, B, \dots$ etc. to $A', B', \dots$ etc s.t.

$$A'=TAT^{-1}, B'=TBT^{-1}, \dots$$

Note. If an arbitrary group G is mapped homomorphically onto a group of operators $D(G)$ in the linear space L , then the *operator group* $D(G)$ is said to be a *Representation of the group G* in the representative space L . In case n is the dimension of L , then the degree of representation is also n or in other words the *representation is n -dimensional*.

If $a, b \in G$, then $D(ab)=D(a)D(b)$

$$D(\alpha^{-1})=[D(a)]^{-1} \text{ and } D(E)=1$$

Matrices and Linear Transformations

Linear operator T is a mapping s.t. $T: L \rightarrow L', L, L'$ being vector space over a field F

Let $\text{Dim. } L=n$ and its basis $B=\{x_1, x_2, \dots, x_n\}$ and let $\text{dim. } L'=m$ and its basis $B'=\{y_1, y_2, \dots, y_n\}$

$\forall x \in L, T(x) \in L'$, and $T: L \rightarrow L' \therefore$ each $T(x)$ is expressible as a linear combination of elements of B' and in particular each $T(x_j)$, $j=1, 2, \dots, n$ is expressible as a linear combination of m vectors in B' .
 Let mn scalars $a_{ij} \in F$, $1 \leq i \leq m, 1 \leq j \leq n$.

$$\text{So } T(x_1)=a_{11}y_1+a_{21}y_2+\dots+a_{m1}y_m=\sum_{i=1}^m a_{i1}y_i$$

$$T(x_2)=a_{12}y_1+a_{22}y_2+\dots+a_{m2}y_m=\sum_{i=1}^m a_{i2}y_i$$

$$\dots\dots\dots$$

$$T(x_n)=a_{1n}y_1+a_{2n}y_2+\dots+a_{mn}y_n=\sum_{i=1}^m a_{in}y_i$$

$$\text{or symbolically } T(x_j)=\sum_{i=1}^m a_{ij}y_i, j=1, 2, \dots, n$$

So that *co-efficient matrix* of these expressions is

$$\begin{bmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ \dots & \dots & \dots & \dots \\ a_{1n} & a_{2n} & \dots & a_{mn} \end{bmatrix} = A \text{ (say)}$$

The matrix of $T: L_1 \rightarrow L_2$ (with respect to basis B, B') is the transpose of the matrix A i.e. matrix of T w.r.t. basis B, B' is

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

which is written symbolically as $[T: B, B']$ or simply $[T]$

In case T is a linear operator s.t. $T: L \rightarrow L$ i.e. $L=L', m=n, B=B'$ the matrix $[a_{ij}]_{m \times n}$ will become $[a_{ij}]_{n \times n}$ and is denoted by $[T, B]$ or $[T]_B$.

$$\text{i.e. } T(x_j)=\sum_{i=1}^n a_{ij}x_i, j=1, 2, \dots, n.$$

Conversely to find T whose matrix w.r.t. basis B is given to be $[a_{ij}]$

$$x \in V \text{ and } B=\{x_1, x_2, \dots, x_n\} \text{ while } x=\sum_{j=1}^n \alpha_j x_j, \alpha_j \in F, \text{ we have}$$

$$T(x)=T\left(\sum_{j=1}^n \alpha_j x_j\right)=\sum_{j=1}^n \alpha_j T(x_j)=\sum_{j=1}^n \alpha_j \left(\sum_{i=1}^m a_{ij}y_i\right)=\sum_{i=1}^m \left(\sum_{j=1}^n a_{ij}\alpha_j\right)y_i$$

$$\left(\sum_{j=1}^n a_{ij}\alpha_j\right)y_i$$

which belongs to L' as B' its basis.

If T and U be two linear operators on a linear space L whose matrices w.r.t. a fixed basis $B=\{x_1, x_2, \dots, x_n\}$ are $[T]$ and $[U]$, then

$$[T+U]=[T]+[U], [\alpha T]=\alpha[T] \text{ and } [TU]=[T][U]$$

since if $[T]=[a_{ij}]_{n \times n}$, $U=[b_{ij}]_{n \times n}$, so that $T(x_j) = \sum_{i=1}^n a_{ij} x_i$, $U(x_j)$

$$= \sum_{i=1}^n b_{ij} x_i$$

then $[T+U](x_j) = T(x_j) + U(x_j) \quad \forall j=1, 2, \dots, n$

$$= \sum_{i=1}^n a_{ij} x_i + \sum_{i=1}^n b_{ij} x_i$$

$$= \sum_{i=1}^n (a_{ij} + b_{ij}) x_i = \sum_{i=1}^n c_{ij} x_i \text{ where } [c_{ij}] = [a_{ij} + b_{ij}]$$

$$\therefore [T+U] = [c_{ij}] = [a_{ij} + b_{ij}] = [a_{ij}] + [b_{ij}] = [T] + [U]$$

$$\text{and } [\alpha T](x_j) = \alpha [T(x_j)] = \alpha \sum_{i=1}^n a_{ij} x_i = \sum_{i=1}^n (\alpha a_{ij}) x_i = \sum_{i=1}^n (c_{ij}) x_i$$

$$\therefore [\alpha T] = [c_{ij}] = \alpha [a_{ij}] = \alpha [T]$$

$$\text{also } [TU](x_j) = T[U(x_j)] = T\left(\sum_{k=1}^n b_{kj} x_k\right) = \sum_{k=1}^n b_{kj} T(x_k)$$

$$= \sum_{k=1}^n b_{kj} \left(\sum_{i=1}^n a_{ik} x_i \right)$$

$$= \sum_{i=1}^n \left(\sum_{k=1}^n a_{ik} b_{kj} \right) x_i = \sum_{i=1}^n (c_{ij}) x_i$$

$$\therefore [TU] = [c_{ij}] = \left[\sum_{k=1}^n a_{ik} b_{kj} \right] = [a_{ij}] [b_{ij}] = [T] [U]$$

If $T: L \rightarrow L'$ and $U: L' \rightarrow L''$ so that $UT: L \rightarrow L''$ where L, L', L'' are finite dimensional linear spaces of dimensions n, m, p (say) and basis B, B', B'' (say) then $[UT; B, B''] = [U; B', B''] [T; B, B']$

If L, L' are two finite dimensional linear spaces of dimensions n and n' and basis B and B' , s.t. the function $T: L \rightarrow L'$ then its matrix w.r.t. B, B' is an isomorphism between the space (L, L') and the space of $n \times n'$ matrices over the field F .

The matrices of identity operator I and zero operator O on a linear space L w.r.t. the basis $B=[x_1, x_2, \dots, x_n]$ are the unit and null matrices respectively, i.e. $[I]=[\delta_{ij}]$ where $\delta_{ij}=0$ for $i \neq j$

$$=1 \text{ for } i=j$$

$$\text{and } [O]=[o]$$

Since if T is a linear operator and $[a_{ij}]$ is the matrix of T w.r.t.

$$\text{basis } B \text{ then } T(x_j) = \sum_{i=1}^n a_{ij} x_i$$

$$\text{and } I(x_i) = x_i = \sum_{i=1}^n \delta_{ij} x_i = 0x_1 + 0x_2 + \dots + 1 \cdot x_j + \dots + 0x_n$$

$$\therefore [I] = [\delta_{ij}] \text{ is a unit matrix.}$$

$$\text{Also } O(x_j) = 0 = 0x_1 + 0x_2 + \dots + 0x_n = \sum_{i=1}^n 0_{ij} x_i$$

i.e. $[O]=[o_{ij}]$ is a null matrix.

If $[T]=[a_{ij}]$ be the matrix of operator T on L w.r.t. basis B , then T is invertible $\Leftrightarrow [T]=[a_{ij}]$ is invertible and in this case

$$[T^{-1}] = [T]^{-1} = [a_{ij}]^{-1}$$

Since T is invertible, $TT^{-1} = T^{-1}T = I$

$$\therefore [TT^{-1}] = [T^{-1}T] = [I]$$

$$\text{or } [T][T^{-1}] = [T^{-1}][T] = [I] = \delta_{ij}$$

which follows that $[T]$ is non-singular and $[T^{-1}] = [T]^{-1} = [a_{ij}]^{-1}$.

If T, U be two linear operators on a linear space L , then T and U are said to be similar if $\exists$ an invertible operator V on L s.t.

$$TV = VU \text{ i.e. } T = VUV^{-1} \text{ or } U = V^{-1}TV$$

The relation of similarity in the set of all $n \times n$ matrices over the field F is an equivalence relation.

Since if A and B be two $n \times n$ matrices and R is a relation of similarity then $\exists$ an invertible $n \times n$ matrix P s.t. $A = PBP^{-1}$ or $B = P^{-1}AP$.

Now R is reflexive since $A = IAI^{-1}$, I being an invertible unit-matrix

R is symmetric since $A = PBP^{-1} \Rightarrow P^{-1}AP = P^{-1}(PBP^{-1})P$

$$\Rightarrow P^{-1}AP = B$$

$$\Rightarrow B = P^{-1}A(P^{-1})^{-1} = CAC^{-1} \text{ where } C = P^{-1}$$

$\Rightarrow B$ is similar to A and hence symmetric.

R is transitive since if $A = PBP^{-1}$ and $B = QCQ^{-1}$ then

$$A = P(QCQ^{-1})P^{-1} = (PQ)C(Q^{-1}P^{-1})$$

$$= (PQ)C(PQ)^{-1}$$

As such a result analogous to it follows:

The relation of similarity in the set of all linear operators on a linear space L is an equivalence relation.

If T is a linear operator on a vector space L and $[T]$ be the matrix of T w.r.t. a basis B , then determinant i.e. $\det T = \det [T]$

Linear functionals. A linear functional f over a vector space L is a mapping which assigns to each member $\alpha \in L$, an element $f(\alpha)$ which in F (field) s.t. f is linear i.e.

$$f(\alpha + \beta) = f(\alpha) + f(\beta), \alpha, \beta \in L$$

$$f(a\alpha) = af(\alpha) \quad a \in F$$

or in one relation, $f(a\alpha + \beta) = af(\alpha) + f(\beta)$

If $f(\alpha)=0 \forall \alpha \in L$, then f is said to be a *zero functional*.

Dual space of vector space. The set of all linear functionals f on L denoted by L^* or $V(L, F)$, F being a field, w.r.t. two compositions s.t.

$$(f_1 + f_2)\alpha = f_1(\alpha) + f_2(\alpha) \quad \forall \alpha \in L$$

$$(af)\alpha = a[f(\alpha)] \quad \forall \alpha \in L, a \in F$$

is said to be a *dual space* of L i.e.

$$L^* = \{f: L \rightarrow F \text{ s.t. } f \text{ is linear}\}.$$

Transpose of Linear Transformation

If T be a linear operator from L to L' (over a field F) s.t.

$$\alpha \in L \Rightarrow T(\alpha) \in L'$$

then if $T: L \rightarrow L'$ induces a linear transformation $T^T: L'^* \rightarrow L^*$ where L^*, L'^* are dual spaces of L, L' ; T^T is called as *transpose* of the linear transformation T .

T^T is linear and unique and $\text{Rank}(T^T) = \text{Rank}(T)$

Adjoint of an Operator

If $T: L \rightarrow L$, L being a vector space over a field F , s.t. $\alpha \in L \Rightarrow T(\alpha) \in L$, induces a linear operator $T^*: L^* \rightarrow L^*$, L^* being dual space of L with its elements as functional on L , then T^* on L^* is called as *adjoint* of the linear operator T on L .

T^* is linear and unique.

Its properties are: (i) $(T_1 + T_2)^* = T_1^* + T_2^*$

(ii) $(kT)^* = kT^*$, k being a scalar

(iii) $(T_1 T_2)^* = T_2^* T_1^*$ (reversal law)

(iv) If T is invertible, $(T^*)^{-1} = (T^{-1})^*$

(v) Zero and identity operators are self adjoint i.e. $O^* = O$ and $I^* = I$

(vi) $(T^*)^* = T^{**} = T$.

Inner Product Vector Space (I.P.V.S.)

A vector space with an inner product defined on it is said to be an inner product space, while the inner product of x (a_1, a_2, a_3) and y (b_1, b_2, b_3) being defined as $(x, y) = a_1 b_1 + a_2 b_2 + a_3 b_3$ with the properties

$$|x| = \sqrt{a_1^2 + a_2^2 + a_3^2}, \quad |x| = 0 \text{ iff } a_1 = 0 = a_2 = a_3$$

$$x \text{ and } y \text{ are orthogonal if } \cos \theta = \frac{(x, y)}{|x||y|} = 0 \text{ i.e. } (x, y) = 0$$

$$(x, x) = 0 \text{ and } (x, y) = (y, x) \text{ also } (ax + by, z) = a(x, z) + b(y, z)$$

Note. $R^n(R)$ is an euclidean space and $C^n(C)$ is an unitary space. As an illustration, the set of all $n \times n$ matrices forms an inner product vector space over a field F (real or complex) if the inner product is defined as $(A, B) = \text{Trace}(AB^0)$, B^0 being transpose conjugate of B .

Here if $B = [b_{ij}]$ then $B^0 = \bar{b}_{ji}$

So $(A + C, B) = \text{trace}[(A + C)B^0]$

$$= \text{tr}(AB^0 + CB^0)$$

$$= \text{tr} AB^0 + \text{tr} CB^0$$

$$= (A, B) + (A, C)$$

Since Trace of sum = sum of traces.

$$(A, B) = \text{tr}(A, B)^0 = \text{sum of diagonal elements of } AB^0$$

$$= \text{Sum of conjugates of diagonal elements of } AB^0$$

$$= \text{Sum of diagonal elements of } A^0 B$$

$$= \text{trace } A^0 B = \text{tr}(BA^0) = (B, A)$$

Similarly $(kA, B) = k(A, B)$ and $\text{tr}(kA, B^0) = k \text{tr}(AB^0) = k(A, B)$ showing that the given set is an inner product vector space.

Unitary and Orthogonal Operator

If $TT^* = T^*T = I$, T is called unitary operator for complex I.P.V.S. and orthogonal for real I.P.V.S.

Normal operator

If $TT^* = T^*T$, then T is called normal operator, T^* being adjoint of T .

Characteristic Vectors and Characteristic Values

If T be a linear operator on a finite dimensional vector space $L(F)$ then a scalar $\lambda \in F$ is said to be a characteristic value of T if $\exists$ a vector $\alpha (\neq 0) \in L(F)$ s.t. $T(\alpha) = \lambda \alpha$. This non-zero vector α associated with characteristic value λ is said to be a characteristic vector of T .

In other words, roots of $|T - \lambda I| = 0$ are characteristic values of T .

If T is invertible and has characteristic root λ , then λ^{-1} is the characteristic value of T^{-1} .

If T is not invertible, then 0 is the characteristic value of T .

If $\lambda \in F$ is a characteristic value of a linear operator T on a vector space $L(F)$ then for a polynomial $p(x)$ over F , $p(\lambda)$ is a characteristic value of $p(T)$.

Hamiltonian group. A non-commutative (i.e. non-abelian) group in which every subgroup is normal is said to be a Hamiltonian group.

Simple group. A simple group is one which contains no other normal subgroup except the two, one itself and the second a unit subgroup which is normal.

Unitary groups. The set of all non-singular square matrices of order n with multiplicative compositions form a group known as a *full linear group*. Its elements are the infinite number linear transformations which change a vector into a new vector and so the order of a full linear group is infinite.

Imposing certain condition on the matrices of its transformation, we may get many subgroups of full linear group. One such type is *discrete group* obtained by excluding all matrices except those whose determinant is ± 1 . The elements of a discrete group can be put into one-one correspondence with the set of positive integers.

The subgroups of a full linear group obtained by expanding all matrices except those whose determinant is ± 1 form a *continuous group* provided its elements are non-denumerable i.e. uncountable. In other words a continuous group contains the elements which can be generated by continuously varying parameters in any region, known as group space. There is one-one correspondence between group-elements and points of group space. Those groups whose elements can be generated by a finite number of continuously varying parameters, are known as *finite continuous groups*.

A subgroup of a full linear group having its elements as square unitary matrices of order 2 with determinant +1 is known as *2-dimensional unimodular unitary group* or *special unitary group*. e.g. a matrix $\begin{bmatrix} \alpha & \beta \\ \gamma & \delta \end{bmatrix} = A$ (say) will be a special unitary group if A is unitary matrix of order 2 i.e. $AA^\dagger = I$ and secondly if $|A| = +1$

Now $A^\dagger = \begin{bmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{bmatrix}$, $\bar{\alpha}, \bar{\beta}, \bar{\gamma}, \bar{\delta}$ being complex conjugate of $\alpha, \beta, \gamma, \delta$ respectively

$$\therefore AA^\dagger = \begin{bmatrix} \alpha & \beta \\ \gamma & \delta \end{bmatrix} \begin{bmatrix} \bar{\alpha} & \bar{\gamma} \\ \bar{\beta} & \bar{\delta} \end{bmatrix} = \begin{bmatrix} \alpha\bar{\alpha} + \beta\bar{\beta} & \alpha\bar{\gamma} + \beta\bar{\delta} \\ \gamma\bar{\alpha} + \delta\bar{\beta} & \gamma\bar{\gamma} + \delta\bar{\delta} \end{bmatrix} = I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \text{ gives}$$

$$\alpha\bar{\alpha} + \beta\bar{\beta} = 1 = \gamma\bar{\gamma} + \delta\bar{\delta} \text{ and } \alpha\bar{\gamma} + \beta\bar{\delta} = 0 = \gamma\bar{\alpha} + \delta\bar{\beta}$$

which yield, $\gamma = -\bar{\beta}$, $\delta = \bar{\alpha}$ and $\alpha\bar{\alpha} + \beta\bar{\beta} = 1$

As such a typical element of special unitary group is

$$U = \begin{bmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{bmatrix}; |U| = \alpha\bar{\alpha} + \beta\bar{\beta} = 1$$

Applying this matrix to the column vector $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ such that

$$UX = X' \text{ we get, } \begin{bmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1' \\ x_2' \end{bmatrix}$$

$$\text{i.e. } \begin{bmatrix} \alpha x_1 + \beta x_2 \\ -\bar{\beta} x_1 + \bar{\alpha} x_2 \end{bmatrix} = \begin{bmatrix} x_1' \\ x_2' \end{bmatrix}$$

or equivalently $\alpha x_1 + \beta x_2 = x_1'$

$$-\bar{\beta} x_1 + \bar{\alpha} x_2 = x_2'$$

which transform any function of x into linear combination of x_1, x_2 ,

$$Uf(x) = f(x') = f(\alpha x_1 + \beta x_2, -\bar{\beta} x_1 + \bar{\alpha} x_2)$$

As such if U operates on a set of $(n+1)$ homogeneous products

$$f_m^{(n)} = x_1^n x_2^{n-m} m = 0, 1, 2, \dots, n$$

then we get $Uf_m^{(n)} = (\alpha x_1 + \beta x_2)^m (-\bar{\beta} x_1 + \bar{\alpha} x_2)^{n-m}$

$$= \sum_{j=0}^n U_{mj}^{(n)} x_1^j x_2^{n-j}$$

$$\text{where } U_{mj}^{(n)} = \sum_k (-1)^k \frac{|m|}{|(m-k)|} \frac{|(n-m)|}{|k|} \frac{|(n-m-j+k)|}{|(j-k)|} \times \alpha^k \beta^{m-k} (\bar{\alpha})^{n-m-j+k} (\bar{\beta})^{j-k}$$

Character of a unitary group. The character (to be defined later) of a special unitary group is found out if a typical matrix by means of unitary transformation is transformed to diagonal form.

Take a unitary matrix V such that

$$V^{-1}UV = \begin{bmatrix} \gamma & 0 \\ 0 & \gamma' \end{bmatrix} = U'$$

Now $|U'| = +1$ is apparently satisfied if $U' = \begin{bmatrix} e^{i\phi/2} & 0 \\ 0 & e^{-i\phi/2} \end{bmatrix}$

All the other matrices of the group will belong to the same class as U and U' since the class constitutes elements obtained by unitary transformation while a unitary matrix remains unitary under such a transformation.

Thus the character (defined in § 4.14) of one element of the class is given by

$$\chi^{(\frac{1}{2})} = e^{i\phi/2} + e^{-i\phi/2} \text{ by using } U' \\ = 2 \cos \phi$$

In general the character (defined in § 4.14) of special unitary group for any value j is given by $\chi^{(j)} = \sum_{m=-j}^j e^{im\phi}$

If $\kappa = e^{i\phi}$, then $\chi^{(j)} = e^{-ij\phi} (1 + \kappa + \kappa^2 + \dots + \kappa^{2j})$

$$= e^{-ij\phi} \left(\frac{1 - \kappa^{2j+1}}{1 - \kappa} \right)$$

$$= \frac{\sin(2j+1)\frac{\phi}{2}}{\sin \frac{\phi}{2}} \text{ on multiplying numerator and denominator by } i \frac{\phi}{2}.$$

n -dimensional Rotation group. A continuous group formed from the set of all orthogonal n -dimensional matrices is said to be an *n -dimensional rotational group*. In fact this is a sub-group of a full linear group provided all elements are real unitary matrices whose determinant is +1.

e.g. if a point $P(x, y, z)$ is taken on the surface of a unit sphere and the sphere is rotated in any manner keeping its centre fixed; then the new coordinates of P say (x', y', z') related to (x, y, z) by some matrix $R(\alpha, \beta, \gamma)$ which is an element of a 3-dimensional rotation group $R^+(3)$ give a rotation factorized as product of three plane rotations described by the Eulerian angles (α, β, γ) (discussed in chapter of classical mechanics) i.e. $R(\alpha, \beta, \gamma) = R_z(\gamma) R_y(\beta) R_x(\alpha)$

where R_x, R_y, R_z are rotations about z, y, x axes respectively.

As an other example if we define a 2-dimensional rotation group $R^+(2)$ as a subgroup of $R^+(3)$, then its elements are obtained by proper rotation in a plane perpendicular to a fixed axis say z -axis. Taking $R(\theta)$ as one element of this group and $T(\theta)$ an operator transforming a vector $\mathbf{x}$ with components x_1, x_2 to another vector $\mathbf{x}'$ with components x_1', x_2' i.e.

$$\mathbf{x}' = T(\theta)\mathbf{x}, \quad 0 \leq \theta \leq 2\pi$$

such that $T(\theta) = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$, we have

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

or equivalently, $x_1' = x_1 \cos \theta + x_2 \sin \theta$

$$x_2' = x_2 \cos \theta - x_1 \sin \theta$$

But if $R(\theta')$ is another element of the group with transformation $T(\theta')$ then

$$T(\theta) T(\theta') = T(\theta + \theta') = T(\theta') T(\theta)$$

which follows that the group is commutative i.e. Abelian.

Point group. The inversion operation in 3-dimensional space is given by the matrix $T = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$

and an identity operation I is given by the unit matrix

$$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Evidently $TI = T$

and $T^2 = I$ i.e. $T = T^{-1}$

Here T and I form a group with matrix multiplication. Such a group is said to be a *point group* since one point (say origin) remains fixed in all operations. The fixed point is sometimes known as *centre of inversion*.

Consider a point group $\{c_n\}$ with operations on a regular polygon of n sides such that there exist

(i) a rotation through an angle $\frac{2\pi}{n}$ about an n -fold axis of rotation properly.

(ii) a rotation through $-\frac{2\pi}{n}$ about an n -fold axis of rotation im-
properly.

(iii) a reflection in a plane given by σ_H, σ_V ; H, V denoting Hori-
zontal and Vertical planes.

(iv) an inversion.

Such operations form a point group $\{c_n\}$.

Quaternion group. If we define a group G of order 8 such that

$$G = \{1, a, a^2, a^3, b, ab, a^2b, a^3b\}$$

with the properties, $ab = ba^3$

$$b^2 = a^2$$

and

$$a^4 = 1$$

whence $ab = ba^3$ and $b^2 = a^2 \Rightarrow ba = a^3b$

	1	a	a ²	a ³	b	ab	a ² b	a ³ b
1	1	a	a ²	a ³	b	ab	a ² b	a ³ b
a	a	a ²	a ³	1	ab	a ² b	a ³ b	b
a ²	a ²	a ³	1	a	a ² b	a ³ b	b	ab
a ³	a ³	1	a	a ²	a ³ b	b	ab	a ² b
b	b	a ³ b	a ² b	ab	a ²	a	1	a ³
ab	ab	b	a ³ b	a ² b	a ³	a ²	a	1
a ² b	a ² b	ab	b	a ³ b	1	a ³	a ²	a
a ³ b	a ³ b	a ² b	ab	b	a	1	a ³	a ²

Since $a^3b = a^2(ba^3) = b^3a^3 = b(a^2a^3) = ba$.

The composition table is as shown here. It is clear from this table that the group of order 8 under consideration does actually exist and defines a group. Such a group is known as *quaternion group*. All of its subgroups are normal, though it is not abelian. clearly a *quaternion group* is also a *Hamiltonian group*.

Torsion, Torsion-free and Mixed Groups

If G be a group such that every element of G than the identity is of infinite order then G is called as *Torsion-free group*. If G is a group such that every element of it is of finite order then G is called as *Torsion group*. Also if G is a group such that it consists of both an element of infinite order and an element (not equal to the identity) of finite order then G is called as *Mixed group*.

p-Primary group or p-group. A group G is said to be a primary group or p -group for some prime p if every element of G is of order which is a power of p . Actually a torsion group is made up of p -groups.

p-Prüfer group. If $\mathbb{Q}/\mathbb{Z}$ represents the additive group of rationals modulo the integers, then $\mathbb{Q}/\mathbb{Z}$ is evidently a torsion group and if

$$(Q/Z)_p = \{x+Z: x+Z \text{ of order a power of } p\}$$

$$= \{x+Z: p^r x \in Z\}$$

$$= \{m/p^r + Z: \text{for various integers } r \text{ and } 0 \leq m < p^{r-1}\}$$

and $Q/Z = \sum_{p \in I} (Q/Z)_p$, I being set of all primes p

then $(Q/Z)_p$ is said to be p -Prüfer group or a group of type P^∞

Clearly p -Prüfer group, $(Q/Z)_p = \bigcup_{r=1}^{\infty} C_r$ where $C_r = gp \left(\frac{1}{p^r} + Z \right)$

and $g \in G$ is of order $p_1^{r_1} p_2^{r_2} \dots p_n^{r_n}$; $p_1, p_2, \dots, p_n$ being distinct primes and $r_1, r_2, \dots, r_n$ positive integers.

Since $(Q/Z)_p = \{m/p^r + Z\}$ for various integers r and $0 \leq m < p^{r-1}$

So $[Q/Z]_p \supset C_r$

and $(Q/Z)_p \subset \bigcup_{r=1}^{\infty} C_r$

Which follow that $(Q/Z)_p = \bigcup_{r=1}^{\infty} C_r$

4.14. PERMUTATIONAL REPRESENTATIONS

Generalization of Cayley's theorem i.e. every group is isomorphic to a group of permutations.

If G be a group and ρ a mapping; $\rho: G \rightarrow G$ s.t. $x \rightarrow xg \forall x \in G$ then $g\rho$ being image of g in G under ρ , we have $g\rho: x \rightarrow xg, x \in G$

Here ρ is an isomorphism of G into a subgroup of S_G if

- (i) $g\rho$ is a permutation of $G \forall g \in G$
- (ii) g is a homomorphism i.e. if $g, h \in G$, then $(gh)\rho = g\rho \cdot h\rho$
- (iii) ρ is an isomorphism i.e. ρ is one-one.

(i) is satisfied since $g\rho$ is one-one mapping of G onto G as

$$x(g\rho) = y(g\rho) \Rightarrow xg = yg, x, y \in G$$

$$\Rightarrow xgg^{-1} = ygg^{-1}$$

$$\Rightarrow x = y$$

showing that the mapping is one-one.

Also if $x \in G$, then $(xg^{-1})g\rho = (xg^{-1})g = x$ showing that $g\rho$ is onto.

(ii) is satisfied since for $x \in G$ we have

$$x((gh)\rho) = x(gh) = (xg)h = [x(g\rho)](h\rho) = x(g\rho \cdot h\rho)$$

i.e. $(gh)\rho = g\rho \cdot h\rho$

showing that ρ is a homomorphism.

(iii) is satisfied since if $g\rho = h\rho$ and $1 \in G$ is the identity element, then $g = 1(g\rho) = 1(h\rho) = h$ showing that ρ is one-one.

For example consider a cyclic group G of order 2 s.t. $G = \{1, a\}$, where $a^2 = 1, 1 \cdot a = a \cdot 1$, then mappings

$$1\rho: 1 \rightarrow 1, a \rightarrow a$$

and $a\rho: 1 \rightarrow a, a \rightarrow 1$ show that ρ is an isomorphism since $a\rho \neq 1\rho$, ρ is one-one.

Definition of a Permutational Representation

A homomorphism of a group G into the symmetric group on a set X is known as a permutational representation of G on X .

If according to Cayley's theorem ρ is the isomorphism for G , then ρ itself is a permutational representation of G on G and known as left regular representation. A mapping μ of G into the symmetric group on the set X is a permutational representation of G if

$$(gh)\mu = g\mu h\mu, \text{ for all } g, h \in G.$$

For example, if G be a dihedral group of degree 4, then G is the group of symmetries of the square. If $g \in G$ takes each vertex, of $ABCD$ to a vertex, then g being one-one, Ag, Bg, Cg, Dg are distinct vertices. Suppose that

$X = \{A, B, C, D\}$ and mapping $\sigma_g: x\sigma_g = xg$
 $\forall x \in X$

so that $\sigma_g \in S_X$

Also if $\tau: G \rightarrow S_X$ s.t. $g\tau = \sigma_g$, then for $x \in X$, $g, h \in G$, we have $x(gh)\tau = x\sigma_{gh} = x(gh)$
 $= (xg)h = (x\sigma_g)\sigma_h = x(\sigma_g\sigma_h) = x(g\tau)(h\tau)$

giving $(gh)\tau = g\tau h\tau$

which shows that τ is a permutational representation.

Degree of a Representation

The number of elements in the set X gives the degree of a permutational representation or simply a representation on X . e.g. the degree of representation in the above example of dihedral group of degree 4, is four (4).

The degree of representation of the symmetric group G on $\{1, 2, 3\}$ is 6 if ρ itself is a representation of G as a permutation group on six elements

$$p_1: 1 \rightarrow 1, 2 \rightarrow 2, 3 \rightarrow 3; p_2: 1 \rightarrow 1, 2 \rightarrow 3, 3 \rightarrow 2; p_3: 1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 1;$$

$$p_4: 1 \rightarrow 2, 2 \rightarrow 1, 3 \rightarrow 3; p_5: 1 \rightarrow 3, 2 \rightarrow 2, 3 \rightarrow 1; p_6: 1 \rightarrow 3, 2 \rightarrow 1, 3 \rightarrow 2$$

So that

$$p_1\rho: p_1 \rightarrow p_1, p_2 \rightarrow p_2, p_3 \rightarrow p_3, p_4 \rightarrow p_4, p_5 \rightarrow p_5, p_6 \rightarrow p_6$$

$$p_2\rho: p_1 \rightarrow p_2, p_2 \rightarrow p_1, p_3 \rightarrow p_5, p_4 \rightarrow p_6, p_5 \rightarrow p_3, p_6 \rightarrow p_4$$

$$p_3\rho: p_1 \rightarrow p_3, p_2 \rightarrow p_4, p_3 \rightarrow p_6, p_4 \rightarrow p_5, p_5 \rightarrow p_2, p_6 \rightarrow p_1$$

$$p_4\rho: p_1 \rightarrow p_4, p_2 \rightarrow p_3, p_3 \rightarrow p_2, p_4 \rightarrow p_1, p_5 \rightarrow p_6, p_6 \rightarrow p_5$$

$$p_5\rho: p_1 \rightarrow p_5, p_2 \rightarrow p_6, p_3 \rightarrow p_4, p_4 \rightarrow p_3, p_5 \rightarrow p_1, p_6 \rightarrow p_2$$

$$p_6\rho: p_1 \rightarrow p_6, p_2 \rightarrow p_5, p_3 \rightarrow p_1, p_4 \rightarrow p_2, p_5 \rightarrow p_4, p_6 \rightarrow p_3$$

giving $(p_i\rho)(p_j\rho) = (p_j h_j)\rho, 1 \leq i, j \leq 6$.

But the degree of representation of the symmetric group on $\{1, 2, 3\}$




Fig. 4.27

is 3 since it is an identity isomorphism as G itself is a permutation group on $\{1, 2, 3\}$.

As another example the degree of representation of a cyclic group G of order n s.t. $G = \{1, a, a^2 \dots a^{n-1}\}$, n being positive integer, is infinite.

Faithful representation. If a representation is one-one then it is called as faithful representation.

e.g. the representations in the above quoted examples are respectively, faithful (dihedral group); faithful; faithful; faithful.

As another example each matrix is its own faithful representation.

The number of rows and columns in a representation matrix is sometimes known as the *dimension* of the representation. As similarity transformations do not change the multiplication properties of matrices so under such transformations, the nature of representation remains invariant. As such new representations known as *equivalent representations* can be obtained from the given representation by means of similarity transformations.

Addition of representations. Consider m square matrices $A_1, A_2, \dots, A_m$ each of order n with their matrix representation of a group as $\mu(A_1), \mu(A_2), \dots, \mu(A_m)$ and similarly take another matrix representation consisting of m square matrices of order p s.t.

$$\mu'(A_1), \mu'(A_2), \dots, \mu'(A_m)$$

Since from the two matrix representations a single row can be found by their addition in which two representations are merely joined into one, therefore a new representation consisting of m square matrices of order $m+n$ can be obtained by adding these representations such as

$$\left[\begin{array}{c|c} \mu & O \\ \hline O & \mu' \end{array} \right] \begin{array}{l} \uparrow n \text{ rows} \\ \uparrow p \text{ rows} \end{array} \quad \dots (1)$$

Columns Columns

whose elements are

$$\left[\begin{array}{c|c} \mu(A_1) & O \\ \hline O & \mu'(A_1) \end{array} \right], \left[\begin{array}{c|c} \mu(A_2) & O \\ \hline O & \mu'(A_2) \end{array} \right], \dots, \left[\begin{array}{c|c} \mu(A_m) & O \\ \hline O & \mu'(A_m) \end{array} \right] \quad \dots (2)$$

Calling the first representation as μ_1 , second as μ_2 and their sum as μ we have

$$\mu_1 + \mu_2 = \mu \quad \dots (3)$$

Reducible representation. A representation arising from the representation (2) by similarity transformation is called as *reducible* representation and clearly these transformations are equivalent to the representation of the form (2). Other representations for which this is not possible are termed as *irreducible representations*.

e.g. A reducible matrix can be put in the form (2), by similarity transformation by means of converting j th row and column into j th row and column. In order to effect this reducible representation take isomorphic linear operator $T: L \rightarrow L'$, L, L' being two linear spaces and matrices, $A, B, \dots \in L, A', B', \dots \in L'$

We have

$$A' = TAT^{-1}, B' = TBT^{-1} \dots \text{etc.}$$

If we choose $T_{\alpha\beta} = \delta_{\alpha\beta}$, then $(T^{-1})_{jk} = \delta_{j'k}$

$$\text{and } \sum_{\beta} T_{\alpha\beta} (T^{-1})_{\beta j} = \sum_{\beta} \delta_{\alpha\beta} \delta_{\beta j} = \delta_{\alpha j}$$

So that the similarity transformation for this T resumes the required numbering such that

$$\bar{A} = T^{-1}AT$$

$$= A_{j'k'} \text{ where } (\bar{A})_{jk} = \sum_{\alpha\beta} \delta_{j'\alpha} A_{\alpha\beta} \delta_{\beta k'}$$

Now we know that every non-singular matrix $\mu(A)$ is invertible and multiplication of any group element A with identity element E gives A , so the multiplication of any representation matrix $\mu(A)$ with the matrix $\mu(E)$ assigned to the identity will yield $\mu(A)$ i.e.

$$\mu(A)\mu(E) = \mu(A) \text{ so that } \mu(E) = I, \text{ a unit matrix}$$

As such the unit matrix may be associated with the identity element of the group and we have

$$\mu(A)\mu(A^{-1}) = \mu(AA^{-1}) = \mu(E) = I$$

$$\text{i.e., } [\mu(A)]^{-1} = \mu(A^{-1}) \quad \dots (4)$$

In case of unitary representation i.e. matrices in representation being unitary, we therefore have

$$[\mu(A)]^{\dagger} = [\mu(A)]^{-1} \quad \dots (5)$$

$$(4) \text{ and } (5) \text{ give } \mu(A^{-1}) = [\mu(A)]^{\dagger} \quad \dots (6)$$

The reduction of a representation. If basis can be found such that all the matrices $\mu(A)$ of an n -dimensional representation may be reduced to the form

$$\mu(A) = \left[\begin{array}{c|c} \mu^{(1)}(A) & R(A) \\ \hline O & \mu^{(2)}(A) \end{array} \right] \begin{array}{l} \uparrow m \text{ rows} \\ \uparrow (n-m) \text{ rows} \end{array} \quad \dots (7)$$

$\begin{array}{c} \longleftrightarrow \\ m \\ \text{columns} \end{array} \quad \begin{array}{c} \longleftrightarrow \\ (n-m) \\ \text{columns} \end{array}$

Where $\mu^{(1)}(A)$ denote $m \times m$ matrices, $\mu^{(2)}(A)$ denote $(n-m) \times (n-m)$ matrices, O is a null matrix of $(n-m)$ rows and m columns and $R(A)$ denotes a rectangular matrix of m rows and $(n-m)$ columns, then the representation $\mu(A)$ is said to be *reducible* and the procedure is said to be a method of *reduction* of the representation $\mu(A)$.

Transforming the basis in m -dimensional space of $\mu^{(1)}$, all the matrices of $\mu^{(1)}(A)$ can be brought to the form (7) i.e.

$$\mu^{(1)}(A) = \left[\begin{array}{c|c} \mu^{(3)}(A) & R^{(1)}(A) \\ \hline O & \mu^{(4)}(A) \end{array} \right] \quad \dots (8)$$

$\mu^{(2)}(A)$ being p -dimensional and $\mu^{(4)}(A)$ being $(n-p)$ dimensional.

Continuing this process, we may get the set of matrices $\mu^{(i)}(A)$,

$\mu^{(2)}(A), \mu^{(3)}(A), \dots, \mu^{(\alpha-1)}(A), \mu^{(\alpha)}(A)$ which can not be further reduced and the dimension of irreducible representations m_j is $\sum_{j=1}^{\alpha} m_j$

In (7) if basis found is such that $R(A)=O$, then (7) becomes

$$\mu(A) = \begin{bmatrix} \mu^{(1)}(A) & \vdots & O \\ O & \vdots & \mu^{(2)}(A) \end{bmatrix} \quad \dots (9)$$

$$\text{i.e.} \quad \mu = \mu^{(1)} + \mu^{(2)} \quad \dots (10)$$

It follows that the representation $\mu(A)$ in this case is fully reducible and the reduction method is the reverse of addition.

If $\mu^{(1)}$ and $\mu^{(2)}$ are also reducible then continuing the process of reduction, the result (10) can be extended in the form

$$\mu = \mu^{(1)} + \mu^{(2)} + \mu^{(3)} + \dots + \mu^{(\alpha)} \quad \dots (11)$$

The irreducible representation $\mu^{(\alpha)}$ may contain several equivalent irreducible representations which are not counted distinctly. As such a representation μ may consist of a particular irreducible representation $\mu^{(k)}$ several times i.e.

$$\begin{aligned} \mu &= i_1 \mu^{(1)} + i_2 \mu^{(2)} + \dots + i_k \mu^{(k)} \\ &= \sum_j i_j \mu^{(j)}, \quad i_j \text{ being positive integers.} \end{aligned} \quad \dots (12)$$

As an illustration, consider a vector x with components x_1, x_2, x_3 , and the elements of the group as operators change x into a new vector x' with the same components in different order. Then the representation μ is a matrix s.t. $x' = \mu x$, rows and columns being labelled with x_1, x_2, x_3 .

Taking E as identity element of the group, $\mu(E)$ is a unit matrix

$$\text{i.e.} \quad \mu(E) = I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Assuming that A replaces x_1 by x_3 but x_1 itself becomes x_2 , that in $\mu(A)$, unity appears at the intersection of x_1 th row and x_2 th column. Also taking similar assumptions with B, C, D, F we have

$$\begin{aligned} \mu(A) &= \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \quad \mu(B) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \quad \mu(C) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \\ \mu(D) &= \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \text{ and } \mu(F) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

Note 1. Since in a space of l -dimensions at the most l orthogonal vectors can exist, therefore sum of squares of dimensions of all inequivalent, irreducible representations is at the most equal to the order of the group of representation i.e. $n_1^2 + n_2^2 + \dots + n_s^2 \leq h$, $n_1, n_2, \dots, n_s$ being dimensions of s inequivalent irreducible representations of a group of order l .

$$\text{So} \quad \sum_{k=1}^s n_k^2 \leq l.$$

Main Features of Reducible and Irreducible Representations

(i) The number of non-equivalent irreducible representations is the same as the number of classes.

(ii) If there are h elements in a group then the number of times the j th irreducible representation occurs in a reducible representation is given by $a_j = \frac{1}{h} \sum X X_j$.

(iii) All the irreducible representations of an abelian group are one-dimensional.

(iv) A representation by non-singular matrices can be transformed into a representation by unitary matrices through a similarity transformation.

(v) Any matrix commuting with all the matrices of an irreducible representation is a constant matrix i.e. a unit matrix multiplied by a constant scalar.

(vi) When a matrix A commutes with every matrix of a given representation of a group, then either A is a scalar matrix or the representation is reducible and the transformation used to diagonalize A wholly or partly reduces the representation.

(vii) If there are two irreducible representations $\mu(A_1), \mu(A_2), \dots, \mu(A_h)$ and $\mu'(A_1), \mu'(A_2), \dots, \mu'(A_h)$ of dimensions d_1 and d_2 respectively and if there exists a matrix M with d_2 rows and d_1 columns such that $\mu(A_j) = \mu'(A_j)M$; $j=1, 2, \dots, h$ then for $d_1 \neq d_2$, the matrix M is a null matrix whereas for $d_1 = d_2$, M is either a null matrix or a singular matrix.

(viii) The direct product of irreducible representations of two different groups is also an irreducible representation of the direct product of the groups.

Orthogonality Theorem for an Irreducible Representation

(i) $\mu(E), \mu(A_1), \mu(A_2), \dots, \mu(A_h)$ and $\mu'(E), \mu'(A_1), \dots, \mu'(A_h)$ be two inequivalent irreducible unitary representations of the same group G , then

$$\sum_R \mu(R)_{jk} \mu'(R)_{pq} = 0 \quad \dots (13)$$

for all elements j, k and p, q , where the summation extends over all group elements $E, A_1, A_2, \dots, A_h$, E being identity element.

Assuming the representation in unitary form as a similarity transformation always leaves multiples of the unit matrix unchanged, a matrix M commutes all the matrices $E(=A_1), A_2, \dots, A_h$ of the representation of the group of order h i.e.

$$A_i M = M A_i, \quad i=1, 2, 3, \dots, h \quad \dots (14)$$

By feature (vi) we have $M \mu(A_i) = \mu'(A_i) M$, $i=1, 2, \dots, h$ $\dots (15)$

Here (13) asserts that a matrix which satisfies (15), must be a null matrix and one which satisfies (14) must be a multiple of the identity matrix.

On account of group property of the representation all matrices of the form

$$M = \sum_R \mu'(R) X \mu(R^{-1}) \quad \dots (16)$$

for arbitrary l -rowed and m -columned matrix X satisfy (15). Also the group property follows that

$\sum_R \mu'(SR) X D\{S(R^{-1})\} = \sum_R \mu'(R) X D(R^{-1}) = M$, since the same matrices appear on the left and right except in different order.

Hence $\mu'(S)M = \sum_R \mu'(S)\mu'(R) X D(R^{-1}) = \sum_R \mu'(SR) X \mu'(SR^{-1})\mu(S)$ or $\mu'(S)M = M\mu'(S)$ (in coincide form), $\dots (17)$

so that by feature (vii) M must be a null matrix i.e. for arbitrary X_{ij} , $M_{pi} = \sum_{ir} \sum_{R} \mu'(R) X_{ir} \mu(R^{-1})_{rj}$; while on setting all matrix elements $X_{ir} = 0$ except one number $X_{qk} = 1$, the generalized form of (15) is $\sum_R \mu'(R)_{pq} \mu'(R^{-1})_{kj}$ where $\mu'(R)$ and $\mu(R)$ must be irreducible, but not necessarily unitary. In case $\mu'(R^{-1})_{kj}$ (R) are unitary $\mu(R^{-1}) = [\mu(R)]^{-1}$ and hence $\mu(R^{-1}) = \mu(R)^*$ so that (M) reduces to unitary representation.

The Character of Representation

Let μ^j with matrices $\mu^j(A)$ be a representation of a group G . Then trace of the matrix $\mu^j(A)$ i.e. the sum of diagonal elements of $\mu^j(A)$ is said to be the *character* of element A in the representation μ and denoted by

$$\chi^j(A) = \sum_k \{\mu^j(A)\}_{kk} \quad \dots (18)$$

The character of an irreducible element is known as **simple** and that of decomposable representation as **composite**. It is worth noting that the equivalent representation has the same set of characters, since if μ and μ' be equivalent representations with matrices $\mu(A)$ and $\mu'(A)$ of the group G , then

$$\mu'(A) = B^{-1} \mu(A) B, \quad B \text{ being some matrix.}$$

$\therefore t_r \mu'(A) = t_r \mu(A)$, trace of a matrix being unaltered under similarity transformation.

Also the character is a class function in the group.

Since if A, B be two conjugate elements of a group G , then $A = U^{-1} B U$,

$$\text{and } \mu(A) = \mu(U^{-1}) \mu(B) \mu(U) = \{\mu(U)\}^{-1} \mu(B) \mu(U)$$

$\mu(A)$ and $\mu(B)$ being equivalent representations (as related to similarity transformations) have same set of characters. As such conjugate elements in a representation correspond to the same character and hence in describing a group by listing the character of its elements in an assumed representation, same character i.e. number is assigned to all elements in a given class. Thus the character is a class function in the group i.e. the character of a single element from each class can

yield the character of the whole group.

Now if $\mu(A) = \mu_1(A) + \mu_2(A)$ (20)

$$\text{then } \chi(A) = t_r[\mu(A)] = t_r[\mu_1(A) + \mu_2(A)] = t_r[\mu_1(A)] + t_r[\mu_2(A)] = \chi_1(A) + \chi_2(A) \quad \dots (19)$$

Also if $\mu(A) = i_1 \mu^{(1)}(A) + i_2 \mu^{(2)}(A) + \dots + i_\alpha \mu^{(\alpha)}(A)$, by (11)

$$\text{then } \chi(A) = i_1 \chi^{(1)}(A) + i_2 \chi^{(2)}(A) + \dots + i_\alpha \chi^{(\alpha)}(A) \quad \dots (20)$$

Other Main Features of Characters

(i) The character of the direct product is the product of the characters i.e. $\chi^j(AB) = \chi^j(A) \cdot \chi^j(B)$.

(ii) The characters form an orthogonal system.

(iii) If both of the representations $\mu(A)$ and $\mu(B)$ are of the first degree then the direct product $\mu(AB)$ is irreducible. In case both are of degree higher than one, $\mu(AB)$ is reducible.

(iv) Two irreducible representations are equivalent if and only if they have the same character.

Character tables. These are the devices to find characters when the complete multiplication table for a group is known. It is effected by calculating first the product of all elements in the class C_i by all elements in C_j and then arranging uniquely the resulting set of elements in classes. Evidently a given class may occur in the products many times or not at all. If h_{ij} , k denotes the number of times the k th class appears, then we can write $C_i C_j = C_k \sum_{i=1}^p h_{ij, k} C_k$

$$\dots (21)$$

$$\text{whence } \sum_{i=1}^p [\chi^{(i)}]^2 = g \text{ (say)} \quad \dots (22)$$

where summation extends over all the numbers of classes p and g is known.

Once $h_{ij, k}$ determined, we can find characters by the use of relations

$$i_1 i_2 \chi^{(i)} \chi^{(j)} = \chi^{(i)} \sum_{k=1}^p h_{ij, k} r_k \chi^{(k)} \quad \dots (23)$$

where r_i is the number of elements in i -th class and r_j the number of elements in j th class.

A table in which we can put all that what we have explained here, is known as a **character table**.

e.g. consider a multiplication group of matrix elements

$$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix},$$

$$C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \quad D = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \quad F = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{ then we have}$$

$$C_2 = A^2, B^2, AB, BA = 2C_1 + C_2$$

$$C_3 = 3C_1 + 3C_2; C_2 C_3 = 2C_3$$

other products are not required since $r_1=1$, $r_2=2$ and $r_3=3$.

Using (24), we therefore have

$$4 [\chi^{(2)}]^2 = \chi^{(1)}(2\chi^{(1)} + 2\chi^{(2)})$$

$$9 [\chi^{(3)}]^2 = \chi^{(1)}(3\chi^{(1)} + 6\chi^{(2)})$$

$$6\chi^{(2)}\chi^{(3)} = 6\chi^{(1)}\chi^{(3)}$$

Since $\chi^{(1)}$ has values 1, 1, 2 [by (23), $g=6$ and $p=3$].

Solving these equations with each of 1, 1, 2 in turn, we get a table as shown here.

Here $\delta_1, \delta_2, \delta_3$ which are themselves matrices form the diagonal elements of the reducible representation.

	C_1	C_2	C_3
δ_1	1	1	1
δ_2	1	1	-1
δ_3	2	-1	0

Important Note. For detailed discussion of character tables in group theory, see Appendix C at the end of the book.

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 39. Show that for a finite group G , every representation is equivalent to a unitary representation.

Consider an arbitrary pair of vectors x, y s.t.

$$\{x, y\} = \sum_{A \in G} \{\mu(A)x, \mu(A)y\}, \mu \text{ being representation.} \quad \dots (1)$$

The sum in (1) being extended over all elements A of G , for any $B \in G$,

$$\begin{aligned} \text{we have } \{\mu(B)x, \mu(B)y\} &= \sum_{A \in G} \{\mu(A)\mu(B)x, \mu(A)\mu(B)y\}, \\ &= \sum_{A \in G} \{\mu(AB)x, \mu(AB)y\} \quad \dots (2) \end{aligned}$$

If A and B are fixed, AB runs through similar elements of G and so R.H.S's of (1) and (2) are identical i.e.

$$\{x, y\} = \{\mu(B)x, \mu(B)y\} \quad \dots (3)$$

For any operator U to be unitary, $(Ux, Uy) = (x, y) \forall x, y$.

So operators $\mu(A)$ are unitary w.r.t. scalar product $\{x, y\}$.

Let u_α be a set of vectors, orthogonal w.r.t. the set of original scalar product and v_α a second set of vectors orthogonal w.r.t. a new scalar product s.t.

$$\{u_\alpha, u_\beta\} = \delta_{\alpha\beta} = \{v_\alpha, v_\beta\}$$

Take an operator T s.t. $v_\alpha = T u_\alpha$

So that $Tx = Tx_\alpha u_\alpha = x_\alpha T u_\alpha = x_\alpha v_\alpha$

$$\therefore \{Tx, Ty\} = x_\alpha y_\alpha = (x, y) \quad \dots (4)$$

If the equivalent representation is given by

$$\mu'(B) = T^{-1} \mu(B) T \quad \dots (5)$$

Then $\{T^{-1} \mu(B) Tx, T^{-1} \mu(B) Ty\} = \{\mu(B)Tx, \mu(B)Ty\}$ by (4)

$$= \{Tx, Ty\} \text{ by (3)}$$

$$= (x, y) \text{ by (4)}$$

which follows that $\mu'(B)$ given by (5) is unitary and hence for finite group G , the representation can be chosen as unitary.

Allter. Let G be a group of order m represented by matrices $A_1, A_2, \dots, A_m$. A 's are distinct if the representation is faithful otherwise A 's are not distinct.

Consider Hermitian Matrix H , obtained by summing over all the group elements s.t.

$$H = \sum_{\alpha} A_{\alpha} A_{\alpha}^{\dagger} \quad \dots (6)$$

Matrix H can be diagonalized as D by unitary matrix U s.t.

$$\begin{aligned} D &= U^{-1} H U = \sum_{\alpha} U^{-1} A_{\alpha} A_{\alpha}^{\dagger} U \\ &= \sum_{\alpha} U^{-1} A_{\alpha} U U^{-1} A_{\alpha}^{\dagger} U \\ &= \sum_{\alpha} U^{-1} A_{\alpha} U (U^{-1} A_{\alpha} U)^{\dagger} \\ &= \sum_{\alpha} \bar{A}_{\alpha} \bar{A}_{\alpha}^{\dagger} U^{-1}, \quad (7) \quad \because \bar{A}_{\alpha} = U^{-1} A_{\alpha} U \quad \dots (7) \end{aligned}$$

All elements of D are real and positive since

$$\begin{aligned} D_{\beta\beta} &= \sum_{\alpha} \sum_{\gamma} (\bar{A}_{\alpha})_{\beta\gamma} (\bar{A}_{\alpha})_{\gamma\beta}^{\dagger} = \sum_{\alpha} \sum_{\gamma} |(\bar{A}_{\alpha})_{\beta\gamma}|^2 \\ &= 0 \text{ if } (\bar{A}_{\alpha})_{\beta\gamma} = 0 \forall \gamma \text{ and } \alpha \text{ in which case an entry row} \end{aligned}$$

of $\bar{A}_{\alpha}$ will be zero so that $|A_{\alpha}| = 0$ thereby contradicting the hypothesis.

Thus $D^{1/2}$ and $D^{-1/2}$ can be uniquely formed from D by taking $\pm 1/2$ power of the diagonal elements and $D^{1/2}$ and $D^{-1/2}$ will be real if

$$(D^{1/2})^{\dagger} = D^{1/2} \text{ and } (D^{-1/2})^{\dagger} = D^{-1/2}$$

But the representation $(\bar{A}_{\alpha}) = D^{-1/2} A_{\alpha} D^{1/2} = D^{-1/2} U^{-1} A_{\alpha} U D^{1/2}$ is unitary and (7) gives $I = D^{-1/2} \sum_{\alpha} \bar{A}_{\alpha} A_{\alpha} D^{-1/2}$

$$\begin{aligned} \therefore \bar{A}_{\lambda} \bar{A}_{\lambda}^{\dagger} &= D^{-1/2} \bar{A}_{\lambda} D^{1/2} (D^{-1/2} \sum_{\alpha} \bar{A}_{\alpha} A_{\alpha} D^{-1/2}) D^{1/2} \bar{A}_{\lambda}^{\dagger} D^{-1/2} \\ &= D^{-1/2} \sum_{\alpha} \bar{A}_{\lambda} \bar{A}_{\alpha} \bar{A}_{\alpha}^{\dagger} \bar{A}_{\lambda}^{\dagger} D^{-1/2} \end{aligned}$$

Also the group axioms for $\bar{A}_{\alpha}$ give that $\bar{A}_{\lambda} \bar{A}_{\alpha}, \alpha=1, 2, \dots, m$ are also $\bar{A}_{\alpha}$ in a different order, therefore

$$\sum_{\alpha} \bar{A}_{\lambda} \bar{A}_{\alpha} (\bar{A}_{\lambda} \bar{A}_{\alpha})^{\dagger} = \sum_{\alpha} \bar{A}_{\alpha} \bar{A}_{\alpha}^{\dagger}$$

$$\text{Thus } \bar{A}_{\lambda} \bar{A}_{\lambda}^{\dagger} = D^{-1/2} \sum_{\alpha} \bar{A}_{\lambda} \bar{A}_{\lambda}^{\dagger} D^{-1/2} = I$$

$$\text{So } \bar{A}_{\lambda}^{\dagger} = (\bar{A}_{\lambda})^{-1}$$

which shows that $\bar{A}_{\alpha}$ is unitary.

Problem 40. If a matrix commutes with all the matrices of an irreducible representation, then show that it is a multiple of unit matrix.

Let $A_1, A_2, \dots, A_m$ be the matrices in representation of a group G in unitary form and B be a matrix which commutes with all of $A_1, A_2, \dots, A_m$

$$i.e. \quad A_\alpha B = B A_\alpha, \alpha = 1, 2, \dots, m \quad \dots (1)$$

$$\therefore (A_\alpha B)^\ominus = (B A_\alpha)^\ominus \text{ (by taking transpose conjugate)}$$

$$i.e. \quad B^\ominus A_\alpha^\ominus = A_\alpha^\ominus B^\ominus \quad \dots (2)$$

$$\text{or} \quad A_\alpha B^\ominus A_\alpha^\ominus = A_\alpha^\ominus A_\alpha^\ominus B^\ominus A_\alpha^\ominus$$

$$\text{or} \quad A_\alpha B^* = B^* A_\alpha \quad \dots (3)$$

$$\therefore A_\alpha A_\alpha^\ominus = A_\alpha^\ominus A_\alpha^\ominus A^\ominus = I, A_\alpha \text{ being unitary. (1) and (3) follow}$$

that B and B^* both commute with all A 's.

$\therefore B + B^* = H_1$ (say), $i(B - B^*) = H_2$, H_1, H_2 being Hermitian; will also commute with all A 's. As such we can conclude that a matrix commutative with all the elements of a unitary representation is Hermitian.

Also assuming that B is unitary, it can be diagonalized as

$$D = U^{-1} B U$$

If A_α is unitary then $C_\alpha = U^{-1} A_\alpha U$ is also unitary and so (1) gives

$$C_\alpha D = D C_\alpha, \alpha = 1, 2, \dots, m$$

Equating j - k th elements on either side,

$$[C_\alpha D]_{jk} = [D C_\alpha]_{jk},$$

$$i.e. \quad (C_\alpha)_{kj} (D)_{jj} = D_{kk} (C_\alpha)_{kj} \quad \dots (4)$$

D being diagonal matrix.

Now $D_{jj} \neq D_{kk} \Rightarrow (C_\alpha)_{kj} = 0$ i.e. representation is reducible which contradicts the hypothesis and hence

$$D_{jj} = D_{kk} \forall j, k$$

which follows that D is a scalar matrix say $D = \lambda I$, λ being a constant and I a unit matrix.

$$\text{As such } B = U D U^{-1} = U \lambda I U^{-1} = \lambda U U^{-1} = \lambda I$$

= a multiple of unit matrix.

Problem 41. If U covers the entire unitary group then show that $\rho(\alpha, \beta, \gamma)$ ranges over all rotations.

$$\text{Take } U_1 = \begin{bmatrix} e^{i\alpha/2} & 0 \\ 0 & e^{-i\alpha/2} \end{bmatrix}$$

$$\text{Then,} \quad \begin{aligned} x' &= x \cos \alpha + y \sin \alpha \\ y' &= -x \sin \alpha + y \cos \alpha \\ z' &= z \end{aligned}$$

represent a rotation through an angle α about Z -axis. Representing it by $r' = \rho_\alpha(\alpha)$ r, r', r' being vectors with components (x, y, z) and (x', y', z') respectively, we have

$$\rho_\alpha(\alpha) = \begin{bmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Similarly $\rho_\alpha(\beta)$ corresponds to a matrix with α replaced by β .

$$\text{Now take } U_2 = \begin{bmatrix} \cos \beta/2 & i \sin \beta/2 \\ i \sin \beta/2 & \cos \beta/2 \end{bmatrix}$$

Then

$$U_2^\ominus T_1 U_2 = T_1$$

$$U_2^\ominus T_2 U_2 = \cos \beta T_2 + \sin \beta T_3$$

$$U_2^\ominus T_3 U_2 = -\sin \beta T_2 + \cos \beta T_3$$

and $r' = \rho_\alpha(B) r$

$$\text{whence } \rho_\alpha(\beta) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \beta & \sin \beta \\ 0 & -\sin \beta & \cos \beta \end{bmatrix}$$

Evidently the product of the unitary matrices $U_1(\alpha), U_2(\beta), U_3(\gamma)$ corresponds to the product of a rotation about z through an angle γ , about y , through β and about x through α i.e. corresponds to a rotation with Euler's angles α, β, γ . Hence the homomorphism is the homomorphism of the unitary group onto the entire 3-dimensional rotation group.

Problem 42. Show that the groups of order 2 and 3 are always cyclic.

Problem 43. Show that a group of order 4 may or may not be a cyclic group.

Problem 44. If $D(A)$ and $D(B)$ denote the determinants of two matrices A and B of any order then show that $D(A) D(B) = D(AB) = D(B) D(A)$

Problem 45. If $D(A)$ is the determinant of the matrix $A = \begin{bmatrix} a & c \\ b & d \end{bmatrix}$ then show that

$$B = \begin{bmatrix} d & -c \\ D(A) & D(A) \\ -b & a \\ D(A) & D(A) \end{bmatrix} \text{ is the inverse of } A, \text{ provided } D(A) \neq 0, \text{ and if } D(A) = ad - bc, \text{ then verify that } AB = BA = I.$$

$$AB = BA = I$$

Problem 46. If we define D_3 group as consisting of six matrix elements

$$E = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, B = \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ \sqrt{3}/2 & 1/2 \end{bmatrix}, C = \begin{bmatrix} -1/2 & -\sqrt{3}/2 \\ -\sqrt{3}/2 & 1/2 \end{bmatrix},$$

$$D = \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ -\sqrt{3}/2 & -1/2 \end{bmatrix}, F = \begin{bmatrix} -1/2 & -\sqrt{3}/2 \\ \sqrt{3}/2 & -1/2 \end{bmatrix}$$

they form the group table and show that this group is non-abelian.

The group table is as shown here and it is easy to show that

$$AB \neq BA \text{ etc.}$$

Also note that order of each of A, B, C is 2 and that of D, F is 3.

Problem 47. Find the classes of D_3 group.

Since in each group identity elements form their own class and no other class contains E , therefore one class of D_3 is E .

Now to find the elements conjugate to A , we have

$$E^{-1}AE = A; A^{-1}AA = A; B^{-1}AB = BAB = FB = C; C^{-1}AC = CAC = DF = B;$$

$$D^{-1}AD = FAD = BD = C; F^{-1}AF = DAF = CF = B.$$

So elements conjugate to A , are B and C and hence A, B, C form a class.

Again consider the elements conjugate to D ,

$$E^{-1}DE = D; A^{-1}DA = ADA = BA = F; B^{-1}DB = BDB = CB = F$$

$$C^{-1}DC = CDC = AC = F; D^{-1}DD = D; F^{-1}DF = DDF = FF = D.$$

i.e. F is the only element conjugate to D and so D, F form a class.

Conclusively D_3 has three classes: $\mathcal{C}_1 = E$

$$\mathcal{C}_2 = A, B, C$$

$$\mathcal{C}_3 = D, F.$$

Problem 48. Defining a commutator of x and y as $x^{-1}y^{-1}xy$, $x, y \in G$ (group) and denoted by $[x, y]$ show that its inverse is also commutator.

We have $[x, y] = x^{-1}y^{-1}xy = z$ (say), then $z^{-1} = y^{-1}x^{-1}yx = [y, x]$.

Problem 49. Define a group. Show that the group D_3 consisting of the symmetry elements which map an equilateral triangle onto itself is isomorphous with the permutation group of three numbers (a, b, c) .

Obtain a two dimensional representation of this group.

(Agra, 1972)

Problem 50. What is meant by the matrix representation of a group. Define the character of a group element in a particular representation. Prove that all the elements of a group constituting a class have the same character.

Obtain the independent irreducible representations of the permutational group of three elements (a, b, c) and hence the character table for the group. [Agra, 1971]

Problem 51. Give example of a group which has a subgroup and construct its multiplication table for its elements. [Agra, 1971]

Problem 52. State and prove the Orthogonality theorem for inequivalent, irreducible, unitary representations of a finite group and hence show that two inequivalent irreducible representations of a group can not have same characters and irreducible representations with equal characters are equivalent. [Rohilkhand, 1971]

See §4.14.

Problem 53. Define the character of representation of a group and show that it is a class function in the group.

Show that the characters of the irreducible representations of a finite group form an orthogonal system in the space of group elements. (Rohilkhand, 1971)

See § 4.14.

CHAPTER 5

COMPLEX VARIABLES

5.1. INTRODUCTION

Cantor, Dedekind and Weierstrass etc., extended the conception of rational numbers to a larger field known as real numbers which constitute rational as well as irrational numbers. Evidently the system of real numbers is not sufficient for all mathematical needs e.g. there is no real number (rational or irrational) which satisfies $x^2 + 1 = 0$. It was therefore felt necessary by Euler, Gauss, Hamilton, Cauchy, Riemann and Weierstrass etc. to extend the field of real numbers to the still larger field of complex numbers. Euler for the first time introduced the symbol i with the property $i^2 = -1$ and then Gauss introduced a number of the form $\alpha + i\beta$ which satisfies every algebraic equation with real coefficients. Such a number $\alpha + i\beta$ with $i = \sqrt{-1}$ and α, β being real, is known as a *complex number*.

5.2. DEFINITIONS

Complex numbers. An ordered pair of real numbers such as (x, y) is termed as a complex number. If we write

$$z = (x, y) \text{ or } x + iy, \text{ where } i = \sqrt{-1}, \text{ then}$$

x is called the *real part* and y the *imaginary part* of the complex number z and denoted by

$$x = R_z \text{ or } R_{(z)} \text{ or } Re_{(z)}$$

$$y = I_z \text{ or } I_{(z)} \text{ or } Im_{(z)}$$

Equality of complex numbers. Two complex numbers (x, y) and (x', y') are equal iff $x = x'$ and $y = y'$.

Modulus of a complex number. If $z = x + iy$ be a complex number then its modulus (or module) is denoted by $|z|$ and given by

$$|z| = |x + iy| = \sqrt{x^2 + y^2}$$

Evidently $|z| = 0$ iff $x = 0, y = 0$.

5.3. OPERATION OF FUNDAMENTAL LAWS OF ALGEBRA ON COMPLEX NUMBERS

Taking three complex numbers $z_1 = (x_1, y_1), z_2 = (x_2, y_2), z_3 = (x_3, y_3)$ we define the following operations:

[1] **Addition.** The sum of two complex numbers $z_1 = (x_1, y_1)$ and $z_2 = (x_2, y_2)$ (say) is defined as a complex number $z = (z_1 + z_2) = (x_1 + x_2, y_1 + y_2)$ such that its real part is the sum of real parts and imaginary part is the sum of imaginary parts of the given numbers.

(i) Addition is commutative. i.e. $z_1 + z_2 = z_2 + z_1$... (1)

Since we have $z_1 + z_2 = (x_1 + iy_1) + (x_2 + iy_2)$
 $= (x_1 + x_2) + i(y_1 + y_2)$
 $= (x_1 + x_2, y_1 + y_2)$
 $= (x_2 + x_1, y_2 + y_1)$, all the numbers being real
 $= (x_2 + iy_2) + (x_1 + iy_1)$
 $= z_2 + z_1$

(ii) Addition is associative. i.e. $z_1 + (z_2 + z_3) = (z_1 + z_2) + z_3$... (2)

Since, we have
 $z_1 + (z_2 + z_3) = x_1 + iy_1 + (x_2 + iy_2 + x_3 + iy_3)$
 $= (x_1 + iy_1 + x_2 + iy_2) + x_3 + iy_3$
 $= (z_1 + z_2) + z_3$

(iii) There exists an additive identity i.e. $z + o = z$... (3)

Since, we have $z + o = (x, y) + (o, o)$
 $= (x + o, y + o)$
 $= (x, y) = z$

(iv) There exists an additive inverse i.e. $z + (-z) = o$... (4)

Since, $z + (-z) = (x, y) + (-x, -y)$
 $= (x - x, y - y)$
 $= (o, o)$
 $= o$

Note. If $z = (x, y)$ then $-z = (-x, -y)$ is called as *additive inverse* of z .

[2] Subtraction. If $z_1 = (x_1, y_1)$ then $-z_1 = (-x_1, -y_1)$ etc.

$\therefore z_1 - z_2 = (x_1, y_1) + (-x_2, -y_2) = x_1 + iy_1 - x_2 - iy_2$
 $= (x_1 - x_2, y_1 - y_2)$... (5)

[3] Multiplication. We have $z_1 z_2 = (x_1 + iy_1)(x_2 + iy_2)$
 $= (x_1 x_2 - y_1 y_2) + i(x_1 y_2 + x_2 y_1)$

i.e. $(x_1, y_1)(x_2, y_2) = (x_1 x_2 - y_1 y_2, x_1 y_2 + x_2 y_1)$... (6)

(i) Multiplication is commutative. i.e. $z_1 z_2 = z_2 z_1$... (7)

Since, $z_1 z_2 = (x_1 x_2 - y_1 y_2, x_1 y_2 + x_2 y_1)$ by (6)
 $= (x_2 x_1 - y_2 y_1, y_2 x_1 + y_1 x_2)$
 $= (x_2 + iy_2)(x_1 + iy_1)$
 $= z_2 z_1$

(ii) Multiplication is associative. i.e. $z_1(z_2 z_3) = (z_1 z_2)z_3 = z_1 z_2 z_3$... (8)

Since $z_1(z_2 z_3) = (x_1, y_1)[x_2 x_3 - y_2 y_3, x_2 y_3 + x_3 y_2]$ by (6)
 $= [x_1(x_2 x_3 - y_2 y_3) - y_1(x_2 y_3 + x_3 y_2),$
 $x_1(x_2 y_3 + x_3 y_2) + y_1(x_2 x_3 - y_2 y_3)]$ by (6)
 $= [(x_1 x_2 - y_1 y_2)x_3 - (x_1 y_2 + x_2 y_1)y_3, (x_1 y_2 + x_2 y_1)x_3$
 $+ (x_1 x_2 - y_1 y_2)y_3]$ (on rearranging)

$$= [(x_1, y_1)(x_2, y_2)](x_3, y_3)$$

$$= (z_1 z_2)z_3$$

(iii) Multiplication is distributive. i.e. $(z_1 + z_2)z_3 = z_1 z_3 + z_2 z_3$... (9)

Since $(z_1 + z_2)z_3 = (x_1 + x_2, y_1 + y_2)(x_3, y_3)$
 $= [(x_1 + x_2)x_3 - (y_1 + y_2)y_3, (x_1 + x_2)y_3$
 $+ (y_1 + y_2)x_3]$ by (6)
 $= [(x_1 x_3 - y_1 y_3) + (x_2 x_3 - y_2 y_3), (x_1 y_3 + x_3 y_1)$
 $+ (x_2 y_3 + x_3 y_2)]$ (on arranging)
 $= (x_1 x_3 - y_1 y_3, x_1 y_3 + x_3 y_1) + (x_2 x_3 - y_2 y_3,$
 $x_2 y_3 + x_3 y_2)$
 $= (x_1, y_1)(x_3, y_3) + (x_2, y_2)(x_3, y_3)$
 $= z_1 z_3 + z_2 z_3$

(iv) There exists a multiplicative identity i.e. $z \cdot 1 = z$... (10)

where $1 = (1, 0)$ is the multiplicative identity known as *unity* for the system of complex numbers.

We have $z \cdot 1 = (x, y)(1, 0)$
 $= (x, y)$
 $= z$

(v) There exists a multiplicative inverse i.e. $z z^{-1} = 1$... (11)

If $z = (x, y)$, then $z^{-1} = (x, y)^{-1}$ so that we have to show that

$(x, y)(x, y)^{-1} = (1, 0)$
 $(x, y)^{-1} = (x', y')$, this becomes
 $(x, y)(x', y') = (1, 0)$

i.e. $(xx' - yy', xy' + yx') = (1, 0)$

which gives $xx' - yy' = 1$ (on equating real and imaginary parts)
 $xy' + yx' = 0$

Solving these equations we get

$$x' = \frac{x}{x^2 + y^2}, y' = \frac{-y}{x^2 + y^2} \text{ provided } x^2 + y^2 \neq 0$$

Hence the complex number (x, y) has a unique multiplicative inverse $\left(\frac{x}{x^2 + y^2}, \frac{-y}{x^2 + y^2}\right)$ which is also a complex number such

that $(x, y)\left(\frac{x}{x^2 + y^2}, \frac{-y}{x^2 + y^2}\right) = (1, 0)$

[4] Division. Consider an equation $z_1 z_2 = z'$

where $z_1 = (x_1, y_1)$, $z_2 = (x_2, y_2)$ and $z' = (x', y')$

Now $z_1 z_2 = (x_1 x_2 - y_1 y_2, x_1 y_2 + x_2 y_1) = z' = (x', y')$

which gives $x_1 x_2 - y_1 y_2 = x'$
 $x_1 y_2 + x_2 y_1 = y'$

Solving $x_2 = \frac{y_1 y' - x_1 x'}{x_1^2 + y_1^2}$, $y_2 = \frac{x_1 y' - x' y_1}{x_1^2 + y_1^2}$... (12)
provided $x_1^2 + y_1^2 \neq 0$ i.e. $|z_1| \neq 0$

Thus we have a unique solution and $z_2 = \frac{z'}{z_1}$ is the quotient.

[5] **Conjugate complex numbers.** If $z = x + iy$, then $x - iy$ is said to be the conjugate of complex number z and denoted by $\bar{z}$

$$\text{Evidently } (z_1 + z_2) = \bar{z}_1 + \bar{z}_2 \quad \dots (13)$$

$$z_1 z_2 = |z_1 z_2| \quad \dots (14)$$

$$\overline{z z} = (x + iy)(x - iy) = x^2 + y^2 = |z|^2 \quad \dots (15)$$

$$z + \bar{z} = 2x = 2R_x \text{ or } 2R(z) \quad \dots (16)$$

$$z - \bar{z} = 2iy = 2iI_x \text{ or } 2iI(z) \quad \dots (17)$$

5.4. GRAPHICAL REPRESENTATION. (ARGAND DIAGRAM)




Fig. 5.1

Here $|z| = |x + iy| = \sqrt{(x^2 + y^2)}$ is the measure of length OP .

If (r, θ) be the polar co-ordinates of the point P , the polar form of the complex number z is

$$z = r(\cos \theta + i \sin \theta) = re^{i\theta}.$$

Here the number r (being taken +ive) is called the **modulus** or **absolute value** of the complex number z and θ is called the **angle** or **argument** of z and usually written as $\arg z$, i.e., $|z| = r$ and $\arg z = \theta$.

Now the co-ordinates of a point P' which is conjugate of z are $\bar{z} = (x, -y)$ or $(r, -\theta)$ in polars.




Fig. 5.3

Argand Plane or Argand diagram or Complex plane or Gaussian plane.

Consider a point P in xy -plane. Let an ordered pair of values of x and y correspond to the co-ordinates of the point P . Then a complex number z may be made to correspond to the point P , where

$$z = x + iy.$$

Here z is called the complex co-ordinate of the point P .

In the adjoining figure, the x -axis is called the **real axis** or **axis of reals** and y -axis is called the **imaginary axis** or the **axis of imaginaries**.




Fig. 5.2

Since $\bar{z} = r\{\cos(-\theta) + i \sin(-\theta)\}$, geometrically the points P and P' represent z and $\bar{z}$ respectively and their situations are symmetrical about the axis of reals, i.e. x -axis. The conjugate of z is called the **reflection** or **image** of z in the real axis.

Note 1. The plane whose points are represented by complex numbers is known as

Note 2. The complex number z representing the point (x, y) is sometimes called as **Affix** of the point (x, y) .

Note 3. The **sum**, **difference**, **product** and **quotient** of complex numbers can be geometrically represented on the Argand plane as follows:

(i) **Sum.** Taking z_1 and z_2 two complex numbers represented by the points P and Q on Argand Plane and completing the parallelogram $OPRQ$, we observe that mid-points of its diagonals OR and PQ coincide, hence they bisect each other i.e.

If $z_1 = (x_1, y_1)$, $z_2 = (x_2, y_2)$, then mid-point of PQ is $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$ which is also the mid-point of OR showing that co-ordinates of R are $(x_1 + x_2, y_1 + y_2)$.

$$\text{But } z_1 + z_2 = (x_1 + x_2, y_1 + y_2)$$

Fig. 5.4

The sum $z_1 + z_2$ corresponds to a vector whose components are $x_1 + x_2$ and $y_1 + y_2$. As such the sum of two complex numbers z_1 and z_2 can be represented by a vector $(z_1 + z_2)$

$$\text{if } \vec{OP} = z_1, \vec{OQ} = z_2 \text{ then } \vec{OR} = \vec{OP} + \vec{PR} = \vec{OP} + \vec{OQ} = z_1 + z_2.$$

Hence the point R on Argand plane corresponds to the sum of two complex numbers z_1 and z_2 as shown in Fig. 5.4.

(ii) **Difference.** Taking $z_1 = (x_1, y_1)$ and $z_2 = (x_2, y_2)$ two complex numbers represented by the points P and Q on Argand Plane and completing the parallelogram $OQPR$, we see that the point R represents the complex number $z_1 - z_2$, since $z_1 - z_2 = (x_1 - x_2, y_1 - y_2)$ being a complex number corresponds to a vector whose components are $x_1 - x_2$ and $y_1 - y_2$ and

$$\text{if } \vec{OP} = z_1, \vec{OQ} = z_2$$

Fig. 5.5

$$\text{then } \vec{QR} = -z_2, \text{ so that}$$

$$\vec{OR} = \vec{OP} - \vec{OQ} = \vec{OP} + \vec{QR} = \vec{OP} + \vec{QP} = \vec{OR}$$



the difference of two complex numbers can be represented by a vector.

(iii) **Product.** If $z_1 = (x_1, y_1)$, $z_2 = (x_2, y_2)$ are two complex numbers, then

$$z_1 z_2 = (x_1 x_2 - y_1 y_2, x_1 y_2 + x_2 y_1) = (x_1 x_2 - y_1 y_2) + i(x_1 y_2 + x_2 y_1)$$

$$\text{Changing to polars by putting } x_1 = r_1 \cos \theta_1, y_1 = r_1 \sin \theta_1$$

$$x_2 = r_2 \cos \theta_2, y_2 = r_2 \sin \theta_2$$



where r_1, r_2 are the moduli and θ_1, θ_2 are arguments of z_1 and z_2 respectively, we have

$$\begin{aligned} z_1 z_2 &= r_1 r_2 [(\cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2) \\ &\quad + i(\cos \theta_1 \sin \theta_2 + \sin \theta_1 \cos \theta_2)] \\ &= r_1 r_2 [\cos (\theta_1 + \theta_2) + i \sin (\theta_1 + \theta_2)] \end{aligned}$$

$$\therefore |z_1 z_2| = r_1 r_2 = |z_1| |z_2|$$

$$\text{and } \arg (z_1 z_2) = \theta_1 + \theta_2 = \arg \theta_1 + \arg \theta_2$$

i.e. the modulus of product of two complex numbers is equal to the product of their moduli and argument of the product of two complex numbers is the sum of their arguments.

In general if there are n complex numbers $z_1, z_2, \dots, z_n$ with moduli $r_1, r_2, \dots, r_n$ and arguments $\theta_1, \theta_2, \dots, \theta_n$ respectively, then repeated application of the above result yields,

$$z_1 z_2 \dots z_n = r_1 r_2 \dots r_n [\cos (\theta_1 + \theta_2 + \dots + \theta_n) + i \sin (\theta_1 + \theta_2 + \dots + \theta_n)]$$

$$\text{so that } |z_1 z_2 \dots z_n| = r_1 r_2 \dots r_n = |z_1| |z_2| \dots |z_n|$$

$$\text{and } \arg (z_1 z_2 \dots z_n) = \theta_1 + \theta_2 + \dots + \theta_n = \arg z_1 + \arg z_2 + \dots + \arg z_n$$

i.e. the modulus of the product of any number of complex quantities is equal to the product of their moduli and the argument of the product of these complex numbers is equal to the sum of their arguments.




Fig. 5.6

In a particular case when $z_1 = z_2 = \dots = z_n = z$ (say), the above results may be summarised as

$$z^n = r^n (\cos n\theta + i \sin n\theta) \text{ under the assumptions}$$

$$r_1 = r_2 = \dots = r_n = r \text{ (say)}$$

$$\theta_1 = \theta_2 = \dots = \theta_n = \theta \text{ (say)}$$

$$\text{i.e. } |z^n| = r^n = |z|^n \quad \dots (3)$$

$$\text{and amp } z^n = n\theta = n \cdot (\text{amp } z) \quad \dots (4)$$

Also if $r=1$, we get the De Moivre's theorem for positive integral exponents such as $z^n = (\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta \quad \dots (5)$

[4] Quotient. Consider two complex numbers z_1 and z_2 such that

$$z_1 = x_1 + iy_1 = r_1 (\cos \theta_1 + i \sin \theta_1)$$

$$z_2 = x_2 + iy_2 = r_2 (\cos \theta_2 + i \sin \theta_2)$$

The quotient of complex numbers z_1 and z_2 is given by

$$\frac{z_1}{z_2} = \frac{r_1 (\cos \theta_1 + i \sin \theta_1)}{r_2 (\cos \theta_2 + i \sin \theta_2)} = \frac{r_1}{r_2} [\cos (\theta_1 - \theta_2) + i \sin (\theta_1 - \theta_2)]$$

$$\therefore \left| \frac{z_1}{z_2} \right| = \frac{r_1}{r_2} = \frac{|z_1|}{|z_2|} \quad \dots (8)$$

$$\text{and } \arg \left(\frac{z_1}{z_2} \right) = \theta_1 - \theta_2 = \arg z_1 - \arg z_2 \quad \dots (9)$$

i.e. the modulus of the quotient of two complex numbers is the quotient of their moduli and the argument of the quotient of two complex numbers is the difference of their arguments.

As a particular case defining the division as the inverse of multiplication we have $\frac{1}{z} = \frac{1}{r} [\cos (-\theta) + i \sin (-\theta)] = \frac{1}{r} [\cos \theta - i \sin \theta]$

$$\text{so that } \frac{1}{z^n} = z^{-n} = \frac{1}{r^n} (\cos n\theta - i \sin n\theta) = \left(\frac{1}{z} \right)^n \quad \dots (10)$$

which shows that De Moivre's theorem is valid when the exponent is any negative integer.

Geometrical representation of $\frac{z_1}{z_2}$ may be shown as below:

Let $\vec{OP}$ and $\vec{OQ}$ represent the vectors z_1 and z_2 in an Argand plane such that $|z_1| = OP$, $|z_2| = OQ$ and $\arg z_1 = \theta_1$, $\arg z_2 = \theta_2$

Rotate the line OP in clockwise direction through an angle $\theta_2 (= \arg z_2)$ such that its new position is OP' and $\angle POP' = \theta_2$. Take $OA = 1$ (unit length) on OX and draw a line AR to meet OP' in R such that $\angle OAR = \angle OQP$.

The point R thus obtained corresponds to the quotient $\frac{z_1}{z_2}$ and it may be justified as follows:

In similar triangles OAR and OQP we have

$$\begin{aligned} \frac{OR}{OA} &= \frac{OP}{OQ} \text{ i.e., } OR = \frac{OP}{OQ}, \because OA = 1 \\ &= \frac{|z_1|}{|z_2|} = \left| \frac{z_1}{z_2} \right| \end{aligned}$$

which shows that the radius vector of the point R is $\left| \frac{z_1}{z_2} \right|$

$$\text{Also } \angle AOR = \angle POR - \angle POX = \theta_2 - \theta_1 = -(\theta_1 - \theta_2)$$




Fig. 5.7

i.e. vectorial angle of R is $-(\theta_1 - \theta_2)$ which, when measured in positive sense is $\theta_1 - \theta_2$.

Hence the point R represents the quotient $\frac{z_1}{z_2}$.

Note 4. Multiplication of a Complex number by i .

Let z be a complex number with its modulus r and amplitude θ

$$\text{i.e. } z = r(\cos \theta + i \sin \theta)$$

$$\text{and } i = \cos \frac{\pi}{2} + i \sin \frac{\pi}{2}$$

$$\begin{aligned} \therefore iz &= \left(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2} \right) r(\cos \theta + i \sin \theta) \\ &= r \left[\cos \left(\frac{\pi}{2} + \theta \right) + i \sin \left(\frac{\pi}{2} + \theta \right) \right] \end{aligned} \quad \dots (11)$$

which follows that iz represents a vector obtained by rotating the vector z through a right angle in the positive direction.

Note 5. Extraction of roots.

Suppose that $z_0^n = z$, n being positive integer $\dots (12)$

We can express,

$$z = r(\cos \theta + i \sin \theta)$$

so that $z_0 = r_0(\cos \theta_0 + i \sin \theta_0)$ provided $z \neq 0$ and r_0, θ_0 are unknown.

$\therefore (12)$ gives, $r_0^n(\cos n\theta_0 + i \sin n\theta_0) = r(\cos \theta + i \sin \theta)$ by (7)

Measuring the angles in radians, we therefore, have

$r_0^n = r$, $n\theta_0 = \theta \pm 2m\pi$, m being zero or any positive integer which follow that r, r_0 being positive, r_0 is the positive n th root of r and $\theta_0 = \frac{\theta}{n} \pm \frac{2m\pi}{n}$ has n distinct values for $m = 0, 1, 2, \dots, n-1$.

As such there are n distinct solutions of (1), given by

$$z^{1/n} = z_0 = r^{1/n} \left(\cos \frac{\theta + 2m\pi}{n} + i \sin \frac{\theta + 2m\pi}{n} \right), m = 0, 1, 2, \dots, n-1. \quad \dots (13)$$

Which are n distinct values of $z^{1/n}$.

Here the length of each of the n vectors $z^{1/n}$ is the positive number $r^{1/n}$ and argument of one of these vectors is $\frac{\theta}{n}$ while the other arguments are obtained by adding multiples of $\frac{2\pi}{n}$ to $\frac{\theta}{n}$.

In particular $z=0$, (12) has the only solution $z_0=0$.

But $1 = \cos 0 + i \sin 0$, then n th roots of unity are given by

$$1^{1/n} = \cos \frac{2\pi m}{n} + i \sin \frac{2\pi m}{n}, m = 0, 1, 2, \dots, n-1 \quad \dots (14)$$

Taking $m=1$, the root of unity being a complex number and denoted by ω , is given by

$$\omega = \cos \frac{2\pi}{n} + i \sin \frac{2\pi}{n} \quad \dots (15)$$

According to De Moivre's theorem, the n th roots of unity are given by

$$1, \omega, \omega^2, \dots, \omega^{n-1} \quad \dots (16)$$

Which are the vertices of a regular polygon in complex plane, of n sides inscribed in the unit circle $|z|=1$ with one vertex at the point $z=1$.




Fig. 5.8

The case (i) of Fig. 5.8 shows for $n=3$ and case (ii) for $n=6$.

Now if ζ is a particular n th root of z , then we have the n roots of z as

$$\zeta, \zeta\omega, \zeta\omega^2, \dots, \zeta\omega^{n-1} \quad \dots (17)$$

since ζ multiplied by ω implies the increment of $\arg \zeta$ by the angle $\frac{2\pi}{n}$.

If m, n be two positive integers prime to each other, then (13) and (17) yield

$$(z^m)^{1/n} = (r^m)^{1/n} \left(\cos \frac{m\theta}{n} + i \sin \frac{m\theta}{n} \right) \omega^q, q = 0, 1, 2, \dots, n-1 \quad \dots (18)$$

$$\begin{aligned} (z^{1/n})^m &= (r^{1/n})^m \left[\left(\cos \frac{\theta}{n} + i \sin \frac{\theta}{n} \right) \omega^l \right]^m \\ &= (r^m)^{1/n} \left(\cos \frac{m\theta}{n} + i \sin \frac{m\theta}{n} \right) \omega^{lm}, l = 0, 1, 2, \dots, n-1 \end{aligned} \quad \dots (19)$$

The two sets of n numbers will be identical if the set ω^q and ω^{lm} coincide and then n numbers in either set can be written as $z^{m/n}$ i.e.

$$z^{m/n} = (r^m)^{1/n} \left[\cos \left\{ \frac{m}{n} (\theta + 2\pi m) \right\} + i \sin \left\{ \frac{m}{n} (\theta + 2\pi m) \right\} \right] m = 0, 1, 2, \dots, n-1 \quad \dots (20)$$

we may similarly define,

$$z^{-m/n} = (z^{1/n})^{-m} = (z^{-m})^{1/n}$$

8.8. PROPERTIES OF MODULI AND ARGUMENTS AND GEOMETRY OF COMPLEX NUMBERS

(A) Properties of Moduli

(i) The modulus of the product of two complex numbers is the product of their Moduli.

If there are two numbers z_1 and z_2 defined by $z_1 = (x_1, y_1)$ or (r_1, θ_1) in polars i.e. $z_1 = r_1(\cos \theta_1 + i \sin \theta_1) = r_1 e^{i\theta_1}$

$z_2 = (x_2, y_2)$ or (r_2, θ_2) in polars i.e. $z_2 = r_2 e^{i\theta_2}$

Then, $z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$

$$\begin{aligned} \text{So that } |z_1 z_2| &= |r_1 r_2 e^{i(\theta_1 + \theta_2)}| \\ &= r_1 r_2 |\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)| \\ &= r_1 r_2 \sqrt{\cos^2(\theta_1 + \theta_2) + \sin^2(\theta_1 + \theta_2)} \\ &= r_1 r_2 \\ &= |z_1| \cdot |z_2| \quad \dots (1) \end{aligned}$$

(2) The modulus of the sum of two complex numbers does never exceed the sum of their moduli.

Let z_1 and z_2 be two complex numbers and $\bar{z}_1$ and $\bar{z}_2$ their conjugates.

We have already mentioned that $|z|^2 = z\bar{z}$.

$$\begin{aligned} \therefore |z_1 + z_2|^2 &= (z_1 + z_2)(\bar{z}_1 + \bar{z}_2) \quad [\because z_1 \bar{z}_2 = r_1 e^{i\theta_1} \cdot r_2 e^{-i\theta_2}] \\ &= z_1 \bar{z}_1 + z_2 \bar{z}_2 + z_1 \bar{z}_2 + z_2 \bar{z}_1 = r_1 r_2 e^{i(\theta_1 - \theta_2)} \\ &= |z_1|^2 + |z_2|^2 + 2R(z_1 \bar{z}_2) \text{ and } z_2 \bar{z}_1 = r_1 r_2 e^{-i(\theta_1 - \theta_2)} \\ &\leq |z_1|^2 + |z_2|^2 + 2|z_1 z_2|, \quad \therefore z_1 \bar{z}_2 + z_2 \bar{z}_1 \\ &\leq \{ |z_1| + |z_2| \}^2. \quad = 2r_1 r_2 \cos(\theta_1 - \theta_2) = 2R(z_1 \bar{z}_1) \\ \therefore |z_1 + z_2| &\leq |z_1| + |z_2|. \end{aligned}$$

$$\begin{aligned} \text{Also if } z &= x + iy, \\ R(z) &= x \\ &\leq \sqrt{x^2 + y^2} \\ &\leq |z|. \end{aligned}$$

$$\begin{aligned} \text{Aliter. } |z_1 + z_2|^2 &= |x_1 + x_2 + i(y_1 + y_2)|^2 \\ &= (x_1 + x_2)^2 + (y_1 + y_2)^2 \quad \because |A + iB|^2 = A^2 + B^2 \\ &= x_1^2 + x_2^2 + y_1^2 + y_2^2 + 2(x_1 x_2 + y_1 y_2) \\ &= |z_1|^2 + |z_2|^2 + 2R(z_1 \bar{z}_2) \\ &\leq |z_1|^2 + |z_2|^2 + 2|z_1 z_2|, \quad \because |\bar{z}_2| = |z_2| \\ |z_1 + z_2| &\leq |z_1| + |z_2|. \quad \dots (2) \end{aligned}$$

(3) The modulus of difference of two complex numbers is greater than or equal to the difference of their moduli.

$$\begin{aligned} \text{Let } z_1 &= x_1 + iy_1 = r_1 (\cos \theta_1 + i \sin \theta_1) = r_1 e^{i\theta_1}, \\ z_2 &= x_2 + iy_2 = r_2 (\cos \theta_2 + i \sin \theta_2) = r_2 e^{i\theta_2}. \end{aligned}$$

$$\text{Then } \bar{z}_1 = r_1 e^{-i\theta_1} \text{ and } \bar{z}_2 = r_2 e^{-i\theta_2}$$

$$\begin{aligned} \text{and } -(z_1 \bar{z}_2 + z_2 \bar{z}_1) &= -[r_1 e^{i\theta_1} \cdot r_2 e^{-i\theta_2} + r_2 e^{-i\theta_2} \cdot r_1 e^{-i\theta_1}] \\ &= -r_1 r_2 [e^{i(\theta_1 - \theta_2)} + e^{-i(\theta_1 - \theta_2)}] \\ &= -r_1 r_2 \cos(\theta_1 - \theta_2) = -2R(z_1 \bar{z}_2). \end{aligned}$$

$$\begin{aligned} \therefore |z_1 - z_2|^2 &= (z_1 - z_2)(\bar{z}_1 - \bar{z}_2) \quad \because |z|^2 = z\bar{z} \\ &= z_1 \bar{z}_1 + z_2 \bar{z}_2 - (z_1 \bar{z}_2 + z_2 \bar{z}_1) \\ &= |z_1|^2 + |z_2|^2 - 2R(z_1 \bar{z}_2). \end{aligned}$$

$$\begin{aligned} \text{But } R(z_1 \bar{z}_2) &= R\{(x_1 + iy_1)(x_2 - iy_2)\} = x_1 x_2 + y_1 y_2, \\ |z_1| + |z_2| &= \sqrt{x_1^2 + y_1^2} \sqrt{x_2^2 + y_2^2}. \end{aligned}$$

$$\text{Now } R(z_1 \bar{z}_2) < |z_1| |z_2|,$$

$$\text{if } x_1 x_2 + y_1 y_2 < \sqrt{(x_1^2 + y_1^2)(x_2^2 + y_2^2)}$$

$$\text{or if } (x_1 x_2 + y_1 y_2)^2 < x_1^2 x_2^2 + y_1^2 y_2^2 + x_1^2 y_2^2 + y_1^2 x_2^2$$

$$\text{or if } 2x_1 x_2 y_1 y_2 < x_1^2 y_2^2 + y_1^2 x_2^2,$$

which is so, since arithmetic mean of two quantities is greater than their geometric mean.

$$\text{As such } -R(z_1 \bar{z}_2) > -|z_1| |z_2|.$$

$$\begin{aligned} \text{Hence } |z_1 - z_2|^2 &> |z_1|^2 + |z_2|^2 - 2|z_1| |z_2| \\ &> [|z_1| - |z_2|]^2. \end{aligned}$$

$$\therefore |z_1 - z_2| > ||z_1| - |z_2||. \quad \dots (3)$$

(B) Properties of Arguments.

(1) The argument of the product of two complex numbers is equal to the sum of their arguments.

Take z_1, z_2 two complex numbers with moduli r_1, r_2 and arguments θ_1, θ_2 respectively, so that

$$z_1 = r_1 (\cos \theta_1 + i \sin \theta_1) = r_1 e^{i\theta_1}, \quad z_2 = r_2 e^{i\theta_2}$$

$$\therefore z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)} = r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)]$$

$$\text{giving } \arg(z_1 z_2) = \theta_1 + \theta_2 = \arg z_1 + \arg z_2. \quad \dots (4)$$

The result may be generalized for any number.

(2) The argument of the quotient of two complex numbers is equal to the difference of their arguments.

Take z_1, z_2 two complex numbers with moduli r_1, r_2 and arguments θ_1, θ_2 , so that $z_1 = r_1 e^{i\theta_1}, z_2 = r_2 e^{i\theta_2}$

$$\text{then } \frac{z_1}{z_2} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)} = \frac{r_1}{r_2} [\cos(\theta_1 - \theta_2) + i \sin(\theta_1 - \theta_2)]$$

$$\therefore \arg\left(\frac{z_1}{z_2}\right) = \theta_1 - \theta_2 = \arg z_1 - \arg z_2 \quad \dots (5)$$

(C) Geometry of Complex Numbers

(1) $\arg \frac{z-a}{z-b}$ gives the angle between the lines joining a to z and b to z taken in the appropriate sense.

Take $\vec{AP} = z - a$ and $\vec{BP} = z - b$ as shown in Fig. 5.9.

$$\begin{aligned} \therefore \angle APB = \theta &= \arg \vec{AP} - \arg \vec{BP} \\ &= \arg(z - a) - \arg(z - b) \\ &= \arg \frac{z-a}{z-b} \quad 0 < \theta \leq \pi \end{aligned}$$

argument being negative as shown in Fig 5.9.




Fig. 5.9

In particular if $\theta = 90^\circ$, $\arg \frac{z-a}{z-b} = \pm \frac{\pi}{2}$ and $\frac{z-a}{z-b}$ is purely imaginary. ... (7)

(2) Equation of a straight line joining two points z_1 and z_2 in Argand plane. Referred to Fig 5.10, arg

$\frac{z-z_1}{z-z_2} = \pi$ or 0 according as z lies inside or outside the line joining A to B .

Fig. 5.10

In either case $\frac{z-z_1}{z-z_2}$ is purely real, so that

$$\frac{z-z_1}{z-z_2} = \left(\frac{\overline{z-z_1}}{\overline{z-z_2}} \right) = \frac{\bar{z}-\bar{z}_1}{\bar{z}-\bar{z}_2} \quad \text{i.e. } z(\bar{z}_1-\bar{z}_2) - \bar{z}(z_1-z_2) + (z_1\bar{z}_2 - z_2\bar{z}_1) = 0 \quad \dots (8)$$

is the required equation of the straight line joining z_1 and z_2 .

General Equation of a Line.

$z_1 - z_2$ being purely imaginary, $\overline{z_1 z_2} = \bar{z}_1 - \bar{z}_2$ is also purely imaginary.

Also $z_1 \bar{z}_2 - z_2 \bar{z}_1$ is purely imaginary since $z_1 \bar{z}_2 = \bar{z}_1 z_2$

Multiplying (8) by i , we have

$$i z(\bar{z}_1 - \bar{z}_2) - i \bar{z}(z_1 - z_2) + i(z_1 \bar{z}_2 - z_2 \bar{z}_1) = 0 \quad \dots (9)$$

each term of which now becomes real.

Setting $i(z_1 \bar{z}_2 - z_2 \bar{z}_1) = \lambda$, and $-i(z_1 - z_2) = \mu$; λ being real and μ a constant, so that

$$[-i(\bar{z}_1 - \bar{z}_2)] = i(\bar{z}_1 - \bar{z}_2) = \mu,$$

the equation (9) becomes

$$\bar{\mu} z + \mu \bar{z} + \lambda = 0 \quad \dots (10)$$

where λ, μ are constants but λ real.

Note 1. In particular if $|z-z_1| = |z-z_2|$ then we have the equivalent form

$$(z-z_1)(\bar{z}-\bar{z}_1) = (z-z_2)(\bar{z}-\bar{z}_2) \quad \text{on using } |z|^2 = z\bar{z} \quad \dots (11)$$

which represents the equation of right bisector of the line joining z_1 and z_2 .

Note 2. The equation (10) i.e. $\bar{\mu} z + \mu \bar{z} + \lambda = 0$ is the necessary and sufficient condition for z_1 to be the reflection of z_2 in the line.

(3) Equation of a circle with centre at z_0 and radius r .

If z is any point on the circle, then

$$|z-z_0| = r \quad \text{i.e. } |z-z_0|^2 = r^2$$

using $|z|^2 = z\bar{z}$, we have

$$(z-z_0)(\bar{z}-\bar{z}_0) = r^2$$

$$\text{or } z\bar{z} - z_0\bar{z} + (z_0\bar{z}_0 - r^2) - z\bar{z}_0 = 0 \quad \dots (12)$$

Setting $-z_0 = \mu - z_0\bar{z}_0 - r^2 = \lambda$, this becomes

$$z\bar{z} + \bar{\mu} z + \mu \bar{z} + \lambda = 0, \lambda \text{ being real} \quad \dots (13)$$

General Equation of a Circle

(13) can be written as $(z+\mu)(\bar{z}+\bar{\mu}) = \mu\bar{\mu} - \lambda$

$$\text{or } |z+\mu|^2 = \mu\bar{\mu} - \lambda$$

So (13) represents the general equation of a circle if λ is real and $\mu\bar{\mu} - \lambda > 0$.

(4) Equation of a circle through three points z_1, z_2, z_3 .

Let A, B, C represent the points z_1, z_2, z_3 respectively. Take a point $P(z)$ on the circle.

$$\therefore \arg \frac{z_3-z_1}{z_3-z_2} - \arg \frac{z-z_1}{z-z_2} = 0 \text{ or } \pi$$

according as case (i) or (ii) of Fig. 5.11 exists.

$$\text{i.e. } \arg \frac{z_3-z_1}{z_3-z_2} \Big/ \arg \frac{z-z_1}{z-z_2} = 0 \text{ or } \pi$$

$$\text{In either case } \operatorname{Im} \left[\frac{z_3-z_1}{z_3-z_2} \Big/ \frac{z-z_1}{z-z_2} \right] = 0$$

i.e. $\left(\frac{z_3-z_1}{z_3-z_2} \right) \Big/ \left(\frac{z-z_1}{z-z_2} \right)$ is purely real

$$\therefore \left(\frac{z_3-z_1}{z_3-z_2} \right) \Big/ \left(\frac{z-z_1}{z-z_2} \right) = \left(\frac{\bar{z}_3-\bar{z}_1}{\bar{z}_3-\bar{z}_2} \right) \Big/ \left(\frac{\bar{z}-\bar{z}_1}{\bar{z}-\bar{z}_2} \right) \quad \dots (14)$$

is the required equation.

Note 3. Condition for four points z_1, z_2, z_3, z_4 to be collinear is that

$$\frac{(z_3-z_1)(z_4-z_2)}{(z_3-z_2)(z_4-z_1)} \text{ is purely real.} \quad \dots (15)$$




Fig. 5.12

(5) Inverse points with respect to a circle.

Two points $P(p), Q(q)$ are said to be the inverse points with respect to a circle with centre $O(z)$ and radius r if

$$OP \cdot OQ = r^2 \quad \dots (16)$$

provided O, P, Q are collinear.

$$\therefore |p-z| |q-z| = r^2$$

Also $\arg(p-z) = \arg(q-z) = -\arg(\overline{q-z})$ gives

$$\arg(p-z) + \arg(\overline{q-z}) = 0$$

i.e. $(p-z)(\overline{q-z})$ is purely real say equal to r^2 so that

$$(p-z)(\overline{q-z})=r^2 \quad \dots (17)$$

gives the required condition for p, q to be inverse points with respect to the circle with centre z and radius r .

Note 4. If $z=0, p, q$ are inverse points provided

$$p\overline{q}=r^2 \quad \dots (18)$$

Note 5. In w plane, $w=0$ and $w=\infty$ are the inverse points with respect to the circle $|w|=1$, since if $w=w_1=0$ and $w=w_2=\infty$ are inverse points then from (18), $w_1\overline{w_2}=1$ or $w_1w_2=1$ i.e. $w_1w_2=1$

$$\therefore w_2 = \frac{1}{w_1} = \frac{1}{0} = \infty$$

Note 6. In z plane, z and $\frac{1}{\overline{z}}$ are inverse points with respect to the unit circle $|z|=1$

Since if z_1, z_2 be inverse points then (18) gives

$$z_1\overline{z_2}=1 \text{ i.e. } z_1=\frac{1}{\overline{z_2}}=\frac{1}{\overline{z}} \text{ when } z_2=z$$

Note 7. In w plane, w and $\frac{1}{\overline{w}}$ are inverse points with respect to $|w|=1$.

Note 8. The equation $\left|\frac{z-p}{z-q}\right|=\lambda, \lambda$ being a positive parameter, represents a family of circles for every member of which p and q are inverse points. If $\lambda=1$, then the locus of z is the right bisector of the join of p and q .

Problem 1. Find moduli and arguments of the following complex numbers:

$$(i) \frac{1-i}{1+i}, (ii) \left(\frac{2+i}{3-i}\right)^2$$

$$(i) \text{ we have } \frac{1-i}{1+i} = \frac{1-i}{1+i} \times \frac{1-i}{1-i} = \frac{-2i}{2} = -i$$

$$\therefore \left|\frac{1-i}{1+i}\right| = |-i| = \sqrt{0^2+(-1)^2}=1$$

$$\text{and arg } \frac{1-i}{1+i} = \arg(-i) = -\frac{\pi}{2} \text{ as } -i = \cos\left(-\frac{\pi}{2}\right) + i \sin\left(-\frac{\pi}{2}\right)$$

$$(ii) \text{ we have } \left(\frac{2+i}{3-i}\right)^2 = \frac{3-4i}{8-6i} \times \frac{8+6i}{8+6i} = \frac{-14i}{28} = -\frac{1}{2}i$$

$$\therefore \left|\left(\frac{2+i}{3-i}\right)^2\right| = \left|-\frac{1}{2}i\right| = \frac{1}{2}|-i| = \frac{1}{2}$$

$$\text{and arg } \left(\frac{2+i}{3-i}\right)^2 = \arg\left(-\frac{1}{2}i\right) = \frac{1}{2} \arg(-i) = \frac{1}{2}\left(-\frac{\pi}{2}\right) = -\frac{\pi}{4}$$

Problem 2. Prove that $\arg(z) - \arg(-z) = \pm\pi$ according as $\arg(z)$ is positive or negative.

when $\arg(z)$ is positive, let $\arg(z)=\theta, -\pi < \theta < \pi$

Then $\arg(-z) = -(\pi-\theta)$ if $0 < \theta < \pi$.

$$= \pi + \theta \text{ if } 0 > \theta > -\pi$$

$\arg(z) - \arg(-z) = \theta + (\pi - \theta)$ or $\theta - (\pi + \theta)$ according as $\arg(z)$ is positive or negative.

$$= \pi \text{ or } -\pi \text{ according as } \arg(z) \text{ is positive or negative.}$$

Problem 3. Prove that $\arg z + \arg \overline{z} = 2n\pi, n$ being an integer including zero.

$$\text{If } z = x + iy \text{ then } \overline{z} = x - iy$$

$$\therefore \arg z + \arg \overline{z} = \arg(z\overline{z})$$

$$= \arg(x+iy)(x-iy)$$

$$= \arg(x^2 + y^2)$$

$$= \arg m, \text{ where } m = x^2 + y^2$$

Obviously m is real and positive.

If $m = r \cos \theta, 0 = r \sin \theta$ then $m = r$ and $\cos \theta = 1, \sin \theta = 0$ so that $\theta = 2n\pi$, where n is an integer including zero.

Hence $\arg z + \arg \overline{z} = 2n\pi, n$ being an integer including zero.

Problem 4. Show that $|z_1 - z_2|^2 + |z_1 + z_2|^2 = 2|z_1|^2 + 2|z_2|^2$ and deduce that $|\alpha + \sqrt{\alpha^2 - \beta^2}| + |\alpha - \sqrt{\alpha^2 - \beta^2}| = |\alpha + \beta| + |\alpha - \beta|$, all the numbers concerned being complex.

We have,

$$\begin{aligned} |z_1 + z_2|^2 + |z_1 - z_2|^2 &= (z_1 + z_2)(\overline{z_1 + z_2}) + (z_1 - z_2)(\overline{z_1 - z_2}) \\ &= 2z_1\overline{z_1} + 2z_2\overline{z_2} \\ &= 2|z_1|^2 + 2|z_2|^2 \end{aligned} \quad \dots (1)$$

$$\begin{aligned} \text{Now } \{|\alpha + \sqrt{\alpha^2 - \beta^2}| + |\alpha - \sqrt{\alpha^2 - \beta^2}|\}^2 &= |\alpha + \sqrt{\alpha^2 - \beta^2}|^2 \\ &+ |\alpha - \sqrt{\alpha^2 - \beta^2}|^2 + 2|\alpha + \sqrt{\alpha^2 - \beta^2}||\alpha - \sqrt{\alpha^2 - \beta^2}| \\ &= 2|\alpha|^2 + 2|\sqrt{\alpha^2 - \beta^2}|^2 + 2|\alpha^2 - (\alpha^2 - \beta^2)| \\ &= 2|\alpha|^2 + 2|\alpha^2 - \beta^2| + 2|\beta^2| \\ &= 2\{|\alpha|^2 + |\beta|^2\} + 2|\alpha^2 - \beta^2| \\ &= |\alpha + \beta|^2 + |\alpha - \beta|^2 + 2|\alpha + \beta||\alpha - \beta| \\ &= \{|\alpha + \beta| + |\alpha - \beta|\}^2 \end{aligned}$$

$$\therefore |\alpha + \sqrt{\alpha^2 - \beta^2}| + |\alpha - \sqrt{\alpha^2 - \beta^2}| = |\alpha + \beta| + |\alpha - \beta|$$

Problem 5. Show that an expression of the form $z^{2n} - 1$ can be expressed as a product of n real quadratic factors.

Assuming $z^{2n} - 1 = 0$ i.e. $z^{2n} = 1 = \cos 2m\pi + i \sin 2m\pi, m=0, 1, 2, \dots, n$

We have $z = \cos \frac{2m\pi}{2n} + i \sin \frac{2m\pi}{2n}$ applying De Moivre's result.

Here $m=0$ corresponds $z=1$ and $m=n \rightarrow z=-1$

Consider $z^{2n}-1=(z^2-1) \times (2n-2)$ factors obtained by putting $m=1, 2, \dots, n-1$ in the expression $z-\cos \frac{2m\pi}{2n} \pm i \sin \frac{2m\pi}{2n}$

$$\begin{aligned} \text{Now } \left(z - \cos \frac{2m\pi}{2n} - i \sin \frac{2m\pi}{2n} \right) \left(z - \cos \frac{2m\pi}{2n} + i \sin \frac{2m\pi}{2n} \right) \\ = \left(z - \cos \frac{2m\pi}{2n} \right)^2 + \sin^2 \frac{2m\pi}{2n} \\ = z^2 - 2z \cos \frac{2m\pi}{2n} + 1 \end{aligned}$$

$$\begin{aligned} \therefore z^{2n}-1 &= (z-1)(z+1) \left(z^2 - 2z \cos \frac{2\pi}{2n} + 1 \right) \left(z^2 - 2z \cos \frac{4\pi}{2n} + 1 \right) \dots \\ &\dots \left(z^2 - 2z \cos \frac{2n-4}{2n}\pi + 1 \right) \left(z^2 - 2z \cos \frac{2n-2}{2n}\pi + 1 \right) \\ &= (z^2-1) \left(z^2 - 2z \cos \frac{\pi}{n} + 1 \right) \left(z^2 - 2z \cos \frac{2\pi}{n} + 1 \right) \dots \\ &\dots \left(z^2 - 2z \cos \frac{n-2}{n}\pi + 1 \right) \left(z^2 - 2z \cos \frac{n-1}{n}\pi + 1 \right) \\ &= (z^2-1) \prod_{m=1}^{n-1} \left\{ z^2 - 2z \cos \frac{m\pi}{n} + 1 \right\}, \text{ II denotes product of} \end{aligned}$$

similar factors.

5.6. REGULAR FUNCTIONS

Before going into the details of Regular or Analytic functions, we first define some terms which are used frequently.

Neighbourhood of a point. Neighbourhood of a point z_0 in the Argand diagram means the set of all points z such that $|z-z_0| < \epsilon$, where ϵ is an arbitrarily chosen small positive number.

Limit point. A point z_0 is said to be a *limit point* of a set of points S in the Argand plane, if every neighbourhood of z_0 contains a point of S other than zero.

The limit points of a set may not necessary be the points of the set. There are two types of limit points:

(i) **Interior points.** A limit point z_0 of the set S is said to be the *interior* or *inner point* if in the neighbourhood of z_0 there exists entirely the points of the set S .

(ii) **Boundary points.** A limit point z_0 is said to be the *boundary point* if all the points in the neighbourhood of z_0 do not belong to the set S .

Closed set. If all the limit points of the set belong to the set, then the set is said to be a *closed set*.

Open set. A set which consists entirely of interior points is known to be an *open set*.

Bounded and unbounded sets. A set of points is said to be *bounded* if there exists a constant number k , such that $|z| \leq k$ for all points

of the set. If there does not exist such number k the set is said to be *unbounded*.

Domain. If every pair of the points of a set of points in the Argand diagram can be joined by a polygonal arc which consists only of the points of the set, then the set of points in the Argand diagram is said to be *connex* (means connected) or *domain* or *region*.

Open domain is an open connex set of points.

Closed domain. When the boundary points of the set are also added to an open domain, it is then called a *closed domain*.

Functions of a complex variable. If $w=u+iv$ and $z=x+iy$ are two complex numbers, then w is said to be the *function* of z and written as $w=f(z)$, if to every value of z in a certain domain D , there correspond one or more values of w . If w takes only one value for each value of z in the domain D , then w is said to be *uniform* or *single-valued* function of z and if it takes more than one values for some or all values of z in the domain D , then w is known as a *many-valued* or a *multiple-valued* function of z .

Since u and v both are functions of x, y

$$w=f(z)=u(x, y)+iv(x, y).$$

It is however notable that the path of a complex variable z is either a straight line or a curve.

Continuity. The function $f(z)$ of a complex variable z is *continuous* at the point z_0 if, given a positive number $\epsilon > 0$, a number δ can be so found that

$$|f(z)-f(z_0)| < \epsilon,$$

for all points z of the domain D satisfying $|z-z_0| < \delta$, where δ depends upon ϵ and also, in general, upon z_0 , i.e.

$$\delta = \phi(\epsilon, z_0).$$

If δ is independent of z_0 or rather say that if a number $h(\epsilon)$ can be found independent of z_0 such that $|f(z)-f(z_0)| < \epsilon$ holds for every pair of points z, z_0 of the domain D for which $|z-z_0| < h$, then $f(z)$ is called *uniformly continuous* in D .

It should be noted that if a function f is continuous at $z=z_0$ i.e. if $f=u+iv$ is continuous at $z=z_0$ then it will be so iff its real and imaginary parts are separately continuous functions of x and y at the point $(x, y)=(x_0, y_0)$

Since if f is continuous at $z=z_0$ then $u(x_0, y_0)$ and $v(x_0, y_0)$ both are uniquely defined such that

$$0 \leq |u(x, y)-u(x_0, y_0)| \leq |f(z)-f(z_0)| \dots (1)$$

$$\text{for, } |f(z)-f(z_0)| = + \{ [u(x, y)-u(x_0, y_0)]^2 + [v(x, y)-v(x_0, y_0)]^2 \}^{1/2} \\ \text{as } z \rightarrow z_0, u(x, y) \rightarrow u(x_0, y_0) \text{ i.e. } u(x_0, y_0) = \lim_{(x, y) \rightarrow (x_0, y_0)} u(x, y) \dots (2)$$

This limit exists independent of the manner in which $x \rightarrow x_0, y \rightarrow y_0$.

(2) shows that $u(x, y)$ is continuous at (x_0, y_0) .

Similarly $v(x, y)$ is continuous at (x_0, y_0)

Thus continuity of u and v for f to be continuous at $z=z_0$, is a necessary condition.

Conversely if $u(x, y)$ and $v(x, y)$ are continuous, then

$$u(x, y) \rightarrow u(x_0, y_0) \text{ and } v(x, y) \rightarrow v(x_0, y_0) \text{ as } z \rightarrow z_0$$

so that $f(z) = u(x, y) + iv(x, y) \rightarrow u(x_0, y_0) + iv(x_0, y_0) = f(z_0)$

So the condition is also sufficient.

Differentiability. If $f(z)$ be a single-valued function defined in a domain D of the Argand diagram, then $f(z)$ is said to be *differentiable* at $z=z_0$ a point of D if $\frac{f(z)-f(z_0)}{z-z_0}$ tends to a unique limit when $z \rightarrow z_0$, provided that z is also a point of D .

A function $f(z)$ is said to be differentiable at a point z_0 , if

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \text{ exists and is a finite quantity provided by}$$

whatever path $z \rightarrow z_0$; then $f(z)$ is differentiable at $z=z_0$. The finite limit when exists is denoted by $f'(z_0)$ and termed as the differential coefficient or derivative of $f(z)$ at $z=z_0$, i.e.

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

Precisely if for a given $\epsilon > 0$, there exists a number δ such that

$$\left| \frac{f(z) - f(z_0)}{z - z_0} - f'(z_0) \right| < \epsilon \text{ whenever } 0 < |z - z_0| < \delta$$

i.e. writing $z - z_0 = \Delta z$ if for an $\epsilon > 0$, there exists a number δ

such that $\left| \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} - f'(z_0) \right| < \epsilon$ whenever $0 < |\Delta z| < \delta$

then $f'(z_0)$ is known as the derivative of $f(z)$ at z_0 .

Clearly the limiting value of $\frac{f(z) - f(z_0)}{z - z_0}$ is independent of the path in D along which $z \rightarrow z_0$.

Consider $f(z) = z^2$, for example, then $f'(z_0) = 2z_0$ at any point z_0

$$\text{since } f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{(z_0 + \Delta z)^2 - z_0^2}{\Delta z} = \lim_{\Delta z \rightarrow 0} (2z_0 + \Delta z) = 2z_0$$

$$\text{In view of the relation } f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} \text{ whenever}$$

$f'(z_0)$ exists at any point z_0 , we have

$$\lim_{\Delta z \rightarrow 0} [f(z_0 + \Delta z) - f(z_0)] = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} \lim_{\Delta z \rightarrow 0} \Delta z = 0$$

which follows $\lim_{z \rightarrow z_0} f(z) = f(z_0)$

i.e. f is necessarily continuous at any point z_0 where its derivative exists. But the converse is not true i.e. if a function is continuous, it is not necessarily differentiable as is evident from the following example.

Consider the function $w = |z|^2$ which is differentiable at every point. It will be shown that its derivative exists only at the point $z=0$ and nowhere else, since

$$\begin{aligned} \frac{\Delta w}{\Delta z} &= \frac{|z_0 + \Delta z|^2 - |z_0|^2}{\Delta z} = \frac{(z_0 + \Delta z)(\bar{z}_0 + \bar{\Delta z}) - z_0 \bar{z}_0}{\Delta z} \\ &= \bar{z}_0 + \bar{\Delta z} + z_0 \frac{\bar{\Delta z}}{\Delta z} \end{aligned} \quad \because |z|^2 = z\bar{z} \quad \dots (3)$$

$$\lim_{\Delta z \rightarrow 0} \frac{\Delta w}{\Delta z} = \frac{dw}{dz} = \lim_{\Delta z \rightarrow 0} \left[\bar{z}_0 + \bar{\Delta z} + z_0 \frac{\bar{\Delta z}}{\Delta z} \right] = 0 \text{ when } z_0 = 0$$

But if $z_0 \neq 0$, then taking $\theta = \arg \Delta z = \arg (z - z_0)$ we have

$$\begin{aligned} \frac{\bar{\Delta z}}{\Delta z} &= \frac{e^{-i\theta}}{e^{i\theta}} = e^{-2i\theta} \\ &= \cos 2\theta - i \sin 2\theta. \end{aligned}$$

$$\text{So that } \lim_{\Delta z \rightarrow 0} \frac{\Delta w}{\Delta z} = \lim_{\Delta z \rightarrow 0} [\bar{z}_0 + \bar{\Delta z} + z_0 (\cos 2\theta - i \sin 2\theta)]$$

Here the $\lim_{\Delta z \rightarrow 0} \frac{\bar{\Delta z}}{\Delta z}$ does not exist as $\Delta z \rightarrow 0$ in any manner, since

if Δz is real, $\Delta z = \Delta x$ i.e. $\bar{\Delta z} = \Delta x = \Delta z$ then limit of (3) is $\bar{z}_0 + z_0$.

Also if Δz is imaginary i.e. $\Delta z = i \Delta y$ so that $\bar{\Delta z} = -\Delta z$, then limit of (3) is $\bar{z}_0 - z_0$. As such the limit does not exist when $z_0 \neq 0$ and hence $|z|^2$ has no derivative at z_0 .

Analytic (or regular or holomorphic or monogenic functions) [Agra 1965]

A function $f(z)$ which is single-valued and differentiable at every point of a domain D , is said to be *regular* in the domain D .

A function may be differentiable in a domain D save possibly for a finite number of points. Such points are called *singularities* or *singular points* of $f(z)$.

The necessary and sufficient conditions for $f(z)$ to be regular.

Necessary conditions.

If $w = f(z)$, where $w = u + iv$ and $z = x + iy$

As such u and v both are the functions of x and y and therefore we can write $w = f(z) = u(x, y) + iv(x, y)$.

Now if $f(z) = u(x, y) + iv(x, y)$ is differentiable at a given point z , the ratio $\frac{f(z + \Delta z) - f(z)}{\Delta z}$ must tend to a certain finite limit as $\Delta z \rightarrow 0$ in any manner.

From the relation $z = x + iy$, we get $\Delta z = \Delta x + i \Delta y$.

If we take Δz to be wholly real, so that $\Delta y = 0$, then

$\lim_{\Delta x \rightarrow 0} \frac{\Delta w}{\Delta x} = \lim_{\Delta x \rightarrow 0} \left[\frac{u(x+\Delta x, y) - u(x, y)}{\Delta x} + i \frac{v(x+\Delta x, y) - v(x, y)}{\Delta x} \right]$
must exist and tend to a definite limit.

$$\therefore \frac{dw}{dx} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \\ = u_x + i v_x \text{ (say),}$$

i.e. the partial derivatives u_x, v_x must exist at the point (x, y) and the limiting value is $u_x + i v_x$.

Similarly again if Δz be taken wholly imaginary, so that $\Delta x = 0$, we find that the partial derivatives u_y, v_y must exist at the point (x, y) and the limiting value is $v_y - i u_y$.

Since the function is differentiable, the two limits so obtained must be identical, i.e. $u_x + i v_x = v_y - i u_y$.

Equating real and imaginary parts, we get

$$u_x = v_y \text{ and } u_y = -v_x$$

or
$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

(Agra, 1967, 69, 71, 73)

These two relations, which are necessary conditions for a function to be analytic, are called the *Cauchy Riemann Differential Equations*.

Sufficient conditions. The continuous single-valued function $f(z)$ is regular in a domain C if the four partial derivatives u_x, u_y, v_x, v_y exist, are continuous and satisfy the Cauchy-Riemann equations at all points of the region D .

Assuming $u_x = v_y$ and $u_y = -v_x$ and these partial derivatives are continuous, we have to show that they exist and are finite.

By the mean value theorem, we have

$$\left. \begin{aligned} f(x+\Delta x, y) - f(x, y) &= \Delta x \frac{f(x+\Delta x\theta, y)}{\Delta x}, \\ &\quad \text{where } 0 < \theta < 1 \\ \text{and } f(x, y+\Delta y) - f(x, y) &= \Delta y \frac{f(x, y+\Delta y\theta')}{\Delta y}, \\ &\quad \text{where } 0 < \theta' < 1. \end{aligned} \right\} \dots (i)$$

Now if $w = f(z)$ and $f(z) = u(x, y) + i v(x, y)$,

$$w + \Delta w = f(z + \Delta z).$$

and
$$f(z + \Delta z) = u(x + \Delta x, y + \Delta y) + i v(x + \Delta x, y + \Delta y).$$

$$\therefore \Delta w = f(z + \Delta z) - f(z),$$

Also if $z = x + i y$, then $\Delta z = \Delta x + i \Delta y$.

Thus

$$\begin{aligned} \frac{\Delta w}{\Delta z} &= \frac{f(z + \Delta z) - f(z)}{\Delta x + i \Delta y} \\ &= \frac{u(x + \Delta x, y + \Delta y) + i v(x + \Delta x, y + \Delta y) - u(x, y) - i v(x, y)}{\Delta x + i \Delta y} \end{aligned}$$

$$\begin{aligned} \text{Now } u(x + \Delta x, y + \Delta y) - u(x, y) &= \{u(x + \Delta x, y + \Delta y) - u(x + \Delta x, y)\} + \{u(x + \Delta x, y) - u(x, y)\} \\ &= \Delta y \frac{\Delta u(x + \Delta x, y + \theta \Delta y)}{\Delta y} + \Delta x \frac{\Delta u(x + \theta' \Delta x, y)}{\Delta x} \end{aligned}$$

[by mean value theorem as stated in (i)]

$$\begin{aligned} &= \Delta x \left[\frac{\Delta u(x, y)}{\Delta x} + \epsilon_1 \right] + \Delta y \left[\frac{\Delta u(x, y)}{\Delta y} + \epsilon_2 \right] \\ &\quad [\because \text{if the function is continuous} \\ &\quad |f(z) - f(z_0)| < \epsilon; \therefore f(z) = f(z_0) + \epsilon_1, \\ &\quad \text{when } |z - z_0| < \delta \text{ where } |\epsilon_1| < \epsilon]. \end{aligned}$$

$$\begin{aligned} &= \Delta x \left[\frac{\partial u}{\partial x} + \epsilon_1 \right] + \Delta y \left[\frac{\partial u}{\partial y} + \epsilon_2 \right] \\ &\quad \left[\because \lim_{\Delta x \rightarrow 0} \frac{\Delta u(x, y)}{\Delta x} = \frac{\partial u}{\partial x} \text{ etc.} \right] \end{aligned}$$

Hence

$$\begin{aligned} \lim_{\Delta z \rightarrow 0} \frac{\Delta w}{\Delta z} &= \frac{dw}{dz} = \lim_{\Delta z \rightarrow 0} \frac{f(z + \Delta z) - f(z)}{\Delta x + i \Delta y} \\ &= \frac{\Delta x \left(\frac{\partial u}{\partial x} + \epsilon_1 \right) + \Delta y \left(\frac{\partial u}{\partial y} + \epsilon_2 \right) + i \Delta x \left(\frac{\partial v}{\partial x} + \epsilon_1' \right) + i \Delta y \left(\frac{\partial v}{\partial y} + \epsilon_2' \right)}{\Delta x + i \Delta y} \\ &= \frac{\Delta x \left(\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \right) + \Delta y \left(\frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) + \Delta x (\epsilon_1 + i \epsilon_1') + \Delta y (\epsilon_2 + i \epsilon_2')}{\Delta x + i \Delta y} \\ &= \frac{(\Delta x + i \Delta y) \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} (\Delta x + i \Delta y) + \eta \Delta x + \eta' \Delta y}{\Delta x + i \Delta y} \end{aligned}$$

[By applying Cauchy-Riemann's equations and putting $\eta = \epsilon_1 + i \epsilon_1'$, $\eta' = \epsilon_2 + i \epsilon_2'$; also $i^2 = -1$].

$$\therefore \frac{dw}{dz} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + \eta \frac{\Delta x}{\Delta x + i \Delta y} + \eta' \frac{\Delta y}{\Delta x + i \Delta y}.$$

$$\begin{aligned} \text{Now } \left| \eta \frac{\Delta x}{\Delta x + i \Delta y} \right| &= \left| \frac{\eta}{\Delta x + i \Delta y} \right| \left| \Delta x \right| \\ &= \frac{|\eta| |\Delta x|}{\sqrt{(\Delta x)^2 + (\Delta y)^2}} \\ &\leq |\eta| \quad \because \Delta x < \sqrt{(\Delta x)^2 + (\Delta y)^2} \\ &\leq \sqrt{(\epsilon_1^2 + \epsilon_1'^2)} \quad \because \eta = \epsilon_1 + i \epsilon_1' \\ &\leq 0 \text{ when } \Delta x \rightarrow 0; \epsilon_1 \text{ and } \epsilon_1' \rightarrow 0. \end{aligned}$$

But by definition the modulus of any quantity is always +ve or zero and it is never negative.

$$\therefore \frac{\eta \Delta x}{\Delta x + i \Delta y} \rightarrow 0, \text{ when } \Delta x \rightarrow 0.$$

$$\text{Similarly } \frac{\eta' \Delta z}{\Delta x + i \Delta y} \rightarrow 0 \text{ when } \Delta y \rightarrow 0.$$

$$\text{Hence } \frac{dw}{dz} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x},$$

i.e., the limit exists and is finite and unique.

Therefore the sufficient conditions for the function $f(z)$ to be regular require the continuity of the four first partial derivatives of u and v .

Polar form of Cauchy-Riemann Equations

The coordinates (x, y) in terms of polar coordinates (r, θ) are given by

$$x = r \cos \theta, y = r \sin \theta$$

So that

$$\frac{\partial r}{\partial x} = \frac{x}{r} = \cos \theta; \frac{\partial r}{\partial y} = \frac{y}{r} = \sin \theta; \frac{\partial \theta}{\partial x} = -\frac{\sin \theta}{r} \text{ and } \frac{\partial \theta}{\partial y} = \frac{\cos \theta}{r}$$

$$\therefore \frac{\partial u}{\partial x} = \frac{\partial u}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial u}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial u}{\partial r} \cos \theta - \frac{\partial u}{\partial \theta} \frac{\sin \theta}{r}.$$

Similarly

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial r} \sin \theta + \frac{\partial u}{\partial \theta} \frac{\cos \theta}{r}; \frac{\partial v}{\partial x} = \frac{\partial v}{\partial r} \cos \theta - \frac{\partial v}{\partial \theta} \frac{\sin \theta}{r};$$

$$\text{and } \frac{\partial v}{\partial y} = \frac{\partial v}{\partial r} \sin \theta + \frac{\partial v}{\partial \theta} \frac{\cos \theta}{r}.$$

Substituting these values in Cauchy-Riemann equations i.e.

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \text{ we get}$$

$$\frac{\partial u}{\partial r} \cos \theta - \frac{\partial u}{\partial \theta} \frac{\sin \theta}{r} = \frac{\partial v}{\partial r} \sin \theta + \frac{\partial v}{\partial \theta} \frac{\cos \theta}{r} \quad \dots (4)$$

$$\text{and } \frac{\partial u}{\partial r} \sin \theta + \frac{\partial u}{\partial \theta} \frac{\cos \theta}{r} = -\frac{\partial v}{\partial r} \cos \theta + \frac{\partial v}{\partial \theta} \frac{\sin \theta}{r} \quad \dots (5)$$

Multiplying (4) by $\cos \theta$, (5) by $\sin \theta$ and adding we get

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}.$$

Again multiplying (4) by $\sin \theta$; (5) by $\cos \theta$ and subtracting, we get

$$\frac{\partial u}{\partial \theta} = -r \frac{\partial v}{\partial r}.$$

Hence Polar form of Cauchy-Riemann conditions are

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}, \frac{\partial u}{\partial \theta} = -r \frac{\partial v}{\partial r} \quad \dots (6)$$

The derivative of the function $w=f(z)$ in the polar form is given as

$$\begin{aligned} \frac{dw}{dz} = \frac{\partial w}{\partial x} &= \frac{\partial w}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial w}{\partial \theta} \frac{\partial \theta}{\partial x} \\ &= \frac{\partial w}{\partial r} \cos \theta - \frac{\sin \theta}{r} \frac{\partial w}{\partial \theta} \\ &= \frac{\partial w}{\partial r} \cos \theta - \frac{\sin \theta}{r} \left(\frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} \right) \quad \because w = u + iv \\ &= \frac{\partial w}{\partial r} \cos \theta - \frac{\sin \theta}{r} \left(-r \frac{\partial v}{\partial r} + ir \frac{\partial u}{\partial r} \right) \text{ using (6)} \\ &= \frac{\partial w}{\partial r} \cos \theta - \sin \theta \left(-\frac{\partial v}{\partial r} + i \frac{\partial u}{\partial r} \right) \\ &= (\cos \theta - i \sin \theta) \frac{\partial w}{\partial r} \\ &= -\frac{i}{r} (\cos \theta - i \sin \theta) \frac{\partial w}{\partial \theta}. \end{aligned}$$

Condition for a Function when it Ceases to be Analytic

If $w=F(\zeta)$ and $\zeta=f(z)$, then w is said to be function of a function of z and we have $\frac{dw}{dz} = \frac{dw}{d\zeta} \frac{d\zeta}{dz}$, $F(\zeta)$ and $f(z)$ both being analytic.

Also if $w=f(z)$ be an analytic function of z such that corresponding to each point w_0 there exists a point w_0 and $z=F(w)$ is such that to each value w_0 of w there corresponds a value z_0 of z , then the function $z=F(w)$ is said to be the *Inverse function* of $w=f(z)$. Clearly, if $f'(z_0) \neq 0$, then w_0 is a regular point of $z=F(w)$ i.e. z is analytic in the neighbourhood of w_0 .

On account of functions being inverse we have $F'(w_0) = \frac{1}{f'(z_0)}$ and hence the function $z=F(w)$ ceases to be analytic where $f'(z)=0$ i.e. $\frac{dw}{dz}=0$ also that $w=f(z)$ where $z=F(w)$ ceases to be regular when

$$\frac{dw}{dz}=0.$$

Evidently when $z=x+iy$, $w=f(z)$, we have $\frac{dw}{dz} = \frac{\partial w}{\partial x}$.

So that if $w=u+iv$, $z=f(w)$, then $\frac{dz}{dw} = \frac{\partial z}{\partial u}$.

As an illustration if $w=e^{-v}(\cos u + i \sin u)$, then $\frac{dz}{dw} = \frac{\partial z}{\partial u}$ gives

$$\frac{dz}{dw} = e^{-v}(-\sin u + i \cos u) = -ie^{-v}(\cos u + i \sin u) = -iz$$

$\therefore w$ ceases to be analytic when $\frac{dz}{dw}=0$ i.e. $z=0$.

To prove that if a function is regular, it is independent of $\bar{z}$ and is function of z .

If $z = x + iy,$
 $\bar{z} = x - iy,$

Adding and subtracting, we get

$$x = \frac{1}{2}(z + \bar{z}) \text{ and } y = \frac{1}{2i}(z - \bar{z}).$$

$$\therefore \frac{\partial x}{\partial \bar{z}} = \frac{1}{2} \text{ and } \frac{\partial y}{\partial \bar{z}} = -\frac{1}{2i} \text{ etc.}$$

Now $w = f(z) = u + iv.$

$$\begin{aligned} \therefore \frac{dw}{d\bar{z}} &= \frac{du}{d\bar{z}} + i \frac{dv}{d\bar{z}} \\ &= \left[\frac{\partial u}{\partial x} \cdot \frac{\partial x}{\partial \bar{z}} + \frac{\partial u}{\partial y} \cdot \frac{\partial y}{\partial \bar{z}} \right] + i \left[\frac{\partial v}{\partial x} \cdot \frac{\partial x}{\partial \bar{z}} + \frac{\partial v}{\partial y} \cdot \frac{\partial y}{\partial \bar{z}} \right] \\ &= \frac{\partial u}{\partial x} \cdot \frac{1}{2} - \frac{1}{2i} \frac{\partial u}{\partial y} + i \left[\frac{\partial v}{\partial x} \cdot \frac{1}{2} - \frac{1}{2i} \frac{\partial v}{\partial y} \right] \\ &= 0 \text{ by Cauchy-Riemann's equations.} \end{aligned}$$

Thus if the function is regular, it is independent of $\bar{z}$, as its differential is zero.

Laplace's Equations

Cauchy-Riemann's equations are

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x};$$

$$\therefore \frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 v}{\partial x \partial y} \text{ and } \frac{\partial^2 u}{\partial y^2} = -\frac{\partial^2 v}{\partial x \partial y} \quad \therefore \frac{\partial^2 v}{\partial x \partial y} = \frac{\partial^2 v}{\partial y \partial x}$$

Adding, $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0,$

i.e., $\nabla^2 u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \dots (II)$

Similarly, $\nabla^2 v = \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = 0 \quad \dots (III)$

These are known as Laplace's equations, in which both u and v satisfy Laplace's equation in two dimensions.

Harmonic Functions. A function of x, y is said to be a harmonic function if it possesses continuous partial derivatives of the first and second orders and satisfies Laplace's equation.

Two harmonic functions u and v as satisfying (ii) and (iii) are known as *Conjugate harmonic functions* or simply *conjugate functions*.

Determination of Conjugate Functions

If $f(z) = u + iv$ is an analytic function such that u and v are conjugate functions then being given one of them say u , we have to determine v .

We have

$$\begin{aligned} dv &= \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy \quad \because v \text{ is a function of } x, y. \\ &= -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy \text{ by Cauchy-Riemann equations.} \end{aligned}$$

The R.H.S. of this equation being of the form $Mdx + Ndy$ will be exact if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} \text{ where } M = -\frac{\partial u}{\partial y} \text{ and } N = \frac{\partial u}{\partial x}$$

$$\text{i.e., } \frac{\partial}{\partial y} \left(-\frac{\partial u}{\partial y} \right) = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial x} \right)$$

$$\text{or } -\frac{\partial^2 u}{\partial y^2} = \frac{\partial^2 u}{\partial x^2}$$

$$\text{i.e., } \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0.$$

As u satisfies Laplace's equation, it is harmonic and hence its conjugate v can be found out by integrating the equation.

$$dv = -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy.$$

As an illustration if $u = y^3 - 3x^2y$, then $\frac{\partial u}{\partial x} = -6xy, \frac{\partial u}{\partial y} = 3y^2 - 3x^2$

$$\frac{\partial^2 u}{\partial x^2} = -6y, \frac{\partial^2 u}{\partial y^2} = 6y \text{ so that } u \text{ satisfies Laplace's equation and}$$

hence is harmonic.

$$\begin{aligned} \text{Now } v &= \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy \\ &= -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy \text{ by Cauchy-Riemann equations} \\ &= -(3y^2 - 3x^2) dx - 6xy dy \\ &= -(3y^2 dx + 6xy dy) + 3x^2 dx \end{aligned}$$

Integrating $v = -3xy^2 + x^3 + c$ which is harmonic conjugate to u .

$$\begin{aligned} \text{Corresponding analytic function } f(z) &= u + iv \\ &= y^3 - 3x^2y + i(-3xy^2 + x^3 + c) \\ &= i(x + iy)^3 + ic \\ &= iz^3 + c' \end{aligned}$$

Using an alternative method, $\frac{\partial u}{\partial x} = -6xy = \frac{\partial v}{\partial y}$ by Cauchy-Riemann equations.

Integrating $\frac{\partial v}{\partial x} = -6xy$ we get $v = -3xy^2 + \phi(x)$, $\phi(x)$ being arbitrary.

But $\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$ gives

$$-3y^2 + \phi'(x) = -3y^2 + 3x^2$$

i.e. $\phi'(x) = 3x^2$ and so $\phi(x) = x^3 + c$

As such $v = -3xy^2 + x^3 + c$

Construction of a function $f(z)$ when one conjugate is given (due to Milne-Thomson)

If $z = x + iy$, $\bar{z} = x - iy$ and so $x = \frac{z + \bar{z}}{2}$, $y = \frac{z - \bar{z}}{2i}$

$\therefore f(z) = u(x, y) + iv(x, y)$

$$= u\left(\frac{z + \bar{z}}{2}, \frac{z - \bar{z}}{2i}\right) + iv\left(\frac{z + \bar{z}}{2}, \frac{z - \bar{z}}{2i}\right)$$

Treating it as a formal identity in two independent variables z and $\bar{z}$ and putting $z = \bar{z}$, we get $x = z$, $y = 0$ so that,

$$f(z) = u(z, 0) + iv(z, 0)$$

Taking $f(z) = u + iv$ to be analytic, we have

$$\begin{aligned} f'(z) &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \\ &= \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} \text{ by Cauchy-Riemann equations.} \end{aligned}$$

Writing $\phi(x, y) \equiv \frac{\partial u}{\partial x}$, $\psi(x, y) \equiv \frac{\partial u}{\partial y}$, we have,

$$\begin{aligned} f'(z) &= \phi(x, y) - i\psi(x, y) \\ &= \phi(z, 0) - i\psi(z, 0) \end{aligned}$$

Integrating, $f(z) = \int \{\phi(z, 0) - i\psi(z, 0)\} dz + c$, c being arbitrary constant.

Similarly if $v(x, y)$ is given then we can find

$$f(z) = \int \{\Phi(z, 0) + i\Psi(z, 0)\} dz + C$$

where $\Phi(x, y) = \frac{\partial v}{\partial y}$ and $\Psi(x, y) = \frac{\partial v}{\partial x}$.

As an illustration if $u = e^x(x \cos y - y \sin y)$, then

$$\frac{\partial u}{\partial x} = e^x(x \cos y - y \sin y + \cos y) = \phi(x, y) \text{ (say)}$$

and $\frac{\partial u}{\partial y} = e^x(-x \sin y - y \cos y - \sin y) = \psi(x, y) \text{ (say)}$

So that $\phi(z, 0) = e^z(z + 1)$ and $\psi(z, 0) = 0$

$$\begin{aligned} \therefore f'(z) &= \phi(z, 0) - i\psi(z, 0) \\ &= e^z(z + 1) \end{aligned}$$

Integrating $f(z) = ze^z + c$.

Problem 6. Prove that the function $u + iv = f(z)$ where

$$f(z) = \frac{x^3(1+i) - y^3(1-i)}{x^2 + y^2} \quad (z \neq 0), \quad f(0) = 0$$

is continuous and that the Cauchy-Riemann equations are satisfied at the origin, yet $f'(0)$ does not exist.

Here $u + iv = f(z)$

$$= \frac{x^3(1+i) - y^3(1-i)}{x^2 + y^2}$$

Equating real and imaginary parts on either side, we get

$$u = \frac{x^3 - y^3}{x^2 + y^2} \text{ and } v = \frac{x^3 + y^3}{x^2 + y^2} \text{ when } z \neq 0.$$

Obviously both u and v are rational and finite for all values of $z \neq 0$. Thus u and v are continuous at all those points for which $z \neq 0$. Hence $f(z)$ is continuous when $z \neq 0$.

Given that $f(0) = 0$, therefore at the origin $u = 0$, $v = 0$. Hence u and v both are continuous at the origin. As such $f(z)$ is continuous at the origin.

Conversely $f(z)$ is continuous everywhere.

Now,

$$\left(\frac{\partial u}{\partial x}\right)_{\substack{\text{at } x=0 \\ y=0}} = \frac{\partial u(x, 0)}{\partial x} \text{ at } x=0$$

$$= \lim_{h \rightarrow 0} \frac{u(h, 0) - u(0, 0)}{h} \because u(h, 0) = \frac{h^3 - 0}{h^2 + 0} = h \text{ etc.}$$

$$= \lim_{h \rightarrow 0} \frac{h}{h} = 1.$$

$$\left(\frac{\partial u}{\partial y}\right)_{\substack{\text{at } x=0 \\ y=0}} = \lim_{y=0} \frac{\partial u(0, y)}{\partial y} = \lim_{k \rightarrow 0} \frac{u(0, k) - u(0, 0)}{k}$$

$$= \lim_{k \rightarrow 0} \frac{-k}{k} = -1 \because u(0, k) = \frac{0 - k^3}{0 + k^2} = -k \text{ etc.}$$

$$\left(\frac{\partial v}{\partial x}\right)_{\substack{\text{at } x=0 \\ y=0}} = \lim_{y=0} \frac{\partial v(x, 0)}{\partial x}$$

$$= \lim_{k \rightarrow 0} \frac{v(h, 0) - v(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

$$\text{and } \left(\frac{\partial v}{\partial y}\right)_{\substack{\text{at } x=0 \\ y=0}} = \lim_{y \rightarrow 0} \frac{\partial v(0, y)}{\partial y}$$

$$= \lim_{k \rightarrow 0} \frac{v(0, k) - v(0, 0)}{k} = \lim_{k \rightarrow 0} \frac{k}{k} = 1.$$

Thus we have found that at the origin

$$\frac{\partial u}{\partial x} = 1 = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -1 = -\frac{\partial v}{\partial x},$$

which clearly satisfy the Cauchy-Riemann equations at $z=0$. Again differential coefficient of $f(z)$ at $z=0$, i.e.

$$\begin{aligned} f'(0) &= \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} \\ &= \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \frac{x^3(1+i) - y^3(1-i)}{(x^2 + y^2)(x + iy)} \quad \because z = x + iy \\ &= \lim_{x \rightarrow 0} \frac{x^3(1+i) - m^3(1-i)x^3}{(x^2 + m^2x^2)(x + imx)} \quad \text{by putting } y = mx \\ &= \frac{(1+i) - m^3(1-i)}{(1+m^2)(1+im)} \end{aligned}$$

which is not unique, as it is different for different values of m . Therefore $f(z)$ is not continuous at $z=0$.

Hence $f'(z)$ does not exist at the origin, i.e., $z=0$.

Problem 7. Show that the function $f(z) = \sqrt{|xy|}$ is not regular at the origin although the Cauchy-Riemann equations are satisfied at that point.

Let the function be

$$f(z) = \sqrt{|xy|} = u(x, y) + iv(x, y).$$

Equating real and imaginary parts, we have

$$u(x, y) = \sqrt{|xy|} \text{ and } v(x, y) = 0.$$

$$\text{Thus, } \left(\frac{\partial u}{\partial x} \right)_{\substack{\text{at } x=0 \\ y=0}} = \frac{\partial u(x, 0)}{\partial x} \text{ at } x=0$$

$$= \lim_{h \rightarrow 0} \frac{u(h, 0) - u(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{0 - 0}{h} = 0.$$

Similarly at $x=0, y=0$,

$$\frac{\partial v}{\partial x} = 0, \frac{\partial u}{\partial y} = 0, \frac{\partial v}{\partial y} = 0.$$

These values clearly satisfy the Cauchy-Riemann equations.

$$\text{Again } f'(0) = \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z} = \lim_{\substack{x \rightarrow 0 \\ y=0}} \frac{\sqrt{|xy|} - 0}{x + iy}$$

$$\begin{aligned} &= \lim_{x \rightarrow 0} \frac{x\sqrt{|m|}}{x(1+im)} \text{ by putting } y = mx \\ &= \frac{\sqrt{|m|}}{1+im}, \end{aligned}$$

which is not unique as its values are different for different values of m . So $f(z)$ is not continuous at $z=0$. Hence it is not regular there.

Problem 8. Prove that the function:

$$u = x^3 - 3xy^2 + 3x^2 - 3y^2 + 1$$

satisfies Laplace's equation and determine the corresponding regular function $u + iv$

$$\text{Here, } \frac{\partial u}{\partial x} = 3x^2 - 3y^2 + 6x, \quad \dots (1)$$

$$\frac{\partial^2 u}{\partial x^2} = 6x + 6, \quad \dots (2)$$

$$\frac{\partial u}{\partial y} = -6xy - 6y, \quad \dots (3)$$

$$\frac{\partial^2 u}{\partial y^2} = -6x - 6. \quad \dots (4)$$

By the additions of (2) and (4) it follows that $\nabla^2 u = 0$, which clearly satisfies the Laplace's equation.

Hence u is a Harmonic function.

Now Cauchy-Riemann equations are

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

$$\begin{aligned} \therefore dv &= \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy \\ &= -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy \text{ (by Cauchy-Riemann equations)} \\ &= (6xy + 6y) dx + (3x^2 - 3y^2 + 6x) dy \\ &= (6xy dx + 3x^2 dy) - 3y^2 dy + (6y dx + 6x dx). \end{aligned}$$

Integrating,

$$v = 3x^2y - y^3 + 6xy + c, \text{ where } c \text{ is an absolute or complex constant.}$$

Thus

$$\begin{aligned} u + iv &= x^3 - 3xy^2 + 3x^2 - 3y^2 + 1 + i [3x^2y - y^3 + 6xy + c] \\ &= (x + iy)^3 + 3(x + iy)^2 + 1 + ic \\ &= z^3 + 3z^2 + 1 + ic \\ &= z^3 + 3z^2 + c'. \end{aligned}$$

Problem 9. Prove that the curves $u = \text{constant}$,
 $v = \text{constant}$

intersect at right angles.

We know that the curves intersect at right angles if the tangents to them at their point of intersection are at right angles.

Differentiating partially the given equations of the curves, we get

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy = 0,$$

$$dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy = 0.$$

Therefore tangents of the angles formed by the tangents drawn to the curves at their point of intersection with the real axes are respectively

$$\left(\frac{dy}{dx}\right)_1 = -\frac{\partial u/\partial x}{\partial v/\partial y} \text{ and } \left(\frac{dy}{dx}\right)_2 = -\frac{\partial v/\partial x}{\partial u/\partial y}.$$

$$\text{Now } \left(\frac{dy}{dx}\right)_1 \left(\frac{dy}{dx}\right)_2 = \frac{\frac{\partial u}{\partial y} \cdot \frac{\partial v}{\partial x}}{\frac{\partial u}{\partial y} \cdot \frac{\partial v}{\partial y}} = -1, \text{ by Cauchy-Riemann equations}$$

which shows that the two tangents are at right angles if the function is regular. It follows that the two curves intersect at right angles.

Problem 10. Show that $\nabla^2 |u|^p = p(p-1) |u|^{p-2} |f'(z)|^2$.

Here if u be positive $|u| = u$ and if u be negative $|-u| = u$.

Taking first u to be positive,

$$\frac{\partial}{\partial x} u^p = p \cdot u^{p-1} \frac{\partial u}{\partial x}.$$

$$\frac{\partial^2}{\partial x^2} u^p = p(p-1) u^{p-2} \left(\frac{\partial u}{\partial y}\right)^2 + p \cdot u^{p-1} \frac{\partial^2 u}{\partial y^2}.$$

$$\text{Similarly, } \frac{\partial^2}{\partial y^2} u^p = p(p-1) u^{p-2} \left(\frac{\partial u}{\partial y}\right)^2 + p \cdot u^{p-1} \frac{\partial^2 u}{\partial x^2}.$$

Adding the last two results,

$$\begin{aligned} \nabla^2 u^p &= pu^{p-1} \nabla^2 u + p(p-1) u^{p-2} \left[\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2 \right], \\ \text{where } \nabla^2 &\equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \\ &= p(p-1) u^{p-2} \left[\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2 \right]. \end{aligned} \quad \dots (1)$$

$$\therefore \nabla^2 u = 0 \text{ (Laplace's equation),}$$

Now if $w = f(z) = u + iv$, then

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}.$$

$$\therefore |f'(z)|^2 = \left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial v}{\partial x}\right)^2 \quad \dots (2)$$

Thus from (1) and (2) it follows that

$$\nabla^2 u^p = p(p-1) u^{p-2} |f'(z)|^2.$$

Again if u be negative say u_1 ,

$$\therefore |-u| = u_1.$$

As before, we have

$$\nabla^2 u_1^p = p(p-1) u_1^{p-2} |f'(z)|^2$$

or

$$\nabla^2 |u_1|^p = p(p-1) |u_1|^{p-2} |f'(z)|^2.$$

8.7. Elementary Functions and Mapping by Them

Exponential functions. If $z = x + iy$ and y is used as radian measure of the angle to define $\cos y$, $\sin y$ etc, then the exponential function in terms of real valued functions is defined by

$$e^z = e^{x+iy} = e^x e^{iy} = e^x (\cos y + i \sin y) \quad \dots (1)$$

$$\text{In case } z \text{ is purely real i.e. } y=0, \text{ we have } e^z = e^x \quad \dots (2)$$

$$\text{and if } z \text{ is purely imaginary i.e. } x=0, \text{ we have } e^{iy} = \cos y + i \sin y \quad \dots (3)$$

As such Maclaurin series representation of e^t on replacing t by iy , given

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(iy)^n}{n!} &= \sum_{n=0}^{\infty} \frac{i^{2n} y^{2n}}{2n!} + \sum_{n=0}^{\infty} \frac{i^{2n+1} y^{2n+1}}{(2n+1)!} \text{ where } |0|=1 \\ &= \sum_{n=0}^{\infty} (-1)^n \frac{y^{2n}}{(2n)!} + i \sum_{n=0}^{\infty} (-1)^n \frac{y^{2n+1}}{(2n+1)!} \end{aligned} \quad \dots (4)$$

which are Maclaurin series for $\cos y$ and $\sin y$ respectively.

The Exponential function given by (1) is an *entire function* since

$$\frac{d}{dz} e^z = e^z \quad \dots (5)$$

$$\text{Similarly } \frac{d}{dz} e^w = e^w \frac{dw}{dz} \quad \dots (6)$$

w being an analytic function of z .

$$\text{Polar form of (1) is } e^z = r (\cos \theta + i \sin \theta) = r e^{i\theta} \quad \dots (7)$$

where $r = e^x$, $\theta = y$.

$$\therefore |e^z| = r = e^x \text{ and } \arg e^z = \theta = y \quad \dots (8)$$

$$\text{Also } |e^z| > 0 \text{ i.e. } e^z \neq 0 \text{ for every value of } z \quad \dots (9)$$

So the range of the exponential function is the entire complex plane excluding the origin where $r=0$.

Now $e^z = -1 = \cos(\pi \pm 2m\pi) + i \sin(\pi \pm 2m\pi)$ gives

$$x=0 \text{ and } y=\pi \pm 2m\pi, m=0, 1, 2, \dots$$

If $e^{z_1} = r_1 (\cos \theta_1 + i \sin \theta_1)$ so that $r_1 = e^{x_1}$, $\theta_1 = y_1$

and $e^{z_2} = r_2 (\cos \theta_2 + i \sin \theta_2)$ so that $r_2 = e^{x_2}$, $\theta_2 = y_2$

then, $e^{z_1} \cdot e^{z_2} = r_1 r_2 \{ \cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) \} = e^{x_1} e^{x_2} [\cos(y_1 + y_2) + i \sin(y_1 + y_2)]$

$$\begin{aligned} &= e^{x_1+x_2} \cdot e^{i(y_1+y_2)} = e^{x_1+x_2+i(y_1+y_2)} \\ &= e^{(x_1+iy_1)+(x_2+iy_2)} = e^{z_1+z_2} \end{aligned} \quad \dots (10)$$

$$\text{Similarly } e^{z_1}/e^{z_2} = e^{z_1-z_2} \quad \dots (11)$$

$$\frac{1}{e^z} = e^{-z} \quad \dots (12)$$

$$(e^z)^n = e^{nz} \quad \dots (13)$$

n being a positive integer.

$$(e^z)^{m/n} = e^{m/n}(z+2\pi pi) \quad \dots (14)$$

$p=0, 1, 2, \dots, x-1$ and m, n are integers ($+ve$).

$$\text{Also } e^{z+2\pi i} = e^z \quad e^{2\pi i} = e^z \text{ as } e^{2\pi i} = 1 \quad \dots (15)$$

(15) follows that the exponential function is *periodic*.

$$\text{Again } e^{\bar{z}} = \overline{(e^z)} \quad \dots (16)$$

In polar form $z = re^{i\theta}$, $\bar{z} = re^{-i\theta}$ so that

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}, \quad \frac{z_1}{z_2} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)} \quad \dots (17)$$

Trigonometric Functions

$e^{iy} = \cos y + i \sin y$ and $e^{-iy} = \cos y - i \sin y$ yield

$$\cos y = \frac{e^{iy} + e^{-iy}}{2}, \quad \sin y = \frac{e^{iy} - e^{-iy}}{2i} \quad \dots (18)$$

$\sin z$ and $\cos z$ are *entire functions* as

$$\frac{d}{dz} \sin z = \cos z, \quad \frac{d}{dz} \cos z = -\sin z \text{ etc.} \quad \dots (19)$$

Now,

$$\begin{aligned} \cos z &= \cos(x+iy) = \frac{1}{2} [e^{i(x+iy)} + e^{-i(x+iy)}] \\ &= \frac{1}{2} [e^{ix-y} + e^{-ix+y}] \\ &= \frac{1}{2} e^{-y} (\cos x + i \sin x) + \frac{1}{2} e^y (\cos x - i \sin x) \\ &= \frac{e^y + e^{-y}}{2} \cos x - i \frac{e^y - e^{-y}}{2} \sin x \quad \dots (20) \end{aligned}$$

Introducing the *hyperbolic functions* with the properties

$$\sinh z = \frac{e^z - e^{-z}}{2}, \quad \cosh z = \frac{e^z + e^{-z}}{2} \quad \dots (21)$$

$$\frac{d}{dz} \sinh z = \cosh z, \quad \frac{d}{dz} \cosh z = \sinh z \text{ etc.} \quad \dots (22)$$

$$\cosh^2 z - \sinh^2 z = 1 \quad \dots (23)$$

$$\cosh iz = \cosh z \text{ and } \sin iz = i \sinh z \quad \dots (24)$$

the relation (20) becomes

$$\cos z = \cos(x+iy) = \cos x \cosh y - i \sin x \sinh y \quad \dots (25)$$

$$\text{Similarly } \sin z = \sin(x+iy) = \sin x \cosh y + i \cos x \sinh y \quad \dots (26)$$

$$\text{Also } \sin \bar{z} = \overline{\sin z} \text{ and } \cos \bar{z} = \overline{\cos z} \quad \dots (27)$$

$$\cos(z+\pi) = -\cos z, \quad \sin(z+\pi) = -\sin z, \text{ etc.} \quad \dots (28)$$

It is easy to show that

$$|\sin z|^2 = \sin^2 x + \sinh^2 y \quad \dots (29)$$

$$|\cos z|^2 = \cos^2 x + \sinh^2 y \quad \dots (30)$$

A value of z for which $f(z)=0$ is known as a *zero* of the function f .

The real zeros of $\sin z$ and $\cos z$ are their only zeros, since

$\sin z=0$ from (26) gives

$$\sin x \cosh y = 0, \quad \cos x \sinh y = 0$$

x and y being real, $\cosh y \geq 1$ and $\sin x=0$ only when $x=0, \pm\pi, \pm 2\pi, \dots$ and for these values of x , $\cos x \neq 0$ and thus $\sinh y=0$ i.e. $y=0$.

Also

$$\begin{aligned} \sin z=0 &\Rightarrow z=0 \text{ or } \pm n\pi, \quad x=1, 2, 3, \dots \\ \cos z=0 &\Rightarrow z=\pm \frac{(2n-1)\pi}{2}, \quad x=1, 2, 3, \dots \end{aligned} \quad \dots (31)$$

we may also show that

$$\cosh z = \cosh(x+iy) = \cosh x \cos y + i \sinh x \sin y \quad \dots (32)$$

$$\sinh z = \sinh x \cos y + i \cosh x \sin y \quad \dots (33)$$

$$\text{So that } |\sinh z|^2 = \sinh^2 x + \sin^2 y \quad \dots (34)$$

$$|\cosh z|^2 = \sinh^2 x + \cos^2 y \quad \dots (35)$$

$\sinh z$ and $\cosh z$ are periodic with *period* $2\pi i$.

Logarithmic Functions and Branch Points

$z = re^{i\theta}$, θ being measured in radian, gives

$$\begin{aligned} \log z &= \log re^{i\theta} = \log r + i\theta \\ &= \log |z| + i \arg \theta \end{aligned} \quad \left. \begin{array}{l} r > 0 \text{ and } -\pi < \theta < \pi \end{array} \right\} \quad \dots (36)$$

If $-\pi < \theta \leq \pi$, then $z = re^{i(\theta \pm 2n\pi)}$, $n=0, 1, 2, \dots$

$$\text{So that } \log z = \log r + i(\theta + i2n\pi), \quad n=0, 1, 2, \quad \dots (37)$$

We write $\log z$ for *principal value* of $\log z$ and $\text{Log } z$ for its *general value*.

In (36) if we put $\log r = u$, $\theta = v$ so that

$$\frac{\partial u}{\partial r} = \frac{1}{r}, \quad \frac{\partial u}{\partial \theta} = 0, \quad \frac{\partial v}{\partial r} = 0, \quad \frac{\partial v}{\partial \theta} = 1, \text{ are all continuous functions}$$

of z .

$$\begin{aligned} \text{Also, } \frac{d}{dz} \log z &= e^{-i\theta} \left(\frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \right) \\ &= \frac{1}{re^{i\theta}} = \frac{1}{z} \quad \dots (38) \end{aligned}$$

where $z \neq 0$ and $-\pi < \arg z < \pi$.

A *branch* F of a many-valued function f is any single-valued function which is analytic in some domain at each point of which the value $F(z)$ is one of the values $f(z)$. The equation (36) gives the *principal branch* $\log z$. Each point of the negative real axis $\theta=\pi$ along with the origin is a *singular point* of the principal branch $\log z$. The n (say) $\theta=\pi$ is said to be a *branch cut* for the principal branch and the singular point $z=0$ common to all branch cuts for the many-valued function $\log z$ is known as a *branch point*.

Now if $w = \log z$, then $e^w = e^{\log z} = e^{(\log r + i\theta)}$

$$= e^{\log r} \cdot e^{i\theta} = re^{i\theta} = z$$

$$\text{i.e. } e^{\log z} = z, \quad z \neq 0 \quad \dots (39)$$

and if $e^z = w$, $\log w = \log e^{x+iy} = \log e^x + \log e^{iy}$.

$$\begin{aligned} &= x + i(y \pm 2m\pi), m=0, 1, 2, 3, \dots \\ &= x + iy \pm 2p\pi i \\ &= z \pm 2p\pi i \end{aligned}$$

So $\log w = z$ when $e^z = w$

and $\log e^z = z$ for appropriate choice of logarithm

Again if $z_1 = r_1 e^{i\theta_1}$, $z_2 = r_2 e^{i\theta_2}$, $r_1 > 0$, $r_2 > 0$, then

$$\begin{aligned} \log z_1 + \log z_2 &= \log r_1 e^{i\theta_1} + \log r_2 e^{i\theta_2} \\ &= \log r_1 + \log r_2 + i(\theta_1 + \theta_2) \\ &= \log r_1 r_2 + i(\theta_1 + \theta_2) \\ &= \log z_1 z_2 \end{aligned}$$

$$\text{Similarly } \log z_1 - \log z_2 = \log \frac{z_1}{z_2}$$

It is easy to verify that

$$\log z^m = m \log z$$

$$\log z^{1/n} = \frac{1}{n} \log z$$

$$z^m/n = e^{m/n \log z}$$

In case of complex exponents, we define

$$z^c = e^{c \log z}, z, c \text{ being complex and } z \neq 0$$

$$\frac{d}{dz} z^c = e^{c \log z} \cdot \frac{c}{z} = c \cdot \frac{e^{c \log z}}{z} = c e^{(c-1) \log z} = c z^{c-1}$$

$$\text{Also } c^z = e^{z \log c}, c \neq 0,$$

$$\frac{d}{dz} c^z = c^z \log c, c \neq 0$$

Inverse Trigonometric Functions

Defining the inverse of sine function as $w = \sin^{-1} z$, we have

$$z = \sin w = \frac{e^{iw} - e^{-iw}}{2i} \text{ i.e. } e^{2iw} - 2ize^{iw} - 1 = 0.$$

Being quadratic in e^{iw} , this gives, $e^{iw} = iz + \sqrt{1-z^2}$

$$\text{So that } w = \sin^{-1} z = -i \log \{iz + \sqrt{1-z^2}\}$$

$$\text{Similarly } \cos^{-1} z = -i \log \{z + \sqrt{z^2-1}\}$$

$$\tan^{-1} z = \frac{i}{2} \log \frac{1-iz}{1+iz} = \frac{i}{2} \log \frac{1+z}{1-z}$$

$$\text{So that } \frac{d}{dz} \sin^{-1} z = \frac{1}{\sqrt{1-z^2}}, \quad \frac{d}{dz} \tan^{-1} z = \frac{1}{1+z^2}$$

Also

$$\left. \begin{aligned} \sinh^{-1} z &= \log \{z + \sqrt{z^2+1}\} \\ \cosh^{-1} z &= \log \{z + \sqrt{z^2-1}\} \\ \tan^{-1} z &= \frac{1}{2} \log \frac{1+z}{1-z} \end{aligned} \right\}$$

... (55)

Mapping. If $w=f(z)$ and corresponding to each point (x, y) in z -plane in a domain of function f , there is a point (u, v) in w -plane where $z=x+iy$ and $w=u+iv$, then this correspondence between the points of two planes is said to be a *mapping* or a *transformation* of points in the z -plane into points of the w -plane by the function f . Corresponding points or set of points are known as *images* of each other. The use of graphic terms as *translation*, *rotation* or *reflection* is rather convenient in mapping, e.g.

The mapping $w=z+c$, c being a complex constant gives the translation of every point z through the vector c i.e. if $z=x+iy$, $w=u+iv$, $c=c_1+ic_2$, then the image of any point (x, y) in z -plane is the point $(x+c_1, y+c_2)$

The mapping $w=Bz$ where $B=be^{i\beta}$ and $z=re^{i\theta}$

$$\text{i.e. } w=br e^{i(\theta+\beta)}$$

maps the point (r, θ) in z -plane into a point $(br, \theta+\beta)$ into w -plane i.e. the mapping consists of a rotation of the radius vector of z about the origin through an angle $\beta=\arg B$ and an extension or contraction of radius vector r by $b=|B|$.

As an illustration the function $w=z^2$ maps the entire first quadrant of the z -plane, $0 \leq \theta \leq \pi/2$, $r > 0$, into the entire upper half of the w -plane.

The transformation

$$T: w = \frac{az+b}{cz+d} \quad \dots (56)$$

a, b, c, d being complex constants is termed as the *linear fractional transformation* or *bilinear transformation* or *Mobius transformation*.

$$\text{Here } T^{-1} \text{ i.e. inverse of } T \text{ is } z = \frac{-dw+b}{cw-a} \quad \dots (57)$$

Any set of elements which satisfies all the following conditions is called a *group*:

(i) There is a rule of combination such that product TT' for each distinct pair T, T' of elements is an element of the set.

(ii) The product is associative i.e. $T(T'T'') = (TT')T''$

(iii) The set contains an identity T_0 such that $TT_0=T_0T=T$ for each element T .

(iv) Each element T has an inverse T^{-1} s.t. $TT^{-1}=T^{-1}T=T_0$.

It may be shown that the set of all linear fractional transformation is a group.

If besides the above four properties a group also satisfies the commutative property then it is called an *Commutative group* or an *Abelian group*.

Problem 11. Show that the set of complex numbers form an *Abelian group* under addition.

Take three complex numbers, $z_1=(x_1, y_1)$, $z_2=(x_2, y_2)$, $z_3=(x_3, y_3)$

belonging to the set C of complex numbers. Then the addition is commutative since

$$z_1 + z_2 = (x_1 + x_2, y_1 + y_2) = (x_2 + x_1, y_2 + y_1) \\ = z_2 + z_1 \quad \forall z_1, z_2 \in C$$

Also $z_1, z_2 \in C \Rightarrow z_1 + z_2 \in C$

The addition is associative since

$$z_1 + (z_2 + z_3) = (x_1, y_1) + (x_2 + x_3, y_2 + y_3) \\ = (x_1 + x_2 + x_3, y_1 + y_2 + y_3) \\ = (x_1 + x_2, y_1 + y_2) + (x_3, y_3) \\ = (z_1 + z_2) + z_3.$$

There exists an additive identity $o = (0, 0)$ such that

$$z + o = (x, y) + (0, 0) = (x, y) = z$$

There exists an additive inverse $(-z)$ for $\forall z$ such that

$$z + (-z) = (x, y) + (-x, -y) = (0, 0) \text{ the identity element.}$$

Hence the set of complex numbers form an abelian group.

Problem 12. Show that the set of complex numbers form an abelian group under multiplication.

It is easy to show that

$$(x_1, y_1), (x_2, y_2) \in C \Rightarrow (x_1, y_1) (x_2, y_2) \text{ i.e. } (x_1 x_2 - y_1 y_2, x_1 y_2 + x_2 y_1) \in C \\ (x_1, y_1) \{ (x_2, y_2) (x_3, y_3) \} = \{ (x_1, y_1) (x_2, y_2) \} (x_3, y_3), (x_1, y_1), (x_2, y_2), (x_3, y_3) \in C$$

$\exists$ multiplicative identity $(1, 0)$ s.t. $(x, y) (1, 0) = (x, y) \quad \forall (x, y) \in C$

$\exists$ multiplicative inverse s.t. $(x, y) \left(\frac{x}{x^2 + y^2}, \frac{-y}{x^2 + y^2} \right) = (1, 0)$

The commutative law holds i.e.

$$(x_1, y_1) + (x_2, y_2) = (x_2, y_2) + (x_1, y_1).$$

Hence the given set is an abelian group.

Complex Differential Operators

If $z = x + iy$ so that $\bar{z} = x - iy$ and F be a continuous differential function, then

$$\frac{\partial F}{\partial x} = \frac{\partial F}{\partial z} \cdot \frac{\partial z}{\partial x} + \frac{\partial F}{\partial \bar{z}} \frac{\partial \bar{z}}{\partial x} = \frac{\partial F}{\partial z} + \frac{\partial F}{\partial \bar{z}} \text{ giving} \\ \frac{\partial}{\partial x} \equiv \frac{\partial}{\partial z} + \frac{\partial}{\partial \bar{z}} \quad \dots (58)$$

$$\text{and } \frac{\partial F}{\partial y} = \frac{\partial F}{\partial z} \frac{\partial z}{\partial y} + \frac{\partial F}{\partial \bar{z}} \frac{\partial \bar{z}}{\partial y} = i \frac{\partial F}{\partial z} - i \frac{\partial F}{\partial \bar{z}} \text{ giving}$$

$$\frac{\partial}{\partial y} \equiv i \left(\frac{\partial}{\partial z} - \frac{\partial}{\partial \bar{z}} \right) \quad \dots (59)$$

$$\text{We then define } \nabla \text{ (Del operator)} \equiv \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} = \frac{\partial}{\partial z} + \frac{\partial}{\partial \bar{z}} \\ + i \left(\frac{\partial}{\partial z} - \frac{\partial}{\partial \bar{z}} \right) = 2 \frac{\partial}{\partial \bar{z}} \quad \dots (60)$$

$$\text{and } \nabla \text{ (Del bar)} \equiv \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} = \frac{\partial}{\partial z} + \frac{\partial}{\partial \bar{z}} - i^2 \left(\frac{\partial}{\partial z} - \frac{\partial}{\partial \bar{z}} \right) = 2 \frac{\partial}{\partial z} \quad \dots (61)$$

Now taking $F(x, y)$ as real continuously differentiable function of scalars x, y and $A(x, y) = P(x, y) + i Q(x, y)$ as complex continuously differentiable function of vectors x and y , then $F(x, y)$

$$= F \left(\frac{z + \bar{z}}{2}, \frac{z - \bar{z}}{2i} \right) = G(z, \bar{z}) \text{ say and } A(x, y) = B(z, \bar{z}) \text{ say } \dots (62)$$

The Gradient of a real scalar function F is defined by

$$\text{grad } F = \nabla F = \frac{\partial F}{\partial x} + i \frac{\partial F}{\partial y} = 2 \frac{\partial G}{\partial \bar{z}} \quad \dots (63)$$

and the gradient of a complex vector function A is defined as

$$\text{grad } A = \nabla A = \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (P + iQ) = \frac{\partial P}{\partial x} - \frac{\partial Q}{\partial y} \\ + i \left(\frac{\partial P}{\partial y} + \frac{\partial Q}{\partial x} \right) = 2 \frac{\partial R}{\partial \bar{z}} \quad \dots (64)$$

Geometrically interpreted, ∇F is a vector normal to the curve $F(x, y) = \text{constant}$, and if B is analytic function of z so that $\frac{\partial B}{\partial \bar{z}} = 0$, then gradient is also zero i.e. $\frac{\partial P}{\partial x} = \frac{\partial Q}{\partial y}$, $\frac{\partial P}{\partial y} = -\frac{\partial Q}{\partial x}$ showing that Cauchy-Riemann equation are satisfied.

The Divergence of a complex vector function A is defined as

$$\text{div } A = \nabla \cdot A = \text{Re} \{ \bar{\nabla} A \} = \text{Re} \left\{ \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right\} (P + iQ) = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} \\ = 2 \text{Re} \left\{ \frac{\partial B}{\partial z} \right\} \text{ where Re denotes real part. } \dots (65)$$

It is notable that the divergence of a real or complex function is always a real function.

The Curl of a complex vector function A is defined as

$$\text{curl } A = \nabla \times A = I_m (\bar{\nabla} A) = I_m \left\{ \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (P + iQ) \right\} \\ = \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} = 2 I_m \left\{ \frac{\partial B}{\partial z} \right\} \quad \dots (66)$$

The *Laplacian operator* is defined as the scalar or dot product of ∇ with itself i.e. $\nabla \cdot \nabla = \nabla^2 = \text{Re} (\bar{\nabla} \nabla) = R \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right)$

$$= \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4 \frac{\partial^2}{\partial z \partial \bar{z}} \quad \dots (67)$$

Herebelow we summarize few identities involving grad, div and curl of two complex differentiable functions A_1 and A_2 .

$$\text{grad} (A_1 + A_2) = \text{grad} A_1 + \text{grad} A_2 \quad \dots (68)$$

$$\text{div} (A_1 + A_2) = \text{div} A_1 + \text{div} A_2 \quad \dots (69)$$

$$\text{curl} (A_1 + A_2) = \text{curl} A_1 + \text{curl} A_2 \quad \dots (70)$$

$$\text{grad} (A_1 A_2) = A_1 (\text{grad} A_2) + (\text{grad} A_1) A_2 \quad \dots (71)$$

$$\text{curl} (\text{grad} A) = 0, \text{ when } A \text{ is real or } I_m(A) \text{ is harmonic} \quad \dots (72)$$

$$\text{div} (\text{grad} A) = 0, \text{ when } A \text{ is imaginary or } \text{Re}(A) \text{ is harmonic} \quad \dots (73)$$

5.8. COMPLEX INTEGRATION

It has been defined that a function of z is said to be *regular* in any domain D if the function is single-valued and differentiable at every point of the domain D . Now in order to show that a regular function possesses a second derivative, we must first of all express the function $f(z)$ of z as a contour integral round any closed contour surrounding the point z .

Let us suppose that the equations $x = \phi(t)$ and $y = \psi(t)$, where $\alpha \leq t \leq \beta$ represent the arc of the plane curve. Let us divide the interval (α, β) by the points $t_0, t_1, t_2, \dots, t_n$ and let these points on the curve be denoted by $P_0, P_1, P_2, \dots, P_n$. Then length of the polygonal line $P_0 P_1 P_2 \dots P_n$ will be the sum of the lengths of the lines $P_0 P_1, P_1 P_2, \dots, P_{n-1} P_n$. If $z_0, z_1, z_2, \dots, z_n$ be the points on the arc corresponding to the values $t_0, t_1, t_2, \dots, t_n$ of t , then length of the polygonal arc $P_0 P_1 P_2 \dots P_n$

$$= \sum_{r=1}^n [(x_r - x_{r-1})^2 + (y_r - y_{r-1})^2].$$

The value of this sum depending upon the mode of subdivision is called the *length of an inscribed polygon*. If the arc is such that the length of all the inscribed polygons have a finite upper bound λ , the curve is known to be **Rectifiable** and λ is called the *length of the curve*.

The necessary and sufficient conditions for the arc to be rectifiable are that the functions $\phi(t), \psi(t)$ must be of bounded variation in the interval (α, β) . In case $\phi'(t), \psi'(t)$ are continuous, the curve denoted by $x = \phi(t), y = \psi(t)$, when $\alpha \leq t \leq \beta$ is rectifiable and its length is

$$s = \int_{\alpha}^{\beta} \sqrt{[\phi'(t)]^2 + [\psi'(t)]^2} dt.$$

Riemann's definition of Integration

Defining the integration as the limit of a sum, Riemann's definition of integration of a complex function is given as below;

Consider a function $w = f(z)$ which is continuous (but not necessarily analytic) along a curve C with end points A and B . Divide the arc C into n arcs by the points $z_0 = \alpha, z_1, z_2, \dots, z_{r-1}, z_r, \dots, z_{n-1}, z_n = \beta$ with z_0 being at A and z_n at B . Take points $\zeta_1, \zeta_2, \dots, \zeta_r, \zeta_n$ such that ζ_n lies on the arc $z_{r-1} z_r$.

Consider the sum $f(\zeta_1)(z_1 - z_0) + f(\zeta_2)(z_2 - z_1) + \dots + f(\zeta_r)(z_r - z_{r-1}) + \dots + f(\zeta_n)(z_n - z_{n-1})$

$$\text{i.e.} \quad \sum_{r=1}^n f(\zeta_r) (z_r - z_{r-1})$$

Fig. 5.13

If this sum tends to a unique limit when $n \rightarrow \infty$ i.e. the number of arcs of the curve C becomes indefinitely large then the definite integral of $f(z)$ along the curve C is given by

$$\int_C f(z) dz = \int_{AB} f(z) dz = \lim_{n \rightarrow \infty} \sum_{r=1}^n f(\zeta_r) (z_r - z_{r-1})$$

Note 1. If $f(z) = u + iv, z = x + iy = \phi(t) + i\psi(t)$ so that $dz = dx + i dy = \{\phi'(t) + i\psi'(t)\} dt, \alpha \leq t \leq \beta$.

We have $\int_C f(z) dz = \int_{\alpha}^{\beta} (u + iv) \{\phi'(t) + i\psi'(t)\} dt$

$$= \int_{\alpha}^{\beta} (u + iv) (dx + i dy) = \int_C (u dx - v dy) + i(v dx + u dy)$$

Note 2. If C consists of a number of arcs C_r ,


$$\int_C f(z) dz = \sum \int_{C_r} f(z) dz.$$

Continuous arc. When $\phi(t)$ and $\psi(t)$ are real continuous functions of the real variable t in the interval (α, β) , the arc is known as *continuous arc*.

Multiple points. If the relations $x = \phi(t)$ and $y = \psi(t)$ are satisfied by two or more values of t in the given integral, then the point $z = (x, y)$ is called a *Multiple point*.

Jordan arc. A continuous arc having no multiple points is called a *Jordan arc*.

Regular arc of a Jordan curve. Considering an arc of a Jordan curve defined by the equation $z = \phi(t) + i\psi(t)$, where $\alpha \leq t \leq \beta$, if z be expressed as single-valued and $\phi(t), \psi(t)$ as well as $\phi'(t), \psi'(t)$ are continuous in



the interval $\alpha \leq t \leq \beta$, the arc is then called *Regular arc of a Jordan Curve*.

Length of the regular Jordan arc is

$$\int_{\alpha}^{\beta} \sqrt{[\phi'(t)]^2 + [\psi'(t)]^2} dt.$$

A *Continuous Jordan Curve* is one which consists of a chain of finite number of continuous arcs.

Contour. By the word 'contour' we mean a continuous Jordan curve consisting of a chain of finite number of regular arcs. It is clear that a contour is rectifiable.

The complex integral of $f(z)$ along the regular arc L is written as

$$\int_L f(z) dz.$$

The integral of $f(z)$ along a contour C , which consists of a finite number of regular arcs L_r is given by

$$\int_C f(z) dz = \sum_r \int_{L_r} f(z) dz.$$

Here $\int_C f(z) dz$ is read as integral $f(z)$ taken over the closed contour C .

Some properties in case of complex integrals are however notable:

1. $\int_C \{f(z) + \phi(z)\} dz = \int_C f(z) dz + \int_C \phi(z) dz.$
2. $\int_C f(z) dz = - \int_{-C} f(z) dz$, where $-C$ represents the direction opposite to that of C .
3. $\int_C f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz$, where C_1 and C_2 are two parts of C .
4. $\int_C k f(z) dz = k \int_C f(z) dz$, k being constant.

An upper bound for a complex integral. If a function of z say $f(z)$ is continuous on a contour L of length l and if the inequality $|f(z)| \leq M$ is satisfied, then

$$|\int_L f(z) dz| \leq Ml.$$

Let the equation to the curve L be

$$x = \phi(t), y = \psi(t)$$

Then the length l of the curve is given by

$$l = \int \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right] dt, \text{ between proper limits. } \dots (1)$$

$$\text{Now } z = x + iy \text{ gives } dz = dx + i dy$$

$$|dz| = |dx + i dy| = \sqrt{(dx)^2 + (dy)^2}$$

$$\text{So that } \int |dz| = \int \sqrt{(dx)^2 + (dy)^2}$$

$$= \int \sqrt{\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2} dt$$

$$= l \text{ by (1)}$$

$$\dots (2)$$

Now the modulus of the sum of n complex numbers being less than or equal to the sum of their moduli, we have

$$\left| \sum_{r=1}^n f(\zeta_r) (z_r - z_{r-1}) \right| \leq \sum_{r=1}^n |f(\zeta_r) (z_r - z_{r-1})|$$

$$\leq \sum_{r=1}^n |f(\zeta_r)| |z_r - z_{r-1}|$$

Proceeding to the limit as $n \rightarrow \infty$, we have

$$\left| \int_C f(z) dz \right| \leq \int_C |f(z)| |dz|$$

$$\leq M \int_C |dz|$$

$$\leq Ml \text{ by (2)} \quad \therefore |f(z)| \leq M$$

3.3. CAUCHY'S THEOREM

If $f(z)$ is a regular function of z and if $f'(z)$ is continuous at each point within and on a closed contour C , then $\int_C f(z) dz = 0$.

(Agra, 1966, 67)

(i.e. the integral of the function round a closed contour is zero).

Elementary Proof. Green's theorem states that if $P(x, y)$, $Q(x, y)$, $\frac{\partial Q}{\partial x}$, $\frac{\partial P}{\partial y}$ are all continuous functions of x and y in the domain D , then

$$\int_C (P dx + Q dy) = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy \quad \dots (i)$$

Let us now assume that $f(z) = u + iv$, where $z = x + iy$.

$$dz = dx + i dy.$$

Substituting these values in $\int_C f(z) dz$, we get

$$\int_C f(z) dz = \int_C (u + iv) (dx + i dy)$$

$$= \int_C (u dx - v dy) + i \int_C (v dx + u dy)$$

$$= \iint_D \left(-\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) dx dy + i \iint_D \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) dx dy$$

[by result (i) of Green's theorem]

$$=0 \because \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \text{ and } \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

(Cauchy-Riemann equations)

Rigorous Proof of Cauchy's Theorem

If $f(z)$ is analytic (regular) at all points within and on the closed contour, C then $\int_C f(z) dz = 0$.

To prove this theorem let us first consider two lemmas.

Lemma I. If C is a closed contour, then we must have

$$\int_C dz = 0 \text{ and also } \int_C z dz = 0.$$

It follows from the definition of integral that

$$\int_C f(z) dz = \lim_{n \rightarrow \infty} \sum_{r=1}^n (z_r - z_{r-1}) f(z).$$

Taking $f(z)=1$, we have

$$\begin{aligned} \int_C dz &= \lim_{n \rightarrow \infty} \sum_{r=1}^n \{(z_r - z_{r-1}) \cdot 1\} \\ &= 0 \text{ as } \max. (z_r - z_{r-1}) \rightarrow 0, \text{ when } n \rightarrow \infty. \end{aligned}$$

$$\begin{aligned} \text{Also } \int_C z dz &= \lim_{n \rightarrow \infty} \sum_{r=1}^n \{(z_r - z_{r-1}) z\} \\ &= \lim_{n \rightarrow \infty} \sum_{r=1}^n \{z_{r-1} (z_r - z_{r-1})\} \\ &= \frac{1}{2} \left[\lim_{n \rightarrow \infty} \sum_{r=1}^n \{z_r (z_r - z_{r-1})\} \right. \\ &\quad \left. + \lim_{n \rightarrow \infty} \sum_{r=1}^n \{z_{r-1} (z_r - z_{r-1})\} \right] \\ &= \frac{1}{2} \lim_{n \rightarrow \infty} \sum \{(z_r + z_{r-1}) (z_r - z_{r-1})\} \\ &= \frac{1}{2} \lim_{n \rightarrow \infty} \sum (z_r^2 - z_{r-1}^2) \\ &= 0 \text{ (for a closed curve).} \end{aligned}$$

Lemma II. (Goursat's Lemma).

Given ϵ , it is possible by suitable transversals, to divide the interior of C into a finite number of meshes, either complete squares or parts of squares, such that within each mesh there is a point z_0 , such that

$$\left| \frac{f(z) - f(z_0)}{z - z_0} - f'(z_0) \right| < \epsilon,$$

$$\text{i.e. } f(z) - f(z_0) = f'(z_0)(z - z_0) + \eta(z - z_0) \quad \dots (1)$$

for all values of z in the mesh where $\eta < \epsilon$.

[Note. Unless the contour is a square, the sum of the meshes will not be a perfect square].




Fig. 5.14




Fig. 5.15

Let us suppose that the lemma is false and however the interior of C be sub-divided. Then there will be at least one mesh for which (1) is not true. We have to show that this necessarily implies the existence of a point within or on C at which $f(z)$ is not differentiable.

Suppose that we enclose C in a large square Γ , of area A and apply the process of repeated quadrisection. When Γ is quadrisectioned there is at least one of the four quarters of the square Γ for which (1) is untrue. Let it be denoted by Γ_1 . We quadrisection Γ_1 and take its quarter for which (1) does not hold. This process is carried on indefinitely. Let the infinite sequence of squares so obtained be $\Gamma_1, \Gamma_2, \Gamma_3, \dots, \Gamma_n, \dots$ each contained in the preceding, for which the lemma is not true. Let this sequence of squares determine a limiting point ζ which clearly lies within C .

Now $f(z)$ being analytic everywhere and so at $z=\zeta$, it is differentiable at ζ and therefore, we have

$$\left| \frac{f(z) - f(\zeta)}{z - \zeta} - f'(\zeta) \right| < \epsilon,$$

i.e. $f(z) - f(\zeta) = f'(\zeta)(z - \zeta) + \eta'(z - \zeta)$, where $|z - \zeta| < \delta$ and $\eta' < \epsilon$.

With centre ζ let us draw a circle of radius $\delta_1 < \delta$ and let

$$|z - \zeta| \leq \delta_1 < \delta,$$

which contradicts our hypothesis, for by taking ζ to be z_0 , (1) is satisfied and thus it follows Goursat's lemma.

Proof of the theorem. It is obvious that some of the meshes obtained by the subdivision of the interior of C will be squares and others will not be squares. Let $C_1, C_2, \dots, C_m, \dots$ be the complete squares and $D_1, D_2, \dots, D_n, \dots$ be the partial squares, then

$$\begin{aligned} \int_C f(z) dz &= \sum \int_{C_m} f(z) dz + \sum \int_{D_n} f(z) dz. \quad \dots (2) \end{aligned}$$




Fig. 5.16

Also we have from (1) of the Goursat's lemma

$$f(z) = f(z_0) + f'(z_0)(z - z_0) + \eta(z - z_0), \text{ where } |\eta| < \epsilon. \quad (1)$$

Now

$$\begin{aligned} \int_{C_n} f(z) dz &= \int_{C_n} [f(z_0) + f'(z_0)(z - z_0) + \eta(z - z_0)] dz \\ &= [f(z_0) - f'(z_0)z_0] \int_{C_n} dz + f'(z_0) \int_{C_n} z dz + \\ &\quad \eta \int_{C_n} (z - z_0) dz \\ &= \eta \int_{C_n} (z - z_0) dz \text{ as } \int_{C_n} dz = \int_{C_n} z dz = 0 \text{ from lemma 1} \end{aligned}$$

$$\begin{aligned} \therefore \left| \int_{C_n} f(z) dz \right| &\leq \eta \int_{C_n} |z - z_0| dz \\ &\leq \epsilon \int_{C_n} \sqrt{2} l_n |dz|, \text{ side of square being } l_n \end{aligned}$$

Max $|z - z_0| = \sqrt{2} l_n$, where $\sqrt{2} l_n$, is the length of the diagonal of square

$$\begin{aligned} &\leq \epsilon \sqrt{2} l_n \int_{C_n} ds, s \text{ being the entire perimeter of the square} \\ &\leq \epsilon \sqrt{2} l_n \cdot 4 l_n \\ &\leq 4 \sqrt{2} \epsilon \cdot l_n^2 \\ &\leq 4 \sqrt{2} \epsilon \cdot A_n, l_n^2 = A_n, \text{ the area of the square.} \end{aligned}$$

Similarly,

$$\begin{aligned} \left| \int_{D_n} f(z) dz \right| &\leq \epsilon \sqrt{2} l_n' \int_{D_n} ds \\ &\leq \epsilon \sqrt{2} l_n' (4 l_n' + s_n), s_n \text{ being length of arc forming the curved boundary of } D_n \\ &\leq \epsilon \cdot 4 \sqrt{2} A_n' + 2 \epsilon s_n l_n', A_n' \text{ is the area of square of side } l_n' \end{aligned}$$

Hence (2) gives

$$\begin{aligned} \left| \int_C f(z) dz \right| &\leq \epsilon \sqrt{2} \sum_{n=1}^{\infty} \{4(A_n + A_n') + s_n l_n'\} \\ &\leq \epsilon \sqrt{2} (4A + SL), S \text{ being perimeter of contour } C, L \end{aligned}$$

the length of a side of some square enclosing C and A the total area ≤ 0 as $\epsilon \rightarrow 0$, S and L both being finite,

$$\text{so that } \int_C f(z) dz = 0.$$

This proves the theorem.

Extension of Cauchy's Theorem

For this purpose, we define some theorems which have not yet been introduced.

Connected region. A region is known as a connected region if any two points of the region D can be connected by a curve lying wholly within the region.

Simply-connected region. A connected region is known as a simply-connected region if all the interior points of a closed curve C described in the region D , are also the point of D .

Multiply-connected region. A connected region is known as a multiply-connected region if all the points enclosed by two or more closed curves described in a region D are also the points of D .

Cross cut or simply cut. The lines drawn in a multiply-connected region, without intersecting any curve, such that the multiply-connected region is converted to a simply-connected region, are said to be cross cuts or cuts.

If the function $f(z)$ is not analytic in the whole region enclosed by a closed contour but it is analytic in the region enclosed between two closed contours then also Cauchy's integral theorem can be applied.

Let the nearly equal and parallel lines AB and $A'B'$ as shown in Fig. 5.17, be used as cross-cuts by connecting the points A and B (very near to each other) on outer contour C with points A' and B' on inner contour γ . Let the simply connected contour so obtained be denoted by Γ . The function $f(z)$ being analytic in this region, the Cauchy's integral formula can be applied for this contour i.e.




Fig. 5.17

$$\int_{\Gamma} f(z) dz = 0 \text{ now gives}$$

$$\begin{aligned} \int_{\Gamma} f(z) dz &= \int_C f(z) dz + \int_{AB} f(z) dz + \int_{A'B'} f(z) dz + \int_{\gamma} f(z) dz = 0 \\ &= \int_C f(z) dz + \int_{\gamma} f(z) dz = 0, \text{ other two integral being equal and opposite in sense, cancel each other.} \end{aligned}$$

$$\text{i.e. } \int_C f(z) dz = - \int_{\gamma} f(z) dz$$

where minus sign shows that the integral is traversed in clockwise direction.

$\therefore$ Taking the integral along γ in anticlockwise direction, we get

$$\int_C f(z) dz = \int_{\gamma} f(z) dz.$$

Note. In general if C be a closed curve and $C_1, C_2, C_3, \dots, C_n$, be the other n closed curves lying inside C and $f(z)$ is analytic within these curves, then $\int_C f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz + \dots + \int_{C_n} f(z) dz$ integrals being taken in anti-clockwise direction.

Moreras' Theorem as Converse of Cauchy's Theorem

If the integral $\int f(z) dz$ of a function $f(z)$ which is continuous in a region D , is zero when taken round any simple closed contour in D , then $f(z)$ is analytic in D .

Taking z_0 as a fixed point and z any variable point in D , the value of the integral.

$$\int_{z_0}^z f(\zeta) d\zeta = F(z) \text{ say} \quad \dots (4)$$

is independent of the curve joining z_0 to z and is dependent of z only:

(4) may be interpreted as:

$$F(z+h) = \int_{z_0}^{z+h} f(\zeta) d\zeta \quad \dots (5)$$

So that (4) and (5) give

$$\begin{aligned} F(z+h) - F(z) &= \int_{z_0}^{z+h} f(\zeta) d\zeta - \int_{z_0}^z f(\zeta) d\zeta \\ &= \int_z^{z+h} f(\zeta) d\zeta \end{aligned}$$

The integration on right being independent of the curve joining z to $z+h$, may be taken along the straight line joining z to $z+h$, so that

$$\begin{aligned} \frac{F(z+h) - F(z)}{h} - f(z) &= \frac{1}{h} \int_z^{z+h} f(\zeta) d\zeta - \frac{1}{h} f(z) \int_z^{z+h} d\zeta \\ &= \frac{1}{h} \int_z^{z+h} [f(\zeta) - f(z)] d\zeta \\ \therefore \left| \frac{F(z+h) - F(z)}{h} - f(z) \right| &= \left| \frac{1}{h} \int_z^{z+h} [f(\zeta) - f(z)] d\zeta \right| \\ &= \frac{1}{|h|} \int_z^{z+h} |f(\zeta) - f(z)| |d\zeta| \\ &\leq \frac{1}{|h|} \int_z^{z+h} |d\zeta| \text{ since } f(z) \text{ being} \\ &\quad \text{continuous in } D, \\ &\quad |f(\zeta) - f(z)| < \epsilon \text{ for } |\zeta - z| < \delta \\ &\leq \frac{1}{|h|} \epsilon \in |h| = \epsilon \rightarrow 0 \text{ as } h \rightarrow 0 \end{aligned}$$

$$\therefore \lim_{h \rightarrow 0} \frac{F(z+h) - F(z)}{h} = F'(z) = f(z)$$

showing that $F(z)$ exists for all z in D i.e. $F(z)$ is analytic in D . As such $F'(z)$ and so $f(z)$ is also analytic in D , for, the derivative of an analytic function is also analytic.

5.10. CAUCHY'S INTEGRAL FORMULA

[Agra, 1974]

If the function $f(z)$ is regular within and on a closed contour C and if ζ be a point within C , then

$$f(\zeta) = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z - \zeta}.$$

Let us describe about the point $z = \zeta$ a small circle γ of radius δ lying entirely within C . Now consider a function

$$\phi(z) = \frac{f(z)}{z - \zeta}$$

which is regular in the region between C and γ .

By making a cross-cut joining any point of γ to any point of C by two almost equal and parallel lines, let us form a closed contour $LMCM'L'\gamma L = \Gamma$ (say) within which the function $\psi(z)$ is regular, so that by Cauchy's theorem, we get

$$\int_{\Gamma} \psi(z) dz = 0, \text{ where } \psi(z) = \frac{f(z)}{z - \zeta}.$$

The function is analytic within and on the boundary of the contour and as points M, M' are very near,

$LM = L'M'$ and $LM \parallel L'M'$ approximately.

It follows that if the contour Γ is described in anti-clockwise direction (i.e., positive sense), the cross-cut is traversed twice, once in each sense, and hence we have

$$\begin{aligned} \int_C \phi(z) dz - \int_{\gamma} \phi(z) dz &= 0 \\ \int_C \phi(z) dz &= \int_{\gamma} \phi(z) dz, \\ \int_C \frac{f(z)}{z - \zeta} dz &= \int_{\gamma} \frac{f(z)}{z - \zeta} dz. \end{aligned} \quad \dots (1)$$

Referring to the adjoining figure,




Fig. 5.19

$$\begin{aligned} \vec{QP} &= \vec{QO} + \vec{OP} \\ &= -\zeta + z = z - \zeta. \end{aligned}$$

$\therefore$ Complex co-ordinate of QP is $z - \zeta = \delta e^{i\theta}$.

$$\begin{aligned} [\because \text{if } z = x + iy, \\ \therefore z = r(\cos \theta + i \sin \theta) \\ = re^{i\theta}], \end{aligned}$$

where θ is the argument and δ is the magnitude of QP which is very small.

Differentiating (2) partially, we get $dz = i\delta e^{i\theta} d\theta$. $\dots (2)$




Fig. 5.18

Now (1) may be written as

$$\begin{aligned} \int_C \frac{f(z)}{z-\zeta} dz &= \int_\gamma \frac{f(\zeta)}{z-\zeta} dz + \int_\gamma \frac{f(z)-f(\zeta)}{z-\zeta} dz \\ &= f(\zeta) \int_0^{2\pi} \frac{i\delta e^{i\theta} d\theta}{\delta e^{i\theta}} + I \text{ (say)} \\ &= 2\pi i f(\zeta) + I, \end{aligned}$$

where

$$\begin{aligned} |I| &= \left| \int_\gamma \frac{f(z)-f(\zeta)}{z-\zeta} dz \right| \\ &\leq \int_\gamma \left| \frac{f(z)-f(\zeta)}{z-\zeta} \right| |dz|. \end{aligned}$$

But from the definition of continuity, the function $f(z)$ is continuous at $z=\zeta$, when

$$|f(z)-f(\zeta)| < \delta_1 \text{ if } |z-\zeta| < \delta.$$

Also if

$$z = x + iy,$$

$$dz = dx + i dy.$$

$$\therefore |dz| = \sqrt{(dx)^2 + (dy)^2} = ds.$$

$$\text{So, } |I| \leq \frac{\delta_1}{\delta} \int_\gamma ds, \text{ where } \int_\gamma ds \text{ means the entire circumference of the circle}$$

$$\leq \frac{\delta_1}{\delta} \times 2\pi\delta$$

$$\leq 2\pi\delta_1$$

$$\leq 0$$

as $\delta_1 \rightarrow 0$ which is so when $\delta \rightarrow 0$

But we know that modulus of any quantity cannot be negative, therefore $|I|=0$ as $\delta_1=0$, i.e., when z coincides with ζ .

Hence

$$\int_C \frac{f(z)}{z-\zeta} dz = 2\pi i f(\zeta),$$

i.e.

$$f(\zeta) = \frac{1}{2\pi i} \int_C \frac{f(z)}{z-\zeta} dz$$

(Agra, 1961)

5.11. SOME RESULTS BASED ON CAUCHY'S INTEGRAL FORMULA

1. If $f(z)$ is regular in a domain D , then its derivatives at any point $z=\zeta$ of the region D is also regular in that domain and is given by

$$f'(\zeta) = \frac{1}{2\pi i} \int_C \frac{f(z)}{(z-\zeta)^2} dz$$

where C is any single closed contour in D surrounding the point $z=\zeta$.

$$\text{Now, } f'(\zeta) = \lim_{h \rightarrow 0} \frac{f(\zeta+h) - f(\zeta)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{1}{2\pi i} \int_C \frac{f(z)}{h} \left\{ \frac{1}{z-\zeta-h} - \frac{1}{z-\zeta} \right\} dz$$

applying Cauchy's integral formula,

$$= \lim_{h \rightarrow 0} \frac{1}{2\pi i} \int_C \frac{f(z)}{(z-\zeta)(z-\zeta-h)} dz$$

$$= \frac{1}{2\pi i} \lim_{h \rightarrow 0} \int_C \frac{f(z)}{(z-\zeta)(z-\zeta-h)} dz$$

$$= \frac{1}{2\pi i} \lim_{h \rightarrow 0} \int_C \left\{ \frac{1}{(z-\zeta)^2} + \frac{h}{(z-\zeta)^3} \right\} f(z) dz$$

$$= \frac{1}{2\pi i} \int_C \frac{f(z)}{(z-\zeta)^2} dz + \frac{1}{2\pi i} \lim_{h \rightarrow 0} \int_C \frac{hf(z)}{(z-\zeta)^3} dz$$

$$= \frac{1}{2\pi i} \int_C \frac{f(z)}{(z-\zeta)^2} dz + I \text{ (say),}$$

where

$$I = \frac{1}{2\pi i} \lim_{h \rightarrow 0} \int_C \frac{hf(z)}{(z-\zeta)^3} dz$$

$$\therefore |I| \leq \lim_{h \rightarrow 0} \frac{|h|}{2\pi} \int_C \frac{|f(z)| |dz|}{|z-\zeta|^3 |z-\zeta-h|}$$

$$\leq \lim_{h \rightarrow 0} \frac{|h|}{2\pi} \cdot \frac{Ml}{d^2(d-h)} \text{ since } f(z) \text{ is regular in and on } C, \text{ it is bounded, so that}$$

$$\leq 0 \text{ as } h \rightarrow 0$$

$$I=0 \text{ as } a \rightarrow 0$$

$|f(z)| \leq M$ on C . Let d be the lower bound of the distance of ζ from C , i.e. $|z-\zeta| > d$ and l be the length of C .

$$\text{Hence } f'(\zeta) = \frac{1}{2\pi i} \int_C \frac{f(z)}{(z-\zeta)^2} dz \quad \dots (1)$$

Similarly the second order derivative of $f(z)$ at $z=\zeta$ of domain D may be found as

$$f''(\zeta) = \lim_{h \rightarrow 0} \frac{f'(\zeta+h) - f'(\zeta)}{h}$$

Applying Cauchy's integral formula, in the form (1)

$$f''(\zeta) = \frac{1}{2\pi i} \lim_{h \rightarrow 0} \frac{1}{h} \left\{ \frac{1}{(z-\zeta-h)^2} - \frac{1}{(z-\zeta)^2} \right\} f(z) dz$$

$$= \frac{1}{2\pi i} 2 \int_C \frac{f(z)}{(z-\zeta)^3} dz + \lim_{h \rightarrow 0} h \int_C R(z) f(z) dz$$

Since

$$\frac{d}{dz} \left[\frac{1}{h} \left\{ \frac{1}{(z-\zeta-h)^2} - \frac{1}{(z-\zeta)^2} \right\} \right]$$

$$= \frac{2}{(z-\zeta)^3} + h R(z) \text{ where } R(z) \text{ is bounded on } C$$

so that

$$\int_C f(z) R(z) \text{ is finite.}$$

$$= \frac{2}{2\pi i} \int_C \frac{f(z)}{(z-\zeta)^3} dz$$

$$= \frac{2}{2\pi i} \int_C \frac{f(z)}{(z-\zeta)^3} dz$$

Proceeding similarly we can show that

$$f''(\zeta) = \frac{2!}{2\pi i} \int_C \frac{f(z)}{(z-\zeta)^4} dz \text{ etc.}$$

Here below the general result follows:

2. If a function $f(z)$ is regular in a domain D then $f(z)$ has at any point $z=\zeta$ of the domain D , derivatives of all orders, values being given by $f^{(n)}(\zeta) = \frac{n!}{2\pi i} \int_C \frac{f(z) dz}{(z-\zeta)^{n+1}}$.

Let us suppose that the theorem is valid for $n=m$ and then consider

$$f^{(m+1)}(\zeta) = \lim_{h \rightarrow 0} \frac{f^{(m)}(\zeta+h) - f^{(m)}(\zeta)}{h}$$

$$= \frac{(m+1)!}{2\pi i} \int_C \frac{f(z) dz}{(z-\zeta)^{m+2}} + I \text{ by the last result,}$$

where $I = \frac{(m+1)!}{2\pi i} \lim_{h \rightarrow 0} \int_C \frac{hf(z) dz}{(z-\zeta)^{m+2} (z-\zeta-h)}$.

It is easy to show as in the previous result that

$$|I| \rightarrow 0 \text{ as } |h| \rightarrow 0.$$

$$\therefore f^{(m+1)}(\zeta) = \frac{(m+1)!}{2\pi i} \int_C \frac{f(z) dz}{(z-\zeta)^{m+2}},$$

which shows that the result is true for $n=m+1$ and hence we have

$$\text{in general } f^{(n)}(\zeta) = \frac{n!}{2\pi i} \int_C \frac{f(z) dz}{(z-\zeta)^{n+1}}.$$

5.12. TAYLOR'S EXPANSION

(Agra, 1966, 68)

If a function $f(z)$ is regular everywhere inside a circle C whose centre is the point $z=a$ and radius is ρ , such that $|z-a| \leq \rho$ and if ζ is a point such that $|\zeta-a| = r < \rho$,

$$\text{then } f(\zeta) = \sum_{n=0}^{\infty} a_n (\zeta-a)^n,$$

$$\text{where } a_n = \frac{f^{(n)}(a)}{n!}$$

We know that

$$\frac{1}{z-\zeta} = \frac{1}{z-a} + \frac{\zeta-a}{(z-a)(z-\zeta)}$$

$$= \frac{1}{z-a} + \frac{\zeta-a}{z-a} \left[\frac{1}{z-a} + \frac{\zeta-a}{(z-a)(z-\zeta)} \right]$$




Fig. 5.20

$$= \frac{1}{z-a} + \frac{\zeta-a}{(z-a)^2} + \frac{(\zeta-a)^2}{(z-a)^3} \left[\frac{1}{z-a} + \frac{\zeta-a}{(z-a)(z-\zeta)} \right]$$

$$\dots \dots \dots \dots \dots \dots \dots \dots$$

$$\dots \dots \dots \dots \dots \dots \dots \dots$$

$$= \frac{1}{z-a} + \frac{\zeta-a}{(z-a)^2} + \frac{(\zeta-a)^2}{(z-a)^3} + \dots + \frac{(\zeta-a)^n}{(z-a)^n (z-\zeta)}.$$

Hence by the Cauchy's Integral formula, we get

$$f(\zeta) = \frac{1}{2\pi i} \int_C \frac{f(z)}{z-\zeta} dz$$

$$= \frac{1}{2\pi i} \int_C \left[\frac{1}{z-a} + \frac{\zeta-a}{(z-a)^2} + \dots + \frac{(\zeta-a)^n}{(z-a)^n (z-\zeta)} \right] f(z) dz$$

$$= \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z-a} + \frac{\zeta-a}{2\pi i} \int_C \frac{f(z) dz}{(z-a)^2} + \dots$$

$$+ \frac{(\zeta-a)^n}{2\pi i} \int_C \frac{f(z) dz}{(z-a)^n (z-\zeta)}$$

$$= f(a) + (\zeta-a) f'(a) + \dots + \frac{(\zeta-a)^{n-1} f^{(n-1)}(a)}{(n-1)!} + R_n.$$

where $R_n = \frac{(\zeta-a)^n}{2\pi i} \int_C \frac{f(z) dz}{(z-a)^n (z-\zeta)}$, the remainder after n terms in Taylor's Theorem.

Now, since $f(\zeta)$ may be written as

$$f(\zeta) = f(a + \zeta - a).$$

the expansion

$$f(\zeta) = f(a) + f'(a) (\zeta-a) + \frac{f''(a)}{2!} (\zeta-a)^2 + \dots$$

represents the Taylor's Theorem.

On assumptions $|\zeta-a| = r < \rho$,

$$|z-a| = \rho' \text{ and } r < \rho' < \rho,$$

as well as taking maximum value of $|f(z)|$ as M , i.e., $|f(z)| \leq M$ on C , we have

$$|R_n| \leq \frac{|\zeta-a|^n}{2\pi} \int_C \frac{|f(z)| |dz|}{|z-a|^n |z-\zeta|}$$

$$\leq \frac{r^n}{2\pi} \cdot \frac{M}{\rho'^{(n)} (\rho'-r)} \int_C ds \left[\because |z-\zeta| = |z-a - (\zeta-a)| \right]$$

$$\leq \frac{r^n}{2\pi} \cdot \frac{M}{\rho'^{(n)} (\rho'-r)} \cdot L, \text{ where } L \text{ is the length of the entire circumference and hence } L=2\pi\rho'$$

$$\leq \frac{r^n}{2\pi} \cdot \frac{M}{\rho'^{(n)} (\rho'-r)} \cdot 2\pi\rho'$$

$$\leq \left(\frac{r}{\rho'}\right)^n \frac{M}{1 - \frac{r}{\rho'}}$$

$$\leq 0 \text{ if } n \rightarrow \infty, \text{ since } \frac{r}{\rho'} < 1.$$

$$\therefore R_n = 0, \text{ when } n \rightarrow \infty.$$

Thus we have the infinite series

$$\begin{aligned} f(\zeta) &= f(a) + f'(a)(\zeta - a) + \dots \\ &= \sum_{n=0}^{\infty} a_n (\zeta - a)^n, \text{ where } a_n = \frac{f^n(a)}{n!} \end{aligned}$$

Note 1. Maclaurin's Expansion as a particular case of Taylor's Expansion.

In Taylor's expansion if $a=0$, we have

$$f(\zeta) = \sum_{n=0}^{\infty} a_n \zeta^n = \sum_{n=0}^{\infty} \frac{f^n(0)}{n!} \zeta^n$$

Note 2. Here $|R_n| \rightarrow 0$ as $n \rightarrow \infty$, it follows that the series $\sum_{n=0}^{\infty} a_n (\zeta - a)^n$ is convergent and has $f(\zeta)$ as its sum-function. Also if the function $f(z)$ is regular in the whole z -plane, the expansion is valid for all values of ζ .

Note 3. If the function $f(\zeta)$ has a maximum $M(r)$ on $|\zeta - a| = r < \rho$, then, $|a_n| \leq \frac{M(r)}{r^n}$, where $a_n = \frac{f^n(a)}{n!}$




Fig. 5.21

$\therefore$ If C' be the circle $|\zeta - a| = r$, then as in Cauchy's theorem,

$$a_n = \frac{1}{2\pi i} \int_{C'} \frac{f(z) dz}{(z - a)^{n+1}}.$$

$$\therefore |a_n| \leq \frac{1}{2\pi} \int_{C'} \frac{|f(z)| |dz|}{|z - a|^{n+1}}.$$

The Taylor's expansion is

$$\begin{aligned} f(\zeta) &= f(a) + f'(a)(\zeta - a) \\ &\quad + \frac{f''(a)}{2!} (\zeta - a)^2 + \dots \\ &\quad \dots + \frac{f^n(a)}{n!} (\zeta - a)^n + \dots \\ &= a_0 + a_1 (\zeta - a) + \dots \\ &\quad \dots + a_n (\zeta - a)^n + \dots \end{aligned}$$

$$\begin{aligned} \text{where } a_n &= \frac{f^n(a)}{n!} \\ &= \frac{1}{2\pi i} \int_C \frac{f(z) dz}{(z - a)^{n+1}} \end{aligned}$$

How if

$$|z - a| = \rho' > r; \frac{1}{\rho'} < \frac{1}{r}.$$

$$\begin{aligned} |a_n| &\leq \frac{M(r)}{2\pi} \int_{C'} \frac{ds}{r^{n+1}} \\ &\leq \frac{M(r)}{2\pi \cdot r^{n+1}} \cdot 2\pi r \\ &\leq \frac{M(r)}{r^n}. \end{aligned}$$

Note 4. (Liouville's theorem). If the function $f(z)$ is regular in the whole z -plane then $f(z)$ must be a constant provided $|f(z)| < k$ for all values of z .

Since $f(z)$ is analytic for all values of z , then by Taylor's theorem, we have

$$f(z) = a_0 + a_1 z + a_2 z^2 + \dots + a_n z^n + \dots = \sum_{n=0}^{\infty} a_n z^n,$$

where $|a_n| \leq \frac{M(r)}{r^n}$ and $|f(z)| = M(r)$ when $|z| = r$.

As given that $|f(z)| < k$ for all values of z including ∞ ,

$$M(r) < k.$$

Hence

$$|a_n| \leq \frac{k}{r^n}$$

$$\leq 0 \text{ when } n \rightarrow \infty \text{ for } n < 0.$$

$$a_n = 0 \text{ for } n > 0.$$

As such $f(z) = a_0$ is a constant.

13. LAURENT'S EXPANSION

(Agra, 1966)

Let there be an annulus between two concentric circles C_1 and C_2 of centre $z = a$ and radii ρ_1 and ρ_2 ($\rho_1 > \rho_2$); then if $f(z)$ be regular within the annulus between C_1 and C_2 , as well as on the circles, and ζ be any point of the annulus,

$$f(\zeta) = \sum_{n=0}^{\infty} a_n (\zeta - a)^n + \sum_{n=1}^{\infty} b_n (\zeta - a)^{-n},$$

$$\text{where } a_n = \frac{1}{2\pi i} \int_{C_1} \frac{f(z) dz}{(z - a)^{n+1}} \text{ and } b_n = \frac{1}{2\pi i} \int_{C_2} (z - a)^{n-1} f(z) dz.$$

We have proved in the Taylor's theorem that

$$\frac{1}{z - \zeta} = \frac{1}{z - a} + \frac{z - a}{(z - a)^2} + \dots + \frac{(\zeta - a)^n}{(z - a)^{n+1} (z - \zeta)} + \dots \quad (1)$$

Interchanging z and ζ with each other we have

$$\frac{1}{\zeta - z} = \frac{1}{\zeta - a} + \frac{\zeta - a}{(\zeta - a)^2} + \dots + \frac{(z - a)^n}{(\zeta - a)^{n+1} (\zeta - z)} + \dots$$

$$\text{i.e. } -\frac{1}{z-\zeta} = \frac{1}{\zeta-a} + \frac{z-a}{(\zeta-a)^2} + \dots + \frac{(z-a)^n}{(\zeta-a)^{n+1}(\zeta-z)} \quad \dots (1)$$




Fig. 5.22

Now, we know that

$$f(\zeta) = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z-\zeta} \text{ by Cauchy's integral}$$

Therefore by making a cross joining any point of the circle C_1 to any point of the circle C_2 , we get

$$\begin{aligned} f(\zeta) &= \frac{1}{2\pi i} \int_{C_1} \frac{f(z) dz}{z-\zeta} - \frac{1}{2\pi i} \int_{C_2} \frac{f(z) dz}{z-\zeta} \\ &= \frac{1}{2\pi i} \int_{C_1} \frac{1}{z-\zeta} f(z) dz + \frac{1}{2\pi i} \int_{C_2} -\frac{1}{z-\zeta} f(z) dz. \end{aligned}$$

Substituting the values of $\frac{1}{z-\zeta}$ and $-\frac{1}{z-\zeta}$ from (1) and (2), we get

$$\begin{aligned} f(\zeta) &= \frac{1}{2\pi i} \int_{C_1} \left[\frac{1}{z-a} + \frac{\zeta-a}{(z-a)^2} + \dots + \frac{(\zeta-a)^n}{(z-a)^{n+1}(\zeta-z)} \right] f(z) dz \\ &\quad + \frac{1}{2\pi i} \int_{C_2} \left[\frac{1}{\zeta-a} + \frac{z-a}{(\zeta-a)^2} + \dots + \frac{(z-a)^n}{(\zeta-a)^{n+1}(z-\zeta)} \right] f(z) dz. \end{aligned}$$

Here term by term integration is possible as the two series are uniformly convergent.

$$\begin{aligned} \therefore f(\zeta) &= f(a) + (\zeta-a) f'(a) + \frac{(\zeta-a)^2}{2!} f''(a) + \dots + P_n \\ &\quad + \frac{b_1}{\zeta-a} + \frac{b_2}{(\zeta-a)^2} + \frac{b_3}{(\zeta-a)^3} + \dots + Q_n, \end{aligned}$$

where

$$\begin{aligned} b_1 &= \frac{1}{2\pi i} \int_{C_2} f(z) dz, \\ b_2 &= \frac{1}{2\pi i} \int_{C_2} f(z-a) f(z) dz, \\ &\dots \dots \dots \dots \\ b_n &= \frac{1}{2\pi i} \int_{C_2} (z-a)^{n-1} f(z) dz \end{aligned}$$

and P_n is the remainder after n th term as in the Taylor's expansion.

It is easy to show (as in Taylor's expansion) that $|P_n| \rightarrow 0$ as $n \rightarrow \infty$.

$$\text{Also } Q_n = \frac{1}{(\zeta-a)^n} \cdot \frac{1}{2\pi i} \int_{C_2} \frac{(z-a)^n}{\zeta-a} f(z) dz.$$

$$\begin{aligned} |Q_n| &\leq \frac{1}{r^n} \cdot \frac{1}{2\pi} \int_{C_2} \frac{|z-a|^n}{|\zeta-z|} |f(z)| |dz| \\ &\leq \frac{\rho_2^n}{r^n} \cdot \frac{M'}{2\pi} \int_{C_2} \frac{ds}{r-\rho_2} \\ &\leq \frac{\rho_2^n}{r^n} \cdot \frac{M'}{2\pi} \cdot \frac{2\pi\rho_2}{r-\rho_2} \\ &\leq \left(\frac{\rho_2}{r}\right)^{n+1} \frac{M'}{1-\frac{\rho_2}{r}} \\ &\leq 0 \text{ as } n \rightarrow \infty, \text{ since } \frac{\rho_2}{r} < 1 \end{aligned}$$

$$\left\{ \begin{array}{l} \because \rho_1 > |\zeta-a| > \rho_2 \\ \text{and Max. value of } |f(z)| \text{ is } M' \\ \text{Also } \zeta-z = \zeta-a-(z-a), \\ \therefore |\zeta-z| > |\zeta-a| - |z-a| > r-\rho_2, \\ \text{i.e. } \frac{1}{|\zeta-z|} > \frac{1}{r-\rho_2}. \end{array} \right.$$

Since modulus of any quantity cannot be negative.

$\therefore Q_n = 0$, when $n \rightarrow \infty$.

$$\begin{aligned} \text{Hence } f(\zeta) &= f(a) + (\zeta-a) f'(a) + \frac{(\zeta-a)^2}{2!} f''(a) + \dots \\ &\quad + \frac{b_1}{\zeta-a} + \frac{b_2}{(\zeta-a)^2} + \frac{b_3}{(\zeta-a)^3} + \dots \\ &= \sum_{n=0}^{\infty} a_n (\zeta-a)^n + \sum_{n=1}^{\infty} b_n (\zeta-a)^{-n}. \quad \dots (3) \end{aligned}$$

Note 1. The integrals giving the values of a_n and b_n as

$$\begin{aligned} a_n &= \frac{1}{2\pi i} \int_{C_1} \frac{f(z) dz}{(z-a)^{n+1}} \text{ and } b_n = \frac{1}{2\pi i} \int_{C_2} \frac{f(z) dz}{(z-a)^{n+1}} \\ &= \frac{1}{2\pi i} \int_{C_2} (z-a)^{n-1} f(z) dz \end{aligned}$$

are analytic every where in the annulus of C_1 and C_2 and therefore by Cauchy's theorem the path of integration can be taken any concentric circle C_0 lying between C_1 and C_2 , so that

$$b_n = \frac{1}{2\pi i} \int_{C_0} (z-a)^{n-1} f(z) dz = \frac{1}{2\pi} \int_{C_0} \frac{f(z) dz}{(z-a)^{-n+1}} = a_{-n}$$

$$\text{Thus } f(\zeta) = \sum_{-\infty}^{\infty} a_n (\zeta-a)^n \text{ where } a_n = \frac{1}{2\pi i} \int_{C_0} \frac{f(z) dz}{(z-a)^{n+1}}.$$

Note 2. Zeros and Singularities of an analytic complex function.

A zero of an analytic function $f(z)$ is defined to be a value of $z = \zeta$ such that $f(\zeta) = 0$.

Taylor's expansion gives

$$f(z) = f(\zeta) + (z-\zeta) f'(\zeta) + \frac{(z-\zeta)^2}{2!} f''(\zeta) + \dots + \frac{(z-\zeta)^n}{n!} f^{(n)}(\zeta) + \dots$$

Here if $f(\zeta) = 0$ and $f'(\zeta) \neq 0$ then $f(z)$ is said to have a **Simple zero** at $z = \zeta$.

If $f(\zeta) = f'(\zeta) = f''(\zeta) = \dots = f^{(n-1)}(\zeta) = 0$ and $f^{(n)}(\zeta) \neq 0$, then the point $z = \zeta$ is known as *zero of order n* of the function $f(z)$, since then

$$\begin{aligned} f(z) &= \frac{(z-\zeta)^n}{n!} f^{(n)}(\zeta) + \frac{(z-\zeta)^{n+1}}{(n+1)!} f^{(n+1)}(\zeta) + \dots \\ &= (z-\zeta)^n \left[\frac{f^{(n)}(\zeta)}{n!} + \frac{(z-\zeta)}{(n+1)!} + \dots \right] \\ &= (z-\zeta)^n \phi(z) \end{aligned} \quad \dots (4)$$

$$\text{where } \phi(z) = \frac{f^{(n)}(\zeta)}{n!} + \frac{z-\zeta}{(n+1)!} + \dots$$

= a non-zero and analytic function at $z = \zeta$.

It follows from (4) that the point $z = \zeta$ may be defined as a *zero of order n* of the function $f(z)$ if $f(z) = (z-\zeta)^n \phi(z)$

for $f(z)$ being analytic and non-zero function at $z = \zeta$.

If we draw a small circle C about ζ as centre then within it $\phi(z) \neq 0$. Also for $z \neq \zeta$ within C , $(z-\zeta)^n \neq 0$ and hence ζ is the only zero of the function $f(z)$ within C . As such the zero point $z = \zeta$ is *isolated*. But ζ being arbitrary may be any zero point of $f(z)$ and therefore all the zeros of an analytic functions $f(z)$ are isolated.

A point at which a function $f(z)$ ceases to be regular (analytic) is termed as a *singular point* of $f(z)$ and the function $f(z)$ is said to have a *singularity* at this point.

In case there is no other singularity in the neighbourhood of a singular point $z = \zeta$, the function $f(z)$ is said to have an *isolated singularity*. We can surround such a singularity by a small circle Γ with ζ as centre such that in the annulus made by Γ with a larger concentric circle C , the function $f(z)$ is analytic. Then we can expand $f(z)$ by Laurent's expansion as

$$f(z) = \sum_{n=0}^{\infty} a_n (z-\zeta)^n + \dots + \frac{b_1}{z-\zeta} + \frac{b_2}{(z-\zeta)^2} + \dots + \frac{b_m}{(z-\zeta)^m} + \dots$$

where the terms containing b 's are termed as *principal part of the expansion* at the singularity $z = \zeta$.

There arise three possibilities:

(i) All the coefficients b 's are zero so that the function is equal to the analytic function except at $z = \zeta$. Such a singularity is defined as *non-essential* or *removable singularity* of $f(z)$.

(ii) The principal part is an infinite series. Then the point $z = \zeta$ is said to be an *essential* or *non-removable singularity* of $f(z)$.

(iii) The principal part contains a finite number of terms i.e.

$$\text{the principal number part} = \frac{b_1}{z-\zeta} + \frac{b_2}{(z-\zeta)^2} + \dots + \frac{b_m}{(z-\zeta)^m}$$

$$\text{so that } b_{m+1} = b_{m+2} = \dots = 0$$

then the function $f(z)$ is said to have a *pole or singularity of order m* . In this case, we have

$$\begin{aligned} f(z) &= \sum_{n=0}^{\infty} a_n (z-\zeta)^n + \sum_{n=1}^m b_n (z-\zeta)^{-n} \\ &= (z-\zeta)^{-m} \left[\sum_{n=0}^{\infty} a_n (z-\zeta)^{m+n} + b_1 (z-\zeta)^{m-1} + b_2 (z-\zeta)^{m-2} \right. \\ &\quad \left. + \dots + b_m \right] \\ &= (z-\zeta)^{-m} \phi(z) = \frac{\phi(z)}{(z-\zeta)^m}, \text{ where } \phi(z) \text{ is an analytic function at } z = \zeta \end{aligned}$$

when $m = 1$, the pole is known as a *simple pole*.

For a *pole of order m* of $z = \zeta$, the limiting value of the function is given by

$$\begin{aligned} \lim_{z \rightarrow \zeta} f(z) &= \lim_{z \rightarrow \zeta} \sum_{n=1}^m b_n (z-\zeta)^{-n} \\ \lim_{z \rightarrow \zeta} |f(z)| &= \lim_{z \rightarrow \zeta} \left| \sum_{n=1}^m b_n (z-\zeta)^{-n} \right| \\ &= \lim_{z \rightarrow \zeta} \left| \frac{b_m}{(z-\zeta)^m} + \frac{b_{m-1}}{(z-\zeta)^{m-1}} + \dots + \frac{b_1}{z-\zeta} \right| \\ &= \lim_{z \rightarrow \zeta} \frac{1}{|z-\zeta|^m} \{ b_m + b_{m-1}(z-\zeta) + \dots + b_1(z-\zeta)^{m-1} \} \\ &\geq \lim_{z \rightarrow \zeta} \frac{1}{|z-\zeta|^m} \{ |b_m| + |b_{m-1}(z-\zeta)| + \dots \\ &\quad + |b_1(z-\zeta)^{m-1}| \} \end{aligned}$$

$$\begin{aligned} &\geq \lim_{z \rightarrow \zeta} \frac{1}{|z-\zeta|^m} |b_m| \\ &\geq \infty \end{aligned}$$

i.e. $|f(z)| \rightarrow \infty$ as $z \rightarrow \zeta$

Conclusively if $f(z)$ is analytic in a given region except at a point $z = \zeta$ and $\lim_{z \rightarrow \zeta} f(z) \rightarrow \infty$, then $f(z)$ is said to have a *pole or singularity* at $z = \zeta$.

For the pole of order m , we have

$$\lim_{z \rightarrow \zeta} (z-\zeta)^m f(z) = b_m = \text{constant } (\neq 0)$$

and the value of the function is given by $\frac{\phi(z)}{(z-\zeta)^m}$.

In case there does not exist a value of m such that $\lim_{z \rightarrow \zeta} (z-\zeta)^m f(z)$

is finite, then the point $z = \zeta$ is an *essential or non-removable singularity* of $f(z)$.

If an analytic function $f(z)$ has a pole of order m at the point $z=\zeta$ then $\frac{1}{f(z)}$ has a zero of order m at this point.

$z=\zeta$ will be a pole of order m of $f(z)$ if $(z-\zeta)^m f(z)$ is analytic and non-zero at $z=\zeta$, so that

$$\phi(z) = \frac{1}{(z-\zeta)^m f(z)} \text{ is analytic and non-zero}$$

$\therefore \frac{1}{f(z)} = (z-\zeta)^m \phi(z)$ follows that $\frac{1}{f(z)}$ has a zero of order m as $z=\zeta$.

A similar procedure will show that if $\frac{1}{f(z)}$ has a pole of order m then $f(z)$ will have a pole of order m .

It also follows that since zeros are isolated, the poles must be isolated.

Note 3. The Point at infinity. In treating the complex variables it is convenient to regard infinity as a point. The point infinity is considered by putting $z = \frac{1}{\zeta}$ in $f(z)$ so that the behaviour of $f(z)$ at infinity is examined by the behaviour of $f\left(\frac{1}{\zeta}\right)$ at $\zeta=0$.

Consequently $f(z)$ is analytic or has a zero or has a simple pole or has an essential singularity etc., at infinity according as $f\left(\frac{1}{\zeta}\right)$ analytic or has a zero or has a simple pole or has an essential singularity at $\zeta=0$.

e.g. if $f\left(\frac{1}{\zeta}\right)$ has a zero of order m at $\zeta=0$ then writing $\phi(\zeta)$ for $f\left(\frac{1}{\zeta}\right)$ its expansion at $\zeta=0$ by Taylor's theorem is $f\left(\frac{1}{\zeta}\right) = \phi(\zeta) = a_m \zeta^m + a_{m+1} \zeta^{m+1} + \dots$ so that corresponding expansion of $f(z)$ at $z=\infty$ is $f(z) = \frac{a_m}{z^m} + \frac{a_{m+1}}{z^{m+1}} + \dots$ which does not contain constant terms and terms containing positive exponent powers of z and hence $f(z)$ has a zero of order m at infinity.

Similarly if $f\left(\frac{1}{\zeta}\right)$ has a pole of order m at $\zeta=0$ then by Laurent's expansion

$$f\left(\frac{1}{\zeta}\right) = \phi(\zeta) = \sum_{n=0}^{\infty} a_n \zeta^n + \sum_{n=1}^m b_n \zeta^{-n}$$

So that corresponding expansion of $f(z)$ at $z=\infty$ is $f(z) = \sum_{n=1}^m b_n z^n + \sum_{n=0}^{\infty} a_n z^{-n}$ which has $\sum_{n=1}^m b_n z^n$ as its principal part as $z=\infty$ and this

principal part being a finite series of positive powers of z at $z=\infty$ shows that $f(z)$ has a pole of order m at infinity.

Problem 13. The function $f(z)$ is regular when $|z| < R'$. Prove that if $|a| < R < R'$, $f(a) = \frac{1}{2\pi i} \int_C \frac{R^2 - a\bar{a}}{(z-a)(R^2 - z\bar{a})} f(z) dz$, where C is the circle $|z|=R$. Deduce Poisson's formula that if $0 < r < R$,

$$f(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \frac{R^2 - r^2}{R^2 - 2Rr \cos(\theta - \phi) + r^2} f(Re^{i\phi}) d\phi.$$

Taking centre as origin and a line OP making an angle θ with the horizontal, let us take a point Q on OP produced, such that $OP \cdot OQ = R^2$, where $|z|=R$; then Q is called the point inverse to P w.r.t the circle of radius R .

Let at P , $z=a$, and at Q , $z=a_1$ be the complex co-ordinates.

Now $|OP| \cdot |OQ| = R^2$.

$$\therefore |OQ| = \frac{R^2}{|OP|} > R \text{ or } R > |OP|.$$

Therefore Q must lie outside the circle.

$$\text{Again } a_1 = OQ \cdot e^{i\theta} = \frac{R^2}{OP} e^{i\theta}$$

$$= \frac{R^2}{OP \cdot e^{-i\theta}} \text{ as } a = OP \cdot e^{i\theta},$$

$$= \frac{R^2}{\bar{a}}$$

$\therefore$ conjugate of a i.e. $\bar{a} = OP \cdot e^{-i\theta}$.

$$\text{By Cauchy's theorem, } f(a) = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z-a} \dots (1)$$

Also the Cauchy's theorem states that if $F(z)$ is a regular function and if $F'(z)$ is continuous at each point within and on a closed contour C , then $\int_C F(z) dz = 0$ and thus as point Q is outside the circle and $f(z)$ is analytic, i.e.,

$$F(z) = \frac{f(z)}{z-a_1} \neq 0$$

therefore

$$0 = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z-a_1} \dots (2)$$

Subtracting (2) from (1), we get

$$f(a) = \frac{1}{2\pi i} \int_C \frac{(a-a_1) f(z) dz}{(z-a)(z-a_1)}$$




Fig. 5.23

$$\begin{aligned}
 &= \frac{1}{2\pi i} \int_C \left(a - \frac{R^2}{\bar{a}} \right) \frac{f(z) dz}{(z-a) \left(z - \frac{R^2}{\bar{a}} \right)} \\
 &= \frac{1}{2\pi i} \int_C \frac{(R^2 - a\bar{a}) f(z) dz}{(z-a) (R^2 - z\bar{a})} \dots (1)
 \end{aligned}$$

To deduce Poisson's formula, if $0 < r < R$.

Put $a = re^{i\theta}$; $\therefore a = re^{-i\theta}$,

$z = Re^{i\phi}$, i.e., $dz = iRe^{i\phi} d\phi$.

Being a circle, limits of ϕ are from 0 to 2π .

$$\begin{aligned}
 \therefore (z-a) (R^2 - \bar{a}z) &= (Re^{i\phi} - re^{i\theta}) (R^2 - re^{-i\theta} \cdot Re^{i\phi}) \\
 &= Re^{i\phi} (Re^{i\phi} - re^{i\theta}) (Re^{-i\phi} - re^{-i\theta}) \\
 &= Re^{i\phi} [R^2 + r^2 - 2Rr \cos(\theta - \phi)]
 \end{aligned}$$

$$\begin{aligned}
 \text{and } R^2 - \bar{a}a &= R^2 - re^{-i\theta} re^{i\theta} [\because e^{i(\theta-\phi)} + e^{-i(\theta-\phi)}] \\
 &= R^2 - r^2 = 2 \cos(\theta - \phi)
 \end{aligned}$$

Substituting these values in result (3), we get

$$f(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \frac{(R^2 - r^2) f(Re^{i\phi}) d\phi}{R^2 - 2Rr \cos(\theta - \phi) + r^2}$$

Problem 14. The function $f(z)$ is regular in $|z-a| < R$. Prove that if $0 < r < R$, $f'(a) = \frac{1}{\pi r} \int_0^{2\pi} P(\theta) e^{-i\theta} d\theta$, where $P(\theta)$ is the real part of $f(a + re^{i\theta})$.

$$\begin{aligned}
 \text{We know that } f'(a) &= \frac{1}{2\pi i} \int_C \frac{f(z) dz}{(z-a)^2} \cdot \text{Put } z-a = re^{i\theta}, \\
 &= \frac{1}{2\pi i} \int_0^{2\pi} f(a + re^{i\theta}) e^{-i\theta} d\theta \dots (1)
 \end{aligned}$$

By Taylor's Theorem,

$$\begin{aligned}
 f(z) &= \sum_{n=0}^{\infty} a_n (z-a)^n = \sum_{n=0}^{\infty} a_n r^n e^{in\theta} \\
 &= \sum_{n=0}^{\infty} R_n e^{i\phi_n} \cdot r^n e^{in\theta}, \text{ where } a_n \text{ is a complex quantity} \\
 &\quad \text{and therefore may be written as} \\
 &\quad \quad \quad a_n = R_n e^{i\phi_n} \\
 &= \sum_{n=0}^{\infty} R_n r^n e^{i(n\theta + \phi_n)}.
 \end{aligned}$$

Let the conjugate of $f(z)$ be $\bar{f}(z)$; then

$$\bar{f}(z) = \sum_{n=0}^{\infty} R_n r^n e^{-i(n\theta + \phi_n)}$$

$$\begin{aligned}
 \therefore \frac{1}{2\pi i} \int_C \frac{\bar{f}(z)}{(z-a)^2} dz &= \frac{1}{2\pi i} \int_0^{2\pi} e^{-i\theta} d\theta \cdot \sum_{n=0}^{\infty} R_n r^n \cdot e^{-i(n\theta + \phi_n)} \\
 &= \frac{1}{2\pi i} \sum_{n=0}^{\infty} R_n \cdot r^n \int_0^{2\pi} e^{-i\{(n+1)\theta + \phi_n\}} d\theta \\
 &= \frac{1}{2\pi i} \sum_{n=0}^{\infty} R_n \cdot r^n \int_0^{2\pi} e^{-i\phi_n} \cdot e^{-i(n+1)\theta} d\theta \\
 &= 0, \dots (2)
 \end{aligned}$$

$$\left[\text{since } \left\{ \frac{e^{-i(n+1)\theta}}{-i(n+1)} \right\}_0^{2\pi} = 0 \text{ or } \left\{ \cos(n+1)\theta - i \sin(n+1)\theta \right\}_0^{2\pi} = 0 \right]$$

Adding (1) and (2), we have

$$\begin{aligned}
 f'(a) &= \frac{1}{2\pi r} \int_0^{2\pi} [f(z) + \bar{f}(z)] e^{-i\theta} d\theta \\
 &= \frac{2}{2\pi r} \int_0^{2\pi} P(\theta) e^{-i\theta} d\theta. \text{ Since } z + \bar{z} = 2R(z), \\
 &= \frac{1}{\pi r} \int_0^{2\pi} P(\theta) e^{-i\theta} d\theta. \therefore f(z) + \bar{f}(z) = 2Rf(z) = 2P(\theta)
 \end{aligned}$$

as given that $P(\theta)$ is the real part of $f(a + re^{i\theta})$.

Problem 15. By using the integral representation of $f^n(a)$, prove that

$$\left(\frac{x^n}{n!} \right)^2 = \frac{1}{2\pi i} \int_C \frac{x^n \cdot e^{xz}}{n! z^{n+1}} dz$$

where C is any closed contour surrounding the origin. Hence prove

$$\text{that } \sum_{n=0}^{\infty} \left(\frac{x^n}{n!} \right)^2 = \frac{1}{2\pi} \int_0^{2\pi} e^{2x \cos \theta} d\theta.$$

We know that

$$f^n(a) = \frac{n!}{2\pi i} \int_C \frac{f(z) dz}{(z-a)^{n+1}} \dots (1)$$

$$\therefore f^n(0) = \frac{n!}{2\pi i} \int_C \frac{f(z) dz}{z^{n+1}} \dots (2)$$

Let us put

$$f(z) = e^{xz}.$$

Differentiating n times w.r.t. z , we have

$$f^n(z) = x^n e^{xz}.$$

$\therefore$

$$f^n(0) = x^n.$$

Substituting its value in (2), we get

$$\frac{x^n}{n!} = \frac{1}{2\pi i} \int_C \frac{e^{xz} dz}{z^{n+1}}.$$

Multiplying both sides by $\frac{x^n}{n!}$, we have

$$\left(\frac{x^n}{n!}\right)^2 = \frac{1}{2\pi i} \int_C \frac{x^n}{n!} \cdot \frac{e^{xz}}{z^{n+1}} dz.$$

Now $\sum_{n=0}^{\infty} \left(\frac{x^n}{n!}\right)^2 = \frac{1}{2\pi i} \int_C \frac{e^{xz}}{z} dz \cdot \sum_{n=0}^{\infty} \frac{x^n}{n! \cdot z^n}$

$$= \frac{1}{2\pi i} \int_C \frac{e^{xz}}{z} dz \cdot \sum_{n=0}^{\infty} \frac{(x/z)^n}{n!}$$

$$= \frac{1}{2\pi i} \int_C \frac{e^{xz}}{z} dz \cdot e^{x/z}$$

$$= \frac{1}{2\pi i} \int_C \frac{e^{x(z+1/z)}}{z} dz.$$

... (3)

If we put

$$z = e^{i\theta}, \quad dz = ie^{i\theta} d\theta$$

and

$$\frac{1}{z} = e^{-i\theta}, \quad \frac{dz}{z} = i d\theta,$$

so that

$$z + \frac{1}{z} = 2 \cos \theta.$$

Substituting its value in (3), we find

$$\sum_{n=0}^{\infty} \left(\frac{x^n}{n!}\right)^2 = \frac{1}{2\pi} \int_0^{2\pi} e^{2x \cos \theta} d\theta.$$

Problem 16. Obtain the expansion

$$f(z) = f(a) + 2 \left\{ \frac{z-a}{2} f' \left(\frac{z+a}{2} \right) + \frac{(z-a)^3}{2^3 \cdot 3!} f''' \left(\frac{z+a}{2} \right) + \frac{(z-a)^5}{2^5 \cdot 5!} f^{(5)} \left(\frac{z+a}{2} \right) + \dots \right\}$$

and determine its range of validity.

We know that

$$f(z) = f \left(\frac{z+a}{2} + \frac{z-a}{2} \right) \text{ and } f(a) = f \left(\frac{z+a}{2} - \frac{z-a}{2} \right).$$

Expanding them by Taylor's expansion, we get

$$f(z) = f \left(\frac{z+a}{2} \right) + \left(\frac{z-a}{2} \right) f' \left(\frac{z+a}{2} \right) + \left(\frac{z-a}{2} \right)^2 \frac{1}{2!} f'' \left(\frac{z+a}{2} \right) + \dots \quad \dots (1)$$

$$\text{and } f(a) = f \left(\frac{z+a}{2} \right) - \left(\frac{z-a}{2} \right) f' \left(\frac{z+a}{2} \right) + \frac{1}{2!} \left(\frac{z-a}{2} \right)^2 f'' \left(\frac{z+a}{2} \right) + \dots \quad \dots (2)$$

Subtracting (2) from (1),

$$f(z) - f(a) = 2 \left\{ \frac{z-a}{2} f' \left(\frac{z+a}{2} \right) + \frac{1}{3!} \left(\frac{z-a}{2} \right)^3 f''' \left(\frac{z+a}{2} \right) + \dots \right\}.$$

$$f(z) = f(a) + 2 \left\{ \frac{z-a}{2} f' \left(\frac{z+a}{2} \right) + \frac{(z-a)^3}{2^3 \cdot 3!} f''' \left(\frac{z+a}{2} \right) + \dots \right\}$$

Range of validity or the condition for the validity of Taylor's expansion is that

$$\left| \frac{z-a}{2} \right| < \delta \text{ or } \left| z - \frac{z+a}{2} \right| < \delta,$$

from which it is obvious that z must be in the neighbourhood of the point a .

Problem 17. Prove that $\cosh \left(z + \frac{1}{z} \right) = a_0 + \sum_1^{\infty} a_n \left(z^n + \frac{1}{z^n} \right)$,

where $a_n = \frac{1}{2\pi} \int_0^{2\pi} \cos n\theta \cosh (2 \cos \theta) d\theta.$

Laurent's expansion yields

$$\cosh \left(z + \frac{1}{z} \right) = a_0 + \sum_1^{\infty} a_n z^n + \sum_1^{\infty} b_n z^{-n}. \quad \dots (1)$$

Putting $\frac{1}{z}$ for z , we get

$$\cosh \left(\frac{1}{z} + z \right) = a_0 + \sum_1^{\infty} \frac{a_n}{z^n} + \sum_1^{\infty} b_n z^n. \quad \dots (2)$$

It is obvious from the two expansions i.e., from (1) and (2) that $b_n = b_n$.

By Laurent's expansion, we know that

$$f(z) = \sum_0^{\infty} a_n (z-a)^n + \sum_1^{\infty} \frac{b_n}{(z-a)^n}.$$

$$\text{then } a_n = \frac{1}{2\pi i} \int_{C_1} \frac{f(z) dz}{(z-a)^{n+1}} \text{ and } b_n = \frac{1}{2\pi i} \int_{C_2} f(z) (z-a)^{n-1} dz.$$

As such in the existing case,

$$a_n = \frac{1}{2\pi i} \int_C \frac{\cosh (z+1/z) dz}{z^{n+1}} \text{ and } b_n = \frac{2}{2\pi i} \int_C \cosh \left(z + \frac{1}{z} \right) dz \cdot z^{n-1}.$$

Putting $z = e^{i\theta}$ by taking a circle of unit radius,

$$\frac{1}{z} = e^{-i\theta}, \text{ so that } z + \frac{1}{z} = 2 \cos \theta$$

and

$$dz = ie^{i\theta} d\theta, \quad \frac{dz}{z} = i d\theta, \text{ we get}$$

$$a_n = \frac{1}{2\pi} \int_0^{2\pi} \frac{\cosh (e^{i\theta} + e^{-i\theta}) e^{i\theta} d\theta}{e^{i(n+1)\theta}}$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \cosh (e^{i\theta} + e^{-i\theta}) e^{-ni\theta} d\theta$$

and

$$b_n = \frac{1}{2\pi} \int_0^{2\pi} \cosh (e^{i\theta} + e^{-i\theta}) e^{ni\theta} d\theta,$$

$$\therefore 2a_n = 2b_n = \frac{1}{2\pi} \int_0^{2\pi} \cosh(2 \cos \theta) \cdot 2 \cos n\theta \, d\theta,$$

$$\text{i.e., } a_n = b_n = \frac{1}{2\pi} \int_0^{2\pi} \cosh(2 \cos \theta) \cdot \cos n\theta \, d\theta.$$

5.14. RESIDUE AND CONTOUR INTEGRATION

Definition of residue. The residue of a function $f(z)$ at the pole $z=a$ is defined to be the coefficient of $(z-a)^{-1}$ in the Laurent's expansion of the function $f(z)$, i.e.,

$$f(z) = \sum_{n=0}^{\infty} a_n (z-a)^n + \sum_{n=1}^m b_n (z-a)^{-n},$$

where $z=a$ is a pole of order m .

If $z=a$ be the pole of order one, then the residue is

$$b_1 = \lim_{z \rightarrow a} (z-a) f(z),$$

i.e., in case of a simple pole $z=a$.

$$\text{Residue} = \lim_{z \rightarrow a} (z-a) f(z).$$

Now consider the integral $b_n = \frac{1}{2\pi i} \int_{C_2} (z-a)^{n-1} f(z) \, dz$ which the value of b_n in Laurent's expansion. Here the circle C_2 is arbitrary and may therefore be replaced by any closed contour C containing within it no other singularities except $z=a$. Thus

$$\frac{1}{2\pi i} \int_{C_2} f(z) \, dz = \frac{1}{2\pi i} \int_C f(z) \, dz,$$

where $n=1$, i.e., pole is of order one,

$$\text{i.e., } b_1 = \frac{1}{2\pi i} \int_C f(z) \, dz.$$

Hence as an alternative, the residue of a function $f(z)$ at the pole $z=a$ is equal to $\frac{1}{2\pi i} \int_C f(z) \, dz$, where C is a closed contour containing within it the only singularity $z=a$ and integration being taken round C in anti-clockwise direction, i.e., the positive sense.

Similarly the residue of $f(z)$ at infinity, i.e., at the point $z=\infty$ is $\frac{1}{2\pi i} \int_C f(z) \, dz$ taken round C in clockwise direction, as it is negative w.r.t. the interior of C and positive w.r.t. the exterior of C provided the value of this integral is definite.

5.15. CAUCHY'S RESIDUE THEOREM (Rohilkhand, 1971)

If the function $f(z)$ be single-valued, continuous and regular within and on a closed contour C , except a finite number of poles (singularities) within C , then

$$\int_C f(z) \, dz = 2\pi i \sum R$$

where $\sum R$ represents the sum of the residues say $R_1, R_2, R_3, \dots, R_n$ of $f(z)$ at the poles $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n$ (say) within A . (Vikram, 1969)

Let us draw a set of circles $\gamma_1, \gamma_2, \dots, \gamma_n$ with centres $\alpha_1, \alpha_2, \dots, \alpha_n$ and radius ρ , such that they do not intersect each other and lie entirely within the closed curve C . Then $f(z)$ is regular within the region enclosed between C and the small circles $\gamma_1, \gamma_2, \dots, \gamma_n$. The entire region A may be deformed to consist of these small circles and the polygon P . Now by Cauchy's theorem, we have

$$\int_C f(z) \, dz = \int_P f(z) \, dz + \sum_{r=1}^n \int_{\gamma_r} f(z) \, dz.$$




Fig. 5.24

But the integral round the polygon P vanishes since $f(z)$ is regular within and on the closed contour P . Therefore

$$\int_C f(z) \, dz = \sum_{r=1}^n \int_{\gamma_r} f(z) \, dz.$$

Let us now consider $z=a$, a pole of order m ; then by Laurent's expansion,

$$f(z) = \sum_{n=0}^{\infty} a_n (z-a)^n + \sum_{s=0}^m \frac{b_s}{(z-a)^s} = \phi(z) + \sum_{s=1}^m \frac{b_s}{(z-a)^s}.$$

where $\phi(z) = \sum_{n=0}^{\infty} a_n (z-a)^n$ is regular within and on γ_r and has no pole.

$$\therefore \int_{\gamma_r} f(z) \, dz = \int_{\gamma_r} \phi(z) \, dz + \sum_{s=1}^m b_s \int_{\gamma_r} \frac{dz}{(z-a)^s}.$$

where $\int_{\gamma_r} \phi(z) \, dz = 0$ by Cauchy's fundamental theorem.

$$\therefore \int_{\gamma_r} f(z) \, dz = \sum_{s=1}^m b_s \int_{\gamma_r} \frac{dz}{(z-a)^s}.$$

Putting $z-a = \rho e^{i\theta}$, where θ varies from 0 to 2π ,
 $dz = \rho i e^{i\theta} d\theta$.

As the point z makes a circuit which consists of the circle γ_r , therefore

$$\int_{\gamma_r} f(z) \, dz = \sum_{s=1}^m b_s \rho i \int_0^{2\pi} \frac{e^{i\theta} d\theta}{\rho^s e^{is\theta}}$$

$$= \sum_{s=1}^m b_s \rho^{(1-s)} i \int_0^{2\pi} e^{i(1-s)\theta} d\theta.$$

$$\text{Now } \int_0^{2\pi} e^{i(1-s)\theta} d\theta = \left[\frac{e^{i(1-s)\theta}}{i(1-s)} \right]_0^{2\pi} = 0 \text{ if } s \neq 1.$$

But if $s=1$, all the terms will be zero except one.

$$\therefore \int_{\gamma_r} f(z) dz = b_1 i \int_0^{2\pi} d\theta = 2\pi i b_1,$$

where b_1 is called the residue for the function.

Let the residues for $r=1, 2, 3, \dots, n$ be respectively $R_1, R_2, R_3, \dots, R_n$.

$$\text{then } \int_{\gamma_1} f(z) dz = 2\pi i R_1,$$

$$\int_{\gamma_2} f(z) dz = 2\pi i R_2,$$

$$\dots \dots \dots$$

$$\int_{\gamma_n} f(z) dz = 2\pi i R_n.$$

$$\text{Hence } \int_C f(z) dz = \sum_{r=1}^m \int_{\gamma_r} f(z) dz = 2\pi i \sum R.$$

5.16. ANOTHER IMPORTANT THEOREM

Let C be the arc $\theta_1 \leq \arg(z-a) \leq \theta_2$ of the circle $|z-a|=r$ and if $\lim_{z \rightarrow a} \{(z-a)f(z)\} = b$, then

$$\lim_{r \rightarrow 0} \int_C f(z) dz = ib(\theta_2 - \theta_1).$$

In order to prove it, let us assume $F(z) = (z-a)f(z)$; then

$$F(a) = \lim_{z \rightarrow a} (z-a)f(z) = b \text{ (given)}.$$

Since $f(z)$ and $(z-a)$ both are individually continuous, therefore their product $(z-a)f(z)$ must also be continuous.

For a given ϵ , $\eta(\epsilon)$ can be so found that if $|z-a| < \eta$, $|\delta| < \epsilon$, where $(z-a)f(z) = b \pm \delta$.

[That's why if $|z-a| < \eta$,

$$|F(z) - F(a)| = \delta < \epsilon,$$

$$\text{i.e., } |F(z) - b| = \delta < \epsilon.$$

$$\text{So } F(z) = b \pm \delta.$$

Let us now take a contour whose centre is $z=a$ and radius is r and $f(z)$ between any two points say A and C , i.e. along the curve AC ; given by




Fig. 5.25

$$\int_{AC} f(z) dz = \int_{AC} \frac{F(z)}{z-a} dz = \int_{AC} \frac{b \pm \delta}{z-a} dz. \quad \dots (1)$$

Let us assume that $z-a = re^{i\theta}$.

$$\therefore dz = ire^{i\theta} d\theta \text{ and } \frac{dz}{z-a} = i d\theta.$$

Substituting these values in (1), we get

$$\begin{aligned} \int_{AC} f(z) dz &= i \int_{\theta_1}^{\theta_2} (b \pm \delta) d\theta \\ &= ib(\theta_2 - \theta_1) \pm i\delta(\theta_2 - \theta_1). \end{aligned}$$

Now $r \rightarrow 0$ as $z \rightarrow a$, $\therefore |F(z) - F(a)| = \delta$, $F(a) = b$ and $z-a = re^{i\theta}$ and $z \rightarrow a$ as $\delta \rightarrow 0$; $\therefore F(z) = b \pm \delta = F(a) \pm \delta$.

Conclusively when $r \rightarrow 0$, $\delta \rightarrow 0$.

$$\text{Hence } \lim_{r \rightarrow 0} \int_{AC} f(z) dz = ib(\theta_2 - \theta_1), \text{ where } \lim_{z \rightarrow a} (z-a)f(z) = b.$$

By similar procedure we can show that

If C be the arc $\theta_1 \leq \arg z \leq \theta_2$ of the circle $|z|=R$ and if $\lim_{R \rightarrow \infty} f(z) = b$ uniformly, then

$$\lim_{R \rightarrow \infty} \int_C f(z) dz = ib(\theta_2 - \theta_1).$$

Cor. If $\theta_1 = 0$, $\theta_2 = 2\pi$, then AC covers the whole circumference of the circle C (say), then

$$\int_C f(z) dz = 2\pi ib.$$

5.17. COMPUTATION OF RESIDUE

(i) Residue of $f(z)$ for a simple pole $z=a$, i.e. a pole of order one. This has been already established that

$$\text{residue of } f(z) \text{ at } z=a \text{ is } = \lim_{z \rightarrow a} (z-a)f(z).$$

If we put $f(z) = \frac{\phi(z)}{\psi(z)}$, where $\psi(z) = (z-a)F(z)$ provided $F(a) \neq 0$ but $\psi(a) = 0$, then

$$\begin{aligned} \lim_{z \rightarrow a} (z-a)f(z) &= \lim_{z \rightarrow a} \frac{(z-a)\phi(z)}{\psi(z)} \\ &= \lim_{z \rightarrow a} \frac{(z-a)\phi\{a + (z-a)\}}{\psi(a + (z-a))} \\ &= \lim_{z \rightarrow a} \frac{(z-a)[\phi(a) + (z-a)\phi'(a) + \dots]}{\psi(a) + (z-a)\psi'(a) + \dots} \end{aligned}$$

expanding by Taylor's theorem

$$= \lim_{z \rightarrow a} \frac{\phi(a) + (z-a)\phi'(a) + \dots}{\psi'(a) + \frac{1}{2}(z-a)\psi''(a) + \dots}$$

$$= \frac{\phi(a)}{\psi'(a)}.$$

$$\therefore \psi(a) = 0$$

Hence for a simple pole, residue at $z=a$ is $\frac{\phi(a)}{\psi'(a)}$.

(ii) Residue at $z=a$, a pole of order m . Let $f(z)$ be of the form

$$f(z) = \frac{\phi(z)}{(z-a)^m}.$$

We have proved that Residue b_1 at $z=a$ is $\frac{1}{2\pi i} \int_C f(z) dz$.

Hence in the existing case,

$$\text{Residue} = \frac{1}{2\pi i} \int_C \frac{\phi(z)}{(z-a)^m} dz = \frac{\phi^{m-1}(a)}{(m-1)!}$$

by Results of Cauchy's Integral.

Hence Residue of $f(z)$ at $z=a$, a pole of order m , is

$$= \frac{\phi^{m-1}(a)}{(m-1)!}.$$

(iii) Residue of $f(z)$ at a pole $z=a$ of any order. Laurent's Expansion

$$\text{is } f(z) = \sum_{n=0}^{\infty} a_n (z-a)^n + \sum_{n=1}^{\infty} \frac{b_n}{(z-a)^n}$$

$$= \sum_{n=0}^{\infty} a_n (z-a)^n + \frac{b_1}{z-a} + \frac{b_2}{(z-a)^2} + \dots + \frac{b_m}{(z-a)^m} + \dots$$

Putting $z-a=t$, i.e. $z=a+t$, we have

$$f(a+t) = \sum_{n=0}^{\infty} a_n t^n + \frac{b_1}{t} + \frac{b_2}{t^2} + \dots + \frac{b_m}{t^m} + \dots,$$

where b_1 the residue of $f(z)$ at $z=a$ the pole of any order is the coefficient of $\frac{1}{t}$. Hence coefficient of $\frac{1}{t}$ is the residue of $f(z)$ at $z=a$, a pole of any order, in the expansion of $f(z)$ after putting $z=a+t$.

(iv) Residue at infinity. We have already defined that residue of $f(z)$ at $z=\infty$ is $\frac{1}{2\pi i} \int_C f(z) dz$ taken clockwise, (i.e. in negative sense) round a large circle C within which all the finite singularities are enclosed.

Putting $z = \frac{1}{x}$, i.e. $dz = -\frac{dx}{x^2}$ the above integral yields

$$\frac{1}{2\pi i} \int_{\gamma} f\left(\frac{1}{x}\right) \left(-\frac{dx}{x^2}\right),$$

round a small circle γ in anti-clockwise direction, (i.e. in positive sense).

$$\text{Thus residue of } f\left(\frac{1}{x}\right) \text{ at } x=0 \text{ is } = \lim_{x \rightarrow 0} \left[x \left\{ -\frac{1}{x^2} f\left(\frac{1}{x}\right) \right\} \right]$$

$$= \lim_{x \rightarrow 0} \left[-\frac{1}{x} f\left(\frac{1}{x}\right) \right]$$

if it has got a definite value.

Hence residue of $f(z)$ at $z=\infty$ is $\lim_{z \rightarrow \infty} \{-zf(z)\}$.

(i) Residue of $f(z)$ at infinity is the negative of the coefficient of $1/z$ in the expansion of $f(z)$ for values of z in the neighbourhood of $z=\infty$. If $f(z)$ has a pole of order m at infinity. Then $f(1/z)$ has a pole of order m at $z=0$. As such $f(1/z)$ can be expanded in a Laurent's series for values of z in the annulus $0 < |z| < \rho$, where ρ is small

$$f\left(\frac{1}{z}\right) = \sum_{n=0}^{\infty} a_n z^n + \sum_{n=1}^m b_n \left(\frac{1}{z}\right)^n. \quad \dots (1)$$

Replacement of $1/z$ by z yields the expansion of $f(z)$ in the neighbourhood of $z=\infty$,

$$f(z) = \sum_{n=1}^m b_n z^n + \sum_{n=0}^{\infty} a_n z^{-n}. \quad \dots (2)$$

that the residue of $f(z)$ at infinity

$$= -\frac{1}{2\pi i} \int_C f(z) dz \text{ (by definition of residue)}$$

$$= -\frac{1}{2\pi i} \int_C \left[\sum_{n=1}^m b_n z^n + \sum_{n=0}^{\infty} a_n z^{-n} \right] dz \text{ from (2)}$$

$$= -\frac{1}{2\pi i} \int_C \frac{a_1}{z} dz, \text{ other integrals vanish, being of the form}$$

$$\int_C \frac{dz}{z^p}, p \neq 1$$

$$= -\frac{1}{2\pi i} a_1 \int_C \frac{dz}{z}$$

$$= -\frac{1}{2\pi i} a_1 \cdot 2\pi i \because \int_C \frac{dz}{z} = 2\pi i$$

$$= -a_1$$

= the negative of the coefficient of $1/z$ in the expansion of $f(z)$ for values of z in the neighbourhood of $z=\infty$.

INTEGRATION ROUND THE UNIT CIRCLE

It is notable that in case of complex integration a curve is generally known as a contour and the process of integration along a contour is known as contour integration.

Consider $\int_0^{2\pi} \phi(\cos \theta, \sin \theta) d\theta$, where $\phi(\cos \theta, \sin \theta)$ is a rational function of $\sin \theta$ and $\cos \theta$.




Fig. 5.26

Let us draw a circle of unit radius and take its centre as origin. Let us now consider a point P on its circumference with complex co-ordinate $z = e^{i\theta}$.

$$\therefore dz = ie^{i\theta} d\theta = iz d\theta \text{ and } \frac{1}{z} = e^{-i\theta}$$

$$\text{i.e.} \quad d\theta = \frac{dz}{iz}$$

$$\text{Also } z + \frac{1}{z} = 2 \cos \theta$$

$$\text{and } z - \frac{1}{z} = 2i \sin \theta.$$

Substituting these values in the given integral,

$$\begin{aligned} \int_0^{2\pi} \phi(\cos \theta, \sin \theta) d\theta &= \frac{1}{i} \int_C \phi \left[\frac{z + \frac{1}{z}}{2}, \frac{z - \frac{1}{z}}{2i} \right] \cdot \frac{dz}{z} \\ &= \frac{1}{i} \int_C \frac{F(z)}{z} dz, \end{aligned}$$

which may be integrated, thereby giving the value of the required complex integral.

5.19. EVALUATION OF $\int_{-\infty}^{\infty} f(x) dx$.

Let $f(z)$ be a function such that

(i) $f(z)$ is analytic throughout the upper half plane except at certain points which are its poles.

(ii) $f(z)$ has no poles on the real axis, i.e., if $R \rightarrow \infty$ (R being radius of semi-circle), then it will cover entire upper half plane.

(iii) $zf(z) \rightarrow 0$ uniformly as $|z| \rightarrow \infty$ for $0 \leq \arg z \leq \pi$.

(iv) $\int_0^{\infty} f(x) dx$ and $\int_{-\infty}^0 f(x) dx$ both converge, then $\int_{-\infty}^{\infty} f(x) dx = 2\pi i \Sigma R^+$, where ΣR^+ denotes the sum of the residues of $f(z)$ at its poles in the upper half plane.

Choosing a semi-circle as contour let the origin be the centre of the semi-circle and R its radius in the upper half plane. Let the semi-circle be represented by Γ and R be chosen large enough in order that the semi-circle may include all the poles of $f(z)$.




Fig. 5.27

If $f(z)$ has no poles on the boundary and we consider the closed contour C , then by residue theorem, we have

$$\int_C f(z) dz = \int_{-R}^R f(x) dx + \int_{\Gamma} f(z) dz,$$

$$\text{since along the axis of } x, z=x, \therefore f(z)=f(x).$$

Assuming $z = Re^{i\theta}$, $dz = iRe^{i\theta} d\theta = iz d\theta$ we have

$$\int_{\Gamma} f(z) dz = i \int_0^{\pi} f(z) z d\theta$$

$$\begin{aligned} &= 0 \text{ since } \left| \int_0^{\pi} f(z) z d\theta \right| \leq \int_0^{\pi} |f(z)| |z| d\theta \\ &\leq 0 \text{ as } z \rightarrow \infty, zf(z) \rightarrow 0 \\ &= 0 \text{ as modulus cannot be -ve.} \end{aligned}$$

Hence by Cauchy's Residue theorem,

$$\begin{aligned} 2\pi i \Sigma R^+ &= \int_C f(z) dz \\ &= \int_{-R}^R f(x) dx, \because \int_C f(z) dz = \int_{-R}^R f(x) dx \\ &= \int_{-\infty}^{\infty} f(x) dx, \text{ as } R \rightarrow \infty. \end{aligned}$$

$$\text{Hence } \int_{-\infty}^{\infty} f(x) dx = 2\pi i \Sigma R^+.$$

$$\text{Now } \int_{-\infty}^{\infty} f(x) dx = \int_0^{\infty} f(x) dx + \int_{-\infty}^0 f(x) dx,$$

where both the integrals on R.H.S. individually converge and hence the result.

Note 1. This theorem utilised for evaluation of the integral of the type $\int_{-\infty}^{\infty} f(x) dx$

by the method of contour integration, may be extended to the case in which $f(x)$ has simple poles on the real axis. In such cases, we can indent the contour by making small semi-circles in the upper half plane to cut out the simple poles on the real axis.




Fig. 5.28

Suppose that $f(z)$ is a quotient function of z , whose numerator and denominator are represented by $N(z)$ and $D(z)$ respectively. Then $D(z)=0$ gives the poles. Let us assume that $D(z)=0$ has only one real root say $z=a$, where a is real. Let the semi-circle, indenting this pole be denoted by γ . Its centre is the point $x=a$ and let its radius be ρ .

If Γ is sufficiently large to enclose all the poles of $f(z)$ in the upper half plane, then the integral round Γ tends to zero as $R \rightarrow \infty$. We therefore have by splitting up the bound (shown in the adjoining

diagram), i.e., $\int_C f(z) dz = 2\pi i \Sigma R^+$ as

$$\int_{-R}^{a-\rho} f(x) dx + \int_{\gamma} f(z) dz + \int_{a+\rho}^R f(x) dx + \int_{\Gamma} f(z) dz = 2\pi i \Sigma R^+,$$

Here $zf(z) \rightarrow 0$ as $z \rightarrow \infty$.

$$\therefore \int_{\Gamma} f(z) dz \rightarrow 0 \quad (\text{as shown above}).$$

Now we have to consider $\int_{\gamma} f(z) dz$.




Fig. 5.29

Let us assume that $z = a + \rho e^{i\theta}$,

where $z = a$ if $\rho \rightarrow 0$;

then $dz = \rho i e^{i\theta} d\theta = i(z-a) d\theta$.

$$\text{So} \quad \int_{\gamma} f(z) dz = i \int_{\pi}^0 f(z) (z-a) d\theta$$

$$\therefore \lim_{\rho \rightarrow 0} \int_{\gamma} f(z) dz = i \int_{\pi}^0 \lim_{z \rightarrow a} (z-a) f(z) d\theta$$

$$= i \Sigma R_0 \int_{\pi}^0 d\theta = -i\pi \Sigma R_0,$$

where R_0 is the residue of $z=a$, the simple pole, on real axis.

Hence, when $R \rightarrow \infty$, we have

$$\int_{-\infty}^{a-\rho} f(x) dx + \int_{a+\rho}^{\infty} f(x) dx = 2\pi i \Sigma R^+ + i\pi \Sigma R_0,$$

where ΣR_0 denotes the sum of residues of $f(z)$ at its simple poles on the real axis, for each pole on the real axis can be treated similarly.

The left hand side of this equation is known as Principal value for dz from $-\infty$ to ∞ and denoted by P , so that

$$P \int_{-\infty}^{\infty} f(x) dx = 2\pi i \Sigma R^+ + i\pi \Sigma R_0,$$

$$\text{where} \quad \lim_{\rho \rightarrow 0} \left[\int_{-\infty}^{a-\rho} f(x) dx + \int_{a+\rho}^{\infty} f(x) dx \right] = P \int_{-\infty}^{\infty} f(x) dx.$$

Note 2. Indented Semi-circular Contour.
When $f(z)$ has simple poles on the real axis, we then indent the contour by making small semi-circles in the upper half-plane in order to cut out the simple poles on the real axis.




Fig. 5.30

5.20. JORDAN'S LEMMA TO EVALUATE INFINITE INTEGRALS

If Γ be a semi-circle with centre as origin and radius R and let $f(z)$ be a function such that

(i) $f(z)$ is analytic in the upper half plane,

(ii) $f(z) \rightarrow 0$ uniformly as $|z| \rightarrow \infty$
for $0 \leq \arg z \leq \pi$,

(iii) m is +ve,

then $\int_{\Gamma} e^{imz} f(z) dz \rightarrow 0$ as $R \rightarrow \infty$.

Let us assume that $|f(z)| = \varepsilon$ when z is sufficiently large and $\varepsilon \rightarrow 0$ as $z \rightarrow \infty$.

Also, let $z = Re^{i\theta}$ and $dz = iRe^{i\theta} d\theta$.




Fig. 5.31

$$\therefore \left| \int_{\Gamma} f(z) e^{imz} dz \right| = \left| \int_0^{\pi} e^{imRe^{i\theta}} f(Re^{i\theta}) iRe^{i\theta} d\theta \right|$$

$$\leq \int_0^{\pi} |e^{imRe^{i\theta}}| |f(Re^{i\theta})| |i| R |e^{i\theta}| d\theta$$

$$\leq \varepsilon \int_0^{\pi} |e^{imRe^{i\theta}}| R d\theta, \text{ since } |e^{i\theta}| = 1, |i| = 1,$$

$$\text{also } |f(Re^{i\theta})| = \varepsilon.$$

Now consider that $e^{imRe^{i\theta}} = e^{imR(\cos \theta + i \sin \theta)}$.

$$\therefore |e^{imRe^{i\theta}}| = |e^{-mR \sin \theta}| |e^{imR \cos \theta}| = |e^{-mR \sin \theta}|$$

$$|e^{imR \cos \theta}| = |\cos(mR \cos \theta) + i \sin(mR \cos \theta)| = 1.$$

$$\text{Thus} \quad \left| \int_{\Gamma} e^{imz} f(z) dz \right| \leq \varepsilon \int_0^{\pi} e^{-mR \sin \theta} R d\theta$$

$$\leq \varepsilon \int_0^{\pi/2} e^{-mR \sin \theta} R d\theta$$

$$+ \int_{\pi/2}^{\pi} e^{-mR \sin \theta} R d\theta \quad \left[\text{on replacing } \phi \text{ by } \theta \right].$$

If we put $\theta = \pi - \phi$, where $\theta = \pi$, $\phi = 0$, $d\theta = -d\phi$, $\theta = \pi/2$, $\phi = \pi/2$.

$$\text{then} \quad \int_{\pi/2}^{\pi} e^{-mR \sin \theta} R d\theta = \int_0^{\pi/2} e^{-mR \sin \phi} R d\phi$$

$$= \int_0^{\pi/2} e^{-mR \sin \theta} R d\theta$$

on replacing ϕ by θ .

$$\text{Hence} \quad \left| \int_{\Gamma} e^{imz} f(z) dz \right| \leq 2\varepsilon \int_0^{\pi/2} e^{-mR \sin \theta} R d\theta.$$

Now we know by Jordan's inequality that if $0 \leq \theta \leq \pi/2$.

$\frac{\sin \theta}{\theta} > \frac{2}{\pi}$ since $\frac{\sin \theta}{\theta}$ steadily decreases from 1 to $\frac{2}{\pi}$ as θ increases from 0 to $\pi/2$.

$$\therefore \sin \theta > \frac{2}{\pi} \theta.$$

$$\text{As such} \quad e^{-mR \sin \theta} < e^{-mR (2/\pi) \theta}.$$

Therefore

$$\begin{aligned} \left| \int_{\Gamma} e^{ims} f(z) dz \right| &\leq 2\epsilon \int_0^{\pi/2} e^{-mR} (\pi/2)^{\theta} R d\theta \\ &\leq 2\epsilon \left[\frac{e^{-mR} (\pi/2)^{\theta}}{-mR (2/\pi)} \right]_0^{\pi/2} \\ &\leq \frac{\pi\epsilon}{m} [1 - e^{-mR}] \\ &\leq 0 \text{ when } z \rightarrow \infty \text{ and } \epsilon \rightarrow 0. \end{aligned}$$

Hence $\int_{\Gamma} e^{ims} f(z) dz \rightarrow 0$ as $R \rightarrow \infty$.




Fig. 5.32

Note 1. Integrals involving many-valued functions. If $z^{\alpha-1}$ be many-valued function, then an integral of the type $\int_0^{\infty} x^{\alpha-1} f(x) dx$, where α is not integer can be evaluated by contour integration by taking the integral round a circle $|z|=R$ and a small circle $|z|=r$, which encloses the branch point $z=0$ and a cross cut along the real axis from 0 to ∞ , i.e. joining the ends of the two circles.

Note 2. There is no special merit in using a particular curve as contour for a particular integration, but in fact any of circle, semi-circle, quadrant of a circle or a rectangle whichever is suitable, can be used as a contour unless otherwise stated, e.g. if the integrand is a periodic function, then rectangular contour is rather convenient.

Problem 18. Find the residue of $\frac{z^2}{z^2+a^2}$ at $z=ia$.

As given $f(z) = \frac{z^2}{z^2+a^2} = \frac{z^2}{(z-ia)(z+ia)}$.

Here $z=ia$ is a pole of order one or say a simple pole.

Residue at $z=ia$ is $\lim_{z \rightarrow ia} (z-ia) f(z)$

$$= \lim_{z \rightarrow ia} (z-ia) \cdot \frac{z^2}{(z+ia)(z-ia)}$$

$$= \lim_{z \rightarrow ia} \frac{z^2}{z+ia} = \frac{(ia)^2}{2ai} = \frac{ia}{2}.$$

Aliter 1. $f(z) = \frac{z^2}{z^2+a^2}$ is of the form $\frac{\phi(z)}{\psi(z)}$,

where $\phi(z) = z^2$, $\psi(z) = z^2+a^2$

$\therefore$ residue at $z=ia$ is $= \frac{\phi(ia)}{\psi'(ia)}$, where $\phi(ia) = (ai)^2$, $\psi'(ia) = 2ai$

$$= \frac{(ai)^2}{2ai} = \frac{ai}{2}.$$

Aliter 2. Put $z=ia+t$.

$$\begin{aligned} \therefore f(ia+t) &= \frac{(ia+t)^2}{(2ia+t)t} \\ &= \frac{(ia)^2 \left(1 + \frac{t}{ia}\right)^2}{2iat \left(1 + \frac{t}{2ia}\right)} = \frac{ia}{2t} \left(1 + \frac{t}{ia}\right)^2 \left(1 + \frac{t}{2ia}\right)^{-1} \\ &= \frac{ia}{2t} \left[1 + \frac{2t}{ia} + \left(\frac{t}{ia}\right)^2 + \dots\right] \left[1 - \frac{t}{2ia} + \left(\frac{t}{2ia}\right)^2 \dots\right] \\ &= \frac{ia}{2t} \left[1 + \left(\frac{2}{ia} - \frac{1}{2i}\right)t + \dots\right] \end{aligned}$$

Residue at $z=ia$ is = coefficient of $\frac{1}{t}$ in $f(ia+t) = \frac{ia}{2}$.

Problem 19. (a). Find the residue of $\frac{z^4}{(z-1)^4(z-2)(z-3)}$ at $z=1$.

(b) Find the residues of $\frac{z}{(z-a)(z-b)}$ and $\frac{z^3-z^2+1}{z^3}$ at infinity.

(a) Here $f(z) = \frac{z^4}{(z-1)^4(z-2)(z-3)}$.

To find residue at $z=1$, which is a pole of order 4.

$$f(z) = \frac{1}{(z-1)^4} \cdot \frac{z^4}{(z-2)(z-3)} = \frac{\phi(z)}{(z-1)^4},$$

where $\phi(z) = \frac{z^4}{(z-2)(z-3)} = z^2 + 5z + 19 - \frac{16}{z-2} + \frac{81}{z-3}$

(breaking into partial fractions).

$$\therefore \phi'(z) = 2z + 5 + \frac{16}{(z-2)^2} - \frac{81}{(z-3)^2},$$

$$\phi''(z) = 2 - \frac{32}{(z-2)^3} + \frac{162}{(z-3)^3},$$

$$\therefore \phi'''(z) = \frac{96}{(z-2)^4} - \frac{486}{(z-3)^4}.$$

$$\text{Thus } \phi'''(1) = \frac{96}{(-1)^4} - \frac{486}{(-2)^4} = 96 - \frac{243}{8} = \frac{525}{8}.$$

The residue at $z=1$ is $= \frac{\phi'''(1)}{3!} = \frac{175}{16}.$

(b) Both the functions are analytic at infinity.

$$(i) \quad f(z) = \frac{z}{(z-a)(z-b)}$$

Residue of $f(z)$ at $z=0 = \lim_{z \rightarrow \infty} \{-zf(z)\}$

$$\begin{aligned} &= \lim_{z \rightarrow \infty} \left\{ -\frac{z^2}{(z-a)(z-b)} \right\} \\ &= \lim_{z \rightarrow \infty} \left\{ -\frac{1}{\left(1-\frac{a}{z}\right)\left(1-\frac{b}{z}\right)} \right\} \\ &= -1. \end{aligned}$$

$$(ii) \quad f(z) = \frac{z^3 - z^2 + 1}{z^3}.$$

Here $\lim_{z \rightarrow \infty} \{-zf(z)\}$ does not exist.

We can write,

$$f(z) = 1 - \frac{1}{z} + \frac{1}{z^3}.$$

Residue of $f(z)$ at $z = \infty$

= the negative of the coefficient of $\frac{1}{z}$ in the expansion of $f(z)$ for values of z in the neighbourhood of $z = \infty$
 $= -(-1) = 1.$

Problem 20. Apply Calculus of residues to prove that

$$(i) \quad \int_0^{2\pi} \frac{d\theta}{2 + \cos \theta} = \frac{2\pi}{\sqrt{3}}. \quad [\text{Vikram, 1969}]$$

$$(ii) \quad \int_0^{2\pi} \frac{\sin^2 \theta d\theta}{a + b \cos \theta} = \frac{2\pi}{b^2} \{a - \sqrt{a^2 - b^2}\}, \text{ if } a > b > 0.$$

$$(iii) \quad \int_0^{2\pi} e^{i n \theta} \cos(\sin \theta - n\theta) d\theta = \frac{2\pi}{|n|} \text{ where } n \text{ is a } +\text{ve integer.}$$

$$(iv) \quad \int_0^{\pi} \frac{a d\theta}{a^2 + \sin^2 \theta} = \frac{\pi}{\sqrt{1+a^2}}, \quad a > 0.$$

$$(v) \quad \int_0^{2\pi} \frac{\cos^2 p\theta d\theta}{1 - 2p \cos 2\theta + p^2} = \pi \cdot \frac{1-p+p^2}{1-p}, \quad (0 < p < 1).$$

$$(vi) \quad \int_0^{2\pi} \frac{d\theta}{(5-4 \cos \theta)^2} = \frac{40}{27} \pi, \text{ and (vii) } \int_0^{2\pi} \frac{\cos 2\theta}{5+4 \cos \theta} d\theta = \frac{\pi}{6}$$

$$(i) \text{ Suppose } I = \int_0^{2\pi} \frac{d\theta}{2 + \cos \theta}. \quad (\text{Agra, 1964})$$

Choosing the contour as a circle of unit radius ($|z|=1$),

$$z = e^{i\theta}; \therefore dz = ie^{i\theta} d\theta = iz d\theta,$$

so that $d\theta = \frac{dz}{iz}$, where θ varies from 0 to 2π .

Also when $z = e^{i\theta} = \cos \theta + i \sin \theta$.

$$\frac{1}{z} = \cos \theta - i \sin \theta.$$




Fig. 5.33

$$\text{Adding, } z + \frac{1}{z} = 2 \cos \theta, \text{ i.e. } \cos \theta = \frac{1}{2} \left(z + \frac{1}{z} \right).$$

$$\text{Thus, } I = \int_C \frac{dz}{iz} \left\{ 2 + \frac{1}{2} \left(z + \frac{1}{z} \right) \right\} = \frac{2}{i} \int_C \frac{dz}{z^2 + 4z + 1},$$

where C denotes the unit circle $|z|=1$.

Poles of the integrand will be given by $z^2 + 4z + 1 = 0$

$$\text{or } z = \frac{-4 \pm \sqrt{(16-4)}}{2} = -2 \pm \sqrt{3},$$

of which the pole $z = -2 - \sqrt{3}$ lies outside the contour and therefore the only pole that lies within the contour is $z = -2 + \sqrt{3}$ (which is of the order one).

$$\therefore \text{ residue at } z = -2 + \sqrt{3} \text{ is } \lim_{z \rightarrow -2 + \sqrt{3}} \{z - (-2 + \sqrt{3})\} f(z)$$

$$\text{where } f(z) = \frac{2}{i(z^2 + 4z + 1)}$$

$$= \frac{2}{i} \lim_{z \rightarrow -2 + \sqrt{3}} (z + 2 - \sqrt{3}) \cdot \frac{1}{(z^2 + 4z + 1)}$$

$$= \frac{2}{i} \lim_{z \rightarrow -2 + \sqrt{3}} \frac{z + 2 - \sqrt{3}}{(z + 2 - \sqrt{3})(z + 2 + \sqrt{3})}$$

$$= \frac{2}{i} \lim_{z \rightarrow -2 + \sqrt{3}} \left(\frac{1}{z + 2 + \sqrt{3}} \right)$$

$$= \frac{2}{i} \cdot \frac{1}{-2 + \sqrt{3} + 2 + \sqrt{3}} = \frac{1}{i\sqrt{3}}.$$

Hence by Cauchy's residue theorem we have

$$I = 2\pi i \Sigma R, \text{ where } \Sigma R \text{ represents sum of the residues inside the contour}$$

$$= 2\pi i \times \frac{1}{i\sqrt{3}} = \frac{2\pi}{\sqrt{3}}.$$

$$(ii) \text{ Suppose } I = \int_0^{2\pi} \frac{\sin^2 \theta d\theta}{a + b \cos \theta}$$

Choosing the contour C as a circle ($|z|=1$) of unit radius,

$$z = e^{i\theta}, \text{ so that } z + \frac{1}{z} = 2 \cos \theta.$$

$$dz = iz d\theta \quad z - \frac{1}{z} = 2i \sin \theta.$$

$$\begin{aligned} \therefore I &= \frac{1}{i} \int_C \left\{ \frac{1}{2i} \left(z - \frac{1}{z} \right) \right\} \frac{dz}{z^2 \left\{ 2a + b \left(z + \frac{1}{z} \right) \right\}} = -\frac{1}{i} \int_C \frac{\left(z - \frac{1}{z} \right)^2}{2z \left\{ 2a + b \left(z + \frac{1}{z} \right) \right\}} dz \\ &= -\frac{1}{2ib} \int_C \frac{(z^2 - 1)^2 dz}{z^2 (2az + bz^2 + b)} = -\frac{1}{2ib} \int_C \frac{(z^2 - 1)^2 dz}{z^2 \left(z^2 + 1 + \frac{2a}{b} z \right)} \\ &= -\frac{1}{2ib} \int_C \frac{(z^2 - 1)^2 dz}{z^2 (z - \alpha)(z - \beta)}, \end{aligned}$$

where $\alpha = \frac{-a + \sqrt{a^2 - b^2}}{b}$ and $\beta = \frac{-a - \sqrt{a^2 - b^2}}{b}$ are the roots of the quadratic $z^2 + \frac{2a}{b}z + 1 = 0$.

$$\text{So } \alpha + \beta = -\frac{2a}{b}, \text{ and } \alpha\beta = 1.$$

As such $|\alpha| |\beta| = 1$, where $|\beta| > |\alpha|$

$\therefore |\beta| = \frac{1}{|\alpha|} > 1$, i.e., $|\alpha| < 1$ and so $z = \alpha$ is the only simple pole within the contour C . $z = \beta$ lies outside the contour. Also $z = 0$ is a pole of order two, which lies inside the contour.

$$\begin{aligned} \text{Residue at } z = \alpha & \text{ is } \lim_{z \rightarrow \alpha} (z - \alpha) \cdot \left(-\frac{1}{2ib} \right) \frac{(z^2 - 1)^2 dz}{z^2 (z - \alpha)(z - \beta)} \\ &= -\frac{1}{2ib} \lim_{z \rightarrow \alpha} \frac{(z^2 - 1)^2}{z^2 (z - \beta)} = -\frac{1}{2ib} \lim_{z \rightarrow \alpha} \frac{\left(z - \frac{1}{z} \right)^2}{z - \beta} \\ &= -\frac{1}{2ib} \frac{\left(\alpha - \frac{1}{\alpha} \right)^2}{\alpha - \beta} = -\frac{(\alpha - \beta)^2}{2ib(\alpha - \beta)} \text{ as } \alpha\beta = 1, \therefore \beta = \frac{1}{\alpha} \\ &= -\frac{\alpha - \beta}{2ib}, \text{ where } (\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta \\ &= \frac{4a^2}{b^2} - 4 = \frac{4}{b^2} (a^2 - b^2) \\ &= \frac{1}{-2ib} \cdot \frac{2}{b} \sqrt{a^2 - b^2} \\ &= \frac{\sqrt{a^2 - b^2}}{-ib^2}. \end{aligned}$$

Again residue at $z = 0$ (a pole of order two) is the coefficient of $\frac{1}{z}$

$$\text{in } -\frac{1}{2ib} \frac{(z^2 - 1)^2}{z^2 \left(z^2 + 1 + \frac{2a}{b} z \right)}, \text{ where } z \text{ is a small quantity near the pole,}$$

$$\text{i.e. in } -\frac{1}{2ib} \frac{\left(1 - \frac{1}{z^2} \right)^2}{\left(1 + \frac{2a}{b} \frac{1}{z} + \frac{1}{z^2} \right)} = -\frac{1}{2ib} \left(1 - \frac{1}{z^2} \right)^2 \left(1 + \frac{2a}{bz} + \frac{1}{z^2} \right)^{-1},$$

$$\text{i.e. in } -\frac{1}{2ib} \left(1 - \frac{2}{z^2} + \dots \right) \left[1 - \left(\frac{2a}{bz} + \frac{1}{z^2} \right) + \dots \right].$$

$$\therefore \text{residue at } z = 0 \text{ is } -\frac{1}{2ib} \cdot \left(-\frac{2a}{b} \right) = \frac{a}{ib^2}.$$

Hence by Cauchy's Residue theorem,

$$I = 2\pi i \Sigma R$$

$$= 2\pi i \left[-\frac{\sqrt{a^2 - b^2}}{ib^2} + \frac{a}{ib^2} \right] = \frac{2\pi}{b^2} [a - \sqrt{a^2 - b^2}].$$

$$(iii) \text{ Suppose } I = \int_0^{2\pi} e^{i\cos\theta} \cos(\sin\theta - n\theta) d\theta.$$

Choosing the contour C as a circle of unit radius, $|z| = 1$.

$$z = e^{i\theta}, \text{ so that } \frac{1}{z} = e^{-i\theta},$$

$$dz = iz d\theta.$$

Now, consider,

$$\begin{aligned} J &= \int_0^{2\pi} e^{i\cos\theta} [\cos(\sin\theta - n\theta) + i \sin(\sin\theta - n\theta)] d\theta \\ &= \int_0^{2\pi} e^{i\cos\theta} \cdot e^{i(\sin\theta - n\theta)} d\theta \end{aligned}$$

$$\text{or } J = \int_0^{2\pi} e^{i\cos\theta + i\sin\theta} \cdot e^{-in\theta} d\theta = \int_0^{2\pi} e^{e^{i\theta}} \cdot (e^{-i\theta})^n d\theta,$$

where I is the real part in J .

On putting the values of $e^{i\theta} = z$ and $e^{-i\theta} = \frac{1}{z}$ etc., we have

$$J = \int_C e^z \frac{1}{z^n} \frac{dz}{iz} = \frac{1}{i} \int_C \frac{e^z}{z^{n+1}} dz.$$

Here the only pole inside the contour C is $z = 0$ of order $(n+1)$.

$$\text{Residue at } z = 0 \text{ a pole of order } (n+1) \text{ is } = \frac{\phi^n(0)}{n!} = \frac{1}{i(n)!},$$

$$\text{where } \phi(z) = \frac{1}{i} e^z, \therefore \phi^n(z) = \frac{e^z}{i}, \phi^n(0) = \frac{1}{i}.$$

Hence by Cauchy's Residue theorem,

$J = 2\pi i \Sigma R$, where ΣR is the sum of residues inside C

$$= \pi i \cdot \frac{1}{i \left| \frac{n}{n} \right|} = \frac{2\pi}{n}$$

Therefore real part in J , i.e.

$$I = \int_0^{2\pi} e^{\cos \theta} \cos(\sin \theta - n\theta) d\theta = \frac{2\pi}{n}$$

(iv) Suppose $I = \int_0^\pi \frac{a d\theta}{a^2 + \sin^2 \theta}$.




Fig. 5.34

Choosing the contour C as a semi-circle of unit radius.

$$z = e^{i\theta}, \therefore dz = iz d\theta$$

or $d\theta = \frac{dz}{iz}$ and $\sin \theta = \frac{1}{2i} \left(z - \frac{1}{z} \right)$,

where θ varies from 0 to π .

$$\begin{aligned} \therefore I &= \frac{a}{i} \int_C \frac{1}{a^2 + \left\{ \frac{1}{2i} \left(z - \frac{1}{z} \right) \right\}^2} \frac{dz}{iz} = \frac{4a}{i} \int_C \frac{dz}{z \left\{ 4a^2 - z^2 - \frac{1}{z^2} + 2 \right\}} \\ &= \frac{4a}{i} \int_C \frac{z dz}{z^2 (4a^2 + 2) - z^4 - 1} = -\frac{4a}{i} \int_C \frac{z dz}{-z^2 (4a^2 + 2) + z^4 + 1} \end{aligned}$$

Poles are given by $-z^2 (4a^2 + 2) + z^4 + 1 = 0$.

Let us assume that

$$z^4 - z^2 (4a^2 + 2) + 1 \equiv (z^2 - \alpha^2) (z^2 - \beta^2).$$

$$\therefore \alpha^2 + \beta^2 = 4a^2 + 2 \text{ and } \alpha^2 \beta^2 = 1.$$

If $\alpha < 1$, then $\beta > 1$, implies that the pole within the semi-circle will be $z = \alpha$, which is the only pole inside C and $z = \beta$ will lie outside the contour.

Residue at $z = \alpha$ is

$$\begin{aligned} &= \lim_{z \rightarrow \alpha} (z - \alpha) \left(-\frac{4a}{i} \right) \cdot \frac{z}{(z^2 - \alpha^2) (z^2 - \beta^2)} \\ &= -\frac{4a}{i} \lim_{z \rightarrow \alpha} \frac{z}{(z + \alpha) (z^2 - \beta^2)} \\ &= -\frac{4a}{i} \frac{\alpha}{2\alpha (\alpha^2 - \beta^2)} = -\frac{2a}{i (\alpha^2 - \beta^2)} \end{aligned}$$

Hence by Cauchy's Residue theorem,

$$\begin{aligned} I &= 2\pi i \Sigma R = 2\pi i \times \frac{2a}{-i (\alpha^2 - \beta^2)} \\ &= \frac{4\pi a}{\beta^2 - \alpha^2}, \text{ where } \beta^2 - \alpha^2 = \sqrt{[(\beta^2 + \alpha^2)^2 - 4\alpha^2 \beta^2]} \\ &= \sqrt{[(4a^2 + 2)^2 - 4]} = 4a \sqrt{1 + a^2} \\ &= \frac{4\pi a}{4a \sqrt{1 + a^2}} = \frac{\pi}{\sqrt{1 + a^2}} \end{aligned}$$

(i) Suppose $I = \int_0^{2\pi} \frac{\cos^2 3\theta d\theta}{1 - 2p \cos 2\theta + p^2}$.

Choosing the contour C as a circle of unit radius,

$$z = e^{i\theta}; \therefore dz = ie^{i\theta} d\theta = iz d\theta,$$

so that $d\theta = \frac{dz}{iz}$.

Also $z^n = e^{in\theta}; \therefore \frac{1}{z^n} = e^{-in\theta}$.

As such, $2 \cos 3\theta = z^3 + \frac{1}{z^3}$ and $2 \cos 2\theta = z^2 + \frac{1}{z^2}$.

$$I = \frac{1}{i} \int_C \frac{\frac{dz}{z} \left\{ \frac{1}{2} \left(z^3 + \frac{1}{z^3} \right) \right\}^2}{1 - p \left(z^2 + \frac{1}{z^2} \right) + p^2} = \frac{1}{i} \int_C \frac{(z^6 + 1)^2 z^2}{4z^6 \{ (1 + p^2) z^2 - pz^4 - p \}} \frac{dz}{z}$$

$$= -\frac{1}{4i} \int_C \frac{(z^6 + 1)^2 dz}{z^5 [pz^4 - (1 + p^2) z^2 + p]} = -\frac{1}{4i} \int_C \frac{(z^6 + 1)^2 dz}{z^5 (pz^2 - 1) (z^2 - p)}.$$

Poles are $z = 0$, a pole of order five, and

$$z = \pm \sqrt{p}, \pm \frac{1}{\sqrt{p}}.$$

If $p > 1$, then $\frac{1}{p} < 1$; so the only poles within the contour are

$z = \pm \sqrt{p}$ and also $z = 0$, a pole of order five.

Residue at $z = 0$ (pole of order 5) is the coefficient of $1/z$ in

$$\begin{aligned} &\frac{(z^6 + 1)^2}{z^5 (pz^2 - 1) (z^2 - p)} \\ &= \text{coeff. of } \frac{1}{z} \text{ in } \frac{(1 + 2z^6 + z^{12})}{-4iz^5(1 - p^2)} \left[\frac{p}{pz^2 - 1} - \frac{1}{z^2 - p} \right] \\ &= \text{,, ,, } \frac{(1 + 2z^6 + z^{12})}{-4iz^5(1 - p^2)} \left[\frac{1}{p \left(1 - \frac{z^2}{p} \right)} - \frac{p}{1 - pz^2} \right] \\ &= \text{coeff. of } \frac{1}{z} \text{ in } \frac{1}{-4i(1 - p^2)} \cdot \frac{1 + 2z^6 + z^{12}}{z^5} \\ &\quad \times \left[\frac{1}{p} \left(1 - \frac{z^2}{p} \right)^{-1} - p(1 - pz^2)^{-1} \right] \\ &= \text{,, ,, } \frac{1}{-4i(1 - p^2)} \cdot \frac{1 + 2z^6 + z^{12}}{z^5} \\ &\quad \times \left[\frac{1}{p} \left(1 + \frac{z^2}{p} + \frac{z^4}{p^2} + \dots \right) - p(1 + pz^2 + p^2z^4 + \dots) \right] \\ &= \frac{1}{-4i(1 - p^2)} \left(\frac{1}{p^3} - p^3 \right) = \frac{1 - p^6}{-4ip^3(1 - p^2)}. \end{aligned}$$

Residue at $z=\sqrt{p}$ is

$$\begin{aligned} &= \frac{-1}{4i} \lim_{z \rightarrow \sqrt{p}} (z-\sqrt{p}) \frac{(z^6+1)^2}{z^5(pz^2-1)(z^2-p)} \\ &= -\frac{1}{4i} \lim_{z \rightarrow \sqrt{p}} \frac{(z^6+1)^2}{z^5(pz^2-1)(z+\sqrt{p})} \\ &= -\frac{1}{4i} \frac{(p^3+1)^2}{p^{5/2}(p^2-1) \cdot 2\sqrt{p}} = \frac{(p^3+1)^2}{8ip^3(1-p^2)} \end{aligned}$$

Similarly, Residue at $z=-\sqrt{p}$ is $-\frac{(p^3+1)^2}{8ip^3(1-p^2)}$.

Hence by Cauchy's Residue theorem,

$$\begin{aligned} I &= 2\pi i \Sigma R = \frac{2\pi i}{4ip^3(1-p^2)} \left[-(1-p^6) + \frac{(p^3+1)^2}{2} + \frac{(p^3+1)^2}{2} \right] \\ &= \frac{\pi}{2p^3(1-p^2)} [-1+p^6+p^6+2p^3+1] \\ &= \frac{\pi}{2p^3(1-p^2)} \cdot 2p^3(1+p^3) = \pi \cdot \frac{(1+p)(1-p+p^2)}{(1-p)(1+p)} \\ &= \pi \frac{(1-p+p^2)}{1-p} \end{aligned}$$

(vi) Suppose $I = \int_0^{2\pi} \frac{d\theta}{(5-4\cos\theta)^2}$.

Choosing the contour C as a circle of unit radius ($|z|=1$).

$$z=e^{i\theta}, dz=iz dz \text{ or } d\theta = \frac{dz}{iz}$$

Also $\frac{1}{z} = e^{-i\theta}$; $\therefore z + \frac{1}{z} = 2\cos\theta$.

$$\begin{aligned} \text{Thus, } I &= \frac{1}{i} \int_C \frac{dz}{z \left\{ 5 - 2 \left(z + \frac{1}{z} \right) \right\}^2} = \frac{1}{i} \int_C \frac{z dz}{(2z^2 - 5z + 2)^2} \\ &= \frac{1}{i} \int_C \frac{z dz}{(z-2)^2 (2z-1)^2} \end{aligned}$$

Poles are $z=2$, a pole of order 2 and $z=\frac{1}{2}$, also a pole of order 2.

But the only pole that lies inside the contour is $z=\frac{1}{2}$ of order 2.

Residue at $z=\frac{1}{2}$ (pole of order 2) is $= \frac{1}{i} \frac{\phi'(\frac{1}{2})}{1} = \frac{20}{27i}$,

where $\phi(z) = \frac{z}{(z-2)^2}$, $\phi'(z) = \frac{(z-2)^2 - 2z(z-2)}{(z-2)^4}$, $\phi'(\frac{1}{2}) = \frac{\frac{1}{4} + \frac{1}{2}}{\frac{81}{16}} = \frac{20}{27}$

Hence by Cauchy's Residue theorem,

$$\begin{aligned} I &= 2\pi i \Sigma R \\ &= 2\pi i \times \frac{20}{27i} = \frac{40}{27} \pi. \end{aligned}$$

Note. The similar procedure will show that

$$\int_0^{2\pi} \frac{d\theta}{(5+4\cos\theta)^2} = \frac{10\pi}{7}$$

and in general $\int_0^{2\pi} \frac{d\theta}{(a+b\cos\theta)^2} = \frac{2\pi a}{(a^2-b^2)^{3/2}}$, $a>b>0$.

(vii) Suppose $I = \int_0^{2\pi} \frac{\cos 2\theta}{5+4\cos\theta} d\theta$

Choosing the contour C as a circle of unit radius i.e. $|z|=1$,

we put, $z=e^{i\theta}$, $dz=iz dz$ or $d\theta = \frac{dz}{iz}$

$$\begin{aligned} \text{Then } I &= \int_0^{2\pi} \frac{2\cos^2\theta - 1}{5+4\cos\theta} d\theta = \int_C \frac{2(z+1/z) - 1}{5+2(z+1/z)} \frac{dz}{iz} = \frac{1}{2i} \int_C \frac{z^2+1}{z^2(2z+1)(z+2)} dz \end{aligned}$$

Within the contour C there are a simple pole at $z=-\frac{1}{2}$ and a pole of order 2 at $z=0$.

Residue at $z=-\frac{1}{2}$, $= \lim_{z \rightarrow -\frac{1}{2}} (z+\frac{1}{2}) \frac{1}{2i} \frac{z^2+1}{z^2(2z+1)(z+2)} = \frac{17}{24}i$

and Residue at $z=0$ (order 2) = coefficient of $\frac{1}{z}$ in $\frac{1}{2i} \frac{z^2+1}{z^2(2z+1)(z+2)}$, z , being small

$$\begin{aligned} &= \text{coeff. of } \frac{1}{z} \text{ in } \frac{1}{4i} \left(1 + \frac{1}{z^4} \right) \left(1 + \frac{1}{2z} \right)^{-1} \left(1 + \frac{2}{z} \right)^{-1} \\ &= \text{,, ,, } \frac{1}{4i} \left(1 + \frac{1}{z^4} \right) \left(1 - \frac{1}{2z} + \dots \right) \left(1 - \frac{2}{z} + \dots \right) \\ &= \frac{1}{4i} \left(-\frac{5}{2} \right) = -\frac{5}{8i}. \end{aligned}$$

Hence by Cauchy's theorem,

$$\begin{aligned} I &= 2\pi i \times \text{sum of residues} = 2\pi \Sigma R \\ &= 2\pi i \left[\frac{17}{24i} - \frac{5}{8i} \right] = \frac{\pi}{6}. \end{aligned}$$

Problem 21. Prove by contour integration that

(i) $\int_0^\infty \frac{dx}{(a+bx^2)^n} = \frac{\pi}{2^n \sqrt{b}} \frac{1.3.5 \dots (2n-3)}{1.2.3 \dots (n-1)} \frac{1}{a^{\frac{1}{2}(2n-1)}}$

(ii) $\int_0^\infty \frac{x \sin ax}{x^2+k^2} dx = \frac{1}{2} \pi e^{-ak}$ where $a>0$;

(iii) $\int_0^\infty \frac{x^6 dx}{(a^4+x^4)^2} = \frac{3\sqrt{2}\pi}{16a}$ where $a>0$; (iv) $\int_0^\infty \frac{dx}{1+x^2} = \frac{\pi}{2}$

$$(v) \int_0^{\infty} \frac{\cos x \, dx}{(x^2+a^2)(x^2+b^2)} = \frac{\pi}{a^2-b^2} \left(\frac{e^{-b}}{b} - \frac{e^{-a}}{a} \right) \text{ where } a > b > 0 \text{ and}$$

$$(vi) \int_0^{\infty} \frac{\log(1+x^2)}{1+x^2} dx = \pi \log 2. \quad (\text{Rohilkhand, 1978})$$




Fig. 5.35

$$\int_C f(z) \, dz = \int_{-R}^R f(x) \, dx + \int_{\Gamma} f(z) \, dz$$

and since $\int_C f(z) \, dz = 2\pi i \Sigma R^+$ where ΣR^+ is the sum of residues,

$$\therefore \int_{-R}^R f(x) \, dx + \int_{\Gamma} f(z) \, dz = 2\pi i \Sigma R^+. \quad \dots (1)$$

Now poles of $f(z)$ are given by

$$(a+bz^2)^n = 0 \text{ or } b^n \left(z+i\sqrt{\frac{a}{b}} \right)^n \left(z-i\sqrt{\frac{a}{b}} \right)^n = 0.$$

i.e. $z = \pm i\sqrt{\frac{a}{b}}$ (poles of order n).

Only the pole $z = i\sqrt{\frac{a}{b}}$ of order n lies within the contour.

Residue at $z = i\sqrt{\frac{a}{b}}$ (a pole of order n) is $= \frac{\phi^{n-1} \left(i\sqrt{\frac{a}{b}} \right)}{(n-1)!}$

where $\phi(z) = \frac{1}{b^n \left(z+i\sqrt{\frac{a}{b}} \right)^n}$

$$\phi^{n-1}(z) = \frac{(-1)^{n-1} \cdot n(n+1)(n+2)\dots(2n-2)}{b^n \left(z+i\sqrt{\frac{a}{b}} \right)^{2n-1}}$$

$$\therefore \phi^{n-1} \left(i\sqrt{\frac{a}{b}} \right) = \frac{(-1)^{n-1} \cdot n(n+1)(n+2)\dots(2n-2)}{b^n \left(2i\sqrt{\frac{a}{b}} \right)^{2n-1}}$$

Thus residue at $z = i\sqrt{\frac{a}{b}}$ (pole of order n) is

$$\frac{(-1)^{n-1}}{(n-1)!} \cdot \frac{1}{b^n} \cdot \frac{n(n+1)(n+2)\dots(2n-2)}{\left(2i\sqrt{\frac{a}{b}} \right)^{2n-1}}$$

Now $\left| \int_{\Gamma} f(z) \, dz \right| = \left| \int_{\Gamma} \frac{dz}{(a+bz^2)^n} \right| \leq \int_{\Gamma} \frac{|dz|}{|a+bz^2|^n}$
which tends to zero as $z \rightarrow \infty$.

Hence when $R \rightarrow \infty$, the relation (1) yields

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{dx}{(a+bx^2)^n} &= 2\pi i \times \frac{(-1)^{n-1}}{(n-1)!} \cdot \frac{n(n+1)(n+2)\dots(2n-2)}{b^n \left(2i\sqrt{\frac{a}{b}} \right)^{2n-1}} \\ &= 2\pi i \times \frac{(-1)^{n-1} (2n-2)!}{(n-1)! \cdot (n-1)! \cdot \sqrt{b} (2i\sqrt{a})^{2n-1}} \\ &\quad [\text{multiplying } N^{\circ} \text{ and } D^{\circ} \text{ by } (n-1)!] \\ &= \frac{2\pi i \times (-1)^{n-1} 1.3.5\dots(2n-3) \times 2.4.6\dots(2n-2)}{(n-1)! \cdot (n-1)! \cdot \sqrt{b} (2i\sqrt{a})^{2n-1}} \\ \int_{-\infty}^{\infty} \frac{dx}{(a+bx^2)^n} &= \frac{2\pi}{2^{2n-1}} \cdot \frac{1.3.5\dots(2n-3) \cdot 2^{n-1} (n-1)!}{(n-1)! \cdot \sqrt{b} (a)^{n-1/2}} \\ \therefore \int_{-\infty}^{\infty} \frac{dx}{(a+bx^2)^n} &= \frac{\pi}{2^n \sqrt{b}} \cdot \frac{1.3.5\dots(2n-3)}{(n-1)!} \cdot \frac{1}{a^{n-1/2}} \end{aligned}$$

(ii) Consider $f(z) = \frac{ze^{iaz}}{z^2+k^2}$, where $\frac{z \sin az}{z^2+k^2}$ is the imaginary part of $f(z)$.

Choosing the closed contour C , which consists of real axis from $-R$ to R and upper half of a large circle $|z|=R$, represented by Γ , we have

$$\int_C f(z) \, dz = \int_{-R}^R f(x) \, dx + \int_{\Gamma} f(z) \, dz = 2\pi i \Sigma R^+. \quad \dots (1)$$

Singularities of $f(z)$ are given by $z^2+k^2=0$, i.e. $z = \pm ik$ of which only the simple pole $z = +ik$ lies within the contour.

Residue at $z = ik$ (a simple pole) $= \lim_{z \rightarrow ik} (z-ik) \cdot \frac{ze^{iaz}}{(z-ik)(z+ik)}$
 $= \frac{ike^{-ak}}{2ik} = \frac{1}{2}e^{-ak}$

Now $\left| \int_{\Gamma} f(z) \, dz \right| = \left| \int_{\Gamma} \frac{ze^{iaz}}{z^2+k^2} \, dz \right| \leq \int_{\Gamma} \frac{|z| |e^{iaz}| |dz|}{|z^2+k^2|}$
 ≤ 0 as $z \rightarrow \infty$.

$\therefore \int_{\Gamma} f(z) \, dz = 0$, since modulus cannot be negative.

Hence (1) yields when $R \rightarrow \infty$

$$\int_{-\infty}^{\infty} \frac{xe^{iaz}}{x^2+k^2} = 2\pi i \times \frac{e^{-ak}}{2} = i\pi e^{-ka}$$

or

$$2 \int_0^{\infty} \frac{x e^{iax}}{x^2 + k^2} dx = i\pi e^{-ak}.$$

Equating the imaginary parts on either side, we get

$$\int_0^{\infty} \frac{x \sin ax}{x^2 + k^2} dx = \frac{\pi}{2} e^{-ak}.$$

$$(iii) \text{ Here } f(z) = \frac{z^6}{(a^4 + z^4)^2}.$$

$zf(z) \rightarrow 0$ as $z \rightarrow \infty$, which satisfies the condition for evaluation of an infinite integral.

Poles of $f(z)$ are given by

$$z^4 = -a^4 = e^{i(2n+1)\pi} a^4$$

$$\therefore e^{i(2n+1)\pi} = \cos(2n+1)\pi + i \sin(2n+1)\pi = \cos \pi = -1,$$

$$\therefore z = ae^{i(2n+1)\pi/4}, \text{ where } n = 0, 1, 2, 3.$$

Thus the poles are $ae^{i\pi/4}$, $ae^{3\pi i/4}$, $ae^{5\pi i/4}$, $ae^{7\pi i/4}$ of which the first two only lie in the upper half plane (coefficient of i being +ve) which is to be chosen as a contour consisting of large semi-circle Γ , along with the real axis from $-R$ to R .

Applying the theorem of residues,

$$\int_{-R}^R f(x) dx + \int_{\Gamma} f(z) dz = 2\pi i \Sigma R^+.$$

where Γ represents the upper half plane.

$$\text{Since } \int_{\Gamma} f(z) dz \rightarrow 0 \text{ as } z \rightarrow \infty.$$

Hence when $R \rightarrow \infty$, we have

$$\int_{-\infty}^{\infty} f(x) dx = 2\pi i \Sigma R^+. \quad \dots (1)$$

Now to find the residue at $z = ae^{i\pi/4} = z_1$ (say), let us put $z = ae^{i\pi/4} + t$, t being small, in the value of $f(z)$, whence we get

$$\begin{aligned} \frac{(z_1 + t)^6}{[a^4 + (z_1 + t)^4]^2} &= \frac{z_1^6 + 6z_1^5 t + \dots}{[a^4 + z_1^4 + 4z_1^3 t + 6z_1^2 t^2 + 4z_1 t^3 + t^4]^2}, \text{ where } z_1^4 = -a^4 \\ &= \frac{z_1^6 + 6z_1^5 t + \dots}{16z_1^6 t^2 \left[1 + \frac{6t}{4z_1} + \dots \right]^2} \\ &= \frac{(z_1^6 + 6z_1^5 t + \dots)}{16z_1^6 t^2} \left[2 + \frac{6t}{4z_1} \dots \right]^2 \\ &= \frac{(z_1^6 + 6z_1^5 t + \dots)}{16z_1^6 t^2} \left[1 - \frac{12t}{4z_1} \dots \right]. \end{aligned}$$

Residue at $z = z_1$ is = coefficient of $\frac{1}{t}$ in $f(t + z_1)$

$$= \frac{6}{16z_1} - \frac{3}{16z_1} = \frac{3}{16z_1}.$$

Similarly, residue at $z = ae^{3\pi i/4} = z_2$ (say) is $\frac{3}{16z_2}$.

Hence by (1),

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{x^6 dx}{(a^4 + x^4)^2} &= 2\pi i \cdot \frac{3}{16} \left(\frac{1}{z_1} + \frac{1}{z_2} \right) \\ &= \frac{3\pi i}{8a} [e^{-\pi i/4} + e^{-3\pi i/4}] \\ &= \frac{3\pi i}{8a} [e^{-\pi i/4} - e^{\pi i/4}], \end{aligned}$$

$$\therefore e^{-3\pi i/4} = e^{-(\pi - \pi/4)} = e^{-\pi i} e^{\pi i/4} = -e^{\pi i/4}$$

$$\text{or } 2 \int_0^{\infty} \frac{x^6 dx}{(a^4 + x^4)^2} = -\frac{3\pi i}{8} \times 2i \sin \frac{\pi}{4} = \frac{3\pi}{4a} \cdot \frac{1}{\sqrt{2}}$$

$$\therefore \int_0^{\infty} \frac{x^6 dx}{(a^4 + x^4)^2} = \frac{3\sqrt{2}\pi}{16a}$$

$$(iv) \text{ Here } f(z) = \frac{1}{1+z^2} \text{ so that}$$

$$zf(z) \rightarrow 0 \text{ as } z \rightarrow \infty.$$

Poles of $f(z)$ are given by $1+z^2=0$ i.e. $z = \pm i$ of which $z = i$ lies in the upper half plane i.e. in the contour chosen as consisting of a large semi-circle Γ along with the real axis from $-R$ to R .




Fig. 5.36

$$\text{Residue of } f(z) \text{ at } z = i \text{ is } \lim_{z \rightarrow i} (z-i) \frac{1}{1+z^2} = \lim_{z \rightarrow i} \frac{1}{z+i} = \frac{1}{2i}.$$

By Cauchy's residue theorem, we have

$$\int_{-R}^R f(x) dx + \int_{\Gamma} f(z) dz = 2\pi i \Sigma R^+ = 2\pi i \times \frac{1}{2i} = \pi \quad \dots (1)$$

$$\begin{aligned} \text{where } \left| \int_{\Gamma} f(z) dz \right| &= \left| \int_{\Gamma} \frac{dz}{1+z^2} \right| \leq \int_{\Gamma} \frac{|dz|}{|1+z^2|} \leq \int_{\Gamma} \frac{|dz|}{|z|^2 - 1} \\ &\text{since } |z^2 + 1| > |z|^2 - 1 \\ &\leq \int_0^{\pi} \frac{R d\theta}{R^2 - 1} \text{ when } z = Re^{i\theta} \text{ and } |dz| = |i Re^{i\theta} d\theta| = R d\theta \\ &\leq \frac{\pi R}{R^2 - 1} \rightarrow 0 \text{ as } R \rightarrow \infty \end{aligned}$$

i.e. $\left| \int_{\Gamma} f(z) dz \right| = 0$ when $R \rightarrow \infty$ and then (1) gives

$$\int_{-\infty}^{\infty} \frac{dx}{1+x^2} = \pi \text{ or } 2 \int_0^{\infty} \frac{dx}{1+x^2} = \pi, \text{ i.e. } \int_0^{\infty} \frac{ax}{1+x^2} = \frac{\pi}{2}$$

$$(v) \text{ Take } f(z) = \frac{e^{iaz}}{(z^2 + a^2)(z^2 + b^2)}$$

and choose the closed contour C as consisting of the upper half Γ of a large circle $|z|=R$ and the real axis $-R$ to R .

Poles of $f(z)$ are given by $(z^2+a^2)(z^2+b^2)=0$ i.e. $z=\pm ia, \pm ib$ which the only poles lying inside C are $z=ia$ and $z=ib$

Residue of $f(z)$ at $z=ia$ is $\lim_{z \rightarrow ia} (z-ia)f(z)$

$$= \lim_{z \rightarrow ia} (z-ia) \frac{e^{iaz}}{(z^2+a^2)(z^2+b^2)} = \frac{-e^{-a}}{2ia(a^2-b^2)}$$

Similarly residue of $f(z)$ at $z=ib$, $= \frac{+e^{-b}}{2ib(a^2-b^2)}$

Hence by Cauchy's residue theorem

$$\int_{\Gamma} f(z) dz + \int_{-R}^R f(x) dx = 2\pi i \Sigma R^+ = \frac{2\pi i}{2i(a^2-b^2)} \left[\frac{e^{-b}}{b} - \frac{e^{-a}}{a} \right]$$

$$\text{where } \left| \int_{\Gamma} \frac{e^{iaz}}{(z^2+a^2)(z^2+b^2)} dz \right| \leq \int_{\Gamma} \frac{|e^{iaz}| |dz|}{(|z|^2-a^2)(|z|^2-b^2)}$$

$$\because |z_1+z_2| > |z_1|-|z_2|$$

$$\leq \int_0^{\pi} \frac{e^{-R \sin \theta} \cdot R d\theta}{(R^2-a^2)(R^2-b^2)}$$

where $z=Re^{i\theta}$, and so $|dz|=Rd\theta$

$$\leq \frac{2R}{(R^2-a^2)(R^2-b^2)} \frac{\pi}{2R} (1-e^{-R})$$

which $\rightarrow 0$ as $R \rightarrow \infty$

$\therefore$ as $R \rightarrow \infty$, we have

$$\int_{-\infty}^{\infty} \frac{e^{iax}}{(x^2+a^2)(x^2+b^2)} dx = \frac{\pi}{a^2-b^2} \left(\frac{e^{-b}}{b} - \frac{e^{-a}}{a} \right)$$

Equating real and imaginary parts we get

$$\int_{-\infty}^{\infty} \frac{\cos x}{(x^2+a^2)(x^2+b^2)} dx = \frac{\pi}{a^2-b^2} \left(\frac{e^{-b}}{b} - \frac{e^{-a}}{a} \right)$$

$$\text{and } \int_{-\infty}^{\infty} \frac{\sin x}{(x^2+a^2)(x^2+b^2)} dx = 0.$$

Note. Similar procedure will show that

$$\int_{-\infty}^{\infty} \frac{dx}{(x^2+a^2)(x^2+b^2)} = \frac{\pi(a+2b)}{2a^3(a+b)^2}, \quad a>0, b>0$$

$$\text{and } \int_{-\infty}^{\infty} \frac{x^3 \sin x}{(x^2+a^2)(x^2+b^2)} dx = \frac{\pi}{(a^2-b^2)} (a^2 e^{-a} - b^2 e^{-b}), \quad a>0, b>0.$$

$$(vi) \text{ take } f(z) = \frac{\log(z+i)}{z^2+1}$$

and choose the contour C as consisting of upper half Γ of circle $|z|=R$, (R being large) and the real axis from $-R$ to R .

Poles of $f(z)$ are given by $x^2+1=0$ i.e. $z=\pm i$

of which the only pole lying within the contour C is $z=i$. Residue of $f(z)$ at $z=i$ is $\lim_{z \rightarrow i} (z-i) \cdot f(z)$

$$= \lim_{z \rightarrow i} (z-i) \frac{\log(z+i)}{z^2+1}$$

$$= \lim_{z \rightarrow i} (z-i) \frac{\log(z+i)}{(z-i)(z+i)} = \frac{\log(2i)}{2i}$$

$$= \frac{\log 2 + i \frac{\pi}{2}}{2i} \because \log i = \log e^{i\pi/2}$$

Hence by Cauchy's Residue theorem, we have

$$\int_{-R}^R f(x) dx + \int_{\Gamma} f(z) dz = 2\pi i \Sigma R^+$$

$$= 2\pi i \times \frac{1}{2i} \left(\log 2 + \frac{i\pi}{2} \right)$$

$$= \pi \left(\log 2 + \frac{i\pi}{2} \right)$$

$$\text{where } \left| \int_{\Gamma} \frac{\log(z+i)}{z^2+1} dz \right| \leq \int_{\Gamma} \frac{|\log(z+i)| |dz|}{|z^2+1|}$$

$$\leq \int_{\Gamma} \frac{|\log|z+i|| + i \arg(z+i)| |dz|}{|z|^2-1}$$

by (36) of §5.7

$$\leq \int_{\Gamma} \frac{|\log(|z|+1)| + i2\pi |dz|}{|z|^2-1}$$

$$\because \arg(z+i) < 2\pi \text{ and } |z|=1$$

$$\leq \frac{\{\log(R+1)+2\pi\}\pi R}{R^2-1} \text{ when } z=Re^{i\theta}$$

$$\leq \left(\log \frac{(R+1)}{R+1} + \frac{2\pi}{R+1} \right) \frac{\pi}{R-1} \rightarrow 0 \text{ as } R \rightarrow \infty$$

and hence when $R \rightarrow \infty$, we have

$$\int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^{\infty} \frac{\log(x+i)}{x^2+1} dx = \pi \left(\log 2 + \frac{i\pi}{2} \right)$$

$$\text{or } \int_{-\infty}^{\infty} \frac{\frac{1}{2} \log(1+x^2) + i \tan^{-1} \frac{1}{x}}{x^2+1} dx = \pi \left(\log 2 + \frac{i\pi}{2} \right)$$

Equating real and imaginary parts, we get

$$\int_{-\infty}^{\infty} \frac{\frac{1}{2} \log(1+x^2)}{1+x^2} dx = \pi \log 2$$

$$\text{i.e. } \int_0^{\infty} \frac{\log(1+x^2)}{1+x^2} dx = \pi \log 2$$

and $\int_{-\infty}^{\infty} \frac{\tan^{-1} \frac{1}{x}}{1+x^2} dx = \frac{\pi^2}{2}$

Problem 22. Prove by contour integration that

$$\int_0^{\infty} \frac{\sin mx}{x} dx = \frac{\pi}{2}$$

(Agra, 19)




Fig. 5.37

no pole within the contour, we have by Cauchy's theorem,

$$\int_{-R}^{-r} f(x) dx + \int_{\gamma} f(z) dz + \int_r^R f(x) dx + \int_{\Gamma} f(z) dz = 0,$$

Residue at $z=0$ of $f(z)$ is $= \lim_{z \rightarrow 0} z f(z)$

$$= \lim_{z \rightarrow 0} z \cdot \frac{e^{imz}}{z} = 1.$$

Now $\lim_{r \rightarrow 0} \int_{\gamma} f(z) dz = -i(\pi-0) \cdot 1 = -\pi i$ by theorem of §5.16 (sign being taken as contour γ is in clockwise direction) and

$$\left| \int_{\Gamma} f(z) dz \right| \leq \int_{\Gamma} |f(z)| |dz| = \int_{\Gamma} \frac{|e^{imz}| |dz|}{|z|}$$

$$= \int_0^{\pi} \frac{e^{-mR \sin \theta}}{R} R d\theta \quad \because z = Re^{i\theta},$$

$$|dz| = R d\theta$$

$$\leq 2 \int_0^{\pi/2} e^{(-2mR\theta)/\pi} d\theta$$

by Jordan's inequality

$$\leq \frac{m\pi}{R} (1 - e^{-mR})$$

which $\rightarrow 0$ as $R \rightarrow \infty$

With these results as $R \rightarrow \infty$ and $r \rightarrow 0$, (1) reduces to

$$\int_{-\infty}^0 f(x) dx - \pi i + \int_0^{\infty} f(x) dx = 0$$

or

$$\int_{-\infty}^{\infty} \frac{e^{imx}}{x} dx = \pi i$$

$$\int_{-\infty}^{\infty} \frac{e^{imx}}{x} dx = \pi i, \text{ i.e., } \int_0^{\infty} \frac{e^{imx}}{x} dx = \frac{\pi}{2} i.$$

Equating imaginary parts on both sides, we have

$$\int_0^{\infty} \frac{\sin mx}{x} dx = \frac{\pi}{2}.$$

Note. Equating real parts, $\int_0^{\infty} \frac{\cos mx}{x} dx = 0.$

Problem 23. Prove by contour integration that

$$\int_0^{\infty} \frac{x^{\alpha-1}}{1-x} dx = \pi \cot \alpha\pi, \quad 0 < \alpha < 1.$$

Let us consider

$$\int_C f(z) dz, \text{ where } f(z) = \frac{z^{\alpha-1}}{1-z}.$$

Choosing the contour C consisting of a large semi-circle $|z|=R$ indented at $z=0$, $z=1$, let the radii of indentation be r_1 and r_2 respectively.

Since there are no poles within the contour, we have

$$\int_{r_1}^R \frac{x^{\alpha-1}}{1-x} dx + \int_{1+r_2}^R \frac{x^{\alpha-1}}{1-x} dx + \int_R^{r_1} \frac{(xe^{i\pi})^{\alpha-1}}{1-xe^{i\pi}} d(xe^{i\pi}) + \int_{\Gamma} f(z) dz + \int_{\gamma_1} f(z) dz + \int_{\gamma_2} f(z) dz = 0 \quad \dots (1)$$

Now residue of $f(z)$ at $z=0$ is $= \lim_{z \rightarrow 0} z \cdot \frac{z^{\alpha-1}}{1-z} = 0$ as $\alpha > 0$.

Residue of $f(z)$ at $z=1$ is $= \lim_{z \rightarrow 1} (z-1) \frac{z^{\alpha-1}}{1-z} = -1.$

Also $\lim_{r_1 \rightarrow 0} \int_{r_1} f(z) dz = 0,$

$$\lim_{r_2 \rightarrow 0} \int_{\gamma_2} f(z) dz = -i(\pi-0)(-1) = \pi i.$$

$$\left| \int_{\Gamma} \frac{z^{\alpha-1}}{1-z} dz \right| \leq \int_{\Gamma} \frac{|z|^{\alpha-1}}{|z-1|} |dz|$$

$$\leq \frac{R^{\alpha-1}}{R-1} \pi R = \frac{\pi R^{\alpha}}{R-1},$$

which $\rightarrow 0$ as $R \rightarrow \infty$, $\because \alpha < 1.$

With these results as $R \rightarrow \infty$ and r_1, r_2 both tending to zero, (1) reduces to




Fig. 5.38

$$\int_0^1 \frac{x^{\alpha-1}}{1-x} dx + \int_0^\infty \frac{x^{\alpha-1}}{1-x} dx + \int_\infty^0 \frac{e^{2\pi i} x^{\alpha-1}}{1+x} dx + \pi i = 0$$

$$\text{or} \quad \int_0^\infty \frac{x^{\alpha-1}}{1-x} dx - e^{2\pi i} \int_0^\infty \frac{x^{\alpha-1}}{1+x} dx + \pi i = 0$$

$$\text{or} \quad \int_0^\infty \frac{x^{\alpha-1}}{1-x} dx - (\cos \alpha\pi + i \sin \alpha\pi) \int_0^\infty \frac{x^{\alpha-1}}{1+x} dx = -\pi i.$$

Equating real and imaginary parts we find

$$\sin \alpha\pi \int_0^\infty \frac{x^{\alpha-1}}{1+x} dx = \pi, \text{ i.e., } \int_0^\infty \frac{x^{\alpha-1}}{1+x} dx = \frac{\pi}{\sin \alpha\pi}$$

$$\text{and } \int_0^\infty \frac{x^{\alpha-1}}{1-x} dx = \cos \alpha\pi \int_0^\infty \frac{x^{\alpha-1}}{1+x} dx = \cos \alpha\pi \cdot \frac{\pi}{\sin \alpha\pi} = \pi \cot \alpha\pi.$$

Problem 24. If $-1 < p < 1$ and $-\pi < \lambda < \pi$, show by contour integration that

$$\int_0^\infty \frac{x^{-p} dx}{1+2x \cos \lambda + x^2} = \frac{\pi}{\sin p\pi} \frac{\sin p\lambda}{\sin \lambda}.$$

Let us consider $\int_C f(z) dz$, where $f(z) = \frac{z^{-p}}{1+2z \cos \lambda + z^2}$




Fig. 5.39

when the contour C is chosen to be consisting of a large circle $|z| = R$, a small circle $|z| = r$, and the cross cut joining the arcs of the two circles along the real axis.

The poles of $f(z)$ are given by

$$1+2z \cos \lambda + z^2 = 0$$

$$\text{or } (z+e^{i\lambda})(z-e^{-i\lambda}) = 0,$$

$$\text{giving } z = -e^{i\lambda} = -e^{i(\pi+\lambda)}$$

$$\text{and } z = -e^{-i\lambda} = e^{i(\pi-\lambda)}$$

Amplitudes of the point being $\pi+\lambda$, and $\pi-\lambda$ follow that $-\pi < \lambda < \pi$. As such the two amplitudes lie between 0 and 2π .

The poles within the contour are $z = -e^{i\lambda}$ and $z = -e^{-i\lambda}$.

Residue of $f(z)$ at $z = -e^{i\lambda}$ is

$$\begin{aligned} &= \lim_{z \rightarrow -e^{i\lambda}} (z+e^{i\lambda}) \frac{z^{-p}}{(z+e^{i\lambda})(z-e^{-i\lambda})} \\ &= (-1)^{-p} \cdot \frac{e^{-ip\lambda}}{-2i \sin \lambda} = \frac{e^{-ip\lambda}}{-2i \sin \lambda}. \end{aligned}$$

$$\text{Similarly residue at } z = -e^{-i\lambda} \text{ is } e^{-ip\lambda} \frac{e^{ip\lambda}}{2i \sin \lambda}.$$

$$\text{Sum of residues} = \frac{e^{-ip\lambda}}{\sin \lambda} \frac{e^{ip\lambda} - e^{-ip\lambda}}{2i} = \frac{e^{-ip\lambda}}{\sin \lambda} \sin p\lambda.$$

Applying Cauchy's residue theorem, we have

$$\begin{aligned} \int_r^R f(x) dx + \int_\Gamma f(z) dz - \int_r^R f(xe^{2\pi i}) dx - \int_\gamma f(z) dz \\ = 2\pi i \times \text{sum of the residues.} \quad \dots (1) \end{aligned}$$

$$\begin{aligned} \text{Here } \left| \int_\Gamma f(z) dz \right| &\leq \int_\Gamma \frac{|z|^{-p} |dz|}{|z+e^{i\lambda}| |z+e^{-i\lambda}|} \\ &\leq \frac{R^{-p} \cdot 2\pi R}{(R-1)(R-1)} = \frac{2\pi}{R^{1+p} \left(1 - \frac{1}{R}\right)^2}, \end{aligned}$$

which $\rightarrow 0$ as $R \rightarrow \infty$ provided $1+p > 0$, i.e. $p > -1$.

$$\text{Similarly } \left| \int_\gamma f(z) dz \right| \leq 2\pi \frac{r^{1-p}}{(1+r)^2} \text{ which } \rightarrow 0 \text{ as } z \rightarrow 0,$$

if $1-p > 0$ or $1 > p$,

Hence there exists a condition $-1 < p < 1$.

With these results as $R \rightarrow \infty$ and $r \rightarrow 0$, (1) reduces to

$$\int_0^\infty f(x) dx - \int_0^\infty f(xe^{2\pi i}) e^{2\pi i} dx = 2\pi i \cdot e^{-p\pi i} \frac{\sin p\lambda}{\sin \lambda}$$

$$\begin{aligned} \int_0^\infty \frac{x^{-p} dx}{1+2x \cos \lambda + x^2} - \int_0^\infty \frac{(xe^{2\pi i})^{-p} \cdot e^{2\pi i} dx}{1+2xe^{2\pi i} \cos \lambda + (xe^{2\pi i})^2} \\ = 2\pi i e^{-p\pi i} \frac{\sin p\lambda}{\sin \lambda} \end{aligned}$$

$$(1 - e^{-2p\pi i}) \int_0^\infty \frac{x^{-p} dx}{1+2x \cos \lambda + x^2} = 2\pi i e^{-p\pi i} \frac{\sin p\lambda}{\sin \lambda} \quad \because e^{2\pi i} = 1$$

$$\int_0^\infty \frac{x^{-p} dx}{1+2x \cos \lambda + x^2} = \frac{2\pi i \cdot e^{-p\pi i} \sin p\lambda}{e^{-p\pi i} (e^{p\pi i} - e^{-p\pi i}) \sin \lambda} = \frac{p}{\sin p\pi} \cdot \frac{\sin p\lambda}{\sin \lambda}$$

Problem 25. By integrating $e^{iz} z^{\alpha-1}$ round a quadrant of a circle of radius R , prove that if $0 < \alpha < 1$,

$$\int_0^\infty x^{\alpha-1} \frac{\cos x}{\sin x} dx = \Gamma(\alpha) \frac{\cos \frac{\pi}{2} \alpha}{\sin \frac{\pi}{2} \alpha}.$$

Consider $\int_C f(z) dz$, where $f(z) = e^{iz} z^{\alpha-1}$.

Choosing the contour C consisting of a quadrant of a large circle $|z| = R$ indented at $z=0$, let r be the radius of indentation. By Cauchy's integral, we have

$$\int_C f(z) dz = 0.$$

Here $z = \infty$ is not a pole, but $z=0$ is a pole which has been excluded from the contour by indentation.

Splitting up the above contour, we get

$$\int_r^R e^{iz} z^{\alpha-1} dz + \int_\Gamma e^{iz} z^{\alpha-1} dz + \int_R^r e^{iz} (iy)^{\alpha-1} i dy + \int_\gamma e^{iz} z^{\alpha-1} dz = 0 \quad \dots (1)$$




Fig. 5.40

Residue of $f(z)$ at $z=0$ is $\lim_{z \rightarrow 0} z \cdot e^{iz} z^{\alpha-1} = 0$

$$\begin{aligned} \text{and } \left| \int_{\Gamma} e^{iz} z^{\alpha-1} dz \right| &= \left| \int_{\Gamma} e^{iR\theta} (Re^{i\theta})^{\alpha-1} iRe^{i\theta} d\theta \right| \text{ as } z = Re^{i\theta} \\ &\leq \int_0^{\pi/2} e^{-R\sin\theta} |e^{iR\cos\theta}| |i| |R^{\alpha}| |e^{i\theta}| d\theta \\ &\leq \int_0^{\pi/2} e^{-R\sin\theta} R^{\alpha} d\theta \\ &\leq \int_0^{\pi/2} e^{-2R\theta/\pi} R^{\alpha} d\theta \text{ as } \frac{\sin\theta}{\theta} > \frac{2}{\pi} \text{ (Jordan's inequality)} \\ &\leq \frac{R^{\alpha}}{2} \cdot \frac{\pi}{R} [1 - e^{-R}] \text{ which tends to zero as } R \rightarrow \infty \end{aligned}$$

$$\begin{aligned} \text{Similarly } \left| \int_{\gamma} e^{iz} z^{\alpha-1} dz \right| &\leq \frac{\pi}{2} \cdot \frac{r^{\alpha}}{r} (1 - e^{-r}) \quad \because z = re^{i\theta} \\ &\leq \frac{\pi}{2} \cdot \frac{r^{\alpha}}{r} \left(r - \frac{r^2}{2!} + \dots \right) \\ &\text{which tends to zero as } r \rightarrow 0. \end{aligned}$$

Hence with these results as $R \rightarrow \infty$ and $r \rightarrow 0$, (1) reduces to

$$\int_0^{\infty} e^{ix} x^{\alpha-1} dx - \int_0^{\infty} e^{-iy} y^{\alpha-1} dy = 0$$

$$\begin{aligned} \text{or } \int_0^{\infty} (\cos x + i \sin x) x^{\alpha-1} dx &= i^{\alpha} \int_0^{\infty} e^{-y} y^{\alpha-1} dy = (e^{i\pi/2})^{\alpha} \Gamma(\alpha) \\ \therefore \int_0^{\infty} e^{-y} y^{\alpha-1} dx &= \Gamma(\alpha) \quad \text{and } i = e^{i\pi/2}. \end{aligned}$$

Equating real and imaginary parts, we get

$$\int_0^{\infty} \cos x \cdot x^{\alpha-1} dx = \Gamma(\alpha) \cos \frac{\pi}{2} \alpha \quad \text{and} \quad \int_0^{\infty} \sin x \cdot x^{\alpha-1} dx = \Gamma(\alpha) \sin \frac{\pi}{2} \alpha.$$

Problem 26. Prove that $\int_0^{\infty} e^{-x^2} \cos 2ax dx = \frac{1}{2} \sqrt{\pi} e^{-a^2}$ by integrating e^{-z^2} round the rectangle whose vertices are $O, R, R+ia, ia$.




Fig. 5.41

Let A be $(R, 0)$, C be $(0, a)$ in the Argand diagram.

$$\text{On } OA, z=x; \quad \therefore dz=dx.$$

$$\text{On } AB, z=R+iy; \quad \therefore dz=i dy.$$

$$\text{On } BC, z=x+ia; \quad \therefore dz=dx$$

$$\text{and on } OC, z=iy; \quad \therefore dz=i dy$$

Now e^{-z^2} has no poles within or on this contour and if the whole contour be denoted by Γ , then by Cauchy's theorem

$$\int_{\Gamma} e^{-z^2} dz = 0.$$

Splitting up the contour

$$\int_0^{\infty} e^{-x^2} dx + \int_0^a e^{-(R+iy)^2} i dy + \int_R^0 e^{-(x+ia)^2} dx + \int_a^0 e^{-(iy)^2} i dy = 0. \quad \dots(1)$$

$$\begin{aligned} \text{Now } \left| \int_0^a e^{-(R+iy)^2} i dy \right| &\leq \int_0^a |e^{-(R^2+2iRy-y^2)}| |i| dy \\ &\leq \int_0^a e^{-R^2} e^{y^2} |e^{-2iRy}| dy \\ &\leq \int_0^a e^{-R^2} e^{y^2} dy \quad \because |e^{-2iRy}| = 1 \\ &\quad \text{and } |i| = 1 \end{aligned}$$

which tends to zero as $R \rightarrow \infty$.

Hence when $R \rightarrow \infty$, (1) reduces to

$$\int_0^{\infty} e^{-x^2} dx - \int_0^{\infty} e^{-(x+ia)^2} dx - i \int_0^a e^{y^2} dy = 0$$

$$\int_0^{\infty} e^{-x^2} dx - e^{-a^2} \int_0^{\infty} e^{-x^2} e^{-2iax} dx - i \int_0^a e^{y^2} dy = 0.$$

$$\int_0^{\infty} e^{-x^2} dx - e^{-a^2} \int_0^{\infty} e^{-x^2} [\cos 2ax - i \sin 2ax] dx - i \int_0^a e^{y^2} dy = 0.$$

$$\text{Equating real parts, } e^{-a^2} \int_0^{\infty} e^{-x^2} \cos 2ax dx = \int_0^{\infty} e^{-x^2} dx = \frac{\sqrt{\pi}}{2}.$$

$$\int_0^{\infty} e^{-x^2} \cos 2ax dx = \frac{1}{2} \sqrt{\pi} e^{-a^2}$$

Problem 27. Using the rectangular contour, prove the following:

$$(i) \int_0^{\infty} \frac{t^{\alpha-1}}{1+t} dt = \pi \operatorname{cosec} \alpha\pi; \quad 0 < \alpha < 1.$$

(For $t=e^x$, Rohilkhand, 1977)

$$(ii) \int_0^{\infty} \frac{\cosh \alpha x}{\cosh \pi x} dx = \frac{1}{2} \sec \frac{\alpha}{2}; \quad -\pi < \alpha < \pi$$

$$(iii) \int_0^{\infty} \frac{\sinh \alpha x}{\cosh \pi x} dx = \frac{1}{2} \tan \frac{\alpha}{2}; \quad 0 < \alpha < \pi.$$

(i) Given integral is

$$I = \int_0^{\infty} \frac{t^{\alpha-1}}{1+t} dt$$

Putting $t=e^x$, we get

$$I = \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{e^x + 1} dx.$$

Consider,

$$\int_C f(z) dz = \int_C \frac{e^{\alpha z}}{e^z + 1} dz$$




Fig. 5.42

where C is the rectangle $ABCD$ with vertices at $R, R+2\pi i, -R+2\pi i, -R$; R being positive.

$$\text{Here, } f(z) = \frac{e^{\alpha z}}{e^z + 1}.$$

The poles of $f(z)$ are given by $e^z + 1 = 0$

$$\text{i.e. } e^z = -1 = e^{(2n+1)\pi i}$$

$$\text{or } z = (2n+1)\pi i \quad \text{where } n=0, \pm 1, \pm 2, \pm 3, \dots$$

Only the simple pole $z = \pi i$, (corresponding to $n=0$) lies inside the contour.

$$\text{Residue of } f(z) \text{ at } z = \pi i = \left[\frac{d}{dx} \frac{e^{\alpha x}}{(1+e^x)} \right]_{z=\pi i} = \frac{e^{\alpha \pi i}}{e^{\pi i}} = -e^{\alpha \pi i}$$

$$\text{as } e^{\pi i} = \cos \pi + i \sin \pi = -1.$$

By Cauchy's residue theorem, we therefore have

$$\begin{aligned} \int_{-R}^R f(x) dx + \int_0^{2\pi} f(R+iy) i dy + \int_R^{-R} f(x+2\pi i) dx \\ + \int_{2\pi}^0 f(-R+iy) i dy = 2\pi i \times \text{sum of residues} \\ = -2\pi i e^{\alpha \pi i}. \quad \dots (1) \end{aligned}$$

$$\begin{aligned} \text{But } \left| \int_0^{2\pi} f(R+iy) i dy \right| &\leq \int_0^{2\pi} |f(R+iy)| |i dy| \\ &\leq \int_0^{2\pi} \left| \frac{e^{\alpha(R+iy)}}{1+e^{R+iy}} \right| dy \quad \text{as } |i| = 1 \\ &\leq \int_0^{2\pi} \left| \frac{e^{\alpha R}}{e^R - 1} \right| dy \quad \text{as } |e^{R+iy} + 1| \leq e^{R+iy} - 1 \\ &\leq \frac{e^{\alpha R}}{e^R - 1} \cdot 2\pi. \\ &\leq \frac{2\pi}{e^{(1-\alpha)R} - e^{-\alpha R}} \text{ which } \rightarrow 0 \text{ as } R \rightarrow \infty \text{ for } 0 < \alpha < 1 \end{aligned}$$

$$\therefore \int_0^{2\pi} f(R+iy) i dy \rightarrow 0 \text{ when } R \rightarrow \infty.$$

$$\text{Similarly } \int_0^{2\pi} f(-R+iy) i dy \rightarrow 0 \text{ when } R \rightarrow \infty,$$

provided $0 < \alpha < 1$.

Hence proceeding to the limit when $R \rightarrow \infty$, (1) gives

$$\int_{-\infty}^{\infty} f(x) dx + \int_{\infty}^{-\infty} f(x+2\pi i) dx = -2\pi i e^{\alpha \pi i}$$

$$\int_{-\infty}^{\infty} \frac{e^{\alpha x}}{e^x + 1} dx - \int_{-\infty}^{\infty} \frac{e^{\alpha(x+2\pi i)}}{e^{x+2\pi i} + 1} dx = -2\pi i e^{\alpha \pi i}$$

$$\int_{-\infty}^{\infty} \frac{e^{\alpha x}}{e^x + 1} dx - \int_{-\infty}^{\infty} \frac{e^{2\alpha \pi i} e^{\alpha x}}{e^x + 1} dx = -2\pi i e^{\alpha \pi i}$$

$$\int_{-\infty}^{\infty} \frac{e^{\alpha x} (1 - e^{2\alpha \pi i})}{e^x + 1} dx = -2\pi i e^{\alpha \pi i}$$

$$\int_{-\infty}^{\infty} \frac{e^{\alpha x}}{e^x + 1} dx = \frac{-2\pi i e^{\alpha \pi i}}{1 - e^{2\alpha \pi i}}$$

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{e^x + 1} dx &= \pi \cdot \frac{2i}{e^{\alpha \pi i} - e^{-\alpha \pi i}} \\ &= \frac{\pi}{\sin \alpha \pi} \end{aligned}$$

$$\int_0^{\infty} \frac{t^{\alpha-1}}{t+1} dt = \pi \operatorname{cosec} \alpha \pi$$

(ii) Given integral is

$$I = \int_0^{\infty} \frac{\cosh \alpha x}{\cosh \pi x} dx.$$

Consider

$$\int_C f(z) dz = \int \frac{e^{\alpha z}}{\cosh \pi z} dz$$

$$\text{where } f(z) = \frac{e^{\alpha z}}{\cosh \pi z}$$

taken round the rectangular

contour which consists of x -axis from $-R$ to R and y -axis from 0 to 1 .

Poles of $f(z)$ are given by $\cosh \pi z = 0$

$$\text{i.e. } \frac{e^{\pi z} + e^{-\pi z}}{2} = 0$$

$$\text{or } e^{\pi z} = -e^{-\pi z} = -\frac{1}{e^{\pi z}}$$

$$\text{or } e^{2\pi z} = -1 = e^{i(2n+1)\pi}$$

$$\text{i.e. } 2z = i(2n+1) \quad \text{where } n=0, \pm 1, \pm 2, \dots$$

$$\text{or } z = \frac{2n+1}{2} i$$

Only the simple pole $z = \frac{i}{2}$ lies within the contour.

$$\text{Residue of } f(z) \text{ at } z = \frac{i}{2} = \left[\frac{d}{dz} \frac{e^{\alpha z}}{\cosh \pi z} \right]_{z=i/2}$$




Fig. 5.43

$$= \frac{e^{\alpha i/2}}{\pi \sinh \frac{\pi i}{2}} = \frac{e^{\alpha i/2}}{\pi i \sin \frac{\pi}{2}} = \frac{e^{\alpha i/2}}{\pi i}.$$

By Cauchy's residue theorem, we have

$$\int_{-R}^R f(x) dx + \int_0^1 f(R+iy) i dy + \int_R^{-R} f(x+i) dx + \int_1^0 f(-R+iy) i dy = 2\pi i \cdot \frac{e^{\alpha i/2}}{\pi i} \dots (1)$$

But

$$\begin{aligned} \left| \int_0^1 f(R+iy) i dy \right| &\leq \int_0^1 |f(R+iy)| |i| dy \\ &\leq \int_0^1 \left| \frac{e^{\alpha(R+iy)}}{\cosh \pi(R+iy)} \right| dy \text{ as } |i|=1 \\ &\leq \int_0^1 \frac{2e^{\alpha R} |e^{\alpha iy}| dy}{|e^{\pi(R+iy)} + e^{-\pi(R+iy)}|} \\ &\leq \int_0^1 \frac{2e^{\alpha R}}{e^{\pi R} - e^{-\pi R}} dy \\ &\leq \frac{2e^{\alpha R}}{e^{\pi R} - e^{-\pi R}} \end{aligned}$$

which $\rightarrow 0$ as $R \rightarrow \infty$ when $\alpha < \pi$

$$\therefore \int_0^1 f(R+iy) i dy \rightarrow 0 \text{ as } R \rightarrow \infty.$$

$$\text{Similarly } \int_0^1 f(-R+iy) i dy \rightarrow 0 \text{ as } R \rightarrow \infty \text{ when } \alpha < \pi.$$

Hence when $R \rightarrow \infty$, (1) becomes

$$\int_{-\infty}^{\infty} f(x) dx + \int_{\infty}^{-\infty} f(x+i) dx = 2e^{i\alpha/2}$$

$$\text{or } \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx + \int_{\infty}^{-\infty} \frac{e^{\alpha(x+i)}}{\cosh \pi(x+i)} dx = 2e^{i\alpha/2}.$$

$$\text{But } \cosh \pi(x+i) = \cosh i\pi(x+i) = \cos(\pi xi - \pi) = -\cos \pi xi = -\cosh \pi x$$

$$\therefore \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx - \int_{\infty}^{-\infty} \frac{e^{\alpha(x+i)}}{\cosh \pi x} dx = 2e^{i\alpha/2}$$

$$\text{or } \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx + \int_{-\infty}^{\infty} \frac{e^{\alpha x} \cdot e^{\alpha i}}{\cosh \pi x} dx = 2e^{i\alpha/2}$$

$$\text{or } (1+e^{i\alpha}) \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx = 2e^{i\alpha/2}$$

$$\text{or } \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx = \frac{2e^{i\alpha/2}}{1+e^{i\alpha}} = \frac{2}{e^{i\alpha/2} + e^{-i\alpha/2}}$$

$$\text{or } \int_{-\infty}^0 \frac{e^{\alpha x}}{\cosh \pi x} dx + \int_0^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx = \frac{2}{2 \cos \frac{\alpha}{2}} = \sec \frac{\alpha}{2}.$$

If we replace x by $-x$ in the first integral this becomes

$$\int_0^{\infty} \frac{e^{-\alpha x}}{\cosh \pi x} dx$$

$$\therefore \int_0^{\infty} \frac{e^{-\alpha x}}{\cosh \pi x} dx + \int_0^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx = \sec \frac{\alpha}{2}$$

$$\text{or } \int_0^{\infty} \frac{e^{\alpha x} + e^{-\alpha x}}{\cosh \pi x} dx = \sec \frac{\alpha}{2}$$

$$\text{or } 2 \int_0^{\infty} \frac{\cosh \alpha x}{\cosh \pi x} dx = \sec \frac{\alpha}{2}$$

$$\text{i.e. } \int_0^{\infty} \frac{\cosh \alpha x}{\cosh \pi x} dx = \frac{1}{2} \sec \frac{\alpha}{2}.$$

Aliter, Consider

$$\int_C f(z) dz = \int_C \frac{e^{\alpha z}}{\cosh \pi z} dz,$$

$$\text{where } f(z) = \frac{e^{\alpha z}}{\cosh \pi z},$$

taken round the rectangular contour with vertices at

$$R, R + \frac{i}{2}, -R + \frac{i}{2} \text{ and } -R,$$




Fig. 5.44

indented at $z = \frac{i}{2}$, since $z = \frac{i}{2}$ is a pole of $f(z)$.

Clearly $f(z)$ is regular within and on the contour, so that Cauchy's theorem gives

$$\begin{aligned} \int_{-R}^R f(x) dx + \int_0^{1/2} f(R+iy) i dy + \int_R^{-R} f(x+\frac{1}{2}i) dx \\ + \int_{\gamma} f(z) dz + \int_{-P}^{-R} f(x+\frac{1}{2}i) dx + \int_{1/2}^0 f(-R+\frac{1}{2}i) i dy = 0. \dots (1) \end{aligned}$$

$$\begin{aligned} \text{But } \lim_{z \rightarrow \frac{1}{2}i} (z - \frac{1}{2}i) f(z) &= \lim_{z \rightarrow \frac{1}{2}i} \frac{(z - \frac{1}{2}i) e^{\alpha z}}{\cosh \pi z} \\ &= \frac{e^{\frac{1}{2}\alpha i}}{\pi \sinh \frac{1}{2}\pi} = \frac{e^{\frac{1}{2}\alpha i}}{\pi i \sin \frac{1}{2}\pi} = \frac{e^{\frac{1}{2}\alpha i}}{\pi i}. \end{aligned}$$

$$\therefore \lim_{\rho \rightarrow 0} \int_{\gamma} f(z) dz = (-\pi i) \cdot \frac{1}{\pi i} e^{\frac{1}{2}\alpha i} = -e^{\frac{1}{2}\alpha i}$$

$$\begin{aligned} \text{and } \left| \int_0^{1/2} f(R+iy) i dy \right| &\leq \int_0^{1/2} \frac{e^{\alpha(R+iy)}}{\cosh \pi(R+iy)} |i| dy \\ &\leq \int_0^{1/2} \frac{e^{\alpha R} dy}{\cosh \pi R \cos \pi y - \sinh \pi R \sin \pi y} \\ &[\because |\cosh \pi R \cos \pi y + i \sinh \pi R \sin \pi y| \\ &\geq \cosh \pi R \cos \pi y - \sinh \pi R \sin \pi y] \\ &\leq \int_0^{1/2} \frac{e^{\alpha R} dy}{e^{\pi R (\cos \pi y - \sin \pi y)}} \\ &\text{writing } e^{\pi R} \text{ for } \cos \pi R \text{ and } \sin \pi R \text{ when} \\ &\quad R \text{ is large} \\ &\rightarrow 0 \text{ when } R \rightarrow \infty \text{ if } \alpha < \pi. \end{aligned}$$

$$\therefore \int_0^{1/2} f(R+iy) i dy \rightarrow 0 \text{ when } R \rightarrow \infty.$$

Similarly $\int_{1/2}^0 f(R+iy) i dy \rightarrow 0$ when $R \rightarrow \infty$ provided $\alpha < \pi$.

Hence in the limit when $R \rightarrow \infty$ and $\rho \rightarrow 0$, (1) becomes

$$\int_{-\infty}^{\infty} f(x) dx + \int_{-\infty}^0 f(x+\frac{1}{2}i) dx - e^{\frac{1}{2}\alpha i} + \int_0^{\infty} f(x+\frac{1}{2}i) dx = 0$$

$$\text{or } \int_{-\infty}^{\infty} f(x) dx - \int_{-\infty}^0 f(x+\frac{1}{2}i) dx = e^{\frac{1}{2}\alpha i}$$

$$\text{or } \int_{-\infty}^{\infty} \left[\frac{e^{\alpha x}}{\cosh \pi x} - \frac{e^{\alpha(x+\frac{1}{2}i)}}{\cosh \pi(x+\frac{1}{2}i)} \right] dx = e^{\frac{1}{2}\alpha i}$$

$$\text{or } \int_{-\infty}^{\infty} \left[\frac{e^{\alpha x}}{\cosh \pi x} - \frac{e^{\alpha x} \cdot e^{\frac{1}{2}\alpha i}}{i \sinh \pi x} \right] dx = \cos \frac{1}{2}\alpha + i \sin \frac{1}{2}\alpha$$

$$\text{or } \int_{-\infty}^{\infty} \left[\frac{e^{\alpha x}}{\cosh \pi x} + \frac{ie^{\alpha x} (\cos \frac{1}{2}\alpha + i \sin \frac{1}{2}\alpha)}{\sinh \pi x} \right] dx = \cos \frac{1}{2}\alpha + i \sin \frac{1}{2}\alpha$$

Equating real and imaginary parts, we get

$$\int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx - \int_{-\infty}^{\infty} \frac{e^{\alpha x} \cdot \sin \frac{1}{2}\alpha}{\sinh \pi x} dx = \cos \frac{1}{2}\alpha. \dots (2)$$

$$\text{and } \int_{-\infty}^{\infty} \frac{e^{\alpha x} \cos \frac{1}{2}\alpha}{\sinh \pi x} dx = \sin \frac{1}{2}\alpha$$

$$\text{or } \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\sinh \pi x} dx = \tan \frac{1}{2}\alpha. \dots (3)$$

(2) gives with the help of (3),

$$\int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx - \sin \frac{1}{2}\alpha \cdot \tan \frac{1}{2}\alpha = \cos \frac{1}{2}\alpha$$

$$\text{or } \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx = \cos \frac{1}{2}\alpha + \frac{\sin^2 \frac{1}{2}\alpha}{\cos \frac{1}{2}\alpha} = \sec \frac{1}{2}\alpha$$

$$\text{or } \int_{-\infty}^0 \frac{e^{\alpha x}}{\cosh \pi x} dx + \int_0^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx = \sec \frac{1}{2}\alpha$$

$$\text{or } \int_0^{\infty} \frac{e^{-\alpha x}}{\cosh \pi x} dx + \int_0^{\infty} \frac{e^{\alpha x}}{\cosh \pi x} dx = \sec \frac{1}{2}\alpha$$

(on replacing x by $-x$ in the first integral),

$$\text{i.e. } \int_0^{\infty} \frac{e^{\alpha x} + e^{-\alpha x}}{\cosh \pi x} dx = \sec \frac{1}{2}\alpha$$

$$\text{or } \int_0^{\infty} \frac{\cosh \alpha x}{\cosh \pi x} dx = \frac{1}{2} \sec \frac{1}{2}\alpha.$$

(iii) This may be proved independently as (ii) or may be derived from (3) of preceding aliter as follows:

$$\text{We have } \int_{-\infty}^{\infty} \frac{e^{\alpha x}}{\sinh \pi x} dx = \tan \frac{1}{2}\alpha$$

$$\text{or } \int_{-\infty}^0 \frac{e^{\alpha x}}{\sinh \pi x} dx + \int_0^{\infty} \frac{e^{\alpha x}}{\sinh \pi x} dx = \tan \frac{1}{2}\alpha$$

$$\text{or } -\int_0^{\infty} \frac{e^{-\alpha x}}{\sinh \pi x} dx + \int_0^{\infty} \frac{e^{\alpha x}}{\sinh \pi x} dx = \tan \frac{1}{2}\alpha$$

(on replacing x by $-x$ in the first integral)

$$\text{or } \int_0^{\infty} \frac{e^{\alpha x} - e^{-\alpha x}}{\sinh \pi x} dx = \tan \frac{1}{2}\alpha, \text{ i.e. } \int_0^{\infty} \frac{\sinh \alpha x}{\sinh \pi x} dx = \frac{1}{2} \tan \frac{1}{2}\alpha.$$

Problem 28. Evaluate the following integrals by contour integration:

$$(i) \int_0^{\infty} \frac{\sin x}{x} dx. \quad (\text{Agra, 1961, 70})$$

$$(ii) \int_0^{2\pi} \frac{d\theta}{\frac{5}{4} + \sin \theta}. \quad (\text{Agra, 1961})$$

$$(iii) \int_0^{\infty} \frac{\cos x}{x^2 + a^2} dx. \quad (\text{Agra, 1962, 65, 71})$$

$$(iv) \int_{-\infty}^{\infty} \frac{x^2 dx}{(x^2 + a^2)^3}. \quad (\text{Agra, 1962, 65})$$

$$(v) \int_0^{\infty} \frac{x^2}{1+x^4} dx.$$

$$(vi) \int_0^{\infty} \frac{1 - \cos x}{x^2} dx. \quad (\text{Agra, 1964})$$

$$(vii) \int_{-\infty}^{\infty} (x^2 + a^2)^{-2} (x^2 + b^2)^{-1} dx. \quad (\text{Agra, 1966; Vikram, 1967})$$

(i) The given integral is

$$\int_0^{\infty} \frac{\sin x}{x} dx.$$

In order to evaluate it, let us consider

$$\int_C f(z) dz = \int_C \frac{e^{iz}}{z} dz,$$

where the contour C is chosen as consisting of the real axis from $-R$ to $+R$,




Fig. 5.45

indented at $z=0$ and the upper half of the semi-circle $|z|=R$ is represented by Γ . Let radius of indentation be R and let it be represented by γ . Here $f(z)$ has got a simple pole at $z=0$.

Since there is no pole within the contour, we have by Cauchy's theorem

$$\int_C f(z) dz = 0, \\ \text{i.e., } \int_{-R}^{-r} f(x) dx + \int_{\gamma} f(z) dz + \int_r^R f(x) dx + \int_{\Gamma} f(z) dz = 0. \quad \dots (1)$$

Residue at $z=0$ of $f(z)$ is $\lim_{z \rightarrow 0} z f(z)$

$$= \lim_{z \rightarrow 0} z \cdot \frac{e^{iz}}{z} = 1.$$

Now $\lim_{r \rightarrow 0} \int_{\gamma} f(z) dz = -i(\pi - 0) = -\pi i$ by theorem of §5.16 (sign being taken as contour γ is in clockwise direction) and

$$\left| \int_{\Gamma} f(z) dz \right| \leq \int_{\Gamma} |f(z)| |dz| = \int_{\Gamma} \frac{|e^{iz}|}{|z|} |dz| \\ = \int_0^{\pi} \frac{e^{-R \sin \theta}}{R} R d\theta, \\ \because z = R e^{i\theta}, |dz| = R d\theta$$

$$\leq 2 \int_0^{\pi/2} e^{-2R\theta/\pi} \text{ by Jordan's inequality}$$

$$\leq 2 \cdot \left(\frac{-\pi}{2R} \right) \left[e^{-2R\theta/\pi} \right]_0^{\pi/2}$$

$$\leq \frac{\pi}{R} (1 - e^{-R}), \text{ which } \rightarrow 0, \text{ as } R \rightarrow \infty.$$

With these results as $R \rightarrow \infty$ and $r \rightarrow 0$, (1) reduces to

$$\int_{-\infty}^{\infty} f(x) dx - \pi i + \int_0^{\infty} (x) dx = 0$$

$$\text{or } \int_{-\infty}^{\infty} f(x) dx = \pi i$$

$$\text{or } 2 \int_0^{\infty} \frac{e^{ix}}{x} dx = \pi i,$$

$$\text{i.e. } \int_0^{\infty} \frac{\cos x + i \sin x}{x} dx = \frac{\pi}{2} i.$$

Equating imaginary parts on either side, we get

$$\int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}.$$

$$(ii) \text{ Given } I = \int_0^{2\pi} \frac{d\theta}{\frac{\pi}{2} + \sin \theta}.$$

Choosing the contour C as a circle ($|z|=1$) of unit radius, we have

$$z = e^{i\theta}, \text{ so that } z - \frac{1}{z} = 2i \sin \theta, \text{ since } \frac{1}{z} = e^{-i\theta}$$

and

$$dz = ie^{i\theta} d\theta = iz d\theta.$$

$$\therefore I = \frac{4}{i} \int_C \frac{1}{5+4 \left\{ \frac{1}{2i} \left(z - \frac{1}{z} \right) \right\}} \frac{dz}{z} = 4 \int_C \frac{dz}{5iz + 2(z^2 - 1)}.$$

$$\text{Here } f(z) = \frac{1}{2z^2 + 5iz - 2}.$$

Its poles are given by $2z^2 + 5iz - 2 = 0$,

$$\text{i.e. } z = \frac{-5i \pm \sqrt{(-25 + 16)}}{2} = \frac{-5 \pm 3i}{4} = -\frac{i}{2}, -2i$$

of which the pole $z = -2i$ lies outside the contour and therefore the only pole that lies within the contour is $z = -i/2$ (which is of order one).

$\therefore$ Residue at $z = -i/2$ is

$$= \lim_{z \rightarrow -i/2} (z + \frac{i}{2}) f(z) = \lim_{z \rightarrow -i/2} (z + \frac{i}{2}) \cdot \frac{1}{\left(z + \frac{i}{2} \right) (z + 2i) \cdot 2}$$

$$\lim_{z \rightarrow -i/2} \frac{1}{2(z + 2i)} = \frac{1}{2(-i/2 + 2i)} = \frac{1}{3i}.$$

Hence by Cauchy's theorem on residues

$I = 4 \times 2\pi i \Sigma R$, where ΣR represents the sum of residues

$$= 8\pi i \cdot \frac{1}{3i} = \frac{8\pi}{3}.$$

$$(iii) \text{ The given integral is } \int_0^{\infty} \frac{\cos x}{x^2 + a^2} dx.$$

$$\text{Let us consider } \int_C f(z) dz = \int_C \frac{e^{iaz}}{z^2 + a^2} dz = I \text{ (say).}$$

Choosing the closed contour C , consisting of real axis from $-R$ to R and upper half of a large circle $|z|=R$, represented by Γ , we have by Cauchy's residue theorem,

$$\int_C f(z) dz = \int_{-R}^R f(x) dx + \int_{\Gamma} f(z) dz \\ = 2\pi i \Sigma R^+. \quad \dots (1)$$

Poles of $f(z)$ are given by $z^2 + a^2 = 0$, i.e. $z = \pm ia$ of which only the simple pole $z = ia$ lies within the contour.




Fig. 5.46

Residue at $z = +ia$ (a simple pole)

$$\begin{aligned} &= \lim_{z \rightarrow ia} (z - ia) \frac{e^{iz}}{(z - ia)(z + ia)} \\ &= \frac{e^{i^2 a}}{2ia} = \frac{e^{-a}}{2ia}. \end{aligned}$$

$$\begin{aligned} \text{Now } \left| \int_{\Gamma} f(z) dz \right| &= \left| \int_{\Gamma} \frac{e^{iz}}{z^2 + a^2} dz \right| \\ &\leq \int_{\Gamma} \frac{|e^{iz}| |dz|}{|z^2 + a^2|} \\ &\leq 0 \text{ as } z \rightarrow \infty. \end{aligned}$$

$\therefore \int_{\Gamma} f(z) dz = 0$, since modulus cannot be negative.

Hence when $R \rightarrow \infty$, (1) reduces to

$$\int_{-\infty}^{\infty} f(x) dx = 2\pi i \times \frac{e^{-a}}{2ia} = \frac{\pi}{a} e^{-a} = \frac{\pi}{ae^a},$$

$$\text{i.e., } 2 \int_0^{\infty} \frac{(\cos x + i \sin x)}{x^2 + a^2} dx = \frac{\pi}{ae^a}.$$

Equating real parts on either side, we get

$$\int_0^{\infty} \frac{\cos x}{x^2 + a^2} = \frac{\pi}{2ae^a}.$$

$$(iv) \text{ Suppose } I = \int_{-\infty}^{\infty} \frac{x^2}{(x^2 + a^2)^3} dx.$$

Here

$$f(z) = \frac{z^2}{(z^2 + a^2)^3}.$$

$zf(z) \rightarrow 0$ as $z \rightarrow \infty$, therefore the condition for the evaluation of an infinite integral is satisfied.




Fig. 5.47

Let us choose the contour consisting of a semi-circle Γ of radius R , large enough to include all the poles of $f(z)$ and the part of the real axis from $-R$ to $+R$.

Poles of $f(z)$ are given by $(z^2 + a^2)^3 = 0$, i.e., $z = \pm ia$ (each pole is of order three).

The only pole $z = \pm ai$ (of order three) lies within the contour C .

Applying the theorem of residues,

$$\int_C f(z) dz = \int_{-R}^R f(x) dx + \int_{\Gamma} f(z) dz = 2\pi i \Sigma R^+. \quad \dots (1)$$

Since $\lim_{z \rightarrow \infty} zf(z) \rightarrow 0$, we have

$$\lim_{R \rightarrow \infty} \int_{\Gamma} f(z) dz = 0.$$

Hence when $R \rightarrow \infty$, (1) becomes

$$\int_{-\infty}^{\infty} f(x) dx = 2\pi i \Sigma R^+. \quad \dots (2)$$

Now residue of $f(z)$ at $z = ai$

= coeff. of $\frac{1}{t}$ in $f(ai+t)$, t being small

$$\begin{aligned} &= \text{coeff. of } \frac{(ai+t)^2}{\{(ai+t)^2 + a^2\}^3} \\ &= \text{coeff. of } \frac{t^2 + 2ait - a^2}{\{-a^2 + 2ait + t^2 + a^2\}^3} \\ &= \text{coeff. of } \frac{1}{(2ait)^3} (t^2 + 2ait - a^2) \left(1 + \frac{t}{2ai}\right)^{-3} \\ &= \text{coeff. of } \frac{-(t^2 + 2ait - a^2)}{8a^3 i^3} \left[1 - \frac{3t}{2ai} - \frac{6t^2}{4a^2} + \dots\right] \\ &= \text{coeff. of } \frac{-(t^2 + 2ait - a^2)}{8a^3 i} \left[\frac{1}{t^3} - \frac{3}{2ait^2} - \frac{6}{4a^2 t} + \dots\right] \\ &= -\frac{1}{8a^3 i} [1 - 3 + \frac{6}{4}] = \frac{1}{16a^3 i}. \end{aligned}$$

Thus (2) gives

$$\int_{-\infty}^{\infty} f(x) dx = 2\pi i \frac{1}{16a^3 i} = \frac{\pi}{8a^3},$$

$$I = \int_{-\infty}^{\infty} \frac{x^2}{(x^2 + a^2)^3} = \frac{\pi}{8a^3}.$$

$$(v) \text{ Suppose } I = \int_0^{\infty} \frac{x^2}{1+x^4} dx.$$

$$\text{Here } f(z) = \frac{z^2}{1+z^4}.$$

$zf(z) \rightarrow 0$ as $z \rightarrow \infty$, thus the condition for evaluation of an infinite integral is satisfied.

Poles of $f(z)$ are given by $1+z^4=0$,

$$z^4 = -1 = e^{i(2n+1)\pi},$$

so that

$$z = e^{i(2n+1)\pi/4}, \text{ where } n = 1, 2, 3,$$

i.e., the poles are $z = e^{i\pi/4}, e^{i3\pi/4}, e^{i5\pi/4}, e^{i7\pi/4}$ of which the first two only lie in the upper half plane, which is to be chosen as a contour consisting of a large semi-circle Γ and the part of the real axis from $-R$ to R .

Applying the theorem of residues,

$$\int_{-R}^R f(x) dx + \int_{\Gamma} f(z) dz = 2\pi i \Sigma R^+,$$

Since $\int_{\Gamma} f(z) dz \rightarrow 0$ as $z \rightarrow \infty$, we have, on proceeding to the limit as $R \rightarrow \infty$,

$$\int_{-\infty}^{\infty} f(x) dx = 2\pi i \sum R^+. \quad \dots (1)$$

To find the residues at the poles, we have $f(z) = \frac{z^2}{1+z^4}$ which is of the form $\frac{\phi(z)}{\psi(z)}$, where $\phi(z) = z^2$ and $\psi(z) = 1+z^4$.

$$\therefore \text{Residue at } z = e^{i\pi/4} \text{ is } = \frac{\phi(e^{i\pi/4})}{\psi'(e^{i\pi/4})} = \frac{e^{i\pi/2}}{4(e^{i\pi/4})^3} = \frac{1}{4} \cdot e^{-\pi i/4}$$

Similarly residue at $z = e^{i3\pi/4}$ is $= \frac{1}{4} e^{-3\pi i/4}$, so that (1) gives

$$\int_{-\infty}^{\infty} f(x) dx = 2\pi i \left[\frac{1}{4} e^{-\pi i/4} + \frac{1}{4} e^{-3\pi i/4} \right]$$

i.e., $2 \int_0^{\infty} \frac{x^2 dx}{1+x^4} = 2\pi i \cdot \frac{1}{4} \left[\cos \frac{\pi}{4} - i \sin \frac{\pi}{4} + \cos \frac{3\pi}{4} - i \sin \frac{3\pi}{4} \right]$

or $I = \int_0^{\infty} \frac{x^2 dx}{1+x^4} = \frac{\pi i}{2} \left(-2i \frac{1}{\sqrt{2}} \right) = \frac{\pi}{\sqrt{2}}$.

(vi) The given integral is $\int_0^{\infty} \frac{1-\cos x}{x^2} dx$.




Fig. 5.48

Consider $f(z) = \frac{1-e^{iz}}{z^2}$.

Choose the contour C , consisting of the upper half of a circle $|z| = R$ and the real axis from $-R$ to R , indented at $z=0$ (a pole of order 2). Let the semi-circle be denoted by Γ and the small semi-circle of indentation by γ . Let the radius of indentation be r .

By Cauchy's theorem, we have

$$\oint_C f(z) dz = \int_{-R}^{-r} f(x) dx + \int_{\gamma} f(z) dz + \int_r^R f(x) dx + \int_{\Gamma} f(z) dz = 0. \quad \dots (1)$$

$$\begin{aligned} \text{Now } \left| \int_{\Gamma} f(z) dz \right| &= \left| \int_{\Gamma} \frac{1-e^{iz}}{z^2} dz \right| \\ &\leq \int_{\Gamma} \frac{|1-e^{iz}|}{|z|^2} |dz| \\ &\leq \int_0^{\pi} \frac{|1-e^{iRe^{i\theta}}|}{|R^2 e^{2i\theta}|} |i| |Re^{i\theta}| |d\theta| \\ &\quad \because z = Re^{i\theta}, dz = iRe^{i\theta} d\theta \\ &\leq \int_0^{\pi} \frac{1+e^{-R \sin \theta}}{R} d\theta, \text{ which } \rightarrow 0 \text{ as } R \rightarrow \infty. \end{aligned}$$

$$\therefore \int_{\Gamma} f(z) dz = 0$$

$$\begin{aligned} \text{and } \int_{\gamma} f(z) dz &= \int_{\pi}^0 \frac{1-eire^{i\theta}}{r^2 e^{2i\theta}} - ire^{i\theta} d\theta \quad \because z = re^{i\theta} \\ &= \int_{\pi}^0 \left\{ 1 - 1 - ire^{i\theta} - \frac{i^2 r^2}{i^2} e^{2i\theta} - \dots \right\} \cdot \frac{i}{r} e^{-i\theta} d\theta \\ &= \int_{\pi}^0 [1 + \text{terms containing powers of } r] d\theta, \end{aligned}$$

$$\lim_{r \rightarrow 0} \int_{\gamma} f(z) dz = \int_{\pi}^0 d\theta = -\pi.$$

Hence when $R \rightarrow \infty$, $r \rightarrow 0$, (1) becomes

$$\int_{-\infty}^{\infty} f(x) dx - \pi + \int_0^{\infty} f(x) dx = 0,$$

$$\text{i.e., } \int_{-\infty}^{\infty} f(x) dx = \pi$$

$$\text{or } \int_{-\infty}^{\infty} \frac{1-e^{ix}}{x^2} dx = \pi.$$

Equating real parts on either side, we get

$$\int_{-\infty}^{\infty} \frac{1-\cos x}{x^2} dx = \pi$$

$$\text{or } 2 \int_0^{\infty} \frac{1-\cos x}{x^2} dx = \pi,$$

$$\text{i.e., } \int_0^{\infty} \frac{1-\cos x}{x^2} dx = \frac{\pi}{2}.$$

(vii) The given integral is $\int_{-\infty}^{\infty} (x^2+a^2)^{-2} (x^2+b^2)^{-1} dx$.

$$\text{Let } f(x) = \frac{1}{(z^2+a^2)^2 (z^2+b^2)}$$

The poles are $z = \pm ai$ (of order 2) and $\pm bi$ (simple poles).

If we choose the contour C , consisting of the upper half of a large circle $|z| = R$ and the real axis from $-R$ to R , then the only poles lying within the contour are

$+ai$ (of order 2) and $+bi$ (simple pole).

Here $zf(z) \rightarrow 0$ as $z \rightarrow \infty$.

Hence by Cauchy's theorem,

$$\oint_C f(z) dz = \int_{-R}^R f(x) dx + \int_{\Gamma} f(z) dz = 2\pi i \sum R^+, \quad \dots (1)$$

where $\sum R^+$ denotes the sum of the residues and Γ denotes the upper half of the semi-circle.

$$\begin{aligned} \text{Now } \left| \int_{\Gamma} f(z) dz \right| &= \left| \int_{\Gamma} \frac{dz}{(z^2+a^2)^2 (z^2+b^2)} \right| \\ &\leq \int_{\Gamma} \frac{|dz|}{|z^2+a^2|^2 |z^2+b^2|} \\ &\leq 0 \text{ as } z \rightarrow \infty, \end{aligned}$$

so that $\int_{\Gamma} f(z) dz = 0$.

$\therefore$ as $R \rightarrow \infty$, (1) reduces to

$$\int_{-\infty}^{\infty} f(x) dx = 2\pi i \Sigma R^+.$$

Residue at $z=ai$ (a pole of order 2)

=coeff. of $\frac{1}{t}$ in $f(ai+t)$

$$= \frac{1}{\{(t+ai)^2 + a^2\}^2 \{(t+ai)^2 + b^2\}}$$

$$= \frac{1}{t^2 (2ai+t)^2 (b^2 - a^2 + 2ait + t^2)}$$

$$= \frac{1}{-4a^2 t^2 (b^2 - a^2)} \left\{ 1 + \frac{t}{2ai} \right\}^{-2} \left\{ 1 + \frac{2ait + t^2}{b^2 - a^2} \right\}^{-1}$$

$$= \frac{1}{-4a^2 t^2 (b^2 - a^2)} \left\{ 1 - \frac{t}{ai} + \dots \right\} \left\{ 1 - \frac{2ait}{b^2 - a^2} - \dots \right\}$$

$$= \frac{1}{4a^2 (b^2 - a^2)} \left\{ \frac{1}{ai} + \frac{2ai}{b^2 - a^2} \right\} = \frac{b^2 - 3a^2}{4a^3 i (b^2 - a^2)^2}$$

Residue at $z=bi$ (simple pole) = $\lim_{z \rightarrow bi} (z - bi) f(z)$

$$= \lim_{z \rightarrow bi} (z - bi) \frac{1}{(z^2 + a^2)^2 (z + bi) (z - bi)}$$

$$= \frac{1}{2bi (b^2 - a^2)^2}.$$

Hence from (2),

$$\int_{-\infty}^{\infty} f(x) dx = 2\pi i \left[\frac{b^2 - 3a^2}{4a^3 i (b^2 - a^2)^2} + \frac{1}{2bi (b^2 - a^2)^2} \right]$$

$$= \frac{\pi}{2a^3 b (b^2 - a^2)^2} [b^3 - 3a^2 b + 2a^3]$$

$$= \frac{\pi (b - a)^2 (b + 2a)}{2a^3 b (b - a)^2 (b + a)^2}$$

$$= \frac{\pi (b + 2a)}{2a^3 b (b + a)^2}.$$




Fig. 5.49

5.21. ANALYTIC CONTINUATION

If there exist two functions $f_1(z)$ and $f_2(z)$, such that they are analytic (regular) in domains D_1 and D_2 respectively and that D_1 and D_2 have a common part, throughout which $f_1(z) = f_2(z)$, then the aggregate of values of $f_1(z)$ and $f_2(z)$ at the interior points of D_1 or D_2 , can be regarded as a single regular function

(say) $F(z)$. It is obvious that $F(z)$ is regular in the common part say of the two domains and $F(z) = f_1(z)$ in domain D_1 and $F(z) = f_2(z)$ in domain D_2 . We thus regard the function $f_2(z)$ as one, extending the domain in which $f_1(z)$ is defined and so it is called an **Analytic continuation** of $f_1(z)$.

The method of analytic continuation. Its standard method is the method of power series, which can be summarised as below.

Suppose that there is a point $P(z_0)$ in the neighbourhood of which $f(z)$ is analytic; then the function $f(z)$ can be expanded by Taylor's theorem, in a series of ascending powers of $(z - z_0)$, the coefficients of which involve the successive derivatives of $f(z)$ at z_0 .

Let there be a singularity S of $f(z)$ which is nearest to P . Then a circle of centre P and radius PS is the circle of convergence within which the Taylor's expansion is valid. If we now take any point P' (not on PS) within this circle, then we can find the values of $f(z)$ and all its derivatives at P' , from the series by applying the method of term by term differentiation. We thus find the Taylor's series for $f(z)$ with P' as origin and this series will define a function which is regular in the circle whose centre is P' . Such a circle will extend as far as the singularity of the function defined by the new series, which is nearest to P' and this may or may not be S . In either case the new circle of convergence may lie partly outside the old circle and for points in the region which is included in the new circle but not in the old one, so that the new series may be utilised in defining the values of $f(z)$ while the old series failed to do so.




Fig. 5.50




Fig. 5.51

In a similar manner, we can take another point P'' in the region for which the values of the function are known and form the Taylor's series with P'' as origin which will, in general, still further extend the domain of definition of the function and so on and so forth.

By this method of continuation, starting from a representation of a function by any one power series, we can find any number of other power series, which between them define the value of the function at all points of a domain, any point of which can be reached from P without passing through a singularity of the function $f(z)$.

It is easy to show that continuation by two different paths PLN and PMN gives the same power series provided the function is analytic and has no singularity inside the closed curve $PLNMP$. To show it, let us suppose that $S, S', S'', \dots$ be the power series with $P, P', P'', \dots$ as origin. Then $S' = S''$ (for, each is equal to S) over a certain domain which contains P' , when P'' is taken sufficiently near

to P' , and therefore S' will be the continuation of S'' . Continuing this process we can deform the path PLN into PMN provided no singular point lies inside the path $PLNMP$.

Note. Weierstrass defined an Analytic function of z as one power series together with all the other power series derivable from it by analytic continuation.

An important remark. There must be at least one singularity of the analytic function on the circle of convergence C_0 of the power series $\sum_{n=0}^{\infty} a_n (z-z_0)^n$.

Assuming that there is no such singularity, we can construct by the method of continuation, a function equal of $f(z)$ within C_0 and regular in a larger concentric circle Γ_0 . The expansion of this function in Taylor series in powers of $(z-z_0)$ would then converge everywhere within the large circle Γ_0 . But this is not possible, since the series would be the original series which has got C_0 as its circle of convergence. Let there be a point z_1 within C_0 and let C_1 be the circle of convergence of the power series,




Fig. 5.52

$$\sum_{n=0}^{\infty} A_n (z-z_1)^n,$$

where

$$A_n = \frac{f^n(z_1)}{n!} \quad (\text{by Taylor's expansion}).$$

Let Γ_1 be the circle of centre z_1 , touching the circle C_0 internally. Then the new power series defined by (1) is certainly convergent within Γ_1 and has the sum $f(z)$ there.

Since the radius of C_1 cannot be less than that of Γ_1 , there are these possibilities:

(i) C_1 has a larger radius, then Γ_1 , in which case C_1 lies partly outside C_0 and the new power series provides an analytic continuation of $f(z)$. Then taking a point z_2 within C_1 and outside C_0 , the process can be repeated.

(ii) C_0 is a natural boundary of $f(z)$. [Note. A closed curve is called a natural boundary of a function if the function is such that it cannot be analytically continued to any point of the same]. In this case we cannot continue $f(z)$ outside C_0 and the circle C_1 touches C_0 internally, wherever the point z lies within C_0 .

(iii) C_1 may touch C_0 internally, though C_0 is not a natural boundary of $f(z)$. In this case the point of contact of C_0 and C_1 is a singularity of the analytic function which has been found by continuation of the original power series; for there is necessarily one singularity on C_1 and this cannot be within C_0 .

Problem 29. Show that the two power series

$$z + \frac{1}{2}z^2 + \frac{1}{3}z^3 + \dots \quad \dots (1)$$

$$i\pi - (z-2) + \frac{1}{2}(z-2)^2 - \dots \quad \dots (2)$$

have no common region of convergence, but they are analytic continuations of the same function.

Here the series (1) defines the function $-\log(1-z)$, whose circle of convergence is $|z|=1$, i.e., a circle with centre $z=0$ and radius 1. The only singularity is $z=+1$.

The series (2) defines the function $i\pi - \log[1+(z-2)]$; where $|z-2| \leq 1$ is a circle with centre $z=2$ and radius 1. It touches the first circle externally as shown in the adjoining figure.




Fig. 5.53

The function $i\pi - \log[1+(z-2)] = i\pi - \log(z-1)$

$$= i\pi - \log[-(1-z)]$$

$$= i\pi - \log(-1) - \log(1-z)$$

$$= i\pi - \log e^{\pi i} - \log(1-z)$$

$$= i\pi - \pi i - \log(1-z)$$

$$= -\log(1-z).$$

It is clear that the two functions defined by the given power series have no common region of convergence, but they are analytic continuations of the same function $-\log(1-z)$.

Problem 30. Show that the series

$$\frac{1}{a} + \frac{z}{a^2} + \frac{z^2}{a^3} + \frac{z^3}{a^4} + \dots$$

represents the function which can be continued analytically outside the circle of convergence.

Let the given series define a function $f(z)$, i.e.

$$f(z) = \frac{1}{a} + \frac{z}{a^2} + \frac{z^2}{a^3} + \frac{z^3}{a^4} + \dots$$

$$= \frac{1}{a} \left[\text{being an infinite geometric series if } \left| \frac{z}{a} \right| > 1 \right]$$

$$= \frac{1}{a-z},$$

only for points within the circle of convergence $|z| = |a|$

If we consider a series

$$\frac{1}{a-b} + \frac{z-a}{(a-b)^2} + \frac{(z-b)^2}{(n-b)^3} + \dots \text{ where } \frac{b}{a} \text{ is real,}$$

then

$$f(z) = \frac{1}{a-b} \left[1 + \frac{z-b}{a-b} + \left(\frac{z-b}{a-b} \right)^2 + \dots \right]$$

$$= \frac{1}{a-b} \left[1 - \frac{z-b}{a-b} \right]^{-1}$$

provided the circle of convergence say C_1 is $|z-b| = |a-b|$, when a and b both are real and positive and also $0 < b < a$, the circle of convergence C_1 of the second series touches the first circle (say) internally at its only singularity $z=a$. It follows that there is no analytic continuation (see figure 5.54).




Fig. 5.54




Fig. 5.55

In case $\frac{b}{a}$ is not real and $|b| < |a|$ and positive, the second series converges at points inside a circle which is partly inside and partly outside the circle $|z|=a$. In fact the two series represent the same function $\frac{1}{a-z}$ at points outside the circle $|z|=|a|$ and hence can be continued analytically as shown in Fig. 5.55.

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 31. Evaluate by the method of contour integration the integral

$$\int_0^{2\pi} \frac{d\theta}{1 + \epsilon \cos \theta}$$

using the transformation.

Ans. $\frac{2\pi}{\sqrt{1-\epsilon^2}}$

Proceed as in Problem 20 (i).

Problem 32. (a) Prove that the n -th derivative of a function $f(z)$ of the complex variable z in the domain of its analyticity is given by

$$f^{(n)}(z) = \frac{n!}{2\pi i} \oint \frac{f(z') dz'}{(z'-z)^{n+1}}$$

where the contour of integration lies within the domain.

(b) using the above formula, show that

$$\left(\frac{d^n e^{-z}}{dz^n} \right)_{z=1} = (-1)^n e^{-1}$$

(Agra, 1968)

Problem 33. Obtain the Cauchy-Riemann condition for a function $f(z)$ to be analytic function.

Problem 34. Evaluate by the theorem of residues

$$\int_{-\infty}^{\infty} \frac{\cos x}{a^2 + x^2} dx, a > 0.$$

See Problem 20 (iii).

Ans. $\frac{\pi}{ae^a}$.

Problem 35. Apply calculus of residue to evaluate

$$\int_0^{\infty} \frac{x^{a-1}}{1+x} dx$$

Ans. $\pi \cot \pi a$

See Problem 23.

Problem 36. State and Prove Cauchy's residue theorem. Evaluate the following integrals by the method of contour integration

(a) $\int_0^{\infty} \frac{\sin x}{\sqrt{x}} dx$

(Agra, 1972, 74)

(b) $\int_0^{\infty} \frac{x^4 dx}{x^6+1}$

(Agra, 1972)

(a) Take $f(z) = \frac{e^{iz}}{\sqrt{z}}$

Choose the contour C as consisting of a positive quadrant of a large circle $|z|=R$ with its bounding radii as two axes, indented at $z=0$, as shown in Fig. 5.56. Let r be the radius of indentation at $z=0$.

$f(z)$ has the only pole $z=0$, of order one and it does not lie inside the contour C indented at $z=0$.

Cauchy's theorem gives

$$\int_r^R f(x) dx + \int_{\Gamma} f(z) dz + \int_R^r f(iy) (idy) + \int_{\gamma} f(z) dz = 0$$

.. (1)

Here $\left| \int_{\Gamma} f(z) dz \right| \leq \int_{\Gamma} |f(z)| |dz| = \int_{\Gamma} \frac{|e^{iz}|}{|\sqrt{z}|} |dz|$

$$\leq \int_0^{\pi/2} \frac{e^{-R \sin \theta}}{\sqrt{R}} R d\theta, \text{ taking } z = Re^{i\theta} \text{ so that } |dz| = R d\theta$$

$$< \frac{R}{\sqrt{R}} \int_0^{\pi/2} e^{-2R\theta/\pi} d\theta \text{ since by Jordan's inequality}$$

$$\text{if } 0 < \theta \leq \frac{\pi}{2} \text{ then } \frac{\sin \theta}{\theta} > \frac{2}{\pi}$$

$$\leq \frac{\pi}{2\sqrt{R}} \left[e^{-2R\theta/\pi} \right]_0^{\pi/2}$$

$$\leq \frac{\pi}{2\sqrt{R}} (1 - e^{-R}) \rightarrow 0 \text{ as } R \rightarrow \infty$$

Also residue of $f(z)$ at $z=0$ being $= \lim_{z \rightarrow 0} \frac{z e^{iz}}{\sqrt{z}} = 0$, by §5.16 we have

$\lim_{R \rightarrow \infty} \int_{\gamma} f(z) dz = -i \left(\frac{\pi}{2} = 0 \right)$. $0=0$, negative sign being taken as γ is traversed in clockwise sense.




Fig. 5.56

Hence as $R \rightarrow \infty$, $r \rightarrow 0$, (1) reduces to

$$\int_0^{\infty} f(x) dx + \int_0^{\infty} f(iy) i dy = 0$$

$$\text{i.e. } \int_0^{\infty} \frac{e^{ix}}{\sqrt{x}} dx + \int_0^{\infty} \frac{e^{i(iy)}}{\sqrt{iy}} i dy = 0$$

$$\text{or } \int_0^{\infty} \frac{e^{ix}}{\sqrt{x}} dx = \sqrt{i} \int_0^{\infty} \frac{e^{-y}}{\sqrt{y}} dy. \text{ Put } \sqrt{y} = \phi$$

$$= 2\sqrt{i} \int_0^{\infty} e^{-\phi^2} d\phi$$

$$= 2\sqrt{i} \frac{\sqrt{\pi}}{2} \quad (\text{See Beta and Gamma functions Equation (3)})$$

$$= \sqrt{\pi} \left(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2} \right)^{1/2}$$

$$= \sqrt{\pi} \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right)$$

$$= \sqrt{\frac{\pi}{2}} [1 + i]$$

Equating real and imaginary parts, we get

$$\int_0^{\infty} \frac{\cos x}{\sqrt{x}} dx = \frac{\sqrt{\pi}}{2} \text{ and } \int_0^{\infty} \frac{\sin x}{\sqrt{x}} dx = \frac{\sqrt{\pi}}{2}$$

Note. Similar procedure will show that $\int_0^{\infty} \frac{\sin x^2}{x} dx = \frac{\pi}{4}$.




Fig. 5.57

$\therefore$ Simple poles of $f(z)$ are

$$e^{\pi i/6}, e^{\pi i/2}, e^{5\pi i/6}, e^{7\pi i/6}, e^{3\pi i/2}, e^{11\pi i/6}$$

of which only the first three lie within the contour C .

Denoting by p , any one of these poles namely $e^{\pi i/6}, e^{\pi i/2}, e^{5\pi i/6}$, we have the

$$\text{residue of } f(z) \text{ at } z=p = \left[\frac{z^4}{\frac{d}{dz}(z^6+1)} \right]_{\text{at } z=p} \quad \text{by §5.17 (i)}$$

$$= \left[\frac{z^4}{6z^5} \right]_{z=p} = \frac{1}{6p} \times \frac{p^5}{p^5}$$

$$= \frac{p^5}{-6} \quad \because p^6 = -1$$

$\therefore$ Sum of residues $= \Sigma R +$

$$= -\frac{1}{6} \left[e^{5\pi i/6} + e^{5\pi i/2} + e^{25\pi i/6} \right]$$

$$(b) \text{ Take } f(z) = \frac{z^4 dz}{z^6+1}$$

and choose the contour C consisting of a large semi-circle Γ of radius R in the upper half plane and the real axis from $-R$ to R .

Poles of $f(z)$ are given by $z^6+1=0$

$$\begin{aligned} \text{i.e. } z &= (-1)^{1/6} \\ &= \{ \cos (2r+1)\pi + i \sin (2r+1)\pi \}^{1/6} \\ &= e^{i(2r+1)\pi/6} \text{ where } r=0, 1, 2, 3, 4, 5. \end{aligned}$$

$$= -\frac{1}{6} \left[e^{5\pi i/6} + e^{2\pi i} e^{\pi i/2} + e^{4\pi i} e^{\pi i/6} \right]$$

$$= -\frac{1}{6} \left[e^{5\pi i/6} + e^{\pi i/2} + e^{\pi i/6} \right]$$

$$\because e^{2\pi i} = e^{4\pi i} = 1$$

$$= -\frac{1}{6} \left[\left(-\frac{\sqrt{3}}{2} + i\frac{1}{2} \right) + i + \left(\frac{\sqrt{3}}{2} + i\frac{1}{2} \right) \right]$$

$$= -\frac{2i}{6} = -\frac{i}{3}.$$

Now by Cauchy's theorem,

$$\int_C f(z) dz = \int_{-R}^R f(x) dx + \int_{\Gamma} f(z) dz = 2\pi i \Sigma R +$$

$$\text{where } \left| \int_{\Gamma} f(z) dz \right| \leq \int_{\Gamma} |f(z)| |dz|$$

$$\leq \int_{\Gamma} \left| \frac{z^4}{z^6+1} \right| |dz|. \text{ Put } z = Re^{i\theta} \text{ so } |dz| = R d\theta$$

$$\leq \int_{\Gamma} \frac{|z^4|}{|z^6+1|} |dz|$$

$$\because |z^6| - 3 < |z^6+1|$$

$$\leq \int_0^{\pi} \frac{R^5 d\theta}{R^6-1} = \frac{R^5 \pi}{R^6-1} \rightarrow 0 \text{ as } R \rightarrow \infty$$

Hence when $R \rightarrow \infty$, we have

$$\int_{-\infty}^{\infty} f(x) dx = 2\pi i \times \left(-\frac{i}{3} \right) = \frac{2\pi}{3}$$

$$\text{i.e. } \int_0^{\infty} f(x) dx = \int_0^{\infty} \frac{x^4 dx}{x^6+1} = \frac{\pi}{3}.$$

Problem 37. Derive the Cauchy-Riemann conditions for a function of a complex variable to be analytic. Test if the functions $\frac{1}{z}$ and $|z|^2$ are analytic.

Prove that if a function is analytic on and inside a closed contour its integral over the contour must vanish. (Agra, 1973)

See § 5.6 for the first part.

It is easy to show that $f(z) = \frac{1}{z}$ (giving $f'(z) = -\frac{1}{z^2}$, $z \neq 0$) is analytic at every point except at $z=0$ where it is not continuous and so $f'(0)$ does not exist ($z=0$ is a pole of $f(z)$).

The function $f(z) = |z|^2$ is not analytic at any point since $f'(z)$ exists only at the point $z=0$ not throughout any neighbourhood. It may be shown as below

$$f(z) = |z|^2 = x^2 + y^2 \quad \because zy = x + iy$$

If $f(z) = u(x, y) + iv(x, y)$, then it is clear on comparison that

$$u(x, y) = x^2 + y^2, v(x, y) = 0$$

$$\therefore \frac{\partial u}{\partial x} = 2x, \frac{\partial u}{\partial y} = 2y, \frac{\partial v}{\partial x} = 0 = \frac{\partial v}{\partial y}$$

Cauchy-Riemann equations are not satisfied except at $z=0$.

For second part see §5.9.

Problem 38. Defining a Meromorphic Function as one which is analytic in a region except at a finite number of poles, if $f(z)$ be an analytic function within and on a closed contour except at a finite number of poles and is not zero on C , then prove that

$$\frac{1}{2\pi i} \int_C \frac{f'(z)}{f(z)} dz = N - P$$

where N is the number of zeros and P is the number of poles inside C (pole of order m being counted m times).

Hence or otherwise deduce the Principle of Argument i.e. $f(z)$ is analytic in C then $P=0$, and

$$N = \frac{1}{2\pi} \Delta_C \arg f(z)$$

Δ_C being the variation of $\log f(z)$ round the closed contour C .

Taking $z=a$ as a zero of order n , we have

$$f(z) = (z-a)^n \phi(z)$$

where $\phi(z)$ is analytic and not zero in C .

$$\text{Thus, } \frac{f'(z)}{f(z)} = \frac{n}{z-a} + \frac{\phi'(z)}{\phi(z)}$$

Since $\phi(z)$ is analytic and so is $\phi'(z)$, therefore $\frac{\phi'(z)}{\phi(z)}$ is analytic at $z=a$, it follows that $\frac{f'(z)}{f(z)}$ has a simple pole at $z=a$ with its residue n .

Similarly if $z=b$ is a pole of order p then $\frac{f'(z)}{f(z)}$ has a simple pole at $z=b$ with its residue $-p$ since then

$$f(z) = (z-b)^{-p} \psi(z)$$

Hence by Cauchy's theorem

$$\begin{aligned} \frac{1}{2\pi i} \int_C \frac{f'(z)}{f(z)} dz &= \text{Sum of all residues within } C \\ &= \Sigma n - \Sigma p \\ &= N - P \end{aligned}$$

Now to deduce the principle of argument, $f(z)$ being analytic in C , we have $P=0$.

Then (1) reduces to

$$\frac{1}{2\pi i} \int_C \frac{f'(z)}{f(z)} dz = N$$

$$\text{or } N = \frac{1}{2\pi i} [\log f(z)]_C$$

$$\begin{aligned} &= \frac{1}{2\pi i} \Delta_C \log f(z) \text{ where } \Delta_C \text{ is given to be the variation of } \log f(z) \text{ round the closed contour } C \\ &= \frac{1}{2\pi i} \Delta_C \{ \log |f(z)| + i \arg f(z) \} \end{aligned}$$

Equating real parts on either side we get

$$N = \frac{1}{2\pi} \Delta_C \arg f(z).$$

Problem 39. (Rouche's theorem) If $f(z)$ and $g(z)$ are analytic within and on a closed contour C and $|g(z)| < |f(z)|$ on C , then $f(z)$ and $f(z)+g(z)$ have same number of zeros inside C .

$\therefore |g(z)| < |f(z)|$ on $C \therefore \left| \frac{g(z)}{f(z)} \right| < 1$ on C which follows that $|f(z)| \neq 0$ otherwise this inequality will not hold.

As such $|f(z)+g(z)| > |f(z)| - |g(z)| \neq 0$ by properties of moduli.

So $|f(z)+g(z)| \neq 0$ implies that neither $f(z)$ nor $f(z)+g(z)$ has a zero on C .

Taking N and N' as number of zeros of $f(z)$ and $f(z)+g(z)$, we have by the principle of argument given in Problem 38,

$$N = \frac{1}{2\pi} \Delta_C \arg f \text{ and } N' = \frac{1}{2\pi} \Delta_C \arg (f+g)$$

$$\text{So that } N' - N = \frac{1}{2\pi} \Delta_C \arg (f+g) - \frac{1}{2\pi} \Delta_C \arg f$$

$$= \frac{1}{2\pi} \left[\Delta_C \arg f + \Delta_C \arg \left(1 + \frac{g}{f} \right) \right] - \frac{1}{2\pi} \Delta_C \arg f$$

$$= \frac{1}{2\pi} \Delta_C \arg \left(1 + \frac{g}{f} \right) \text{ by the properties of arguments.}$$

$$\text{If } 1 + \frac{g}{f} = Q, \text{ then } |Q-1| = \left| \frac{g}{f} \right| < 1$$

$$\therefore |g| < |f|$$

showing that Q is always an internal point of the circle with centre $Q=1$ and radius unity. This circle being wholly to the right of imaginary axis, the join of any point of this circle to the origin makes an angle θ with x -axis such that $0 \leq \theta < \frac{\pi}{2}$. As such putting $Q = Re^{i\theta}$ we have $\arg \left(1 + \frac{g}{f} \right) = \arg Q = \theta$ which

returns to its initial value when z describes C and hence

$$\Delta_C \arg \left(1 + \frac{g}{f} \right) = 0$$

Hence, $N' - N = 0$ or $N = N'$

which proves the proposition.

Problem 40. Evaluate the following integrals by applying the method of contour integration.

$$(i) \int_0^\infty \frac{x \sin x}{x^2 + a^2} dx, a > 0$$

$$\text{Ans. } \frac{\pi}{2} e^{-a}$$

$$(ii) \int_0^\infty \frac{\sinh ax}{\sinh \pi x} dx, -\pi < a < \pi,$$

$$\text{Ans. } \frac{1}{2} \tan \frac{a}{2}$$

$$(iii) \int_0^\infty \frac{\sin mx}{x^4 + a^4} dx$$

$$\text{Ans. } \frac{\pi}{2a^2} e^{-\frac{me}{\sqrt{2}}} \sin \frac{ma}{\sqrt{2}}$$

$$(iv) \int_{-\infty}^\infty \frac{a \cos x + x \sin x}{x^2 + a^2} dx$$

$$\text{Ans. } 2\pi e^{-a}$$

$$\text{[Hint: take } f(z) = \frac{e^{iz}}{z-ia} \text{]}$$

$$(v) \int_0^\infty \frac{x dx}{\sinh x}$$

$$\text{Ans. } \frac{\pi^2}{4}$$

$$(vi) \int_0^\pi \frac{a d\theta}{a^2 + \sin^2 \theta}, a > 0$$

$$\text{Ans. } \frac{\pi}{\sqrt{1+a^2}}$$

$$(vii) \int_0^\infty \frac{\cos 2ax - \cos 2bx}{x^2} dx, a \geq 0, b \geq 0,$$

$$\text{Ans. } \pi (b-a)$$

$$(viii) \int_0^\infty e^{-x^2} \cos 2\alpha \cos (x^2 \sin 2\alpha) dx, \alpha \leq 4$$

$$\text{Ans. } \frac{\sqrt{\pi}}{2} \cos \alpha$$

$$\text{and } \int_0^\infty e^{-x^2} \cos 2\alpha \sin (x^2 \sin 2\alpha) dx, \alpha \leq \pi/4$$

$$\text{Ans. } \frac{\sqrt{\pi}}{2} \sin \alpha$$

$$(ix) \int_0^\infty e^{-x^2} \cos 2ax dx \text{ and } \int_0^\infty e^{-x^2} \sin 2ax dx$$

$$\text{Ans. } \frac{e^{-a^2}}{2} \sqrt{\pi}; e^{-a^2} \int_0^a e^{y^2} dy$$

$$(x) \int_{-\infty}^{\infty} \frac{\sin \pi x}{x(1-x^2)} dx,$$

$$(xi) \int_0^{\infty} \frac{e^{ax}}{1+e^x} dx, (0 < a < f) \text{ [see Problem 27(i)],}$$

Problem 41. The function $w(z)$ is analytic in a region R , except at two poles A and B . Find: $\oint_C w(z) dz$ where C is a contour in the region R (i) enclosing A and B (ii) enclosing B only.

Problem 42. Evaluate the following integrals:

$$(i) \int_0^{2\pi} e^{\cos \theta} \cos(n\theta - \sin \theta) d\theta,$$

$$(ii) \int_{-\infty}^{\infty} \frac{\cos x}{x^2 + a^2} dx, \quad (iii) \int_{-\infty}^{\infty} \frac{\sin x}{x} dx$$

$$\text{Ans. (i) } \frac{2\pi}{n}, \quad (ii) \frac{\pi}{2a} e^{-a}, \quad (iii) \pi.$$

Problem 43. Evaluate the following integrals:

$$(a) \int_0^{2\pi} (a + b \cos \phi)^{-1} \sin^2 \phi d\phi, (a > b > 0)$$

$$(b) \int_0^{\infty} (1+x^2)^{-1} \log(1+x^2) dx \text{ [see Problem 21 (vi)]}$$

$$\text{Ans. (a) } \frac{2\pi}{b^2} \left\{ a - \sqrt{a^2 - b^2} \right\}, (b) \pi \log 2$$

Problem 44. (a) Derive the Cauchy-Riemann condition for complex function to be analytic. Are the conditions sufficient?

(b) Evaluate the following integrals:

$$(i) \int_0^{\infty} \frac{\sin x}{x} dx \quad (ii) \int_{-\infty}^{\infty} \frac{dx}{1+x^2}$$

$$\text{Ans. (a) yes, (b) (i) } \frac{\pi}{2}, \quad (ii) \pi.$$

Problem 45. State and prove Cauchy's Residue theorem for a complex function. Explain how it is extended for the case of an isolated first order pole lying on the contour of integration. Using this theorem show that

$$\int_{-\infty}^{\infty} \frac{e^{ax}}{1+e^x} dx = \frac{\pi}{\sin \pi a}; \text{ where } 0 < a < 1$$

see § 5.15 and Problem 27(i).

Problem 46. State and prove Cauchy's residue theorem for a complex function and using it evaluate the following integrals.

$$(i) \int_0^{\infty} \frac{\log(1+x^2)}{1+x^2} dx; \quad (ii) \int_0^{2\pi} \frac{d\theta}{H \epsilon \cos \theta}$$

See § 5.15; (i) see Problem 21 (vi); (ii) see Problem 31.

CHAPTER 6

BETA, GAMMA AND ERROR FUNCTIONS

6.1. DEFINITIONS

Under the study of Definite Integrals, we come across two very important integrals known as Eulerian Integrals which are of the type

$$\int_0^1 x^{m-1} (1-x)^{n-1} dx \text{ and } \int_0^{\infty} e^{-x} x^{n-1} dx,$$

where the quantities m and n are supposed to be positive. These integrals are so fundamental and hold such an important place that they are widely applied in different branches of mathematical analysis like mechanics and physics etc.

The first Eulerian integral is generally known as *Beta Function* and defined as $\beta(m, n) \equiv \int_0^1 x^{m-1} (1-x)^{n-1} dx$ where m and n are positive.

The second Eulerian integral is known as *Gamma Function* and is defined as $\Gamma(n) = \int_0^{\infty} e^{-x} x^{n-1} dx$, where n is positive. (Agra, 1961)

6.2. FUNDAMENTAL PROPERTY OF GAMMA FUNCTIONS

$$\Gamma(n+1) = n\Gamma(n).$$

In order to prove this relation, let us consider the integral

$$\int_0^{\infty} e^{-x} x^n dx = \Gamma(n+1).$$

Integrating it by parts taking e^{-x} as second function, we get

$$\begin{aligned} \int_0^{\infty} e^{-x} x^n dx &= \left[-e^{-x} x^n \right]_0^{\infty} - n \int_0^{\infty} -e^{-x} x^{n-1} dx \\ &= n \int_0^{\infty} e^{-x} x^{n-1} dx, \\ \therefore \Gamma(n+1) &= n\Gamma(n) \end{aligned} \quad \left\{ \begin{array}{l} \text{since } \frac{x^n}{e^x} \text{ vanishes for both} \\ \text{the limits as } \lim_{x \rightarrow 0} \frac{x^n}{e^x} = 0 \text{ and} \\ \lim_{x \rightarrow \infty} \frac{x^n}{e^x} = \lim_{x \rightarrow \infty} \frac{x^n}{1+x+\frac{x^2}{2}+\dots} = 0. \end{array} \right. \quad (1)$$

From (1) it is evident that if the value of $\Gamma(n)$ is known for n bet-

ween two successive positive integers, then the value Γn for any positive value of n can be determined by the successive application of (1).

Now (1) can be written as

$$\Gamma(n) = \frac{\Gamma(n+1)}{n} \quad \dots (2)$$

If $-1 < n < 0$ then (2) gives Γn , since $n+1$ is positive. As such the value of Γn may be determined if $-2 < n < -1$ since then $\Gamma(n+1)$ on the R.H.S. of (2) is known. Similarly Γn may be determined when $-3 < n < -2$ and so on so forth.

Hence $\Gamma n = \int_0^\infty e^{-x} x^{n-1} dx = \frac{\Gamma(n+1)}{n}$ define Γn completely for all value of n except $n=0, -1, -2, -3, \dots$

Now replacing n by $n-1$ in (1) we get

$$\Gamma n = (n-1) \Gamma(n-1)$$

Similarly $\Gamma(n-1) = (n-2) \Gamma(n-2)$ etc.

Hence (1) yields

$$\Gamma(n+1) = n(n-1)(n-2)\dots 3.2.1 \Gamma(1)$$

$$\text{But by definition } \Gamma(1) = \int_0^\infty e^{-x} dx = \left[-e^{-x} \right]_0^\infty = 1$$

$$\therefore \Gamma(n+1) = n(n-1)(n-2)\dots 3.2.1 = \Gamma(n) \quad \text{provided } n \text{ is a positive integer}$$

Putting $n=0$ in (3) we get

$$\Gamma(1) = \Gamma(0) = 1 \quad \because \Gamma(0) = 1$$

$$\therefore \Gamma(1) = 1 \quad \dots (4)$$

Also if we put $n=0$ in (2), then we find

$$\Gamma(0) = \frac{\Gamma(1)}{0} = \infty \quad \dots (5)$$

By repeated application of (2), it may be shown that the gamma function becomes infinite when x is zero or any negative integer i.e.

$$\Gamma(-n) = \infty \quad \dots (6)$$

when $n=0$ or a positive integer.

But the function has finite value for negative values of n which are not integer.

Note. Gauss's Pi function in terms of gamma function is defined by

$$\Pi(n) = \Gamma(n+1) = \Gamma\left(\frac{n}{2}\right) \quad \dots (7)$$

6.3. THE VALUE OF $\Gamma\left(\frac{1}{2}\right)$ AND GRAPH OF THE GAMMA FUNCTION

We have by definition

$$\Gamma n = \int_0^\infty e^{-x} x^{n-1} dx, \quad n > 0$$

Putting $x = \phi^2$ i.e. $dx = 2\phi d\phi$, we get

$$\Gamma(n) = 2 \int_0^\infty \phi^{2n-1} e^{-\phi^2} d\phi$$

when $n = \frac{1}{2}$, this yields,

$$\Gamma\left(\frac{1}{2}\right) = 2 \int_0^\infty e^{-\phi^2} d\phi \quad \dots (1)$$

$$\text{Suppose } I = \int_0^\infty e^{-\phi^2} d\phi$$

Putting $\phi = \lambda\psi$ so that $d\phi = \lambda d\psi$

$$\text{We have } I = \int_0^\infty e^{-\lambda^2\psi^2} \cdot \lambda d\psi$$

Multiplying both sides by $e^{-\lambda^2}$, we find

$$I \cdot e^{-\lambda^2} = \int_0^\infty e^{-\lambda^2(1+\psi^2)} \lambda d\psi$$

Integrating both sides w.r.t. λ within the limits 0 to ∞ ,

$$I \int_0^\infty e^{-\lambda^2} d\lambda = \int_0^\infty \int_0^\infty e^{-\lambda^2(1+\psi^2)} \lambda d\lambda d\psi$$

$$\text{or } I \int_0^\infty e^{-\phi^2} d\phi = \int_0^\infty \left[-\frac{1}{2} \cdot \frac{e^{-\lambda^2(1+\psi^2)}}{1+\psi^2} \right]_0^\infty d\psi$$

$$I^2 = \frac{1}{2} \int_0^\infty \frac{d\psi}{1+\psi^2} = \frac{1}{2} \left[\tan^{-1}\psi \right]_0^\infty = \frac{1}{2} \cdot \frac{\pi}{2} = \frac{\pi}{4}$$

$$I = \int_0^\infty e^{-\phi^2} d\phi = \frac{\sqrt{\pi}}{2} \quad \dots (2)$$

From (1) and (2), we get

$$\Gamma\left(\frac{1}{2}\right) = 2 \cdot \frac{\sqrt{\pi}}{2} = \sqrt{\pi} \quad \dots (3)$$

Now putting $n = -\frac{1}{2}$ in (2) of §6.2, we find

$$\Gamma\left(-\frac{1}{2}\right) = \frac{\Gamma\left(\frac{1}{2}\right)}{-\frac{1}{2}} = -2\sqrt{\pi} \text{ by (3),} \quad \dots (4)$$

$$\text{Similarly } \Gamma\left(-\frac{3}{2}\right) = \frac{\Gamma\left(-\frac{1}{2}\right)}{-\frac{3}{2}} = -\frac{2}{3}(-2\sqrt{\pi}) = 4\frac{\sqrt{\pi}}{3} \text{ etc.} \quad \dots (5)$$

The graph of Γn may be shown as below under the definition that the function becomes continuous function of n except when $n=0$ or any negative integer.




Fig. 6.1

6.4. TRANSFORMATION OF GAMMA FUNCTION

By definition

$$\Gamma(n) = \int_0^{\infty} e^{-x} x^{n-1} dx \quad \dots (1)$$

Putting $x = \lambda y$, $dx = \lambda dy$ in (1), we get

$$\Gamma(n) = \int_0^{\infty} e^{-\lambda y} \lambda^n \cdot y^{n-1} dy$$

or

$$\frac{\Gamma(n)}{\lambda^n} = \int_0^{\infty} e^{-\lambda y} y^{n-1} dy. \quad \dots (2)$$

If we put $e^{-x} = y$ in (1), then we get

$$\Gamma(n) = - \int_1^0 y \left(\log \frac{1}{y} \right)^{n-1} \frac{1}{y} dy = \int_0^1 \left(\log \frac{1}{y} \right)^{n-1} dy \quad \dots (3)$$

$$\left[\because x = \log \frac{1}{y} \text{ and } dx = \frac{1}{1/y} \left(-\frac{1}{y^2} \right) dy = -\frac{1}{y} dy \right].$$

Again if we write $x = y^{1/n}$ in (1), we get

$$\Gamma(n) = \frac{1}{n} \int_0^{\infty} e^{-y^{1/n}} (y)^{(n-1)/n} \cdot (y)^{(1-n)/n} dy$$

$$\left[\because dx = \frac{1}{n} y^{(1-n)/n} dy \right]$$

$$= \frac{1}{n} \int_0^{\infty} e^{-y^{1/n}} dy.$$

$$n\Gamma(n) = \Gamma(n+1) = \int_0^{\infty} e^{-y^{1/n}} dy. \quad \dots (4)$$

COROLLARY. If we replace n by $\frac{1}{2}$ in (4), we find

$$\frac{1}{2}\Gamma\left(\frac{1}{2}\right) = \int_0^{\infty} e^{-y^2} dy,$$

which is the same as (1) of § 6.3.

§ 6.5. TO SHOW THAT, $\beta(m, n) = \beta(n, m)$.

By definition

$$\beta(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx.$$

Replacing x by $1-x$, we get

$$\begin{aligned} \beta(m, n) &= \int_1^0 (1-x)^{m-1} \{1-(1-x)\}^{n-1} (-dx) \\ &= \int_0^1 (x)^{n-1} (1-x)^{m-1} dx \\ &= \beta(n, m). \end{aligned}$$

§ 6.6. DIFFERENT FORMS OF BETA FUNCTION

Substituting $\frac{y}{1+y}$ for x , we have

$$\beta(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx \left[\because dx = \frac{(1+y)-y}{(1+y)^2} dy = \frac{1}{(1+y)^2} dy \right]$$

$$\text{and } 1-x = \frac{1}{1+y}$$

$$\begin{aligned} &= \int_0^{\infty} \frac{y^{m-1}}{(1+y)^{m-1}} \cdot \frac{1}{(1+y)^{n-1}} \cdot \frac{dy}{(1+y)^2} \\ &= \int_0^{\infty} \frac{y^{m-1}}{(1+y)^{m+n}} dy. \end{aligned} \quad \dots (1)$$

Also, since $\beta(m, n) = \beta(n, m)$,

$$\therefore \beta(m, n) = \int_0^{\infty} \frac{y^{n-1}}{(1+y)^{n+m}} dy. \quad \dots (2)$$

§ 6.7. TO FIND THE RELATION BETWEEN BETA AND GAMMA FUNCTIONS

$$\beta(m, n) = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}. \quad (\text{Agra, 1962, 65, 70, 71, 72, 74})$$

From equation (2) of § 6.4, we have

$$\frac{\Gamma m}{\lambda^m} = \int_0^{\infty} e^{-\lambda x} x^{m-1} dx,$$

i.e.,

$$\Gamma m = \int_0^{\infty} \lambda^m e^{-\lambda x} x^{m-1} dx.$$

Multiplying both sides by $e^{-\lambda\lambda^{n-1}}$ and integrating w.r.t. λ within the limits 0 to ∞ , we get

$$\Gamma m \int_0^\infty e^{-\lambda} \cdot \lambda^{n-1} d\lambda = \int_0^\infty \left[\int_0^\infty e^{-\lambda(1+x)} \cdot \lambda^{m+n-1} d\lambda \right] x^{m-1} dx$$

or $\Gamma m \cdot \Gamma n = \int_0^\infty \frac{\Gamma(m+n)}{(1+x)^{m+n}} \cdot x^{m-1} dx$ by equ. (2) of §6.4
 $= \Gamma(m+n) \beta(m, n)$ by §6.5.

$$\therefore \beta(m, n) = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}.$$

COROLLARY. If we put $m+n=1$ in

$$\Gamma m \Gamma n = \int_0^\infty \frac{\Gamma(m+n)}{(1+x)^{m+n}} \cdot x^{m-1} dx,$$

we have $\Gamma m \Gamma (1-m) = \int_0^\infty \frac{x^{m-1}}{1+x} dx$ [$\because \Gamma 1=1$]

$$= \frac{\pi}{\sin m\pi} \quad (\text{Agra, 1964})$$

$$\left[\because \int_0^\infty \frac{x^{2m}}{1+x^{2n}} dx = \frac{\pi}{2n \sin \frac{2m+1}{2n} \pi} \right]$$

when m and n are +ive integers and $n > m$ as will be shown in §6.9

Replacing m by $\frac{1}{2}$, we get

$$\Gamma \frac{1}{2} \Gamma \frac{1}{2} = \pi \text{ or } \Gamma \frac{1}{2} = \sqrt{\pi}.$$

(Nagpur, 1965)

6.8. REDUCTION OF DEFINITE INTEGRALS TO GAMMA FUNCTIONS

[1] To show that

$$\int_0^\infty \frac{y^{m-1}}{(1+y)^{m+n}} dy = \int_0^1 \frac{x^{m-1} + x^{n-1}}{(1+x)^{m+n}} dx = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$$

we know that

$$\int_0^\infty \frac{y^{m-1}}{(1+y)^{m+n}} dy = \int_0^1 \frac{x^{m-1}}{(1+y)^{m+n}} dy + \int_1^\infty \frac{y^{m-1}}{(1+y)^{m+n}} dy.$$

Substituting $\frac{1}{x}$ for y in the second integral on R.H.S., we get

$$\int_1^\infty \frac{y^{m-1}}{(1+y)^{m+n}} dy = \int_1^0 \frac{\left(\frac{1}{x}\right)^{m-1}}{\left(1+\frac{1}{x}\right)^{m+n}} \left(-\frac{1}{x^2}\right) dx,$$

$$\therefore y = \frac{1}{x}, dy = -\frac{1}{x^2} dx$$

$$= \int_0^1 \frac{x^{n-1}}{(1+x)^{m+n}} dx.$$

$$\therefore \int_0^\infty \frac{y^{m-1}}{(1+y)^{m+n}} dy = \int_0^1 \frac{x^{m-1}}{(1+x)^{m+n}} dx + \int_0^1 \frac{x^{n-1}}{(1+x)^{m+n}} dx$$

[$\because$ change of variable does not change the value of integral]

$$\beta(m, n) = \int_0^1 \frac{x^{m-1} + x^{n-1}}{(1+x)^{m+n}} dx = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}.$$

[2] If we substitute $\frac{ay}{b}$ for x , we get

$$\int_0^\infty \frac{x^{m-1}}{(1+x)^{m+n}} a^m b^n \int_0^\infty \frac{y^{m-1}}{(ay+b)^{m+n}} dy$$

Since $\int_0^\infty \frac{x^{m-1}}{(1+x)^{m+n}} dx = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$ by §§6.6 and 6.7

$$\therefore a^m b^n \int_0^\infty \frac{y^{m-1}}{(ay+b)^{m+n}} dy = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$$

$$\int_0^\infty \frac{y^{m-1}}{(ay+b)^{m+n}} dy = \frac{\Gamma m \Gamma n}{a^m b^n \Gamma(m+n)}.$$

COROLLARY. Substituting $y = \tan^2 \theta$, this relation transforms to

$$\int_0^{\pi/2} \frac{\sin^{2m-1} \theta \cos^{2n-1} \theta d\theta}{(a \sin^2 \theta + b \cos^2 \theta)^{m+n}} = \frac{\Gamma m \Gamma n}{2a^m b^n \Gamma(m+n)}.$$

[3] If we put $x = \sin^2 \theta$, we get

$$\int_0^1 x^{m-1} (1-x)^{n-1} dx = 2 \int_0^{\pi/2} \sin^{2(m-1)} \theta \cos^{2(n-1)} \theta \cdot \sin \theta \cos \theta d\theta$$

$$[\because dx = 2 \sin \theta \cos \theta d\theta]$$

$$\frac{\Gamma m \Gamma n}{\Gamma(m+n)} = 2 \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta.$$

$$\therefore \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta = \frac{\Gamma m \Gamma n}{2\Gamma(m+n)}.$$

COROLLARY. Replacing $2m-1$ by p and $2n-1$ by q , this relation

$$\text{reduces to } \int_0^{\pi/2} \sin^p \theta \cos^q \theta d\theta = \frac{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{q+1}{2}\right)}{2\Gamma\left(\frac{p+q+2}{2}\right)}.$$

Putting $p=0$ and $q=0$ in succession, we get

$$\int_0^{\pi/2} (\cos \theta)^q d\theta = \frac{\Gamma\left(\frac{q+1}{2}\right) \sqrt{\pi}}{2\Gamma\left(\frac{q}{2}+1\right)} \text{ and } \int_0^{\pi/2} (\sin \theta)^p d\theta = \frac{\Gamma\left(\frac{p+1}{2}\right) \sqrt{\pi}}{\Gamma\left(\frac{p}{2}+1\right)}.$$

[4] By putting $x = \sin^2 \theta$ in $\frac{\Gamma m \Gamma n}{\Gamma(m+n)} = \int_0^1 x^{m-1} (1-x)^{n-1} dx$.

We have just proved that

$$\int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta = \frac{\Gamma m \Gamma n}{2\Gamma(m+n)} \quad \dots(1)$$

Now if we put $2n=1$, we have

$$\int_0^{\pi/2} \sin^{2m-1} \theta d\theta = \frac{\sqrt{\pi}}{2} \frac{\Gamma m}{\Gamma(m+\frac{1}{2})} \quad \dots(2)$$

Again putting $m=n$ in (1), we find

$$\int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2m-1} \theta d\theta = \frac{(\Gamma m)^2}{2\Gamma(2m)}$$

$$\begin{aligned} \text{or } \frac{\{\Gamma(m)\}^2}{2\Gamma(2m)} &= \frac{1}{2^{2m-1}} \int_0^{\pi/2} \sin^{2m-1} 2\theta d\theta \quad [\because 2 \sin \theta \cos \theta = \sin 2\theta] \\ &= \frac{1}{2^{2m}} \int_0^{\pi} \sin^{2m-1} \phi d\phi \quad \text{put } 2\theta = \phi, d\theta = \frac{1}{2} d\phi \\ &= \frac{1}{2^{2m}} \cdot 2 \int_0^{\pi/2} \sin^{2m-1} \phi d\phi \quad [\because \sin(\pi-\phi) = \sin \phi] \\ &\quad \text{(Prop. of definite integral)} \\ \therefore \int_0^{\pi/2} \sin^{2m-1} \theta d\theta &= \frac{2^{2m-2} (\Gamma m)^2}{\Gamma(2m)} \quad \dots(3) \\ &\quad \text{(replacing } \phi \text{ by } \theta) \end{aligned}$$

From (2) and (3) it is obvious that

$$\frac{2^{2m-2} (\Gamma m)^2}{\Gamma(2m)} = \frac{\Gamma m}{\Gamma(m+\frac{1}{2})} \frac{\sqrt{\pi}}{2}$$

$$\text{or } \Gamma m \Gamma(m+\frac{1}{2}) = \frac{\sqrt{\pi}}{2^{2m-1}} \Gamma(2m).$$

This may also be put in the form

$$\Gamma(2m) = \frac{2^{2m-1}}{\sqrt{\pi}} \Gamma m \Gamma(m+\frac{1}{2}). \quad (\text{Agra, 1962, 65})$$

6.9. MISCELLANEOUS PROPOSITIONS

[1] **Infinite and Improper Integrals : their Principal and General values.**

Riemann introduced the definition of definite integral $\int_a^b f(x) dx$ under two assumptions :

- Integral $f(x)$ is bounded, and
- the range of integration i.e. (a, b) is finite.

Cauchy modified this definition to include the cases:

- The range of integration is infinite, e.g.

$$\int_1^{\infty} \frac{dx}{x^{2/3}}, \quad \int_{-\infty}^{\infty} \frac{dx}{1+x^2}, \quad \text{and}$$

- the integrand becomes infinite, i.e. the function is unbounded in the range e.g., $\int_{-1}^1 \frac{dx}{x^{2/3}}$ (unbounded at $x=0$).

The former is known as *Infinite integral of first kind* and the latter as *infinite integral of second kind*. But by convention we are accustomed of speaking the first type as **infinite integral** and the second kind as **improper integral**.

If there is an integral $\int_a^b f(x) dx$, where $f(x)$ becomes infinite at $x=c$ ($a < c < b$), then in order to exclude c , the point $x=c$ may be enclosed in a small neighbourhood $(c-\mu\epsilon, c+\nu\epsilon)$, μ and ν being any arbitrary constants, so that

$$\int_a^b f(x) dx = \lim_{\epsilon \rightarrow 0} \left[\int_a^{c-\mu\epsilon} f(x) dx + \int_{c+\nu\epsilon}^b f(x) dx \right].$$

This limit is known to be the **General value** of the given integral. In case $\mu=\nu$, the value of this limit is said to be the **Principal value** of the integral.

The integrals having infinite limits may be expressed as follows:

$$\int_a^{\infty} f(x) dx = \lim_{\epsilon \rightarrow 0} \int_a^{1/\epsilon} f(x) dx,$$

$$\int_{-\infty}^b f(x) dx = \lim_{\epsilon \rightarrow 0} \int_{-1/\epsilon}^b f(x) dx,$$

$$\int_{-\infty}^{\infty} f(x) dx = \lim_{\epsilon \rightarrow 0} \int_{-1/\mu\epsilon}^{1/\nu\epsilon} f(x) dx \text{ etc.}$$

[2] **Evaluation of the Integrals of the Type** $\int_{-\infty}^{\infty} \frac{f(x)}{F(x)} dx$,

where $\frac{f(x)}{F(x)}$ is rational algebraic function such that the degree of $f(x)$ is at least two lower than that of $F(x)$ and all the roots of the equation $F(x)=0$ are imaginary.

Since imaginary roots occur in pairs and all the roots of $F(x)=0$ are imaginary, it follows that $F(x)$ must be of an even number in degree say $2n$. Further it follows that $F(x)$ cannot become zero for any real value of x and consequently the integrand $\frac{f(x)}{F(x)}$ is always finite for all real values of x .

Let us suppose that any pair of imaginary roots is $a \pm ib$, so that the factors of $F(x)$ corresponding to these roots are $(x-a-ib)$ and $(x-a+ib)$. Let the corresponding partial fractions be of the form

$$\frac{A-ib}{x-a-ib} \quad \text{and} \quad \frac{A+ib}{x-a+ib}.$$

$$\text{Then } \frac{A-ib}{(x-a)-ib} + \frac{A+ib}{(x-a)+ib} = \frac{2A(x-a)+2Bb}{(x-a)^2+b^2}.$$

$$\text{Now, } \int_{-\infty}^{\infty} \frac{2A(x-a)+2Bb}{(x-a)^2+b^2} dx$$

$$= \lim_{\epsilon \rightarrow 0} \left[\int_{-1/\mu\epsilon}^{1/\nu\epsilon} \frac{2A(x-a)}{(x-a)^2+b^2} dx + \int_{-1/\mu\epsilon}^{1/\nu\epsilon} \frac{2Bb}{(x-a)^2+b^2} dx \right]$$

$$\begin{aligned}
&= \lim_{\epsilon \rightarrow 0} \left[A \log \{(x-a)^2 + b^2\} + 2B \tan^{-1} \frac{x-a}{b} \right]^{1/\nu\epsilon} \\
&= \lim_{\epsilon \rightarrow 0} \left[A \log \frac{(1-a\mu\epsilon)^2 + c^2\nu^2\epsilon^2}{(-1-a\mu\epsilon)^2 + b^2\mu^2\epsilon^2} \cdot \frac{\mu^2}{\nu^2} \right. \\
&\quad \left. + 2B \left\{ \tan^{-1} \left(\frac{1-a}{b\mu\epsilon} \right) + \tan^{-1} \left(\frac{1+a}{b\mu\epsilon} \right) \right\} \right] \\
&= A \log \frac{\mu^2}{\nu^2} + 2\pi B = 2 \left(A \log \frac{\mu}{\nu} + \pi B \right). \quad \dots (1)
\end{aligned}$$

As we have already supposed that $F(x)$ is of degree $2n$, so it will consist of n pairs of imaginary roots. Let these pairs be $a_1 \pm ib_1, a_2 \pm ib_2, \dots, a_n \pm ib_n$ and let the corresponding constants in the partial fractions be $(A_1, B_1), (A_2, B_2), \dots, (A_n, B_n)$; then we have

$$\begin{aligned}
\int_{-\infty}^{\infty} \frac{f(x)}{F(x)} dx &= \int_{-\infty}^{\infty} \frac{2A_1(x-a_1) + 2B_1b_1}{(x-a_1)^2 + b_1^2} dx \\
&+ \int_{-\infty}^{\infty} \frac{2A_2(x-a_2) + 2B_2b_2}{(x-a_2)^2 + b_2^2} dx + \dots + \int_{-\infty}^{\infty} \frac{2A_n(x-a_n) + 2B_nb_n}{(x-a_n)^2 + b_n^2} dx \\
&= 2 \left[A_1 \log \frac{\mu}{\nu} + \pi B_1 \right] + 2 \left[A_2 \log \frac{\mu}{\nu} + \pi B_2 \right] + \dots + 2 \left[A_n \log \frac{\mu}{\nu} + \pi B_n \right] \quad \text{from (1)} \\
&= 2 (A_1 + A_2 + \dots + A_n) \log \frac{\mu}{\nu} + 2\pi (B_1 + B_2 + \dots + B_n). \quad \dots (2)
\end{aligned}$$

$$\text{Also } \frac{f(x)}{F(x)} = \frac{2A_1(x-a_1) + 2B_1b_1}{(x-a_1)^2 + b_1^2} + \dots + \frac{2A_n(x-a_n) + 2B_nb_n}{(x-a_n)^2 + b_n^2}.$$

$$\therefore f(x) = \frac{2A_1(x-a_1) + 2B_1b_1}{(x-a_1)^2 + b_1^2} \cdot F(x) + \dots$$

Since $(x-a_1)^2 + b_1^2$ is one of the factors of $F(x)=0$, therefore $F(x)$ is completely divisible by $(x-a_1)^2 + b_1^2$ etc....

$$\therefore f(x) = \{2A_1(x-a_1) + 2B_1b_1\} [x^{2n-2} + \dots] + \dots$$

But $f(x)$ being at most of degree $2n-2$, on equating the coefficients of x^{2n-1} on either side, we get

$$0 = 2(A_1 + A_2 + \dots + A_n).$$

Hence from (2), we get

$$\int_{-\infty}^{\infty} \frac{f(x)}{F(x)} dx = 2\pi(B_1 + B_2 + \dots + B_n).$$

Applications of this result.

(i) Evaluation of $\int_0^{\infty} \frac{x^{2m}}{1+x^{2n}} dx$, where m and n are positive integers and $n > m$.

Here $1+x^{2n}=0$ gives $x=(-1)^{1/2n}$, i.e., all the roots imaginary.

If α be a root of the equation $x^{2n}+1=0$, then $\alpha=(-1)^{1/2n}$.

Putting $-1=k \cos \theta$ and $0=k \sin \theta$ so that $k=1, \theta=\pi$, we get

$1 = [\cos(2r\pi + \pi) + i \sin(2r\pi + \pi)]^{1/2n}$ where $r=0, 1, 2, \dots$

$$= \cos \frac{2r+1}{2n} \pi + i \sin \frac{2r+1}{2n} \pi \text{ by De Moivre's theorem}$$

where $r=0, 1, 2, \dots, 2n-1$.

Since imaginary roots occur in pairs, α is of the form

$$\alpha = \cos \frac{2r+1}{2n} \pi \pm i \sin \frac{2r+1}{2n} \pi, \text{ where } r=0, 1, 2, \dots, (n-1).$$

Now α being one of the roots of $x^{2n}+1=0$, $(x-\alpha)$ is one of the factors of $(x^{2n}+1)$. Let $\frac{A-Bi}{x-\alpha}$ be the partial fraction corresponding

to any root α of the type $\cos \frac{2r+1}{2n} \pi + i \sin \frac{2r+1}{2n} \pi$. Then

$$\frac{A-Bi}{x-\alpha} = \frac{\alpha^{2m}}{2n\alpha^{2n-1}(x-\alpha)}$$

[Note. If $(x-a)$ is a factor of $F(x)$ in $\frac{f(x)}{F(x)}$, then the partial fraction corresponding to $(x-a)$ is $\frac{f(a)}{F'(a)(x-a)}$].

$$\therefore A-Bi = \frac{\alpha^{2m+1}}{2n\alpha^{2n}} = \frac{\alpha^{2m-1}}{-2n}, \text{ since } \alpha^{2n} = -1,$$

α being root of $x^{2n}+1=0$

$$= -\frac{1}{2n} \left[\cos \frac{2r+1}{2n} \pi + i \sin \frac{2r+1}{2n} \pi \right]^{2m-1}$$

$$= -\frac{1}{2n} \left[\cos(2r+1) \frac{(2m+1)}{2n} \pi + i \sin(2r+1) \frac{(2m+1)}{2n} \pi \right]$$

$$= -\frac{1}{2n} \left[\cos(2r+1)\theta + i \sin(2r+1)\theta \right], \text{ where } \theta = \frac{2m+1}{2n} \pi.$$

Equating imaginary parts on either side, we get

$$B = \frac{1}{2n} \sin(2r+1)\theta \text{ where } r=0, 1, 2, \dots, (n-1).$$

Hence if $B_1, B_2, \dots, B_n$ be the constants corresponding to $r=0, 1, \dots, (n-1)$, then

$$\int_{-\infty}^{\infty} \frac{x^{2m}}{1+x^{2n}} dx = 2\pi [B_1 + B_2 + \dots + B_n]$$

$$\therefore \int_0^{\infty} \frac{x^{2m}}{1+x^{2n}} dx = \frac{\pi}{2n} [\sin \theta + \sin 3\theta + \dots + \sin(2n-1)\theta]$$

$$= \frac{\pi}{4n \sin \theta} [2 \sin^2 \theta + 2 \sin \theta \sin 3\theta + \dots$$

$$+ 2 \sin \theta \sin(2n-1)\theta]$$

$$\begin{aligned}
&= \frac{\pi}{4n \sin \theta} [(1 - \cos 2\theta) + (\cos 2\theta - \cos 4\theta) + \dots \\
&\quad + \{\cos (2n-2)\theta - \cos 2n\theta\}] \\
&= \frac{\pi}{4n \sin \theta} [1 - \cos 2n\theta] = \frac{\pi}{4n \sin \theta} \cdot 2 \sin^2 n\theta \\
&= \frac{\pi \sin^2 n\theta}{2n \sin \theta}.
\end{aligned}$$

But $\sin n\theta = \sin \frac{2m+1}{2} \pi = \pm 1$, so that $\sin^2 n\theta = 1$.

$$\text{Hence } \int_0^\infty \frac{x^{2m} dx}{1+x^{2n}} = \frac{\pi}{2n \sin \theta} = \frac{\pi}{2n} \operatorname{cosec} \frac{2m+1}{2n} \pi.$$

(ii) Evaluation of $\int_0^\infty \frac{x^{2m}}{1-x^{2n}} dx$, where m and n are positive integers and $n > m$.

Here the equation $1-x^{2n}=0$ has got only two real roots ± 1 , corresponding to the factors $(1-x)$ and $(1+x)$.

Now $(x-1)$ corresponds to the partial fraction

$$\frac{(1)^{2m}}{-2n(+1)^{2n-1}(x-1)}, \text{ i.e., } \frac{1}{-2n(x-1)}$$

and $(x+1)$ corresponds to the partial fraction

$$\frac{(-1)^{2m}}{-2n(-1)^{2n-1}(x+1)}, \text{ i.e., } \frac{1}{2n(x+1)}.$$

$$\therefore \frac{1}{2n(x+1)} - \frac{1}{2n(x-1)} + \frac{1}{2n} \left[\frac{x-1-x-1}{x^2-1} \right] = \frac{1}{n(1-x^2)}.$$

$$\text{But } \int_0^\infty \frac{dx}{1-x^2} = \int_0^1 \frac{dx}{1-x^2} + \int_1^\infty \frac{dx}{1-x^2}.$$

If we put $x = \frac{1}{z}$ in the last integral, we have

$$\int_0^\infty \frac{dx}{1-x^2} = - \int_1^0 \frac{(1/z^2) dz}{1-(1/z^2)} = \int_0^1 \frac{dz}{z^2-1} = - \int_0^1 \frac{dx}{1-x^2}$$

(by the properties of definite integrals),

$$\text{so that } \int_0^\infty \frac{dx}{1-x^2} = \int_0^1 \frac{dx}{1-x^2} - \int_0^1 \frac{dx}{1-x^2} = 0.$$

It follows that the part of the definite integral corresponding to real roots of $1-x^{2n}=0$ vanishes and as such we have to consider only the integral corresponding to $(n-1)$ conjugate pairs of imaginary roots.

Now $1-x^{2n}=0$ gives $x=(1)^{1/2n}$

or $x=(\cos 2r\pi + i \sin 2r\pi)^{1/2n}$ on putting $1=k \cos \theta$, $0=k \sin \theta$,
so that $k=1$, $\theta=0$,

$$= \cos \frac{2r\pi}{2n} \pm i \sin \frac{2r\pi}{2n}.$$

Leaving the roots corresponding to $r=0$ and $r=n$, which are real, let α be any of the $(n-1)$ pairs of imaginary roots, so that

$$\alpha = \cos \frac{2r\pi}{2n} + i \sin \frac{2r\pi}{2n} \text{ where } r=1, 2, 3, \dots, (n-1).$$

If $\frac{A-Bi}{x-\alpha}$ be the partial fraction corresponding to any root α ,

$$\text{then } A-Bi = \frac{\alpha^{2m}}{-2n\alpha^{2n-1}} = -\frac{1}{2n} \alpha^{2m+1} \quad \text{since } \alpha^{2n}=1$$

$$= -\frac{1}{2n} \left[\cos \frac{2r\pi}{2n} + i \sin \frac{2r\pi}{2n} \right]^{2m+1}$$

$$= -\frac{1}{2n} \left[\cos \frac{2r(2m+1)\pi}{2n} + i \sin \frac{2r(2m+1)\pi}{2n} \right]$$

$$= -\frac{1}{2n} [\cos 2r\theta + i \sin 2r\theta] \text{ where } \theta = \frac{2m+1}{2n} \pi$$

and $r=1, 2, 3, \dots, (n-1)$.

Equating imaginary parts on either side, we get

$$B = \frac{1}{2n} \sin 2r\theta.$$

Hence if $B_1, B_2, \dots, B_n$ be the constants corresponding to $r=1, 2, \dots, (n-1)$, then

$$\int_0^\infty \frac{x^{2m}}{1-x^{2n}} dx = \pi [B_1 + B_2 + \dots + B_n]$$

$$= \frac{\pi}{2n} [\sin 2\theta + \sin 4\theta + \dots + \sin (2n-2)\theta]$$

$$= \frac{\pi}{4n \sin \theta} (2 \sin \theta \sin 2\theta + 2 \sin \theta \sin 4\theta + \dots$$

$$+ 2 \sin \theta \sin (2n-2)\theta]$$

$$= \frac{\pi}{4n \sin \theta} [(\cos \theta - \cos 3\theta) + (\cos 3\theta - \cos 5\theta) + \dots \\ + \{\cos (2n-3)\theta - \cos (2n-1)\theta\}]$$

$$= \frac{\pi}{4n \sin \theta} [\cos \theta - \cos (2n-1)\theta]$$

$$= \frac{\pi}{4n \sin \theta} \cdot 2 \sin n\theta \cdot \sin (n-1)\theta.$$

$$\text{But } \sin (n-1)\theta = \sin (n-1) \frac{2m+1}{2n} \pi$$

$$= \sin \left[\frac{2m+1}{2} \pi - \frac{2m+1}{2n} \pi \right] = \pm \cos \frac{2m+1}{2n} \pi$$

and

$$\sin n\theta = \frac{2m+1}{2} \pi = \pm 1.$$

Hence

$$\int_0^\infty \frac{x^{2m}}{1-x^{2n}} dx = \frac{\pi}{2n} \cot \frac{2m+1}{2n} \pi.$$

Deductions. We have proved

$$\int_0^{\infty} \frac{x^{2m}}{1+x^{2n}} dx = \frac{\pi}{2n} \operatorname{cosec} \frac{2m+1}{2n} \pi \quad \dots (1)$$

$$\text{and} \quad \int_0^{\infty} \frac{x^{2m}}{1-x^{2n}} dx = \frac{\pi}{2n} \cot \frac{2m+1}{2n} \pi \quad \dots (2)$$

If we put $x^{2n}=z$ and then $\frac{2m+1}{2n}=\alpha$, we can easily show that (1) and (2) give respectively

$$\int_0^{\infty} \frac{z^{\alpha-1} dz}{1+z} = \pi \operatorname{cosec} \pi \alpha \quad \text{and} \quad \int_0^{\infty} \frac{z^{\alpha-1} dz}{1-z} = \pi \cot \pi \alpha.$$

[iii] Evaluation of $\Gamma \frac{1}{n} \Gamma \frac{2}{n} \Gamma \frac{3}{n} \dots \Gamma \left(\frac{n-1}{n} \right)$, where n is a positive integer.

$$\begin{aligned} \text{Let } P &= \Gamma \frac{1}{n} \Gamma \frac{2}{n} \Gamma \frac{3}{n} \dots \Gamma \left(\frac{n-1}{n} \right) \\ &= \Gamma \frac{1}{n} \Gamma \frac{2}{n} \Gamma \frac{3}{n} \dots \Gamma \left(1 - \frac{1}{n} \right). \end{aligned} \quad \dots (3)$$

By reversing the order of the factors, this may be written as

$$\begin{aligned} P &= \Gamma \left(1 - \frac{1}{n} \right) \Gamma \left(1 - \frac{2}{n} \right) \Gamma \left(1 - \frac{3}{n} \right) \dots \Gamma \frac{2}{n} \Gamma \frac{1}{n} \\ &= \Gamma \left(1 - \frac{1}{n} \right) \Gamma \left(1 - \frac{2}{n} \right) \Gamma \left(1 - \frac{3}{n} \right) \dots \Gamma \left(1 - \frac{n-2}{n} \right) \Gamma \left(1 - \frac{n-1}{n} \right) \end{aligned} \quad \dots (4)$$

Multiplying (1) and (2), we get

$$\begin{aligned} P^2 &= \left\{ \Gamma \left(\frac{1}{n} \right) \Gamma \left(1 - \frac{1}{n} \right) \right\} \left\{ \Gamma \left(\frac{2}{n} \right) \Gamma \left(1 - \frac{2}{n} \right) \right\} \\ &\quad \dots \left\{ \Gamma \left(\frac{n-1}{n} \right) \Gamma \left(1 - \frac{n-1}{n} \right) \right\} \\ &= \frac{\pi}{\sin \frac{\pi}{n}} \cdot \frac{\pi}{\sin \frac{2\pi}{n}} \dots \frac{\pi}{\sin \frac{n-1}{n} \pi}, \text{ since } \Gamma m \Gamma(1-m) = \frac{\pi}{\sin \pi m} \end{aligned}$$

$$= \frac{\pi^{n-1}}{\sin \frac{\pi}{n} \sin \frac{2\pi}{n} \dots \sin \frac{n-1}{n} \pi} \quad \dots (5)$$

$$\text{Suppose that } D = \sin \frac{\pi}{n} \sin \frac{2\pi}{n} \dots \sin \frac{n-1}{n} \pi.$$

In order to calculate the value of D , let us consider the equation $1-x^{2n}=0$, i.e., $x=(1)^{1/2n}$

$$= (\cos 2r\pi + i \sin 2r\pi)^{1/2n}$$

$$= \cos \frac{2r\pi}{2n} + i \sin \frac{2r\pi}{2n}, \quad \text{where } r=0, 1, 2, \dots, (2n-1),$$

so that we have the identity

$$\begin{aligned} 1-x^{2n} &= (1-x)(1+x) \left\{ x - \left(\cos \frac{\pi}{n} + i \sin \frac{\pi}{n} \right) \right\} \\ &\quad \left\{ x - \left(\cos \frac{\pi}{n} - i \sin \frac{\pi}{n} \right) \right\} \times \left\{ x - \left(\cos \frac{2\pi}{n} + i \sin \frac{2\pi}{n} \right) \right\} \\ &\quad \left\{ x - \left(\cos \frac{2\pi}{n} - i \sin \frac{2\pi}{n} \right) \right\} \dots \\ &\quad \left\{ x - \left(\cos \frac{n-1}{n} \pi + i \sin \frac{n-1}{n} \pi \right) \right\} \left\{ x - \left(\cos \frac{n-1}{n} \pi - i \sin \frac{n-1}{n} \pi \right) \right\}, \\ 1-x^{2n} &= \left(1-2x \cos \frac{\pi}{n} + x^2 \right) \left(1-2x \cos \frac{2\pi}{n} + x^2 \right) \dots \\ &\quad \dots \left(1-2x \cos \frac{n-1}{n} \pi + x^2 \right) \end{aligned}$$

Putting $x=1$ and $x=-1$ in turn in the above identity, we have

$$\begin{aligned} n &= \left(2-2 \cos \frac{\pi}{n} \right) \left(2-2 \cos \frac{2\pi}{n} \right) \dots \left(2-2 \cos \frac{n-1}{n} \pi \right) \\ &= \left(2 \sin \frac{\pi}{2n} \right)^2 \left(2 \sin \frac{2\pi}{2n} \right)^2 \dots \left(2 \sin \frac{n-1}{n} \pi \right)^2 \quad \dots (4) \end{aligned}$$

$$\begin{aligned} n &= \left(2+2 \cos \frac{\pi}{n} \right) \left(2+2 \cos \frac{2\pi}{n} \right) \dots \left(2+2 \cos \frac{n-1}{n} \pi \right) \\ &= \left(2 \cos \frac{\pi}{2n} \right)^2 \left(2 \cos \frac{2\pi}{2n} \right)^2 \dots \left(2 \cos \frac{n-1}{2n} \pi \right)^2. \quad \dots (5) \end{aligned}$$

$$\left[\text{Since } \lim_{x \rightarrow 1} \frac{1-x^{2n}}{1-x^2} = \lim_{x \rightarrow 1} \frac{-2nx^{2n-1}}{-2x} = n, \text{ also } \lim_{x \rightarrow -1} \frac{1-x^{2n}}{1-x^2} = n \right]$$

Multiplying (4) and (5) we get

$$n^2 = 2^{2n-2} \sin^2 \frac{\pi}{n} \sin^2 \frac{2\pi}{n} \dots \sin^2 \frac{n-1}{n} \pi = 2^{2(n-1)} \cdot D^2$$

so that

$$D = \frac{n}{2^{n-1}}.$$

[Note. The value of D may be calculated by using the identity

$$\sin^2 n\theta = 2^{n-1} \sin \left(\theta + \frac{\pi}{n} \right) \sin \left(\theta + \frac{2\pi}{n} \right) \dots \sin \left(\theta + \frac{n-1}{n} \pi \right)$$

if we put $\theta=0$, $\lim_{\theta \rightarrow 0} \frac{\sin n\theta}{\sin \theta} = \lim_{\theta \rightarrow 0} \frac{n \cos n\theta}{\cos \theta} = n$, so that

$$n = 2^{n-1} \cdot \sin \frac{\pi}{2} \sin \frac{2\pi}{n} \dots \sin \frac{n-1}{n} \pi \quad]$$

Thus we have from (3)

$$P^2 = \frac{\pi^{n-1}}{n} \cdot 2^{n-1}$$

i.e. $P = \Gamma \frac{1}{n} \Gamma \frac{2}{n} \dots \Gamma \left(\frac{n-1}{n} \right) = \frac{(2\pi)^{(n-1)/2}}{\sqrt{n}}$, which is the required value

Problem 1. Show that $\int_0^1 \frac{x^{m-1} (1-x)^{n-1} dx}{(a+x)^{m+n}} = \frac{\Gamma m \Gamma n}{a^n (1+a)^m \Gamma(m+n)}$

Let
$$I = \int_0^1 \frac{x^{m-1} (1-x)^{n-1} dx}{(a+x)^{m+n}}$$

Put
$$\frac{x(1+a)}{(a+x)} = y, \text{ so that } x = \frac{ay}{1+a-y}.$$

and $(1+a) \cdot \frac{a+x-x}{(a+x)^2} dx = dy$, i.e. $dx = \frac{(a+x)^2}{a(1+a)} dy$.

As such $1-x = 1 - \frac{ay}{1+a-y} = \frac{1+a-y-ay}{1+a-y} = \frac{(1+a)(1-y)}{1+a-y}$

Also $a+x = a + \frac{ay}{1+a-y} = \frac{a+a^2-ay+ay}{1+a-y} = \frac{a(1+a)}{1+a-y}$

and therefore, $dx = \frac{a^2(1+a)^2}{(1+a-y)^2 a(1+a)} dy = \frac{a(1+a)}{(1+a-y)^2} dy$.

Thus

$$I = \int_0^1 \frac{a^{m-1} y^{m-1} (1+a)^{n-1} (1-y)^{n-1} (1+a-y)^{m+n} a(1+a)}{(1+a-y)^{m-1} (1+a-y)^{n+1} a^{m+n} (1+a)^{m+n} (1+a-y)^2} dy$$

$$= \frac{1}{a^n (1+a)^m} \int_0^1 y^{m-1} (1-y)^{n-1} dy$$

or $I = \frac{1}{a^n (1+a)^m} \beta(m, n) = \frac{1}{a^n (1+a)^m} \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$

Problem 2. Show that $2^n \Gamma(n+\frac{1}{2}) = 1 \cdot 3 \cdot 5 \dots (2n-1) \sqrt{\pi}$.

(Agra, 1972)

We know that $\Gamma(n+1) = n\Gamma n$.

$$\therefore \Gamma(n+\frac{1}{2}) = (n-\frac{1}{2}) \Gamma(n-\frac{1}{2})$$

$$= (n-\frac{1}{2}) (n-\frac{3}{2}) \Gamma(n-\frac{3}{2})$$

$$\dots \dots \dots = (n-\frac{1}{2}) (n-\frac{3}{2}) (n-\frac{5}{2}) (n-\frac{7}{2}) \dots \frac{5}{2} \cdot \frac{3}{2} \cdot \frac{1}{2} \Gamma \frac{1}{2}$$

$$= (n-\frac{1}{2}) (n-\frac{3}{2}) (n-\frac{5}{2}) \dots \frac{5}{2} \cdot \frac{3}{2} \cdot \frac{1}{2} \sqrt{\pi}$$

$$= \frac{\sqrt{\pi}}{2^n} (2n-1) (2n-3) \dots 5 \cdot 3 \cdot 1$$

or $2^n \Gamma(n+\frac{1}{2}) = 1 \cdot 3 \cdot 5 \dots (2n-1) \sqrt{\pi}$.

Problem 3. Show that $\int_0^1 \frac{dx}{\sqrt{(1-x^n)}} = \frac{\Gamma(\frac{1}{n})}{\Gamma(\frac{1}{2} + \frac{1}{n})} \frac{\sqrt{\pi}}{n}$. (Agra, 1972)

Suppose
$$I = \int_0^1 \frac{dx}{\sqrt{(1-x^n)}}$$

Put $x^n = \sin^2 \theta$, i.e. $x = \sin^{2/n} \theta$.

$\therefore dx = \frac{2}{n} \sin^{(2-n)/n} \theta \cos \theta d\theta$.

Thus
$$I = \frac{2}{n} \int_0^{\pi/2} \frac{\sin^{(2-n)/n} \theta \cos \theta d\theta}{\cos \theta}$$

$$= \frac{2}{n} \int_0^{\pi/2} \sin^{(2-n)/n} \theta d\theta$$

$$= \frac{2}{n} \cdot \frac{\Gamma(\frac{1}{2}) \Gamma(\frac{1}{n})}{2\Gamma(\frac{1}{2} + \frac{1}{n})} = \frac{\Gamma(\frac{1}{n})}{\Gamma(\frac{1}{2} + \frac{1}{n})} \frac{\sqrt{\pi}}{n}$$

Problem 4. Prove the relation

$$\int_0^{\pi/2} \frac{d\theta}{\sqrt{(\sin \theta)}} \times \int_0^{\pi/2} \sqrt{(\sin \theta)} d\theta = \pi.$$

L.H.S. = $\int_0^{\pi/2} (\sin \theta)^{-1/2} d\theta \times \int_0^{\pi/2} (\sin \theta)^{1/2} d\theta$

$$= \frac{\Gamma \frac{1}{2} \Gamma \frac{1}{2}}{2\Gamma \frac{3}{2}} \times \frac{\Gamma \frac{3}{2} \Gamma \frac{3}{2}}{2\Gamma \frac{5}{2}}$$

$$= \frac{\pi}{4} \cdot \frac{\Gamma \frac{1}{2}}{\frac{1}{2} \Gamma \frac{1}{2}} = \pi$$

$\because \Gamma \frac{5}{2} = \frac{1}{2} \Gamma \frac{3}{2}$

Problem 5. Prove that $\int_0^{\pi/2} \sqrt{(\tan \theta)} d\theta = \frac{\Gamma \frac{1}{2} \Gamma \frac{3}{2}}{2}$. (Nagpur, 1965)

L.H.S. = $\int_0^{\pi/2} (\sin \theta)^{1/2} \times (\cos \theta)^{-1/2} d\theta$

$$= \frac{\Gamma \frac{3}{2} \Gamma \frac{1}{2}}{2\Gamma 1} = \frac{\Gamma \frac{1}{2} \Gamma \frac{3}{2}}{2}$$

Problem 6. Express the definite integral

$$\int_0^{\pi/2} \frac{d\theta}{\sqrt{(1-k^2 \sin^2 \theta)}}$$

in the form of series, where $k < 1$.

$$I = \int_0^{\pi/2} \frac{d\theta}{\sqrt{(1-k^2 \sin^2 \theta)}} = \int_0^{\pi/2} (1-k^2 \sin^2 \theta)^{-1/2} d\theta.$$

Expanding it by Binomial Theorem for any index,

$$I = \int_0^{\pi/2} \left[1 + \frac{1}{2} k^2 \sin^2 \theta + \frac{1 \cdot 3}{2 \cdot 4} k^4 \sin^4 \theta + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} k^6 \sin^6 \theta + \dots \right] d\theta$$

$$= \left[\theta \right]_0^{\pi/2} + \frac{1}{2} k^2 \frac{1}{2} \cdot \frac{\pi}{2} + \frac{1.3}{2.4} k^4 \cdot \frac{1.3}{2.4} \cdot \frac{\pi}{2} + \frac{1.3.5}{2.4.6} k^6 \cdot \frac{1.3.5}{2.4.6} \cdot \frac{\pi}{2} + \dots$$

$$= \frac{\pi}{2} \left[1 + \left(\frac{1}{2} \right)^2 k^2 + \left(\frac{1.3}{2.4} \right)^2 k^4 + \left(\frac{1.3.5}{2.4.6} \right)^2 k^6 + \dots \right]$$

Problem 7. Prove $\int_0^1 \frac{35x^3}{32\sqrt{(1-x)}} dx = 1$ (Nagpur, 1963)

Suppose $I = \int_0^1 \frac{35x^3}{32\sqrt{(1-x)}} dx$ put $x = \sin^2 \theta$

$$\therefore dx = 2 \sin \theta \cos \theta d\theta$$

$$= \int_0^{\pi/2} \frac{35 \sin^6 \theta \cdot 2 \sin \theta \cos \theta d\theta}{32 \cos \theta}$$

$$= \frac{35}{16} \int_0^{\pi/2} \sin^7 \theta d\theta$$

$$= \frac{35}{16} \cdot \frac{\Gamma(4) \Gamma(\frac{1}{2})}{2\Gamma(\frac{9}{2})} = \frac{35}{16} \cdot \frac{3.2.1 \sqrt{\pi}}{2 \cdot \frac{7}{2} \cdot \frac{5}{2} \cdot \frac{3}{2} \cdot \frac{1}{2} \sqrt{\pi}} = 1.$$

Problem 8. Prove that $\int_0^1 \frac{x^2 dx}{(1-x^4)^{1/2}} \times \int_0^1 \frac{dx}{(1+x^4)^{1/2}} = \frac{\pi}{4\sqrt{2}}$

Let $I_1 \equiv \int_0^1 \frac{x^2 dx}{(1-x^4)^{1/2}}$ and $I_2 \equiv \int_0^1 \frac{dx}{(1+x^4)^{1/2}}$

If we put $x^2 = \sin \theta$ in I_1 , i.e. $2x dx = \cos \theta d\theta$, we have

$$I_1 = \frac{1}{2} \int_0^{\pi/2} \frac{\sqrt{(\sin \theta) \cdot \cos \theta} d\theta}{\cos \theta}$$

$$= \frac{1}{2} \int_0^{\pi/2} \sin^{1/2} \theta d\theta = \frac{1}{2} \cdot \frac{\Gamma(\frac{3}{4}) \Gamma(\frac{1}{2})}{2\Gamma(\frac{5}{4})} = \frac{1}{4} \cdot \frac{\Gamma(\frac{3}{4}) \Gamma(\frac{1}{2})}{\Gamma(\frac{5}{4})} = \frac{\Gamma(\frac{3}{4}) \Gamma(\frac{1}{2})}{\Gamma(\frac{5}{4})}$$

Again putting $x^2 = \tan \phi$ in I_2 , i.e. $2x dx = \sec^2 \phi d\phi$, we get

$$I_2 = \frac{1}{2} \int_0^{\pi/4} \frac{\sec^2 \phi d\phi}{\sqrt{(\tan \phi) \cdot \sec \phi}} = \frac{1}{2} \int_0^{\pi/4} \frac{\sqrt{2} d\phi}{\sqrt{2 \sin \phi \cos \phi}}$$

$$= \frac{1}{\sqrt{2}} \int_0^{\pi/4} \frac{d\phi}{\sqrt{(\sin 2\phi)}} \text{ put } 2\phi = \psi, \text{ i.e. } 2 d\phi = d\psi$$

$$= \frac{1}{2\sqrt{2}} \int_0^{\pi/2} \sin^{-1/2} \psi d\psi$$

$$= \frac{1}{2\sqrt{2}} \cdot \frac{\Gamma(\frac{1}{4}) \Gamma(\frac{1}{2})}{2\Gamma(\frac{3}{4})} = \frac{\Gamma(\frac{1}{4}) \Gamma(\frac{1}{2})}{4\sqrt{2} \Gamma(\frac{3}{4})}$$

$$\text{Hence } I_1 \times I_2 = \frac{\Gamma(\frac{3}{4}) \Gamma(\frac{1}{2})}{\Gamma(\frac{5}{4})} \times \frac{\Gamma(\frac{1}{4}) \Gamma(\frac{1}{2})}{4\sqrt{2} \Gamma(\frac{3}{4})}$$

$$= \frac{(\Gamma(\frac{1}{2}))^2}{4\sqrt{2}} = \frac{\pi}{4\sqrt{2}}$$

Problem 9. Show that $\int_0^1 \frac{x^n dx}{\sqrt{(1-x^2)}} = \frac{1.3.5 \dots (n-1)}{2.4.6 \dots n} \cdot \frac{\pi}{2}$

$$\text{or } \frac{2.4.6 \dots (n-1)}{1.3.5 \dots n}$$

according as n is even or odd positive integer.

$$\text{L.H.S} = \int_0^1 \frac{x^n dx}{\sqrt{(1-x^2)}} \quad \text{put } x = \sin \theta, dx = \cos \theta d\theta$$

$$= \int_0^{\pi/2} \frac{\sin^n \theta \cdot \cos \theta d\theta}{\cos \theta}$$

$$= \int_0^{\pi/2} \sin^n \theta d\theta = \frac{\Gamma(\frac{n+1}{2}) \Gamma(\frac{1}{2})}{2\Gamma(\frac{n+2}{2})}$$

$$= \frac{\frac{n-1}{2} \cdot \frac{n-3}{2} \dots \frac{3}{2} \cdot \frac{1}{2} \Gamma(\frac{1}{2}) \Gamma(\frac{1}{2})}{2 \cdot \frac{n}{2} \cdot (\frac{n}{2} - 1) \dots 2.1}$$

$$= \frac{\frac{n-1}{2} \cdot \frac{n-3}{2} \dots 2.1 \Gamma(\frac{1}{2})}{2 \cdot \frac{n}{2} \cdot (\frac{n-2}{2}) \cdot (\frac{n-4}{2}) \dots \frac{3}{2} \cdot \frac{1}{2} \Gamma(\frac{1}{2})}$$

according as n is even or odd

$$= \frac{1.3.5 \dots (n-3)(n-1)}{2^{n/2} \cdot 1.2.3 \dots \frac{n}{2}} \cdot \frac{\pi}{2} \text{ or } \frac{1.2 \dots \frac{n-1}{2} \cdot 2^{(n-1)/2}}{1.3.5 \dots n}$$

according as n is even or odd

$$= \frac{1.3.5 \dots (n-1)}{2.4.6 \dots n} \cdot \frac{\pi}{2} \text{ or } \frac{2.4.6 \dots (n-1)}{1.3.5 \dots n}$$

according as n is even or odd.

Problem 10. Prove that $\int_0^1 (1-x^n)^{1/n} dx = \frac{1}{n} \frac{[\Gamma(1/n)]^2}{2\Gamma(2/n)}$

Let $I = \int_0^1 (1-x^n)^{1/n} dx$, put $x^n = \sin^2 \theta$, i.e., $x = \sin^{2/n} \theta$,

so that $dx = \frac{2}{n} \sin^{(2-n)/2} \theta \cos \theta d\theta$

$$= \frac{2}{n} \int_0^{\pi/2} \cos^{(2/n)+1} \theta \sin^{(2/n)-1} \theta d\theta$$

$$= \frac{2}{n} \cdot \frac{\Gamma(\frac{1}{n} + 1) \Gamma(\frac{1}{n})}{2\Gamma(\frac{2}{n} + 1)}$$

$$= \frac{1}{n} \cdot \frac{1/n \Gamma(1/n) \Gamma(1/n)}{2/n \Gamma(2/n)} = \frac{1}{n} \cdot \frac{\{\Gamma(1/n)\}^2}{\Gamma(2/n)}$$

Problem 11. Show that $\int_0^1 x^{m-1} (1-x^a)^n dx = \frac{1}{a} \frac{\Gamma(m/a)}{\Gamma(\frac{m}{a} + n + 1)}$

L.H.S. = $\int_0^1 x^{m-1} (1-x^a)^n dx$ put $x^a = \sin^2 \theta$, i.e. $x = \sin^{2/a} \theta$,
 so that $dx = \frac{2}{a} \sin^{(2/a)-1} \theta \cos \theta d\theta$

$$= \frac{2}{a} \int_0^{\pi/2} \sin^{2(m-1)/a} \theta \cdot \cos^{2n} \theta \cdot \sin^{(2/a)-1} \theta \cos \theta d\theta$$

$$= \frac{2}{a} \int_0^{\pi/2} \sin^{(2m/a)-1} \theta \cos^{(2n+1)} \theta d\theta$$

$$= \frac{2}{a} \cdot \frac{\Gamma(m/a) \Gamma(n+1)}{2\Gamma(\frac{m}{a} + n + 1)}$$

$$= \frac{2}{a} \cdot \frac{\Gamma(m/a) \Gamma(n+1)}{2\Gamma(\frac{m}{a} + n + 1)} = \frac{1}{a} \cdot \frac{\Gamma(m/a)}{\Gamma(\frac{m}{a} + n + 1)}$$

Problem 12. The equation of motion of a particle moving from rest towards a centre of attraction point, situated at distance 'a' cms. apart from it, is given by

$$\frac{d^2x}{dt^2} + \frac{k}{x} = 0, \quad \text{where } k \text{ is a constant}$$

Applying your knowledge of gamma functions, show that it will reach the centre of attraction point in a time given by

$$T = a \sqrt{\left(\frac{\pi}{2k}\right)} \quad (\text{Nagpur, 1961})$$

The given equation of motion is

$$\frac{d^2x}{dt^2} + \frac{k}{x} = 0$$

or

$$\frac{d^2x}{dt^2} = -\frac{k}{x}$$

Multiplying both sides by $2 \frac{dx}{dt} dt$, this equation may be written as

$$2 \frac{d^2x}{dt^2} \frac{dx}{dt} dt = -\frac{2k}{x} dx.$$

Integrating, $\left(\frac{dx}{dt}\right)^2 = -2k \log x + A$, A being a constant.

Initially when $x=a$, $\frac{dx}{dt} = 0$; $\therefore A = 2k \log a$.

We thus have

$$\left(\frac{dx}{dt}\right)^2 = -2k \log x + 2k \log a = 2k \log \frac{a}{x}$$

$$\frac{dx}{dt} = \sqrt{(2k)} \sqrt{\left(\log \frac{a}{x}\right)},$$

so that $\int_0^T dt = \frac{1}{\sqrt{(2k)}} \int_0^a \left(\log \frac{a}{x}\right)^{-1/2} dx$

$$T = \frac{a}{\sqrt{(2k)}} \int_0^a \left(\log \frac{a}{x}\right)^{-1/2} dx \quad \text{put } \log \frac{a}{x} = p$$

$$= -\frac{1}{\sqrt{(2k)}} \int_0^\infty (p)^{-1/2} e^{-p} dp \quad \text{or } x = ae^{-p}$$

$$= -\frac{a}{\sqrt{(2k)}} \int_0^\infty p^{(1/2)-1} e^{-p} dp \quad \therefore dx = -ae^{-p} dp$$

$$= -\frac{a}{\sqrt{(2k)}} \Gamma\left(\frac{1}{2}\right) \text{ by the definition of Gamma function}$$

$$= -\frac{a}{\sqrt{(2k)}} \sqrt{\pi}$$

$$= a \sqrt{\left(\frac{\pi}{2k}\right)}$$

6.10. ERROR FUNCTION OR PROBABILITY INTEGRAL

The error function denoted by $\text{erf}(x)$ is defined as

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-n^2} dn \quad \dots (1)$$

which arises in the solution of certain partial differential equations.

The definition (1) follows the properties of error function as

$$\text{erf}(-x) = -\text{erf}(x) \quad \dots (2)$$

$$\text{erf}(0) = 0 \quad \dots (3)$$

$$\text{erf}(\infty) = \frac{2}{\sqrt{\pi}} \int_0^\infty e^{-n^2} dx = \frac{2}{\sqrt{\pi}} \cdot \frac{\sqrt{\pi}}{2} \text{ by §6.3 (2)}$$

$$= 1 \quad \dots (4)$$

$$\text{and } \text{erf}(ix) = \frac{2i}{\sqrt{\pi}} \int_0^x e^{-n^2} dx, \text{ when } i = \sqrt{-1} \quad \dots (5)$$

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 13. Prove the following (i) $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$ (Agra, 1970)

$$(ii) \int_0^{\pi/2} \sqrt{\tan \theta} d\theta = \frac{1}{2} \Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{3}{4}\right) \text{ and (iii) } \int_0^1 \frac{35x^3}{32\sqrt{1-x}} dx = 1$$

(Nagpur, 1965)

Problem 14. Express the product $2.5 \dots 8(3n-1)$ in terms of the gamma function (Agra, 1966)

using $\Gamma(n+1) = n \Gamma(n)$ we have

$$\Gamma\left(n + \frac{2}{3}\right) = \left(n - \frac{1}{3}\right) \Gamma\left(n - \frac{1}{3}\right) = \left(n - \frac{1}{3}\right) \Gamma\left(n - \frac{4}{3}\right) \Gamma\left(n - \frac{7}{3}\right) = \dots$$

$$= \left(n - \frac{1}{3}\right) \left(n - \frac{4}{3}\right) \dots \frac{8}{3} \cdot \frac{5}{3} \cdot \frac{2}{3} \Gamma \frac{2}{3}$$

$$\text{Hence } 2.5.8 \dots (3n-1) = 3^n \Gamma\left(n + \frac{2}{3}\right) / \Gamma \frac{2}{3}$$

Problem 15. (a) Evaluate the following integrals (i) $\int_0^\infty \frac{\cos(xy)}{x^2 + b^2} dx$

$$(ii) \int_0^\infty e^{-by} \cos(xy) dy, b > 0.$$

(b) From the above results show that

$$\int_0^\infty \cos(xy) \cos(xy') dx = \frac{\pi}{2} \delta(y - y')$$

where $y > 0, y' > 0$ and $\delta(y - y')$ being the Dirac delta function. (Agra, 1968)

$$(a) (i) \text{ We have } \int_0^\infty 2z e^{-(b^2 + x^2)z^2} dz = -\frac{1}{b^2 + x^2} \left[e^{-(b^2 + x^2)z^2} \right]_0^\infty = \frac{1}{b^2 + x^2}$$

Multiplying both sides by $\cos xy$ and integrating within limits from $x=0$ to ∞ , we get

$$\int_0^\infty \frac{\cos xy}{b^2 + x^2} dx = \int_0^\infty \int_0^\infty \cos xy \cdot 2z e^{-(b^2 + x^2)z^2} dz dx = \int_0^\infty 2z e^{-b^2 z^2} \left[\int_0^\infty e^{-x^2 z^2} \cos xy dx \right] dz$$

$$\text{Here } I = \int_0^\infty e^{-x^2 z^2} \cos xy dx \text{ gives } \frac{dI}{dy} = - \int_0^\infty e^{-x^2 z^2} x \sin xy dx \\ = \left[\frac{e^{-x^2 z^2} \sin xy}{2z^2} \right]_0^\infty - \frac{y}{2z^2} \int_0^\infty e^{-x^2 z^2} \cos xy dx, \text{ (integrating by parts)} \\ = 0 - \frac{y}{2z^2} I$$

$$\therefore \frac{dI}{I} = - \frac{y}{2z^2} dy$$

Integrating $\log I = -\frac{y^2}{4z^2} + \log A$ i.e. $I = A e^{-y^2/4z^2}$ whence A can be determined by putting $y=0$ so that $A = \left[I \right]_{y=0} = \left[\int_0^\infty e^{-x^2 z^2} \cos xy dx \right]_{y=0}$

$$= \int_0^\infty e^{-x^2 z^2} dx = \frac{\sqrt{\pi}}{2z}$$

$$\text{As such } I = \int_0^\infty e^{-x^2 z^2} \cos xy dx = \frac{\sqrt{\pi}}{2z} e^{-y^2/4z^2}$$

$$\text{Thus } \int_0^\infty \frac{\cos xy}{b^2 + x^2} dx = \int_0^\infty 2z e^{-b^2 z^2} \cdot \frac{\sqrt{\pi}}{2z} e^{-y^2/4z^2} dz = \sqrt{\pi} \int_0^\infty e^{-b^2 z^2} e^{-\frac{y^2}{4z^2}} dz \\ = \sqrt{\pi} \int_0^\infty e^{-b^2 \left(z^2 + \frac{y^2}{4b^2 z^2} \right)} dz = \sqrt{\pi} J \text{ (say)}$$

$$\text{Now consider } J = \int_0^\infty e^{-b^2 \left(z^2 + \frac{y^2}{4b^2 z^2} \right)} dz$$

$$\text{so that } \frac{df}{dy} = -\frac{1}{2} \int_0^\infty e^{-b^2 \left(z^2 + \frac{y^2}{4b^2 z^2} \right)} \cdot \frac{y}{z^2} dz. \text{ Put } \frac{y}{2bz} = p \text{ so that } dp = -\frac{y}{2bz^2} dz$$

$$= +b \int_0^\infty e^{-b^2 \left(p^2 + \frac{y^2}{4b^2 p^2} \right)} dp = -b \int_0^\infty e^{-b^2 \left(z^2 + \frac{y^2}{4b^2 z^2} \right)} dz, \\ \text{(on replacing } p \text{ by } z)$$

$$= -b J$$

$$\frac{dJ}{J} = -b dy \text{ which gives on integration } J = B e^{-by} \text{ where } B \text{ is determined by}$$

$$\text{putting } y=0, \text{ so that } B = \left[J \right]_{y=0} = \int_0^\infty e^{-b^2 z^2} dz = \frac{\sqrt{\pi}}{2b} \text{ and hence } J = \frac{\sqrt{\pi}}{2b} e^{-by}$$

$$\text{Thus } \int_0^\infty \frac{\cos xy}{b^2 + x^2} dx = \sqrt{\pi} J = \sqrt{\pi} \cdot \frac{\sqrt{\pi}}{2b} e^{-by} = \frac{\pi}{2b} e^{-by}$$

$$(ii) \text{ (say) } I = \int_0^\infty e^{-by} \cos xy dy = \left[-\frac{e^{-by}}{b} \cos xy \right]_0^\infty - \frac{x}{b} \int_0^\infty e^{-by} \sin xy dy \\ \text{(integrating by parts)}$$

$$= \frac{1}{b} - \frac{x}{a} \left[-\frac{e^{-by}}{b} \sin xy \right]_0^\infty - \frac{x^2}{b^2} \int_0^\infty e^{-by} \cos xy dy = \frac{1}{b} - 0 - \frac{x^2}{b^2} I$$

$$\text{so that } I \left[1 + \frac{x^2}{b^2} \right] = \frac{1}{b} \text{ or } I = \frac{1}{b} \int_0^\infty e^{-by} \cos xy dy = \frac{b}{x^2 + b^2}$$

(b) This follows from the definition of Dirac delta function given in §8.10.

Problem 16. (a) Evaluate the following integrals

$$(i) \int_{-\infty}^\infty e^{-\alpha |x| + i kx} dx, \alpha > 0 \text{ and } |x| \text{ is the absolute value of } x$$

$$(ii) \frac{1}{\beta} \int_0^\infty (x^\beta - 1) e^{-\alpha x} dx$$

(b) In the second integral given above, take the limit $\beta \rightarrow 0$ and hence evaluate integral $\int_0^\infty \log x \cdot e^{-\alpha x} dx$

(Use the approximations $\alpha^\beta \cong 1 + \beta \log \alpha$ and $\Gamma(1 + \beta) \cong 1 + \xi \beta$, $\xi = -0.577218$ for small β . (Agra, 1967))

$$(i) \text{ Say } I = \int_{-\infty}^\infty e^{-\alpha |x| + i kx} dx = \int_{-\infty}^\infty e^{-\alpha |x|} (\cos kx + i \sin kx) dx \\ = \int_{-\infty}^\infty e^{-\alpha |x|} \cos kx dx + i \int_{-\infty}^\infty e^{-\alpha |x|} \sin kx dx = I_1 + i I_2 \text{ (say)} \\ \dots (1)$$

Now $|x|$ is absolutely positive if x is negatively large or positively large and α is always positive also the values of $\cos kx$ and $\sin kx$ are always finite lying between -1 and $+1$ and hence both the integrals I_1 and I_2 being convergent exist.

$$\text{We have } I_2 = \int_{-\infty}^0 e^{-\alpha |x|} \sin kx dx + \int_0^\infty e^{-\alpha |x|} \sin kx dx = I_{21} + I_{22} \text{ (say)}$$

$$\text{Replacing } x \text{ by } -x, I_{21} = - \int_0^\infty e^{-\alpha |x|} \sin kx dx.$$

$$\text{So that } I_2 = I_{21} + I_{22} = - \int_0^\infty e^{-\alpha |x|} \sin kx dx + \int_0^\infty e^{-\alpha |x|} \sin kx dx = 0$$

Hence $I = I_1 = \int_{-\infty}^\infty e^{-\alpha |x|} \cos kx dx = 2 \int_0^\infty e^{-\alpha |x|} \cos kx dx$, the integrand being even function of x

$$= 2 \int_0^\infty e^{-\alpha x} \cos kx dx = \frac{2b}{x^2 + b^2} \text{ by Problem 15 (a-ii)}$$

(ii) We have $\frac{1}{\beta} \int_0^\infty (x^\beta - 1) e^{-\alpha x} dx = \frac{1}{\beta} \int_0^\infty e^{-\alpha x} x^{\beta+1-1} dx - \frac{1}{\beta} \int_0^\infty e^{-\alpha x} dx$

$$= \frac{1}{\beta} \left[\frac{\Gamma(\beta+1)}{\alpha^{\beta+1}} - \frac{1}{\alpha} \right] \therefore \int_0^\infty e^{-\alpha x} x^{\beta-1} dx = \frac{\Gamma\beta}{\alpha^\beta} \text{ and for } \beta=1, \text{ this gives}$$

$$\int_0^\infty e^{-\alpha x} dx = \frac{1}{\alpha}$$

$$= \frac{1}{\alpha} \left[\frac{\Gamma(\beta+1)}{\beta \alpha^\beta} - \frac{1}{\beta} \right]$$

(b) In (ii) we have shown that $\frac{1}{\beta} \int_0^\infty (x^\beta - 1) e^{-\alpha x} dx = -\frac{1}{\alpha} \left[\frac{\Gamma(\beta+1)}{\beta \alpha^\beta} - \frac{1}{\beta} \right]$

$$= -\frac{1}{\alpha} \left[\frac{\Gamma(\beta+1) - \alpha^\beta}{\beta \alpha^\beta} \right]$$

when $\beta \rightarrow 0$, both sides of this relation are of the form $\frac{0}{0}$ and hence taking limit as $\beta \rightarrow 0$, we have

$$\frac{\lim_{\beta \rightarrow 0} \frac{d}{d\beta} (x^\beta - 1) e^{-\alpha x} dx}{\lim_{\beta \rightarrow 0} \frac{d}{d\beta} (\beta)} = \lim_{\beta \rightarrow 0} \left[\frac{(1+\xi\beta) - (1+\beta \log \alpha)}{\beta(1+\beta \log \alpha)} \right] \cdot \frac{1}{\alpha}$$

$$\text{or } \int_0^\infty \log x e^{-\alpha x} dx = \lim_{\beta \rightarrow 0} \frac{1}{\alpha} \left[\frac{\beta(\xi - \log \alpha)}{\beta(1+\beta \log \alpha)} \right] = \lim_{\beta \rightarrow 0} \left[\frac{\xi - \log \alpha}{1+\beta \log \alpha} \cdot \frac{1}{\alpha} \right]$$

$$= \frac{\xi - \log \alpha}{\alpha}$$

Problem 17. Show by integrating by parts that

$$\int_0^\infty e^{-n^2} dn = \frac{\sqrt{\pi}}{2} - \frac{e^{-x^2}}{2x} \left(1 - \frac{1}{2x^2} + \frac{1 \times 3}{2^2 x^4} \right) + \frac{1 \times 3 \times 5}{2^3} \int_x^\infty \frac{e^{-n^2}}{n^6} dx$$

(Agra, 1971)

Show how this expression may be used to compute the value of $\text{erf}(x)$ for large values of x .

Problem 18. Define the error function $\text{erf}(x)$.

Problem 19. Define Gamma and Beta functions.

Problem 20. If $\Gamma z = \int_0^\infty e^{-t} t^{z-1} dt$, evaluate

$$\int_0^{\pi/2} \cos^{m-1} x \sin^{n-1} x dx \text{ in terms of gamma function.} \quad (\text{Agra, 1975})$$

Problem 21. Define Beta and Gamma functions. Show that

$$(a) \Gamma(n+1) = n\Gamma n \quad (b) \beta(m, n) = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$$

$$(c) \Gamma \frac{1}{2} = \sqrt{\pi} \quad (d) \Gamma \frac{1}{2} \Gamma \frac{3}{2} = \pi 2^{1/2} \quad (\text{Agra, 1976})$$

Problem 22. (a) Derive the recurrence relation $\Gamma(z+1) = z\Gamma z$ from Euler's integral $\Gamma z = \int_0^\infty e^{-t} t^{z-1} dt$; $R.P. (z) > 0$

(b) Show by means of beta function that

$$\int_0^1 \frac{dx}{(z-x)^{1-\alpha} (x-t)^\alpha} = \frac{\pi}{\sin \pi \alpha}; 0 < \alpha < 1 \quad (\text{Rohilkhand, 1976})$$

(i) See §6.2.

$$(ii) \text{ Let } I = \int_t^z \frac{dx}{(z-x)^{1-\alpha} (x-t)^\alpha} = \int_t^z (x-t)^{-\alpha} (z-x)^{\alpha-1} dx$$

Put $x = t \sin^2 \theta + z \cos^2 \theta$, so that $dx = -2(z-t) \sin \theta \cos \theta d\theta$

$$x-t = (z-t) \cos^2 \theta \text{ and } z-x = (z-t) \sin^2 \theta$$

Also when $x=z$, $\sin^2 \theta = 0 \Rightarrow \theta = 0$ and $x=t$, $\cos^2 \theta = 0 \Rightarrow \theta = \frac{\pi}{2}$. With these substi-

tutions we have

$$I = \int_0^{\pi/2} \frac{(z-t)^{-\alpha} \cos^{-2\alpha} \theta \cdot (z-t)^{\alpha-1} \sin^{2\alpha-2} \theta \cdot 2(z-t) \sin \theta \cos \theta d\theta}{(z-t)^{1-\alpha} \cos^{2\alpha} \theta}$$

$$= 2 \int_0^{\pi/2} \sin^{2\alpha-2} \theta \cos^{1-2\alpha} \theta d\theta$$

$$= 2 \cdot \frac{\Gamma \alpha \Gamma \{1-\alpha\}}{\Gamma \{ \alpha + (1-\alpha) \}} \text{ by property of Gamma function}$$

$$= \frac{\Gamma \alpha \Gamma (1-\alpha)}{\Gamma \{ \alpha + (1-\alpha) \}} = \beta(\alpha, 1-\alpha) \therefore \beta(m, n) = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$$

$$= \int_0^\infty \frac{y^{\alpha-1}}{1+y} dy \text{ by (2) of §6.5.}$$

$$\text{But we have } \int_0^\infty \frac{y^{2m}}{1+y^{2n}} = \frac{\pi}{2n} \operatorname{cosec} \frac{2m+1}{2n} \pi \text{ by §6.8.}$$

On putting $n = \frac{1}{2}$, $m = \frac{\alpha-1}{2}$, this yields

$$I = \int_0^\infty \frac{y^{\alpha-1}}{1+y} dy = \pi \operatorname{cosec} \pi \alpha = \frac{\pi}{\sin \pi \alpha} \text{ when } 0 < \alpha < 1.$$

After, We have, by definition of Beta function,

$$\beta(m, n) = \int_0^1 y^{m-1} (1-y)^{n-1} dy$$

$$\text{Put } y = \frac{x-t}{z-t} \text{ so that } dy = \frac{dx}{z-t}$$

$$\beta(m, n) = \int_t^z \frac{(x-t)^{m-1}}{(z-t)^{m-1}} \cdot \frac{(z-x)^{n-1}}{(z-t)^{n-1}} \cdot \frac{dx}{z-t}$$

$$= \frac{1}{(z-t)^{m+n-1}} \int_t^z (x-t)^{m-1} (z-x)^{n-1} dx$$

$$\text{or } \int_t^z (x-t)^{m-1} (z-x)^{n-1} dx = (z-t)^{m+n-1} \beta(m, n)$$

If we put $n = \alpha$, $m = 1-\alpha$, this yields

$$\int_t^z \frac{dx}{(z-x)^{1-\alpha} (x-t)^\alpha} = \beta(\alpha, 1-\alpha)$$

$$= \frac{\Gamma \alpha \Gamma (1-\alpha)}{\Gamma(\alpha+1-\alpha)} = \Gamma \alpha \Gamma (1-\alpha).$$

$$= \frac{\pi}{\sin \pi \alpha} \therefore \Gamma n \Gamma (1-n) = \frac{\pi}{\sin \pi n}$$

DIFFERENTIAL EQUATIONS

(ORDINARY AND PARTIAL WITH FINITE DIFFERENCE
i.e. INTERPOLATION)

7.1. INTRODUCTION

In mathematics we call the changing entities as *variables* and the rate of change of one variable with respect to another as a *derivative*. Equations expressing a relationship among these variables and their derivatives are known as *Differential equations*. In other words differential equations originate whenever a universal law is expressed by means of variables and their derivatives.

A relationship between two variables x and y is expressed as $f(x, y) = 0$ and read as " f of x, y equals zero" or as "a function of x, y equals zero". A function defining y as a function of x in the form $f(x, y) = 0$ is said to be an *implicit function* of x . A special class of constant functions is referred as *elementary functions*. An equation involving x , the function $f(x)$ which defines a function of x and one or more of its derivatives is called an *ordinary differential equation*. In other words an ordinary differential equation involves only one independent variable say x such as $\frac{dy}{dx} = \frac{1}{x}$ while a *partial differential equation* is one which involves two or more independent variables x and y such as $\frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} = x^2$. The *order* of a differential equation is the order of the highest derivative involved in that equation e.g.,

$\frac{d^2y}{dx^2} = \frac{1}{x}$ is of order 2.
 $\frac{d^3y}{dx^3} = \frac{1}{x^3}$ is of order 3.

If $y = f(x)$ defines y as a function of x on any interval, then $f(x)$ is said to be an *explicit solution* or simply a *solution* of an ordinary differential equation involving $x, f(x)$ and its derivatives. Whenever it can be shown that an implicit function does satisfy a given differential equation on any interval then $f(x, y) = 0$ is said to be an *implicit solution* of the differential equation.

The solution of a differential equation of order n contains n arbitrary constants say $c_1, c_2, \dots, c_n$ and called an *n -parameter family of solutions*, referring to the constants as parameters.

An n -parameter family of solutions of an n th order differential equation is called its *general solution* and the function obtained by giving a definite set of values to the constants $c_1, c_2, \dots, c_n$ in the family is said to be the *Particular Integral* of the differential equation.

There are differential equations which have solutions not obtainable from the n -parameter family whatsoever values are given to the constants, such solutions are called *Singular solutions*.

The n conditions which enable us to find the values of n arbitrary constants in an n -parameter family if given in terms of one value of the independent variable, are called *Initial conditions* or *boundary conditions*.

The reader of this book is presumed to have passed through the degree level where the methods of solving ordinary differential equations of first order are discussed in details. In the next section we discuss some methods of solving ordinary differential equations of second order which are of physical interest.

7.2. SOLUTION IN SIMPLE CASES OF ORDINARY DIFFERENTIAL EQUATIONS OF SECOND ORDER

Generally an ordinary equation of second order is of the form

$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = X$$

where P, Q, X are functions of x only.

1. Linear Equations with Constant Coefficients.

The general solution is found by usual methods of finding the complementary function and particular integral of the equation. Although the student is presumed to have a sound knowledge of the methods to be employed for finding the complementary function and particular integral, but still then, we summarize them as below:

Let there be a differential equation of the type

$$\frac{d^2y}{dx^2} + p_1 \frac{dy}{dx} + p_2 y = X$$

where p_1, p_2 are constants and X is a function of x .

In terms of the D operator, it may be written

$$(D^2 + p_1 D + p_2)y = X \text{ where } D \text{ stands for } \frac{d}{dx} \text{ i.e., } D \equiv \frac{d}{dx}$$

or we may write thus, $f(D)y = X$.

To find the *complementary function* (C.F.). The X is removed and replaced by zero. Then an auxiliary equation is written either by putting $y = e^{mx}$ whence, we get

$$m^2 + p_1 m + p_2 = 0,$$

or simply writing $D^2 + p_1 D + p_2 = 0$ i.e., $f(D) = 0$.

In either case, we get the roots of the quadratic.

Case I. If the roots are of the type m_1 and m_2 (real and distinct) C.F. is $C_1 e^{m_1 x} + C_2 e^{m_2 x}$.

Case II. If $m_1 = m_2$ i.e., both the roots are real and equal, C.F. is $(C_1 + C_2 x) e^{m_1 x}$.

Case III. If the roots are imaginary i.e., of the type $\alpha \pm i\beta$, C.F. is $e^{\alpha x} [C_1 \cos \beta x + C_2 \sin \beta x]$ or $C_1 e^{\alpha x} \cos (\beta x + C_2)$.

Case IV. If the roots are of the type $\alpha \pm \sqrt{\beta}$, C.F. is $C_1 e^{\alpha x} \cosh (\sqrt{\beta} x + C_2)$.

Note. The number of the arbitrary constants will be the same as the order of the equation.

To find the Particular Integral (P.I.)

We have $P.I. = \frac{X}{f(D)}$.

Case I. If $X = e^{ax}$ where a is any constant.

$$P.I. = \frac{e^{ax}}{f(D)} = \frac{e^{ax}}{f(a)} \text{ if } f(a) \neq 0.$$

Case II. If $X = x^m$, where m is a positive integer

$$P.I. = \frac{x^m}{f(D)} = [f(D)]^{-1} x^m.$$

Expand $[f(D)]^{-1}$ binomially upto m th power of D and then operate x^m on every term.

Case III. If $X = \sin ax$ or $\cos ax$.

$$P.I. = \frac{\sin ax \text{ or } \cos ax}{f(D^2)} = \frac{\sin ax \text{ or } \cos ax}{f(-a^2)} \text{ provided } f(-a^2) \neq 0.$$

In case $f(-a^2) = 0$, $\frac{\sin ax}{f(D^2)} = \text{Imaginary part of } \frac{e^{iax}}{f(D^2)}$ and $\frac{\cos ax}{f(D^2)} = \text{Real part of } \frac{e^{iax}}{f(D^2)}$, which is case I.

Case IV. If $X = e^{ax} V$, where V is any function of x , then

$$P.I. = \frac{e^{ax} V}{f(D)} = e^{ax} \cdot \frac{1}{f(D+a)} \cdot V.$$

Case V. If $X = x \cdot V$, where V is any function of x , then

$$P.I. = \frac{xV}{f(D)} = x \frac{1}{f(D)} \cdot V - \frac{1}{f(D)} \cdot f'(D) \frac{1}{f(D)} V \\ = \left\{ x - \frac{1}{f(D)} f'(D) \right\} \frac{1}{f(D)} V$$

Hence general solution = C.F. + P.I.

Problem 1. Solve $\frac{d^2 y}{dx^2} - y = x \sin x + (1+x^2) e^x$
or $(D^2 - 1)y = x \sin x + (1+x^2) e^x$

Now its complementary function and particular integral may be found thus:

For complementary function, the auxiliary equation is

$$D^2 - 1 = 0 \text{ or } D = \pm 1.$$

$$\therefore \text{C.F. is } C_1 e^x + C_2 e^{-x}.$$

$$\text{Particular integral} = \frac{x \sin x}{D^2 - 1} + \frac{(1+x^2) e^x}{D^2 - 1}$$

$$= \text{imaginary part in } \frac{x e^{ix}}{D^2 - 1} + e^x \frac{1}{(D+1)^2 - 1} \cdot (1+x^2)$$

$$= \text{imaginary part in } e^{ix} \cdot \frac{1}{(D+i)^2 - 1} \cdot x + e^x \cdot \frac{1}{D^2 + 2D} (1+x^2)$$

$$= \text{imaginary part in } e^{ix} \cdot \frac{1}{D^2 + 2iD - 2} \cdot x + \frac{e^x \cdot 1}{2D \left(1 + \frac{D}{2} \right)} (1+x^2)$$

$$= \text{imaginary part in } \frac{e^{ix}}{-2} \left[1 - \frac{2iD + D^2}{2} \right]^{-1} x \\ + \frac{e^x}{2} \cdot D^{-1} \left\{ 1 + \frac{D}{2} \right\}^{-1} \cdot (1+x^2)$$

$$= \text{imaginary part in } \frac{e^{ix}}{-2} \left[1 + \frac{2iD + D^2}{2} + \dots \right] \cdot x \\ + \frac{e^x}{2} D^{-1} \left[1 - \frac{D}{2} + \frac{D^2}{4} - \dots \right] (1+x^2)$$

$$= \text{imaginary part in } \frac{e^{ix}}{-2} [x + i] + \frac{e^x}{2} D^{-1} [1 + x^2 - x + \frac{1}{2}]$$

$$= \text{imaginary part in } \left(\frac{\cos x + i \sin x}{-2} \right) (x + i) \\ + \frac{e^x}{2} \int (x^2 - x + \frac{1}{2}) dx$$

$$= -\frac{1}{2} (x \sin x + \cos x) + \frac{e^x}{2} \left[\frac{x^3}{3} - \frac{x^2}{2} + \frac{3}{2} x \right]$$

Hence the general solution is

$$y = C_1 e^x + C_2 e^{-x} - \frac{1}{2} (x \sin x + \cos x) + \frac{x e^x}{12} (2x^2 - 3x + 9).$$

II. Linear Equations with Variable Coefficients (Homogeneous Linear Equations).

Consider, $p_0 x^2 \frac{d^2 y}{dx^2} + p_1 x \frac{dy}{dx} + p_2 y = X$.

Put $x = e^z$, i.e., $z = \log x$; $\therefore \frac{dz}{dx} = \frac{1}{x}$.

Then $\frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx} = \frac{1}{x} \frac{dy}{dz}$,

$$\frac{d^2 y}{dx^2} = \frac{1}{x} \frac{d}{dz} \left(\frac{1}{x} \frac{dy}{dz} \right) = \frac{1}{x^2} \frac{d^2 y}{dz^2} - \frac{1}{x^3} \frac{dx}{dz} \frac{dy}{dz} \\ = \frac{1}{x^2} \left(\frac{d^2 y}{dz^2} - \frac{dy}{dz} \right)$$

If we put $\frac{d}{dz} \equiv D$, we have

$$x \frac{dy}{dx} = Dy, \quad x^2 \frac{d^2 y}{dx^2} = D(D-1)y.$$

Now substituting these values in (1), we get

$$p_0 D(D-1)y + p_1 Dy + p_2 y = X,$$

which may be solved by the method employed in I.

Problem 2. Solve $x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + y = 2 \log x$.

Put $x = e^z$ and denote $\frac{d}{dz}$ by D ; we have

$$D(D-1)y - Dy + y = 2z \text{ or } (D^2 - 2D + 1)y = 2z.$$

Auxiliary equation is

$$D^2 - 2D + 1 = 0$$

or $(D-1)^2 = 0$; $\therefore D=1$ (repeated twice).

$$\text{C.F.} = (c_1 + c_2 z)e^z.$$

$$\text{P.I.} = \frac{2z}{D^2 - 2D + 1} = 2[1 - (2D - D^2)]^{-1} \cdot z$$

$$= 2(1 + 2D + \dots)z = 2(z + 2) = 2z + 4.$$

$\therefore$ General solution is $y = (c_1 + c_2 z)e^x + 2z + 4$

$$= (c_1 + c_2 \log x)x + 2 \log x + 4.$$

Note. Any equation of the type

$$(a+bx)^n \frac{d^n y}{dx^n} + P_1 (a+bx)^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots$$

$$\dots + P_{n-1} (a+bx) \frac{dy}{dx} + P_n y = F(x)$$

can be reduced to the homogeneous linear form by putting $z = ax + b$ or this can be solved by putting $ax + b = e^z$ as above.

III. Exact Differential Equations and Equations of other Special types.

The equation of the type

$$P_0 \frac{d^2 y}{dx^2} + P_1 \frac{dy}{dx} + P_2 y = 0,$$

where P_0 , P_1 and P_2 are the functions of x , is said to be exact if

$$P_2 - P_1' + P_0'' = 0$$

or in general an equation of order n (say),

$$P_0 \frac{d^n y}{dx^n} + P_1 \frac{d^{n-1} y}{dx^{n-1}} + \dots + P_n y = Q(x)$$

is exact if $P_n - P_{n-1}' + \dots + (-1)^n P_0^{(n)} = 0$,

where P' , P'' , ..., $P^{(n)}$ are the successive derivatives of P .

In case the equation is exact, its first integral is

$$P_0 y_{n-1} + (P_1 - P_0') y_{n-2} + (P_2 - P_1' + P_0'') y_{n-3} + \dots$$

$$\dots + \{P_{n-1} - P_{n-2}' + \dots + (-1)^n P_0^{(n-1)}\} y = \int Q(x) dx + C,$$

where y_n stands for $\frac{d^n y}{dx^n}$ etc.

Problem 3. Solve $\sin^2 x \frac{d^2 y}{dx^2} = 2y$

$$\frac{d^2 y}{dx^2} - 2y \operatorname{cosec}^2 x = 0$$

$$\sin x \frac{d^2 y}{dx^2} - 2y \cot x \operatorname{cosec}^2 x = 0,$$

which is exact and hence its first integral is

$$\sin x \frac{dy}{dx} + y \operatorname{cosec}^2 x = c_1$$

$$\frac{dy}{dx} + \frac{1}{\sin x \cos x} y = c_1 \tan x,$$

which is a linear differential equation of the first order.

$$\begin{aligned} \text{Integrating factor} &= e^{\int \frac{dx}{\sin x \cos x}} = e^{\int 2 \operatorname{cosec} 2x dx} \\ &= e^{\log \tan x} = \tan x. \end{aligned}$$

$$\text{The solution is } y \tan x = \int c_1 \tan^2 x dx + c_2$$

$$= c_1 \int (\sec^2 x - 1) dx + c_2$$

$$= c_1 (\tan x - x) + c_2.$$

Note 1. Sometimes the equation becomes exact by multiplying an integrating factor x^m , where m can be found by applying the condition of exactness.

Note 2. Equations of the form $\frac{d^2 y}{dx^2} = f(y)$ can be integrated by

multiplying by $2 \frac{dy}{dx}$, whence we get

$$\left(\frac{dy}{dx}\right)^2 = 2 \int f(y) dy + c_1$$

$$\frac{dy}{dx} = \sqrt{c_1 + 2 \int f(y) dy}$$

which may further be integrated by any of the standard methods.

Note 3. Equations not containing x directly can be integrated by

$$\text{putting } \frac{dy}{dx} = p.$$

$$\text{Problem 4. } \frac{d^2 y}{dx^2} + \frac{dy}{dx} + \left(\frac{dy}{dx}\right)^3 = 0.$$

$$\text{Put } \frac{dy}{dx} = p; \therefore \frac{d^2 y}{dx^2} = p \frac{dp}{dy}.$$

$$\text{So we get } p \frac{dp}{dy} + p + p^3 = 0.$$

or $\frac{dp}{1+p^2} = -dy$.

Integrating $\tan^{-1} p = c - y$, i.e., $\frac{dy}{dx} = p = \tan(c - y)$,

which can further be integrated by standard methods.

Similarly equations not containing y can be integrated by

$$\frac{dy}{dx} = p \left(\because \frac{d^2y}{dx^2} = \frac{dp}{dx} = p \frac{dp}{dy} \text{ etc.} \right).$$

Note. 4. Equations in which y appears in only two derivatives of orders differ by unity can be integrated as follows:

Problem 5. $a \frac{d^2y}{dx^2} = \left[1 + \left(\frac{dy}{dx} \right)^2 \right]^{1/2}$

Put $\frac{dy}{dx} = q$; $\therefore \frac{d^2y}{dx^2} = \frac{dq}{dx}$. We have

$$a \frac{dq}{dx} = (1 + q^2)^{1/2}$$

or $\frac{dq}{\sqrt{1+q^2}} = \frac{dx}{a}$.

Integrating, $\sinh^{-1} q = \frac{x}{a} + c_1$.

$\therefore q = \frac{dy}{dx} = \sinh \left(\frac{x}{a} + c_1 \right)$.

Integrating again, $y = a \cosh \left(\frac{x}{a} + c_1 \right) + c_2$.

IV. The Complete Solutions in Terms of a Known Integral.

Let $y = y_1$ be a known integral in the complementary function

$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = X.$$

Put $y = vy_1$; so that $\frac{dy}{dx} = y_1 \frac{dv}{dx} + v \frac{dy_1}{dx}$

and $\frac{d^2y}{dx^2} = y_1 \frac{d^2v}{dx^2} + 2 \frac{dv}{dx} \cdot \frac{dy_1}{dx} + v \frac{d^2y_1}{dx^2}$.

With these substitutions, (1) gives

$$y_1 \left(\frac{d^2v}{dx^2} \right) + \frac{dv}{dx} \left(2 \frac{dy_1}{dx} + P y_1 \right) + v \left(\frac{d^2y_1}{dx^2} + P \frac{dy_1}{dx} + Q y_1 \right) = X$$

or $\frac{d^2v}{dx^2} + \left(P + \frac{2}{y_1} \frac{dy_1}{dx} \right) \frac{dv}{dx} = \frac{X}{y_1}$,

since $\frac{d^2y_1}{dx^2} + P \frac{dy_1}{dx} + Q y_1 = 0$ by hypothesis.

Putting $\frac{dv}{dx} = p$, (2) becomes

$$\frac{dp}{dx} + \left(P + \frac{2}{y_1} \frac{dy_1}{dx} \right) p = \frac{X}{y_1},$$

which is a linear equation of first order and hence its integrating

factor $= e^{\int \left(P + \frac{2}{y_1} \frac{dy_1}{dx} \right) dx} = e^{\int P dx + 2 \log y_1} = y_1^2 e^{\int P dx}$

Hence solution of (3) in p is

$$p \cdot y_1^2 e^{\int P dx} = \left[\int y_1^2 \cdot \frac{X}{y_1} e^{\int P dx} dx \right] + c_1$$

or $p = \frac{dy}{dx} = \frac{c_1 e^{-\int P dx}}{y_1^2} + \frac{e^{-\int P dx}}{y_1^2} \int (y_1 \cdot X \cdot e^{\int P dx}) dx$

Integrating,

$$y = c_2 + c_1 \int \frac{e^{-\int P dx}}{y_1^2} dx + \int \frac{e^{-\int P dx}}{y_1^2} \int (y_1 \cdot X \cdot e^{\int P dx} (dx)^2) dx$$

Hence the required solution of (1) is

$$y = v y_1 = c_2 y_1 + c_1 y_1 \int \frac{e^{-\int P dx}}{y_1^2} dx + y_1 \int \frac{e^{-\int P dx}}{y_1^2} \times \int (y_1 \cdot X \cdot e^{\int P dx}) (dx)^2.$$

Problem 6. Solve $x^2 \frac{d^2y}{dx^2} - (x^2 + 2x) \frac{dy}{dx} + (x + 2)y = x^3 \cdot e^x$.

Obviously $y = x$ is a part of the complementary.

Putting $y = vx$; the given equation gives

$$\frac{d^2y}{dx^2} + \left(\frac{2}{x} - \frac{x^2 + 2x}{x^2} \right) \frac{dv}{dx} = \frac{x \cdot e^x}{x}.$$

Now put $\frac{dv}{dx} = p$; then we get

$$\frac{dp}{dx} - p = e^x.$$

Being a linear equation of first order in p ,

Integrating factor $= e^{-\int dx} = e^{-x}$.

Thus $p \cdot e^{-x} = \int e^x \cdot e^{-x} dx + C_1$

$$= x + C_1$$

$$p = \frac{dv}{dx} = (x + C_1) e^x.$$

$\therefore$ Required solution is $y = vx = (x^2 - x + C_1 x) e^x + C_2 x$.

Note. It will be helpful to find that

$y = x$	"	"	"	$P + Qx = 0,$
$y = x^2$	"	"	"	$2 + 2Px + Qx^2 = 0,$
$y = e^x$	"	"	"	$P + Q + 1 = 0,$
$y = e^{-x}$	"	"	"	$1 - P + Q = 0,$
$y = e^{ax}$	"	"	"	$1 + \frac{P}{a} + \frac{Q}{a^2} = 0 \text{ etc.}$

V. Transformation of Equation

(i) By Changing the Dependent Variable

$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = X. \quad \dots(4)$$

Putting $y = vy_1$ this becomes

$$\frac{d^2v}{dx^2} + \left(P + \frac{2}{y_1} \frac{dy_1}{dx} \right) \frac{dv}{dx} + \frac{v}{y_1} \left(\frac{d^2y_1}{dx^2} + P \frac{dy_1}{dx} + Qy_1 \right) = \frac{X}{y_1}. \quad \dots(5)$$

Choosing v_1 such that $\frac{d^2y_1}{dx^2} + P \frac{dy_1}{dx} + Qy_1 = 0$,

$$(5) \text{ reduces to } \frac{d^2v}{dx^2} + P_1 \frac{dv}{dx} = \frac{X}{y_1}, \quad \dots(6)$$

where

$$P_1 = P + \frac{2}{y_1} \frac{dy_1}{dx}.$$

Now equation (6) may be solved as in previous method.

Problem 7.

$$\frac{d^2y}{dx^2} + \frac{2}{x} \frac{dy}{dx} = n^2y.$$

Here

$$P_1 = \frac{2}{x}.$$

$$\text{Take } y_1 = e^{-\frac{1}{2} \int \frac{2}{x} dx} = e^{-\log x} = 1/x.$$

Now put $y = vy_1$ whence the given equation becomes

$$\frac{d^2v}{dx^2} - n^2v = 0.$$

Its solution is $v = C_1 e^{nx} + C_2 e^{-nx}$.

The required solution is $y = vy_1 = \frac{1}{x} (C_1 e^{nx} + C_2 e^{-nx})$

(ii) By Changing the Independent Variable

$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = X. \quad \dots(7)$$

We know that

$$\frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx} \text{ and } \frac{d^2y}{dx^2} = \frac{d^2y}{dz^2} \left(\frac{dz}{dx} \right)^2 + \frac{dy}{dz} \cdot \frac{d^2z}{dx^2}$$

Substituting these values in (7), we get

$$\frac{d^2y}{dz^2} \left(\frac{dz}{dx} \right)^2 + \frac{dy}{dz} \cdot \frac{d^2z}{dx^2} + P \frac{dy}{dz} \cdot \frac{dz}{dx} + Qy = X$$

or

$$\frac{d^2y}{dz^2} + \frac{\frac{d^2z}{dx^2} + P \frac{dz}{dx}}{\left(\frac{dz}{dx} \right)^2} \cdot \frac{dy}{dz} + \frac{Q}{\left(\frac{dz}{dx} \right)^2} y = \frac{X}{\left(\frac{dz}{dx} \right)^2}$$

or

$$\frac{d^2y}{dz^2} + P_1 \frac{dy}{dz} + Q_1 y = X_1, \quad \dots(8)$$

where

$$P_1 = \frac{\frac{d^2z}{dx^2} + P \frac{dz}{dx}}{\left(\frac{dz}{dx} \right)^2}, Q_1 = \frac{Q}{\left(\frac{dz}{dx} \right)^2} \text{ and } X_1 = \frac{X}{\left(\frac{dz}{dx} \right)^2}$$

If z be chosen such that $\frac{d^2z}{dx^2} + P \frac{dz}{dx} = 0$,

i.e.,

$$z = \int e^{-\int P dx} dx,$$

then the given equation changes into

$$\frac{d^2y}{dz^2} + Q_1 y = X_1,$$

which can be solved if Q_1 is a constant or a constant multiplied by $\frac{1}{z^2}$.

Again if z be chosen that Q_1 be a^2 (a constant), then

$$a^2 \left(\frac{dz}{dx} \right)^2 = Q,$$

i.e.,

$$az = \int \sqrt{Q} dx.$$

With this substitution (7) reduces to

$$\frac{d^2y}{dz^2} + P_1 \frac{dy}{dz} + a^2 y = X,$$

which can be solved if P_1 is a constant.

Problem 8. Solve $\frac{d^2y}{dx^2} + \frac{2}{x} \frac{dy}{dx} + \frac{a^4}{x^4} y = 0$.

We can choose z such that

$$\left(\frac{dz}{dx} \right)^2 = Q = \frac{a^2}{x^4},$$

giving

$$z = + \frac{a}{x}.$$

Now changing the independent variable from x to z when $z = \frac{a}{x}$, we have

$$P_1 = \frac{\left(\frac{d^2z}{dx^2} + P \frac{dz}{dx} \right)}{\left(\frac{dz}{dx} \right)^2} = 0.$$

The given equation reduces to

$$\frac{d^2y}{dz^2} + y = 0.$$

Its solution is $y = C_1 \cos z + C_2 \sin z$.

Required solution is $y = C_1 \cos \frac{a}{x} + C_2 \sin \frac{a}{x}$.

7.3. SIMPLE PROBLEMS FROM PHYSICS

Problem 9. Solve $L \frac{di}{dt} + \frac{1}{C} \int i dt = 0$, the differential equation which means that the self-induction and capacity in a circuit neutralize each other. Determine the constants in such a way that I is the maximum current and $i=0$ when $t=0$.

Given
$$L \frac{di}{dt} + \frac{1}{C} \int i dt = 0.$$

or
$$\frac{di}{dt} + \frac{1}{LC} \int i dt = 0.$$

Differentiating, we get
$$\frac{d^2 i}{dt^2} + \frac{i}{LC} = 0.$$

Its auxiliary equation is $(D^2 + \frac{1}{LC}) = 0$, when $D \equiv \frac{d}{dt}$

or
$$D = \pm \sqrt{-1} \frac{1}{\sqrt{LC}}.$$

Solution is $i = A \cos \frac{t}{\sqrt{LC}} + B \sin \frac{t}{\sqrt{LC}}$, (1)
where A and B are two arbitrary constants.

Applying the given conditions, when $i=0$, $t=0$, (1) gives
 $0 = A.$

Again $i=I$, when $\frac{di}{dt}=0$.

$$\therefore \frac{di}{dt} = -\frac{A}{\sqrt{LC}} \sin \frac{t}{\sqrt{LC}} + \frac{B}{\sqrt{LC}} \cos \frac{t}{\sqrt{LC}}$$

For maximum or minimum of i , $\frac{di}{dt}=0$ gives

$$\begin{aligned} \tan \frac{t}{\sqrt{LC}} &= \frac{B}{A} \\ &= \infty \text{ when } A=0 \\ &= \tan \frac{\pi}{2}. \end{aligned}$$

$$\therefore t = \frac{\pi}{2} \sqrt{LC}.$$

Putting in (1), $i=I$ when $t = \frac{\pi}{2} \sqrt{LC}$, we get

$$I = A \cos \frac{\pi}{2} + B \sin \frac{\pi}{2},$$

$$B = I.$$

Hence putting the values of the constants A and B in (1), the required solution is $i = I \sin \frac{t}{\sqrt{LC}}.$

Problem 10. The relation between the small horizontal deflection θ of a bar magnet under the action of the earth's magnetic field is

$$A \frac{d^2 \theta}{dt^2} + MH\theta = 0,$$

when A is the moment of inertia of the magnet about the axis, M the magnetic moment of the magnet and H the horizontal component of the intensity of the field due to the Earth. Find the time of a complete vibration.

$$[\text{Ans. } T = 2\pi \sqrt{\left(\frac{A}{MH}\right)}]$$

Problem 11. In the case of the simple pendulum of length l , the equation connecting the acceleration due to gravity and the angle θ

through which the pendulum swings is $l \frac{d^2 \theta}{dt^2} + g\theta = 0$ where θ is small.

Determine the time of an oscillation.

$$[\text{Ans. } T = 2\pi \sqrt{\left(\frac{l}{g}\right)}]$$

Problem 12. Solve $\frac{d^2 i}{dt^2} + \frac{R}{L} \frac{di}{dt} + \frac{i}{LC} = 0$, where $R^2 C = 4L$.

$$[\text{Ans. } i = e^{-(Rt/2L)} \cdot (C_1 + C_2 t).]$$

Problem 13. The differential equation for a circuit containing resistance R , self-inductance L and capacitance C , in terms of current i and the time t is

$$\frac{d^2 i}{dt^2} + \frac{R}{L} \frac{di}{dt} + \frac{i}{LC} = \frac{1}{L} \frac{d}{dt} E(t)$$

$E(t)$ being the electromotive force which is the sum of voltage drops in a closed circuit according to Kirchhoff's second law. Find the current i and interpret the result physically.

$$\text{Given equation is } \frac{d^2 i}{dt^2} + \frac{R}{L} \frac{di}{dt} + \frac{i}{LC} = \frac{1}{L} \frac{d}{dt} E(t) \quad \dots (1)$$

Assume, $E(t) = F \sin(\omega t + \beta)$

$$\text{So that } \frac{d}{dt} E(t) = F \omega \cos(\omega t + \beta)$$

$$\text{Then (1) becomes, } \frac{d^2 i}{dt^2} + \frac{R}{L} \frac{di}{dt} + \frac{i}{LC} = \frac{F\omega}{L} \cos(\omega t + \beta) \quad \dots (2)$$

The solution may easily be found as

$$i = A e^{-\frac{R}{2L}t} \sin \left(\frac{\sqrt{4CL - R^2C^2}}{2CL} t + \epsilon \right) + F\omega C \left[\frac{R\omega C \sin(\omega t + \beta) + (1 - CL\omega^2) \cos(\omega t + \beta)}{(R\omega C)^2 + (1 - CL\omega^2)^2} \right]$$

$$\text{Putting } \sin \alpha = \frac{1 - CL\omega^2}{\sqrt{(R\omega C)^2 + (1 - CL\omega^2)^2}}$$

$$\text{and } \cos \alpha = \frac{R\omega C}{\sqrt{(R\omega C)^2 + (1 - CL\omega^2)^2}} \quad \dots (3)$$

The second term on the R.H.S. of (3)

$$= \frac{F\omega C}{\sqrt{(R\omega C)^2 + (1 - CL\omega^2)^2}} \sin [\omega t + \beta] \cos \alpha + \cos (\omega t + \beta) \sin \alpha \\ = \frac{F\omega C}{\sqrt{(R\omega C)^2 + (1 - CL\omega^2)^2}} \sin (\omega t + \beta + \alpha)$$

Hence the solution is

$$i = Ae^{-\frac{R}{2L}t} \sin \left(\frac{\sqrt{4CL - R^2C^2}}{2CL} t + \epsilon \right) + \frac{F\omega C}{\sqrt{(R\omega C)^2 + (1 - CL\omega^2)^2}} \sin (\omega t + \beta + \alpha) \quad \dots (4)$$

The current in the circuit consists of two parts namely a *damped harmonic motion* due to the first part and a *simple harmonic motion* due to the second part. The first part consists of the *damping factor* $e^{-\frac{R}{2L}t}$ and may be called as *transient current* while the second part as *steady state current*. The function $E(t)$ or $\frac{d}{dt} E(t)$ is called the *input* of the system and the solution as *output* of the system.

Amplitude of steady state current is

$$A_m = \frac{F\omega C}{\sqrt{(R\omega C)^2 + (1 - CL\omega^2)^2}} = \frac{F}{\sqrt{R^2 + \left(\frac{1}{\omega C} - L\omega\right)^2}}$$

whose denominator is called as *Impedance Z* of the circuit i.e.

$$Z = \sqrt{R^2 + \left(\frac{1}{\omega C} - L\omega\right)^2}$$

$$\text{So that } \frac{dz}{d\omega} = 0 \text{ gives } 2 \left(\frac{1}{\omega C} - L\omega \right) \left(-\frac{1}{C\omega^2} - L \right) = 0$$

$$\text{i.e. } \omega^2 = \frac{1}{CL} \text{ or } \omega = \frac{1}{\sqrt{CF}}$$

showing that the impedance z of the current is minimum for $\omega = \frac{1}{\sqrt{CL}}$ and then the amplitude A_m will be maximum and in such case we say that the electromotive force is in *Resonance* with the circuit.

$$\text{Putting } \omega = \frac{1}{\sqrt{CL}} \text{ in (5) we have } \text{Max } A_m = \frac{F}{R}$$

i.e. $\text{Max } A_m \propto \frac{1}{R}$ when Resonance occurs.

Consequently R should be sufficiently large to prevent a circuit breakdown.

Problem 14. A particle weighing 8 lbs distant 12 ft. apart from a fixed point O is given a velocity of 16 ft/sec in a direction perpendicular to x -axis. The particle is attracted to the fixed point by a force F whose magnitude is inversely proportional to the cube of its distance from O . Taking constant of proportionality as 10, find the distance of the particle from O as a function of time t .

Let (r, θ) be the polar coordinates of the position of the particle at time t . Then we have

$$F \propto \frac{1}{r^3} \text{ i.e. } F_r = -\frac{10}{r^3} = ma_r \quad \dots (1)$$

a_r being component of acceleration along r i.e. radial axis

$$\text{and } F_\theta = 0 = m a_\theta \quad \dots (2)$$

a_θ being component of acceleration in a direction perpendicular to radial axis

$$\text{Given } mg = 8 \text{ so that } m = \frac{8}{32} = \frac{1}{4}.$$

Since the particle subject to a central force moves in a plane, we have

$$a_r = \ddot{r} - r \dot{\theta}^2 \text{ and } a_\theta = 2\dot{r}\dot{\theta} + r\ddot{\theta}$$

$$\therefore (1) \text{ and } (2) \text{ give } \frac{1}{4}(\ddot{r} - r \dot{\theta}^2) = -\frac{10}{r^3} \text{ i.e. } \ddot{r} - r \dot{\theta}^2 = -\frac{40}{r^3} \quad \dots (3)$$

$$\text{and } \frac{1}{4}(2\dot{r}\dot{\theta} + r\ddot{\theta}) = 0 \quad \dots (4)$$

$$(4) \text{ multiplied by } 2r \text{ is equivalent to } \frac{d}{dt}(r^2\dot{\theta}) = 0.$$

Integrating $r^2 \dot{\theta} = A$, A being constant of integration.

Initially when $t=0$, $r=12$, $v_\theta=16$ (velocity component perpendicular to r i.e. x -axis since when $\theta=0$, r is along x -axis).

$$\text{Also } v_\theta = r \dot{\theta} = 16, \text{ so that } r^2 \dot{\theta} = A \text{ gives } A = r(r \dot{\theta}) = 12 \times 16 = 192$$

$$\therefore \dot{\theta} = \frac{A}{r^2} = \frac{192}{r^2} \quad \dots (5)$$

From (5) and (3), we have

$$\ddot{r} = \frac{(192)^2 - 40}{r^3} = \frac{36824}{r^3} \quad \dots (6)$$

$$\text{Put } u = \frac{dr}{dt} = r \text{ so that } \dot{u} = \frac{36824}{r^3} \quad \dots (7)$$

Multiplying both sides of (7) by $2u$, we get

$$2u \dot{u} = 2 \times \frac{36824}{r^3} u = 2 \times \frac{36824}{r^3} r;$$

$$\text{or } d(u^2) = 2 \times \frac{36824}{r^3} dr \text{ which gives on integration, } u^2 = -\frac{36824}{r^2} + B$$

$$\text{Initially when } t=0, r=12, u=r=0, \text{ so that } B = \frac{36824}{144}$$

$$\therefore u^2 = \dot{r}^2 = -36824 \left(\frac{1}{r^2} - \frac{1}{144} \right) = 36824 \left(\frac{r^2 - 144}{144r^2} \right)$$

$$\text{or } \left(\frac{dr}{dt} \right)^2 = \frac{144r^2}{36824(r^2 - 144)}$$

$$\text{i.e. } dt = \frac{12r}{191.9\sqrt{r^2 - 144}} dr \text{ which yields on integration}$$

$$t = \frac{1}{15.99} \sqrt{r^2 - 144} + C$$

$$\text{Initially when } t=0, r=12 \therefore C=0$$

$$\text{Hence } (15.99 t)^2 = r^2 - 144$$

$$\text{or } r^2 = 144 + 255.68 t^2 \text{ i.e. } r = \sqrt{144 + 255.68 t^2}$$

7.4. PARTIAL DIFFERENTIAL EQUATIONS

These are the equations containing one or more partial derivatives and are concerned with at least two independent variables. The order of a partial differential equation is the order of its highest derivative appearing in the equation:

[A] **The Partial Differential Equation of the first Order.** Let a relation $\phi(x, y, z, a, b) = 0$ $\dots (1)$

be derived from the partial differential equation

$$F(x, y, z, p, q) = 0 \quad \dots (2)$$

$$\text{where } p = \frac{\partial z}{\partial x} \text{ and } q = \frac{\partial z}{\partial y}.$$

The solution (1) consisting of as many arbitrary constants as the number of independent variables is called the *Complete Integral* of (2). If we give particular values to a and b in (1), then it becomes *Particular Integral*.

Since the envelope of all the surfaces given by (1) is touched at each of its points by some one of these surfaces, the coordinates of any point on the envelope with p and q belonging to the envelope at

that point must satisfy (2). Hence the relation found by eliminating a and b between $\phi(x, y, z, a, b) = 0, \frac{d\phi}{da} = 0, \frac{d\phi}{db} = 0$

is called the *Singular Integral*.

If $b=f(a)$ then (1) becomes $\phi[x, y, z, a, f(a)] = 0$.

The elimination of (a) between this equation and $\frac{d\phi}{da} = 0$ gives the *Singular Integral*.

Methods of Solution

(i) Lagrange's Method.

Lagrange's equation is of the form $Pp + Qq = R$ $\dots (3)$

P, Q, R being functions of x, y, z .

If $u=f(x, y, z)=a$ satisfies (3), then we get on differentiating (3) partially w.r.t. x and y ,

$$\frac{\partial u}{\partial x} + \frac{\partial u}{\partial z} p = 0 \text{ and } \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} q = 0 \text{ giving } p = -\frac{\partial u}{\partial x} / \frac{\partial u}{\partial z},$$

$$q = -\frac{\partial u}{\partial y} / \frac{\partial u}{\partial z} \text{ so that}$$

$$(3) \text{ yields, } P \frac{\partial u}{\partial x} + Q \frac{\partial u}{\partial y} + R \frac{\partial u}{\partial z} = 0 \quad \dots (4)$$

Obviously $u=a$ satisfy (4) and hence comparing (3) and (4) we get equations known as *Lagrange's Subsidiary equations*, i.e.

$$\frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R} \quad \dots (5)$$

which are also satisfied by $u=a$.

If $v=b$ be another integral of (5), then $\phi(u, v)=0$ or $u=\phi(v)$, ϕ , being arbitrary function, is an integral of (3).

Problem 15. Solve $(y^3x - 2x^4)p + (2y^4 - x^3y)q = 9z(x^3 - y^3)$

Lagrange's subsidiary equations are

$$\frac{dx}{y^3x - 2x^4} = \frac{dy}{2y^4 - x^3y} = \frac{dz}{9z(x^3 - y^3)} \quad \dots (1)$$

First two fractions give $\frac{dy}{dx} = \frac{2y^4 - x^3y}{y^3x - 2x^4}$

which being a homogeneous equation may be solved by putting $y=vx$ whence we get

$$\frac{dx}{x} = \frac{v^3 - 2}{v(v+1)(v^2 - v + 1)} dv = \left(-\frac{2}{v} + \frac{1}{v+1} + \frac{2v-1}{v^2 - v + 1} \right) dv$$

Integrating $\log x + \log A = -2 \log v + \log(v+1) + \log(v^2 - v + 1)$

$$\text{i.e. } Ax = \frac{(v+1)(v^2 - v + 1)}{v^2} = \left(\frac{y}{x} + 1 \right) \left(\frac{y^2}{x^2} - \frac{y}{x} + 1 \right) / \frac{y^2}{x^2}$$

$$\text{or } Ax^2y^2 = x^3 + y^3 \text{ i.e. } \frac{x^2y^2}{x^3 + y^3} = A$$

Again from (1) we have

$$\frac{\frac{dx}{x}}{y^3-2x^3} = \frac{\frac{dy}{y}}{2y^3-x^3} = \frac{\frac{dz}{9z(x^3-y^3)}}{-3(x^3-y^3)} = \frac{\frac{dx}{x} + \frac{dy}{y}}{-3(x^3-y^3)}$$

The last two fractions give $\frac{dx}{x} + \frac{dy}{y} = -\frac{dz}{-3z}$

Integrating $x^3y^3 = \frac{1}{Bz}$ or $z = \frac{1}{x^3y^3} \phi\left(\frac{x^3+y^3}{x^2y^2}\right)$ by taking

$$\frac{1}{B} = \phi\left(\frac{1}{A}\right)$$

or

$$z = \frac{1}{x^3y^3} \phi\left(\frac{x}{y^2} + \frac{y}{x^2}\right)$$

which is the required solution.

(ii) **Standard Methods:**

Standard I. Equation involving p and q only as $F(p, q) = 0$... (6)
have their complete integrals $z = ax + by + c$... (7)

where a and b are connected by the relation $F(a, b) = 0$.

Problem 16. Solve $3p^2 - 2q^2 = 4pq$.

Its solution is $z = ax + by + c$ provided $3a^2 - 2b^2 = 4ab$

$$\text{i.e., } b = \frac{-4a \pm \sqrt{16a^2 + 24a^2}}{4} = a\left(-1 \pm \frac{\sqrt{10}}{2}\right)$$

Hence the complete integral is

$$z = a\left[x + y\left(-1 \pm \frac{\sqrt{10}}{2}\right)\right] + c$$

Standard II. The equation analogous to Clairaut's form (i.e., $y =$

$x \frac{dy}{dx} + f\left(\frac{dy}{dx}\right)$ which has its solution as $y = xc + f(c)$), such that

$$z = px + qy + f(p, q)$$

has for its complete integral, $z = ax + by + f(a, b)$.

Problem 17. Solve $z = px + qy - 2\sqrt{pq}$

Its complete integral is $z = ax + by - 2\sqrt{ab}$

Standard III. The equations of the form $F(z, p, q) = 0$ are solved by putting $q = ap$, (a being an arbitrary constant) and changing p into $\frac{dz}{dX}$ where $X = x + ay$ and then solving the resulting ordinary differential equations between z and X .

Problem 18. Solve $9(p^2z + q^2) = 4$.

Putting $q = ap$ where $p = \frac{dz}{dX}$ and $X = x + ay$, we have

$$p = \frac{dz}{dX} = \pm \frac{1}{3\sqrt{z+a^2}} \text{ i.e., } dX = \frac{3}{2}(z+a^2)^{\frac{1}{2}} dz, \text{ taking +ve sign}$$

Integrating $X + b = (z + a^2)^{3/2}$ or $(x + ay + b)^2 = (z + a^2)^3$

Standard IV. Equations of the form $f_1(x, p) = f_2(y, q)$ are solved by putting $f_1(x, p) = f_2(y, q) = a$ (an arbitrary constant).

These equations give p and q which when substituted in $dz = p dx + q dy$ give the complete integral.

Problem 19. Solve $q = 2yp^2$.

We have $p^2 = \frac{q}{2y} = a^2$ (say).

When $p^2 = a^2$, $p = \frac{dz}{dx} = a$ we have $z = ax + \text{constant}$

and when $q = \frac{dz}{dy} = 2a^2y$ we have $z = a^2y^2 + \text{constant}$.

The complete integral is $z = ax + a^2y^2 + b$.

(iii) **Charpit's Method.** Let the partial differential equation be

$$F(x, y, z, p, q) = 0 \quad \dots (8)$$

Since z depends upon x and y both, therefore

$$dz = \frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy = p dx + q dy \quad \dots (9)$$

Forming Charpit's auxiliary equations (proof is not required)

$$\frac{dp}{\frac{\partial F}{\partial x} + p \frac{\partial F}{\partial z}} = \frac{dq}{\frac{\partial F}{\partial y} + q \frac{\partial F}{\partial z}} = \frac{dz}{-p \frac{\partial F}{\partial p} - q \frac{\partial F}{\partial q}} = \frac{dx}{-\frac{\partial F}{\partial p}} = \frac{dy}{-\frac{\partial F}{\partial q}} \quad \dots (10)$$

We may find a relation $f(p, q) = 0$

between p and q . (8) and (10) will yield p and q which when substituted in (9) give the required solution.

Problem 20. Solve $2zx - px^2 - 2qxy + pq = 0$

Charpit's auxiliary equations are

$$\frac{dp}{2z - 2qy} = \frac{dq}{0} = \frac{dz}{px^2 + 2xyq - 2pq} = \frac{dx}{x^2 - q} = \frac{dy}{2xy - p}$$

Whence $dq = 0$ gives $q = a$ (constant)

Putting $q = a$ in the given equation we get $p = \frac{2x(z - ay)}{x^2 - a}$.

Now substituting values of p and q in $dz = p dx + q dy$ we have

$$dz = \frac{2x(z - ay)}{x^2 - a} dx + a dy$$

$$\text{or } \frac{dx - a dy}{z - ay} = \frac{2x}{x^2 - a} dx$$

which gives on integration, $z - ay = c(x^2 - a)$ i.e., $z = ay + c(x^2 - a)$

(iv) **Partial Differential Equations of the Second and Higher Orders.**

Such an equation of second order is of the form

$$Rr + Ss + Tt = V$$

where $r = \frac{\partial^2 z}{\partial x^2}$, $s = \frac{\partial^2 z}{\partial x \partial y}$, $t = \frac{\partial^2 z}{\partial y^2}$ and R, S, T, V are functions of x, y, z, p, q , where $p = \frac{\partial z}{\partial x}$, $q = \frac{\partial z}{\partial y}$.

The complete solution of such equation will contain two arbitrary function as $z = f(x+ay) + \phi(x-ay)$.

Methods of Solution:

(i) **By Inspection.** Method is clear from the following Problem.

Problem 21. Solve $az = xy$ i.e., $a \frac{\partial^2 z}{\partial x^2} = xy$.

Integrating with regard to x , $a \frac{\partial z}{\partial x} = \frac{x^2}{2} y + \phi(y)$, constant of integration with regard to x being possibly a function of y .

Integrating again with regard to x ,

$$az = \int \frac{x^2}{2} y dy + \int \phi(y) dx + \text{const.}$$

$$= \frac{x^3 y}{6} + x\phi(y) + \psi(y).$$

(ii) **Monge's Method.**

The equation is $Rr + Ss + Tt = V$

Total differentials of p and q being

$$dp = \frac{\partial p}{\partial x} dx + \frac{\partial p}{\partial y} dy = r dx + s dy$$

and
$$dq = \frac{\partial q}{\partial x} dx + \frac{\partial q}{\partial y} dy = s dx + t dy$$

We have $r = \frac{dp - s dy}{dx}$, $t = \frac{dq - s dx}{dy}$.

Substituting these values of r and t in (1) we may get

$$(R dp dy + T dq dx - V dx dy) - s(R dy^2 - S dx dy + T dx^2) = 0 \quad \dots (3)$$

If any relation between x, y, z, p, q makes each of the bracketed expressions in (2) vanish, this relation will also satisfy (2). Hence from (2) we have so-called *Monge's subsidiary equations* as

$$R dy^2 - S dx dy + T dx^2 = 0 \quad \dots (3)$$

$$R dp dy + T dq dx - V dx dy = 0 \quad \dots (4)$$

Let (3) resolve into two linear equations

$$dy - m_1 dx = 0 \quad \dots (5)$$

$$dy - m_2 dx = 0 \quad \dots (6)$$

Combining (5) with (4) and with $dz = p dx + q dy$ if necessary, we may get two integrals $u_1 = a$, $v_1 = b$ giving an *intermediary integral* $u_1 = f_1(v_1)$, f , being an arbitrary function.

Similarly from (6) and (4) alongwith $dz = p dx + q dy$, we may find another intermediary integral $u_2 = f_2(v_2)$, f_2 being arbitrary.

These two intermediary integrals can yield p and q which when substituted in $dz = p dx + q dy$ will yield the complete integral on integration.

In case $m_1 = m_2$ either of the intermediary integral may be integrated to give the complete integral.

Problem 22. Solve $r + (a+b)s + abt = xy$.

Putting $r = \frac{dp - s dy}{dx}$ and $t = \frac{dq - s dx}{dy}$ in the given equation we get

$$\text{Monge's subsidiary equations as } dy^2 - (a+b) dx dy + ab dx^2 = 0 \dots (1)$$

$$dp dy + abd q dx - xy dx dy = 0 \dots (2)$$

(1) yields, $dy - a dx = 0$ and $dy - b dx = 0$

Integrating them, $y - ax = c_1$ and $y - bx = c_2$

Comparing these with (2) we get $adp + abd q - ax(c_1 + ax) dx = 0$

$$bdp + abd q - bx(c_2 + ax) dx = 0$$

Their integration yields $p + bq - c_1 \frac{x^2}{2} - \frac{ax^3}{3} = k_1$

$$p + aq - c_2 \frac{x^2}{2} - \frac{bx^3}{3} = k_2$$

$$p + bq - (y - ax) \frac{x^2}{2} - \frac{ax^3}{3} = \phi_1(c_1) = \phi_1(y - ax) \dots (3)$$

$$p + aq - (y - bx) \frac{x^2}{2} - \frac{bx^3}{3} = \phi_2(c_2) = \phi_2(y - bx) \dots (4)$$

Solving (3) and (4),

$$p = \frac{1}{a-b} \left[\frac{yx^2}{2} (a-b) - (a^2 - b^2) \frac{x^3}{6} + a\phi_1(y-ax) - b\phi_2(y-bx) \right]$$

and $q = \frac{1}{a-b} \left[\frac{x^3}{6} (a-b) - \phi_1(y-ax) + \phi_2(y-bx) \right]$

Substituting these values of p and q in $dz = p dx + q dy$, we find

$$dz = \left[\frac{yx^2}{2} - (a+b) \frac{x^3}{6} + \frac{a\phi_1(y-ax)}{a-b} - \frac{b\phi_2(y-bx)}{a-b} \right] dx$$

$$+ \left[\frac{x^3}{6} - \frac{\phi_1(y-ax)}{a-b} + \frac{\phi_2(y-bx)}{a-b} \right] dy$$

$$= -(a+b) \frac{x^3}{6} dx + \frac{3x^2 y dx + x^3 dy}{6} - \frac{1}{a-b} \left[\phi_1(y-ax) \right.$$

$$\left. (dy - a dx) \right] + \frac{1}{a-b} \left[\phi_2(y-bx) (dy - b dx) \right].$$

Integrating, $z = -(a+b) \frac{x^4}{24} + \frac{yx^3}{6} + \psi_1(y-ax) + \psi_2(y-bx)$

(iii) **Monge's Method of Integrating** $Rr + Ss + Tt + U(rt - s^2) = V$, R, S, T, U, V being functions of x, y, z, p, q .

Putting $r = \frac{dp - s dy}{dx}$ and $t = \frac{dq - s dx}{dy}$ in the given equation we get

$$(Rdp \, dy + Tdq \, dx + Udp \, dq - Vdx \, dy) - s(Rdy^2 - Sdx \, dy + Tdx^2 + Udp \, dx + Udq \, dy) = 0$$

$$\text{Say } N - sM = 0$$

$$\text{Consider, } M + \lambda N = Rdy^2 + Tdx^2 - (S + \lambda V)dxdy + Udpdx + Udqdy + \lambda Rdpdy + \lambda Tdqdx + \lambda Udpdq$$

$$\equiv (A dy + B dx + C dp) (E dy + F dx + G dq) \text{ (say)}$$

Then equating the coefficients of dy^2 , dx^2 , $dxdy$ we get

$$AE = R, BF = T, GC = \lambda U$$

$$\text{Also taking } A = R, E = 1, B = kT, F = \frac{1}{k}, C = mU, G = \frac{\lambda}{m} \text{ and}$$

equating the coefficients of the other five terms, we may find

$$kT + \frac{R}{k} = -(S + \lambda V) \quad \dots (14)$$

$$\frac{\lambda R}{m} = U \quad \dots (15)$$

$$\frac{kT\lambda}{m} = \lambda T, \quad \dots (16)$$

$$mU = \lambda R \quad \dots (17)$$

$$\frac{mU}{k} = U \quad \dots (18)$$

From (18), $m = k$ which satisfies (16).

From (15) or (17) $m = \frac{\lambda R}{U} = k$ and hence from (14),

$$\lambda^2 (RT + UV) + \lambda US + U^2 = 0 \quad \dots (19)$$

So if λ is a root of (19), the required factors of $M + \lambda N$ are

$$\left(R dy + \lambda \frac{RT}{U} du + \lambda R dp \right) \left(dy + \frac{U}{\lambda R} dx + \frac{U}{R} dq \right)$$

$$\text{i.e. } \frac{R}{U} (Udy + \lambda Tdx + \lambda Udp) \frac{1}{\lambda R} (\lambda Rdy + Udx + \lambda Udq)$$

we thus obtain integrals from the linear equations

$$Udy + \lambda Tdx + \lambda Udp = 0 \quad \dots (20)$$

$$\lambda Rdy + Udx + \lambda Udq = 0 \quad \dots (21)$$

If $u_1 = f_1(v_1)$ and $u_2 = f_2(v_2)$ be the two intermediary integrals so obtained for finding p and q and substituting in $dz = pdx + qdy$ we get the required solution after integration.

$$\text{Problem 23. Solve } z(I + q^2)r - 2pqzs + z(I + p^2)t - z^2(s^2 - rt) + I + p^2 + q^2 = 0.$$

$$\text{Here } R = z(1 + q^2), S = -2pqz, T = (1 + p^2)z, U = z^2V, V = -(1 + p^2 + q^2)$$

$$\therefore \text{quadratic in } \lambda \text{ is } (RT + UV)\lambda^2 + \lambda US + U^2 = 0 \text{ i.e. } (pq\lambda - z)^2 = 0$$

$$\text{giving } \lambda = \frac{z}{pq}.$$

Now the system of intermediary integrals is

$$Udy + \lambda Tdx + \lambda Udp = 0, Udx + \lambda Rdy + \lambda Udq = 0$$

$$\text{i.e. } pqdy + (1 + p^2)dp + zdp = 0, \quad \dots (1)$$

$$pq \, dn + (1 + q^2)dz + zdq = 0 \quad \dots (2)$$

$$\text{Also } dz = pdx + qdy \quad \dots (3)$$

$$\text{So (1) can be written as } dx + p(pdx + qdy) + zdp = 0$$

$$\text{or } dx + pdz + zdp = 0 \text{ giving } x + pz = A$$

$$\text{Similarly from (2), } y + qz = B$$

$$\text{So that } p = \frac{A-x}{z} \text{ and } q = \frac{B-y}{z}$$

Putting in $dz = pdx + qdy$ and integrating we get the required solution,

$$dz = \frac{A-x}{z} dx + \frac{B-y}{z} dy \text{ or } -zdz = (A-x)(-dx) + (B-y)(-dy)$$

$$-\frac{z^2}{2} = \frac{(A-x)^2}{2} + \frac{(B-y)^2}{2} + \text{const. i.e. } z^2 + (x-A)^2 + (y-B)^2 = C^2$$

10. General Linear Partial Differential Equations of an order higher than the first.

Such equations are of the form

$$A_0 \frac{\partial^n z}{\partial x^n} + A_1 \frac{\partial^n z}{\partial x^{n-1} \partial y} + \dots + A_n \frac{\partial^n z}{\partial y^n} + B_0 \frac{\partial^{n-1} z}{\partial x^{n-1}} + \dots + M \frac{\partial z}{\partial x} + N \frac{\partial z}{\partial y} + Pz = f(x, y)$$

$$\text{or } [A_0 D^n + A_1 D^{n-1} D' + \dots + A_n D'^n + B_0 D^{n-1} + \dots + MD + ND' + P]z = f(x, y)$$

$$\text{where } D \equiv \frac{\partial}{\partial x} \text{ and } D' \equiv \frac{\partial}{\partial y}$$

$$\text{i.e. } F(D, D')z = f(x, y).$$

Its complete solution consists of two parts: (i) Complementary function (C.F.) (ii) Particular Integral (P.I.).

The complementary function is obtained from $F(D, D') = 0$.

Methods of Solution.

(i) Complementary function of a homogeneous partial differential equation with constant coefficients.

Such an equation is of the form

$$(A_0 D^n + A_1 D^{n-1} D' + A_2 D^{n-2} D'^2 + \dots + A_n D'^n)z = f(x, y)$$

$$\text{Taking the trial solution as } z = \phi(y + mx), \text{ so that } D^n(z)$$

$$= m^n \phi^{(n)}(y + mx), D^n(z) = \phi^{(n)}(y + mx)$$

$$\text{and in general } D^r D'^s = m^r \phi^{(r+s)}(y + mx)$$

the auxiliary equation of the given equation becomes

$$A_0 m^n + A_1 m^{n-1} + \dots + A_n = 0$$

If the n roots given by it be $m_1, m_2, \dots, m_n$, then the required complementary function is $z = \phi_1(y + m_1x) + \phi_2(y + m_2x) + \dots + \phi_n(y + mx)$

Problem 24. Solve $\frac{\partial^2 z}{\partial x^2} - a^2 \frac{\partial^2 z}{\partial y^2} = 0$ i.e. $(D^2 - a^2 D'^2)z = 0$.

Putting $z = \phi(z + mx)$, the auxiliary equation is $m^2 - a^2 = 0$ giving $m = \pm a$

Hence the solution is $z = \phi(y + ax) + \psi(y - ax)$

Note. If an equation has repeated roots such as $(D - mD')^2 z = 0$, the solution is $z = x\phi(y + mx) + \psi(y + mx)$

If a root is repeated thrice then $z = x^2\phi(y + mx) + x\psi(y + mx) + \chi(y + mx)$ and so on

(ii) **Particular integral of homogeneous equations.**

Let the equation be $F(D, D')z = \phi(x, y)$. Then

$$P.I. = \frac{1}{F(D, D')} \phi(x, y).$$

Case I. If $\phi(x, y) = e^{ax+by}$ then $\frac{1}{F(D, D')} e^{ax+by} = \frac{1}{F(a, b)} e^{ax+by}$ provided $F(a, b) \neq 0$

Case II. If $\phi(x, y) = \sin ax$ or $\cos ax$ then express $F(D, D')$ in $F(D^2, D D', D'^2)$ and write

$$\frac{\sin ax \text{ or } \cos ax}{F(D^2, D D', D'^2)} = \frac{\sin ax \text{ or } \cos ax}{F(-a^2, -ab, -b^2)} \text{ provided } F(-a^2, -ab, -b^2) \neq 0$$

Case III. If $\phi(x, y) = x^m y^n$, then $\frac{1}{F(D, D')} x^m y^n = [F(D, D')]^{-1} x^m y^n$, expanding binomially and operating $x^m y^n$ on every term.

Case IV. If $\phi(x, y) = e^{ax+by} V$ which may also arise in case of failures of cases I and II.

$\frac{1}{F(D, D')} e^{ax+by} V = e^{ax+by} \frac{1}{F(D+a, D'+b)} V$ which reduces to any of the above cases.

Problem 25. Find the Particular Integral of $(D^2 - 2DD' + D'^2)z = 12xy$

$$\begin{aligned} \text{We have, } P.I. &= \frac{1}{(D^2 - 2DD' + D'^2)} 12xy = 12 \frac{1}{(D - D')^2} xy \\ &= 12 \left(1 - \frac{D'}{D} \right)^{-1} xy \\ &= 12 \left(1 + \frac{2D'}{D} + \dots \right) xy = 12 \cdot \frac{1}{D^2} \left(xy + \frac{2}{D} x \right) \\ &= 12 \left[\frac{1}{D^2} xy + \frac{2}{D^3} x \right] = 12 \left[\frac{x^3 y}{6} + \frac{x^4}{12} \right] = 2x^3 y + x^4 \end{aligned}$$

Complete Integral is $z = x\phi(y+x) + \psi(y+x) + 2x^3 y + x^4$.

COROLLARY 1. In case $\phi(x, y) = f(ax+by)$ and $F(D, D')$ is a homogeneous function of D, D' of degree n . Then

$$D^n f(ax+by) = a^n f^n(ax+by)$$

$$D'^n f(ax+by) = b^n f^n(ax+by)$$

So that $F(D, D')f(ax+by) = F(a, b)f^n(ax+by)$

$$\therefore \frac{f^n(ax+by)}{F(D, D')} = \frac{f(ax+by)}{F(a, b)} \text{ when } F(a, b) \neq 0$$

Hence $\frac{f(ax+by)}{F(D, D')}$ may be evaluated by integrating $f(ax+by)$, n times with regard to $(ax+by)$ and then dividing by $F(a, b)$

Problem 26. Solve $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 12(x+y)$

i.e. $(D^2 + D'^2)V = 12(x+y)$.

Auxiliary equation is $m^2 + 1 = 0$ i.e. $m = \pm i$ so that C.F. is $\phi(y+ix) + \psi(y-ix)$

$$\text{and } P.I. = \frac{12(x+y)}{D^2 + D'^2} = \frac{12 \cdot \frac{(x+y)^3}{2 \cdot 3}}{1^2 + 1^2} = (x+y)^3$$

Hence the solution is $V = (x+y)^3 + \phi(y+ix) + \psi(y-ix)$.

COROLLARY 2. In case, method of Cor. 1 fails i.e. $F(a, b) = 0$, then consider

$$(D - mD')z = p - mq = x^r \psi(x+mx) \dots (22)$$

Lagrange's subsidiary equations are $\frac{dx}{1} = \frac{dy}{-m} = \frac{dz}{x^r \psi(y+mx)}$

of which first two fractions, give $dy + mdx = 0$ i.e. $y + mx = \text{const.}$ = c (say). From first and third fractions, we have

$$dx = \frac{dz}{x^r \psi(y+mx)} = \frac{dz}{x^2 \psi(c)} \text{ or } dz = x^r \psi(c) dx$$

$$\text{Integrating } z = \frac{x^{r+1}}{r+1} \psi(c) = \frac{x^{r+1}}{r+1} \psi(y+mx)$$

Thus (22) yields $(D - mD') \frac{x^{r+1}}{r+1} \psi(y+mx) = x^r \psi(y+mx)$

$$\text{i.e. } \frac{1}{D - mD'} x^r \psi(y+mx) = \frac{x^{r+1}}{r+1} \psi(y+mx)$$

$$\begin{aligned} \text{Hence } \frac{1}{(D - mD')^n} \psi(y+mx) &= \frac{1}{(D - mD')^{n-1}} \cdot \frac{x}{1} \psi(y+mx) \\ &= \frac{1}{(D - mD')^{n-2}} \cdot \frac{x^2}{1 \cdot 2} \psi(y+mx) \\ &\dots \dots \dots \\ &= \frac{x^n}{n!} \psi(y+mx) \end{aligned}$$

Problem 27. Solve $(D^2 - 6D + 9D')z = 6x + 2y$. i.e. $(D - 3D')^2 z = 6x + 2y$. C.F. is clearly, $x\phi(y+3x) + \psi(y+3x)$

$$\begin{aligned} P.I. &= \frac{1}{(D-3D')^2} \cdot (6x+2y) = \frac{2}{(D-3D')^2} \cdot (y+3x) \\ &= 2 \cdot \frac{x^2}{2} (y+3x) = x^2(y+3x). \end{aligned}$$

Solution is $z = x^2(3x+y) + x\phi(y+3x) + \psi(y+3x)$.

COROLLARY 3. General rule.

Consider $(D - mD')z = p - mq = f(x, y)$

Lagrange's subsidiary equations are $\frac{dx}{1} = \frac{dy}{-m} = \frac{dz}{f(x, y)}$

First two fractions give $y + mx = c$

and then first and third fractions yield $dz = f(x, y) dx = f(x, c - mx) dx$
 $z = \int f(x, c - mx) dx + \text{const.}$

$\therefore$ Hence, $\frac{f(x, y)}{D - mD'} = \int f(x, c - mx) dx$, where c is replaced by $y + mx$ after integration.

Problem 28. Solve $(2D^2 - DD' - 3D'^2)z = 5e^{x-y}$ i.e., $(2D - 3D')(D + D')z = 5e^{x-y}$. Clearly C.F. is $\phi(y-x) + \psi(2y+3x)$

$$\begin{aligned} P.I. &= \frac{1}{(D+D')(2D-3D')} 5e^{x-y} = \frac{1}{D+D'} \cdot \frac{5e^{x-y}}{2-3(-1)} \text{ by Cor. 1} \\ &= \frac{1}{D+D'} e^{x-y} = \int e^{x-(c+x)} dx = \int e^{-c} dx = xe^{-c} = xe^{x-y}. \end{aligned}$$

Solution is $z = xe^{x-y} + \phi(y-x) + \psi(2y+3x)$.

(iii) **Non-homogeneous linear equations** (complementary function)

Consider $(D - mD' - n)z = 0$ i.e., $p - mq = nz$

Lagrange's subsidiary equations are $\frac{dx}{1} = \frac{dy}{-m} = \frac{dz}{nz}$

First two fractions give $y + mx = \text{const.}$

First and third fractions give $n \frac{dx}{z} = \frac{dz}{z}$ i.e., $nx = \log z - \log k$ or $z = k e^{nx}$.

Hence the integral of given equation is $z = e^{nx} \phi(y + mx)$.

Note 1. If factors are repeated say $(D - mD' - n)^2 z = 0$, then $z = xe^{nx} \phi(y + mx) + e^{nx} \psi(y + mx)$.

Note 2. The Particular Integral is obtained by the methods already discussed.

Problem 29. Solve

$$\frac{\partial^2 z}{\partial x^2} - \frac{\partial z}{\partial x} \frac{\partial z}{\partial y} - 2 \frac{\partial^2 z}{\partial y^2} + 2 \frac{\partial z}{\partial y} + 2 \frac{\partial z}{\partial x} = e^{2x+3y} + \sin(2x+y) + xy.$$

Given equation is $(D^2 - DD' + 2D'^2 + 2D' + 2D)z = e^{2x+3y} + \sin(2x+y) + xy$
 $= e^{2x+3y} + \sin(2x+y) + xy$

C.F. is $\phi(y-x) + e^{-2x}\psi(y+2x)$ and by usual methods

$$P.I. = -\frac{1}{10} e^{2x+3y} - \frac{1}{6} \cos(2x+y) + \frac{x}{24} (6xy + 9x - 2x^2 - 6y - 12)$$

Hence the solution.

(iv) **Equations Reducible to Linear Form.**

Consider an equation all of whose terms are of the form

$$Ax^m y^n \frac{\partial^{m+n} z}{\partial x^m \partial y^n}$$

Put $x = e^u$ i.e. $u = \log x$ and $y = e^v$ i.e. $v = \log y$, then it is easy to verify that if $D \equiv \frac{\partial}{\partial u} \equiv x \frac{\partial}{\partial x}$; $D' \equiv \frac{\partial}{\partial v} \equiv y \frac{\partial}{\partial y}$,

$$\text{then } x^2 \frac{\partial^2}{\partial x^2} = D(D-1), \quad x^3 \frac{\partial^3}{\partial x^3} = D(D-1)(D-2) \text{ etc.}$$

$$y^2 \frac{\partial^2}{\partial y^2} = D'(D'-1), \quad y^3 \frac{\partial^3}{\partial y^3} = D'(D'-1)(D'-2) \text{ etc.}$$

$$\text{in general } x^m y^n \frac{\partial^{m+n}}{\partial x^m \partial y^n} = x^m \frac{\partial^m}{\partial x^m} \cdot y^n \frac{\partial^n}{\partial y^n} = D(D-1)\dots(D-m+1) \cdot D'(D'-1)\dots(D'-n+1).$$

so that the given equation will reduce to $F(D, D')z = V$ which can be integrated by usual methods.

$$\text{Problem 30. Solve } x^2 \frac{\partial^2 z}{\partial x^2} - y^2 \frac{\partial^2 z}{\partial y^2} - y \frac{\partial z}{\partial y} + x \frac{\partial z}{\partial x} = 0.$$

Putting $u = \log x$, $v = \log y$ with $D \equiv \frac{\partial}{\partial u}$, $D' \equiv \frac{\partial}{\partial v}$, this becomes

$$(D(D-1) - D'(D'-1) - D' + D)z = 0 \text{ i.e. } (D^2 - D'^2)z = 0 \text{ or } (D - D')(D + D')z = 0$$

$$\text{so that solution is } z = f(v+u) + F(v-u) = f(\log xy) + F\left(\log \frac{y}{x}\right) = \phi(xy) + \psi\left(\frac{y}{x}\right).$$

(v) **Equations in which Linear Factors of $F(D, D') = 0$ are not possible.**

Consider $(D^2 - D')z = 0$.

Assume $z = Ae^{\alpha x + \beta y}$, giving $D^2 z = A\beta^2 e^{\alpha x + \beta y}$ and $D^2 z = A\alpha^2 e^{\alpha x + \beta y}$ so that the given equation, yields $\alpha^2 - \beta = 0$ i.e. $\alpha^2 = \beta$

Hence the Complementary function is $z = Ae^{\alpha x + \alpha^2 y} = Ae^{\alpha(x + \alpha y)}$ in general, $z = \Sigma Ae^{\alpha(x + \alpha y)}$

7.5. SPECIAL TYPES OF DIFFERENTIAL EQUATIONS ARISING IN PHYSICS

[A] Hyperbolic, Parabolic and Elliptic Equations

Consider $f_1(x, y, z) \frac{\partial^2 z}{\partial x^2} + f_2(x, y, z) \frac{\partial^2 z}{\partial x \partial y} + f_3(x, y, z) \frac{\partial^2 z}{\partial y^2} + F(x, y, z, p, q) = 0$... (1)

Writing it as $Rr + Ss + Tt + V = 0$,

where $r = \frac{\partial^2 z}{\partial x^2}$, $s = \frac{\partial^2 z}{\partial x \partial y}$, and $t = \frac{\partial^2 z}{\partial y^2}$ etc.

(i) if $S^2 > 4RT$, the equation (1) is said to be *Hyperbolic*

(ii) if $S^2 = 4RT$, " " *Parabolic*

(iii) if $S^2 < 4RT$, " " *Elliptic*.

e.g. $\frac{\partial^2 z}{\partial x^2} = \frac{\partial^2 z}{\partial y^2}$ i.e. $r = t$ is an hyperbolic equation

$\frac{\partial^2 z}{\partial x^2} = \frac{\partial z}{\partial y}$ i.e. $r = q$ is a parabolic equation

and $\frac{\partial^2 z}{\partial x^2} = -\frac{\partial^2 z}{\partial y^2}$ i.e. $r = -t$ is an elliptic equation.

[B] Diffusion Equation or Fourier Equation of Heat Flow

A parabolic equation of the type $\nabla^2 V = \frac{1}{k} \frac{\partial V}{\partial t}$ is said to be a *diffusion equation*.

Here $\nabla^2 \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ (in 3-D); k is the diffusivity and $V(x, y, z, t)$ is the temperature at any point (x, y, z) of a solid at time t .

One-dimensional diffusion equation is $\frac{\partial^2 V}{\partial x^2} = \frac{1}{k} \frac{\partial V}{\partial t}$

Two-dimensional " $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = \frac{1}{k} \frac{\partial V}{\partial t}$

Three-dimensional " $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial z^2} = \frac{1}{k} \frac{\partial V}{\partial t}$

COROLLARY. In steady-state heat flow, (1) yields $\nabla^2 V = 0$

Here $\nabla^2 V \equiv \frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial z^2} = 0$ is known as *Laplace's equation*

Its cylindrical form is $\nabla^2 V \equiv \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial V}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 V}{\partial \theta^2} + \frac{\partial^2 V}{\partial z^2} = 0$

and spherical polar form is $\nabla^2 V \equiv \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial V}{\partial r} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 V}{\partial \phi^2} + \left(\sin \theta \frac{\partial}{\partial \theta} \right) \left(\frac{1}{\sin \theta} \frac{\partial V}{\partial \theta} \right) = 0$

(C) Wave Equation

A hyperbolic equation of the type $\square^2 V = 0$... (4)

where $\square^2 \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}$, c being a constant is said to be a *wave equation*.

One-dimensional wave equation is $\frac{\partial^2 V}{\partial x^2} = \frac{1}{c^2} \frac{\partial^2 V}{\partial t^2}$

Two-dimensional " $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = \frac{1}{c^2} \frac{\partial^2 V}{\partial t^2}$

Three-dimensional " $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial z^2} = \frac{1}{c^2} \frac{\partial^2 V}{\partial t^2}$

Methods of Solution

(i) Separation of Variables.

The equations introduced in this section may conveniently be solved by the method of separation of variables. This method is illustrated here.

Consider $\frac{\partial^2 z}{\partial x^2} = \frac{1}{k} \frac{\partial z}{\partial t}$, where $z = z(x, t)$... (5)

Suppose that $z = X(x) T(t)$... (6)

$X(x)$ being function of x only and $T(t)$ function of t only.

We have from (6), $\frac{\partial^2 z}{\partial x^2} = \frac{\partial^2 X}{\partial x^2} T$ and $\frac{\partial z}{\partial t} = X \frac{dT}{dt}$.

Their substitutions in (5) yield $\frac{1}{X} \frac{d^2 X}{dx^2} = \frac{1}{kT} \frac{dT}{dt}$... (7)

Assuming that L.H.S. of (7) is independent of t and R.H.S. independent of x , each side of (7) can be equated to some constant known as *Constant of separation*.

According as this constant is negative or positive, there arise two cases.

Case I. Let the constant of separation be $-n^2$. Then (7) gives

$\frac{1}{X} \frac{d^2 X}{dx^2} = -n^2$ i.e. $\frac{d^2 X}{dx^2} + n^2 X = 0$... (8)

and $\frac{1}{kT} \frac{dT}{dt} = -n^2$ i.e. $\frac{dT}{dt} + n^2 kT = 0$... (9)

(8) gives on integration $X = a \cos(nx + \alpha)$, a, α being arbitrary constants and (9) gives on integration $T = be^{-kn^2 t}$, b being an arbitrary constant.

The two solutions when combined in accordance with (6) i.e. $z = XT$, yield

$z = A \cos(nx + \alpha) e^{-kn^2 t}$... (10)

where $A (=ab)$, n and α are constants and they can be determined by the boundary conditions.

Case II. Let the constant of separation be n^2 . Then (7) gives

$$\frac{d^2 X}{dx^2} - n^2 X = 0 \text{ and } \frac{dT}{dt} - n^2 k T = 0.$$

Which give on integration, $X = a \cosh (nx + \alpha)$ and $T = be^{kn^2 t}$

So that the combined solution of (5) is $z = A \cosh (nx + \alpha) e^{kn^2 t}$... (11)

The solutions (10) and (11) can be expressed as

$$z = A_n \cos (nx + \alpha_n) e^{-kn^2 t} \quad \dots (12)$$

$$z = A_n \cosh (nx + \alpha_n) e^{kn^2 t} \quad \dots (13)$$

where the constant A_n corresponds to a particular value of n .

Since these are the solutions of (5) for all n , therefore summing over all values of n , the solution of (5) is

$$z(x, t) = \sum_{n=0}^{\infty} A_n \cos (nx + \alpha_n) e^{-kn^2 t} \quad \dots (14)$$

$$\text{or } z(x, t) = \sum_{n=0}^{\infty} A_n \cosh (nx + \alpha_n) e^{kn^2 t} \quad \dots (15)$$

according as the constant of separation is negative or positive.

Problem 31. Solve $\frac{\partial^2 z}{\partial x^2} = \frac{I}{k} \frac{\partial z}{\partial t}$ with the boundary conditions, $z=0$ when $x=0$ and $x=l$ for all values of t .

Using (10), the solution is $z(x, t) = A \cos (nx + \alpha) e^{-kn^2 t}$

Initially $z=0$ when $x=0$ gives $z(0, t) = A \cos \alpha e^{-kn^2 t} = 0$

and $x=l$ for all t gives $z(l, t) = A \cos (nl + \alpha) e^{-kn^2 t} = 0$

So that $A \cos \alpha = 0$ and $A \cos nl \cos \alpha - A \sin nl \sin \alpha = 0$.

Which yield $\alpha = \frac{\pi}{2}$ and $n = \frac{\pi}{l}, \frac{2\pi}{l}, \frac{3\pi}{l}, \dots$

With $n = \frac{\pi}{l}$, the solution is $z(x, t) = A \sin nxe^{-kn^2 t}$

Note. Here $n = \frac{\pi}{l}, \frac{2\pi}{l}, \frac{3\pi}{l}, \dots$ are called **eigen values**.

(ii) **D'Alemberts' Method.**

Consider the equation of vibrating cords

$$\frac{\partial^2 u}{\partial t^2} = h^2 \frac{\partial^2 u}{\partial x^2} \quad \dots (16)$$

which is also known as *D'Alembert's equation*.

Suppose that $v = x + ht$
 $w = x - ht$

So that $\frac{\partial v}{\partial x} = \frac{\partial w}{\partial x} = 1, \frac{\partial v}{\partial t} = h, \frac{\partial w}{\partial t} = -h$

$$\text{and } \frac{\partial u}{\partial x} = \frac{\partial u}{\partial v} \frac{\partial v}{\partial x} + \frac{\partial u}{\partial w} \frac{\partial w}{\partial x} = \frac{\partial u}{\partial v} + \frac{\partial u}{\partial w} \text{ i.e. } \frac{\partial}{\partial x} \equiv \frac{\partial}{\partial v} + \frac{\partial}{\partial w}$$

$$\therefore \frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial x} \right) = \left(\frac{\partial}{\partial v} + \frac{\partial}{\partial w} \right) \left(\frac{\partial u}{\partial v} + \frac{\partial u}{\partial w} \right) = \frac{\partial^2 u}{\partial v^2} + 2 \frac{\partial^2 u}{\partial v \partial w} + \frac{\partial^2 u}{\partial w^2} \quad \dots (18)$$

$$\text{Also } \frac{\partial u}{\partial t} = \frac{\partial u}{\partial v} \frac{\partial v}{\partial t} + \frac{\partial u}{\partial w} \frac{\partial w}{\partial t} = h \left(\frac{\partial u}{\partial v} - \frac{\partial u}{\partial w} \right) \text{ i.e. } \frac{\partial}{\partial t} \equiv h \left(\frac{\partial}{\partial v} - \frac{\partial}{\partial w} \right)$$

$$\therefore \frac{\partial^2 u}{\partial t^2} = h^2 \left(\frac{\partial}{\partial v} - \frac{\partial}{\partial w} \right) \left(\frac{\partial u}{\partial v} - \frac{\partial u}{\partial w} \right) = h^2 \left(\frac{\partial^2 u}{\partial v^2} - 2 \frac{\partial^2 u}{\partial v \partial w} + \frac{\partial^2 u}{\partial w^2} \right) \quad \dots (19)$$

(16) yields with the help of (18) and (19), $\frac{\partial^2 u}{\partial v \partial w} = 0$.

Integrating with regard to w , we get $\frac{\partial u}{\partial v} = \text{const.} = f(v)$ (say)

Integrating it again with regard to v , $u = \int f(v) dv + \text{const.}$
 $= \phi(v) + \psi(w)$ (say)

$$\text{i.e. } u(x, t) = \phi(x + ht) + \psi(x - ht) \quad \dots (20)$$

Physically interpreted (20) represents two plane waves travelling in opposite directions with same period.

Aliter. Writing $D_x \equiv \frac{\partial}{\partial x}$, $D_x^2 \equiv \frac{\partial^2}{\partial x^2}$, (16) becomes

$$\frac{\partial^2 u}{\partial t^2} = (h D_x)^2 u \quad \dots (21)$$

Treating (20) as an ordinary differential equation with constant coefficients such as

$$\frac{d^2 u}{dt^2} = a^2 u \quad \dots (22)$$

where

$$a = h D_x \quad \dots (23)$$

We have the solution of (22) as $u = A_1 e^{at} + A_2 e^{-at}$... (24)

where A_1, A_2 are arbitrary constants.

Now (22) is formally satisfied by

$$u = e^{h D_x t} \phi(x) + e^{-h D_x t} \psi(x) \quad \dots (25)$$

Since (22) has been integrated with regard to t and so the arbitrary constants A_1, A_2 appearing in (24) can be replaced by arbitrary functions $\phi(x)$ and $\psi(x)$ of x . Now we know that Taylor's expansion is

$$f(x+h) = f(x) + \frac{h}{1!} f'(x) + \frac{h^2}{2!} f''(x) + \dots + \frac{h^n}{n!} f^n(x) + \dots$$

Using the symbolic operator $D_x^n = \frac{d^n}{dx^n}$ for $n=1, 2, 3, \dots$

$$\text{so that } D_x^n f(x) = \frac{d^n}{dx^n} f(x)$$

$$\begin{aligned} \text{This becomes } f(x+h) &= f(x) + \frac{h}{1} D_x f(x) + \frac{h^2}{2} D_x^2 f(x) + \dots \\ &\quad + \frac{h^n}{n} D_x^n f(x) + \dots \\ &= \left(1 + \frac{hD_x}{1} + \frac{h^2 D_x^2}{2} + \dots + \frac{h^n D_x^n}{n} + \dots \right) f(x) \\ &= e^{hD_x} f(x) \end{aligned} \quad \dots (26)$$

Using (26), and replacing h by ht and by $-ht$ in succession, (25) yields

$$u(x, t) = \phi(x+ht) + \psi(x-ht)$$

which is the same as (20).

7.6. FINITE DIFFERENCES

If $f(x)$ be a function of an independent variable x , then in calculus we study the change in $f(x)$ corresponding to an infinitely small change in x and denote this change by δx or dx or Δx . Here we shall use the last notation.

The change Δx in x may be constant for all values of x or may vary for different values of x . Taking $\Delta x = h$ (a constant) the increment in function $f(x)$ will be $f(x+\Delta x) - f(x)$ which is represented by $\Delta f(x)$ i.e. the increment Δx in x corresponds an increment $\Delta f(x)$ in $f(x)$. This $\Delta f(x)$ is called as the *first difference* of $f(x)$ and the interval h as the interval of differencing. In other words the *first difference* of a function $f(x)$ denoted by $\Delta f(x)$ and read as 'Delta of x ' is defined as

$$\Delta f(x) = f(x+h) - f(x) \quad \dots (1)$$

where h is a fixed constant. In fact it is the difference in values of the function for two neighbouring values of x , h units apart.

The *second difference* $\Delta^2 f(x)$ is defined as the difference of the first difference of $f(x)$ for two neighbouring values of x , h units apart and written as

$$\Delta^2 f(x) = \Delta[\Delta f(x)] = \Delta[f(x+h) - f(x)] = \Delta f(x+h) - \Delta f(x) \quad \dots (2)$$

Similarly the *third difference* $\Delta^3 f(x)$ is the difference of the second difference of $f(x)$ for two neighbouring values of x , h units apart i.e.

$$\Delta^3 f(x) = \Delta[\Delta^2 f(x)] \quad \dots (3)$$

and in general the *nth difference* is $\Delta^n f(x) = \Delta[\Delta^{n-1} f(x)]$, $n=1, 2, \dots$ $\dots (4)$

In finding $\Delta f(x)$ by (1) it is clear that $f(x)$ is shifted to $f(x+h)$ and then $f(x)$ is subtracted from $f(x+h)$. The operation of shifting $f(x)$ to $f(x+h)$ is sometimes denoted by $Ef(x)$ i.e.,

$$f(x+h) = Ef(x) \quad \dots (5)$$

$$\text{As such } f(x+2h) = Ef(x+h) = E\{Ef(x)\} = E^2 f(x) \quad \dots (6)$$

$$f(x+3h) = Ef(x+2h) = E\{E^2 f(x)\} = E^3 f(x) \quad \dots (7)$$

and in general $f(x+nh) = E^n f(x) \quad \dots (8)$
It should be noted that Δ^n and E^n do not bear n as an exponent but they simply represent the n times operations.

Relation between Δ and E .

$$\begin{aligned} \Delta f(x) &= f(x+h) - f(x) \\ &= Ef(x) - f(x) \end{aligned}$$

$$\begin{aligned} Ef(x) &= \Delta f(x) + f(x) = (\Delta + 1)f(x), \\ E &\equiv \Delta + 1. \end{aligned} \quad \dots (9)$$

i.e. operators E and Δ are symbolically connected with this identity.

Difference table. The successive differences $\Delta f(x), \Delta^2 f(x), \dots, \Delta^r f(x)$ of $f(x)$ for different values of x are easily calculated from a table, known as *difference table*, by subtraction. In any difference table the independent variable (say x) is called *argument* and the corresponding value of the function [as $f(x)$] is said to be the *entry*. A difference table can be constructed as shown below.

DIFFERENCE TABLE

Argument x	Entry $f(x)$	First difference $\Delta f(x)$	Second difference $\Delta^2 f(x)$	Third difference $\Delta^3 f(x)$	Fourth difference $\Delta^4 f(x)$
a	$f(a)$	$\Delta f(a) = f(a+h) - f(a)$			
$a+h$	$f(a+h)$	$\Delta f(a+h) = f(a+2h) - f(a+h)$	$\Delta^2 f(a)$		
$a+2h$	$f(a+2h)$	$\Delta f(a+2h) = f(a+3h) - f(a+2h)$	$\Delta^2 f(a+h)$	$\Delta^3 f(a)$	
$a+3h$	$f(a+3h)$	$\Delta f(a+3h)$	$\Delta^2 f(a+2h)$		$\Delta^4 f(a)$
$a+4h$	$f(a+4h)$				

Clearly each of the $\Delta f(x)$, $\Delta^2 f(x)$, or $\Delta^3 f(x)$ etc. is obtained by subtracting the corresponding value from its succeeding value i.e.

$$\begin{aligned} \Delta^r f(x+kh) &= \Delta^{r-1} \Delta f(x+kh) \\ &= \Delta^{r-1} \{f(x+k+1)h - f(x+kh)\} \\ \Delta^r f(x+kh) &= \Delta^{r-1} f(x+k+1)h - \Delta^{r-1} f(x+kh) \end{aligned} \quad \dots (10)$$

for every value of k and r .

Giving different values to k and r , we get different results of the difference table.

In difference table its first given term i.e. $f(a)$ is called the *leading term* and the differences, $\Delta f(a)$, $\Delta^2 f(a)$, $\Delta^3 f(a)$ etc. are called *leading differences* of $f(a)$.

Difference of any order can be expressed in terms of the function alone.

$$(i) \Delta f(a) = f(a+h) - f(a). \quad \dots (11)$$

$$(ii) \quad f(a) = \Delta \{ \Delta f(a) \} = \Delta f \{ (a+h) - f(a) \} = \Delta f(a+h) - \Delta f(a) \\ = [f(a+2h) - f(a+h)] - [f(a+h) - f(a)] \\ = f(a+2h) - 2f(a+h) + f(a).$$

$$(iii) \quad \Delta^3 f(a) = \Delta \{ (\Delta^2 f(a)) \} = \Delta \{ f(a+2h) - 2f(a+h) + f(a) \} \\ = \Delta f(a+2h) - 2\Delta f(a+h) + \Delta f(a) \\ = \{ f(a+3h) - f(a+2h) \} - 2\{ f(a+2h) - f(a+h) \} + \{ f(a+h) - f(a) \} \\ = f(a+3h) - 3f(a+2h) + 3f(a+h) - f(a)$$

or $\Delta^3 f(a) = f(a+3h) - 3f(a+2h) + 3f(a+h) - f(a)$
Properties of the operators Δ and E .

(i) If a is a constant $\Delta a = 0$ and $Ea = a$.

(ii) $\Delta \{ af(x) \} = a \Delta f(x)$ and $E \{ af(x) \} = a Ef(x)$

(iii) $\Delta^r \Delta^s f(x) = \Delta^{r+s} f(x)$ and $E^r E^s f(x) = E^{r+s} f(x)$
 $[\because E \{ af(x) \} = af(x+h) = a \Delta f(x) + af(x)]$
 $[\because \Delta^r \Delta^s f(x) = (\Delta \Delta \dots r \text{ times}) (\Delta \Delta \dots s \text{ times}) f(x) = (\Delta \Delta \dots r+s \text{ times}) f(x) = \Delta^{r+s} f(x)]$

(iv) $\Delta \{ f(x) + \phi(x) \} = \Delta f(x) + \Delta \phi(x)$

and $E \{ f(x) + \phi(x) \} = Ef(x) + E\phi(x)$

$[\because E \{ f(x) + \phi(x) \} = f(x+h) + \phi(x+h) = Ef(x) + E\phi(x)]$

Formulation for n th difference

If we call $y_0, y_1, y_2, \dots, y_n$ the values of $f(x)$ corresponding to $f(x_0), f(x_0+h), \dots, f(x_0+nh)$ then by (1) we have

$$\Delta y_0 = y_1 - y_0$$

$$\Delta^2 y_0 = \Delta(\Delta y_0) = \Delta(y_1 - y_0) = \Delta y_1 - \Delta y_0 \\ = (y_2 - y_1) - (y_1 - y_0) = y_2 - 2y_1 + y_0$$

$$\Delta^3 y_0 = \Delta(\Delta^2 y_0) = \Delta(y_2 - 2y_1 + y_0) = \Delta y_2 - 2\Delta y_1 + \Delta y_0 \\ = (y_3 - y_2) - 2(y_2 - y_1) + (y_1 - y_0) \\ = y_3 - 3y_2 + 3y_1 - y_0$$

$$\text{Similarly } \Delta^4 y_0 = y_4 - 4y_3 + 6y_2 - 4y_1 + y_0$$

and so on.

Clearly, the coefficients in the expansion of $\Delta^n y_0$ will be the same as the coefficients in the expansion of $(x-1)^n$ i.e. $1, {}^nC_1, {}^nC_2, \dots, {}^nC_n$ and therefore it will follow by the induction method that

$$\Delta^n y_0 = \sum_{k=0}^n \frac{(-1)^k |n|}{|k| |n-k|} y_{n-k} \text{ or } \sum_{k=0}^n (-1)^k \left(\frac{n}{k} \right) y_{n-k} \quad (1)$$

where $\Delta^0 y_0 = y_0$ and in terms of $f(x)$ it becomes

$$\Delta^n f(x_0) = \sum_{k=0}^n \frac{(-1)^k |n|}{|k| |n-k|} f[(x_0 + (n-k)h)]$$

Problem 32. Evaluate

$$(a) \Delta x(x+1)(x+2)(x+3)$$

$$(b) \Delta^n (ab^{cx})$$

$$(c) \Delta^2 \left[\frac{5x+12}{x^2+5x+6} \right]$$

$$(d) \Delta^n (e^{a+bx})$$

$$(e) \Delta^n (ax^n + bx^{n-1})$$

$$(f) \Delta^n \left(\frac{1}{x} \right)$$

$$(g) \Delta (x!)$$

$$(a) \Delta x(x+1)(x+2)(x+3)$$

$$= (x+1)(x+2)(x+3)(x+4) - x(x+1)(x+2)(x+3)$$

$$= (x+1)(x+2)(x+3)\{(x+4)-x\}$$

$$= 4(x+1)(x+2)(x+3).$$

$$(b) \Delta^n (ab^{cx}) = \Delta^{n-1} (\Delta ab^{cx}).$$

$$\text{Now } \Delta (ab^{cx}) = ab^{c(x+1)} - ab^{cx} = ab^{cx} \{b^c - 1\}.$$

$$\therefore \Delta^2 (ab^{cx}) = \Delta (ab^{cx}) (b^c - 1) = (b^c - 1) \Delta (ab^{cx})$$

$$= (b^c - 1)^2 (ab^{cx}) \text{ and so on.}$$

$$\therefore \Delta^n (ab^{cx}) = (b^c - 1)^n (ab^{cx}).$$

$$(c) \Delta^2 \left[\frac{5x+12}{x^2+5x+6} \right] = \Delta^2 \left[\frac{2(x+3)+3(x+2)}{(x+2)(x+3)} \right]$$

$$= \Delta^2 \left[\frac{2}{(x+2)} + \frac{3}{(x+3)} \right]$$

$$= \Delta \cdot \Delta \left[\frac{2}{(x+2)} + \frac{3}{(x+3)} \right]$$

$$\text{Also } \Delta \left[\frac{2}{(x+2)} + \frac{3}{(x+3)} \right] = \left[\frac{2}{(x+3)} - \frac{2}{(x+2)} \right]$$

$$+ \left[\frac{3}{(x+4)} - \frac{3}{(x+3)} \right]$$

$$= \text{either } \frac{-2}{(x+2)(x+3)} = \frac{3}{(x+4)(x+3)} \text{ or } \frac{3}{(x+4)} - \frac{1}{(x+3)} - \frac{2}{(x+2)} \dots (1)$$

So that

$$\Delta^2 \left[\frac{5x+12}{x^2+5x+6} \right] = \Delta \left[-\frac{2}{(x+2)(x+3)} - \frac{3}{(x+3)(x+4)} \right]$$

$$= -2 \left[\frac{1}{(x+3)(x+4)} - \frac{1}{(x+2)(x+3)} \right]$$

$$- 3 \left[\frac{1}{(x+4)(x+5)} - \frac{1}{(x+3)(x+4)} \right]$$

$$= -2 \left[\frac{(x+2) - (x+4)}{(x+2)(x+3)(x+4)} \right]$$

$$- 3 \left[\frac{(x+3) - (x+5)}{(x+3)(x+4)(x+5)} \right]$$

$$\text{or } \Delta^2 \left[\frac{5x+12}{x^2+5x+6} \right] = \frac{4}{(x+2)(x+3)(x+4)} + \frac{6}{(x+3)(x+4)(x+5)}$$

$$(d) \Delta^n (e^{a+bx}) = \Delta^{n-1} (\Delta e^{a+bx}).$$

$$\therefore \Delta e^{a+bx} = e^{a+b(x+1)} - e^{a+bx} = e^{a+bx} \{e^b - 1\}$$

$$\therefore \Delta^2 e^{a+bx} = (e^b - 1)^2 e^{a+bx} \text{ and so on.}$$

$$\text{Hence } \Delta^n e^{a+bx} = (e^b - 1)^n e^{a+bx}.$$

$$(e) \Delta^n (ax^n + bx^{n-1}) = \Delta^n (ax^n) + \Delta^n (bx^{n-1}).$$

$$\text{But } \Delta^n (bx^{n-1}) = 0$$

(as bx^{n-1} is a polynomial of $(n-1)$ degree only)

$$\therefore \Delta^n (ax^n) = a \Delta^{n-1} (\Delta x^n) = a \Delta^{n-1} \{(x+1)^n - x^n\}$$

$$= a \Delta^{n-1} \{ \Delta x^{n-1} + \text{terms with powers less than } (n-1) \text{ in } x \}$$

$$= a \Delta^{n-1} nx^{n-1} = a \Delta^{n-2} \cdot n \{(n-1)x^{n-2} + \text{terms with powers less than } (n-2) \text{ in } x\}$$

$$= a \Delta^{n-3} n(n-1)(n-2)x^{n-3} \text{ and so on.}$$

$$\text{Ultimately, } \Delta^n (ax^n) = a(n!).$$

$$(f) \Delta^n \left(\frac{1}{x} \right) = \Delta^{n-1} \cdot \Delta \left(\frac{1}{x} \right).$$

$$\therefore \Delta \left(\frac{1}{x} \right) = \frac{1}{x+1} - \frac{1}{x} = \frac{(-1)}{x(x+1)}.$$

$$\therefore \Delta^2 \left(\frac{1}{x} \right) = \frac{(-1)^2}{x(x+1)(x+2)} \text{ and so on.}$$

$$\Delta^n \left(\frac{1}{x} \right) = \frac{(-1)^n}{x(x+1)(x+2)\dots(x+n)} - \frac{(-1)^n}{(x+1)^{(n+1)}} \\ = (-1)^n [(x+n) - (n+1)]^{(n+1)} = (-1)^n [x-1]^{(n+1)}$$

$$(g) \Delta |x| = |(x+1)| - |x| = |x| \{(x+1) - 1\} = x |x|.$$

7.7. POLYNOMIAL INTERPOLATION

The process of estimating the value of the function for any intermediate value of the variable with the help of certain given values of the function corresponding to a number of variable values, is called *Interpolation*.

In other words if a function $y=f(x)$ is known for values $x_1, x_2, x_3, \dots, x_n$ as $f(x_1), f(x_2), \dots, f(x_n)$, then the process of finding the value of $f(x)$ for some other value of x (lying between the values, $x_1, x_2, \dots, x_n$) is called *Interpolation*.

When we estimate $f(x)$ for some such variable value which lies outside the given values, the process is called *extrapolation*.

If the exact form of the function $f(x)$ is not known, or, known, in a complicated form, then to solve the problem of interpolation we assume that the function $f(x)$ can always be taken as a polynomial in x . Only after making this assumption the calculus of finite differences can be applied to obtain the value of function $f(x)$, for any unknown value of variable x . Moreover the polynomials are the simplest functions.

Validity of this assumption of replacing $f(x)$ by a polynomial function lies in an important theorem due to Weierstrass, which states that if $f(x)$ is continuous between x_1 and x_n , then it can be replaced by a Polynomial of suitable degree in that interval with as small an error as we please.

The degree of polynomial depends upon the number of given values of the function. In general if n values are given, we can fit a polynomial of degree $(n-1)$, no matter whatever be the exact degree of the polynomial.

This limitation of fitting a polynomial of degree $(n-1)$ only, when n values are known, creates two problems :

(i) If $f(x)$ is of a degree higher than $n-1$, then we shall interpolate the values of $f(x)$ only approximately.

(ii) If $f(x)$ is not actually of polynomial form but of a different form like exponential etc., then also the interpolated values will not be exactly the same. Such functions as exponential can be expanded upto infinity, but we can assume them only of degree $(n-1)$, if n values are given.

7.8. INTERPOLATION WITH EQUAL INTERVALS, (i.e., when function-values are given at equidistant intervals).

(i) Newton-Gregory Formula.

If n is a positive integer, then

$$f(x+nh) = f(x) + {}^nC_1 \Delta^2 f(x) + {}^nC_2 \Delta^2 f(x) + {}^nC_3 \Delta^3 f(x) + \dots + {}^nC_n \Delta^n f(x).$$

$$\text{We have } f(x+nh) = E^n f(x) = (1 + \Delta)^n f(x)$$

$$= \{1 + {}^nC_1 \Delta + {}^nC_2 \Delta^2 + \dots + {}^nC_n \Delta^n\} f(x)$$

$$= f(x) + {}^nC_1 \Delta f(x) + {}^nC_2 \Delta^2 f(x) + \dots + {}^nC_n \Delta^n f(x) \dots (1)$$

which is Newton-Gregory formula.

Here we get $f(x+nh)$ in terms of $f(x)$ and its leading differences.

Note 1. This formula can be applied, when there are n equidistant terms, and

(i) we want to obtain an intermediate value.

(ii) out of those one value is missing which we want to obtain,

(iii) out of those r values are missing which we want to determine from the $(n-r)$ known, values.

Note 2. The Gregory's expansion is convergent. But its nature of convergence may finish if n becomes either a negative quantity or a fractional one, and the formula may take a divergent shape. So the results obtained will be very much approximate and sometimes inaccurate also.

If the function remains a polynomial one, then Newton's advancing difference formula, where $f(x)$ is expressed in terms of $f(0)$ and its leading differences, can be used for all values of x , positive or negative, integral or fractional. Also this formula gives highly approximate results if $f(x)$ is of any other form than a polynomial.

[2] Newton's Advancing Difference Formula. By (8) of §7.6, taking $h=1$ we get, $f(x) = E^x f(0)$

$$= (1 + \Delta)^x f(0) = \{1 + {}^xC_1 \Delta + {}^xC_2 \Delta^2 + \dots\} f(0)$$

or $f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(0) + \dots$ (1)
which is Newton's advancing difference formula.

So long as $f(x)$ is a polynomial in x , this binomial expansion is valid for all values of x if starting value of variable is 0 and interval of differencing is unity.

[This formula can also be deduced from the Gregory formula by putting $a=0$ and $h=1$]

Problem 33. Show that Newton's formula

$$u_x = u_0 + x_{(1)} \Delta u_0 + x_{(2)} \Delta^2 u_0 + x_{(3)} \Delta^3 u_0 + \dots$$

can be put into the form

$$u_x = u_0 + x \Delta u_0 - xa \Delta^2 u_0 + xab \Delta^3 u_0 - xabc \Delta^4 u_0 + \dots$$

where $a = 1 - \frac{1}{2}(x+1)$, $b = 1 - \frac{1}{3}(x+1)$, $c = 1 - \frac{1}{4}(x+1)$, etc.

Hence show that the successive coefficients converge slowly and tend eventually to numerical equality.

$$\therefore a = 1 - \frac{1}{2}(x+1), \therefore -a = \frac{1}{2}(x+1) - 1 = \frac{1}{2}(x-1)$$

$$\text{Also } b = 1 - \frac{1}{3}(x+1), \therefore -b = \frac{1}{3}(x+1) - 1 = \frac{1}{3}(x-2)$$

Similarly

$$-c = \frac{1}{4}(x+1) - 1 = \frac{1}{4}(x-3)$$

$$-d = \frac{1}{5}(x+1) - 1 = \frac{1}{5}(x-4), \text{ etc.}$$

$$\text{Given } u_x = u_0 + \frac{x}{1} \Delta u_0 + \frac{x(x-1)}{1.2} \Delta^2 u_0 + \frac{x(x-1)(x-2)}{1.2.3} \Delta^3 u_0 + \frac{(x-1)(x-2)(x-3)}{2.3.4} \Delta^4 u_0 + \dots$$

$$\text{or } u_x = u_0 + x \Delta u_0 + x \left(\frac{x-1}{2} \right) \Delta^2 u_0 + x \left(\frac{x-1}{2} \right) \left(\frac{x-2}{3} \right) \Delta^3 u_0 + x \left(\frac{x-1}{2} \right) \left(\frac{x-2}{3} \right) \left(\frac{x-3}{4} \right) \Delta^4 u_0 + \dots$$

$$= u_0 + x \Delta u_0 + x(-a) \Delta^2 u_0 + x(-a)(-b) \Delta^3 u_0 + x(-a)(-b)(-c) \Delta^4 u_0 + \dots$$

$$\text{or } u_x = u_0 + x \Delta u_0 - xa \Delta^2 u_0 + xab \Delta^3 u_0 - xabc \Delta^4 u_0 + xabcd \Delta^5 u_0 - \dots$$

It is clear from the values of a, b, c and d etc. that successive coefficients in the expression converge.

Problem 34. Given the values $\sin 45^\circ = .7071$, $\sin 50^\circ = .7660$, $\sin 55^\circ = .8192$, $\sin 60^\circ = .8660$, find $\sin 52^\circ$.

θ	$\sin \theta$	$\Delta \sin \theta$	$\Delta^2 \sin \theta$	$\Delta^3 \sin \theta$
45	.7071			
50	.7660	.0589		
55	.8192	.0532	-.0057	
60	.8660	.0468	-.0068	-.0007

$$\sin 52^\circ = \sin (45 + 5.1 \cdot 4) = E^{1.4} \sin 45^\circ = (1 + \Delta)^{1.4} \sin 45^\circ$$

$$= \sin 45^\circ + 1.4 \Delta \sin 45^\circ + \frac{(1.4)(.4)}{2!} \Delta^2 \sin 45^\circ + \frac{(1.4)(.4)(1.4-2)}{3!} \Delta^3 \sin 45^\circ$$

$$= .7071 + (1.4)(.0589) + (.7)(.4)(-.0057) + (.7)(.4)(-.2)(-.0007)$$

$$= .7071 + .08246 - .001596 + 0.0000392$$

$$= .7880032 = 0.7880 \text{ approx.}$$

Problem 35. Given $\log x$ for $x=310, 320, 330, 340, 350$ and 360 according to the following table. Find the value of $\log 337.5$.

DIFFERENCE TABLE

	$\log x$	$\Delta \log x$	$\Delta^2 \log x$	$\Delta^3 \log x$	$\Delta^4 \log x$	$\Delta^5 \log x$
310	2.4913617					
320	2.5051500	.0137883				
330	2.5185139	.0133639	-.0004244			
340	2.5314781	.0129642	-.0003989	.0000255		
350	2.5440680	.0125891	-.0003751	.0000230	-.0000025	
360	2.5563025	.0122345	-.0003546	.0000205	-.0000025	0

We shall first evaluate $\log 337.5$.

$$\log 337.5 = \log (310 + 10 \times 2.75) = E^{2.7.5} \log 310$$

$$= \log 310 + 2.75 \Delta \log 310 + \frac{2.75(1.75)}{2!} \Delta^2 \log 310$$

$$+ \frac{(2.75)(1.75)(.75)}{3!} \Delta^3 \log 310$$

$$+ \frac{(2.75)(1.75)(.75)(-.25)}{4!} \Delta^4 \log 310$$

(taking approximation upto five places)

$$\log 337.5 = 2.49136 + 2.75(.01379) + \frac{2.75(1.75)}{2}(-.00042)$$

$$+ \frac{(2.75)(1.75)(.75)}{6}(.00003) \text{ {leaving other terms}}$$

$$= 2.49136 + 2.75(.01379) - (2.75)(1.75)(.00021)$$

$$+ (1.75)(1.75)(.75)(.00003)$$

$$= 2.49136 + .03792 - .00101 + .00005 = 2.52832$$

$$\text{Now } \log 3375 = \log 10 \times 337.5 = \log 10 + \log 337.5 = 1 + 2.52832.$$

$$\log 3375 = 3.52832.$$

Problem 36. The following table is given :

x	0	1	2	3	4
$f(x)$	3	6	11	18	27

What is the form of the function $f(x)$?

x	$f(x)$	$\Delta f(x)$	$\Delta^2 f(x)$	$\Delta^3 f(x)$
0	3			
1	6	3		
2	11	5	2	
3	18	7	2	0
4	27	9	2	0

$$\therefore f(x) = f(0) + {}^xC_1 \Delta f(0) + {}^xC_2 \Delta^2 f(0)$$

$$= 3 + x \cdot 3 + \frac{x(x-1)}{1 \cdot 2} \cdot 2 \quad \{\text{as third differences are zero}\}$$

$$\text{or } f(x) = 3 + 3x + x(x-1) = 3 + 3x + x^2 - x$$

$$\text{or } f(x) = x^2 + 2x + 3.$$

Problem 37. Given $f(0) = 1$, $f(1) + f(2) = 10$ and $f(3) + f(4) + f(5) = 65$, find $f(4)$.

Since we are given three relations, so we can assume $f(x)$ a polynomial of second degree.

$$\text{Suppose } f(x) = A + Bx + Cx^2.$$

$$\therefore f(0) = 1 \text{ gives } A = 1.$$

$$\text{and } f(1) + f(2) = 10 \text{ gives } (A + B + C) + (A + 2B + 4C) = 10$$

$$\text{or } 2A + 3B + 5C = 10$$

$$\text{or } 3B + 5C = 8$$

$$\text{and } f(3) + f(4) + f(5) = 65 \text{ gives}$$

$$(A + 3B + 9C) + (A + 4B + 16C) + (A + 5B + 25C) = 65$$

$$\text{or } 12B + 50C = 65 - 3 = 62.$$

$$(2) \text{ and } (3) \text{ give } C = 1 \text{ and } B = 1.$$

$$\therefore f(x) = x^2 + x + 1.$$

$$\text{Hence } f(4) = 21.$$

[3] Equidistant Terms with Terms Missing.

In general if p out of n equidistant values of the function are unknown, i.e., only $(n-p)$ function-values are known, then we shall

assume the function of degree $(n-p-1)$. Thus $(n-p)$ th difference of the function will be zero for every value of the variable. Equating these differences to zero, we may have as many equations as we like (in finding out missing terms) by giving different values to x .

Problem 38. Given $f(1) = 386$, $f(3) = 530$, $f(5) = 810$, find $f(2)$ and $f(4)$.

Since only three values of the function are known, so $f(x)$ may be assumed of second degree. Thus $\Delta^3 f(x) = 0$ for every value of x .

$$\Delta^3 f(1) = 0 \text{ gives } (E-1)^3 f(1) = 0$$

$$f(4) - 3f(3) + 3f(2) - f(1) = 0$$

$$f(4) + 3f(2) = 3f(3) + f(1) = 1976. \quad \dots (1)$$

$$\Delta^3 f(2) = 0 \text{ gives } f(5) - 3f(4) + 3f(3) - f(2) = 0$$

$$3f(4) + f(2) = f(5) + 3f(3) = 810 + 1590 = 2400. \quad \dots (2)$$

Solving (1) and (2), we get

$$f(2) = 441, f(4) = 653.$$

Problem 39. Estimate the missing terms in the following table where $f(x) = 3^x$.

x	1	2	3	4	5	6
$f(x)$	3	9	—	81	—	729

Explain why $f(3)$ differs from 3^3 and $f(5)$ from 3^5 .

As only four values are given so $\Delta^4 f(x) = 0$ for every x .

As above, $\Delta^4 f(1)$ gives

$$f(5) - 4f(4) + 6f(3) - 4f(2) + f(1) = 0 \quad \dots (1)$$

and $\Delta^4 f(2)$ gives

$$f(6) - 4f(5) + 6f(4) - 4f(3) + f(2) = 0 \quad \dots (2)$$

Substituting the given values in (1) and (2), we get

$$f(5) + 6f(3) = 357 \quad \dots (3)$$

$$f(5) + f(3) = 306 \quad \dots (4)$$

Solving (3) and (4), $f(3) = 10.2$ and $f(5) = 295.8$.

When estimating $f(3)$ and $f(5)$, we have assumed here that the function $f(x)$ is a polynomial of degree 3, whereas it is an exponential function. That is why $f(3)$ and $f(5)$ are differing from 3^3 and 3^5 , i.e. from 27 and 243 respectively.

19. INTERPOLATION WITH UNEQUAL INTERVALS

In §7.8 we have assumed that values of $f(x)$ are given at equidistant values of x , say for $a, a+h, a+2h$ etc, but it may happen that the values of $f(x)$ are not given at equal distances but are given at points say $x_0, x_1, x_2, x_3, \dots, x_n$, where $(x_1 - x_0), (x_2 - x_1), (x_3 - x_2), \dots, (x_n - x_{n-1})$ are not necessarily equal. We define divided differences to explain the interpolation formulae in such cases.

[1] Divided Differences

Suppose $f(x_0), f(x_1), \dots, f(x_n)$ are the values of $f(x)$ corresponding to values $x_0, x_1, \dots, x_n$ which are not necessarily equally spaced. Then

(i) $\frac{f(x_1)-f(x_0)}{x_1-x_0}, \frac{f(x_2)-f(x_1)}{x_2-x_1}, \dots, \text{etc.} \frac{f(x_n)-f(x_{n-1})}{x_n-x_{n-1}}$ are called *divided differences* of x for the variable-values $(x_1, x_0), (x_2, x_1), \dots, (x_n, x_{n-1})$ respectively. These differences may be denoted as $f(x_1, x_0), f(x_2, x_1), \dots, f(x_n, x_{n-1})$.

(ii) Similarly *second divided differences* are

$$\frac{f(x_2, x_1)-f(x_1, x_0)}{x_2-x_0}, \frac{f(x_3, x_2)-f(x_2, x_1)}{x_3-x_1} \text{ etc.}$$

denoted by $f(x_2, x_1, x_0), f(x_3, x_2, x_1)$ etc.

(iii) And similarly *third divided differences* are

$$\frac{f(x_3, x_2, x_1)-f(x_2, x_1, x_0)}{x_3-x_0}$$

denoted as $f(x_3, x_2, x_1, x_0)$ etc.

[2] The Divided Differences remain Unaffected by Changing the Permutations of their Variable Values.

$$\begin{aligned} \therefore (x_1, x_0) &= \frac{f(x_1)-f(x_0)}{x_1-x_0} = \frac{f(x_1)}{(x_1-x_0)} + \frac{f(x_0)}{(x_0-x_1)} \\ &= \frac{f(x_0)-f(x_1)}{x_0-x_1} = f(x_0, x_1). \end{aligned}$$

$$\text{Similarly } f(x_2, x_1, x_0) = \frac{f(x_2, x_1)-f(x_1, x_0)}{(x_2-x_0)}$$

$$\begin{aligned} &= \frac{\left\{ \frac{f(x_2)}{(x_2-x_1)} + \frac{f(x_1)}{(x_1-x_2)} \right\} - \left\{ \frac{f(x_1)}{(x_1-x_0)} + \frac{f(x_0)}{(x_0-x_1)} \right\}}{x_2-x_0} \\ &= \frac{f(x_2)}{(x_2-x_0)(x_2-x_1)} + \frac{f(x_1)}{(x_1-x_0)(x_1-x_2)} + \frac{f(x_0)}{(x_0-x_1)(x_0-x_2)} \\ &= f(x_0, x_1, x_2) = f(x_1, x_0, x_2) \text{ etc.} \end{aligned}$$

The n th divided difference is sometimes written as:

$$\begin{aligned} \Delta_{a_1, a_2, \dots, a_n}^n f(a_0) &= f(x_0, x_1, \dots, x_n) \\ &= \Sigma \frac{f(x_0)}{(x_0-x_1)(x_0-x_2) \dots (x_0-x_n)} \end{aligned}$$

Σ is the summation of all x 's.

[3] The n th divided differences of a polynomial of the n th degree are also constant. If $a_0 x^n$ is the leading term of the polynomial then n th divided difference will be a_0 .

Problem 40. Find $f(2, 4, 9, 10)$ where $f(x) = x^4 + x^2 + 1$.

$\therefore \Delta_{4, 9, 10}^3 f(2)$ or $f(2, 4, 9, 10) = 25$. [See Table on the next page]

$f(x)$	$\Delta f(x)$	$\Delta^2 f(x)$	$\Delta^3 f(x)$
21	$\frac{273-21}{4-2} = 126$	$\frac{1274-126}{9-2} = 164$	
273	$\frac{6643-273}{9-4} = 1274$	$\frac{3458-1274}{10-4} = 364$	$\frac{364-164}{10-2} = 25$
6643	$\frac{10101-6643}{10-9} = 3458$	$\frac{10101-6643}{10-9} = 3458$	$\frac{10101-6643}{10-9} = 3458$
10101			

[4] Newton's Divided Difference Formula

Suppose $f(x_0), f(x_1), \dots, f(x_n)$ are the $(n+1)$ values corresponding to values $x_0, x_1, \dots, x_n$. With these $(n+1)$ values we can fit a polynomial of degree n . Thus n th divided difference $f(x_0, x_1, x_2, \dots, x_n) = \text{constant}$ and $(n+1)$ th divided difference $f(x, x_0, x_1, \dots, x_n) = 0$, where x is some value other than $x_0, x_1, \dots, x_n$.

$$f(x, x_0, x_1, \dots, x_n) = \frac{f(x, x_0, x_1, \dots, x_{n-1}) - f(x_0, x_1, \dots, x_n)}{x - x_n} = 0$$

$$f(x, x_0, x_1, \dots, x_{n-1}) = f(x_0, x_1, \dots, x_n)$$

$$f(x, x_0, x_1, \dots, x_{n-2}) - f(x_0, x_1, \dots, x_{n-1}) = f(x_0, x_1, \dots, x_n)$$

$$f(x, x_0, x_1, \dots, x_{n-2}) = f(x_0, x_1, \dots, x_{n-1}) + (x - x_{n-1}) f(x_0, x_1, \dots, x_n)$$

$$f(x, x_0, x_1, \dots, x_{n-3}) = f(x_0, x_1, \dots, x_{n-2}) + (x - x_{n-2}) f(x_0, x_1, \dots, x_{n-1})$$

$$f(x, x_0, x_1, \dots, x_{n-4}) = f(x_0, x_1, \dots, x_{n-3}) + (x - x_{n-3}) f(x_0, x_1, \dots, x_{n-2})$$

and so on.

The last but one step will be

$$\begin{aligned} f(x, x_0) &= f(x_0) + (x - x_1) f(x_0, x_1, x_2) \\ &\quad + (x - x_1)(x - x_2) f(x_0, x_1, x_2, x_3) + \dots \\ &\quad + (x - x_1)(x - x_2) \dots (x - x_{n-1}) f(x_0, x_1, \dots, x_n) \\ &= \frac{f(x) - f(x_0)}{x - x_0} \\ f(x) &= f(x_0) + (x - x_0) f(x_0, x_1) + (x - x_0)(x - x_1) f(x_0, x_1, x_2) \\ &\quad + \dots + (x - x_0) \dots (x - x_{n-1}) f(x_0, \dots, x_n) \\ f(x) &= f(x_0) + (x - x_0) \Delta_{x_1} f(x_0) + (x - x_0)(x - x_1) \Delta_{x_1, x_2}^2 f(x_0) + \dots \end{aligned}$$

$$+ \dots + (x-x_0)(x-x_1)\dots(x-x_{n-2}) \frac{\Delta^{n-1} f(x_0)}{x_1, x_2, \dots, x_{n-1}} \\ + (x-x_0)(x-x_1)\dots(x-x_{n-1}) \frac{\Delta^n f(x_0)}{x_1, x_2, \dots, x_n}$$

which is Newton's divided difference formula.

[5] Newton's Gregory Formula as a Particular Case of Newton's divided Difference Formula.

In the above Newton's divided difference formula, if we put $x = x_0 + nh$, $x_1 = x_0 + h$, $x_2 = x_0 + 2h$, $x_3 = x_0 + 3h$ etc., we get the various terms as under:

$$(x-x_0)f(x_0, x_1) = nh \frac{f(x_1) - f(x_0)}{x_1 - x_0} = nh \frac{f(x_0+h) - f(x_0)}{h} \\ = nh \frac{\Delta f(x_0)}{h} = n \Delta f(x_0) = {}^nC_1 \Delta f(x_0) \\ (x-x_0)(x-x_1)f(x_0, x_1, x_2) = (nh)(n-1)h \frac{f(x_1, x_2) - f(x_0, x_1)}{x_2 - x_0} \\ = (nh)(n-1)h \frac{\Delta f(x_1) - \Delta f(x_0)}{2h} \\ = \frac{n(n-1)}{2} \Delta \{f(x_0+h) - f(x_0)\} \\ = \frac{n(n-1)}{2} \Delta^2 f(x_0) = {}^nC_2 \Delta^2 f(x_0)$$

Similarly,

$$(x-x_0)(x-x_1)(x-x_2)f(x_0, x_1, x_2, x_3) = \frac{n(n-1)(n-2)}{(3)} \Delta^3 f(x_0), \\ \text{and finally, } (x-x_0)(x-x_1)\dots(x-x_{n-1})f(x_0, x_1, x_2, \dots, x_n) \\ = \frac{n(n-1)(n-2)\dots(n-n-1)}{(n)} \Delta^n f(x_0) \\ = {}^nC_n \Delta^n f(x_0).$$

Substituting these values in the Newton's divided difference formula, we get

$$f(x_0 + nh) = f(x_0) + {}^nC_1 \Delta f(x_0) + {}^nC_2 \Delta^2 f(x_0) + \dots + {}^nC_n \Delta^n f(x_0).$$

Problem 41. Find the polynomial of the lowest possible degree which assumes the values 1245, 33, 5, 9 and 1355 when x has the values -4, -1, 0, 2 and 5 respectively. Also obtain the value of the polynomial at the abscissa 1.

We have to find a polynomial satisfied by $(-4, 1245)$, $(-1, 33)$, $(0, 5)$, $(2, 9)$, and $(5, 1355)$ and the value for '1' is to be interpolated. The difference table is given on the next page.

$$\therefore f(x) = f(x_0) + (x-x_0)f(x_0, x_1) + (x-x_0)(x-x_1)f(x_0, x_1, x_2) \\ + (x-x_0)(x-x_1)(x-x_2)f(x_0, x_1, x_2, x_3) \\ + (x-x_0)(x-x_1)(x-x_2)(x-x_3)f(x_0, x_1, x_2, x_3, x_4) \\ = 1245 - (x+4)404 + (x+4)(x+1)94 - (x+4)(x+1)x.14 \\ + (x+4)(x+1)x(x-2)3$$

$f(x)$	$\Delta f(x)$	$\Delta^2 f(x)$	$\Delta^3 f(x)$	$\Delta^4 f(x)$
1245	$\frac{1245-33}{-4-(-1)} = -404$	$\frac{-404-(-28)}{-4-0} = 94$	$\frac{94-10}{-4-2} = -14$	$\frac{-14-13}{-4-5} = 3$
33	$\frac{33-5}{-1-(0)} = -28$	$\frac{-28-2}{-1-2} = 10$	$\frac{10-88}{-1-5} = 13$	
5	$\frac{5-9}{0-2} = 2$	$\frac{2-442}{0-5} = 88$		
9	$\frac{9-1355}{2-5} = 442$			
1355				

$$f(x) = 5 - 14x + 6x^2 - 5x^3 + 3x^4.$$

$$\text{When } x=1, f(x) = 5 - 14 + 6 - 5 + 3 = -5.$$

[6] Lagrange's Formula.

Suppose $(n+1)$ values of $f(x)$ are known as $f(x_0), f(x_1), \dots, f(x_n)$ at points $x_0, x_1, \dots, x_n$, which are not necessarily equally spaced. Thus $f(x)$ can be taken as a polynomial of degree n and one of the possible forms of $f(x)$ can be taken as under

$$f(x) = A_0(x-x_1)(x-x_2)\dots(x-x_n) + A_1(x-x_0)(x-x_2)\dots(x-x_n) \\ + A_2(x-x_0)(x-x_1)(x-x_3)\dots(x-x_n) + \dots \\ + A_n(x-x_0)\dots(x-x_{n-1})$$

At the point $x=x_0$, $f(x)=f(x_0)$. Putting $x=x_0$ and $f(x)=f(x_0)$, we get

$$A_0 = \frac{f(x_0)}{(x_0-x_1)(x_0-x_2)\dots(x_0-x_n)}.$$

Also at the point $x=x_1$, $f(x)=f(x_1)$. This substitution gives us

$$A_1 = \frac{f(x_1)}{(x_1-x_0)(x_1-x_2)\dots(x_1-x_n)}.$$

$$\text{Similarly, } A_2 = \frac{f(x_2)}{(x_2-x_0)(x_2-x_1)\dots(x_2-x_n)}.$$

$$\dots \dots \dots \dots \dots \dots$$

$$A_n = \frac{f(x_n)}{(x_n-x_0)(x_n-x_1)\dots(x_n-x_{n-1})}.$$

With these values, we get

$$f(x) = \frac{(x-x_1)(x-x_2)\dots(x-x_n)}{(x_0-x_1)(x_0-x_2)\dots(x_0-x_n)} f(x_0) + \dots$$

$$+ \frac{(x-x_0)(x-x_2)\dots(x-x_n)}{(x_1-x_0)(x_1-x_2)\dots(x_1-x_n)} f(x_1) + \dots$$

$$+ \frac{(x-x_0)(x-x_1)(x-x_2)\dots(x-x_{n-1})}{(x_n-x_0)(x_n-x_1)(x_n-x_2)\dots(x_n-x_{n-1})} f(x_n)$$

which is Lagrange's Formula.

Problem 42. By means of Lagrange's formula, prove that

(i) $y_1 = y_3 - 3(y_5 - y_3) + 2(y_3 - y_5)$ approximately.

(ii) $y_0 = \frac{1}{2}(y_1 + y_{-1}) - \frac{1}{8}[\frac{1}{2}(y_3 - y_1) - \frac{1}{2}(y_{-1} - y_{-3})]$ approximately.

(i) Here we are given y_{-5}, y_{-3}, y_3 and y_5 and y_1 is to be determined

$$\therefore y_1 = \frac{(1+3)(1-3)(1-5)}{(-5+3)(-5-3)(-5-5)} y_{-5}$$

$$+ \frac{(1+5)(1-3)(1-5)}{(-3+5)(-3-3)(-3-5)} y_{-3} + \frac{(1+5)(1+3)(1-5)}{(3+5)(3+3)(3-5)} y_3$$

$$+ \frac{(1+5)(1+3)(1-3)}{(5+5)(5+3)(5-3)} y_5$$

$$\text{or } y_1 = \frac{4(-2)(-4)}{(-2)(-8)(-10)} y_{-5} + \frac{(6)(-2)(-4)}{(2)(-6)(-8)} y_{-3}$$

$$+ \frac{(6)(4)(-4)}{(8)(6)(-2)} y_3 + \frac{(6)(4)(-2)}{(10)(8)(2)} y_5$$

$$\text{or } y_1 = -\frac{2}{15} y_{-5} + \frac{1}{5} y_{-3} + y_3 - \frac{3}{15} y_5 - 2y_{-5} + 5y_{-3} - 3y_5$$

$$= y_3 - 3(y_5 - y_{-3}) + 2(y_{-3} - y_{-5})$$

(ii) Here we are given the values of y_{-3}, y_{-1}, y_1 and y_3 and y_0 is to be determined.

$$y_0 = \frac{(0+1)(0-1)(0-3)}{(-3+1)(-3-1)(-3-3)} y_{-3} + \frac{(0+3)(0-1)(0-3)}{(-1+3)(-1-1)(-1-3)} y_{-1}$$

$$+ \frac{(+3)(+1)(-3)}{(1+3)(1+1)(1-3)} y_1 + \frac{(+3)(+1)(-1)}{(3+3)(3+1)(3-1)} y_3$$

$$\text{or } y_0 = -\frac{3}{15} y_{-3} + \frac{9}{15} y_{-1} + \frac{9}{15} y_1 - \frac{3}{15} y_3$$

$$= \frac{3}{15} (y_1 + y_{-1}) - \frac{1}{15} [(y_3 - y_1) - (y_{-1} - y_{-3})]$$

$$\text{or } y_0 = \frac{1}{2} (y_1 + y_{-1}) - \frac{1}{8} [\frac{1}{2} (y_3 - y_1) - \frac{1}{2} (y_{-1} - y_{-3})]$$

7.10. CENTRAL DIFFERENCE FORMULAE

When applying Newton's Gregory formula with $(n+1)$ known values of the variable, we assumed the function as n th degree polynomial for finding out any intermediate value, but actually the polynomial happens to be different from n th degree, so Newton's formula gives an approximate result only. For better results central difference formulae are introduced.

Suppose function $f(x)$ takes the values $f(0), f(1), f(2), f(3), f(4), f(5), f(6), f(7), \dots$, equally spaced having unit interval.

If we shift our origin to the 4th point then $f(x)$ will become $f(-3), f(-2), f(-1), f(0), f(1), f(2), f(3), \dots$

(i) Gauss' Forward Formula.

Newton's advancing difference formula (in which interval is h and zero is the starting value of the variable) is

$$f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(0) + {}^x C_3 \Delta^3 f(0) + \dots + {}^x C_r \Delta^r f(0)$$

$$\Delta^3 f(0) = \Delta^3 f(-1) + \Delta^2 f(-1)$$

$$[\because \Delta^r f(-1) = \Delta^{r-1} \{f(0) - f(-1)\}]$$

$$\Delta^3 f(0) = \Delta^4 f(-1) + \Delta^3 f(-1)$$

$$\text{Similarly, } \Delta^4 f(0) = \Delta^5 f(-1) + \Delta^4 f(-1) + \Delta^3 f(-1)$$

$$= \Delta^4 f(-2) + \Delta^3 f(-2),$$

$$\Delta^4 f(-1) = \Delta^5 f(-2) + \Delta^4 f(-2) \text{ etc.}$$

In general $\Delta^r f(p) = \Delta^{r+1} f(p-1) + \Delta^r (p-1)$, for every integral values of r and p .

Substituting these transformations in advancing difference formula we get

$$f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(0) + {}^x C_3 \Delta^3 f(0) + {}^x C_4 \Delta^4 f(0)$$

$$+ {}^x C_5 \Delta^5 f(0) + \dots$$

$$= f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \{\Delta^2 f(-1) + \Delta^3 f(-1)\}$$

$$+ {}^x C_3 \{\Delta^3 f(-1) + \Delta^4 f(-1)\} + {}^x C_4 \{\Delta^4 f(-1) + \Delta^5 f(-1)\}$$

$$+ {}^x C_5 \{\Delta^5 f(-1) + \Delta^6 f(-1)\} \text{ and so on}$$

$$f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(-1) + ({}^x C_2 + {}^x C_3) \Delta^3 f(-1)$$

$$+ ({}^x C_3 + {}^x C_4) \Delta^4 f(-1) + ({}^x C_4 + {}^x C_5) \Delta^5 f(-1)$$

$$+ ({}^x C_5 + {}^x C_6) \Delta^6 f(-1) \text{ and so on}$$

$$f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(-1) + {}^{x+1} C_3 \Delta^3 f(-1)$$

$$+ {}^{x+1} C_4 \Delta^4 f(-1) + {}^{x+1} C_5 \Delta^5 f(-1) + {}^{x+1} C_6 \Delta^6 f(-1) + \dots$$

$$\text{and so on.}$$

(Now we have to transform Δ 's of $f(-1)$ into Δ 's of $f(-2)$ after 6th term, and then Δ 's of $f(-2)$ into Δ 's of $f(-3)$ after 6th term and then after 8th term Δ 's of $f(-3)$ into Δ 's of $f(-4)$ and so on.)

$$f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(-1) + {}^{x+1} C_3 \Delta^3 f(-1)$$

$$+ {}^{x+1} C_4 \{\Delta^4 f(-2) + \Delta^5 f(-2)\} + {}^{x+1} C_5 \{\Delta^5 f(-2)$$

$$+ \Delta^6 f(-2)\} + {}^{x+1} C_6 \{\Delta^6 f(-2) + \Delta^7 f(-2)\} + \dots$$

$$f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(-1) + {}^{x+1} C_3 \Delta^3 f(-1)$$

$$+ {}^{x+1} C_4 \Delta^4 f(-2) + \{{}^{x+1} C_4 + {}^{x+1} C_5\} \Delta^5 f(-2)$$

$$+ \{{}^{x+1} C_5 + {}^{x+1} C_6\} \Delta^6 f(-2) + \{{}^{x+1} C_6 + {}^{x+1} C_7\} \Delta^7 f(-2) + \dots$$

$$f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(-1) + {}^{x+1} C_3 \Delta^3 f(-1)$$

$$+ {}^{x+1} C_4 \Delta^4 f(-2) + {}^{x+2} C_5 \Delta^5 f(-2) + {}^{x+2} C_6 \Delta^6 f(-2)$$

$$+ {}^{x+2} C_7 \Delta^7 f(-2) + \dots$$

$$f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(-1) + {}^{x+1} C_3 \Delta^3 f(-1)$$

$$+ {}^{x+1} C_4 \Delta^4 f(-2) + {}^{x+2} C_5 \Delta^5 f(-2) + {}^{x+2} C_6 \Delta^6 f(-3)$$

$$+ {}^{x+3} C_7 \Delta^7 f(-3) + \dots$$

$$\{\because \Delta^6 f(-2) = \Delta^7 f(-3) + \Delta^6 f(-3)\}$$

which is Gauss' forward formula.

If origin is not 0 and interval is not unity then, in general,

$$f(a+xh) = f(a) + {}^xC_1 \Delta f(a) + {}^xC_2 \Delta^2 f(a-h) + {}^{x+1}C_3 \Delta^3 f(a-h) \\ + {}^{x+1}C_4 \Delta^4 f(a-2h) + {}^{x+2}C_5 \Delta^5 f(a-2h) + \dots$$

This formula contains the odd differences just below the central line through $f(0)$ and even differences on the central line. The following table will make it clear :

[2] A central difference table can be prepared as follows :

x	$f(x)$	$\Delta f(x)$	$\Delta^2 f(x)$	$\Delta^3 f(x)$	$\Delta^4 f(x)$	$\Delta^5 f(x)$
-3	$f(-3)$					
-2	$f(-2)$	$\Delta f(-3)$	$\Delta^2 f(-3)$			
-1	$f(-1)$	$\Delta f(-2)$	$\Delta^2 f(-2)$	$\Delta^3 f(-3)$		
0	$f(0)$	$\Delta f(-1)$	$\Delta^2 f(-1)$	$\Delta^3 f(-2)$	$\Delta^4 f(-3)$	$\Delta^5 f(-3)$
1	$f(1)$	$\Delta f(0)$	$\Delta^2 f(0)$	$\Delta^3 f(-1)$	$\Delta^4 f(-1)$	$\Delta^5 f(-1)$
2	$f(2)$	$\Delta f(1)$	$\Delta^2 f(1)$	$\Delta^3 f(0)$	$\Delta^4 f(0)$	
3	$f(3)$	$\Delta f(2)$	$\Delta^2 f(2)$	$\Delta^3 f(1)$		
4	$f(4)$	$\Delta f(3)$				

[3] Gauss' Backward Formula.

Here Newton's advancing difference formula is so transformed as to have a formula involving odd differences just about the central line through $f(0)$ and even differences as before i.e. on the line.

$$\therefore f(x) = f(0) + {}^xC_1 \Delta f(0) + {}^xC_2 \Delta^2 f(0) + {}^xC_3 \Delta^3 f(0) + {}^xC_4 \Delta^4 f(0) + \dots \\ = f(0) + {}^xC_1 \{ \Delta f(-1) + \Delta^2 f(-1) \} + {}^xC_2 \{ \Delta^2 f(-1) \\ + \Delta^3 f(-1) \} + {}^xC_3 \{ \Delta^3 f(-1) + \Delta^4 f(-1) \} + \dots \\ = f(0) + {}^xC_1 \Delta f(-1) + ({}^xC_1 + {}^xC_2) \Delta^2 f(-1) + ({}^xC_2 + {}^xC_3) \Delta^3 f(-1) \\ + ({}^xC_3 + {}^xC_4) \Delta^4 f(-1) + \dots$$

$$\text{or } f(x) = f(0) + {}^xC_1 \Delta f(-1) + {}^{x+1}C_2 \Delta^2 f(-1) \\ + {}^{x+1}C_3 \{ \Delta^3 f(-1) + \Delta^4 f(-1) \} + {}^{x+1}C_4 \Delta^5 f(-1) + \dots$$

$$\text{or } f(x) = f(0) + {}^xC_1 \Delta f(-1) + {}^{x+1}C_2 \Delta^2 f(-1) + {}^{x+1}C_3 \{ \Delta^3 f(-2) \\ + \Delta^4 f(-1) + \Delta^5 f(-2) \} + \dots$$

$$\text{or } f(x) = f(0) + {}^xC_1 \Delta f(-1) + {}^{x+1}C_2 \Delta^2 f(-1) \\ + {}^{x+1}C_3 \Delta^3 f(-2) + {}^{x+2}C_4 \Delta^4 f(-2) + \dots$$

Thus the backward formula is

$$f(x) = f(0) + {}^xC_1 \Delta f(-1) + {}^{x+1}C_2 \Delta^2 f(-1) + {}^{x+1}C_3 \Delta^3 f(-2) \\ + {}^{x+2}C_4 \Delta^4 f(-2) + {}^{x+3}C_5 \Delta^5 f(-3) + \dots$$

In case origin is a and interval of differencing not unity, then
(taking h negative)

$$f(a-nh) = f(a+n(-h))$$

$$\Delta f(a) = f(a+h) - f(a) = f(a-h) - f(a) = -\Delta f(a-h).$$

$$\Delta^2 f(a-h) = f(a+h) - 2f(a) + f(a-h)$$

$$= f(a-h) - 2f(a) + f(a+h) = \Delta^2 f(a-h).$$

Thus even differences remain as such whereas odd ones change their signs.

$$f(a-nh) = f(a) - {}^nC_1 \Delta f(a-h) + {}^nC_2 \Delta^2 f(a-h) \\ - {}^{(n+1)}C_3 \Delta^3 f(a-2h) + {}^{(n+1)}C_4 \Delta^4 f(a-2h) - {}^{(n+1)}C_5 \Delta^5 f(a-3h) + \dots$$

[4] Newton's Stirling Formula.

This formula is obtained directly from forward formulae.

$$f(a+nh) = f(a) + {}^nC_1 \Delta f(a) + {}^nC_2 \Delta^2 f(a-h) \\ + {}^{n+1}C_3 \Delta^3 f(a-h) + {}^{n+1}C_4 \Delta^4 f(a-2h) \\ + {}^{n+2}C_5 \Delta^5 f(a-2h) + {}^{n+2}C_6 \Delta^6 f(a-3h) + \dots$$

$$= f(a) + n \Delta f(a) + \frac{n^2-n}{2} \Delta^2 f(a-h) + \frac{n(n^2-1^2)}{3} \Delta^3 f(a-h)$$

$$+ \frac{(n+1)n(n-1)(n-2)}{4} \Delta^4 f(a-2h)$$

$$+ \frac{(n^2-2^2)(n^2-1^2)n}{5} \Delta^5 f(a-2h) + \dots$$

$$= f(a) + n \frac{2 \Delta f(a) - \Delta^2 f(a-h)}{2} + \frac{n^2}{2} \Delta^2 f(a-h)$$

$$+ \frac{n(n^2-1^2)}{3} \frac{2 \Delta^3 f(a-h) - \Delta^4 f(a-2h)}{2}$$

$$+ \frac{n^2(n^2-1^2)}{4} \Delta^4 f(a-2h)$$

$$+ \frac{n(n^2-1^2)(n^2-2^2)}{5} \frac{2 \Delta^5 f(a-2h) - \Delta^6 f(a-3h)}{2}$$

$$+ \frac{n^2(n^2-1^2)(n^2-2^2)}{6} \Delta^6 f(a-3h) + \dots$$

$$\therefore f(a+nh) = f(a) + n \frac{\Delta f(a) + \Delta f(a-h)}{2} + \frac{n^2}{2} \Delta^2 f(a-h)$$

$$+ \frac{n(n^2-1^2)}{3} \frac{\Delta^3 f(a-h) + \Delta^3 f(a-2h)}{2}$$

$$+ \frac{n^2(n^2-1^2)}{4} \Delta^4 f(a-2h)$$

$$+ \frac{n(n^2-1^2)(n^2-2^2)}{5} \frac{\Delta^5 f(a-2h) + \Delta^5 f(a-3h)}{2}$$

$$+ \frac{n^2(n^2-1^2)(n^2-2^2)}{6} \Delta^6 f(a-3h) + \dots$$

which is Stirling's formula.

If $a=0$, $h=1$, and $n=x$, we get the formula as

$$f(x) = f(0) + \frac{x}{1} \frac{\Delta f(0) + \Delta f(-1)}{2} + \frac{x^2}{2} \Delta^2 f(-1) \\ + \frac{x(x^2-1^2)}{3} \frac{\Delta^3 f(-1) + \Delta^3 f(-2)}{2} \\ + \frac{x^2(x^2-1^2)}{4} \Delta^4 f(-2) + \frac{x(x^2-1^2)(x^2-2^2)}{5} \\ \times \frac{\Delta^5 f(-2) + \Delta^5 f(-3)}{2} + \dots$$

The central difference table shows that this formula involves the mean of the odd differences above and below the central line and even differences on the central line. This gives very good results in successive approximations as it decreases or increases uniformly.

[5] Bessel's Formula.

Gauss backward formula is

$$f(x) = f(0) + {}^x C_1 \Delta f(-1) + {}^{x+1} C_2 \Delta^2 f(-1) \\ + {}^{x+1} C_3 \Delta^3 f(-2) + {}^{x+2} C_4 \Delta^4 f(-2) + \dots \quad \dots (1)$$

And forward formula is

$$f(x) = f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(0) \\ + {}^{x+1} C_3 \Delta^3 f(-1) + {}^{x+1} C_4 \Delta^4 f(-2) + \dots \quad \dots (2)$$

Shifting the origin to 1 in backward formula (1) i.e. at 4 in the original data, we get

$$f(x) = f(1) + {}^{x-1} C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(0) \\ + {}^x C_3 \Delta^3 f(-1) + {}^{x+1} C_4 \Delta^4 f(-1) + \dots \quad \dots (3)$$

Taking mean of (2) and (3), we get

$$f(x) = \frac{1}{2} [f(0) + f(1)] + \frac{1}{2} [{}^x C_1 + {}^{x-1} C_1] \Delta f(0) \\ + {}^x C_2 \frac{\Delta^2 f(0) + \Delta^2 f(-1)}{2} + \frac{1}{2} [{}^{x+1} C_3 + {}^x C_3] \Delta^3 f(-1) \\ + {}^{x+1} C_4 \frac{\Delta^4 f(-1) + \Delta^4 f(-2)}{2} + \frac{1}{2} [{}^{x+1} C_5 + {}^{x+2} C_5] \Delta^5 f(-2) + \dots$$

$$\text{or } f(x) = \frac{1}{2} [f(0) + f(1)] + (x - \frac{1}{2}) \Delta f(0) + \frac{x(x-1)}{2} \frac{\Delta^2 f(0) + \Delta^2 f(-1)}{2} \\ + \frac{(x-\frac{1}{2})x(x-1)}{3} \Delta^3 f(-1) + \frac{(x+1)x(x-1)(x-2)}{4} \\ \times \frac{\Delta^4 f(-1) + \Delta^4 f(-2)}{2} + \frac{(x-\frac{1}{2})(x+1)x(x-1)(x-2)}{5} \\ \times \Delta^5 f(-2) + \dots$$

which is Bessel's formula.

If we replace $x - \frac{1}{2}$ by t , i.e. x by $t + \frac{1}{2}$, we get

$$f(t + \frac{1}{2}) = \frac{1}{2} [f(0) + f(1)] + t \Delta f(0) \\ + \frac{t^2 - (\frac{1}{2})^2}{2} \frac{\Delta^2 f(0) + \Delta^2 f(-1)}{2} + \frac{t \{t^2 - (\frac{1}{2})^2\}}{3} \Delta^3 f(-1) \\ + \frac{\{t^2 - (\frac{1}{2})^2\} \{t^2 - (\frac{3}{2})^2\}}{4} \frac{1}{2} \{\Delta^4 f(-1) + \Delta^4 f(-2)\} \\ + \frac{t \{t^2 - (\frac{1}{2})^2\} \{t^2 - (\frac{5}{2})^2\}}{5} \Delta^5 f(-2) + \dots$$

which is rather convenient shape of Bessel's formula for practical purposes {but only if $x > \frac{1}{2}$ }.

Here we have alternately mean of the differences falling on both sides of central line and differences lying upon central line.

[6] Laplace-Everett's Formula.

Gauss forward formula is

$$f(x) = f(0) + x \Delta f(0) + {}^x C_2 \Delta^2 f(-1) + {}^{x+1} C_3 \Delta^3 f(-1) + {}^{x+1} C_4 \Delta^4 f(-2) \\ + {}^{x+2} C_5 \Delta^5 f(-2) + \dots \quad \dots (4)$$

$$\Delta^{2k+1} f(a) = \Delta^{2k} f(a+h) - \Delta^{2k} f(a).$$

With the help of this relation, eliminating odd differences from (4), we get

$$f(x) = f(0) + {}^x C_1 \{f(1) - f(0)\} + {}^x C_2 \Delta^2 f(-1) + {}^{x+1} C_3 \{\Delta^2 f(0) \\ - \Delta^2 f(-1)\} + {}^{x+1} C_4 \Delta^4 f(-2) + {}^{x+2} C_5 \{\Delta^4 f(-1) - \Delta^4 f(-2)\} \\ + {}^{x+2} C_6 \Delta^6 f(-3) + \dots$$

$$\text{or } f(x) = (1-x) f(0) + {}^x C_1 f(1) + ({}^x C_2 - {}^{x+1} C_3) \Delta^2 f(-1) \\ + {}^{x+1} C_3 \Delta^2 f(0) + ({}^{x+1} C_4 - {}^{x+2} C_5) \Delta^4 f(-2) + {}^{x+2} C_5 \Delta^4 f(-1) \\ + ({}^{x+2} C_6 - {}^{x+3} C_7) \Delta^6 f(-3) + {}^{x+3} C_7 \Delta^6 f(-2) \\ + ({}^{x+3} C_8 - {}^{x+4} C_9) \Delta^8 f(-4) + \dots$$

$$\text{or } f(x) = (1-x) f(0) + x f(1) + \left[\frac{x^{(2)}}{2} - \frac{(x+1)^{(3)}}{3} \right] \Delta^2 f(-1) \\ + \frac{(x+1)^{(3)}}{3} \Delta^2 f(0) + \left(\frac{(x+1)^{(4)}}{4} - \frac{(x+2)^{(5)}}{5} \right) \Delta^4 f(-2) \\ + \frac{(x+2)^{(5)}}{5} \Delta^4 f(-1) + \dots$$

$$\text{or } f(x) = \left[(1-x) f(0) + \left\{ \frac{x^{(2)}}{2} + \frac{(x+1)^{(3)}}{3} \right\} \Delta^2 f(-1) \right. \\ \left. + \left\{ \frac{(x+1)^{(4)}}{4} - \frac{(x+2)^{(5)}}{5} \right\} \Delta^4 f(-2) + \dots \right] \\ + \left[x f(1) + \frac{(x+1)^{(3)}}{3} \Delta^2 f(-1) + \frac{(x+2)^{(5)}}{5} \Delta^4 f(-1) + \dots \right]$$

If $(1-x) = t$ in the 1st part of the formula, then we get

$$f(x) = \left[tf(0) + \frac{t(t^2-1^2)}{|3|} \Delta^2 f(-1) + \frac{t(t^2-1^2)(t^2-2^2)}{|5|} \Delta^4 f(-2) + \dots \right] \\ + \left[xf(1) + \frac{x(x^2-1^2)}{|5|} \Delta^2 f(0) + \frac{x(x^2-1^2)(x^2-2^2)}{|5|} \Delta^4 f(1) + \dots \right]$$

which is the Everett's formula.

Problem 43. Given $u_0, u_1, u_2, u_3, u_4, u_5$ (fifth differences constant), prove that $u_{2\frac{1}{2}} = \frac{1}{2}c + \frac{25(c-b)+3(a-c)}{256}$ by any central differences formula, where $a=u_0+u_5, b=u_1+u_4, c=u_2+u_3$.

Bessel's formula is

$$u_{(t+\frac{1}{2})} = \frac{1}{2} [u_0 + u_1] + t \Delta u_0 + \frac{t^2 - (\frac{1}{2})^2}{|2|} \frac{\Delta^2 u_0 + \Delta^2 u_{-1}}{2} + \frac{t \{t^2 - (\frac{1}{2})^2\}}{|3|} \Delta^3 u_{-\frac{1}{2}} \\ + \frac{\{t^2 - (\frac{1}{2})^2\} \{t^2 - (\frac{3}{2})^2\}}{|4|} \frac{\Delta^4 u_{-1} + \Delta^4 u_{-2}}{2} \\ + \frac{t \{t^2 - (\frac{1}{2})^2\} \{t^2 - (\frac{3}{2})^2\}}{|5|} \Delta^5 u_{-1}$$

{fifth differences constant means that differences above fifth are zero}.

Putting $t=0$ in the formula

$$u_{\frac{1}{2}} = \frac{1}{2} [u_0 + u_1] - \frac{1}{8} \frac{\Delta^2 u_0 + \Delta^2 u_{-1}}{2} + \frac{(-\frac{1}{2})(-\frac{9}{4})}{24} \frac{\Delta^4 u_{-1} + \Delta^4 u_{-2}}{2} \\ = \frac{1}{2} (u_0 + u_1) - \frac{1}{8} \frac{\Delta^2 u_0 + \Delta^2 u_{-1}}{2} + \frac{3}{128} \frac{\Delta^4 u_{-1} + \Delta^4 u_{-2}}{2} \\ = \frac{1}{2} (u_0 + u_1) - \frac{1}{128} \{u_2 - 2u_1 + u_0 + u_{-1} - 2u_0 + u_{-1}\} \\ + \frac{3}{8192} \{u_3 - 4u_2 + 6u_1 - 4u_0 + u_{-1} + u_{-2} - 4u_{-1} + 6u_0 - 4u_{-1} + u_{-3}\} \\ = \frac{1}{2} (u_0 + u_1) - \frac{1}{128} \{u_2 - u_1 - u_0 + u_{-1} + \frac{3}{8192} \{u_3 - 3u_2 + 2u_1 + 2u_0 - 3u_{-1} + u_{-3}\}\}$$

$$\text{or } u_{\frac{1}{2}} = \frac{1}{2}(u_0 + u_1) + \frac{1}{8192} \{3(u_3 + u_{-2}) - 9(u_2 + u_{-1}) + 6(u_1 + u_0) + 16(u_0 + u_1) - 16(u_2 + u_{-1})\}$$

Shifting the origin to -2 , we get

$$u_{2\frac{1}{2}} = \frac{1}{2} (u_2 + u_3) + \frac{1}{8192} \{3(u_5 + u_0) - 9(u_4 + u_1) + 6(u_3 + u_2) + 16(u_2 + u_3) + 6(u_4 + u_1)\}$$

$$\text{or } u_{2\frac{1}{2}} = \frac{1}{2}c + \frac{1}{8192} \{3a - 9b + 6c + 16c - 16b\}$$

$$\text{or } u_{2\frac{1}{2}} = \frac{1}{2}c + \frac{1}{8192} \{3a + 22c - 25b\}$$

$$\text{or } u_{2\frac{1}{2}} = \frac{1}{2}c + \frac{25(c-b)+3(a-c)}{256}$$

11. APPROXIMATE INTEGRATION

The approximate integration or numerical quadrature deals with the evaluation of approximate area under a curve by determining the value of a definite integral with the help of a given set of numerical values of the function under integration. For this purpose we first approximate the function into the form of a polynomial applying some formula of interpolation. The degree of the polynomial depends upon the number of known values of the function. With n known values the function can be supposed as a polynomial of degree $(n-1)$. Now approximating the form of the function, we can integrate it within desired limits.

(1) Trapezoidal Rule

Suppose there are two values of the function $f(x)$, $f(0)$ and $f(1)$, represented by H and K on the curve. There can be unlimited curves passing through these two points; the simplest of them will be one degree polynomial, i.e. a straight line, say $f(x) = A + Bx$.

As such area under the curve between $(0, f(0))$, $(1, f(1))$ and the axis of x is




Fig. 7.1

$$\int_0^1 f(x) dx = \int_0^1 (A + Bx) dx \\ = \left[Ax + \frac{Bx^2}{2} \right]_0^1 = A + \frac{B}{2} \quad \dots (1)$$

$$\text{Suppose } \int_0^1 f(x) dx = lf(0) + mf(1) \quad \dots (2) \\ = l(A) + m(A+B) \quad [\text{by } f(x) = A + Bx] \\ = (1+m)A + mB. \quad \dots (3)$$

Comparing (1) and (3), we get

$$l + m = 1, m = \frac{1}{2}, \\ l = \frac{1}{2}.$$

Substituting these values of l and m in (2), we have

$$\int_0^1 f(x) dx = \frac{1}{2} [f(0) + f(1)].$$

(2) Effect of changing Origin and Scale upon Trapezoidal Rule.

Suppose $x = \frac{z-a}{h}$ {where a and h are constant and z is a new variate}.

$$\text{So that } z = a + xh, \text{ giving } dz = h dx \text{ or } dx = \frac{dz}{h}.$$

When $x=0$, $z=a$ and when $x=1$, $z=a+h$.

Also we have $\int_0^1 f(x) dx = \frac{1}{2} [f(0) + f(1)]$.

$$\therefore \int_a^{a+h} f(z) \frac{dz}{h} = \frac{1}{2} [f(a) + f(a+h)]$$

$$\text{or } \int_a^{a+h} f(z) dz = \frac{h}{2} [f(a) + f(a+h)] = \int_a^{a+h} f(x) dx.$$

[3] Extension of Trapezoidal Rule.

We have by taking n intervals each of length h ,

$$\begin{aligned} \int_a^{a+nh} f(x) dx &= \int_a^{a+h} f(x) dx + \int_{a+h}^{a+2h} f(x) dx + \int_{a+2h}^{a+3h} f(x) dx + \dots + \int_{a+(n-1)h}^{a+nh} f(x) dx \\ &= h \left[\frac{1}{2} \{f(a) + f(a+nh)\} + \{f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)\} \right] \\ &= \text{distance between two consecutive ordinates} \\ &\quad \times [\text{mean of the first and the last ordinates} \\ &\quad + \text{sum of all the intermediate terms}] \end{aligned}$$




Fig. 7.2

$$f(x) = A + Bx + Cx^2.$$

$$\therefore f(0) = A,$$

$$f(1) = A + B + C$$

$$\text{and } f(2) = A + 2B + 4C.$$

The area between the curve HKL and the axis of X is

$$\begin{aligned} \int_0^2 f(x) dx &= \int_0^2 (A + Bx + Cx^2) dx \\ &= \left[Ax + \frac{1}{2} Bx^2 + \frac{1}{3} Cx^3 \right]_0^2 = 2A + 2B + \frac{8}{3}C \end{aligned} \quad \dots (7)$$

$$\text{Suppose } \int_0^2 f(x) dx = lf(0) + mf(1) + nf(2)$$

$$\begin{aligned} &= lA + m(A + B + C) + n(A + 2B + 4C) \\ &\quad [\text{by (4), (5) and (6)}] \\ &= A(l + m + n) + B(m + 2n) + C(m + 4n) \end{aligned} \quad \dots (8)$$

Comparing (7) and (8), we get

$$\left. \begin{aligned} l + m + n &= 2 \\ m + 2n &= 2 \\ m + 4n &= \frac{8}{3} \end{aligned} \right\}, \therefore l = \frac{1}{3}, m = \frac{4}{3}, n = \frac{1}{3}.$$

$$\text{So that } \int_0^2 f(x) dx = \frac{1}{3} f(0) + \frac{4}{3} f(1) + \frac{1}{3} f(2)$$

$$\text{i.e. } \int_0^2 f(x) dx = \frac{1}{3} \{f(0) + 4f(1) + f(2)\} \quad \dots (9)$$

which is Simpson's one-third rule.

$$\begin{aligned} \text{After. } \int_0^2 f(x) dx &= \int_0^2 \{f(0) + {}^x C_1 \Delta f(0) + {}^x C_2 \Delta^2 f(0)\} dx \\ &= \int_0^2 [f(0) + x\{\Delta f(0) - \frac{1}{2} \Delta^2 f(0)\} + \frac{x^2}{2} \Delta^2 f(0)] dx \\ &= \left[x f(0) + \frac{x^2}{2} \{\Delta f(0) - \frac{1}{2} \Delta^2 f(0)\} + \frac{x^3}{6} \Delta^2 f(0) \right]_0^2 \\ &= 2f(0) + 2\Delta f(0) + \frac{1}{3} \Delta^2 f(0) \\ &= 2f(0) + 2\{f(1) - f(0)\} + \frac{1}{3} \{f(2) - 2f(1) + f(0)\} \end{aligned}$$

$$\text{or } \int_0^2 f(x) dx = \frac{1}{3} \{f(0) + 4f(1) + f(2)\}.$$

[5] Effect of Changing Unit and Origin on Simpson's $\frac{1}{3}$ Rule.

$$\text{Suppose } x = \frac{z-a}{h}; \therefore dx = \frac{dz}{h}.$$

$$\text{Thus } \int_0^2 f(x) dx = \frac{1}{3} \{f(0) + 4f(1) + f(2)\}, \text{ changes to}$$

$$\int_a^{a+2h} f(z) dz = \frac{h}{3} \{f(a) + 4f(a+h) + f(a+2h)\}.$$

$$\text{In particular. } \int_{-1}^1 f(x) dx = \frac{1}{3} \{f(-1) + 4f(0) + f(1)\} \quad \dots (10)$$

Also when $a=0$ this yields

$$\int_0^{2h} f(z) dz = \frac{h}{3} \{f(0) + 4f(h) + f(2h)\} \quad \dots (11)$$

[6] Extension of Simpson's $\frac{1}{3}$ rule (for all even values of n).

$$\begin{aligned} \text{We have } \int_a^{a+nh} f(x) dx &= \int_a^{a+2h} f(x) dx + \int_{a+2h}^{a+4h} f(x) dx + \dots \\ &\quad + \int_{a+(n-2)h}^{a+nh} f(x) dx \end{aligned}$$

$$\begin{aligned} &= \frac{h}{3} [\{f(a) + 4f(a+h) + f(a+2h)\} + \{f(a+2h) + 4f(a+3h) + f(a+4h)\} \\ &\quad + \dots + \{f(a+(n-2)h) + 4f(a+(n-1)h) + f(a+nh)\}] \end{aligned}$$

$$= \frac{h}{3} [f(a) + f(a+nh) + 4\{f(a+h) + f(a+3h) + \dots + f(a+(n-1)h)\}]$$

$$+2 \{ f(a+2h) + f(a+4h) + \dots + f(a+n-2h) \} \\ = (\text{one-third of the interval}) \{ (\text{sum of the 1st and the last terms}) \\ + 4 (\text{sum of the odd number of terms}) \\ + 2 (\text{sum of the even number of terms}) \} \quad (12)$$

[7] Simpson's $\frac{3}{2}$ th rule.

Consider four points $\{0, f(0)\}$, $\{1, f(1)\}$, $\{2, 2f(2)\}$ and $\{3, f(3)\}$ of instead three as taken in [4]

$$\text{Then Required area} = \int_0^3 f(x) dx. \quad \dots (13)$$

For the sake of convenience let us double the unit of interval and then shift the origin to 3. By this transformation we have the four points $\{-3, f(-3)\}$, $\{-1, f(-1)\}$, $\{1, f(1)\}$ and $\{3, f(3)\}$ as given and so we have to find out the value of the integral $\int_{-3}^3 f(x) dx$, where $f(x) = A + Bx + Cx^2 + Dx^3$, a third degree polynomial with four given values.

$$\text{Suppose } \int_{-3}^3 f(x) dx = m \{ f(-3) + f(3) \} + n \{ f(-1) + f(1) \}$$

$$\text{or } \int_{-3}^3 \{ A + Bx + Cx^2 + Dx^3 \} dx = m \{ 2A + 18C \} + n \{ 2A + 2C \} \\ \{ \because f(-3) = A - 3B + 9C - 27D \text{ etc.} \}$$

$$\text{or } \left[Ax + \frac{Bx^2}{2} + \frac{Cx^3}{3} + \frac{Dx^4}{4} \right]_{-3}^3 = A (2m + 2n) + C (18m + 2n)$$

$$\text{or } 6A + 18C = (2m + 2n) A + (18m + 2n) C.$$

$$\therefore (2m + 2n) = 6 \text{ and } 18m + 2n = 18,$$

$$\text{giving } 16m = 12 \text{ or } m = \frac{3}{4}$$

$$\text{and } 2n = 6 - 2m = 6 - \frac{3}{2} = \frac{9}{2} \text{ or } n = \frac{9}{4}.$$

Substituting these values of m and n in (13) we get

$$\int_{-3}^3 f(x) dx = \frac{3}{4} \{ f(-3) + f(3) \} + \frac{9}{4} \{ f(-1) + f(1) \}$$

$$\text{or } \int_{-3}^3 f(x) dx = \frac{3}{8} [2 \{ f(-3) + f(3) \} + 6 \{ f(-1) + f(1) \}] \quad \dots (14)$$

which is the form of the Simpson's $\frac{3}{2}$ th rule.

Shifting the origin to -3 , we get

$$\int_0^6 f(x) dx = \frac{3}{8} [2 \{ f(0) + f(6) \} + 6 \{ f(2) + f(4) \}] \quad \dots (15)$$

which is another form of the formula.

Making the interval half of the present one which is 2, i.e., making the interval unity, we have

$$\int_0^3 f(x) dx = \frac{3}{8} [\{ f(0) + f(3) \} + 3 \{ f(1) + f(2) \}] \quad \dots (16)$$

Note. Like the $\frac{1}{2}$ rd rule, the $\frac{3}{2}$ th rule, can also be obtained by substituting the integral.

$$\int_a^b f(x) dx = \int_0^3 E^x f(0) dx = \int_0^3 (1 + \Delta)^x f(0) dx \\ = \int_0^3 \left\{ 1 + x\Delta + \frac{x(x-1)}{2} \Delta^2 + \frac{x(x-1)(x-2)}{6} \Delta^3 \right\} f(0) dx \\ = \int_0^3 \left\{ 1 + x\Delta + \left(\frac{x^2}{2} - \frac{x}{2} \right) \Delta^2 + \left(\frac{x^3}{6} - \frac{x^2}{2} + \frac{x}{3} \right) \Delta^3 \right\} f(0) dx \\ = \left[x + \frac{x^2}{2} \Delta + \left(\frac{x^3}{6} - \frac{x^2}{4} \right) \Delta^2 + \left(\frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{6} \right) \Delta^3 \right]_0^3 f(0) \\ = 3f(0) + \frac{9}{2} \Delta f(0) + \left(\frac{9}{2} - \frac{9}{4} \right) \Delta^2 f(0) + \left(\frac{27}{8} - \frac{9}{2} + \frac{9}{6} \right) \Delta^3 f(0) \\ = 3f(0) + \frac{9}{2} \{ f(1) - f(0) \} + \frac{9}{4} \{ f(2) - 2f(1) + f(0) \} \\ + \frac{9}{24} \{ f(3) - 3f(2) + 3f(1) - f(0) \} \\ = \left(3 - \frac{9}{2} + \frac{9}{4} - \frac{9}{24} \right) f(0) + \left\{ \frac{9}{2} - \frac{9}{2} + \frac{9}{8} \right\} f(1) + \left\{ \frac{9}{4} - \frac{9}{8} \right\} \\ (2) + \frac{3}{8} f(3) \\ = \frac{3}{8} [\{ f(0) + f(3) \} + 3 \{ f(1) + f(2) \}].$$

[8] Effect of Changing Origin and Scale Upon $\frac{3}{2}$ th Rule.

$$\text{We have } \int_0^3 f(x) dx = \frac{3}{8} [\{ f(0) + f(3) \} + 3 \{ f(1) + f(2) \}].$$

$$\text{Suppose } \frac{z-a}{h} = x, \text{ so that } dz = h dx$$

$$\int_a^{a+3h} f(z) dz = \frac{3h}{8} [\{ f(a) + f(a+3h) \} + 3 \{ f(a+h) + f(a+2h) \}] \quad \dots (17)$$

[9] Extension of Simpson's $\frac{3}{2}$ th rule. (When n is a multiple of 3.)

$$\int_a^{a+n h} f(x) dx = \int_a^{a+3h} f(x) dx + \int_{a+3h}^{a+6h} f(x) dx + \int_{a+6h}^{a+9h} f(x) dx + \dots \\ + \int_{a+(n-3)h}^{a+n h} f(x) dx$$

$$= \frac{3h}{8} [\{ f(a) + f(a+n h) \} + 2 \{ f(a+3h) + f(a+6h) + \dots + f(a+(n-3)h) \} \\ + 3 \{ f(a+h) + f(a+2h) + f(a+4h) + f(a+5h) + \dots \}] \quad \dots (18)$$

$$= \frac{3h}{8} [(\text{sum of the first and last term})$$

$$+ 2 (\text{sum of 4th, 7th, 10th etc. terms}) + 3 \{ \text{sum of the remaining} \\ \text{2nd, 3rd, 5th, 6th, 8th, 9th, etc. terms} \}]$$

[10] Weddle's Rule.

If $f(x)$ is given for certain equidistant values of x say $x_0, x_0+h, x_0+2h, \dots$ and the range (a, b) is divided into n equal parts each of width h i.e. $b-a=nh$ and if $x_0=a, x_1=x_0+h, \dots, x_n=a+nh=b$ assuming that $(n+1)$ ordinates $y_0, y_1, \dots, y_n$ are equidistant then by

change of scale, $u = \frac{x-x_0}{h}, dx=hdu$, the general quadrature formula is

$$\int_a^b y dx = \int_{x_0}^{x_0+nh} y_x dx = \int_0^n y_{x_0+x_u} hdu = h \int_0^n \left[y_0 + u \Delta y_0 + \frac{u(u-1)}{2} \Delta^2 y_0 + \dots \right] du$$

$$= h \left[ny_0 + \frac{n^2}{2} \Delta y_0 + \left(\frac{n^3}{3} - \frac{n^2}{2} \right) \frac{\Delta^2 y_0}{2} + \left(\frac{n^4}{4} - n^3 + n^2 \right) \frac{\Delta^3 y_0}{6} + \dots \text{upto } (n+1) \text{ terms} \right] \dots (19)$$

Putting $n=6$, and neglecting all differences above the sixth, we find

$$\int_{x_0}^{x_0+6h} y dx = h \left[6y_0 + 18\Delta y_0 + 27\Delta^2 y_0 + 24\Delta^3 y_0 + \frac{123}{10}\Delta^4 y_0 + \frac{33}{10}\Delta^5 y_0 + \frac{41}{140}\Delta^6 y_0 \right]$$

Replacing the last term by $\frac{3}{10}\Delta^6 y_0$ and neglecting the error

$$\Delta^6 y_0 \text{ made, we have}$$

$$\int_{x_0}^{x_0+6h} y dx = \frac{3h}{10} \left[y_0 + 5y_1 + y_2 + 6y_3 + y_4 + 5y_5 + y_6 \right]$$

$$\text{Similarly, } \int_{x_0+6h}^{x_0+12h} y dx = \frac{3h}{10} \left[y_6 + 5y_7 + y_8 + 6y_9 + y_{10} + 5y_{11} + y_{12} \right]$$

$$\dots \dots \dots$$

$$\int_{x_0+(n-6)h}^{x_0+nh} y dx = \frac{3h}{10} \left[y_{n-6} + 5y_{n-5} + y_{n-4} + 6y_{n-3} + y_{n-2} + 5y_{n-1} + y_n \right], n \text{ being multiple of 6}$$

$$\text{Adding altogether, } \int_{x_0}^{x_0+nh} y dx = \frac{3h}{10} \left[y_0 + 5y_1 + y_2 + 6y_3 + y_4 + 5y_5 + 2y_6 + 5y_7 + y_8 + \dots \right] \dots (20)$$

which is Weddle's rule.

Note. The trapezoidal rule, Simpson's $\frac{1}{3}$ rd rule and Simpson's $\frac{1}{4}$ th rule can successively be obtained by putting $n=1, n=2$ and $n=3$ respectively in (19).

Problem 44. Evaluate the following:

$$(i) \int_0^1 \frac{x^2}{1+x^2} dx \text{ and hence find the value of } \log_e 2.$$

$$(ii) \int_2^{10} \frac{dx}{1+x} \text{ by dividing the range into eight equal parts.}$$

$$(i) \text{ We have } \int_0^1 f(x) dx = \frac{1}{2} \cdot \frac{1}{3} \{ f(0) + 4f(\frac{1}{2}) + f(1) \}. \dots (1)$$

$$\therefore \int_0^1 \frac{x^2}{1+x^3} dx = \frac{1}{2} \cdot \frac{1}{3} \left\{ \frac{0}{1+0} + 4 \cdot \frac{(\frac{1}{2})^2}{1+(\frac{1}{2})^3} + \frac{1^2}{1+1^3} \right\}$$

$$= \frac{1}{6} \left\{ 0 + \frac{8}{5} + \frac{1}{2} \right\} = \frac{2.5}{6}. \dots (2)$$

Now putting $x^3=t$ in the given integral we get

$$\int_0^1 \frac{x^2}{1+x^3} dx = \frac{1}{3} \int_0^1 \frac{dt}{1+t} = \frac{1}{3} \left\{ \log_e(1+t) \right\}_0^1 = \frac{1}{3} \log_e 2 \dots (3)$$

$\therefore$ from (2) and (3), $\log_e 2 = 3 \times \frac{2.5}{6} = \frac{2.5}{2} = .6944$.

(ii) Dividing the range into eight parts by means of 9 points of the x -values as 2, 3, 4, 5, 6, 7, 8, 9, 10, the extended form of Simpson's $\frac{1}{3}$ rd rule gives.

$$\int_2^{10} f(x) dx = \frac{1}{3} \{ \{ f(2) + f(10) \} + 2 \{ f(4) + f(6) + f(8) \} + 4 \{ f(3) + f(5) + f(7) + f(9) \} \} \dots (4)$$

$$\text{But } f(x) = \frac{1}{1+x}.$$

$$\therefore f(2) = .333, f(3) = .25, f(4) = .20, f(5) = .166, f(6) = .143, f(7) = .125, f(8) = .111, f(9) = .10, f(10) = .091.$$

Substituting these values in (4) we get

$$\int_2^{10} f(x) dx = \frac{1}{3} \{ \{ .333 + .091 \} + 2 \{ .20 + .143 + .111 \} + 4 \{ .25 + .166 + .125 + .10 \} \}$$

$$= \frac{1}{3} [.422 + .908 + 2.564] = 1.298.$$

$$\text{Hence } \int_2^{10} \frac{dx}{1+x} = 1.298.$$

Problem 45. If u_x is a function whose fifth differences are constant,

$\int u_x dx$ can be expressed in the form $pu_{-\alpha} + qu_0 + pu_{\alpha}$. Find the values of p, q and α .

Use this formula, after making necessary changes in the origin and scale, to find the value of $\log_e 2$ to four places of decimals from the equation $\int_0^1 \frac{I}{1+x} dx = \log_e 2$.

(i) As fifth differences of the function are constant, so u_x must be of fifth degree such as

$$u_x = a + bx + cx^2 + dx^3 + ex^4 + fx^5.$$

$$\therefore \int_{-1}^1 u_x dx = \int_{-1}^1 \{ a + bx + cx^2 + dx^3 + ex^4 + fx^5 \} dx$$

$$= 2 \left\{ a + \frac{c}{3} + \frac{e}{5} \right\}. \dots (1)$$

$$\text{Now } pu_{-\alpha} + qu_0 + pu_{\alpha} = 2p \{ a + \alpha^2 c + \alpha^4 e \} + qa. \dots (2)$$

But $\int_{-1}^1 u_x dx = pu_{-1} + qu_0 + pu_1.$

Now equating the coefficients of a , c and e in (1) and (2), we get
 $2=2p+q...$ (3), $\frac{2}{3}=2p\alpha^2...$ (4) and $\frac{2}{3}=2p\alpha^4$... (5)

$$\therefore \alpha^2 = \frac{2p\alpha^4}{2p\alpha^2} = \frac{\frac{2}{3}}{\frac{2}{3}} = \frac{2}{3}.$$

Substituting this value of α^2 in (4), we get $p = \frac{3}{8}.$

And then substituting this value of p in (3), we get $q = \frac{3}{8}.$

Thus $\int_{-1}^1 u_x dx = \frac{3}{8} \{5(u_{-1} + u_1) + 8u_0\}$, where $\alpha^2 = \frac{2}{3}.$

(ii) $\therefore \int_{-1}^1 u_x dx = \frac{3}{8} \{5(u_{-1} + u_1) + 8u_0\}.$

$\therefore \int_0^2 u_x dx = \frac{3}{8} \{5(u_{1+\alpha} + u_{1-\alpha}) + 8u_1\}$ (shifting the origin to -1)

$\therefore \int_0^2 u_x dx = \frac{3}{8} \{5 u_{(1+\alpha)/2} + u_{(1-\alpha)/2} + 8u_{1/2}\}.$... (6)

(Dividing the unit by 2)

$$\therefore u_x = \frac{1}{1+x}, \therefore u_{1/2} = \frac{1}{1+\frac{1}{2}} = \frac{2}{3}$$

and $u_{(1+\alpha)/2} + u_{(1-\alpha)/2} = \frac{1}{1+\frac{1+\alpha}{2}} + \frac{1}{1+\frac{1-\alpha}{2}} = \frac{12}{9-\alpha^2} = \frac{12}{9-\frac{2}{3}} = \frac{12}{\frac{25}{3}} = \frac{36}{25}$

Substituting these values in (6), we get

$$\int_0^2 \frac{1}{1+x} dx = \frac{3}{8} \left\{ 5 \times \frac{36}{25} + \frac{8 \times 2}{3} \right\} = \frac{131}{189} = .6931.$$

But $\int_0^2 \frac{1}{1+x} dx = \log_e 2$, $\therefore \log_e 2 = .6931.$

Problem 46. Calculate $\int_0^{\pi/2} \sin x dx$ by

(i) trapezoidal rule

(ii) Simpson's rule (using 11 ordinates)

Dividing $\left(0, \frac{\pi}{2}\right)$ into 10 equal parts so that $h = \frac{\pi}{20}$ we have from trigonometric tables.

$$y_1 = \sin 0 = 0.0000 \quad y_7 = \sin \frac{3\pi}{10} = 0.8090$$

$$y_2 = \sin \frac{\pi}{20} = 0.1564 \quad y_8 = \sin \frac{7\pi}{20} = 0.8910$$

$$y_3 = \sin \frac{\pi}{10} = 0.3090 \quad y_9 = \sin \frac{2\pi}{5} = 0.9511$$

$$y_4 = \sin \frac{3\pi}{10} = 0.4540 \quad y_{10} = \sin \frac{9\pi}{20} = 0.9877$$

$$y_1 = \sin \frac{\pi}{5} = 0.5878$$

$$y_{11} = \sin \frac{\pi}{2} = 1.0000$$

$$y_4 = \sin \frac{\pi}{4} = 0.7071$$

(i) Required value of the integral

$$= h \left\{ \frac{1}{2} (y_1 + y_{11}) + (y_2 + y_3 + \dots + y_{10}) \right\}$$

$$= \frac{\pi}{20} \left\{ 0.5 + 5.8531 \right\} = 0.9981.$$

(ii) Required value of the integral

$$= \frac{h}{3} [y_1 + y_{11} + 2(y_2 + y_4 + y_6 + y_8 + y_{10}) + 4(y_3 + y_5 + y_7 + y_9)]$$

$$= \frac{\pi}{60} [1 + 5.3138 + 12.7848] = 19.0986 \times 0.0524 = 1.0006.$$

Problem 47. Use Weddle's rule to find the value of $\int_4^{5.2} \log_e x dx$

Total interval = $5.2 - 4 = 1.2.$

Dividing 1.2 into six equal parts, $h = 0.2.$

By Table we have

x	4.0	4.2	4.4	4.6	4.8	5.0	5.2
$\log_e x$	1.3863	1.4351	1.4816	1.5261	1.5686	1.6094	1.6487

Required value of the integral

$$= (0.2) \times \frac{3}{10} \left[1.3863 + 5(1.4351) + 1.4816 + 6(1.5261) \right.$$

$$\left. + 1.5686 + 5(1.6094) + 2(1.6487) \right]$$

$$= 1.8278 \text{ approx.}$$

[11] Picard's Method

Consider $\frac{dy}{dx} = f(x, y)$... (21)

Suppose it is required to find its solution under the condition $y = y_0$ when $x = x_0.$

Integration of (1) gives $y = \int f(x, y) dx + C$

When $x = x_0, y = y_0 \quad \therefore C = y_0 - \int f(x_0, y) dx$

So that $y = y_0 + \int_{x_0}^x f(x, y) dx - \int_{x_0}^x f(x_0, y) dx$

$$= y_0 + \int_{x_0}^x f(x, y) dx \quad \dots (22)$$

Now (22) is integrated by successive approximation.

The first approximation is found by putting $y = y_0$ in the integrand of (22), whence we get

$$y_1 = y_0 + \int_{x_0}^x f(x, y_0) dx$$

The second approximation is obtained by replacing y by y_1 in R.H.S. of (22)

$$i.e. \quad y_2 = y_0 + \int_{x_0}^x f(x, y_1) dx$$

and so on.

In general the n th approximation is

$$y_n = y_0 + \int_{x_0}^x f(x, y_{n-1}) dx.$$

Problem 48. Solve $\frac{dy}{dx} = x + y^2$ where $y=0$ when $x=0$.

Here $x_0=0$, $y_0=0$ and $f(x, y) = x + y^2$

$\therefore$ First approximation is $y_1 = y_0 + \int_{x_0}^x f(x, y_0) dx = 0 + \int_0^x x dx = \frac{x^2}{2}$
 $\therefore f(x, y_0) = x + 0^2 = x$

Second approximation is

$$y_2 = y_0 + \int_{x_0}^x f(x, y_1) dx, \quad f(x, y_1) = x + \left(\frac{x^2}{2}\right)^2 = x + \frac{x^4}{4}$$

$$= 0 + \int_0^x \left(x + \frac{x^4}{4}\right) dx = \frac{x^2}{2} + \frac{x^5}{20}$$

Third approximation is

$$y_3 = y_0 + \int_{x_0}^x f(x, y_2) dx, \quad f(x, y_2) = x + \left(\frac{x^2}{2} + \frac{x^5}{20}\right)^2$$

$$= x + \frac{x^4}{4} + \frac{x^7}{20} + \frac{x^{10}}{400}$$

$$= 0 + \int_0^x \left(x + \frac{x^4}{4} + \frac{x^7}{20} + \frac{x^{10}}{400}\right) dx$$

$$= \frac{x^2}{2} + \frac{x^5}{20} + \frac{x^8}{160} + \frac{x^{11}}{4400}$$

Which gives fairly good approximation.

[12] Extension of Picard's Method.

Consider simultaneous equations

$$\frac{dy}{dx} = f(x, y, z), \quad \frac{dz}{dx} = g(x, y, z) \quad \dots (23)$$

subject to the conditions $y=y_0$, $z=z_0$ when $x=x_0$

The first approximations of (23) are

$$y_1 = y_0 + \int_{x_0}^x f(x, y_0, z_0) dx, \quad z_1 = z_0 + \int_{x_0}^x g(x, y_0, z_0) dx \quad \dots (24)$$

In general the n th approximations are

$$y_n = y_0 + \int_{x_0}^x f(x, y_{n-1}, z_{n-1}) dx, \quad z_n = z_0 + \int_{x_0}^x g(x, y_{n-1}, z_{n-1}) dx$$

Problem 49. Solve $\frac{dy}{dx} = x + z$, $\frac{dz}{dx} = x - y^2$ where $y=2$, $z=1$ when $x=0$.

Here $x_0=0$, $y_0=2$, $z_0=1$, $f(x, y, z) = x + z$ and $g(x, y, z) = x - y^2$

First approximations are $y_1 = 2 + \int_0^x (x+1) dx = 2 + x + \frac{x^2}{2}$

$$z_1 = 1 + \int_0^x (x-4) dx = 1 - 4x + \frac{x^2}{2}$$

Similarly second and third approximations can be evaluated.

[13] Runge-Kutta Method.

Consider $\frac{dy}{dx} = f(x, y)$, $y=y_0$ when $x=x_0$

Assume that $y=F(x)$. Then Taylor's theorem yields

$$F(x_0+h) = F(x_0) + h F'(x_0) + \frac{h^2}{2} F''(x_0) + \frac{h^3}{6} F'''(x_0) + \dots \quad (25)$$

Here $F'(x) = \frac{dy}{dx} = f(x, y) = f(\text{say})$.

Take $p = \frac{\partial f}{\partial x}$, $q = \frac{\partial f}{\partial y}$, $r = \frac{\partial^2 f}{\partial y^2}$, $s = \frac{\partial^2 f}{\partial x \partial y}$, $t = \frac{\partial^2 f}{\partial y^2}$

At $y=y_0$ when $x=x_0$, let $p=p_0$, $q=q_0$, $r=r_0$, $s=s_0$, $t=t_0$.

$$\text{Now } F''(x) = \frac{df}{dx} = \left(\frac{\partial}{\partial x} + \frac{dy}{dx} \frac{\partial}{\partial y} \right) f = p + fq$$

$$\therefore df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$

$$F'''(x) = \frac{d^2 f}{dx^2} = \frac{d}{dx} \left(\frac{df}{dx} \right) = \left(\frac{\partial}{\partial x} + \frac{dy}{dx} \frac{\partial}{\partial y} \right) (p + fq)$$

$$= \frac{\partial p}{\partial x} + q \frac{\partial f}{\partial x} + f \frac{\partial q}{\partial x} + \frac{dy}{dx} \frac{\partial p}{\partial y} + \frac{dy}{dx} q \frac{\partial f}{\partial y} + \frac{dy}{dx} f \frac{\partial q}{\partial y}$$

$$= r + pq + fs + f(s + q^2 + ft)$$

$$\text{Hence } F(x_0+h) - F(x_0) = hf_0 + \frac{h^2}{2} (p_0 + f_0 q_0) + \frac{h^3}{6} (r_0 + 2f_0 s_0 + f_0^2 t_0 + p_0 q_0 + f_0 q_0^2) + \dots \quad (26)$$

Here the first term represents the first approximation i.e.,

$$F(x_0+h) - F(x_0) = hf_0 \text{ i.e. } y = y_0 + hf(x_0, y_0) \quad \dots (27)$$

The second approximation may be taken as

$$y - y_0 = hf \left(x_0 + \frac{h}{2}, y_0 + \frac{1}{2} hf_0 \right) = k_1 \text{ (say)} \quad \dots (28)$$

whence by Taylor's expansion, we have

$$f \left(x_0 + \frac{h}{2}, y_0 + \frac{1}{2} hf_0 \right) = f_0 + \frac{1}{2} h p_0 + \frac{1}{2} h f_0 q_0 + \frac{1}{2} \left(\frac{1}{4} h^2 r_0 + \frac{1}{2} h^2 f_0 s_0 + \frac{1}{4} h^2 f_0^2 t_0 \right)$$

$$\text{giving } k_1 = hf_0 + \frac{1}{2} h^2 (p_0 + f_0 q_0) + \frac{1}{8} h^3 (r_0 + 2f_0 s_0 + f_0^2 t_0) \quad \dots (29)$$

From (26) it is observed that the difference in the two values of k found from (28) and (29) is in the coefficient of h^3 .

In order to find the extra terms in h^3 , Runge-Kutta replaces

$$hf(x_0+h, y_0+hf_0) \text{ by } k''' = hf(x_0+h, y_0+k'') \text{ where } k'' = hf(x_0+h, y_0+hf_0)$$

The modified formula therefore becomes $\frac{1}{3}\{k' + 4k_1 + k'''\}$ where $k' = hf_0 = \frac{2}{3}k_1 + \frac{1}{3}k_2 = k_1 + \frac{1}{3}(k_2 - k_1)$ and $k_2 = \frac{1}{2}(k' + k''')$.

Hence this method is applied in the sequence:

$$k' hf_0; k'' = hf(x_0+h, y_0+k'); k''' = hf(x_0+h, y_0+k'')$$

$$k_1 = hf\left(x_0 + \frac{h}{2}, y_0 + \frac{1}{2}k'\right); k_2 = \frac{1}{2}(k' + k''') \text{ and } k = k_1 + \frac{1}{3}(k_2 - k_1)$$

Problem 50. Apply Runge-Kutta method to find y when $x=0.3$ from $\frac{dy}{dx} = x + y^2$ where $y=0$ when $x=0$.

Here $x_0=0, y_0=0, h=0.3, f(x, y)=x+y^2$ and $f_0=f(x_0, y_0)=0$.

$$\therefore k' = hf_0 = 0$$

$$k'' = hf(x_0+h, y_0+k') = 0.3 \times f\{0.3, 0\} = 0.3 \times 0.3 = 0.09$$

$$k''' = hf(x_0+h, y_0+k'') = 0.3 \times f\{0.3, 0.09\} = 0.3 \times \{0.3 + 0.0081\} = 0.0924$$

$$k_1 = hf\left(x_0 + \frac{h}{2}, y_0 + \frac{k'}{2}\right) = 0.3 \times f\{0.15, 0\} = 0.3 \times 0.15 = 0.045$$

$$k_2 = \frac{1}{2}(k' + k''') = \frac{1}{2} \times 0.0924 = 0.0462$$

$$k = k_1 + \frac{1}{3}(k_2 - k_1) = 0.045 + 0.0004 = 0.0454.$$

7.12. PERTURBATION METHOD

[A] First Order Differential Equations.

When dealing with physical problems, we frequently encounter a differential equation of the type

$$\frac{dy}{dx} + y^2 = 0 \text{ with } y(1) = 1 \text{ (say)} \quad \dots (1)$$

which has been disturbed by a small effect. We thus modify (1) as

$$\frac{dy}{dx} + y^2 = \epsilon x, y(1) = 1 \quad \dots (2)$$

where ϵ is arbitrarily small.

We have now to determine by how much the solution of (1) has been changed on account of the presence of the disturbing factor ϵx . This change is known as a *Perturbation*.

Suppose $y_0(x)$ is a solution of (1) satisfying $y(1)=1$ and assume the solution of (2) as

$$y(x) = y_0(x) + p(x), p(x) \text{ being perturbation} \quad \dots (3)$$

Expanding $y(x)$ in a series in powers of ϵ , such that

$$y(x) = y_0(x) + \epsilon y_1(x) + \epsilon^2 y_2(x) + \epsilon^3 y_3(x) + \dots \quad \dots (4)$$

and comparing (3) and (4) we find

$$p(x) = \epsilon y_1(x) + \epsilon^2 y_2(x) + \epsilon^3 y_3(x) + \dots \quad \dots (5)$$

Here the first term $\epsilon y_1(x)$ is known as *first order perturbation*, second term $\epsilon^2 y_2(x)$ as *second order perturbation* etc.

With the substitution of (4), (2) yields

$$y_0' + \epsilon y_1' + \epsilon^2 y_2' + \epsilon^3 y_3' + \dots + (y_0 + \epsilon y_1 + \epsilon^2 y_2 + \epsilon^3 y_3 + \dots)^2 = \epsilon x \quad \dots (6)$$

$$\text{or } (y_0' + y_0^2) + (y_1' + 2y_0 y_1) \epsilon + (y_2' + 2y_0 y_2 + y_1^2) \epsilon^2 + \dots = \epsilon x \quad \dots (7)$$

Equating like powers of ϵ on either side, we get

$$y_0' + y_0^2 = 0, y_1' + 2y_0 y_1 = x, y_2' + 2y_0 y_2 + y_1^2 = 0, \text{ etc.} \quad \dots (8)$$

Solving these equations in succession, we may find $y_1(x), y_2(x), \dots$ in (4) and each of these functions satisfies the initial condition $y(1)=1$. But y_0 is a solution of (1) and hence we have

$$y_0(1)=1, y_1(1)=0, y_2(1)=0 \quad \dots (9)$$

Now in order to solve (1), substitute (4) in (2) to give

$$(y_0' + y_0^2) + (y_1' + 2y_0 y_1) \epsilon + (y_2' + 2y_0 y_2 + y_1^2) \epsilon^2 + \dots = \epsilon x \quad \dots (10)$$

Equating coefficients of $\epsilon^0, \epsilon, \epsilon^2$, on either side of (10), we get

$$y_0' + y_0^2 = 0, y_1' + 2y_0 y_1 = x, y_2' + 2y_0 y_2 + y_1^2 = 0 \quad \dots (11)$$

of which first equation gives for $y_0(1)=1, y_0 = \frac{1}{x}$, and then second one gives $y_1' + \frac{2}{x} y_1 = x$ which with $y_1(1)=0$, yields,

$$y_1 = \frac{1}{4} \left(x^2 - \frac{1}{x^2} \right)$$

With these values the third equation of (11) gives

$$y_2' + \frac{2}{x} y_2 = -\frac{1}{16} \left(x^4 - 2 + \frac{1}{x^4} \right)$$

$$\text{which with } y_2(1)=0 \text{ yields } y_2 = -\frac{1}{16} \left(\frac{x^5}{7} - \frac{2x}{3} - \frac{1}{x^3} \right) - \frac{2}{21x^3}.$$

With these substitutions (4) yields

$$y = \frac{1}{x} + \frac{\epsilon}{4} \left(x^2 - \frac{1}{x^2} \right) - \frac{\epsilon^2}{336} \left(3x^5 - 14x - \frac{21}{x^3} + \frac{32}{x^2} \right) \quad \dots (12)$$

This follows that the solution of (1) is $\frac{1}{x}$; but where the disturbing function ϵx is present, the first and second order perturbation terms are respectively the second and third terms in (12).

[B] Second Order Differential Equations.

$$\text{Consider an equation } \frac{d^2 y}{dx^2} + y = 0 \quad \dots (13)$$

$$\text{With } y(0)=0, y'(0)=1.$$

(13) is the differential equation of S.H.M.

Taking the disturbing function as $-2\epsilon(y')^2$, ϵ being small, (13) can be modified as

$$y'' + y = -2\epsilon(y')^2 \quad \dots (15)$$

with $y(0)=0$, $y'(0)=1$.

If $y_0(x)$ is a solution of (13), then suppose that

$$y = y_0 + \epsilon y_1 + \epsilon^2 y_2 + \dots \quad \dots (16)$$

giving $y'' = y_0'' + \epsilon y_1'' + \epsilon^2 y_2'' + \dots$

So that (16) will satisfy (14) under the assumptions

$$y_0(0)=0, y_1(0)=0, y_2(0)=0, \dots; y_0'(0)=1, y_1'(0)=0, y_2'(0)=0, \dots \quad \dots (17)$$

Substituting (16) in (13) and using the terms upto ϵ^2 , we find

$$y_0'' + \epsilon y_1'' + \epsilon^2 y_2'' + y_0 + \epsilon y_1 + \epsilon^2 y_2 = -2\epsilon[(y_0')^2 + \epsilon^2(y_1')^2 + \epsilon^4(y_2')^2 + 2\epsilon y_0' y_1' + 2\epsilon^2 y_0' y_2' + 2\epsilon^3 y_1' y_2'] \quad \dots (18)$$

Collecting the coefficients of like powers of ϵ , we have,

$$(y_0'' + y_0) + (y_1'' + y_1) \epsilon + (y_2'' + y_2) \epsilon^2 = -2(y_0')^2 \epsilon - 4y_0' y_1' \epsilon^2 \quad \dots (19)$$

Equating the coefficients of ϵ^0 , ϵ , ϵ^2 on either side, we find

$$y_0'' + y_0 = 0, y_1'' + y_1 = -2(y_0')^2, y_2'' + y_2 = -4y_0' y_1' \quad \dots (20)$$

of which the first equation in view of $y_0(0)=0$, $y_0'(0)=1$ gives a solution $y_0 = \sin x$, $y_0' = \cos x$ $\dots (21)$

and then the second equation gives $y_1'' + y_1 = -2 \cos^2 x$ $\dots (22)$

So that a general solution of (22) is

$$y_1 = A_1 \sin x + A_2 \cos x - \frac{1}{3} \sin^2 x - \frac{1}{3} \cos^2 x; y_1' = A_1 \cos x - A_2 \sin x - \frac{2}{3} \sin x \cos x \quad \dots (23)$$

$\therefore$ A particular solution of (23) satisfying $y_1(0)=0$, $y_1'(0)=0$, is

$$y_1 = \frac{2}{3} \cos x - \frac{4}{3} \sin^2 x - \frac{2}{3} \cos^2 x; y_1' = -\frac{2}{3} \sin x - \frac{4}{3} \sin x \cos x + \frac{4}{3} \sin x \cos x = -\frac{2}{3} \sin x - \frac{4}{3} \sin x \cos x \quad \dots (24)$$

With these substitutions in the third equation of (20) [i.e., y_0' from (21) and y_1 from (24)], yields

$$y_2'' + y_2 = -4 \left(-\frac{2}{3} \sin x \cos x - \frac{4}{3} \sin x \cos^2 x \right) = \frac{8}{3} \sin x \cos x + \frac{16}{3} \sin x \cos^2 x \quad \dots (25)$$

Its Complementary function is

$$y_c = A_1 \sin x + A_2 \cos x \quad \dots (26)$$

While a particular solution;

$$\left. \begin{aligned} \text{of } y_2'' + y_2 = \frac{8}{3} \sin x \cos x \text{ is } y_2 = -\frac{8}{9} \sin x \cos x \\ \text{of } y_2'' + y_2 = \frac{16}{3} \sin x \cos^2 x \text{ is } y_2 = -\frac{16}{9} \cos^3 x \\ \text{of } y_2'' + y_2 = \frac{16}{3} \sin^3 x \text{ is } y_2 = \frac{16}{9} \sin^3 x - 2x \cos x \end{aligned} \right\} \quad \dots (27)$$

Hence with the help of (26) and (27), a general solution of (25) is

$$\left. \begin{aligned} y_2 = A_1 \sin x + A_2 \cos x - \frac{8}{9} \sin x \cos x - \frac{16}{9} \cos^3 x - \frac{16}{9} \sin^3 x + 2x \cos x \\ y_2' = A_1 \cos x - A_2 \sin x + \frac{8}{9} (\sin^2 x - \cos^2 x) + \frac{2}{3} (x \sin x - \cos x) - \frac{1}{3} \cos 3x \end{aligned} \right\} \quad \dots (28)$$

Using the initial conditions of (17), (28) yields,

$$A_2 = 0, A_1 = \frac{8}{9} + \frac{2}{9} + \frac{1}{2} = \frac{17}{9} \quad \dots (29)$$

Hence using (28) and (29), a particular solution of (25) satisfying

$$y_2(0)=0, y_2'(0)=0 \text{ is} \quad \dots (30)$$

$$y_2 = \frac{8}{9} \sin x - \frac{8}{9} \sin x \cos x - \frac{16}{9} \cos^3 x + \frac{16}{9} \sin^3 x + 2x \cos x$$

Making substitutions for y_0 , y_1 and y_2 from (21), (24) and (30) to

(16) we find the solution as

$$y = \sin x + \epsilon \left(\frac{2}{3} \cos x - \frac{4}{3} \sin^2 x - \frac{2}{3} \cos^2 x \right) + \epsilon^2 \left(\frac{17}{9} \sin x - \frac{8}{9} \sin x \cos x + \frac{1}{3} \cos x - \frac{1}{3} \sin 3x \right) \quad \dots (31)$$

This follows that in the absence of disturbing function the solution of (15) is $\sin x$, while in the presence of disturbing function $-2\epsilon(y')^2$, the first and second order perturbation terms are respectively the second and third terms in (31).

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 51. Transform the equation $\Delta^2 \psi = \frac{1}{c^2} \frac{\partial^2 \psi}{\partial t^2}$ to spherical polar coordinates. If ψ depends on r and t only, show that the equation can be written in the form

$$\frac{\partial^2}{\partial r^2} (r\psi) = \frac{1}{c^2} \frac{\partial^2}{\partial t^2} (r\psi)$$

Hence show that the general solution is of the form

$$\psi(r, t) = \frac{1}{r} \{ f(r-ct) + g(r+ct) \}$$

(Bombay, 1965)

Explain the physical meaning of this solution

Hint. Use the transformations $x = r \sin \theta \cos \phi$, $y = r \sin \theta \sin \phi$, $z = r \cos \theta$.

Problem 52. Obtain Newton's formula for interpolation. Using this or some other formula find the value of the derivative at a point.

The following table gives the corresponding values of x and y

x	3	4	5	6	7	8
y	205	240	259	262	250	224

Find the value of x for which y is maximum.

(Bombay, 1970)

HARMONICS (With Special Functions)

8.1. INTRODUCTION

In the previous chapter on Differential Equations we have already mentioned that an equation of the type $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial z^2} = 0$ or in terms of 'del' operator ∇ , $\nabla^2 V = 0$

is known as *Laplace's equation*. It frequently appears in applied mathematics when discussions are made on mechanics, sound, electricity, heat etc, wherever the theory of potential is involved e.g. if V is the Newtonian potential due to an attracting mass at a point (x, y, z) not forming a part of the mass itself then V satisfies (1). Similarly if V be the electric potential at (x, y, z) where the electric density is zero then V satisfies (1). Moreover if a body is in a state of equilibrium as to temperature, V being the temperature at (x, y, z) ,

$$\frac{dV}{dt} = 0 \text{ and } V \text{ satisfies (1).}$$

The equation (1) can be transformed to spherical polar form by using the transformation $x = r \sin \theta \cos \phi$, $y = r \sin \theta \sin \phi$, $z = r \cos \theta$ so that $r^2 = x^2 + y^2 + z^2$, $\phi = \tan^{-1} \frac{y}{x}$, $\theta = \tan^{-1} \frac{\sqrt{x^2 + y^2}}{z}$

$$\text{and } \frac{\partial r}{\partial x} = \frac{x}{r} = \sin \theta \cos \phi, \frac{\partial r}{\partial y} = \frac{y}{r} = \sin \theta \sin \phi, \frac{\partial r}{\partial z} = \frac{z}{r} = \cos \theta,$$

$$\frac{\partial \theta}{\partial x} = \frac{\cos \theta \cos \phi}{r}, \frac{\partial \theta}{\partial y} = \frac{\cos \theta \sin \phi}{r}, \frac{\partial \theta}{\partial z} = -\frac{\sin \theta}{r},$$

$$\frac{\partial \phi}{\partial x} = -\frac{\sin \phi}{r \sin \theta}, \frac{\partial \phi}{\partial y} = \frac{\cos \phi}{r \sin \theta}, \frac{\partial \phi}{\partial z} = 0$$

$$\text{whence } \frac{\partial u}{\partial x} = \frac{\partial u}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial u}{\partial \theta} \frac{\partial \theta}{\partial x} + \frac{\partial u}{\partial \phi} \frac{\partial \phi}{\partial x} = \frac{\partial u}{\partial r} \sin \theta \cos \phi$$

$$+ \frac{\partial u}{\partial \theta} \frac{\cos \theta \cos \phi}{r} + \frac{\partial u}{\partial \phi} \left(-\frac{\sin \phi}{r \sin \theta} \right)$$

$$\text{giving } \frac{\partial}{\partial x} \equiv \sin \theta \cos \phi \frac{\partial}{\partial r} + \frac{\cos \theta \cos \phi}{r} \frac{\partial}{\partial \theta} - \frac{\sin \phi}{r \sin \theta} \frac{\partial}{\partial \phi}$$

$$\begin{aligned} \frac{\partial^2 V}{\partial x^2} &= \frac{\partial}{\partial x} \left(\frac{\partial V}{\partial x} \right) = \left(\sin \theta \cos \phi \frac{\partial}{\partial r} + \frac{\cos \theta \cos \phi}{r} \frac{\partial}{\partial \theta} - \frac{\sin \phi}{r \sin \theta} \frac{\partial}{\partial \phi} \right) \\ &\quad \left(\sin \theta \cos \phi \frac{\partial V}{\partial r} + \frac{\cos \theta \cos \phi}{r} \frac{\partial V}{\partial \theta} - \frac{\sin \phi}{r \sin \theta} \frac{\partial V}{\partial \phi} \right) \\ &= \sin^2 \theta \cos^2 \phi \frac{\partial^2 V}{\partial r^2} + \frac{2 \sin \theta \cos \theta \cos^2 \phi}{r} \frac{\partial^2 V}{\partial r \partial \theta} \\ &\quad - \frac{2 \sin \theta \cos \theta \cos^2 \phi}{r^2} \frac{\partial V}{\partial \theta} - \frac{2 \sin \phi \cos \phi}{r} \frac{\partial^2 V}{\partial r \partial \phi} \\ &\quad + \frac{\sin \phi \cos \phi}{r^2} \frac{\partial V}{\partial \phi} + \frac{\cos^2 \theta \cos^2 \phi}{r} \frac{\partial^2 V}{\partial r} \\ &\quad + \frac{\cos^2 \theta \cos^2 \phi}{r^2} \frac{\partial^2 V}{\partial \theta^2} - \frac{2 \cos \theta \sin \phi \cos \phi}{r^2 \sin \theta} \frac{\partial^2 V}{\partial \theta \partial \phi} \\ &\quad + \frac{\cos^2 \theta \sin \phi \cos \phi}{r^2 \sin^2 \theta} \frac{\partial V}{\partial \phi} + \frac{\sin^2 \phi}{r} \frac{\partial V}{\partial r} + \frac{\cos \theta \sin^2 \phi}{r^2 \sin \theta} \frac{\partial V}{\partial \theta} \\ &\quad + \frac{\sin^2 \phi}{r^2 \sin^2 \theta} \frac{\partial^2 V}{\partial \phi^2} + \frac{\sin \phi \cos \phi}{r^2 \sin \phi} \frac{\partial V}{\partial \phi} \end{aligned}$$

with similar expression for $\frac{\partial^2 V}{\partial y^2}$ by replacing ϕ by $\frac{\pi}{2} + \phi$ in this expression

$$\begin{aligned} \text{and } \frac{\partial^2 V}{\partial z^2} &= \frac{\partial}{\partial z} \left(\frac{\partial V}{\partial z} \right) = \left(\cos \theta \frac{\partial}{\partial r} - \frac{\sin \theta}{r} \frac{\partial}{\partial \theta} \right) \left(\cos \theta \frac{\partial V}{\partial r} - \frac{\sin \theta}{r} \frac{\partial V}{\partial \theta} \right) \\ &= \cos^2 \theta \frac{\partial^2 V}{\partial r^2} - \frac{2 \sin \theta \cos \theta}{r} \frac{\partial^2 V}{\partial r \partial \theta} + \frac{2 \sin \theta \cos \theta}{r} \frac{\partial V}{\partial \theta} + \frac{\sin^2 \theta}{r} \frac{\partial V}{\partial r} \\ &\quad + \frac{\sin^2 \theta}{r^2} \frac{\partial^2 V}{\partial \theta^2} \end{aligned}$$

With these substitutions, the spherical polar form of (1) is

$$\frac{\partial^2 V}{\partial r^2} + \frac{2}{r} \frac{\partial V}{\partial r} + \frac{1}{r^2} \frac{\partial^2 V}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial V}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 V}{\partial \phi^2} = 0 \quad \dots (2)$$

Which may be put as

$$\frac{1}{r^2} \left\{ \frac{\partial}{\partial r} \left(r^2 \frac{\partial V}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial V}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2 V}{\partial \phi^2} \right\} = 0 \quad \dots (3)$$

Also if $\mu = \cos \theta$, it takes the form

$$r \frac{\partial^2 (Vr)}{\partial r^2} + \frac{\partial}{\partial \mu} \left\{ (1 - \mu^2) \frac{\partial V}{\partial \mu} \right\} + \frac{1}{1 - \mu^2} \frac{\partial^2 V}{\partial \phi^2} = 0 \quad \dots (4)$$

The subject matter of *Spherical Harmonics* is partly concerned with the development of the functions which will satisfy this equation.

A homogeneous rational integral algebraic function of (x, y, z) of degree n in the form $r^n f(\theta, \phi)$ in spherical polar coordinates which is a value of V satisfying (1) is said to be a *solid spherical Harmonics* of n th degree and the function $f(\theta, \phi)$ is said to be a *surface spherical Harmonics* of n th degree. Since Laplace employed these spherical

Harmonics in determining V , these are also known as *Laplace's coefficients*.

In case V is independent of ϕ i.e. $\frac{\partial V}{\partial \phi} = 0$, and so $\frac{\partial^2 V}{\partial \phi^2} = 0$, then (1) reduces to

$$r \frac{\partial^2(rV)}{\partial r^2} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial V}{\partial \theta} \right) = 0 \quad \dots (4)$$

If we put $V = r^n P$, P being a function of θ only and then change the independent variable θ by the transformation $\mu = \cos \theta$, the equation (5) yields

$$\frac{d}{d\mu} (1 - \mu^2) \frac{dP}{d\mu} + n(n+1) P = 0 \quad \dots (6)$$

which is known as Legendre's equation and will be discussed a bit later.

A function satisfying (5) or (6) is said to be a *Surface Zonal Harmonics*. A special class of zonal harmonics is sometimes known as *Legendrean Coefficients*.

There are equations belonging to such a form which cannot be solved by any of the methods discussed in the previous chapter then we claim to find a convergent series arranged according to powers of the independent variable, which will approximately express the value of the dependent variable. For this purpose the series solution of linear differential equations will be discussed in the next section and then in the subsequent sections the series integration method will be used to discuss the solutions of very important equations that often occur in investigations in applied mathematics such as the equations of Riccati, Bessel, Legendre, Hermite, Laguerre and the Hypergeometric series.

Note. Bessel's functions are known as *Cylindrical Harmonics*.

8.2. METHODS OF INTEGRATION IN SERIES

[A] Power Series Solutions of Linear Differential Equations.

We know that a linear differential equation is one in which both the dependent variables and their derivatives are of the first degree and these do not occur as products of dependent variables and/or their derivatives.

A series of the form

$$a_0 + a_1(x-x_0) + a_2(x-x_0)^2 + a_3(x-x_0)^3 + \dots \quad \dots (1)$$

where $a_0, a_1, a_2, \dots, x_0$ are constants and x is a variable, is said to be a *power series*, and it may converge

(i) only for the single value $x = x_0$

(ii) absolutely for $|x - x_0| < \epsilon$ i.e. for values of x in the neighbourhood of x_0 and diverge for $|x - x_0| < \epsilon$ while at the end points $x_0 \pm \epsilon$, it may either converge or diverge.

(iii) absolutely for all x i.e. for $-\infty < x < \infty$.

The set of values of x for which the power series converges is said to be the *Interval of Convergence* and denoted by I .

In case a power series converges on an interval $I: |x - x_0| < R$, R being a positive constant, then the power series defines a function $f(x)$ which is continuous for each x in I .

If $f(x)$ be a function defined by a power series i.e.

$$f(x) = a_0 + a_1(x-x_0) + a_2(x-x_0)^2 + a_3(x-x_0)^3 + \dots \text{for } I: |x-x_0| < R \quad \dots (2)$$

then the power series obtained by differentiating each term of (2) defines the derivative of $f(x)$ on the same interval I i.e.

$$f'(x) = a_1 + 2a_2(x-x_0) + 3a_3(x-x_0)^2 + \dots \text{for } I: |x-x_0| < R \quad \dots (3)$$

We can thus define the successive derivatives $f''(x), f'''(x), \dots$ and so on.

It is clear that $f(x_0) = a_0, f'(x_0) = a_1, \frac{f''(x_0)}{2!} = a_2, \dots$ etc.

As such (2) yields

$$f(x) = f(x_0) + f'(x_0)(x-x_0) + \frac{f''(x_0)}{2!}(x-x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x-x_0)^n + \dots; |x-x_0| < R \quad \dots (4)$$

which is *Taylor's series expansion* in powers of $(x-x_0)$.

If $x_0 = 0$ this becomes

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots, |x| < R \quad \dots (5)$$

which is *Maclaurin's series expansion* in power of x .

Two functions $f(x)$ and $g(x)$ defined by power series such that

$$f(x) = a_0 + a_1(x-x_0) + a_2(x-x_0)^2 + \dots, |x-x_0| < R$$

$$g(x) = b_0 + b_1(x-x_0) + b_2(x-x_0)^2 + \dots, |x-x_0| < R$$

will be equal if and only if $a_0 = b_0, a_1 = b_1, a_2 = b_2$ etc.

A function $f(x)$ is said to be *analytic* or *regular* at a point $x = x_0$ if it can be expanded as a Taylor's series expansion in powers of $(x-x_0)$ for every x in the neighbourhood of x_0 and it is *analytic on the interval I* if it is analytic at every point of the interval.

Now consider a linear differential equation with variable coefficients, such that

$$y^{(n)} - f_{n-1}(x)y^{(n-1)} + \dots + f_1(x)y' + f_0(x)y = Q(x) \quad \dots (6)$$

Where $y^{(n)} = \frac{d^n y}{dx^n}$ etc..., then a sufficient condition for the existence of a power series solution of (6) is that each function $f_0(x), f_1(x), \dots, f_{n-1}(x), Q(x)$ is analytic at $x = x_0$ and the solution which is unique must satisfy the n initial conditions

$$y(x_0)=a_0, y'(x_0)=a_1, \dots, y^{(n-1)}(x_0)=a_{n-1}.$$

There are two methods for solving such equations by series integrations.

(1) *First Series Method* in which the series solution is found by successive differentiation.

(2) *Second Series Method* in which the series solution is obtained by undetermined coefficients.

We explain these methods applying on the following Problem.

Problem 1. Apply Power-series method to find the solution of the following linear equation:

$$\frac{d^2y}{dx^2} - (x+1)\frac{dy}{dx} + x^2y = x$$

i.e. $y'' - (x+1)y' + x^2y = x$ with the initial conditions $y(0)=1$, $y'(0)=1$

We have $y'' - (x+1)y' + x^2y = x$... (1)

Its comparison with (6) of §8.2 gives

$$f_0(x) = x^2, f_1(x) = -x-1, Q(x) = x.$$

All these functions being polynomials, the series solution of (1) is valid for all x , since a polynomial is a finite series and therefore the functions $f_0(x)$, $f_1(x)$, $Q(x)$ each being polynomial, every solution of (1) has a Taylor series expansion valid for all x .

First Method. Suppose we seek a solution in the form of Maclaurin series. Then replacing $f(x)$ by $y(x)$ in (5) of §8.2, we get

$$y(x) = y(0) + y'(0)x + \frac{y''(0)}{2!}x^2 + \frac{y'''(0)}{3!}x^3 + \frac{y^{(4)}(0)}{4!}x^4 + \dots \quad \dots (2)$$

Initially when $x=0$, $y=1$, $y'=1$, so that (1) gives $y''(0)=1$... (3)

We thus know the values $y(0)$, $y'(0)$, $y''(0)$ and in order to find the values of succeeding coefficients, let us take successive derivatives of (1) and evaluate them at $x=0$, $y=1$. Differentiating (1) w.r.t x , we get

$$y''' - (x+1)y'' - y' + x^2y' + 2xy = 1 \quad \dots (4)$$

which yields $y'''(0)=3$... (5)

when $x=0$, $y=1$, $y'=1$, $y''=1$.

Again differentiating (4) w.r.t x , we find

$$y^{(4)} - (x+1)y''' - 2y'' + x^2y'' + 4xy' + 2y = 0.$$

Which yields $y^{(4)}(0)=3$... (6)

when $x=0$, $y=y'=y''=1$, $y'''=3$.

Similarly $y^{(5)}(0)$, $y^{(6)}(0)$, ... etc. can be evaluated.

Substituting these values in (2) we get the required series solution as

$$y(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{2} + \frac{x^4}{8} + \dots, -\infty < x < 8 \quad \dots (7)$$

Second Method. Suppose the series solution in powers of x has the form

$$y(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + \dots \quad \dots (8)$$

Since $y(x)$ is valid for all x , therefore its two successive derivatives will also be valid for all x . The two successive derivatives of (8) are

$$y'(x) = a_1 + 2a_2x + 3a_3x^2 + 4a_4x^3 + \dots \quad \dots (9)$$

$$y''(x) = 2a_2 + 6a_3x + 12a_4x^2 + \dots \quad \dots (10)$$

Substituting the values of (8), (9) and (10) in (1) we get

$$2a_2 + 6a_3x + 12a_4x^2 + \dots - (x+1)(a_1 + 2a_2x + 3a_3x^2 + 4a_4x^3 + \dots) + x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + \dots) = x$$

$$\therefore (2a_2 - a_1) + (6a_3 - 2a_2 - a_1 - 1)x + (12a_4 - 3a_3 - 2a_2 + a_0)x^2 + \dots = 0.$$

This being an identity in x , the coefficients of powers of x vanish separately i.e. $2a_2 - a_1 = 0$ i.e. $a_2 = \frac{a_1}{2}$

$$\left. \begin{aligned} 6a_3 - 2a_2 - a_1 - 1 &= 0 \text{ i.e., } a_3 = \frac{2a_2 + a_1 + 1}{6} \\ 12a_4 - 3a_3 - 2a_2 + a_0 &= 0 \text{ i.e., } a_4 = \frac{3a_3 + 2a_2 - a_0}{12} \end{aligned} \right\} \dots (11)$$

Comparison of (8) with Taylor series i.e. (4) of §8.2 yields, $f(x_0) = y(0) = a_0 = 1$, $f'(x_0) = y'(0) = a_1 = 1$.

As such (11) give $a_2 = \frac{1}{2}$, $a_3 = \frac{1}{2}$, $a_4 = \frac{1}{8}$.

Substituting these values in (8) we get the required series solution as

$$y(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{2} + \frac{x^4}{8} + \dots, -\infty < x < \infty.$$

Problem 2. Find by power series methods a particular solution of the linear differential equation:

$$y''' + \frac{1}{x}y' - \frac{1}{x^2}y = 0, x \neq 0 \text{ and } y(1)=1, y'(1)=0, y''(1)=1$$

Here $f_0(x) = -\frac{1}{x^2}$, $f_1(x) = \frac{1}{x}$, $Q(x) = 0$ etc.

$$\text{Ans. } y(x) = 1 + \frac{(x-1)^2}{2} + \frac{(x-1)^3}{6} - \frac{(x-1)^4}{8} + \frac{(x-1)^5}{15} - \dots, 0 < x < 2$$

[B] Ordinary Points and Singularities of a Linear Differential Equation.

Consider a linear differential equation of the form

$$y^{(n)} + F_{n-1}(x)y^{(n-1)} + \dots + F_1(x)y' + F_0(x)y = Q(x) \quad \dots (7)$$

A point $x=x_0$ is said to be an *ordinary point* of (7) if each function $F_0, F_1, \dots, F_{n-1}$ and Q is analytic at $x=x_0$ i.e. each function has a Taylor series expansion in powers of $(x-x_0)$ valid in the neighbourhood of x_0 .

Whenever $x=x_0$ is an ordinary point then (7) has a solution which is also analytic at $x=x_0$ i.e., the solution has a Taylor series

representation in powers of $(x-x_0)$ in the neighbourhood of x_0 .

A point $x=x_0$ is said to be a *singularity* of (7) if one or more of the functions $F_0(x)$, ..., $F_{n-1}(x)$, $Q(x)$ are not analytic at $x=x_0$.

If we now consider a second order linear equation

$$y'' + F_1(x)y' + F_2(x)y = 0 \quad \dots (8)$$

F_1, F_2 being continuous functions of x , then if $x=x_0$ is a singularity of (8) and the product functions $F_1(x)(x-x_0)$, and $F_2(x)(x-x_0)^2$ both are analytic at $x=x_0$, then the point $x=x_0$ is said to be a *regular singularity*. In case one or both of the product functions are not analytic at $x=x_0$ then the point $x=x_0$ is said to be an *irregular singularity*.

[C] Frobenius Method for Solving a Homogeneous Linear Differential Equation.

Consider,

$$y'' + F_1(x)y' + F_2(x)y = 0 \quad \dots (9)$$

In case (9) has an irregular singularity at $x=x_0$ then it is too difficult to find the series solution of (9) here, but if it has a regular singularity at $x=x_0$ then the series solution of (9) can be found out in the neighbourhood of x_0 . For the purpose of finding the solution in later case, Frobenius introduced a series solution i.e.,

$$y = (x-x_0)^m [a_0 + a_1(x-x_0) + a_2(x-x_0)^2 + a_3(x-x_0)^3 + \dots], a_0 \neq 0 \quad \dots (10)$$

which is known as a *Frobenius Series*.

When $m=0$, (10) reduces to the usual Taylor series and hence Taylor's series is a special case of Frobenius series.

In case m is negative or has non-integral positive values then (10) is not a Taylor series.

Assuming that x_0 is a regular singularity of (9) and F_1 or F_2 or both are not analytic at $x=x_0$ but $(x-x_0)F_1$ and $(x-x_0)^2F_2$ both are analytic at $x=x_0$, it follows that F_1 has $(x-x_0)$ in its denominator and/or F_2 has $(x-x_0)^2$ in its denominator, thereby showing that either $F_1(x) = f_1(x)/(x-x_0)$ or $F_2(x) = f_2(x)/(x-x_0)^2$ or both exists. In either case if (9) is multiplied by $(x-x_0)^2$, it will transform to the form

$$(x-x_0)^2 y'' + (x-x_0) f_1(x) y' + f_2(x) y = 0 \quad \dots (11)$$

where $f_1(x)$ and $f_2(x)$ both now become analytic at $x=x_0$.

Existence of a Frobenius Series Solution.

If x_0 is a regular singularity of (11), then it has at least one Frobenius series solution of the form (10) if it is valid in the common interval of convergence of $f_1(x)$ and $f_2(x)$ of (11) except possibly for $x=x_0$.

Without loss of generality, taking $x_0=0$, (11) reduces to

$$x^2 y'' + x f_1(x) y' + f_2(x) y = 0 \quad \dots (12)$$

where $f_1(x)$ and $f_2(x)$ are analytic at $x=0$ and therefore each function has a Taylor series expansion in powers of x valid in the neighbourhood of $x=0$. Let us assume

$$f_1(x) = b_0 + b_1 x + b_2 x^2 + \dots \quad \dots (13)$$

$$f_2(x) = c_0 + c_1 x + c_2 x^2 + \dots \quad \dots (14)$$

For $x_0=0$, Frobenius series i.e., (10) and its next two derivatives become

$$y = a_0 x^m + a_1 x^{m+1} + a_2 x^{m+2} + \dots + a_n x^{m+n} + \dots, a_0 \neq 0, \quad \dots (15)$$

$$y' = a_0 m x^{m-1} + a_1 (m+1) x^m + a_2 (m+2) x^{m+1} + \dots + a_n (m+n) x^{m+n-1} + \dots \quad (16)$$

$$y'' = a_0 m(m-1) x^{m-2} + a_1 m(m+1) x^{m-1} + a_2 (m+1)(m+2) x^m + \dots + a_n (m+n-1)(m+n) x^{m+n-2} + \dots \quad (17)$$

$y(x)$ will be a solution of (12) if (13), (14), (15), (16), (17) identically satisfy (12) i.e., if

$$x^2 [a_0 m(m-1) x^{m-2} + a_1 m(m+1) x^{m-1} + \dots + a_n (m+n-1)(m+n) x^{m+n-2} + \dots] + x [b_0 + b_1 x + b_2 x^2 + \dots] [a_0 m x^{m-1} + a_1 (m+1) x^m + \dots + a_n (m+n) x^{m+n-1} + \dots] + (c_0 + c_1 x + c_2 x^2 + \dots) [a_0 x^m + a_1 x^{m+1} + \dots + a_n x^{m+n} + \dots] \equiv 0 \quad \dots (18)$$

If we expand (18) and collect the coefficient of like power of x , then we get

$$a_0 [m(m-1) + b_0 m + c_0] x^m + [a_1 \{(m+1)m + b_0(m+1) + c_0\} + a_0 \{b_1 m + c_1\}] x^{m+1} + \dots + [a_n \{(m+n)(m+n-1) + b_0(m+n) + c_0\} + a_{n-1} \{(b_n m + c_n)\}] x^{m+n} \equiv 0 \quad \dots (19)$$

This relation being an identity, each of the coefficients of x^k , $k=m, m+1, \dots, m+n$, is zero. But since $a_0 \neq 0$, therefore the first term in (19) gives

$$m(m-1) + b_0 m + c_0 = 0 \quad \dots (20)$$

which is known as *Indicial Equation*. This is quadratic in m and hence has two roots say m_1 and m_2 . There arise three cases according as $m_1 \neq m_2$ and $m_1 - m_2 = \text{not an integer}$; $m_1 \neq m_2$ but $m_1 - m_2 = \text{an integer}$; $m_1 = m_2$.

Case I. When m_1, m_2 are distinct and their difference is not an integer.

Let us first put $m=m_1$ in all the coefficients of (19) except the first one and then equate each equal to zero. We are thus capable of finding $a_1, a_2, \dots, a_n, \dots$ in terms of a_0 . Substituting these values of $a_1, a_2, \dots, a_n$, in (15) we get the solution $y(x)$.

With similar procedure taking $m=m_2$, we may get another solution.

Problem 3. Solve $2x^2 y'' - xy' + (1-x^2)y = 0$.

Let the series solution be

$$y = a_0 x^m + a_1 x^{m+1} + a_2 x^{m+2} + \dots + a_n x^{m+n} + \dots, a_0 \neq 0 \quad \dots (1)$$

$$\text{so that } y' = m a_0 x^{m-1} + (m+1) a_1 x^m + (m+2) a_2 x^{m+1} + \dots + (m+n) a_n x^{m+n-1} + \dots$$

$$y'' = m(m-1) a_0 x^{m-2} + (m+1)m a_1 x^{m-1} + \dots + (m+n)(m+n-1) a_n x^{m+n-2} + \dots$$

Substituting these values in the given equation, we get an identity
 $(m-1)(2m-1)a_0x^m + m(2m+1)a_1x^{m+1} + \{m+1\}(2m+3)a_2x^{m+2} + \dots + \{m+n-1\}(2m+2n-1)a_n - a_{n-2}\}x^{m+n} = 0$... (2)

Since $a_0 \neq 0$, the coefficient of first term vanishes if $x=1$, $\frac{1}{2}$ i.e., roots of the indicial equation are 1 and $\frac{1}{2}$ which are distinct and not differ by an integer.

The coefficients of x^{m+n} equated to zero gives the *recursion formula*

$$a_n = \frac{1}{(m+n-1)(2m+2n-1)} a_{n-2} \text{ for } n \geq 2 \quad \dots (3)$$

since all terms except the first two in (2) vanish if $a_2, a_3, \dots$ satisfy (3)

The second term of (2) gives $a_1 = 0$ since $m(2m+1) \neq 0$ for $m=1, \frac{1}{2}$

When $m=1$, (3) gives $a_n = \frac{1}{n(2n+1)} a_{n-2}$.

Putting $n=2, 4, 6, 8, \dots$ we get (since $a_1=a_3=a_5=\dots=0$)

$$a_2 = \frac{1}{2} a_0$$

$$a_4 = \frac{1}{4 \cdot 9} a_2 = \frac{1}{2 \cdot 5 \cdot 4 \cdot 9} a_0 = \frac{1}{2 \cdot 4 \cdot 5 \cdot 9} a_0$$

$$a_6 = \frac{1}{6 \cdot 13} a_4 = \frac{1}{2 \cdot 4 \cdot 6 \cdot 5 \cdot 9 \cdot 13} a_0 \text{ etc.}$$

Hence from (1), say $y_1 = a_0 x \left[1 + \frac{x^2}{2 \cdot 5} + \frac{x^6}{2 \cdot 4 \cdot 5 \cdot 9} + \frac{x^6}{2 \cdot 4 \cdot 6 \cdot 5 \cdot 9 \cdot 13} + \dots \right]$

Again when $n = \frac{1}{2}$, (3) gives $a_n = \frac{1}{\left(n - \frac{1}{2}\right) \cdot 2n} a_{n-2}$

Here also $a_1 = a_3 = a_5 = \dots = 0$

and

$$a_2 = \frac{1}{3 \cdot 4} a_0 = \frac{1}{2 \cdot 3} a_0$$

$$a_4 = \frac{1}{2 \cdot 4 \cdot 3 \cdot 7} a_0$$

$$a_6 = \frac{1}{2 \cdot 4 \cdot 6 \cdot 3 \cdot 7 \cdot 11} a_0 \text{ etc.}$$

So that from (1), say $y_2 = a_0 \sqrt{x} \left[1 + \frac{x^2}{2 \cdot 3} + \frac{x^6}{2 \cdot 4 \cdot 3 \cdot 7} + \frac{x^6}{2 \cdot 4 \cdot 6 \cdot 3 \cdot 7 \cdot 11} + \dots \right]$

The complete solution is

$$y = Ay_1 + By_2$$

$$= Ax \left[1 + \frac{x^2}{2 \cdot 5} + \frac{x^4}{2 \cdot 4 \cdot 5 \cdot 9} + \dots \right] + B \sqrt{x} \left[1 + \frac{x^2}{2 \cdot 3} + \frac{x^4}{2 \cdot 4 \cdot 3 \cdot 7} + \dots \right]$$

where a_0 has been incorporated into the arbitrary constants A and B .

Problem 4. Solve $x^2 y'' + x(x + \frac{1}{2})y' - (x^2 + \frac{1}{2})y = 0$.

Ans. $y = Ax \left[1 - \frac{2}{5}x + \frac{9}{35}x^2 - \frac{82}{945}x^3 + \dots \right] + Bx^{-1/2} \left[1 - x + \frac{3}{2}x^2 - \frac{13}{18}x^3 + \dots \right]$

Problem 5. Solve $4xy'' + 2(1-x)y' - y = 0$.

Ans. $y = A \left(1 + \frac{1}{2}x + \frac{1}{2^2 \cdot 2}x^2 + \frac{1}{2^3 \cdot 3}x^3 + \dots \right) + B \sqrt{x} \left[1 + \frac{x}{1 \cdot 3 \cdot 5} + \frac{x^2}{1 \cdot 3 \cdot 5} + \frac{x^3}{1 \cdot 3 \cdot 5 \cdot 7} + \dots \right]$

Case II. When $m_1 \neq m_2$ and their difference is an integer.

If m_1, m_2 differ by a non-zero integer, we can write them as m and $m+N$ where N is a positive integer. Then the indicial equation (20) can be written as

$$(m+N)(m+N-1) + b_0(m+N) + c_0 = 0 \quad \dots (21)$$

since $m+N$ is a root of (20).

Comparison of L.H.S. of (21) with the coefficient of a_n in (19) shows that both of them are exactly the same except that n is replaced by N . This follows that the use of smaller root m in (19) to find a 's, will make each coefficient of x^k zero and so we shall be stopped at the term in which a_N appears and whose coefficient is zero. Consequently the equation for a_N can not be solved in terms of previous a 's unless accidentally the remaining terms also add to zero. In such cases, the equation will be satisfied for all arbitrary values of a_N . We may then continue to find values of $a_{N+1}, a_{N+2}, \dots$ in terms of a_0 and a_N .

Now there arise two possibilities:

Possibility (1). In (19), the coefficient of a_N is zero and the remaining terms in the coefficient of x^{m+N} also add to zero. Then the larger root $m+N$ will determine a set of values of a 's in terms of a_0 and the smaller root m will determine two sets of values of a 's; one in terms of a_0 and the other in terms of a_N , whose linear combination will yield the general solution.

Problem 6. Solve $x^2 y'' + xy' + \left(x^2 - \frac{1}{2^2}\right)y = 0$, (which is Bessel equation of index $\frac{1}{2}$).

Clearly $x=0$ is a regular singularity of the equation

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{2^2}\right)y = 0 \quad \dots (1)$$

Let its Frobenius series solution be

$$y = a_0 x^m + a_1 x^{m+1} + a_2 x^{m+2} + \dots \quad (1)$$

Derive y' and y'' and put these values in (1). Proceeding first like in Problem 3, we shall find, the indicial equation as

$$m(m-1) + m - \frac{1}{4} = 0 \text{ i.e. } m^2 - \frac{1}{4} = 0 \quad (2)$$

Giving $m = \pm \frac{1}{2}$ which differ by an integer unity.

The indicial equation may also be found by comparing (1) with (12) of § 8.2 [C] whence we get $f_1(x) = 1$, $f_2(x) = -\frac{1}{4} + x^2$

So that $f_1(x) = 1 = b_0 + b_1 x + b_2 x^2 + \dots$ which is (13) of § 8.2 (C) and $f_2(x) = -\frac{1}{4} + x^2 = c_0 + c_1 x + c_2 x^2 + \dots$ which is (14) of § 8.2 (C)

Comparison gives, $b_0 = 1$, $c_0 = -\frac{1}{4}$, $c_1 = 0$, $c_2 = 1$, remaining b 's

and c 's being zero.

Substituting these values in the indicial equation (20) we get (1) giving $m = -\frac{1}{2}$, $\frac{1}{2}$ so that $N = \frac{1}{2} - (-\frac{1}{2}) = 1$.

Taking smaller root $-\frac{1}{2}$ and setting the second coefficient of (19) to zero, we find $a_1 [\frac{1}{2}(-\frac{1}{2}) + \frac{1}{2} - \frac{1}{4}] + a_0(0) = 0$ i.e. $0 \cdot a_1 + 0 \cdot a_0 = 0$.

Since the coefficient of a_1 is zero and the other term in its coefficient is also zero, therefore we have the first possibility. Thus a_0 and a_1 are arbitrary.

In case of $m = -\frac{1}{2}$, the recursion formula by setting the coefficient of x^{m+n} as zero is $a_n \left[\left(-\frac{1}{2} + n\right) \left(-\frac{3}{2} + n\right) + \left(-\frac{1}{2} + n\right) - \frac{1}{4} \right] + a_{n-1} = 0$

$$(0) + a_{n-2} = 0$$

$$\text{i.e. } a_n = -\frac{a_{n-2}}{n^2 - n}, \text{ for } n \geq 2$$

which yields,

$$\begin{aligned} a_2 &= -\frac{1}{2} a_0 & \text{and} & & a_3 &= -\frac{1}{6} a_1 \\ a_4 &= -\frac{1}{12} a_2 = \frac{1}{24} a_0 & a_5 &= -\frac{1}{20} a_3 = -\frac{1}{120} a_1 \\ a_6 &= -\frac{1}{30} a_4 = -\frac{1}{720} a_0 & a_7 &= -\frac{1}{42} a_5 = -\frac{1}{5040} a_1 \\ \text{etc...} & & \text{etc...} & & \end{aligned}$$

Substituting these values in (2), we get the solution for $m = -\frac{1}{2}$,

$$y = a_0 x^{-1/2} \left(1 - \frac{1}{2} x^2 + \frac{1}{24} x^4 - \frac{1}{720} x^6 + \dots \right) + a_1 x^{1/2} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + \dots \right)$$

Possibility (2). In (19) the coefficient of a_N is zero and the remaining terms in the coefficient of x^{m+N} do not add to zero. Then the larger root $m+N$ will determine a set of values of a 's in terms of a_0 . In this case there will be only one Frobenius series solution of (12) and a second independent solution will be of the form

$$y_2(x) = u(x) - b_N y_1(x) \log x, \quad n > 0 \quad (22)$$

where N is the positive integral difference between m_1 and m_2 ; $y_1(x)$ is a Frobenius series solution of (12) with the larger root $m+N$ and $u(x)$ is a Frobenius series of the form

$$u(x) = x^m (b_0 + b_1 x + b_2 x^2 + \dots) \quad (23)$$

where m is the smaller root of (20). If we substitute (22) and (23) with their necessary derivatives in (12), we shall find that $y_2(x)$ is a solution of it provided.

$$x^2 u'' + x f_1 u' + f_2 u = b_N [2x y_1' + (f_1 - 1) y_1] \quad (24)$$

Substituting for u , u' , u'' , y , and y_1' in it, we get the values of b 's in (23). These solutions are known as *Logarithmic solutions*.

Problem 7. Solve $x^2 y'' - x(2-x)y' + (2+x^2)y = 0$ (1)

If we divide (1) by x^2 , then we observe that $x=0$ is a regular singularity.

Let the Frobenius series solution be

$$y = x^m (a_0 + a_1 x + a_2 x^2 + \dots) \quad (2)$$

$$\text{Comparing it with (12), } f_1(x) = -2 + x, f_2(x) = 2 + x^2 \quad (3)$$

So that (13) and (14) give on comparison

$$b_0 = -2, b_1 = 1, c_0 = 2, c_1 = 0, c_2 = 1 \quad (4)$$

the remaining b 's and c 's being zero.

Hence from (20), the indicial equation is

$$m(m-1) - 2m + 2 = 0 \text{ i.e. } m^2 - 3m + 2 = 0 \quad (5)$$

giving $m = 1, 2$ i.e. $m_1 = 1, m_2 = 2$ which differ by $N = 1$.

Taking $m = 1$ and setting the second coefficient of (19) as zero, we have

$$a_1 (2 - 4 + 2) + a_0 (1 + 0) = 0 \text{ i.e. } 0 \cdot a_1 + a_0 = 0 \quad (6)$$

So that $a_0 \neq 0$ but the coefficient of $a_N = a_1$ is zero and hence $m = 1$ cannot give a solution.

Now taking $m = 2$ and setting the second coefficient of (19) equal to zero, we find

$$a_1 (3 \cdot 2 - 2 \cdot 3 + 2) + 2a_0 = 0 \text{ i.e. } a_1 = -a_0 \quad (7)$$

Again by setting the coefficient of x^{m+n} of (19) equal to zero when $m = 2$, the recursion formula is

$$a_n [(2+n)(1+n) - 2(2+n) + 2] + (1+n) a_{n-1} + a_{n-2} = 0$$

$$\text{i.e. } (n^2 + n) a_n = -(n+1) a_{n-1} - a_{n-2} \text{ for } n \geq 2 \quad (8)$$

When $n = 2$, (8) gives $6a_2 = -3a_1 - a_0 = 3a_0 - a_0$ by (7)

$$\text{i.e. } a_2 = \frac{a_0}{3} \quad (9)$$

When $n=3$, (8) gives $12a_3 = -4a_2 - a_1 = -\frac{4}{3}a_0 + a_0$ by (7) and (9)

i.e. $a_3 = -\frac{a_0}{36}$ and so on.

Hence from (2), the solution is

$$y_1 = a_0 x^2 \left[1 - x + \frac{x^2}{2} - \frac{x^3}{36} + \dots \right] \quad \dots (10)$$

A second solution of (1) will have the form

$$y_2(x) = u(x) - b_1 y_1(x) \log x, \quad x > 0 \text{ as } N=1,$$

where $u(x) = x(b_0 + b_1 x + b_2 x^2 + \dots)$.

Case III. When $m_1 = m_2$.

In this case only one Frobenius series solution is possible.

Problem 8. Solve $x^2 y'' + xy' + x^2 y = 0$ (which is Bessel equation for index 0).

Given equation is $x^2 y'' + xy' + x^2 y = 0 \quad \dots (1)$

Dividing it by x^2 we observe that $x=0$ is a regular singularity of it.

Let the series solution of it be

$$y = x^m (a_0 + a_1 x + a_2 x^2 + \dots) \quad \dots (2)$$

Comparing it with (12) of §8.2 (C), we find

$$f_1(x) = 1, f_2(x) = x^2$$

So that (13) and (14) of §8.2 (C) give on comparison

$$b_0 = 1, c_0 = 0, c_1 = 0, c_2 = 1 \quad \dots (3)$$

other b 's and c 's being zero.

The indicial equation (20) becomes here

$$m(m-1) + m = 0 \text{ i.e. } m^2 = 0 \text{ giving } m=0, 0 \quad \dots (4)$$

Taking $m=0$ and setting the second coefficient in (19) equal to zero, we have

$$a_1(1+0) + a_0(0) = 0 \text{ i.e. } a_1 = 0 \quad \dots (5)$$

Again setting the coefficient of x^{m+n} in (19) equal to zero, we find the recursion formula $a_n[n(n-1) + (1)(n)] + a_{n-1}(0) + a_{n-2}(1) = 0$

i.e. $a_n = -\frac{a_{n-2}}{n^2}, n \geq 2 \quad \dots (6)$

So that

$$a_2 = -\frac{1}{2} a_0$$

$$a_3 = -\frac{1}{3^2} a_1 = 0 = a_5 = a_7 = a_9 = \dots$$

$$a_4 = -\frac{1}{4^2} a_2 = \frac{1}{2^2 \cdot 4^2} a_0$$

$$a_6 = -\frac{1}{6^2} a_4 = -\frac{1}{2^2 \cdot 4^2 \cdot 6^2} a_0 \text{ etc.}$$

Hence from (2), a solution is

$$y = (\text{say}) y_1 = a_0 \left(1 - \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} - \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots \right) \quad \dots (7)$$

A second solution may be found by (22) with $N=0$ as

$$y = c_1 y_1 + c_2 \left(\frac{x^2}{2^2} - \frac{1+\frac{1}{2}}{2^2 \cdot 4^2} x^4 + \frac{1+\frac{1}{2}+\frac{1}{6}}{2^2 \cdot 4^2 \cdot 6^2} x^6 + \dots + y_1 \log x \right), x > 0.$$

Another Method. If the indicial equation has equal roots say $m=\alpha$, the two independent solutions are obtained by putting $m=\alpha$ in y and $\frac{\partial y}{\partial m}$ both. The second solution will always consist of the product of the first solution or a numerical multiple of it and $\log_e x$, added to another series.

Problem 9. Solve $x(1-x^2)y'' + (1-3x^2)y' - xy = 0 \quad \dots (1)$

Let the series solution be $y = \sum_{n=0}^{\infty} a_n x^{m+n} \quad \dots (2)$

Then substituting for y, y', y'' from (2) to (1), and equating to zero the coefficient of x^m we get the indicial equation as $m^2 = 0$ i.e. $m=0$, 0 if $a_0 \neq 0$

Also equating to zero the coefficient of x^{m+n} , we shall find the recursion formula

$$a_n = \frac{(m+n-1)^2}{(m+n)^2} a_{n-2} \text{ for } n \geq 2 \text{ which yields when } m=0,$$

$$a_n = \frac{(n-1)^2}{n^2} a_{n-2} \text{ for } n \geq 2 \quad \dots (3)$$

While the coefficient of x^{m+1} equated to zero will give $a_1 = 0$

$$a_2 = \frac{1^2}{2^2} a_0 \text{ and } a_3 = \frac{2^2}{3^2} a_1 = 0 = a_5 = a_7 = \dots$$

$$a_4 = \frac{3^2}{4^2} a_2 = \frac{1^2 \cdot 3^2}{2^2 \cdot 4^2} a_0 \text{ and } a_6 = \frac{1^2 \cdot 3^2 \cdot 5^2}{2^2 \cdot 4^2 \cdot 6^2} a_0 \text{ etc.}$$

Hence a solution is

$$y_1 = a_0 \left[1 + \frac{1^2}{2^2} x^2 + \frac{1^2 \cdot 3^2}{2^2 \cdot 4^2} x^4 + \dots \right] \quad \dots (4)$$

But from $a_n = \frac{(m+n-1)^2}{(m+n)^2} a_{n-2}$ we find

$$y = a_0 x^m \left[1 + \left(\frac{m+1}{m+2} \right)^2 x^2 + \left(\frac{m+1}{m+2} \right)^2 \left(\frac{m+3}{m+4} \right)^2 x^4 + \dots \right] \quad \dots (5)$$

If it is substituted in L.H.S. of (1), we get a single term $a_0 m^2 x^{m-1}$ ($m \neq 0$)

Its partial differential coefficient w.r.t. m is

$$2a_0 m x^{m-1} + a_0 m^2 x^{m-1} \log x \text{ which is zero when } m=0.$$

It means $\frac{\partial}{\partial m} \left[x(1-x^2) \frac{d^2}{dx^2} + (1-3x^2) \frac{d}{dx} - x \right] y = 2a_0 m x^{m-1} + a_0 m^2 x^{m-1} \log x$

$$i.e. \left\{ x(1-x^2) \frac{d^2}{dx^2} + (1-3x^2) \frac{d}{dx} - x \right\} \frac{\partial y}{\partial m} = 2a_0 m x^{m-1} + a_0 m^2 x^{m-1} \log x$$

(the differential operators being commutative)

So $\frac{\partial y}{\partial m}$ is another solution and hence from (5)

$$\frac{\partial y}{\partial m} = y \log x + a_0 x^m \left[2 \left(\frac{m+1}{m+2} \right) \frac{1}{(m+2)^3} x^2 + \dots \right]$$

$$\text{When } m=0, y=y_1=a_0 \left[1 + \frac{1^2}{2^2} x^2 + \frac{1^2 \cdot 3^2}{2^2 \cdot 4^2} x^4 + \dots \right]$$

$$\text{and so } \frac{\partial y}{\partial m} = y_1 \log x + a_0 \left[\frac{1}{2^2} x^2 + \frac{2 \cdot 3}{2^2 \cdot 4^2} \cdot \frac{7}{4} x^4 + \dots \right] \quad (\text{when } m=0) \quad \dots (6)$$

When $m=0$ let in (5) $a_0=a_1$ and in (6) $a_0=b$, then

$$y_1 = a \left(1 + \frac{1^2}{2^2} x^2 + \frac{1^2 \cdot 3^2}{2^2 \cdot 4^2} x^4 + \dots \right) = au \text{ (say)}$$

$$\text{and } \frac{\partial y}{\partial m} = bu \log x + b \left\{ \frac{1}{4} x^2 + \frac{21}{128} x^4 + \dots \right\} = bv \text{ (say)}$$

Therefore the complete primitive is

$$y = Au + Bv$$

$$= (a+b \log x) \left[1 + \frac{1^2}{2^2} x^2 + \frac{1^2 \cdot 3^2}{2^2 \cdot 4^2} x^4 + \dots \right] + b \left[\frac{x^2}{4} + \frac{21x^4}{128} + \dots \right]$$

Problem 10. Solve $xy'' + y' + xy = 0$.

$$\text{Ans. } y = (a+b \log x) \left[1 - \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} - \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots \right] + b \left[\frac{x^2}{2^2} - \frac{x^4}{2^2 \cdot 4^2} \left(1 + \frac{1}{2} \right) + \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} \left(1 + \frac{1}{2} + \frac{1}{3} \right) \dots \right]$$

Problem 11. Solve $x^2(x+1)y'' + x(x+1)y' - y = 0$.

$$\text{Ans. } y = ax \left(1 - \frac{x}{3} + \frac{x^2}{6} - \frac{x^3}{10} + \dots \right) + bx^{-1}(1-x).$$

Problem 12. Solve in series $xy'' + 2y' + xy = 0$.

$$\text{Ans. } y = a \left[1 - \frac{x^2}{3} + \frac{x^4}{5} \dots \right] + bx^{-1} \left[1 - \frac{x^2}{2} + \frac{x^4}{4} \dots \right]$$

[D] Particular Integral (P.I.).

$$\text{Consider } x^4 y'' + xy' + y = \frac{1}{x} \quad \dots (1)$$

Complementary function is the solution of $x^4 y'' + xy' + y = 0$ and will be obtained as above to give

$$C.F. = a_0 \left(1 - \frac{1}{3} x^{-2} - \frac{1}{5} x^{-4} \dots \right) + b_0 \left(x - \frac{1}{x} \right).$$

To find P.I. put $y = C_0 x^m$ in (1), so that $m(m-1) C_0 x^{m+2} = x^{-1}$ which yields

$$m+2 = -1 \text{ and } m(m-1) C_0 = 1 \text{ i.e. } m = -3 \text{ and then } C_0 = \frac{1}{12}.$$

With $m = -3$, find the recursion formula by the usual method, and this is here

$$C_n = \frac{2n}{(2n+3)(2n+4)} C_{n-1} \text{ giving } C_1 = \frac{1}{5 \cdot 6} C_0, C_2 = \frac{2 \cdot 4}{5 \cdot 6 \cdot 7 \cdot 8} C_0, C_3 = \frac{2 \cdot 4 \cdot 6}{5 \cdot 6 \cdot 7 \cdot 8 \cdot 9 \cdot 10} C_0 \text{ etc.} \dots$$

$$\text{Hence P.I.} = \frac{x^{-3}}{12} \left(1 + \frac{2}{5 \cdot 6} x^{-2} + \frac{2 \cdot 4}{5 \cdot 6 \cdot 7 \cdot 8} x^{-4} + \dots \right) = 2x^{-3} \left(\frac{1}{4} + \frac{2}{6} x^{-2} + \frac{2 \cdot 4}{8} x^{-4} + \dots \right)$$

Thus the complete integral is $y = C.F. + P.I.$

8.3. LEGENDRE DIFFERENTIAL EQUATION, FUNCTIONS AND POLYNOMIALS WITH PROPERTIES

[A] Legendre's Differential Equation.

(Agra, 1961, 63)

This equation is of the form

$$(1-x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + n(n+1)y = 0. \quad \dots (1)$$

The equation of such type can be solved in series of ascending or descending powers of x . Suppose, we have to integrate it in a series of descending powers of x . There is no singularity at $x=0$, so the solution of the equation can be obtained in the form of a series developed about $x=0$.

Let us assume the solution of the given equation in the form of series:

$$y = \sum_{r=0}^{\infty} a_r x^{k-r}.$$

$$\therefore \frac{dy}{dx} = \sum_{r=0}^{\infty} (k-r) a_r x^{k-r-1}$$

$$\text{and } \frac{d^2 y}{dx^2} = \sum_{r=0}^{\infty} (k-r)(k-r-1) a_r x^{k-r-2}.$$

Substituting these values in (1), we have

$$\sum_{r=0}^{\infty} [(1-x^2)(k-r)(k-r-1)x^{k-r-2} - 2x(k-r)x^{k-r-1} + n(n+1)x^{k-r}] a_r = 0$$

$$\text{or } \sum_{r=0}^{\infty} [(k-r)(k-r-1)x^{k-r-2} + \{n(n+1)-2(k-r) - (k-r)(k-r-1)\}x^{k-r}] a_r = 0$$

$$\text{or } \sum_{r=0}^{\infty} [(k-r)(k-r-1)x^{k-r-2} + \{n(n+1) - (k-r)(k-r+1)\}x^{k-r}] a_r = 0 \quad \dots (2)$$

The relation (2) is an identity and therefore the coefficients of various powers of x can be equated to zero.

Let us first equate the coefficient of x^k the highest power of x (by putting $r=0$ in (2)) to zero; then we get

$$a_0 \{n(n+1) - k(k+1)\} = 0,$$

where a_0 being the coefficient of the first term of the series cannot be zero, i.e. $a_0 \neq 0$ and thus

$$n(n+1) - k(k+1) = 0$$

$$\text{or } n^2 + n - k^2 - k = 0$$

$$\text{or } n^2 - k^2 + (n-k) = 0$$

$$\text{or } (n-k)(n-k+1) = 0$$

which gives

$$k = n \text{ or } -n-1 \quad \dots (3)$$

Again equating the coefficient of x^{k-1} to zero, by putting $r=1$ in (2), we have

$$\{n(n+1) - (k-1)k\} a_1 = 0 \quad \dots (4)$$

$$\text{From (3), } \{n(n+1) - k(k-1)\} \neq 0$$

and therefore

$$a_1 = 0.$$

Let us now equate the coefficient of x^{k-r} , the general term in (2), to zero,

$$(k-r+2)(k-r+1)a_{r-2} + \{n(n+1) - (k-r)(k-r+1)\}a_r = 0$$

$$\text{or } a_r = -\frac{(k-r+2)(k-r+1)}{n(n+1) - (k-r)(k-r+1)} a_{r-2} \quad \dots (5)$$

Putting $k=n$, the recurrence formula is

$$\begin{aligned} a_r &= -\frac{(n-r+2)(n-r+1)}{n^2+n-(n-r)(n-r+1)} a_{r-2} \\ &= -\frac{(n-r+2)(n-r+1)}{n^2+n-n^2+nr-n+nr-r^2+r} a_{r-2} \\ &= -\frac{(n-r+2)(n-r+1)}{r(2n-r+1)} a_{r-2} \quad \dots (6) \end{aligned}$$

Again putting $k=-n-1$ in (5), we have the recurrence formula

$$\begin{aligned} \text{as } a_r &= -\frac{(-n-r+1)(-n-r)}{n^2+n-(-n-r-1)(-n-r)} a_{r-2} \\ &= -\frac{(n+r-1)(n+r)}{n^2+n-(n+r+1)(n+r)} a_{r-2} \\ &= -\frac{(n+r-1)(n+r)}{r(2n+r+1)} a_{r-2} \quad \dots (7) \end{aligned}$$

Case I. When $k=n$, we have by putting $r=2, 3, \dots$ in (6),

$$a_2 = -\frac{n(n-1)}{2(2n-1)} a_0$$

$$a_3 = -\frac{(n-1)(n-2)}{3(2n-2)} a_1$$

$$= 0 \text{ since } a_1 = 0.$$

Similarly $a_5, a_7, a_9, \dots$ etc. all the a 's having odd suffixes are zero.

$$a_4 = -\frac{(n-2)(n-3)}{4(2n-3)} a_2 = -\frac{n(n-1)(n-2)(n-3)}{2 \cdot 4(2n-1)(2n-3)} a_0 \text{ etc.}$$

(by putting value of a_2)

$$\text{In general, } a_{2r} = (-1)^r \frac{n(n-1)(n-2)(n-3)\dots(n-2r+1)}{2 \cdot 4 \dots 2r(2n-1)(2n-3)\dots(2n-2r+1)} a_0$$

(by putting value of a_2 etc.)

Hence the series solution when $k=n$, is

$$y = a_0 \left[x^n - \frac{n(n-1)}{2(2n-1)} x^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 4(2n-1)(2n-3)} x^{n-4} \dots \right] \quad \dots (8)$$

where a_0 is an arbitrary constant and is equal to

$$\frac{1 \cdot 3 \cdot 5 \dots (2n-1)}{|n|} \quad \dots (9)$$

where n is a positive integer.

This solution of Legendre's equation is known as $P_n(x)$, i.e.

$$P_n(x) = \frac{1 \cdot 3 \cdot 5 \dots (2n-1)}{|n|} \left[x^n - \frac{n(n-1)}{2(2n-1)} x^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 4(2n-1)(2n-3)} x^{n-4} - \dots \right] \quad \dots (10)$$

Case II. When $k=-n-1$, we have by putting $r=2, 3, \dots$ in (7),

$$a_2 = \frac{(n+1)(n+2)}{2(n+3)} a_0.$$

Now a_3 will contain a_1 and hence is zero. As such $a_5, a_7, a_9, \dots$ all are zero.

$$\begin{aligned} a_4 &= \frac{(n+3)(n+4)}{4(2n+5)} a_2 \\ &= \frac{(n+1)(n+2)(n+3)(n+4)}{2 \cdot 4(2n+3)(2n+5)} a_0, \text{ by putting value of } a_2 \text{ etc.} \end{aligned}$$

$$\text{In general, } a_{2r} = \frac{(n+1)\dots(n+2r)}{2 \cdot 4 \dots 2r(2n+3)\dots(2n+2r+1)} a_0.$$

Hence the series solution is

$$y = a_0 \left[x^{-n-1} + \frac{(n+1)(n+2)}{2(2n+3)} x^{-n-3} + \frac{(n+1)(n+2)(n+3)(n+4)}{2 \cdot 4(2n+3)(2n+5)} x^{-n-5} + \dots \right] \dots (11)$$

With $a_0 = \frac{|n|}{1 \cdot 3 \cdot 5 \dots (2n+1)} \dots (12)$

this solution is known as $Q_n(x)$.

$$\therefore Q_n(x) = \frac{|n|}{1 \cdot 3 \cdot 5 \dots (2n+1)} \left[x^{-n-1} + \frac{(n+1)(n+2)}{2(2n+3)} x^{-n-3} + \frac{(n+1)(n+2)(n+3)(n+4)}{2 \cdot 4(2n+3)(2n+5)} x^{-n-5} + \dots \right] \dots (13)$$

The most general solution of the Legendre's equation is

$$y = AP_n(x) + BQ_n(x) \dots (14)$$

where A and B are arbitrary constants.

Note 1. If the infinite series as the solution of a given differential equation is reduced into a finite series, then the solution is called as polynomial. $P_n(x)$ gives Legendre's polynomials or zonal harmonics or Legendre coefficients (n being a positive integer) of first kind and $Q_n(x)$ gives Legendre's polynomials of the second kind for positive integral n .

Note 2. The importance of a solution lies in its convergence and the convergency may be seen by ratio test e.g., $\lim_{n \rightarrow \infty} \left| \frac{u_{n+2}}{u_n} \right| = \frac{1}{x^2}$ from (8) or (11).

Thus the series (8) or (11) will be convergent if $|x| > 1$ i.e., the above solutions for Legendre equation are not convergent in the interval $-1 < x < 1$. In order to find the convergent solution of (1) we seek for solution in descending powers of x .

Suppose a series solution of (1) is

$$y = \sum_{r=0}^{\infty} a_r x^{k+r}, a_0 \neq 0 \dots (15)$$

So that $y' = \sum_{r=0}^{\infty} (k+r) a_r x^{k+r-1}$

$$y'' = \sum_{r=0}^{\infty} (k+r)(k+r-1) a_r x^{k+r-2}$$

Substituting these values of y , y' and y'' in (1) we get the identity

$$\sum_{r=0}^{\infty} [(k+r)(k+r-1) x^{k+r-2} - (k+r-n)(k+r+n+1) x^{k+r}] a_r = 0 \dots (16)$$

Equating to zero the coefficient of x^{k-2} (when $r=0$) the first term in (16) under the assumption $a_0 \neq 0$, yields $k(k-1)=0$ giving $k=0, 1$.

Now equating the coefficient of second term i.e. x^{k+1} to zero, we have $a_1(k+1)k=0$ giving $a_1 \neq 0$ for $k=-1$ while a_1 may or may not be zero for $k=0$.

Equating the coefficient of general term i.e. x^{k+r} to zero, we find the recursion formula $a_{r+2} = \frac{(k+r-n)(k+r+n+1)}{(k+r+2)(k+r+1)} a_r \dots (17)$

Putting $k=0$, (17) gives $a_{r+2} = \frac{(r-n)(r+n+1)}{(r+2)(r+1)} a_r \dots (18)$

And putting $k=1$, (17) gives $a_{r+2} = \frac{(1+r-n)(2+r+n)}{(3+r)(2+r)} a_r \dots (19)$

Case I. When $k=0$, we have by putting $r=0, 1, 2, 3, 4, 5, \dots$ in (18)

$$a_2 = -\frac{n(n+1)}{2} a_0 \quad \text{and} \quad a_3 = -\frac{(n-1)(n+2)}{3} a_1$$

$$a_4 = \frac{n(n-2)(n+1)(n+3)}{4} a_0 \quad a_5 = \frac{(n-1)(n-3)(n+2)(n+4)}{5} a_1$$

and in general $a_{2r} = (-1)^r \frac{n(n-2) \dots (n-2r+1)}{2r} a_0$

$$a_{2r+1} = \frac{(-1)^r (n-1)(n-3) \dots (n-2r+1)(n+2) \dots (n+2r)}{2r+1} a_1$$

Hence the series solution for $k=0$, is

$$y = a_0 \left[1 - \frac{n(n+1)}{2} x^2 + \frac{n(n-2)(n+1)(n+3)}{4} x^4 + \dots \right]$$

$$+ a_1 x \left[1 - \frac{(n-1)(n+2)}{3} x^2 + \frac{(n-1)(n-3)(n+2)(n+4)}{5} x^4 + \dots \right] \dots (19)$$

Case II. When $k=1$, we have from (19),

$$a_2 = -\frac{(n-1)(n+2)}{3} a_0 \quad \text{and} \quad a_1 = a_3 = a_5 = a_{2r+1} = \dots = 0$$

$$a_4 = \frac{(n-1)(n-3)(n+2)(n+4)}{5} a_0$$

$$a_{2r} = \frac{(-1)^r (n-1)(n-3) \dots (n-2r+1)(n+2) \dots (n+2r)}{2r+1} a_0$$

Hence the series solution for $k=1$ is

$$y = a_0 x \left[1 - \frac{(n-1)(n+2)}{3} x^2 + \frac{(n-1)(n-3)(n+2)(n+4)}{5} x^4 + \dots \right] \dots (20)$$

The solution (20) is included in (19) in the coefficient of a_1 except that a_1 is to be replaced by a_0 . Hence setting $a_1=0$ for $k=0$ also, the solution (19) reduces to

$$y = a_0 \left[1 - \frac{n(n+1)}{2} x^2 + \frac{n(n-2)(n+1)(n+3)}{4} x^4 + \dots \right] \dots (21)$$

It may be shown by ratio test that the solutions (20) and (21) are convergent in the interval $-1 < x < 1$.

Calling the solution (21) as $S_n(x)$ and (20) as $T_n(x)$, the general solution of Legendre equation in ascending powers of x is

$$y = A S_n(x) + B T_n(x) \dots (22)$$

where A and B are arbitrary constants.

[B] Legendre Polynomials.

If we put $n=2r$ (say) i.e. if n be taken as an even positive integer then (21) gives the Legendre polynomial as

$$y = a_0 \left[n - \frac{n(n+1)}{2} x^2 + \dots + (-1)^{n/2} \frac{n(n-2) \dots 4 \cdot 2 (n+1)(n+3) \dots (2n-1)}{n} x^n \right] \dots (23)$$

While (8) gives

$$y = a_0 x^n \left[1 - \frac{n(n-1)}{2(2n-1)} x^{-2} + \dots + (-1)^{n/2} \frac{n}{n(n-2) \dots 2(n+1)(n+3) \dots (2n-1)} x^{-n} \right] \dots (24)$$

(23) and (24) will be identical if (24) is multiplied by

$$(-n)^{n/2} \frac{n(n-2) \dots 4 \cdot 2 (n+1)(n+3) \dots (2n-1)}{n}$$

and then the solution (8), (21) will become identical.

Again if we take, n as an odd negative integer then (21) is identical with (11). Also if n is an odd positive integer then (20) reduces to

$$y = a_0 \left[x - \frac{(n-1)(n+2)}{3} x^3 + \dots + (-1)^{(n-1)/2} \frac{(n-1)(n-3) \dots 2(n+2) \dots (2n-1)}{n} x^n \right] \dots (25)$$

and (8) reduces to

$$y = a_0 x^n \left[1 - \frac{n(n-1)}{2(2n-1)} x^{-2} + \dots + (-1)^{(n-1)/2} \frac{n}{2 \cdot 4 \dots (n-1)(n-3) \dots (2n-1)} x^{-n+1} \right] \dots (26)$$

which become identical when multiplied by the coefficient of last term in (26).

Further if n is an even negative integer, (20) and (11) become identical.

These discussions follow the conclusions :

(i) For integral values for n , the solutions (8) and (11) have great utility of them.

(ii) For positive integral n , (8) is a polynomial but (11) is an infinite series and the complete integral is a linear combination of them,

(iii) For negative integral n , (8) is an infinite series and (11) is a polynomial.

(iv) For positive integral n , in (9) or (10) we have chosen

$$a_0 = \frac{1 \cdot 3 \cdot 5 \dots (2n+1)}{n}$$

(v) For negative integral n , in (12) or (13) we have chosen

$$a_0 = \frac{n}{1 \cdot 3 \cdot 5 \dots (2n+1)}$$

(vi) For positive integral n , the polynomial $P_n(x)$ has the expansion given by (10) ending with term free from x i.e.

$$\begin{aligned} P_n(x) &= \sum_{r=0}^{n/2} (-1)^r \frac{1 \cdot 3 \cdot 5 \dots (2n-1)}{n} \frac{n(n-1) \dots (n-2r+1)}{2 \cdot 4 \dots 2r(2n-1)(2n-3) \dots (2n-2r+1)} x^{n-2r} \\ &= \sum_{r=0}^{n/2} (-1)^r \frac{1 \cdot 3 \cdot 5 \dots (2n-2r-1)}{2^r r! (n-2r)} x^{n-2r} \dots (27) \end{aligned}$$

or more concisely

$$P_n(x) = \sum_{r=0}^{n/2} (-1)^r \frac{1 \cdot 2n-2r}{2^n r! (n-r)! (n-2r)!} x^{n-2r} \dots (28)$$

Evaluation of the values of $P_0(\mu)$, $P_1(\mu)$, $P_2(\mu)$, $P_3(\mu)$, $P_4(\mu)$, $P_5(\mu)$, $P_6(\mu)$, and $P_7(\mu)$ etc.

We have

$$\begin{aligned} P_n(\mu) &= \frac{1 \cdot 3 \cdot 5 \dots (2n-1)}{n!} \left[\mu^n - \frac{n(n-1)}{2(2n-1)} \mu^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 4 \cdot (2n-1)(2n-3)} \mu^{n-4} - \dots \right] \\ &= \frac{(2n)!}{(2 \cdot 4 \cdot 6 \dots 2n)n!} \left[\mu^n - \frac{n(n-1)}{2(2n-1)} \mu^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 4 \cdot (2n-1)(2n-3)} \mu^{n-4} - \dots \right] \end{aligned}$$

$$= \frac{(2n)!}{2^n n! n!} \left[\mu^n - \frac{n(n-1)}{2(2n-1)} \mu^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 4 \cdot (2n-1)(2n-3)} \mu^{n-4} \dots \right]$$

Putting $n=0, 1, 2, 3, 4, 5, 6, 7$ etc., we have

$$P_0(\mu)=1,$$

$$P_1(\mu) = \frac{2!}{2 \cdot 1! \cdot 1!} [\mu], \text{ other terms vanishing} \\ = \mu,$$

$$P_2(\mu) = \frac{4!}{2^2 \cdot 2! \cdot 2!} [\mu^2 - \frac{1}{2}] \text{ other terms vanishing} \\ = \frac{4 \cdot 3 \cdot 2}{2 \cdot 2 \cdot 2 \cdot 2} \frac{3\mu^2 - 1}{3} = \frac{3\mu^2 - 1}{2}.$$

$$P_3(\mu) = \frac{6!}{2^3 \cdot 3! \cdot 3!} \left[\mu^3 - \frac{3 \cdot 2}{3 \cdot 5} \mu \right] = \frac{5\mu^3 - 3\mu}{2}.$$

$$\text{Similarly } P_4(\mu) = \frac{1}{8} (35\mu^4 - 30\mu^2 + 3),$$

$$P_5(\mu) = \frac{1}{8} (63\mu^5 - 70\mu^3 + 15\mu),$$

$$P_6(\mu) = \frac{1}{8} (231\mu^6 - 315\mu^4 + 105\mu^2 - 5)$$

$$P_7(\mu) = \frac{1}{8} (429\mu^7 - 693\mu^5 + 315\mu^3 - 35\mu) \text{ etc.}$$

[B] Generating Functions for $P_n(x)$

$P_n(\mu)$ is the coefficient of h^n in $(1-2\mu h+h^2)^{-1/2}$.

We have $(1-2\mu h+h^2)^{-1/2}$

$$= [1-h(2\mu-h)]^{-1/2}$$

$$= 1 + \frac{1}{2}h(2\mu-h) + \frac{1.3}{2.4}h^2(2\mu-h^2) + \frac{1.3.5}{2.4.6}h^3(2\mu-h)^3 + \dots$$

$$+ \frac{1.3.5 \dots (2n-1)}{2.4.6 \dots 2n} h^n (2\mu-h)^n + \dots$$

(by Binomial expansion)

The coefficient of h^n in the expansion

$$= \frac{1.3.5 \dots (2n-1)}{2.4.6 \dots 2n} (2\mu)^n = \frac{1.3.5 \dots (2n-3)}{2.4.6 \dots (2n-2)} (2\mu)^{n-2} \cdot {}^{n-1}C_1$$

$$+ \frac{1.3.5 \dots (2n-5)}{2.4.6 \dots (2n-4)} (2\mu)^{n-4} \cdot {}^{n-2}C_2 + \dots$$

$$= \frac{1.3.5 \dots (2n-1)}{n!} \left[\mu^n - \frac{n(n-1)}{2(2n-1)} \mu^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 2 \cdot 2! (2n-1)(2n-4)} \mu^{n-4} - \dots \right]$$

$$= \frac{1.3.5 \dots (2n-1)}{n!} \left[\mu^n - \frac{n(n-1)}{2(2n-1)} \mu^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 4 \cdot (2n-1)(2n-3)} \mu^{n-4} + \dots \right],$$

which is $P_n(\mu)$.

$$\text{Hence } \sum_{n=0}^{\infty} h^n P_n(\mu) = (1-2\mu h+h^2)^{-1/2}. \quad \dots (30)$$

(Agra, 1967)

COROLLARY 1. $P_n(1)$ is the coefficient of h^n in $(1-2h+h^2)^{-1/2}$, i.e., in $(1-h)^{-1}$.

As such $P_n(1)$ = coeff. of h^n in $\{1+h+h^2+\dots+h^n+\dots\}=1$, which is however a distinguishing property of Legendre's Polynomials.

COROLLARY 2. Putting $\mu = \cos \theta$, we have

$$\sum_{n=0}^{\infty} h^n P_n(\cos \theta) = \{1-2(\cos \theta)h+h^2\}^{-1/2} \\ = \{1-(e^{i\theta}+e^{-i\theta})h+e^{i\theta} \cdot e^{-i\theta} h^2\}^{-1/2} \\ = (1-e^{i\theta})^{-1/2} (1-he^{-i\theta})^{-1/2} \\ = \left[1 + \frac{1}{2}he^{i\theta} + \frac{1.3}{2.4}h^2e^{2i\theta} + \frac{1.3.5}{2.4.6}h^3e^{3i\theta} + \dots \right. \\ \left. + \frac{1.3 \dots (2n-1)}{2.4 \dots 2n} h^n e^{in\theta} + \dots \right] \\ \times \left[1 + \frac{1}{2}he^{-i\theta} + \frac{1.3}{2.4}h^2e^{-2i\theta} + \frac{1.3.5}{2.4.6}h^3e^{-3i\theta} + \dots \right. \\ \left. + \frac{1.3.5 \dots (2n-1)}{2.4 \dots 2n} h^n e^{-ni\theta} + \dots \right]$$

Equating coefficients of h^n on either side,

$$P_n(\cos \theta) = \frac{1.3.5 \dots (2n-1)}{2.4.6 \dots 2n} (e^{ni\theta} + e^{-ni\theta}) \\ + \frac{1}{2} \frac{1.3.5 \dots (2n-3)}{2.4.6 \dots (2n-2)} \{e^{(n-2)i\theta} + e^{-(n-2)i\theta}\} \\ + \frac{1.2}{2.4} \frac{1.3.5 \dots (2n-5)}{2.4.6 \dots (2n-4)} \{e^{(n-4)i\theta} + e^{-(n-4)i\theta} + \dots \\ = 2 \cdot \frac{1.3.5 \dots (2n-1)}{2.4 \dots 2n} \left[\cos n\theta + \frac{1}{2} \frac{2}{2n-1} \cos (n-2)\theta \right. \\ \left. + \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{2n(2n-2)}{(2n-1)(2n-3)} \cos (n-4)\theta + \dots \right] \\ = 2 \cdot \frac{1.3.5 \dots (2n-1)}{2.4.6 \dots 2n} \left[\cos n\theta + \frac{n}{2n-1} \cos (n-2)\theta \right. \\ \left. + \frac{1.3}{2!} \frac{n(n-1)}{(2n-1)(2n-3)} \cos (n-4)\theta + \dots \right]. \quad \dots (31)$$

COROLLARY 3. We know that

$$\begin{aligned}
 & (1-2h \cos \theta + h^2)^{-1/2} \\
 &= \{(1-h)^2 + 2h(1-\cos \theta)\}^{-1/2} \\
 &= \left\{ (1-h)^2 + 4h \sin^2 \frac{\theta}{2} \right\}^{-1/2} \\
 &= \frac{1}{1-h} \left\{ 1 - \frac{(-1) 4h \sin^2 \theta/2}{(1-h)^2} \right\}^{-1/2} \\
 &= \frac{1}{1-h} \left[1 + \frac{1}{2} \left\{ -\frac{4h \sin^2 \theta/2}{(1-h)^2} \right\} + \frac{1.3}{2.4} \left\{ -\frac{4h \sin^2 \theta/2}{(1-h)^2} \right\}^2 + \dots \right. \\
 &\quad \left. \dots + \frac{1.3.5 \dots (2r-1)}{2.4 \dots 2r} \left\{ -\frac{4h \sin^2 \theta/2}{(1-h)^2} \right\}^r + \dots \right] \\
 &= \frac{1}{1-h} + \frac{1}{2} \left[\left\{ -\frac{4h \sin^2 \theta/2}{(1-h)^2} \right\} \frac{1}{1-h} + \dots \right. \\
 &\quad \left. \dots + \frac{1.3.5 \dots (2r-1) (-1) 4^r h^r \sin^{2r} \theta/2}{2.4 \dots 2r (1-h)^{2r+1}} + \dots \right] \\
 &= \frac{1}{1-h} + \sum_{r=1}^{\infty} (-1)^r \frac{1.3.5 \dots (2r-1)}{2.4 \dots 2r} \frac{2^r \cdot 2^r h^r \sin^{2r} \theta/2}{(1-h)^{2r+1}}
 \end{aligned}$$

Equating coefficients of h^n on either side,

$$\begin{aligned}
 P_n(\cos \theta) &= 1 + \sum_{r=1}^{\infty} (-1)^r \frac{1.3.5 \dots (2r-1)}{2.4 \dots 2r} \cdot 2^r \cdot 2^r \sin^{2r} \frac{\theta}{2} \\
 &\quad \times \frac{(2r+1)(2r+2) \dots \{2r+1+(n-r-1)\}}{(n-1)!}
 \end{aligned}$$

$$\left[\text{since coeff. of } h^{n-1} \text{ in } (1-h)^{-(2r+1)} \text{ is } \frac{(2r+1)(2r+2) \dots (2r+1+n-r-1)}{(n-r)!} \text{ etc.} \right]$$

$$\begin{aligned}
 \therefore P_n(\cos \theta) &= 1 + \sum_{r=1}^{\infty} (-1)^r \frac{1.3.5 \dots (2r-1)}{2.4 \dots 2r} \cdot \frac{(n+r)!}{(2r)!(n-r)!} \cdot 2^r \cdot 2^r \sin^{2r} \frac{\theta}{2} \\
 &= 1 + \sum_{r=1}^{\infty} (-1)^r \frac{(2r)!(n-r)!(n-r+1)(n-r+2) \dots (n+r)}{2^r r! \cdot 2^r r! (2r)!(n-r)!} \\
 &\quad \times 2^r \cdot 2^r \sin^{2r} \frac{\theta}{2} \\
 &= 1 + \sum_{r=1}^{\infty} (-1)^r \frac{(n-r+1)(n-r+2) \dots (n+r)}{r! r!} \sin^{2r} \frac{\theta}{2} \\
 &= 1 - \frac{n(n+1)}{1!1!} \sin^2 \frac{\theta}{2} + \frac{(n-1)n(n+1)(n+2)}{2!2!} \sin^4 \frac{\theta}{2} \\
 &\quad - \frac{(n-2)(n-1)n(n+1)(n+2)(n+3)}{3!3!} \sin^6 \frac{\theta}{2} + \dots
 \end{aligned}$$

$$\begin{aligned}
 &= 1 + \frac{(-n)(n+1)}{1.1} \sin^2 \frac{\theta}{2} + \frac{(-n)(-n+1)(n+1)(n+2)}{1.2.1(1+1)} \left(\sin^2 \frac{\theta}{2} \right)^2 \\
 &\quad + \frac{(-n)(-n+1)(-n+2)(n+1)(n+2)(n+3)}{1.2.3.1(1+1)(1+2)} \left(\sin^2 \frac{\theta}{2} \right)^3 + \dots \\
 &= F\left(-n, n+1, 1, \sin^2 \frac{\theta}{2}\right) \dots (32)
 \end{aligned}$$

Replacing θ by $\theta + \pi$.

$$P_n(-\cos \theta) = (-1)^n P_n(\cos \theta) = F\left(-n, n+1, 1, \cos^2 \frac{\theta}{2}\right) \dots (33)$$

i.e. $P_n(\cos \theta) = (-1)^n \cdot F(-n, n+1, 1, \cos^2 \theta/2)$.

(C) Integral for $P_n(\mu)$.

$$I. P_n(\mu) = \frac{1}{\pi} \int_0^\pi \{\mu \mp \sqrt{(\mu^2-1) \cos \phi}\}^n d\phi. \quad (\text{Vikram, 1962})$$

It is easy to show that

$$\int_0^\pi \frac{d\phi}{a \pm b \cos \phi} = \frac{\pi}{\sqrt{a^2 - b^2}}, \text{ when } a^2 > b^2.$$

Let us put $a = 1 - \mu h$, $b = h\sqrt{(\mu^2-1)}$, so that $a^2 - b^2 = 1 - 2\mu h + h^2$.

$$\begin{aligned}
 \text{Then } \frac{\pi}{(1-2\mu h + h^2)^{-1/2}} &= \int_0^\pi \frac{d\phi}{1 - h\mu \pm h\sqrt{(\mu^2-1) \cos \phi}} \\
 \text{or } (1-2\mu h + h^2)^{-1/2} &= \frac{1}{\pi} \int_0^\pi [1 - h\{\mu \mp \sqrt{(\mu^2-1) \cos \phi}\}]^{-1} d\phi \\
 &= \frac{1}{\pi} \int_0^\pi \left[\sum_{n=0}^{\infty} h^n \{\mu \mp \sqrt{(\mu^2-1) \cos \phi}\}^n \right] d\phi \\
 &\quad \text{where } h \text{ is sufficiently small} \\
 &= \frac{1}{\pi} \sum_{n=0}^{\infty} h^n \int_0^\pi [\mu \mp \sqrt{(\mu^2-1) \cos \phi}]^n d\phi.
 \end{aligned}$$

Equating the coefficients of h^n on either side, we have

$$P_n(\mu) = \frac{1}{\pi} \int_0^\pi \{\mu \mp \sqrt{(\mu^2-1) \cos \phi}\}^n d\phi, \dots (34)$$

which is known as first of the Laplace's Integrals.

COROLLARY 4. If we put $\mu = \cos \theta$.

$$P_n(\cos \theta) = \frac{1}{\pi} \int_0^\pi (\cos \theta \mp i \sin \theta \cos \phi)^n d\phi. \dots (35)$$

$$\text{COROLLARY 5. } P_n(\mu) = \frac{1}{\pi} \int_0^\pi \frac{d\phi}{\{\mu \pm \sqrt{(\mu^2-1) \cos \phi}\}^{n+1}}.$$

We have as above

$$\int_0^\pi \frac{d\phi}{a - b \cos \phi} = \frac{\pi}{\sqrt{a^2 - b^2}}, \text{ where } a^2 > b^2.$$

Putting $a = \mu h - 1$, $b = \mp h\sqrt{(\mu^2-1)}$, we get

$$\frac{\pi}{\sqrt{(1-2\mu h + h^2)}} = \int_0^\pi \frac{d\phi}{h \{\mu \pm \sqrt{(\mu^2-1) \cos \phi}\} - 1}$$

$$\begin{aligned}
 \text{or } & \frac{\pi}{h \sqrt{\left(1 - \frac{2\mu}{h} + \frac{1}{h^2}\right)}} \\
 &= \int_0^\pi \frac{d\phi}{h \{\mp \sqrt{(\mu^2 - 1) \cos \phi} \left[1 - \frac{1}{h \{\mu \pm \sqrt{(\mu^2 - 1) \cos \phi}\}} \right]}} \\
 \text{or } & \sum_{n=0}^{\infty} \frac{1}{h^{n+1}} P_n(\mu) = \frac{1}{\pi} \int_0^\pi \left[\frac{1 - \frac{1}{h \{\mu \pm \sqrt{(\mu^2 - 1) \cos \phi}\}}}{h \{\mu \pm \sqrt{(\mu^2 - 1) \cos \phi}\}} \right] d\phi \\
 &= \frac{1}{\pi} \int_0^\pi \left[\sum_{n=0}^{\infty} \frac{1}{h^{n+1} \{\mu \pm \sqrt{(\mu^2 - 1) \cos \phi}\}^{n+1}} \right] d\phi \\
 &= \frac{1}{\pi} \sum_{n=0}^{\infty} \int_0^\pi \frac{1}{h^{n+1} \{\mu \pm \sqrt{(\mu^2 - 1) \cos \phi}\}^{n+1}} d\phi
 \end{aligned}$$

Equating coefficients of $\frac{1}{h^{n+1}}$ on either side, we get

$$P_n(\mu) = \frac{1}{\pi} \int_0^\pi \frac{d\phi}{\{\mu \pm \sqrt{(\mu^2 - 1) \cos \phi}\}^{n+1}} \quad \dots (36)$$

which is known as **Second of the Laplace's Integrals**.

COROLLARY 6. It is obvious from the above two Laplace's integrals that $P_n(\mu) = P_{-(n+1)}(\mu)$. $\dots (37)$

[D] Recurrence Formulae for $P_n(\mu)$.

$$I. nP_n(\mu) = (2n-1)\mu P_{n-1}(\mu) - (n-1)P_{n-2}(\mu). \quad \dots (38)$$

(Agra, 1953, 61, 74; Vikram, 1963)

$$\text{Suppose } V = (1 - 2\mu h + h^2)^{-1/2} = \sum_{n=0}^{\infty} h^n P_n(\mu)$$

$$\text{or } V^2(1 - \mu h + h^2) = 1.$$

Differentiating w.r.t. h .

$$2V \cdot \frac{dV}{dh} (1 - 2\mu h + h^2) + V^2(2h - 2\mu) = 0$$

$$\text{or } \frac{dV}{dh} (1 - 2\mu h + h^2) + V(h - \mu) = 0$$

$$\text{or } (1 - 2\mu h + h^2) \sum_{n=0}^{\infty} nh^{n-1} P_n(\mu) + (h - \mu) \sum_{n=0}^{\infty} h^n P_n(\mu) = 0$$

$$\left[\because \frac{dV}{dh} = \sum_{n=0}^{\infty} nh^{n-1} P_n(\mu) \right]$$

The coefficient of h^{n-1} equated to zero gives

$$nP_n(\mu) - 2\mu(n-1)P_{n-1}(\mu) + (n-2)P_{n-2}(\mu) + P_{n-2}(\mu) - \mu P_{n-1}(\mu) = 0$$

$\dots (38)$

$$\text{or } nP_n(\mu) = (2n-1)\mu P_{n-1}(\mu) - (n-1)P_{n-2}(\mu).$$

Note. If $n-1=m$, then this result reduces to $(m+1)P_{m+1}(\mu) = (2m+1)\mu P_m(\mu) - mP_{m-1}(\mu)$ $\dots (39)$
(Agra, 65; Vikram, 63)

$$II. (\mu^2 - 1) \frac{dP_n(\mu)}{d\mu} = n \{\mu P_n(\mu) - P_{n-1}(\mu)\} \\ = -(n+1) \{\mu P_n(\mu) - P_{n+1}(\mu)\}.$$

We have $\mu P_n(\mu) - P_{n+1}(\mu)$

$$\begin{aligned}
 &= \frac{\mu}{\pi} \int_0^\pi \{\mu + \sqrt{(\mu^2 - 1) \cos \phi}\}^n d\phi - \frac{1}{\pi} \int_0^\pi \{\mu + \sqrt{(\mu^2 - 1) \cos \phi}\}^{n+1} d\phi \\
 &= \frac{1}{\pi} \int_0^\pi \{\mu + \sqrt{(\mu^2 - 1) \cos \phi}\}^{n-1} [\mu \{\mu + \sqrt{(\mu^2 - 1) \cos \phi}\} - 1] d\phi \\
 &= \frac{\mu^2 - 1}{\pi} \int_0^\pi \{\mu + \sqrt{(\mu^2 - 1) \cos \phi}\}^{n-1} \left[1 + \frac{\mu \cos \phi}{\sqrt{(\mu^2 - 1)}} \right] d\phi \\
 &= \frac{\mu^2 - 1}{\pi} \frac{d}{d\mu} \int_0^\pi \frac{\{\mu + \sqrt{(\mu^2 - 1) \cos \phi}\}^n}{n} d\phi \\
 &= \frac{\mu^2 - 1}{n} \frac{d}{d\mu} P_n(\mu).
 \end{aligned}$$

$$\therefore (\mu^2 - 1) \frac{dP_n(\mu)}{d\mu} = n \{\mu P_n(\mu) - P_n(\mu)\}. \quad \dots (40)$$

(Agra, 1965)

Replacing n by $-(n-1)$, as $P_n(\mu) = P_{-(n+1)}(\mu)$ and as such $P_{-(n+2)}(\mu) = P_{n+1}(\mu)$, we get

$$(\mu^2 - 1) \frac{dP_n(\mu)}{d\mu} = -(n+1) \{\mu P_n(\mu) - P_{n+1}(\mu)\} \quad \dots (41)$$

III. By Legendre's equation, we have

$$(\mu^2 - 1) \frac{d^2 P_n(\mu)}{d\mu^2} + 2\mu \frac{dP_n(\mu)}{d\mu} - n(n+1)P_n(\mu) = 0.$$

This may be written as

$$\frac{d}{d\mu} \left\{ (\mu^2 - 1) \frac{dP_n(\mu)}{d\mu} \right\} = n(n+1)P_n(\mu)$$

$$\text{or } \frac{d}{d\mu} [n \{\mu P_n(\mu) - P_{n-1}(\mu)\}] = n(n+1)P_n(\mu) \text{ from (40)}$$

$$\text{or } \mu \frac{dP_n(\mu)}{d\mu} + P_n(\mu) - \frac{dP_{n-1}(\mu)}{d\mu} = (n+1)P_n(\mu)$$

$$\text{or } \mu \frac{dP_n(\mu)}{d\mu} - \frac{dP_{n-1}(\mu)}{d\mu} = nP_n(\mu). \quad \dots (41)$$

(Agra, 1954; Vikram, 62)

IV. Replacing n by $-(n+1)$ and applying $P_{-n-1}(\mu) = P_n(\mu)$, $P_{-n-2}(\mu) = P_{n+1}(\mu)$ etc., in (41) we get

$$-\mu \frac{dP_n(\mu)}{d\mu} + \frac{dP_{n+1}(\mu)}{d\mu} = (n+1)P_n(\mu). \quad \dots (42)$$

Addition of (41) and (42) yields

$$(2n+1)P_n(\mu) = \frac{d}{d\mu}P_{n+1}(\mu) - \frac{d}{d\mu}P_{n-1}(\mu). \quad \dots (43)$$

(Agra, 1952; 55, 61, 74; Vikram, 62, 63)

Now (43) may be written as

$$\frac{d}{d\mu}P_{n+1}(\mu) = (2n+1)P_n(\mu) + \frac{d}{d\mu}P_{n-1}(\mu).$$

Replacing n by $(n-1)$, we get

$$\begin{aligned} \frac{d}{d\mu}P_n(\mu) &= (2n-1)P_{n-1}(\mu) + \frac{d}{d\mu}P_{n-2}(\mu) \\ &= (2n-1)P_{n-1}(\mu) + (2n-5)P_{n-3}(\mu) + \frac{d}{d\mu}P_{n-4}(\mu) \quad \dots (44) \end{aligned}$$

[since by putting $n-2$ for n ,

$$\frac{dP_{n-2}(\mu)}{d\mu} = (2n-5)P_{n-3}(\mu) + \frac{d}{d\mu}P_{n-4}(\mu)]$$

The repeated application of this replacement will give

$$\frac{d}{d\mu}P_n(\mu) = (2n-1)P_{n-1}(\mu) + (2n-5)P_{n-3}(\mu) + (2n-9)P_{n-5}(\mu) + \dots$$

ending with $3P_1(\mu)$ or $P_0(\mu)$ according as n is even or odd. $\dots (48)$

This is known as *Christoffel's Expansion*.

Again multiplying (40) by $(n+1)$, (41) by n and adding, we have

$$(2n+1)(\mu^2-1)\frac{dP_n(\mu)}{d\mu} = n(n+1)\{P_{n+1}(\mu) - P_{n-1}(\mu)\} \quad \dots (49)$$

[E] Some Important Results.

$$[e_1]. P_n(-\mu) = (-1)^n P_n(\mu). \quad \dots (50)$$

(Agra, 1967)

We know that

$$\begin{aligned} P_n(\mu) &= \frac{1.3.5\dots(2n-1)}{2^n |n|} \left[\mu^n - \frac{n(n-1)}{2 \cdot (2n-1)} \mu^{n-2} + \dots \right] \\ &= \frac{|(2n)|}{2.4.6\dots 2n} |n| \left[\mu^n - \frac{n(n-1)}{2(2n-1)} \mu^{n-2} + \dots \right] \\ &= \frac{|(2n)|}{2^n |n|} \left[\mu^n - \frac{n(n-1)}{2(2n-1)} \mu^{n-2} + \dots \right] \end{aligned}$$

$$\therefore P_n(-\mu) = \frac{|(2n)|}{2^n |n|} \left[(-\mu)^n - \frac{n(n-1)}{2(2n-1)} (-\mu)^{n-2} + \dots \right]$$

$$\begin{aligned} \text{or } P_n(-\mu) &= \frac{|(2n)|}{2^n |n|} \left[(-1)^n \mu^n - \frac{n(n-1)}{2(2n-1)} (-1)^{n-2} \mu^{n-2} + \dots \right] \\ &= \frac{|(2n)|}{2^n |n|} \left[(-1)^n \mu^n - \frac{n(n-1)}{2(2n-1)} (-1)^n \mu^{n-2} + \dots \right] \end{aligned}$$

$$\begin{aligned} &= (-1)^n \cdot \frac{|(2n)|}{2^n |n|} \left[\mu^n - \frac{n(n-1)}{2(2n-1)} \mu^{n-2} + \dots \right] \\ &= (-1)^n P_n(\mu). \end{aligned}$$

Note. If $n=2m$, this gives $P_{2m}(-\mu) = P_{2m}(\mu)$. (Agra, 1965)

If $n=2m+1$, we have

$$P_{2m+1}(-\mu) = -P_{2m+1}(\mu). \quad \dots (51)$$

(Agra, 65)

[e₂] Rodrigue's formula

$$P_n(\mu) = \frac{1}{2^n |n|} \left(\frac{d}{d\mu} \right)^n (\mu^2-1)^n. \quad \dots (52)$$

(Rodrigue's formula)
(Agra, 1656, 60, 63; Vikram 63, 64)

Let $y = (\mu^2-1)^n$.

$$\therefore \frac{dy}{d\mu} = n(\mu^2-1)^{n-1} \cdot 2\mu = \frac{n \cdot y \cdot 2\mu}{\mu^2-1},$$

$$\text{i.e.} \quad (\mu^2-1) \frac{dy}{d\mu} = 2ny\mu.$$

Differentiating this $(n+1)$ times by Leibnitz's theorem, we get

$$\begin{aligned} (\mu^2-1) \frac{d^{n+2}y}{d\mu^{n+2}} + {}^{n+1}C_1 (2\mu) \cdot \frac{d^{n+1}y}{d\mu^{n+1}} + {}^{n+1}C_2 (2) \cdot \frac{d^ny}{d\mu^n} \\ = 2n \left[\mu \frac{d^{n+1}y}{d\mu^{n+1}} + {}^{n+1}C_1 (1) \frac{d^ny}{d\mu^n} \right] \end{aligned}$$

$$\text{or } (\mu^2-1) \frac{d^{n+2}y}{d\mu^{n+2}} + 2\mu \frac{d^{n+1}y}{d\mu^{n+1}} - n(n+1) \frac{d^ny}{d\mu^n} = 0.$$

Putting $\frac{d^ny}{d\mu^n} = z$, we have

$$(\mu^2-1) \frac{d^2z}{d\mu^2} + 2\mu \frac{dz}{d\mu} - n(n+1)z = 0,$$

which is Legendre's equation, one of whose solution is z and hence it is satisfied by Cz or $C \frac{d^ny}{d\mu^n}$, where C is an arbitrary constant.

$$\begin{aligned} \text{Hence } P_n(\mu) &= C \frac{d^ny}{d\mu^n} = C \left(\frac{d}{d\mu} \right)^n (\mu^2-1)^n \\ &= C \left(\frac{d}{d\mu} \right)^n (\mu-1)^n (\mu+1)^n \\ &= C [\mu(\mu+1)^n + \text{terms containing } (\mu-1) \text{ as one of the factors}] \end{aligned}$$

on differentiating n times by Leibnitz's theorem.

$$\therefore P_n(1) = C.2^n |n|, \text{ where } P_n(1) = 1; \therefore C = \frac{1}{2^n |n|}$$

$$\text{Hence } P_n(\mu) = \frac{1}{2^n |n|} \left(\frac{d}{d\mu} \right)^n (\mu^2-1)^n.$$

We know that

$$P_n(\mu) = \frac{1.3.5 \dots (2n-1)}{n!} \left[\mu^n - \frac{n(n-1)}{2(2n-1)} \mu^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2.4.(2n-1)(2n-3)} \mu^{n-4} + \dots \right]$$

$$\therefore P_{2m+1}(\mu) = \frac{1.3.5 \dots \{2(2m+1)-1\}}{(2m+1)!} \times \left[\mu^{2m+1} - \frac{(2m+1)(2m+1-1)}{2 \{2(2m+1)-1\}} \mu^{2m-1} + \dots \right]$$

$$\text{so that } P_{2m+1}(0) = \frac{1.3.5 \dots (4m+1)}{(2m+1)!} [0] = 0.$$

Again, we have

$$1 - 2\mu h + h^2)^{-1/2} = \sum_{r=0}^{\infty} h^r P_r(\mu)$$

$$= P_0(\mu) + h P_1(\mu) + h^2 P_2(\mu) + \dots + h^{2m} P_{2m}(\mu) + \dots$$

Putting $\mu=0$, we get

$$\begin{aligned} P_0(0) + h P_1(0) + h^2 P_2(0) + \dots + h^{2m} P_{2m}(0) \\ = (1 + h^2)^{-1/2} \\ = \{1 - (-h^2)\}^{-1/2} \end{aligned}$$

$$= 1 + \frac{1}{2} (-h^2) + \frac{\frac{1}{2} \cdot \frac{3}{2} (-h^2)^2}{2} + \dots + \frac{\frac{1}{2} \cdot \frac{3}{2} \dots \left(\frac{2r-1}{2} \right)}{r!} (-h^2)^r + \dots$$

expanding by Binomial Theorem.

Equating the coefficients of x^{2m} on either side.

$$P_{2m}(0) = (-1)^m \frac{1.3.5 \dots (2m-1)}{2^m m!} = (-1)^m \frac{1.3.5 \dots (2m-1)}{2.4.6 \dots 2m}$$

Problem 14. Show that the Legendre's equation

$$(1-\mu^2) \frac{d^2 y}{d\mu^2} - 2\mu \frac{dy}{d\mu} + n(n+1)y = 0$$

changes into the hypergeometric form

$$x(1-x) \frac{d^2 y}{dx^2} + \left(\frac{1}{2} - \frac{3x}{2} \right) \frac{dy}{dx} + \frac{n(n+1)}{4} y = 0.$$

Hence show by comparison that its complete primitive is

$$AF\left(-\frac{n}{2}, \frac{n+1}{2}, \frac{1}{2}, \mu^2\right) + B\mu F\left(-\frac{n-1}{2}, \frac{n+2}{2}, \frac{3}{4}, \mu^2\right)$$

$$\text{Given } \mu^2 = x; \therefore 2\mu = \frac{dx}{d\mu}.$$

$$\text{Now } \frac{dy}{d\mu} = \frac{dy}{dx} \cdot \frac{dx}{d\mu} = 2\mu \cdot \frac{dy}{dx}.$$

$$\begin{aligned} \frac{d^2 y}{d\mu^2} &= \frac{d}{d\mu} \left(2\mu \frac{dy}{dx} \right) \\ &= 2 \frac{dy}{dx} + 2\mu \frac{d^2 y}{dx^2} \frac{dx}{d\mu} \\ &= 2 \frac{dy}{dx} + 4\mu^2 \frac{d^2 y}{dx^2} \end{aligned}$$

Substituting these values in the given Legendre's equation, we

$$(1-x) \left(4x \frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} \right) - 4x \frac{dy}{dx} + n(n+1)y = 0$$

$$x(1-x) \frac{d^2 y}{dx^2} + \frac{1-x-2x}{2} \frac{dy}{dx} + \frac{n(n+1)}{4} y = 0$$

$$x(1-x) \frac{d^2 y}{dx^2} + \left(\frac{1}{2} - \frac{3x}{2} \right) \frac{dy}{dx} + \frac{n(n+1)}{4} y = 0.$$

Again, we have already proved in $E(e_3)$ of § 8.3 that the complete primitive of the hypergeometric series

$$x(1-x) \frac{d^2 y}{dx^2} + [\gamma - (\alpha + \beta + 1)] \frac{dy}{dx} - \alpha\beta y = 0$$

$$AF(\alpha, \beta, \gamma, x) + Bx^{1-\gamma} F(\alpha+1-\gamma, \beta+1-\gamma, 2-\gamma, x).$$

In the existing case, by comparison, we have

$$\gamma = \frac{1}{2}, \quad \alpha + \beta + 1 = \frac{3}{2}, \quad \alpha\beta = -\frac{n(n+1)}{4}$$

$$\gamma = \frac{1}{2}, \quad \alpha + \beta = \frac{1}{2}, \quad \alpha\beta = -\frac{n(n+1)}{4}$$

Now

$$(\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$$

$$= \frac{1}{4} + n(n+1)$$

$$= \frac{1}{4} + n^2 + n = \left(n + \frac{1}{2} \right)^2$$

$$\alpha - \beta = n + \frac{1}{2} \text{ and } \alpha + \beta = \frac{1}{2}$$

Solving

$$\alpha = \frac{n+1}{2}, \quad \beta = -\frac{n}{2}.$$

Hence by comparison the complete primitive of the transformed hypergeometric series is

$$AF\left(\frac{n+1}{2}, -\frac{n}{2}, \frac{1}{2}, \mu^2\right) + B\mu F\left(\frac{n+2}{2}, -\frac{n-1}{2}, \frac{3}{2}, \mu^2\right)$$

Problem 15. Show that

$$P_n(\cos \theta) = \cos^{2n} \frac{\theta}{2} F\left(-n, -n, 1, -\tan^2 \frac{\theta}{2}\right)$$

We know from the first Laplace's Integral that

$$P_n(\mu) = \frac{1}{\pi} \int_0^\pi \{\mu \pm \sqrt{\mu^2 - 1} \cos \phi\}^n d\phi.$$

$\therefore P_n(\cos \theta)$

$$= \frac{1}{\pi} \int_0^\pi (\cos \theta + i \sin \theta \cos \phi)^n d\phi \text{ (taking positive sign)}$$

$$= \frac{1}{\pi} \int_0^\pi \left[\cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2} + 2i \sin \frac{\theta}{2} \cos \frac{\theta}{2} \left(\frac{e^{i\phi} + e^{-i\phi}}{2} \right) \right]^n d\phi$$

$$= \frac{1}{\pi} \int_0^\pi \left[\cos^2 \frac{\theta}{2} + i \sin \frac{\theta}{2} \cos \frac{\theta}{2} e^{i\phi} + i^2 \sin^2 \frac{\theta}{2} e^{i\phi} e^{-i\phi} + i \sin \frac{\theta}{2} \cos \frac{\theta}{2} e^{-i\phi} \right]^n d\phi$$

$$= \frac{1}{\pi} \int_0^\pi \left[\cos \frac{\theta}{2} \left\{ \cos \frac{\theta}{2} + i \sin \frac{\theta}{2} e^{i\phi} \right\} + i \sin \frac{\theta}{2} e^{-i\phi} \left\{ \cos \frac{\theta}{2} + i \sin \frac{\theta}{2} e^{i\phi} \right\} \right]^n d\phi$$

$$= \frac{1}{\pi} \int_0^\pi \left[\left(\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} e^{i\phi} \right) \left(\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} e^{-i\phi} \right) \right]^n d\phi$$

$$= \frac{1}{\pi} \cos^{2n} \frac{\theta}{2} \int_0^\pi \left(1 + i \tan \frac{\theta}{2} e^{i\phi} \right)^n \left(1 + i \tan \frac{\theta}{2} e^{-i\phi} \right)^n d\phi$$

$$= \frac{1}{\pi} \cos^{2n} \frac{\theta}{2} \int_0^\pi \left\{ 1 + ni \tan \frac{\theta}{2} e^{i\phi} - \frac{n(n-1)}{2} \tan^2 \frac{\theta}{2} e^{i\phi} e^{-i\phi} + \dots \right\} \times \left\{ 1 + ni \tan \frac{\theta}{2} e^{-i\phi} - \frac{n(n-1)}{2} \tan^2 \frac{\theta}{2} e^{-i\phi} e^{i\phi} + \dots \right\} d\phi.$$

Here the diagonal product will give cosines of multiples of ϕ and when they will be taken in groups, they vanish in limits of integration. We have only to consider the column products, i.e.

$$P_n(\cos \theta) = \frac{1}{\pi} \cos^{2n} \frac{\theta}{2} \int_0^\pi \left[1 - \tan^2 \frac{\theta}{2} + \frac{n^2 (n-1)^2}{2 \cdot 2} \tan^4 \frac{\theta}{2} - \dots \right] d\phi$$

$$= \frac{1}{\pi} \cos^2 \frac{\theta}{2} \left[\pi - \pi n^2 \tan^2 \frac{\theta}{2} + \pi \frac{n^2 (n-1)^2}{2 \cdot 2} \tan^4 \frac{\theta}{2} - \dots \right]$$

$$= \cos^{2n} \frac{\theta}{2} \left[1 + \frac{(-n)(-n)}{1 \cdot 1} \left(-\tan^2 \frac{\theta}{2} \right) \right]$$

$$+ \frac{(-n)(-n+1)(-n)(-n+1)}{1 \cdot 2 \cdot 1 \cdot (1+1)} \left(-\tan^2 \frac{\theta}{2} \right)^2 - \dots]$$

$$= \cos^{2n} \frac{\theta}{2} F\left(-n, -n, 1, -\tan^2 \frac{\theta}{2}\right).$$

Another form of Problem 15.

Show that

$$P_n(\cos \theta) = \cos^n \theta F\left(-\frac{n}{2}, \frac{1}{2} - \frac{n}{2}, 1, -\tan^2 \theta\right).$$

The first of Laplace's integrals gives

$$P_n(\cos \theta) = \frac{1}{\pi} \int_0^\pi (\cos \theta + i \sin \theta \cos \phi)^n d\phi$$

$$= \frac{1}{\pi} \cos^n \theta \int_0^\pi (1 + i \tan \theta \cos \phi)^n d\phi$$

$$= \frac{1}{\pi} \cos^n \theta \int_0^\pi \left\{ 1 + ni \tan \theta \cos \phi - \frac{n(n-1)}{2} \tan^2 \theta \cos^2 \phi + \frac{n(n-1)(n-2)}{6} \tan^3 \theta \cos^3 \phi + \dots \right\} d\phi.$$

In limits from 0 to $\pi/2$ all the cosine terms having odd powers vanish.

$$\therefore P_n(\cos \theta) = \frac{2}{\pi} \cos^n \theta \int_0^{\pi/2} \left[1 - \frac{n(n-1)}{2} \tan^2 \theta \cos^2 \phi + \dots \right] d\phi$$

$$= \frac{2}{\pi} \cos^n \theta \left[\frac{2}{\pi} - \frac{n(n-1)}{2} \tan^2 \theta \cdot \frac{1}{2} \cdot \frac{\pi}{2} + \frac{n(n-1)(n-2)(n-3)}{4} \tan^4 \theta \cdot \frac{3 \cdot 1 \pi}{4 \cdot 1 \cdot 2} + \dots \right]$$

$$= \cos^n \theta \left[1 + \frac{(-\frac{1}{2}n)(-\frac{1}{2}n+\frac{1}{2})}{1 \cdot 1} (-\tan^2 \theta) \right]$$

$$+ \frac{(-\frac{1}{2}n)(-\frac{1}{2}n+1)(-\frac{1}{2}n+\frac{3}{2})(-\frac{1}{2}n+\frac{5}{2})}{1 \cdot 2 \cdot 1 \cdot (1+1)} (-\tan^2 \theta)^2 + \dots]$$

$$= \cos^n \theta F\left[-\frac{1}{2}n, \frac{1}{2} - \frac{1}{2}n, 1, -\tan^2 \theta\right].$$

[F] Orthogonal Properties of Legendre's Polynomials of the First Kind

To show that

$$\int_{-1}^1 P_m(\mu) P_n(\mu) d\mu = 0 \text{ unless } m=n \quad (\text{Agra, 1963})$$

$$= \frac{2}{2n+1} \text{ if } m=n$$

or

$$= \frac{2}{2n+1} \delta_{m,n} \text{ in Kronecker delta symbol } m, n$$

being positive integers.

(Agra, 1963; Kurukshetra, 1965)

Rodrigue's formula is

$$P_n(\mu) = \frac{1}{2^n |n|} \left(\frac{d}{d\mu} \right)^n (\mu^2 - 1)^n.$$

With its application, we have

$$\begin{aligned} & \int_{-1}^1 P_m(\mu) P_n(\mu) d\mu \\ &= \frac{1}{2^{m+n}} \frac{1}{|m| |n|} \int_{-1}^1 \left(\frac{d}{d\mu}\right)^m (\mu^2-1)^m \cdot \left(\frac{d}{d\mu}\right)^n (\mu^2-1)^n d\mu \\ &= \frac{1}{2^{m+n}} \frac{1}{|m| |n|} \left[\left\{ \left(\frac{d}{d\mu}\right)^m (\mu^2-1)^m \cdot \left(\frac{d}{d\mu}\right)^{n-1} (\mu^2-1)^n \right\}_{-1}^1 \right. \\ & \quad \left. - \int_{-1}^1 \left(\frac{d}{d\mu}\right)^{m+1} (\mu^2-1)^m \cdot \left(\frac{d}{d\mu}\right)^{n-1} (\mu^2-1)^n d\mu \right] \\ &= -\frac{1}{2^{m+n}} \frac{1}{|m| |n|} \left[\int_{-1}^1 \left(\frac{d}{d\mu}\right)^{m+1} (\mu^2-1)^m \cdot \left(\frac{d}{d\mu}\right)^{n-1} (\mu^2-1)^n d\mu \right] \end{aligned}$$

(integrating by parts)

(the first term vanishing for both the limits)

Continuing the process of integration by parts m times on R.H.S., we get

$$\begin{aligned} & \int_{-1}^1 P_m(\mu) P_n(\mu) d\mu \\ &= \frac{(-1)^m}{2^{m+n}} \frac{1}{|m| |n|} \int_{-1}^1 \left(\frac{d}{d\mu}\right)^{2m} (\mu^2-1)^m \left(\frac{d}{d\mu}\right)^{n-m} (\mu^2-1)^n d\mu \end{aligned}$$

Here $\left(\frac{d}{d\mu}\right)^{2m} (\mu^2-1)^m = \frac{d^{2m} (\mu^2-1)^m}{d\mu^{2m}} = |2m|$

$$\begin{aligned} \therefore \int_{-1}^1 P_m(\mu) P_n(\mu) d\mu &= \frac{(-1)^m |2m|}{2^{m+n}} \frac{1}{|m| |n|} \int_{-1}^1 \left(\frac{d}{d\mu}\right)^{n-m} (\mu^2-1)^n d\mu \end{aligned} \quad \dots (63)$$

$$\begin{aligned} &= \frac{(-1)^m |2m|}{2^{m+n}} \frac{1}{|m| |n|} \left[\left(\frac{d}{d\mu}\right)^{n-m-1} (\mu^2-1)^n \right]_{-1}^1 \\ &= \frac{(-1)^m |2m|}{2^{m+n}} \frac{1}{|m| |n|} [0] = 0 \text{ when } m \neq n \text{ and } n > m. \end{aligned} \quad \dots (64)$$

Again when $m=n$, from (63), we have

$$\int_{-1}^1 P_n^2(\mu) d\mu = \frac{(-1)^n |2n|}{2^{n+n}} \frac{1}{|n|} \int_{-1}^1 (\mu^2-1)^n d\mu$$

$$\begin{aligned} \text{or } \int_{-1}^1 P_n^2(\mu) d\mu &= \frac{(-1)^n |2n|}{2^{n+n}} \frac{1}{|n|} (-1)^n \int_{-1}^1 (1-\mu^2)^n d\mu \\ &= \frac{1.3.5 \dots (2n-1)}{2^n |n|} \int_{-1}^1 (1-\mu^2)^n d\mu \end{aligned}$$

put $\mu = \cos \theta$, $d\mu = -\sin \theta d\theta$

$$\begin{aligned} &= -\frac{1.3.5 \dots (2n-1)}{2^n |n|} \int_{\pi}^0 \sin^{2n+1} \theta d\theta \\ &= \frac{2 \times 1.3.5 \dots (2n-1)}{2^n |n|} \int_0^{\pi/2} \sin^{2n+1} \theta d\theta \\ &= 2 \cdot \frac{1.3.5 \dots (2n-1)}{2^n |n|} \times \frac{2n(2n-2) \dots 2}{(2n+1)(2n-1) \dots 3.1} \\ &= 2 \cdot \frac{2^n |n|}{2^n |n| (2n+1)} = \frac{2}{2n+1} \end{aligned} \quad \dots (65)$$

Problem 16. Show that $\int_{-1}^1 \left\{ \frac{dP_n(\mu)}{d\mu} \right\}^2 d\mu = n(n+1)$.

We have proved that

$$\frac{dP_n(\mu)}{d\mu} = (2n-1) P_{n-1}(\mu) + (2n-5) P_{n-3}(\mu) + \dots \text{ ending with } 3P_1(\mu) \text{ or } P_0(\mu) \text{ according as } n \text{ is even or odd.}$$

$$\begin{aligned} \therefore \int_{-1}^1 \left\{ \frac{dP_n(\mu)}{d\mu} \right\}^2 d\mu &= \int_{-1}^1 [(2n-1) P_{n-1}(\mu) + (2n-5) P_{n-3}(\mu) + \dots \text{ ending with } \\ & \quad 3P_1(\mu) \text{ or } P_0(\mu) \text{ according as } n \text{ is even or odd}]^2 d\mu. \end{aligned}$$

Here on the R.H.S. the product terms vanish in limits and only the square terms are left.

$$\begin{aligned} \therefore \int_{-1}^1 \left\{ \frac{dP_n(\mu)}{d\mu} \right\}^2 d\mu &= (2n-1)^2 \cdot \frac{2}{2(n-1)+1} + (2n-5)^2 \cdot \frac{2}{2(n-3)+1} + \dots \\ & \quad \text{ending with } 3^2 \cdot \frac{2}{2+1} \text{ or } 1^2 \cdot \frac{2}{2 \times 0 + 1} \text{ according as } n \\ & \quad \text{is even or odd} \\ &= 2[(2n-1) + (2n-5) + (2n-9) + \dots + 3] \text{ when } n \text{ is even} \quad \dots (1) \\ &= 2[(2n-1) + (2n-5) + \dots + 1] \text{ when } n \text{ is odd.} \quad \dots (2) \end{aligned}$$

Now we know that in an arithmetical progression having a as its first term, l the last term, d the common difference and N the total number of terms,

$$l = a + (N-1)d, \text{ i.e. } N = \frac{l-a}{d} + 1.$$

In (1), $N = \frac{2n-1-3}{4} + 1 = \frac{n}{2}.$

$$\therefore \text{sum} = 2 \cdot \frac{N}{2} [a+l] = \frac{2n}{4} [3+2n-1] = n(n+1).$$

$$\text{In (2).} \quad N = \frac{2n-1-1}{4} + 1 = \frac{n+1}{2}.$$

$$\therefore \text{sum} = 2 \cdot \left(\frac{n+1}{4} \right) \{1 + 2n-1\} = n(n+1).$$

We see that the sum in two cases is the same whether n is even or odd.

$$\text{Hence} \quad \int_{-1}^1 \left\{ \frac{dP_n(\mu)}{d\mu} \right\}^2 d\mu = n(n+1).$$

$$\text{Problem 17. Show that} \quad \int_{-1}^1 (1-x^2) \left\{ \frac{dP_n(x)}{dx} \right\}^2 dx = \frac{2n(n+1)}{2n+1}.$$

From recurrence formulae (45) and (44), we have

$$(1-x^2) \frac{dP_n(x)}{dx} = \frac{n(n+1)}{2n+1} \{P_{n-1}(x) - P_{n+1}(x)\}, \quad \dots (1)$$

$$\text{and} \quad \frac{dP_n(x)}{dx} = (2n-1)P_{n-1}(x) + (2n-5)P_{n-3}(x)$$

$$+ (2n-9)P_{n-5}(x) \dots \text{ending within } 3P_1(x)$$

or $P_0(x)$ according as n is even or odd. $\dots (2)$

Multiplying (1) and (2) together and integrating within the limits -1 and 1 , we get

$$\int_{-1}^1 (1-x^2) \left\{ \frac{dP_n(x)}{dx} \right\}^2 dx = \int_{-1}^1 \left[\frac{n(n+1)}{2n+1} \{P_{n-1}(x) - P_{n+1}(x)\} \{ (2n-1)P_{n-1}(x) + (2n-5)P_{n-3}(x) \} \right] dx = 0$$

$$\begin{aligned} \text{or} \quad \int_{-1}^1 (1-x^2) \left\{ \frac{dP_n(x)}{dx} \right\}^2 dx &= \frac{n(n+1)}{2n+1} (2n-1) \int_{-1}^1 P_{n-1}^2(x) dx \\ &\quad \text{other terms vanishing} \\ &= \frac{n(n+1)(2n-1)}{2n+1} \cdot \frac{2}{2(n-1)+1} \\ &= \frac{2n(n+1)}{2n+1}. \end{aligned}$$

Problem 18. If (u, ϕ, z) and (r, θ, ϕ) be the cylindrical and polar co-ordinates of the same point and if $\mu = \cos \theta$, show that

$$P_n(\mu) = (-1)^n \cdot \frac{r^{n+1}}{n!} \left(\frac{\partial}{\partial z} \right)^n \left(\frac{1}{r} \right).$$

$$\text{Here} \quad r = \sqrt{(x^2 + y^2 + z^2)} = (u^2 + z^2).$$

$$\therefore \frac{1}{r} = (u^2 + z^2)^{-1/2} = \phi(u, z) \text{ say,}$$

so that, by Taylor's theorem, we have

$$\phi(u, z-k) = \{u^2 + (z-k)^2\}^{-1/2}$$

$$\begin{aligned} &= \phi(u, z) - k \frac{\partial}{\partial z} \phi(u, z) + \frac{k^2}{2!} \left(\frac{\partial}{\partial z} \right)^2 \phi(u, z) \dots \\ &\dots + (-1)^n \frac{k^n}{n!} \left(\frac{\partial}{\partial z} \right)^n \phi(u, z) + \dots \quad \dots (1) \end{aligned}$$

$$\begin{aligned} \text{Also, } \phi(u, z-k) &= \{u^2 + (z-k)^2\}^{-1/2} \\ &= \{u^2 + z^2 - 2zk + k^2\}^{-1/2} \\ &= \{r^2 - 2r \cos \theta \cdot k + k^2\}^{-1/2} \\ &= (r^2)^{-1/2} \left[1 - 2 \cos \theta \cdot \frac{k}{r} + \frac{k^2}{r^2} \right]^{-1/2} \\ &= \frac{1}{r} \sum_{n=0}^{\infty} \left(\frac{k}{r} \right)^n P_n(\cos \theta). \quad \dots (2) \end{aligned}$$

Equating the coefficients of k^n in (1) and (2), we get

$$\frac{1}{r} \cdot \frac{1}{r^n} P_n(\cos \theta) = \frac{(-1)^n}{n!} \left(\frac{\partial}{\partial z} \right)^n \phi(u, z)$$

$$\text{or} \quad P_n(\cos \theta) = (-1)^n \cdot \frac{r^{n+1}}{n!} \left(\frac{\partial}{\partial z} \right)^n \cdot \frac{1}{r}.$$

Note. Relations between polar and cartesian co-ordinates:

$$x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi, \quad z = r \cos \theta.$$

Relations between cylindrical and cartesian co-ordinates:

$$x = u \cos \phi, \quad y = u \sin \phi, \quad z = z.$$

[G] The Associated Legendre Polynomials

Laplace's equation in spherical co-ordinates may be expressed as

$$r \frac{\partial^2(rV)}{\partial r^2} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial V}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2 V}{\partial \phi^2} = 0. \quad \dots (66)$$

Assuming that V is a product of single-valued function, let $V = R \cdot \Theta \cdot \Phi$;

where R is the function of r only, Θ is the function of θ only and Φ is the function of ϕ only.

$$\text{Then} \quad \frac{\partial V}{\partial r} = \Theta \cdot \Phi \frac{dR}{dr}; \quad \frac{\partial^2 V}{\partial r^2} = \Theta \cdot \Phi \frac{d^2 R}{dr^2},$$

$$\frac{\partial V}{\partial \theta} = R \cdot \Phi \frac{\partial \Theta}{\partial \theta}$$

$$\text{and} \quad \frac{\partial V}{\partial \phi} = R \cdot \Theta \frac{d\Phi}{d\phi}; \quad \frac{\partial^2 V}{\partial \phi^2} = R \cdot \Theta \cdot \frac{d^2 \Phi}{d\phi^2}.$$

Substituting these values in (66), we get

$$r \Theta \cdot \Phi \frac{d^2}{dr^2} (rR) + \frac{1}{\sin \theta} R \cdot \Phi \frac{d}{d\theta} \left(\sin \theta \frac{\partial \Theta}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} R \cdot \Theta \frac{d^2 \Phi}{d\phi^2} = 0$$

$$\text{or} \quad \frac{r}{R} \frac{d^2(rR)}{dr^2} + \frac{1}{\Theta \sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{\partial \Theta}{\partial \theta} \right) + \frac{1}{\Phi \sin^2 \theta} \frac{d^2 \Phi}{d\phi^2} = 0 \quad \dots (67)$$

(on dividing throughout by $R \cdot \Theta \cdot \Phi$)

or $\frac{r \sin^2 \theta}{R} \frac{d^2(rR)}{dr^2} + \frac{\sin \theta}{\Theta} \cdot \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) = -\frac{1}{\Phi} \frac{d^2\Phi}{d\phi^2} \dots (68)$

Consider the equation (67) whose L.H.S. contains ϕ , but the L.H.S. does not. It follows that $-\frac{1}{\Phi} \frac{d^2\Phi}{d\phi^2} = \text{constant (say)} n^2$, so that

$$\frac{d^2\Phi}{d\phi^2} + n^2\Phi = 0. \dots (69)$$

The roots of this equation being $\pm in$, its solution is $\Phi = A \cos n\phi + B \sin n\phi$.

... (70)

Now the equation (67) by the substitution $-\frac{1}{\Phi} \frac{d^2\Phi}{d\phi^2} = n^2$ becomes

$$\frac{r}{R} \frac{d^2(rR)}{dr^2} + \frac{1}{\Theta \sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) - \frac{n^2}{\sin^2 \theta} = 0. \dots (71)$$

It is obvious that the first term of the equation (71) does not contain θ and the second and third terms do not contain r . It follows that the first term i.e. $\frac{r}{R} \frac{d^2(rR)}{dr^2}$ must be constant.

Let $\frac{r}{R} \frac{d^2(rR)}{dr^2} = m(m+1).$... (72)

[Its solution may easily be shown to be $R = Cr^m + Dr^{-m-1}$.]

Putting this value in (71), we get

$$m(m+1) + \frac{1}{\Theta \sin \theta} \cdot \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) - \frac{n^2}{\sin^2 \theta} = 0$$

i.e. $\frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) + \left[m(m+1) - \frac{n^2}{\sin^2 \theta} \right] \Theta = 0. \dots (73)$

Taking $\Theta (\cos \theta) = z(x)$, i.e. replacing Θ by z and $\cos \theta$ by x , (73) becomes

$$(1-x^2) \frac{d^2z}{dx^2} - 2x \frac{dz}{dx} + \left[m(m+1) - \frac{n^2}{1-x^2} \right] z = 0. \dots (74)$$

If we now put $z = (1-x^2)^{n/2} y$

i.e. $\frac{dz}{dx} = -nx (1-x^2)^{(n/2)-1} y + (1-x^2)^{n/2} \frac{dy}{dx}$

and $\frac{d^2z}{dx^2} = -2nx (1-x^2)^{(n/2)-1} \frac{dy}{dx} - ny [(1-x^2)^{(n/2)-1} -$

$$-(n-2) x^2 (1-x^2)^{(n/2)-2}] + (1-x^2)^{n/2} \frac{d^2y}{dx^2},$$

then the equation (74) transforms to

$$(1-x^2) \frac{d^2y}{dx^2} - 2(n+1)x \frac{dy}{dx} + [m(m+1) - n(n+1)] y = 0. \dots (75)$$

In order to find the series solution of this differential equation, let us suppose that its series solution is

$$y = \sum a_k x^k,$$

so that $\frac{dy}{dx} = \sum k a_k x^{k-1}$ and $\frac{d^2y}{dx^2} = \sum k(k-1) a_k x^{k-2}$.

With these substitutions, (10) gives

$$\sum [(1-x^2) k(k-1) x^{k-2} - 2(n+1)x k x^{k-1} + \{m(m+1) - n(n+1)\} x^k] a_k = 0$$

or $\sum [k(k-1) x^{k-2} + \{m(m+1) - n(n+1) - 2k(n+1) - k(k-1)\} x^k] a_k = 0.$

This is an identical equation. We therefore, on equating the coefficients of x^k on either side, get

$$\{(k+2)(k+2-1) a_{k+2} + \{m(m+1) - n(n+1) - 2k(n+1) - k(k-1)\} a_k = 0$$

or $(k+1)(k+2) a_{k+2} + (m^2 - n^2 + m - n - 2nk - k - k) a_k = 0$

or $(k+1)(k+2) a_{k+2} + (m - n - k)(m + n + 1 + k) a_k = 0$

or $a_k = - \frac{(k+1)(k+2)}{(m-n-k)(m+n+1+k)} a_{k+2}. \dots (76)$

If $a_k = 0$, then it is obvious that $a_{k-2} = a_{k-4} = a_{k-6} = \dots = 0$.

But $a_k = 0$ if $k = -1$ or $k = -2$.

Now $a_{k+2} = - \frac{(m-n-k)(m+n+1+k)}{(k+1)(k+2)} a_k$ gives the sequence of coefficients

$$a_2 = - \frac{(m-n)(m+n+1)}{2!} a_0,$$

$$a_4 = - \frac{(m-n-2)(m+n+3)}{3.4} a_2$$

$$= \frac{(m-n)(m-n-2)(m+n+1)(m+n+3)}{4!} a_0$$

... etc.

and $a_3 = - \frac{(m-n-1)(m+n+2)}{3!} a_0$

$$a_5 = - \frac{(m-n-3)(m+n+4)}{4.5} a_0$$

$$= \frac{(m-n-1)(m-n-3)(m+n+2)(m+n+4)}{5!} a_0$$

Taking a_0 and a_1 as unity, since they are arbitrary, we have the two series solutions of (74) as

$$P_m^n(x) = (1-x^2)^{n/2} \left[x - \frac{(m-n)(m+n+1)}{2} x^2 + \frac{(m-n)(m-n-2)(m+n+1)(m+n+3)}{4} x^4 + \dots \right]$$

$$\text{and } Q_m^n(x) = (1-x^2)^{n/2} \left[x - \frac{(m-n-1)(m+n+2)}{3} x^3 + \frac{(m-n-1)(m-n-3)(m+n+2)(m+n+4)}{5} x^5 - \dots \right];$$

where $P_m^n(x)$ and $Q_m^n(x)$ are called as *associated function* of the n th order and m th degree.

In case when $(m-n)$ is a positive integer, $P_m^n(x)$ or $Q_m^n(x)$ will terminate involving x^{m-n} and in that case

$$z = (-x^2)^{n/2} \left[x^{m-n} - \frac{(m-n)(m-n-1)}{2 \cdot (2m-1)} x^{m-n-2} + \frac{(m-n)(m-n-1)(m-n-2)(m-n-3)}{2 \cdot 4 \cdot (2m-1)(2m-3)} x^{m-n-4} - \dots \right] \dots (77)$$

where the bracketed expression ends with a term involving x^0 if $(m-n)$ is even and x if $(m-n)$ is odd.

If m and n both are integers, then

$$z = \frac{2^m |m| |m-n|}{(2m)!} P_m^n(x).$$

With the substitution of this value of z along with $x = \cos \theta$ (77) becomes

$$P_m^n(x) = \frac{(2m)! \sin^n \theta}{2^m |m| |m-n|} \left[x^{m-n} - \frac{(m-n)(m-n-1)}{2 \cdot (2m-1)} x^{m-n-2} + \frac{(m-n)(m-n-1)(m-n-2)(m-n-3)}{2 \cdot 4 \cdot (2m-1)(2m-3)} x^{m-n-4} - \dots \right] \dots (78)$$

where the bracketed expression ends with a term involving x^0 if $(m-n)$ is even and x if $(m-n)$ is odd.

Further if we differentiate (75) w.r.t. x , then we get

$$(1-x^2) \frac{d^3 y}{dx^3} - [2x + 2(n+1)x] \frac{d^2 y}{dx^2} + [-2(n+1) + m(m+1) - n(n+1)] \frac{dy}{dx} = 0$$

$$\text{i.e. } (1-x^2) \frac{d^3 y}{dx^3} - 2(n+2)x \frac{d^2 y}{dx^2} + [m(m+1) - (n+1)(n+2)] \frac{dy}{dx} = 0. \dots (79)$$

Clearly if y satisfies (75) for n , then $\frac{dy}{dx}$ satisfies (79) for $n+1$. Thus

for $n=0$, y becomes identical with $P_m(x)$ and so $\frac{d^n}{dx^n} P_m(x)$ satisfies (75) for $n=n$ and the solution of (74) is

$$P_m^n(x) = (1-x^2)^{n/2} \frac{d^n P_m(x)}{dx^n}. \dots (80)$$

These functions are called the *associated Legendre Polynomials* or *associated harmonics*.

Note. The associated Legendre functions of the second kind are

$$Q_m^n(x) = (1-x^2)^{n/2} \frac{d^n Q_m(x)}{dx^n}. \dots (81)$$

Now (80) yields on putting $n=0$,

$$P_m^0(x) = P_m(x) \dots (82)$$

From (80) the values of associated Legendre polynomials can be found with the help of (79) as

$$\left. \begin{aligned} P_1^1 &= (1-x^2)^{1/2} = \sin \theta \text{ when } x = \cos \theta \\ P_2^1 &= 3x(1-x^2)^{1/2} = 3 \cos \theta \sin \theta \text{ when } x = \cos \theta \\ P_2^2 &= \frac{3}{2}(5x^2-1)(1-x^2)^{1/2} = \frac{3}{2}(5 \cos^2 \theta - 1) \sin \theta \end{aligned} \right\} \text{etc.} \dots (83)$$

A generating function for associated Legendre polynomials may be derived from that of $P_n(x)$, in the form

$$\sum_{k=0}^{\infty} P_{n+k}^n(\mu) h^k = \frac{2^n (1-\mu^2)^{n/2}}{2^n |n| (1-2\mu h + h^2)^{n+1}} \dots (84)$$

Now there being two indices in Associated Legendre polynomials, several recurrence formulae can be derived, but we summarize here below a few of them. We have from (43)

$$(2n+1) P_n(\mu) = \frac{d}{d\mu} P_{n+1}(\mu) - \frac{d}{d\mu} P_{n-1}(\mu)$$

Its differentiation, m times w.r.t. μ yields

$$(2n+1) \frac{d^m}{dx^m} P_n(\mu) = \frac{d^{m+1}}{d\mu^{m+1}} P_{n+1}(\mu) - \frac{d^{m+1}}{d\mu^{m+1}} P_{n-1}(\mu)$$

Multiplying throughout by $(1-\mu^2)^{(m-1)/2}$, it becomes with the help of (80)

$$(2n+1)(1-\mu^2)^{1/2} P_n^m(\mu) = P_{n+1}^{m+1}(\mu) - P_{n-2}^{m+1}(\mu) \dots (85)$$

$$= (n+m)(n+m-1) P_{n-1}^{m-1}(\mu) - (n-m+1)(n-m+2) P_{n-1}^{m-1}(\mu) \dots (86)$$

and with similar few results

$$(2n+1)\mu P_n^m(\mu) = (n+m) P_{n-1}^m(\mu) + (n-m+1) P_{n+1}^m(\mu) \dots (87)$$

$$P_n^{m+1}(\mu) - \frac{2m\mu}{(1-\mu^2)^{1/2}} P_n^m(\mu) + [n(n+1) - m(m-1)] P_n^{m-1}(\mu) = 0 \quad \dots (88)$$

$$(1-\mu^2)^{1/2} \left(P_n^m \right)' = \frac{1}{2} P_n^{m+1}(\mu) - \frac{1}{2} (n+m)(n-m+1) P_n^{m-1}(\mu) \quad \dots (89)$$

Further using Rodrigue's formula, orthogonality of Legendre associated functions may be shown as

$$\int_{-1}^1 P_p^m(\mu) P_q^m(\mu) d\mu = 0 \quad \text{or} \quad \frac{2}{2q+1} \frac{|q+m|}{|q-m|} \quad \dots (90)$$

according as $q \neq p$ or $q = p$.

If it is transformed by the transformation $\mu = \cos \theta$, it becomes

$$\left. \begin{aligned} \int_0^\pi P_p^m(\cos \theta) P_q^m(\cos \theta) \sin \theta d\theta &= 0 \quad \text{when } q \neq p \\ &= \frac{2}{2q+1} \frac{|q+m|}{|q-m|} \quad \text{when } q = p \end{aligned} \right\} \quad \dots (91)$$

or
$$= \frac{2}{2q+1} \frac{|q+m|}{|q-m|} \delta_{q,p} \quad \text{in Kronecker delta symbol.}$$

If we now define a function

$$\bar{P}_l^m(\cos \theta) = \left(\sqrt{\frac{2l+1}{2}} \frac{|l-m|}{|l+m|} \right) P_l^m(\cos \theta), \quad -l \leq m \leq l \quad \dots (92)$$

$$\text{then } \int_{-1}^1 \bar{P}_l^m(\mu) \bar{P}_{l'}^m(\mu) d\mu = \delta_{ll'} \quad \dots (93)$$

So that $\bar{P}_l^m$ is orthogonal w.r.t. θ .

Also we introduce a function
$$\Phi_m(\phi) = \frac{1}{\sqrt{2\pi}} e^{im\phi}$$

$$\text{then } \int_0^{2\pi} \Phi_{m_1}^*(\phi) \Phi_{m_2}(\phi) d\phi = \frac{1}{2\pi} \int_0^{2\pi} e^{-im_1\phi} e^{im_2\phi} d\phi = \delta_{m_1, m_2} \quad \dots (94)$$

Showing that $\Phi_m(\phi)$ is an orthonormal function w.r.t. azimuthal angle.

As such the spherical harmonics

$$Y_l^m(\theta, \phi) = \bar{P}_l^m(\cos \theta) \Phi_m(\phi) = \sqrt{\frac{2l+1}{4\pi}} \left(\frac{|l-m|}{|l+m|} \right) P_l^m(\cos \theta) e^{im\phi} \quad \dots (95)$$

The spherical harmonics $Y_l^m(\theta, \phi)$ are a complete orthonormal set of functions i.e.

$$(Y_{l'}^{m'}, Y_l^m) = \delta_{ll'} \delta_{mm'} \quad \dots (96)$$

$$\text{We can also show that } P_l^m(\mu) = (-1)^m \frac{|l-m|}{|l+m|} P_l^m(\mu) \quad \dots (97)$$

$$Q_l^m = \text{constant} \times r^l Y_l^m(\theta, \phi) \quad \dots (98)$$

$$Y_l^m = (-1)^m Y_l^{m*} \quad \dots (99)$$

Also few of the spherical Harmonics are

$$\left. \begin{aligned} Y_0^0 &= \frac{1}{\sqrt{4\pi}}, Y_1^0 = \sqrt{\frac{3}{4\pi}} \cos \theta, \\ Y_1^1 &= -\sqrt{\frac{3}{8\pi}} e^{i\phi} \sin \theta \\ Y_2^0 &= \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1), \\ Y_2^1 &= -\sqrt{\frac{15}{8\pi}} e^{i\phi} \sin \theta \cos \theta \\ Y_2^2 &= \sqrt{\frac{15}{32\pi}} e^{2i\phi} \sin^2 \theta \text{ etc.} \end{aligned} \right\} \quad \dots (100)$$

Addition theorem of spherical Harmonics is

$$Y_l(0, \phi) = \sqrt{\frac{2l+1}{4\pi}} (Y_l^0, Y_l) = \frac{2l+1}{4\pi} (P_l(\cos \theta), Y_l) \quad \dots (101)$$

[H] Legendre's function of second kind i.e. Neumann's Integral for $Q_n(x)$ where $x > 1$.

We have from (13)

$$\begin{aligned} Q_n(x) &= \frac{|n|}{1.3.5 \dots (2n+1)} \left[x^{-n-1} + \frac{(n+1)(n+2)}{2(2n+3)} x^{-n-3} \right. \\ &\quad \left. + \frac{(n+1)(n+2)(n+3)(n+4)}{2.4.(2n+3)(2n+5)} x^{-n-5} + \dots \right] \\ &= \frac{2^n |n|}{|2n+1|} \sum_{r=0}^{\infty} \frac{|n+2r|}{2^r |r(2n+3)(2n+5) \dots (2n+2r-1)} x^{-(n+2r+1)} \quad \dots (102) \end{aligned}$$

Neumann's formula for Legendre's function is defined as

$$\int_{-1}^1 \frac{P_n(x) dx}{y-x} = 2 Q_n(y) \quad \dots (103)$$

where $|y| > 1$ and n is a positive integer.

The recurrence formulae for $Q_n(x)$ can be summarised without proof as they have no utility in physics:

If $a_0 = \frac{1}{2^n \Gamma(n+1)}$, this solution is called as $J_n(x)$.

$$\begin{aligned} \text{Thus } J_n(x) &= \frac{x^n}{2^n \Gamma(n+1)} \cdot \sum_{r=0}^{\infty} \frac{(-1)^r x^{2r}}{2^{2r} (r)! (n+1)(n+2)\dots(n+r)} \\ &= \sum_{r=0}^{\infty} (-1)^r \left(\frac{x}{2}\right)^{n+2r} \cdot \frac{1}{r! \Gamma(n+r+1)} \end{aligned} \quad \dots (11)$$

Case II. When $k = -n$.

The series solution is obtained by replacing n by $-n$ in the value of $J_n(x)$, whence, we get

$$J_{-n}(x) = \sum_{r=0}^{\infty} (-1)^r \cdot \left(\frac{x}{2}\right)^{-n+2r} \cdot \frac{1}{r! \Gamma(-n+r+1)} \quad \dots (12)$$

The complete primitive of Bessel's equation is

$$A J_n(x) + B J_{-n}(x), \quad (\text{Agra, 1961})$$

where n is not an integer, A, B being two arbitrary constants.

COROLLARY. Bessel's equation for $n=0$ is

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + y = 0. \quad (\text{Nagpur, 1961})$$

Its series solution by the same substitution $y = \sum_{r=0}^{\infty} a_r x^{k+r}$ (as above) is obtained to be

$$y = a_0 \left(1 - \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} - \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots \right).$$

If $a_0 = 1$, this solution is denoted by $J_0(x)$, i.e.

$$J_0(x) = 1 - \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} - \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots \quad \dots (9)$$

where $J_0(x)$ is called *Bessel function of zeroth order*.

In fact $J_0(x)$ is that solution of Bessel's equation for $n=0$, which is equal to unity for $x=0$.

Note. $J_n(x)$ is called *Bessel's function of the first kind of order n* .

[B] **Generating Function for $J_n(x)$** , i.e. to show that

$$e^{(x/2)(t-1/t)} = \sum_{n=-\infty}^{\infty} t^n J_n(x).$$

We know that

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$= \sum_{r=0}^{\infty} \frac{x^r}{r!}.$$

$$\therefore e^{xt/2} = \sum_{r=0}^{\infty} \frac{x^r t^r}{2^r r!}. \quad \dots (10)$$

$$\text{Similarly, } e^{-x/2t} = \sum_{s=0}^{\infty} \frac{(-1)^s x^s}{2^s t^s s!}. \quad \dots (11)$$

Multiplying (10) and (11), we get

$$e^{x/2(t-1/t)} = \sum_{r=0}^{\infty} \frac{x^r t^r}{2^r r!} \times \sum_{s=0}^{\infty} \frac{(-1)^s x^s}{2^s t^s s!}.$$

In order to find the (t^n) th term, we should replace r by $n+s$ and coefficient of t^n is

$$\begin{aligned} \frac{x^{n+s}}{2^{n+s} (n+s)!} \times \frac{(-1)^s x^s}{2^s s!} &= \sum_{s=0}^{\infty} (-1)^s \left(\frac{x}{2}\right)^{n+2s} \frac{1}{(n+s)! s!} \\ &= J_n(x) \quad \dots (12) \end{aligned}$$

Again the coefficient of t^{-n} is obtained by putting $s=n+r$ and

$$\begin{aligned} \text{coefficient of } t^{-n} &= \sum_{r=0}^{\infty} \frac{x^r}{2^r r!} \times \frac{(-1)^{n+r} x^{n+r}}{2^{n+r} (n+r)!} \\ &= (-1)^n \sum_{r=0}^{\infty} (-1)^r \left(\frac{x}{2}\right)^{n+2r} \frac{1}{r! (n+r)!} \\ &= (-1)^n \cdot J_n(x) \\ &= J_{-n}(x) \quad \dots (13) \end{aligned}$$

$J_{-n}(x) = (-1)^n J_n(x)$, where n is a positive integer.

It may be shown as below:

$$J_{-n}(x) = \sum_{r=0}^{\infty} (-1)^r \left(\frac{x}{2}\right)^{-n+2r} \cdot \frac{1}{r! \Gamma(-n+r+1)},$$

which tends to zero if $-n+r+1=0$, i.e. $r=n-1$ ($\because \Gamma 0 = \infty$).

Hence all the terms upto n th, vanish and therefore the limit $r=0$ may be changed to $r=n$.

$$J_{-n}(x) = \sum_{r=0}^{\infty} (-1)^r \left(\frac{x}{2}\right)^{-n+2r} \cdot \frac{1}{r! \Gamma(-n+r+1)}.$$

Now putting $r=n+s$, where s is a positive integer, we have

$$(1) = \sum_{s=0}^{\infty} (-1)^{n+s} \left(\frac{x}{2}\right)^{n+2s} \frac{1}{(n+s)! \Gamma(-n+s+n+1)}$$

$$= \sum_{s=0}^{\infty} (-1)^{n+s} \left(\frac{x}{2}\right)^{n+2s} \frac{1}{(n+s)! \Gamma(s+1)}$$

$$\begin{aligned} J_{-n}(x) &= (-1)^n \sum_{s=0}^{\infty} (-1)^s \left(\frac{x}{2}\right)^{n+2s} \frac{1}{s! \Gamma(n+s+1)} \\ &= (-1)^n J_n(x). \quad \dots (14) \end{aligned}$$

(Agra, 1961)

Hence from (12) and (13), we have

$$e^{x/2(t-1/t)} = \sum_{n=-\infty}^{\infty} t^n J_n(x). \quad \dots (15)$$

COROLLARY. Putting $t = e^{i\phi}$ and $\frac{1}{t} = e^{-i\phi}$, we get

$$e^{i\omega} \frac{(e^{i\phi} - e^{-i\phi})}{2i} = \sum_{n=-\infty}^{\infty} e^{ni\phi} J_n(x)$$

$$\begin{aligned} \text{or } e^{i\omega \sin \phi} &= J_0(x) + \{J_1(x) e^{i\phi} + J_{-1}(x) e^{-i\phi}\} \\ &\quad + \{J_2(x) e^{2i\phi} + J_{-2}(x) e^{-2i\phi}\} + \dots \\ \text{or } \cos(x \sin \phi) + i \sin(x \sin \phi) &= J_0(x) + J_1(x) \{e^{i\phi} - e^{-i\phi}\} + J_2(x) \{e^{2i\phi} + e^{-2i\phi}\} + \dots \\ &\quad \text{[since } J_n(x) = J_{-n}(x) \text{ when } n \text{ is even]} \\ &= J_0(x) + 2i \sin \phi J_1(x) + 2 \cos 2\phi J_2(x) + \dots \end{aligned}$$

Equating real and imaginary parts, we get

$$\begin{aligned} \cos(x \sin \phi) &= J_0(x) + 2 \cos 2\phi J_2(x) + 2 \cos 4\phi J_4(x) + \dots \\ \text{and } \sin(x \sin \phi) &= 2J_1(x) \sin \phi + 2J_3(x) \sin 3\phi + 2J_5(x) \sin 5\phi + \dots \end{aligned}$$

Replacing ϕ by $\frac{\pi}{2} - \phi$, we have, from (16) and (17)

$$\begin{aligned} \cos(x \cos \phi) &= J_0(x) - 2 \cos 2\phi J_2(x) + 2 \cos 4\phi J_4(x) - \dots \\ \text{and } \sin(x \cos \phi) &= 2 \cos \phi J_1(x) - 2 \cos 3\phi J_3(x) + 2 \cos 5\phi J_5(x) - \dots \end{aligned}$$

[C] Integrals for $J_0(x)$ and $J_n(x)$.

$$\text{I. } J_0(x) = \frac{1}{\pi} \int_0^\pi \cos(x \sin \phi) d\phi.$$

We have by (16),

$$\cos(x \sin \phi) = J_0(x) + 2 \cos 2\phi J_2(x) + 2 \cos 4\phi J_4(x) + \dots$$

If we integrate both the sides of this relation with respect to ϕ from the limits 0 to ϕ , then we see that all the integrals except the first of the R.H.S. vanish, thereby giving

$$\int_0^\pi \cos(x \sin \phi) d\phi = J_0(x) \int_0^\pi d\phi = \pi J_0(x).$$

$$\text{II. } J_n(x) = \frac{1}{\pi} \int_0^\pi \cos(n\phi - x \sin \phi) d\phi.$$

We have already proved that

$$\begin{aligned} \cos(x \sin \phi) &= J_0(x) + 2 \cos 2\phi J_2(x) + 2 \cos 4\phi J_4(x) + \dots \\ \text{and } \sin(x \sin \phi) &= 2J_1(x) \sin \phi + 2 \sin 3\phi J_3(x) \\ &\quad + 2 \sin 5\phi J_5(x) + \dots \end{aligned}$$

If we multiply (21) by $\cos n\phi$, (22) by $\sin n\phi$ and integrate between the limits 0 to π , we have

$$\int_0^\pi \cos(x \sin \phi) \cos n\phi d\phi = 0 \quad \text{or} \quad \pi J_n(x)$$

according as n is odd or even

$$\text{and } \int_0^\pi \sin(x \sin \phi) \sin n\phi d\phi = \pi J_n(x) \quad \text{or} \quad 0$$

according as n is odd or even.

Adding (23) and (24), we find

$$\int_0^\pi [\cos(x \sin \phi) \cos n\phi + \sin(x \sin \phi) \sin n\phi] d\phi = \pi J_n(x)$$

whether n is odd or even

$$\text{or } \int_0^\pi \cos(n\phi - x \sin \phi) d\phi = \pi J_n(x),$$

$$J_n(x) = \frac{1}{\pi} \int_0^\pi \cos(n\phi - x \sin \phi) d\phi. \quad \dots (25)$$

$$\text{III. } J_n(x) = \frac{1}{\sqrt{\pi} \Gamma(n + \frac{1}{2})} \left(\frac{x}{2}\right)^n \int_0^\pi \cos(x \sin \phi) \cos^{2n} \phi d\phi \quad \dots (26)$$

If we expand $\cos(x \sin \phi)$ in the powers of $x \sin \phi$, the general term is

$$(-1)^r \frac{x^{2r}}{(2r)!} \sin^{2r} \phi.$$

General term of R.H.S. of (26)

$$= \frac{2}{\sqrt{\pi} \Gamma(n + \frac{1}{2})} \left(\frac{x}{2}\right)^n (-1)^r \int_0^\pi \frac{x^{2r}}{(2r)!} \sin^{2r} \phi \cos^{2n} \phi d\phi, \dots (27)$$

$$\text{where } \int_0^\pi \sin^{2r} \phi \cos^{2n} \phi d\phi$$

$$\begin{aligned} &= 2 \int_0^{\pi/2} \sin^{2r} \phi \cos^{2n} \phi d\phi \quad \left\{ \begin{array}{l} \text{Put } \sin^2 \phi = t, \\ \therefore 2 \sin \phi \cos \phi d\phi = dt \\ \text{or } d\phi = \frac{dt}{2\sqrt{t(1-t)}} \end{array} \right. \\ &= \int_0^1 t^r (1-t)^n \frac{dt}{t^{1/2} (1-t)^{1/2}} \\ &= \int_0^1 t^{(2r-1)/2} (1-t)^{(2n-1)/2} dt \\ &= \beta\left(\frac{2r+1}{2}, \frac{2n+1}{2}\right) \\ &= \frac{\Gamma\left(\frac{2r+1}{2}\right) \Gamma\left(\frac{2n+1}{2}\right)}{\Gamma\left\{\left(\frac{2r+1}{2}\right) + \left(\frac{2n+1}{2}\right)\right\}} \quad \because \beta(m, n) = \frac{\Gamma m \Gamma n}{\Gamma(m+n)} \\ &= \frac{2r-1}{2} \cdot \frac{2r-3}{2} \dots \frac{1}{2} \sqrt{\pi} \Gamma\left(n + \frac{1}{2}\right) \\ &\quad \Gamma(n+r+1) \end{aligned}$$

Substituting this value in R.H.S. of (27) we have

$$\begin{aligned} &= \frac{2}{\sqrt{\pi} \Gamma(n + \frac{1}{2})} \left(\frac{x}{2}\right)^n (-1)^r \int_0^\pi \frac{x^{2r}}{(2r)!} \sin^{2r} \phi \cos^{2n} \phi d\phi \\ &= \frac{2}{\sqrt{\pi} \Gamma(n + \frac{1}{2})} \left(\frac{x}{2}\right)^n (-1)^r \frac{x^{2r}}{(2r)!} \times \frac{2r-1}{2} \cdot \frac{2r-3}{2} \dots \frac{1}{2} \sqrt{\pi} \Gamma\left(n + \frac{1}{2}\right) \\ &\quad \Gamma(n+r+1) \end{aligned}$$

$$\begin{aligned}
 &= 2 \left(\frac{x}{2} \right)^n (-1)^r \cdot \frac{x^{2r}}{\Gamma(n+r+1)} \cdot \frac{(2r-1)(2r-3) \dots 1}{2^r \cdot 2r(2r-1)(2r-2) \dots 1} \\
 &= \left(\frac{x}{2} \right)^{n+2r} (-1)^r \frac{1}{r! \Gamma(n+r+1)} \\
 &= (-1)^r \left(\frac{x}{2} \right)^{n+2r} \frac{1}{r! (n+r)!} = \text{general term in } J_n(x).
 \end{aligned}$$

$$\text{Hence } J_n(x) = \frac{1}{\sqrt{\pi} \Gamma(n+\frac{1}{2})} \left(\frac{x}{2} \right)^n \int_0^\pi \cos(x \sin \phi) \cos^{2n} \phi \, d\phi.$$

[D] Recurrence Formulae for $J_n(x)$.

I. We know that

$$J_n(x) = \sum_{r=0}^{\infty} (-1)^r \frac{1}{r! (n+r)!} \left(\frac{x}{2} \right)^{n+2r},$$

where n is a positive integer.

Differentiating it w.r.t x , we get

$$J_n'(x) = \sum_{r=0}^{\infty} (-1)^r \frac{1}{r! (n+r)!} (n+2r) \left(\frac{x}{2} \right)^{n+2r-1} \cdot \frac{1}{2}.$$

Multiplying both sides by x , we have

$$\begin{aligned}
 x J_n'(x) &= \sum_{r=0}^{\infty} (-1)^r \frac{1}{r! (n+r)!} \left(\frac{x}{2} \right)^{n+2r} \\
 &= n \sum_{r=0}^{\infty} (-1)^r \frac{(n+2r)}{r! (n+r)!} \left(\frac{x}{2} \right)^{n+2r} + x \sum_{r=0}^{\infty} (-1)^r \cdot \frac{r}{r! (n+r)!} \left(\frac{x}{2} \right)^{n+2r-1}
 \end{aligned}$$

$$\text{or } x J_n'(x) = n J_n(x) + x \sum_{r=1}^{\infty} (-1)^r \cdot \frac{1}{(r-1)! (n+r)!} \left(\frac{x}{2} \right)^{n+2r-1}$$

[since on R.H.S. the second term vanishes for $r=1$ and hence limit of $r=0$ may be replaced by $r=1$]

Putting $r-1=s$, we have

$$\begin{aligned}
 x J_n'(x) &= n J_n(x) + x \sum_{s=0}^{\infty} (-1)^{s+1} \frac{1}{s! (n+1+s)!} \left(\frac{x}{2} \right)^{n+2s-1} \\
 &= n J_n(x) - x J_{n+1}(x). \quad \dots (19)
 \end{aligned}$$

$$\text{II. Again } x J_n'(x) = \sum_{r=0}^{\infty} (-1)^r \cdot \frac{(n+2r)}{r! (n+r)!} \left(\frac{x}{2} \right)^{n+2r-1} \frac{x}{2} \text{ may be}$$

written as

$$x J_n'(x) = \sum_{r=0}^{\infty} (-1)^r \frac{-n+2(n+r)}{r! (n+r)!} \left(\frac{x}{2} \right)^{n+2r}$$

$$\begin{aligned}
 &= -n \sum_{r=0}^{\infty} (-1)^r \frac{1}{r! (n+r)!} \left(\frac{x}{2} \right)^{n+2r} \\
 &\quad + x \sum_{r=0}^{\infty} (-1)^r \frac{1}{r! (n+r-1)!} \left(\frac{x}{2} \right)^{n-1+2r} \\
 &= -n J_n(x) + x J_{n-1}(x). \quad (\text{Agra, 1962, 64}) \quad \dots (30)
 \end{aligned}$$

Sum and difference of (29) and (30) give

$$\text{III. } 2J_n'(x) = J_{n-1}(x) - J_{n+1}(x) \quad \dots (31)$$

$$\text{IV. } 2n J_n(x) = x \{J_{n+1}(x) + J_{n-1}(x)\}. \quad (\text{Agra, 1964}) \quad \dots (32)$$

$$\text{V. } \frac{d}{dx} \{x^n J_n(x)\} = x^n J_{n-1}(x). \quad \dots (33)$$

$$\text{Here } \frac{d}{dx} \{x^n J_n(x)\} = n x^{n-1} J_n(x) + x^n J_n'(x)$$

$$\begin{aligned}
 &= x^{n-1} \{n J_n(x) + x J_n'(x)\} \\
 &= x^{n-1} \{n J_n(x) - n J_n(x) + x J_{n-1}(x)\} \text{ by (30)} \\
 &= x^n J_{n-1}(x).
 \end{aligned}$$

VI. Similarly it is easy to show that

$$\frac{d}{dx} \{x^{-n} J_n(x)\} = -x^{-n} J_{n+1}(x). \quad \dots (34)$$

Problem 19. Show that $y = \frac{1}{\pi} \int_0^\pi \cos(x \cos \phi) \, d\phi$ satisfies the differential equation $\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + y = 0$ and that y is no other than $J_0(x)$.

$$\text{Given } y = \frac{1}{\pi} \int_0^\pi \cos(x \cos \phi) \, d\phi \quad \dots (1)$$

If we differentiate it w.r.t. x under the sign of integration, we find

$$\frac{dy}{dx} = \frac{1}{\pi} \int_0^\pi -\cos \phi \sin(x \cos \phi) \, d\phi \quad \dots (2)$$

$$\text{and } \frac{d^2 y}{dx^2} = \frac{1}{\pi} \int_0^\pi -\cos^2 \phi \cos(x \cos \phi) \, d\phi. \quad \dots (3)$$

Now from (2), we have

$$\begin{aligned}
 \frac{dy}{dx} &= -\frac{1}{\pi} \left[\int_0^\pi \sin(x \cos \phi) \sin \phi \, d\phi \right. \\
 &\quad \left. - \int_0^\pi -x \sin \phi \cos(x \cos \phi) \cdot \sin \phi \, d\phi \right] \text{ (integrating by parts)} \\
 &= -\frac{x}{\pi} \int_0^\pi \sin^2 \phi \cos(x \cos \phi) \, d\phi \\
 &= -\frac{x}{\pi} \int_0^\pi (1 - \cos^2 \phi) \cos(x \cos \phi) \, d\phi
 \end{aligned}$$

$$= -\frac{x}{\pi} \int_0^\pi \cos(x \cos \phi) d\phi + \frac{x}{\pi} \int_0^\pi \cos^2 \phi \cos(x \cos \phi) d\phi$$

$$= -xy - x \frac{d^2 y}{dx^2} \text{ from (1) and (3)}$$

or
$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + y = 0.$$

Hence $y = \frac{1}{\pi} \int_0^\pi \cos(x \cos \phi) d\phi$ satisfies $\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + y = 0,$

which is the Bessel's equation for $n=0$.

Since $y=1$ when $x=0$, therefore y is no other than $J_0(x)$, as $J_0(x)$ being the solution of Bessel's equation is unity for $x=0$.

Problem 20. Prove that $J_{1/2}(x) = \sqrt{\left(\frac{2}{\pi x}\right)} \sin x$.
(Agra, 1962, 66; Kanpur, 68)

We know that

$$J_n(x) = \frac{x^n}{2^n \Gamma(n+1)} \left[1 - \frac{x^2}{2 \cdot 2(n+1)} + \frac{x^4}{2 \cdot 4 \cdot 2(n+1)(n+2)} - \dots \right]$$

$$\therefore J_{1/2}(x) = \left(\frac{x}{2}\right)^{1/2} \frac{1}{\Gamma(\frac{3}{2})} \left[1 - \frac{x^2}{2 \cdot 2 \cdot \frac{3}{2}} + \frac{x^4}{2 \cdot 4 \cdot 2 \cdot \frac{3}{2} \cdot \frac{5}{2}} - \dots \right]$$

$$= \sqrt{\left(\frac{x}{2}\right)} \cdot \frac{1}{\frac{1}{2}\sqrt{\pi}} \left[1 - \frac{x^2}{2 \cdot 3} + \frac{x^4}{2 \cdot 3 \cdot 4 \cdot 5} - \dots \right]$$

$$= \sqrt{\left(\frac{2x}{\pi}\right)} \cdot \frac{1}{x} \left[x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \right]$$

$$= \sqrt{\left(\frac{2}{\pi x}\right)} \sin x.$$

Problem 21. Show that

(i) $J_0'(x) = -J_1(x)$ and (ii) $2J_0''(x) = J_2(x) - J_0(x)$.

From recurrence formula (29) we have

$$xJ_n'(x) = nJ_n(x) - xJ_{n+1}(x).$$

Putting $n=0$, we get $J_0'(x) = -J_1(x)$,

which proves the first result.

Now differentiating it and multiplying throughout by 2, we get

or
$$2J_0''(x) = -2J_1'(x)$$

$$2J_0''(x) = -[J_0(x) - J_2(x)]$$

by recurrence formula III

$$= J_2(x) - J_0(x).$$

Problem 22. Prove that $J_{n+3} + J_{n+5} = \frac{2}{x}(n+4)J_{n+4}$.

From recurrence formula IV, we have

$$2nJ_n = \lambda(J_{n-1} + J_{n+1}).$$

Replacing n by $n+1$, we get

$$\frac{2}{x}(n+4)J_{n+4} = J_{n+3} + J_{n+5}.$$

Problem 23. Prove that

$$\cos x = J_0(x) - 2J_2(x) + 2J_4(x) \dots$$

$$\sin x = 2J_1(x) - 2J_3(x) + 2J_5(x) \dots$$

We know that

$$\cos(x \cos \phi) = J_0(x) - 2 \cos 2\phi J_2(x) + 2 \cos 4\phi J_4(x) \dots$$

$$\text{and } \sin(x \sin \phi) = 2 \cos \phi J_1(x) - 2 \cos 3\phi J_3(x) + 2 \cos 5\phi J_5(x) \dots$$

Putting $\phi=0$, we get

$$\cos x = J_0(x) - 2J_2(x) + 2J_4(x) \dots$$

$$\sin x = 2J_1(x) - 2J_3(x) + 2J_5(x) \dots$$

Problem 24. Establish the relation

$$J_n(x) J_{-n}'(x) - J_n'(x) J_{-n}(x) = -\frac{2 \sin n\pi}{\pi x}.$$

We know that $J_n(x)$ and $J_{-n}(x)$ are the two solutions of Bessel's

equation $\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + \left(1 - \frac{n^2}{x^2}\right) y = 0. \dots (1)$

Hence, if $y = J_n(x)$, $\frac{dy}{dx} = J_n'(x)$ and $\frac{d^2 y}{dx^2} = J_n''(x)$, we have from (1)

$$J_n''(x) + \frac{1}{x} J_n'(x) + \left(1 - \frac{n^2}{x^2}\right) J_n(x) = 0. \dots (2)$$

Similarly putting $y = J_{-n}(x)$, we get from (1)

$$J_{-n}''(x) + \frac{1}{x} J_{-n}'(x) + \left(1 - \frac{n^2}{x^2}\right) J_{-n}(x) = 0. \dots (3)$$

Multiplying (2) by $J_{-n}(x)$, (3) by $J_n(x)$, and then subtracting (3) from (2), we have

$$J_n'(x) J_{-n}(x) - J_{-n}''(x) J_n(x) + \frac{1}{x} \{J_n'(x) J_{-n}(x) - J_{-n}'(x) J_n(x)\} = 0.$$

Put $z = J_n'(x) J_{-n}(x) - J_{-n}'(x) J_n(x)$.

$$\therefore z' = J_n''(x) J_{-n}(x) + J_n'(x) J_{-n}'(x) - J_{-n}''(x) J_n(x) - J_{-n}'(x) J_n'(x)$$

$$= J_n''(x) J_{-n}(x) - J_{-n}''(x) J_n(x).$$

Thus $z' + \frac{z}{x} = 0$ or $\frac{z'}{z} = -\frac{1}{x}.$

Integrating, $\log z = -\log x + \log C$, where C is some arbitrary constant

$$= \log \frac{C}{x}.$$

$$z = \frac{C}{x}.$$

$$J_n'(x) J_{-n}(x) - J_{-n}'(x) J_n(x) = \frac{C}{x}.$$

Equating the coefficients of $\frac{1}{x}$ on either side, we get

$$C = \frac{1}{2^n \Gamma(n+1)} \cdot \frac{1}{2^{-n} \Gamma(-n+1)} \{n - (-n)\} = C$$

$$C = \frac{2n}{\Gamma(n+1) \Gamma(-n+1)} = \frac{2}{\Gamma n \Gamma(1-n)}$$

$$= \frac{2}{\pi / \sin n\pi} = \frac{2 \sin n\pi}{\pi}; \quad \therefore \Gamma n \Gamma(1-n) = \frac{\pi}{\sin n\pi}$$

(Gamma functions).

$$\text{Hence } J_n'(x) J_{-n}(x) - J_{-n}'(x) J_n(x) = \frac{2 \sin n\pi}{\pi}.$$

[E] Orthogonal Properties of Bessel's Polynomials.

To prove that $\int_0^a J_n(\mu r) J_n(\mu' r) r dr = 0$ where μ and μ' are different roots of $J_n(\mu a) = 0$.

Since $J_n(x)$ is a solution of Bessel's equation,

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + \left(1 - \frac{n^2}{x^2}\right) y = 0 \quad \dots (35)$$

therefore putting $x = \mu r$ and calling $y = u$ in (35) we get

$$\frac{1}{\mu^2} \frac{d^2 u}{dr^2} + \frac{1}{\mu r} \frac{du}{dr} + \left(1 - \frac{n^2}{\mu^2 r^2}\right) u = 0,$$

$$\left[\because \frac{dy}{dx} = \frac{du}{dr} \cdot \frac{dr}{dx} = \frac{1}{\mu} \frac{du}{dr} \right]$$

$$\text{and } \frac{d^2 y}{dx^2} = \frac{1}{\mu} \frac{d}{dr} \left(\frac{1}{\mu} \frac{du}{dr} \right) = \frac{1}{\mu^2} \frac{d^2 u}{dr^2}.$$

Multiplying throughout by $\mu^2 r^2$,

$$r^2 \frac{d^2 u}{dr^2} + r \frac{du}{dr} + (\mu^2 r^2 - n^2) u = 0. \quad \dots (36)$$

Similarly putting $x = \mu' r$ and calling $y = v$ in (36), we have

$$r^2 \frac{d^2 v}{dr^2} + r \frac{dv}{dr} + (\mu'^2 r^2 - n^2) v = 0. \quad \dots (37)$$

If we multiply (36) by $\frac{v}{r}$, (37) by $\frac{u}{r}$ and subtract

$$(r \cdot v u'' - r u v'') + (v u' - u v') + (\mu^2 - \mu'^2) r u v = 0,$$

$$\text{where } u' = \frac{du}{dr} \text{ and } v' = \frac{dv}{dr} \text{ etc.}$$

$$\text{or } \frac{d}{dr} \{r (v u' - u v')\} + (\mu^2 - \mu'^2) r u v = 0, \quad \dots (38)$$

$$\text{where } u = J_n(\mu r) \text{ and } v = J_n(\mu' r).$$

Integrating (38) w.r.t. r between the limits 0 and a , we get

$$\left[r \{J_n(\mu r) \cdot J_n'(\mu' r) \mu' - J_n(\mu' r) J_n'(\mu r) \mu\} \right]_0^a$$

$$+ \int_0^a (\mu^2 - \mu'^2) J_n(\mu r) J_n(\mu' r) r dr = 0.$$

The first term vanishes for both the limits since

$$J_n(\mu a) = 0, \quad J_n(\mu' a) = 0.$$

$$\text{Hence } \int_0^a (\mu^2 - \mu'^2) J_n(\mu r) J_n(\mu' r) r dr = 0,$$

$$\text{i.e., } \int_0^a J_n(\mu r) J_n(\mu' r) r dr = 0 \quad \text{as } \mu^2 - \mu'^2 \neq 0. \quad \dots (39)$$

[F] Bessel's Functions of the Second Kind (Neumann functions).

Bessel's equation of zero order is

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + y = 0. \quad \dots (40)$$

One of its solution is $J_0(x)$ so that $J_0'' + \frac{1}{x} J_0' + J_0 = 0$

Let the other solution of (40) be

$$y = u J_0(x) + v$$

u, v being functions of x .

$$\text{Then } y' = u J_0' + u' J_0 + v'$$

$$y'' = u J_0'' + 2u' J_0' + J_0 u'' + v''$$

Substituting these values in (40) and using $J_0'' + \frac{1}{x} J_0' + J_0 = 0$,

$$\text{we get } \left(v'' + \frac{1}{x} v' + v\right) + J_0 \left(u'' + \frac{1}{x} u'\right) + 2u' J_0' = 0$$

$$\text{Choosing } u \text{ such that } u'' + \frac{1}{x} u' = 0 \text{ i.e. } \frac{u''}{u'} = -\frac{1}{x}.$$

$$\text{Integration gives } \log u' = -\log x \text{ i.e. } u' = \frac{1}{x}$$

$$\text{giving } u = \log x$$

then the above transformed equation becomes

$$v'' + \frac{1}{x} v' + v + \frac{2}{x} J_0' = 0$$

$$\text{or } v'' + \frac{1}{x} v' + v = \frac{2}{x} J_0', \quad \therefore J_0' = -J_1 \text{ by problem 21.}$$

But the recurrence formula $2n J_n = x (J_{n-1} + J_{n+1})$ gives on replacing n by $n+1$

$$2(n+1) J_{n+1} = x (J_n + J_{n+2})$$

$$\text{i.e. } \frac{x}{2} J_n = (n+1) J_{n+1} - \frac{1}{2} x J_{n+2}.$$

Replacing n by $n+2$, this gives

$$\frac{x}{2} J_{n+2} = (n+3) J_{n+3} - \frac{x}{2} J_{n+4}$$

$$\text{i.e. } \frac{x}{2} J_n = (n+1) J_{n+1} - (n+3) J_{n+3} + \frac{x}{2} J_{n+4}$$

Repeating the process we find

$$\frac{x}{2} J_n = (n+1) J_{n+1} - (n+3) J_{n+3} + (n+5) J_{n+5} \dots$$

that if $n=1$, $\frac{x}{2} J_1 = 2J_2 - 4J_4 + 6J_6 \dots$

$$\begin{aligned} \text{Thus } v'' + \frac{1}{x} v' + v &= \frac{4}{x^2} [2J_2 - 4J_4 + 6J_6 \dots] \\ &= \frac{4}{x^2} \sum (-1)^{n/2-1} n J_{n-1} \quad n \text{ being even integer} \\ &= \sum \frac{n^2}{x^2} \cdot \frac{4(-1)^{n/2-1}}{n} J_n(x) \end{aligned}$$

But $\frac{4(-1)^{n/2-1}}{n} J_n(x)$ is the particular solution of

$$y'' + \frac{1}{x} y' + y = \frac{4(-1)^{n/2-1}}{n} J_n(x) \cdot \frac{n^2}{x^2}.$$

Thus $v = \sum \frac{4(-1)^{n/2-1}}{n} J_n(x)$ (by comparison).

Hence the solution of (40) is

$$y = J_0(x) \log x + \sum \frac{4}{n} (-1)^{n/2-1} J_n = Y_0 \text{ (say)}$$

But Neumann gave the solution of (40) as

$$N_0(x) = J_0(x) \log x - \sum_{p=1}^{\infty} (-1)^p \frac{\left(\frac{x}{2}\right)^{2p}}{(\frac{p}{2})^2} \left[1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{p} \right] \quad \dots (41)$$

where $N_0(x)$ is said to be *Neumann functions of the second kind of order zero*.

Weber gave the solution of (40) as

$$Y_0(x) = \frac{2}{\pi} \left[\log \frac{x}{2} + \gamma \right] J_0(x) - \frac{2}{\pi} \sum_{p=1}^{\infty} (-1)^p \frac{(\frac{1}{2}x)^{2p}}{(\frac{p}{2})^2}$$

$$\left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{p} \right), \text{ where } \gamma \text{ is Euler's constant defined as}$$

$$\gamma = \lim_{m \rightarrow \infty} [1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{m} - \log m]. \quad \dots (42)$$

$$\dots (43)$$

From (2) and (3) it is easy to show that

$$Y_0(x) = \frac{2}{\pi} \{N_0(x) - (\log 2 - \gamma) J_0(x)\}. \quad \dots (44)$$

The complete primitive of (40) is

$$Y = A J_0(x) + B Y_0(x). \quad \dots (45)$$

In case of Bessel's equation of n th order, i.e.

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + \left[1 - \frac{n^2}{x^2} \right] y = 0, \quad \dots (46)$$

the complete primitive is

$$y = A' J_n(x) + B' Y_n(x), \quad \dots (47)$$

where $Y_n(x) = \frac{2}{\pi} \left[\gamma + \log \frac{x}{2} \right] J_n(x) - \frac{1}{\pi} \sum_{p=0}^{n-1} \frac{\Gamma(n-p)}{p!} \left[\frac{2}{p} \right]^{n-2p}$

$$- \frac{1}{\pi} \sum_{p=0}^{\infty} (-1)^p \frac{(\frac{1}{2}x)^{n+2p}}{p! \Gamma(n+p+1)} \left\{ 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{p} \right.$$

$$\left. + 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n+p} \right\}, \quad \dots (48)$$

when n is integral.

But if n is not an integer, then

$$Y_n = \frac{J_n(x) \cos n\pi - J_{-n}(x)}{\sin n\pi}. \quad \dots (49)$$

[G] **Bessel's Functions of the third kind (Hankel functions).**

These are defined as $H_n^{(1)}(x) = J_n(x) + iY_n(x). \quad \dots (50)$

$$H_n^{(2)}(x) = J_n(x) - iY_n(x). \quad \dots (51)$$

Thus addition and subtraction of (50) and (51) give

$$J_n(x) = \frac{1}{2} \{ H_n^{(1)}(x) + H_n^{(2)}(x) \}$$

and

$$Y_n(x) = -\frac{1}{2i} \{ H_n^{(1)}(x) - H_n^{(2)}(x) \}$$

[H] **The Modified Bessel's Functions.**

Bessel's differential equation of n th order is

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + \left(1 - \frac{n^2}{x^2} \right) y = 0. \quad \dots (52)$$

Put $x = iz$, so that $\frac{dx}{dz} = i$, i.e., $\frac{dz}{dx} = \frac{1}{i}$.

$$\therefore \frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx} = \frac{1}{i} \frac{dy}{dz}$$

and $\frac{d^2 y}{dx^2} = \frac{d}{dx} \left(\frac{1}{i} \frac{dy}{dz} \right) = \frac{d}{dz^2} \cdot \frac{dz}{dx} = \frac{1}{i^2} \frac{d^2 y}{dz^2} = -\frac{d^2 y}{dz^2}.$

With these substitutions, (52) becomes

$$-\frac{d^2y}{dz^2} + \frac{1}{iz} \cdot \frac{1}{i} \frac{dy}{dz} + \left\{ 1 - \frac{n^2}{(iz)^2} \right\} y = 0,$$

$$\text{i.e.,} \quad \frac{d^2y}{dz^2} + \frac{1}{z} \frac{dy}{dz} - \left(1 + \frac{n^2}{z^2} \right) y = 0. \quad \dots (53)$$

Since the complete primitive of (52) is $y = AJ_n(x) + BY_n(x)$, therefore the solution of (53) is obtained by putting $x = iz$ and that is

$$y = AJ_n(iz) + BY_n(iz). \quad \dots (54)$$

Now we know that

$$J_n(x) + \sum_{r=0}^{\infty} (-1)^r \frac{(x/2)^{n+2r}}{|r| (n+r)}$$

Putting $x = iz$, this becomes

$$J_n(iz) = \sum_{r=0}^{\infty} (-1)^r \frac{(iz)^{n+2r}}{2^{n+2r} |r| (n+r)}$$

$$= i^n \sum_{r=0}^{\infty} \frac{z^{n+2r}}{2^{n+2r} |r| (n+r)}$$

$$\therefore (i^2)^r = (i^2)^r = (-1)^r \text{ and therefore } (-1)^r \cdot i^{2r} = (-1)^r = 1$$

$$\text{or} \quad i^{-n} J_n(iz) = \sum_{r=0}^{\infty} \frac{z^{n+2r}}{2^{n+2r} |r| (n+r)} \quad \dots (55)$$

Here $i^{-n} J_n(iz)$ is denoted by $I_n(z)$, which is known as the *modified function of the first kind* of order n . Thus replacing z by x , we get

$$I_n(x) = i^{-n} J_n(ix) = \sum_{r=0}^{\infty} \frac{x^{n+2r}}{2^{n+2r} |r| \Gamma(n+r+1)} \quad \dots (56)$$

$$\therefore |r| (n+r) = \Gamma(n+r+1)$$

If we put $n=0$, its particular case is found as

$$\begin{aligned} I_0(x) &= \sum_{r=0}^{\infty} \frac{(x/2)^{2r}}{(|r|)^2} \\ &= 1 + \left(\frac{x}{2} \right)^2 + \frac{(x/2)^4}{(|2|)^2} + \frac{(x/2)^6}{(|3|)^2} + \dots \end{aligned} \quad \dots (57)$$

when n is integral, the *modified function of the second kind*, i.e. $K_n(x)$ is related with the modified function as

$$\begin{aligned} K_n(x) &= \frac{1}{2} \sum_{r=0}^{n-1} (-1)^r \frac{|(n-r-1)| \left(\frac{2}{x} \right)^{n-2r}}{|r|} \\ &\quad + (-1)^{n+1} \sum_{r=0}^{\infty} \frac{(x/2)^{n+2r}}{|r| (n+r)} \left[\log \frac{x}{2} - \frac{1}{2} \{ \psi(r+1) \right. \\ &\quad \left. + \psi(n+r+1) \} \right] \end{aligned} \quad \dots (58)$$

$$\text{where} \quad \psi(r+1) = (1 + \frac{1}{2} + \frac{1}{3} + \dots + 1/r) - \gamma \quad \dots (59)$$

$$\text{and} \quad \psi(n+r+1) = \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n+r} \right) - \gamma. \quad \dots (60)$$

But if n is not an integer, then

$$K_n(x) = \frac{\pi}{2} \frac{[I_{-n}(x) - I_n(x)]}{\sin n\pi} \quad \dots (61)$$

In particular when n is an integer, we have

$$K_0(x) = - \left\{ \gamma + \log \frac{x}{2} \right\} I_0(x) + \sum_{r=1}^{\infty} \frac{(\frac{1}{2}x)^{2r}}{(|r|)^2} \left\{ 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{r} \right\} \quad \dots (62)$$

Hence the complete primitive of the equation of the type

$$\frac{d^2y}{dx^2} + \frac{1}{x} \frac{dy}{dx} - \left(1 - \frac{n^2}{x^2} \right) y = 0$$

$$y = AI_n(x) + BK_n(x). \quad \dots (63)$$

1) The Ber, Bei, Ker, Kei Functions.

Suppose we have an equation of the form

$$x \frac{d^2y}{dx^2} + \frac{dy}{dx} - ip^2xy = 0. \quad \dots (64)$$

This may be written as

$$\frac{d^2y}{dx^2} + \frac{1}{x} \frac{dy}{dx} - p^2 y = 0.$$

Obviously this is a modified Bessel equation of the order zero and therefore its solution is

$$y = AI_0(xp\sqrt{i}) + BK_0(xp\sqrt{i}). \quad \dots (65)$$

In order to express this solution into real and imaginary parts, we introduce four new functions, namely, Ber (i.e. Bessel real); Bei (i.e. Bessel imaginary); Ker and Kei are their analogues.

These are defined as

$$I_0(x\sqrt{i}) = \text{Ber}(x) + i \text{Bei}(x), \quad \dots (66)$$

$$K_0(x\sqrt{i}) = \text{Ker}(x) + i \text{Kei}(x), \quad \dots (67)$$

$$\text{where} \quad \text{Ber}(x) = \frac{(\frac{1}{2}x)^4}{(|2|)^2} + \frac{(\frac{1}{2}x)^8}{(|4|)^2} \quad \dots (68)$$

$$\text{and} \quad \text{Bei}(x) = \left(\frac{x}{2} \right)^2 - \frac{(\frac{1}{2}x)^6}{(|3|)^2} + \frac{(\frac{1}{2}x)^{10}}{(|5|)^2} - \dots \quad \dots (69)$$

Problem 25. Derive Bessel's equation from that of Legendre.

We know that Legendre's equation is

$$(1-x^2) \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + n(n+1)y = 0 \quad \dots (1)$$

Differentiating it N times, we have by Leibnitz's theorem,

$$(1-x^2) \frac{d^{N+2}y}{dx^{N+2}} + N(-2x) \frac{d^{N+1}y}{dx^{N+1}} + \frac{N(N-1)}{1 \cdot 2} (-2) \cdot \frac{d^N y}{dx^N} \\ - 2x \frac{d^{N+1}y}{dx^{N+1}} + N(-2) \cdot \frac{d^N y}{dx^N} + n(n+1) \cdot \frac{d^N y}{dx^N} = 0,$$

$$\text{i.e., } (1-x^2) \frac{d^{N+2}y}{dx^{N+2}} - 2(N+1)x \frac{d^{N+1}y}{dx^{N+1}} \\ + \{n(n+1) - N(N+1)\} \frac{d^N y}{dx^N} = 0.$$

If we put $z = \frac{d^N y}{dx^N}$, the equation (2) becomes

$$(1-x^2) \frac{d^2 z}{dx^2} - 2(N+1)x \frac{dz}{dx} + \{n(n+1) - N(N+1)\} z = 0 \dots (1)$$

$$\text{Now put } Y = (1-x^2)^{\frac{1}{2}N} z$$

$$\text{i.e., } z = Y (1-x^2)^{-\frac{1}{2}N}$$

$$\text{so that } \frac{dz}{dx} = (1-x^2)^{-\frac{1}{2}N} \frac{dY}{dx}$$

$$+ Y \left(-\frac{N}{2} \right) (1-x^2)^{-\frac{1}{2}N-1} (-2x)$$

$$= (1-x^2)^{-\frac{1}{2}N} \frac{dY}{dx} + Nx (1-x^2)^{-\frac{1}{2}N-1} Y$$

$$\text{and } \frac{d^2 z}{dx^2} = (1-x^2)^{-\frac{1}{2}N} \frac{d^2 Y}{dx^2} + 2Nx (1-x^2)^{-\frac{1}{2}N-1} \frac{dY}{dx} \\ + N \left\{ (1-x^2)^{-\frac{1}{2}N-1} + x (1-x^2)^{-\frac{1}{2}N-2} \left(-\frac{N}{2} - 1 \right) (-2x) \right\} Y$$

$$\text{or } \frac{d^2 z}{dx^2} = (1-x^2)^{-\frac{1}{2}N} \frac{d^2 Y}{dx^2} + 2Nx (1-x^2)^{-\frac{1}{2}N-1} \frac{dY}{dx} \\ + N \{ 1 + (N+1)x^2 \} (1-x^2)^{-\frac{1}{2}N-2} Y.$$

$$\text{Substituting these values in (3), the transformed equation is} \\ (1-x^2) \frac{d^2 Y}{dx^2} + \{2Nx - 2(N+1)x\} \frac{dY}{dx} + \left[n(n+1) - N(N+1) \right. \\ \left. - \frac{2Nx^2(N+1)}{1-x^2} + \frac{N\{1+(N+1)x^2\}}{1-x^2} \right] Y = 0$$

[on dividing throughout by $(1-x^2)^{-\frac{1}{2}N}$]

$$\text{or } (1-x^2) \frac{d^2 Y}{dx^2} - 2x \frac{dY}{dx} + \left\{ n(n+1) - \frac{N}{(1-x^2)} \right\} Y = 0. \dots (4)$$

Now in order to change the independent variable, put

$$X^2 = n^2 (1-x^2), \text{ i.e. } 1-x^2 = \frac{X^2}{n^2}.$$

$$\text{and } X = n\sqrt{1-x^2} \text{ gives } \frac{dX}{dx} = \frac{-nx}{\sqrt{1-x^2}},$$

$$\text{so that } \frac{dY}{dx} = \frac{dY}{dX} \cdot \frac{dX}{dx} = \frac{-nx}{\sqrt{1-x^2}} \frac{dY}{dX}$$

$$\text{and } \frac{d^2 Y}{dx^2} = \frac{d}{dx} \left(\frac{dY}{dx} \right) = \frac{d}{dx} \left(\frac{-nx}{\sqrt{1-x^2}} \frac{dY}{dX} \right)$$

$$\text{or } \frac{d^2 Y}{dx^2} = -n \left[\left\{ \frac{\sqrt{1-x^2} + \frac{x^2}{\sqrt{1-x^2}}}{1-x^2} \right\} \frac{dY}{dX} + \frac{x}{\sqrt{1-x^2}} \frac{d^2 Y}{dX^2} \frac{dX}{dx} \right] \\ = -n \left[\frac{1}{(1-x^2)^{3/2}} \frac{dY}{dX} - \frac{nx^2}{1-x^2} \frac{d^2 Y}{dX^2} \right]$$

With these substitutions (4) becomes

$$n^3 x^2 \frac{d^2 Y}{dX^2} + \left[\frac{-n}{\sqrt{1-x^2}} + \frac{2nx^2}{\sqrt{1-x^2}} \right] \frac{dY}{dX} + \left\{ n(n+1) - \frac{N^2}{1-x^2} \right\} Y = 0$$

$$\text{i.e. } \left(1 - \frac{X^2}{n^2} \right) \frac{d^2 Y}{dX^2} + \frac{(2x^2-1)}{X} \frac{dY}{dX} + \left\{ 1 + \frac{1}{n} - \frac{N^2}{X^2} \right\} Y = 0 \\ \therefore X^2 = n^2 (1-x^2)$$

$$\text{or } \left(1 - \frac{X^2}{n^2} \right) \frac{d^2 Y}{dX^2} + \left(1 - \frac{2x^2}{n^2} \right) \frac{1}{X} \frac{dY}{dX} + \left(1 + \frac{1}{n} - \frac{N^2}{X^2} \right) Y = 0.$$

Proceeding to the limit when $n \rightarrow \infty$, we have

$$\frac{d^2 Y}{dX^2} + \frac{1}{X} \frac{dY}{dX} + \left(1 - \frac{N^2}{X^2} \right) Y = 0$$

which is clearly the Bessel's differential equation.

Problem 26. Integrate in series the Bessel's equation of zeroeth order. (Nagpur, 1965)

Bessel's equation is

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + \left(1 - \frac{n^2}{x^2} \right) y = 0.$$

If $n=0$, this equation reduces to

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + y = 0, \dots (1)$$

which is known as Bessel's differential equation of zeroeth order.

In order to find the series solution of (1), let us assume that its series solution is of the form

$$y = \sum_{r=0}^{\infty} a_r x^{k+r},$$

$$\text{so that } \frac{dy}{dx} = \sum_{r=0}^{\infty} a_r (k+r) x^{k+r-1}$$

$$\text{and } \frac{d^2 y}{dx^2} = \sum_{r=0}^{\infty} a_r (k+r) (k+r-1) x^{k+r-2}.$$

Putting these values in (1), we get

$$\sum_{r=0}^{\infty} a_r (k+r) (k+r-1) x^{k+r-2} + \sum_{r=0}^{\infty} a_r (k+r) x^{k+r-2} + \sum_{r=0}^{\infty} a_r x^{k+r} = 0$$

$$\text{or } \sum_{r=0}^{\infty} [(k+r)(k+r-1) + k+r] x^{k+r-2} + x^{k+r} a_r = 0 \quad \dots (1)$$

The equation (2) being an identity, we can equate the coefficients of various powers of x to zero.

Equating to zero the coefficient of lowest power of x , i.e., x^{k-2} by putting $r=0$ in (2), we have

$$\{k(k-1) + k\} a_0 = 0, \text{ i.e., } k^2 a_0 = 0.$$

Being the coefficient of first term $a_0 \neq 0$ and therefore $k=0$.

Now equating to zero the coefficient of x^{k-1} by putting $r=1$ in (2), we get

$$\{(k+1)k + k+1\} a_1 = 0, \text{ i.e., } (k+1)^2 a_1 = 0,$$

which gives $a_1 = 0$, since $(k+1)^2 \neq 0$ by virtue of (3).

Equating to zero the coefficient of general term, i.e., x^{k+r} in (2), we get

$$\{(k+r+2)(k+r+1) + k+r+2\} a_{r+2} + a_r = 0,$$

$$\text{i.e., } (k+r+2)^2 a_{r+2} + a_r = 0$$

$$\text{or } a_{r+2} = -\frac{a_r}{(r+2)^2}, \quad \dots (4)$$

since by (3) $k=0$.

Now $a_1 = 0$ and putting $r=1, 3, 5, \dots$ in (4), we have

$$a_3 = -\frac{1}{2^2} a_1 = 0.$$

Similarly, $a_5 = 0, a_7 = 0,$

$$\text{i.e., } a_1 = a_3 = a_5 = a_7 = \dots = 0.$$

Again putting $r=0, 2, 4, \dots$ in (4), we get

$$a_2 = -\frac{a_0}{2^2}, a_4 = -\frac{1}{4^2} a_2 = -\frac{1}{2^2 \cdot 4^2} a_0,$$

$$a_6 = -\frac{a_0}{6^2 \cdot 4^2 \cdot 2^2}, a_8 = -\frac{a_0}{8^2 \cdot 6^2 \cdot 4^2 \cdot 2^2} \text{ etc...}$$

Substituting these values in $y = \sum_{r=0}^{\infty} a_r x^{k+r}$, we get the required series solution as

$$y = a_0 \left[1 - \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} - \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots \right]$$

Note. This solution is denoted by $J_0(x)$ when $a_0 = 1$.

Problem 27. Solve the equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + \left(x^2 - \frac{1}{9} \right) y = 0 \quad (\text{Agra, 1964})$$

The given equation may be written as

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + \left(1 - \frac{1}{x^2} \right) y = 0$$

(on dividing throughout by x^2).

This is the same differential equation as discussed in § 11.4, [A] under the head Bessel's differential equation. Here $n^2 = \frac{1}{9}$.

Problem 28. Solve the differential equation

$$\frac{d^2 \psi}{dx^2} + (E - x^2) \psi = 0 \text{ such that } \psi \rightarrow 0 \text{ as } |x| \rightarrow \infty.$$

(Agra, 1962)

The given differential equation is

$$\frac{d^2 \psi}{dx^2} + (E - x^2) \psi = 0 \quad \dots (1)$$

Put

$$\psi = v e^{-x^2/2}$$

so that

$$\frac{d\psi}{dx} = e^{-x^2/2} \frac{dv}{dx} - vx e^{-x^2/2}$$

and

$$\frac{d^2 \psi}{dx^2} = e^{-x^2/2} \frac{d^2 v}{dx^2} - 2x e^{-x^2/2} \frac{dv}{dx} - v e^{-x^2/2} + vx^2 e^{-x^2/2}.$$

Substituting these values in (1) and dividing throughout by $e^{-x^2/2}$, we get

$$\frac{d^2 v}{dx^2} - 2x \frac{dv}{dx} + (E - 1) v = 0. \quad \dots (2)$$

Let its series solution be

$$v = \sum_{r=0}^{\infty} a_r x^{k-r}, \quad \dots (3)$$

then

$$\frac{dv}{dx} = \sum_{r=0}^{\infty} a_r (k-r) x^{k-r-1}$$

and

$$\frac{d^2 v}{dx^2} = \sum_{r=0}^{\infty} a_r (k-r)(k-r-1) x^{k-r-2}.$$

Substituting these values in (3), we get

$$\sum_{r=0}^{\infty} a_r (k-r)(k-r-1) x^{k-r-2} - 2 \sum_{r=0}^{\infty} a_r (k-r) x^{k-r-1} + (E-1) \sum_{r=0}^{\infty} a_r x^{k-r} = 0.$$

$$\text{or } \sum_{r=0}^{\infty} \{[(k-r)(k-r-1)] x^{k-r-2} + \{(-1)(k-r) + (E-1)\} x^{k-r}\} a_r = 0. \quad \dots (4)$$

The relation (4) being an identity, the coefficients of various powers of x can be equated to zero.

Let us first equate the coefficient of x^k the highest power of x [by putting $r=0$ in (2)], to zero; then we get

$$(E-1-2k)a_0=0.$$

This gives $k = \frac{E-1}{2}, \dots (3)$

since $a_0 \neq 0$, being the coefficient of first term of the series.

Now equating the coefficient of x^{k-1} to zero, by putting $r=1$ in (4), we get

$$\{E-1-2(k-1)a_1\}=0.$$

Here $E-1-2(k-1) \neq 0$ by virtue of (5) and therefore, we have $a_1=0$.

Further equating to zero the coefficient of x^{k-r} , the general term, in (4), we have

$$(k-r+2)(k-r+1)a_{r-2} + \{(E-1)-2(k-r)\}a_r = 0$$

or
$$a_r = \frac{(k-r+2)(k-r+1)}{E-1-2(k-r)} a_{r-2}$$

$$= \frac{\left(\frac{E-1}{2}-r+2\right)\left(\frac{E-1}{2}-r+1\right)}{E-1-2\left(\frac{E-1}{2}-r\right)} a_{r-2} \text{ from (5)}$$

$$= -\frac{(E+3-2r)(E+1-2r)}{8r} a_{r-2}.$$

Putting $r=2, 3, 4, 5, 6 \dots$, we get

$$a_2 = -\frac{(E-1)(E-3)}{8 \cdot 2} a_0.$$

$$a_3 = -\frac{(E-3)(E-5)}{8 \cdot 3} a_1 = 0 \text{ as } a_1 = 0.$$

Similarly $a_5 = a_7 = a_9 = \dots = 0$

and
$$a_4 = -\frac{(E-5)(E-7)}{8 \cdot 4} a_2$$

$$= \frac{(E-1)(E-3)(E-5)(E-7)}{8^2 \cdot 2 \cdot 4} a_0,$$

$$a_6 = -\frac{(E-9)(E-11)}{8 \cdot 6} a_4$$

$$= -\frac{(E-1)(E-3)(E-5)(E-7)(E-9)(E-11)}{8^3 \cdot 2 \cdot 4 \cdot 6} a_0 \text{ etc.}$$

Substituting the values of these coefficients in (3), we get

$$v = a_0 \left[x^{(E-1)/2} - \frac{(E-1)(E-3)}{8 \cdot 2} x^{(E-5)/2} + \frac{(E-1)(E-3)(E-5)(E-7)}{8 \cdot 2 \cdot 4} x^{(E-9)/2} - \dots \right]$$

Putting this value of v in $\psi = ve^{-x^2/2}$, the series solution of the given differential equation (1) is

$$\psi = a_0 e^{-x^2/2} \left[x^{(E-1)/2} - \frac{(E-1)(E-3)}{8 \cdot 2} x^{(E-5)/2} + \frac{(E-1)(E-3)(E-5)(E-7)}{8^2 \cdot 2 \cdot 4} x^{(E-9)/2} - \dots \right]$$

Problem 29. When calculating the dependence of the current density on the distance ρ from the axis, we come across a scalar equation of the following form in cylindrical co-ordinates:

$$\frac{\partial^2 u}{\partial \rho^2} + \frac{1}{\rho} \frac{\partial u}{\partial \rho} = \frac{4\pi\sigma\mu}{c^2} \frac{\partial u}{\partial t}.$$

Find a solution for u which is a periodic function of the time.

Suppose that $u = \phi(\rho) e^{i\omega t}$ is a solution; then

$$\frac{\partial u}{\partial \rho} = \frac{d\phi}{d\rho} e^{i\omega t}$$

$$\frac{\partial^2 u}{\partial \rho^2} = \frac{d^2 \phi}{d\rho^2} e^{i\omega t}$$

$$\frac{\partial u}{\partial t} = i\omega \phi(\rho) e^{i\omega t}.$$

Substituting these values in the given equation, we get

$$\frac{d^2 \phi}{d\rho^2} + \frac{1}{\rho} \frac{d\phi}{d\rho} = i\omega \frac{4\pi\sigma\mu}{c^2} \phi$$

$$\frac{d^2 \phi}{d\rho^2} + \frac{1}{\rho} \frac{d\phi}{d\rho} + k^2 \phi = 0, \dots (1)$$

$$\text{where } k^2 = -\frac{4\pi\sigma\mu\omega}{c^2}.$$

Now in order to change the independent variable, let us put

$$x = k\rho, \text{ so that } \frac{dx}{d\rho} = k.$$

$$\frac{d\phi}{d\rho} = \frac{d\phi}{dx} \cdot \frac{dx}{d\rho} = k \frac{d\phi}{dx}$$

$$\frac{d^2 \phi}{d\rho^2} = \frac{d}{d\rho} \left(\frac{d\phi}{d\rho} \right) = \frac{d}{d\rho} \left(k \frac{d\phi}{dx} \right) = k \frac{d}{dx} \left(\frac{d\phi}{dx} \right) = k^2 \frac{d^2 \phi}{dx^2}.$$

With these substitutions (1) reduces to

$$\frac{d^2 \phi}{dx^2} + \frac{1}{x} \frac{d\phi}{dx} + \phi = 0,$$

which is the Bessel's differential equation of zeroth order, as we have already discussed in Problem 26. Its solution is

$$\phi = a_0 J_0(x) = a_0 \left[1 - \frac{x^2}{2} + \frac{x^4}{2 \cdot 4} - \frac{x^6}{2 \cdot 4 \cdot 6^2} + \dots \right].$$

Hence $u = a_0 J_0(x) \cdot e^{i\omega t}$, where a_0 is the amplitude factor
 $= a_0 J_0(k\rho) e^{i\omega t}$

$$= a_0 J_0 \left\{ \sqrt{(-i)} \sqrt{(4\pi\sigma\mu\omega)} \cdot \frac{\rho}{c} \right\} e^{i\omega t}$$

$$= a_0 \left[\text{ber} \left\{ \sqrt{(4\pi\sigma\mu\omega)} \cdot \frac{\rho}{c} \right\} + i \text{bei} \left\{ \sqrt{(4\pi\sigma\mu\omega)} \cdot \frac{\rho}{c} \right\} \right] e^{i\omega t}$$

which is the required solution.

Note. *Ber and Bei functions.*

The real and imaginary parts of the function $J_0(x\sqrt{-i})$, are known as 'ber (x)' and 'bei (x)', i.e.,

$$R[J_0 x \sqrt{-i}] = \text{ber } x,$$

$$I[J_0 x \sqrt{-i}] = \text{bei } x.$$

Problem 30. Find the integral of the differential equation of cylindrical wave

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} = \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2}.$$

Suppose that

$$u = v(r) e^{i\omega t}.$$

$$\therefore \frac{\partial u}{\partial r} = \frac{dv}{dr} e^{i\omega t} \text{ and } \frac{\partial^2 u}{\partial r^2} = \frac{d^2 v}{dr^2} e^{i\omega t}.$$

Also

$$\frac{\partial u}{\partial t} = i\omega v(r) e^{i\omega t}$$

and

$$\frac{\partial^2 u}{\partial t^2} = -\omega^2 v(r) e^{i\omega t}.$$

Substituting these values in the given wave equation, we get

$$\frac{d^2 v}{dr^2} + \frac{1}{r} \frac{dv}{dr} + \frac{\omega^2}{c^2} v = 0. \quad \dots (1)$$

$$\text{Now put } r = \frac{cx}{\omega}, \text{ so that } \frac{dx}{dr} = \frac{\omega}{c}.$$

$$\therefore \frac{dv}{dr} = \frac{dv}{dx} \cdot \frac{dx}{dr} = \frac{\omega}{c} \frac{dv}{dx}$$

$$\text{and } \frac{d^2 v}{dr^2} = \frac{d}{dr} \left(\frac{\omega}{c} \frac{dv}{dx} \right) = \frac{\omega^2}{c^2} \frac{d^2 v}{dx^2}.$$

With these substitutions the equation (1) transforms to

$$\frac{d^2 v}{dx^2} + \frac{1}{x} \frac{dv}{dx} + v = 0,$$

which is the Bessel's differential equation of zeroth order and therefore its solution is

$$v = a_0 J_0(x) \text{ (as in the previous problem)}$$

Hence the required solution is

$$u = a_0 J_0 \left(\frac{r\omega}{c} \right) e^{i\omega t}.$$

Problem 31. Prove that $\sqrt{\left(\frac{\pi x}{2}\right)} J_{3/2}(x) = \frac{\sin x}{x} \cos x.$

(Agra, 1966)

We have

$$J_n(x) = \sum_{s=0}^{\infty} \frac{(-1)^s}{s! (n+s)!} \left(\frac{x}{2}\right)^{n+2s}$$

$$= \frac{x^n}{2^n n!} - \frac{x^{n+2}}{2^{n+2} 1! (n+1)!} + \frac{x^{n+4}}{2^{n+4} 2! (n+2)!} - \dots \text{since } 0! = 1$$

$$= \frac{x^n}{2^n n!} \left[1 - \frac{x^2}{2 \cdot 1 \cdot (n+1)} + \frac{x^4}{2^2 \cdot 1 \cdot 2 \cdot (n+1)(n+2)} - \dots \right]$$

$$= \frac{x^n}{2^n \Gamma(n+1)} \left[1 - \frac{x^2}{2 \cdot (2n+2)} + \frac{x^4}{2 \cdot 4 \cdot (2n+2)(2n+3)} - \dots \right],$$

$$\therefore \Gamma(n+1) = n!$$

Putting $n = \frac{3}{2}$ on either side, we get

$$J_{3/2}(x) = \frac{x^{3/2}}{2^{3/2} \Gamma_{\frac{3}{2}}} \left[1 - \frac{x^2}{2 \cdot 5} + \frac{x^4}{2 \cdot 4 \cdot 5 \cdot 7} - \frac{x^6}{2 \cdot 4 \cdot 6 \cdot 5 \cdot 7 \cdot 9} + \dots \right]$$

$$= \frac{x^{3/2}}{2^{3/2} \cdot \frac{1}{2} \sqrt{\pi}} \left[1 - \frac{x^2}{2 \cdot 5} + \frac{x^4}{2 \cdot 4 \cdot 5 \cdot 7} - \frac{x^6}{2 \cdot 4 \cdot 6 \cdot 5 \cdot 7 \cdot 9} + \dots \right],$$

$$\sqrt{\left(\frac{\pi x}{2}\right)} \cdot J_{3/2}(x) = \frac{1}{3} \left[x^2 - \frac{x^4}{2 \cdot 5} + \frac{x^6}{2 \cdot 4 \cdot 5 \cdot 7} - \frac{x^8}{2 \cdot 4 \cdot 6 \cdot 5 \cdot 7 \cdot 9} + \dots \right]$$

$$= \frac{x^2}{3} - \frac{x^4}{2 \cdot 3 \cdot 5} + \frac{x^6}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 7} - \frac{x^8}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 9} + \dots$$

$$= \frac{2x^2}{|3|} - \frac{4x^4}{|5|} + \frac{6x^6}{|7|} - \frac{8x^8}{|9|} + \dots$$

$$= \frac{(3-1)x^2}{|3|} - \frac{(5-1)x^4}{|5|} + \frac{(7-1)x^6}{|7|} - \frac{(9-1)x^8}{|9|} + \dots$$

$$= \frac{x^2}{|2|} - \frac{x^4}{|3|} + \frac{x^6}{|4|} - \frac{x^8}{|5|} + \frac{x^{10}}{|6|} - \frac{x^{12}}{|7|} + \dots$$

$$= -\frac{x^2}{|3|} + \frac{x^4}{|5|} - \frac{x^6}{|7|} + \dots + \frac{x^2}{|2|} - \frac{x^4}{|4|} + \frac{x^6}{|6|} - \dots$$

$$= \frac{1}{x} \left[x - \frac{x^3}{|3|} + \frac{x^5}{|5|} - \frac{x^7}{|7|} + \dots \right]$$

$$= \frac{1}{x} \left[1 - \frac{x^2}{|2|} + \frac{x^4}{|4|} - \frac{x^6}{|6|} + \dots \right]$$

$$= \frac{1}{x} \sin x - \cos x.$$

Problem 32. Prove that

$$\frac{J_{n+1}(x)}{J_n(x)} = \frac{x/2}{(n+1) - (n+2)} - \frac{(x/2)^2}{(n+3) - \dots}$$

We have the recurrence relation,

$$J_{n-1}(x) + J_{n+1}(x) = \frac{2n}{x} J_n(x)$$

or

$$J_{n-1}(x) = \frac{2n}{x} J_n(x) - J_{n+1}(x).$$

Replacing n by $(n+1)$, this becomes

$$J_n(x) = \frac{2(n+1)}{x} J_{n+1}(x) - J_{n+2}(x).$$

Now $\frac{J_{n+1}(x)}{J_n(x)} = \frac{1}{\frac{J_n(x)}{J_{n+1}(x)}} = \frac{1}{\frac{2(n+1)}{x} J_{n+1}(x) - J_{n+2}(x)} \quad \text{from (1)}$

$$= \frac{1}{\frac{2(n+1)}{x} - \frac{J_{n+2}(x)}{J_{n+1}(x)}} = \frac{1}{\frac{2(n+1)}{x} - \frac{1}{\frac{J_{n+1}(x)}{J_{n+2}(x)}}}$$

$$= \frac{1}{\frac{2(n+1)}{x} - \frac{1}{\frac{2(n+2)}{x} J_{n+2}(x) - J_{n+3}(x)}} \quad \text{by (1)}$$

$$= \frac{1}{\frac{2(n+1)}{x} - \frac{1}{\frac{2(n+2)}{x} - \frac{1}{\frac{J_{n+2}(x)}{J_{n+3}(x)}}}}$$

$$= \frac{1}{\frac{2(n+1)}{x} - \frac{1}{\frac{2(n+2)}{x} - \frac{1}{\frac{2(n+3)}{x} - \dots}}}$$

$$= \frac{x/2}{(n+1) - \frac{x/2}{2(n+2) - \frac{1}{2(n+3) - \dots}}}$$

$$\text{by repeated application of (1)}$$

$$= \frac{x/2}{(n+1) - \frac{(x/2)^2}{(n+2) - \frac{x/2}{2(n+3) - \dots}}}$$

$$= \frac{x/2}{(n+1) - \frac{(x/2)^2}{(n+2) - \frac{(x/2)^3}{(n+3) - \dots}}}$$

8.5. HYPERGEOMETRIC OR GAUSS EQUATION AND FUNCTIONS

[A] Hypergeometric or Gauss' Differential Equation.

This equation is of the form

$$x(1-x) \frac{d^2 y}{dx^2} + \{\gamma - (\alpha + \beta + 1)x\} \frac{dy}{dx} - \alpha\beta y = 0 \quad \dots (1)$$

where α, β, γ are parameteric constants.

Here $x=0, x=1$ and $x=\infty$ are the singularities, since on dividing (1) by $x(x-1)$ we observe that coefficients of $\frac{dy}{dx}$ and y become infinite when $x=0, 1$ or ∞ . Thus we can integrate (1) in series about $x=0$ or $x=1$ or $x=\infty$. We therefore discuss the series integration in three cases.

Case [a₁]. When $x=0$, then taking the series solution of (1) as

$$y = \sum_{r=0}^{\infty} a_r x^{k+r} \quad \dots (2)$$

We have already discussed the solution in §8.3 $E[e_3]$ and obtained the solution as

$$y = AF(\alpha, \beta, \gamma, x) + Bx^{1-\gamma} F(\alpha+1-\gamma, \beta+1-\gamma, 2-\gamma, x) \quad \dots (3)$$

where A and B are arbitrary constants,

$$\text{and } F(\alpha, \beta, \gamma, x) = 1 + \frac{\alpha\beta}{1\cdot\gamma} x + \frac{\alpha(\alpha+1)(\beta+1)}{1\cdot2\cdot\gamma(\gamma+1)} x^2 + \frac{\alpha(\alpha+1)(\alpha+2)\beta(\beta+1)(\beta+2)}{1\cdot2\cdot3\cdot\gamma(\gamma+1)(\gamma+2)} x^3 + \dots$$

Case [a₂]. When $x=1$, is the singularity, then the series solution is obtained by developing the series about $x=1$, by making a substitution

$$X = 1 - x \quad \text{in (1)} \quad \dots (4)$$

$$\therefore \frac{dy}{dx} = \frac{dy}{dX} \frac{dX}{dx} = -\frac{dy}{dX}, \quad \text{since by (1) } \frac{dX}{dx} = -1$$

$$\text{and } \frac{d^2 y}{dx^2} = \frac{d}{dX} \left(-\frac{dy}{dX} \right) \frac{dX}{dx} = \frac{d^2 y}{dX^2}$$

Substituting these values in (1) and arranging we get

$$(X^2 - X) \frac{d^2 y}{dX^2} + [(1 + \alpha + \beta)X + (\gamma - 1 - \alpha - \beta)] \frac{dy}{dX} + \alpha\beta y = 0 \quad \dots (5)$$

which is similar to (1) except that γ is replaced by $1 + \alpha + \beta - \gamma$ and x by $1 - x$ and hence by (3) the solution in this case becomes,

$$y = AF(\alpha, \beta, 1 + \alpha + \beta - \gamma, 1 - x) + B(1 - x)^{\gamma - \alpha - \beta} F(\gamma - \alpha, \gamma - \beta, \gamma - \alpha - \beta + 1, 1 - x) \quad \dots (6)$$

Case [a₃]. When $x=\infty$ gives a singularity, then the series solution of (1) is obtained by developing the series about $x=\infty$, by making a substitution

$$x = \frac{1}{X} \text{ in (1)} \quad \dots (7)$$

$$\text{So that } \frac{dy}{dx} = \frac{dy}{dX} \frac{dX}{dx} = -\frac{1}{x^2} \frac{dy}{dX} = -X^2 \frac{dy}{dX} \text{ etc.}$$

$\therefore$ (1)-becomes

$$X^2(1-X) \frac{d^2y}{dX^2} + [2X(1-X) - (1+\alpha+\beta)X + \gamma X^2] \frac{dy}{dX} + \alpha\beta y = 0 \quad \dots (8)$$

Let its series solution be

$$y = \sum_{r=0}^{\infty} a_r X^{k+r}, a_0 \neq 0 \quad \dots (9)$$

$$\text{So that } \frac{dy}{dX} = \sum_{r=0}^{\infty} a_r (k+r) X^{k+r-1}$$

$$\frac{d^2y}{dX^2} = \sum_{r=0}^{\infty} a_r (k+r) (k+r-1) X^{k+r-2}$$

Substituting these values in (8) we get the identity

$$\sum_{r=0}^{\infty} [(k+r)(k+r-1) + (2-1-\alpha-\beta)(k+r) + \alpha\beta] X^{k+r} - \{(k+r)(k+r-1) + (2-\gamma)(k+r)\} X^{k+r+1} a_r \equiv 0 \quad \dots (10)$$

Equating to zero the coefficient of X^k (the first term) in (10), we get

$$[k(k-1) + (1-\alpha-\beta)k + \alpha\beta] a_0 = 0$$

$$\therefore a_0 \neq 0, \therefore k(k-1) + (1-\alpha-\beta)k + \alpha\beta = 0$$

$$\text{or } k^2 - (\alpha+\beta)k + \alpha\beta = 0$$

$$\text{or } (k-\alpha)(k-\beta) = 0 \text{ giving } k = \alpha, \beta \quad \dots (11)$$

Again equating to zero the coefficient of X^{k+r} in (10), we find the recurrence relation as

$$\{(k+r)(k+r-1) + (1-\alpha-\beta)(k+r) + \alpha\beta\} a_r - \{(k+r-1)(k+r-2) + (2-\gamma)(k+r-1)\} a_{r-1} = 0$$

$$\text{i.e. } (k+r-\alpha)(k+r-\beta) a_r - (k+r-1)(k+r-\gamma) a_{r-1} = 0$$

$$\text{or } a_r = \frac{(k+r-1)(k+r-\gamma)}{(k+r-\alpha)(k+r-\beta)} a_{r-1} \quad \dots (12)$$

$$\text{when } k = \alpha, (10) \text{ gives } a_r = \frac{(\alpha+r-1)(\alpha+r-\gamma)}{r(\alpha+r-\beta)} a_{r-1}$$

$$\text{So that } a_1 = \frac{\alpha(\alpha+1-\gamma)}{1(\alpha+1-\beta)} a_0$$

$$a_2 = \frac{\alpha(\alpha+1)(\alpha+1-\gamma)(\alpha+2-\gamma)}{1.2(\alpha+1-\beta)(\alpha+2-\beta)} a_0$$

$$a_3 = \frac{\alpha(\alpha+1)(\alpha+2)(\alpha+1-\gamma)(\alpha+2-\gamma)(\alpha+3-\gamma)}{1.2.3(\alpha+1-\beta)(\alpha+2-\beta)(\alpha+3-\beta)} a_0 \text{ etc.}$$

$\therefore$ Solution is $y = a_0 X^\alpha F(\alpha, \alpha+1-\gamma, \alpha+1-\beta, X)$

$$= a_0 x^{-\alpha} F\left(\alpha, \alpha+1-\gamma, \alpha+1-\beta, \frac{1}{x}\right) \quad \dots (13)$$

and similarly when $k = \beta$, the solution is

$$y = a_0 x^{-\beta} F\left[\beta, \beta+1-\gamma, \beta+1-\alpha, \frac{1}{x}\right] \quad \dots (14)$$

Hence the complete integral is

$$y = Ax^{-\alpha} F(\alpha, \alpha+1-\gamma, \alpha+1-\beta, \frac{1}{x}) + Bx^{-\beta} F\left[\beta, \beta+1-\gamma, \beta+1-\alpha, \frac{1}{x}\right] \quad \dots (15)$$

where A and B are arbitrary constants.

We thus have found all the three possible solutions of Hypergeometric or Gauss equation, i.e.,

(i) for $x=0$, exponents are $0, 1-\gamma$ by (3)

(ii) for $x=\infty$, exponents are $-\alpha, -\beta$ by (13)

(iii) for $x=1$, exponents are $0, \gamma-\alpha-\beta$ by (6)

These results may be shown by a scheme as follows:

$$y = P \begin{bmatrix} 0 & \infty & 1 \\ 0 & \alpha & 0 \\ 1-\gamma & \beta & \gamma-\alpha-\beta \end{bmatrix} x \quad \dots (16)$$

where the R.H.S. is said to be the *Riemann-P function of the equation*.

$$\text{In symbolic form, } F(\alpha, \beta, \gamma; x) = F \begin{bmatrix} \alpha, \beta \\ \gamma \end{bmatrix} x \quad \dots (17)$$

which is known as *Hypergeometric function*.

$$\text{We also denote } F(\alpha, \beta, \gamma, x) \text{ by } \sum_{k=0}^{\infty} \frac{(\alpha)_k (\beta)_k}{(\gamma)_k k!} x^k \quad \dots (18)$$

[B] Particular Cases of Hypergeometric Series.

$$(i) \text{ we have } (1+x)^n = 1 + nx + \frac{n(n-1)}{1.2} x^2 + \frac{n(n-1)(n-2)}{1.2.3} x^3 + \dots = F(-n, 1, 1, -x)$$

$$(ii) \log(1+x) = x - \frac{1}{2} x^2 + \frac{x^3}{3} - \frac{x^4}{4} + \dots = x \left[1 - \frac{1}{2} x + \frac{1}{3} x^2 - \frac{1}{4} x^3 + \dots \right] = x F(1, 1, 2, -x)$$

$$(iii) \sin^{-1} x = x + \frac{x^3}{|3|} + \frac{3^2 x^5}{|5|} + \frac{3^2.5^2}{|7|} x^7 + \dots = x \left[1 + \frac{\frac{1}{2}. \frac{1}{2}}{1. \frac{3}{2}} x^2 + \frac{\frac{1}{2}. \frac{3}{2}. \frac{1}{2}. \frac{3}{2}}{1.2. \frac{5}{2}. \frac{3}{2}} x^4 + \dots \right]$$

$$= xF\left[\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, x^2\right]$$

$$(iv) \tan^{-1}x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$$

$$= x\left[1 - \frac{x^2}{3} + \frac{x^4}{5} - \frac{x^6}{7} + \dots\right]$$

$$= x\left[1 + \frac{1 \cdot \frac{1}{2}}{1 \cdot \frac{3}{2}}(-x^2) + \frac{1 \cdot 2 \cdot \frac{1}{2} \cdot \frac{3}{2}}{1 \cdot 2 \cdot \frac{5}{2} \cdot \frac{3}{2}}(-x^2)^2 + \dots\right]$$

$$= xF\left[1, \frac{1}{2}, \frac{3}{2}, -x^2\right]$$

$$(v) e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$= \lim_{n \rightarrow \infty} \left\{ 1 + \frac{n \cdot 1}{1 \cdot 1} \frac{x}{n} + \frac{n(n+1) \cdot 1 \cdot 2}{1 \cdot 2 \cdot 1 \cdot 2} \left[\frac{x}{n}\right]^2 + \dots \right\}$$

$$= \lim_{n \rightarrow \infty} F\left[n, 1, 1, \frac{x}{n}\right].$$

[C] Simple Properties of Hypergeometric Function.

(1) Symmetry property. The value of a hypergeometric function does not change by the interchange of parameters α and β .

$$\begin{aligned} \therefore F(\alpha, \beta, \gamma, x) &= \sum_{k=0}^{\infty} \frac{(\alpha)_k (\beta)_k}{(\gamma)_k k!} x^k \\ &= \sum_{k=0}^{\infty} \frac{(\beta)_k (\alpha)_k}{(\gamma)_k k!} x^k \\ &= F(\beta, \alpha, \gamma, x) \end{aligned}$$

$$\therefore F\left[\begin{matrix} \alpha, \beta \\ \gamma \end{matrix} x\right] = F\left[\begin{matrix} \beta, \alpha \\ \gamma \end{matrix} x\right]$$

... (19)

(2) Differentiation of Hypergeometric Functions. We have

$$F(\alpha, \beta, \gamma, x) = \sum_{k=0}^{\infty} \frac{(\alpha)_k (\beta)_k}{(\gamma)_k k!} x^k$$

Its differentiation w.r.t. x , yields,

$$\frac{d}{dx} F(\alpha, \beta, \gamma, x) = \sum_{k=0}^{\infty} k \frac{(\alpha)_k (\beta)_k}{k! (k-1)! (\gamma)_k} x^{k-1} = \sum_{k=1}^{\infty} \frac{(\alpha)_k (\beta)_k}{(k-1)! (\gamma)_k} x^{k-1}$$

$\therefore$ it vanishes for $k=0$

$$= \sum_{p=0}^{\infty} \frac{(\alpha)_{p+1} (\beta)_{p+1}}{p! (\gamma)_{p+1}} \text{ on putting } k-1=p.$$

$$= \sum_{p=0}^{\infty} \frac{\alpha(\alpha+1) \dots \alpha(\alpha+p) \beta(\beta+1) \dots \beta(\beta+p)}{p! \gamma(\gamma+1) \dots \gamma(\gamma+p+1)} \therefore (\alpha)_{p+1} = \alpha(\alpha+1) \dots \alpha(\alpha+p) \\ = \alpha(\alpha+1) \dots \alpha(\alpha+p-1) \alpha(\alpha+p) \\ = \alpha(\alpha+1)_p \text{ etc.}$$

$$= \frac{\alpha\beta}{\gamma} \sum_{p=0}^{\infty} \frac{(\alpha+1)_p (\beta+1)_p}{p! (\gamma+1)_p}$$

$$= \frac{\alpha\beta}{\gamma} F(\alpha+1, \beta+1, \gamma+1, x)$$

... (20)

$$\frac{d^2}{dx^2} F(\alpha, \beta, \gamma, x) = \frac{d}{dx} \left[\frac{\alpha\beta}{\gamma} F(\alpha+1, \beta+1, \gamma+1, x) \right]$$

$$= \frac{\alpha\beta}{\gamma} \frac{d}{dx} F(\alpha+1, \beta+1, \gamma+1, x)$$

$$= \frac{\alpha\beta}{\gamma} \cdot \frac{(\alpha+1)(\beta+1)}{\gamma+1} F(\alpha+1+1, \beta+1+1, \gamma+1+1, x)$$

by applying (20)

$$= \frac{\alpha(\alpha+1) \beta(\beta+1)}{\gamma(\gamma+1)} F(\alpha+2, \beta+2, \gamma+2, x)$$

... (21)

Repeating the process m times, we may have

$$\begin{aligned} \frac{d^m}{dx^m} F(\alpha, \beta, \gamma, x) &= \frac{\alpha(\alpha+1) \dots (\alpha+m-1) \beta(\beta+1) \dots (\beta+m-1)}{\gamma(\gamma+1) \dots (\gamma+m-1)} F(\alpha+m, \beta+m, \gamma+m, x) \\ &= \frac{(\alpha)_m (\beta)_m}{(\gamma)_m} F(\alpha+m, \beta+m, \gamma+m, x) \end{aligned}$$

... (22)

In symbolic form (22) can be written as

$$\frac{d^m}{dx^m} F\left[\begin{matrix} \alpha, \beta \\ \gamma \end{matrix} x\right] = \frac{(\alpha)_m (\beta)_m}{(\gamma)_m} \left[\begin{matrix} \alpha+m, \beta+m \\ \gamma+m \end{matrix} x\right]$$

COROLLARY 1. When $x=0$, $F(\alpha, \beta, \gamma, 0) = \lim_{x \rightarrow 0} \sum_{k=0}^{\infty} \frac{(\alpha)_k (\beta)_k}{k! (\gamma)_k} x^k$

$$= \lim_{x \rightarrow 0} \left[1 + \frac{\alpha\beta}{\gamma} x + \frac{\alpha(\alpha+1) \beta(\beta+1)}{1 \cdot 2 \cdot \gamma(\gamma+1)} x^2 + \dots \right]$$

$= 1$, since all the terms except the first vanish.

Similarly $F(\alpha+1, \beta+1, \gamma+1, 0) = 1$

Hence from (18), it follows that

$$\left[\frac{d}{dx} F(\alpha, \beta, \gamma, x) \right]_{x=0} = \frac{\alpha\beta}{\gamma} \quad \dots (23)$$

COROLLARY 2. Had the parameter α been a negative integer say $-N$, then we should have

$$F(-N, \beta, \gamma, x) = \sum_{r=0}^N \frac{(-N)_r (\beta)_r}{r! (\gamma)_r} x^r \quad \dots (24)$$

since it vanishes for $k=N, N+1, N+2, \dots$ etc.

Similarly if β were a negative integer say $-M$, then

$$F(\alpha, -M, \gamma, x) = \sum_{k=0}^M \frac{(\alpha)_k (-M)_k}{k! (\gamma)_k} x^k \quad \dots (25)$$

In case $\alpha = -N$ and $\gamma = -(N+M)$, N, M being positive integers, we have

$$F(-N, \beta, -N-M, x) = \sum_{k=0}^{\infty} \frac{(-N)_k (\beta)_k}{(-N-M)_k k!} x^k \quad \dots (26)$$

where $(-N)_k = (-N)(-N+1)\dots(-N+k-1) = (-1)^k \frac{|N|}{|N-k|}$

$$\text{Similarly } (-N-M)_k = (-1)^k \frac{|N+M|}{|N+M-k|}$$

$$\begin{aligned} \text{So that } \frac{(-N)_k}{(-N-M)_k} &= \frac{|N|}{|N-k|} \times \frac{|N+M-k|}{|N+M|} = \frac{|N|}{|N+M|} \\ &= \left[\left(1 - \frac{k}{N+M}\right) \left(1 - \frac{k}{N+M-1}\right) \dots \left(1 - \frac{k}{N+1}\right) \right] \end{aligned}$$

Hence (26) yields

$$F(-N, \beta, -N-M, x) = \sum_{k=0}^{\infty} \left[\left(1 - \frac{k}{N+M}\right) \left(1 - \frac{k}{N+M-1}\right) \dots \left(1 - \frac{k}{N+1}\right) \right] \frac{(\beta)_k}{k!} x^k \quad \dots (27)$$

Here it is notable that on the R.H.S. the terms do not vanish for $k=0, 1, 2, \dots, N$ and they vanish for $k=N+1, N+2, \dots, N+M$, but the terms do not vanish for $k=N+M+1, N+M+2, \dots$, since the term corresponding to $k=N+M+1$ is

$$\left(1 - \frac{N+M+1}{N+M}\right) \dots \left(1 - \frac{N+M+1}{N+1}\right) \cdot \frac{(\beta)_{N+M+1}}{|N+M+1|} x^{N+M+1}$$

Conclusively the series which stopped for $\alpha = -N$ or $\beta = -M$ at N th and M th terms respectively, starts again when $\alpha = -N$ and $\gamma = -(N+M)$ or likewise when $\beta = -M$ and $\gamma = -(N+M)$.

(3) Integral Formula for the Hypergeometric Functions.

$$\text{We have } F(\alpha, \beta, \gamma, x) = \sum_{k=0}^{\infty} \frac{(\alpha)_k (\beta)_k}{k! (\gamma)_k} x^k$$

$$\text{where } (\alpha)_k = \alpha(\alpha+1)(\alpha+2)\dots(\alpha+k-1) = \frac{|\alpha+k-1|}{|\alpha-1|} = \frac{\Gamma(\alpha+k)}{\Gamma(\alpha)}$$

$$\text{Similarly } (\beta)_k = \frac{\Gamma(\beta+k)}{\Gamma\beta}, (\gamma)_k = \frac{\Gamma(\gamma+k)}{\Gamma\gamma}$$

$$\text{Also we have } B(m, n) = \frac{\Gamma m \Gamma n}{\Gamma(m+n)} \text{ (Beta function)}$$

$$\begin{aligned} \therefore \frac{(\beta)_k}{(\gamma)_k} &= \frac{\Gamma(\beta+k)}{\Gamma\beta} \times \frac{\Gamma\gamma}{\Gamma(\gamma+k)} = \frac{\Gamma\gamma}{\Gamma\beta} \cdot \frac{\Gamma(\beta+k)}{\Gamma(\gamma+k)} \times \frac{\Gamma(\gamma-\beta)}{\Gamma(\gamma-\beta)} \\ &= \frac{\Gamma\gamma}{\Gamma\beta\Gamma(\gamma-\beta)} \cdot \frac{\Gamma(\beta+k)\Gamma(\gamma-\beta)}{\Gamma(\gamma+k)} \end{aligned}$$

$$= \frac{\Gamma\gamma}{\Gamma\beta\Gamma(\gamma-\beta)} \frac{\Gamma(\beta+k)\Gamma(\gamma-\beta)}{\Gamma(\beta+k)+(\gamma-\beta)} = \frac{\Gamma\gamma}{\Gamma\beta\Gamma(\gamma-\beta)} \cdot B(\gamma-\beta, \beta+k)$$

$$= \frac{\Gamma\gamma}{\Gamma\beta\Gamma(\gamma-\beta)} \int_0^1 (1-t)^{\gamma-\beta-1} t^{\beta+k-1} dt$$

$$\therefore B(P, Q) = \int_0^1 (1-x)^{P-1} x^{Q-1} dx$$

$$= \frac{1}{B(\beta, \gamma-\beta)} \int_0^1 (1-t)^{\gamma-\beta-1} t^{\beta+k-1} dt$$

With these substitutions, we therefore, have

$$F(\alpha, \beta, \gamma, x) = \frac{1}{B(\beta, \gamma-\beta)} \sum_{k=0}^{\infty} \frac{(\alpha)_k}{k!} x^k \int_0^1 (1-t)^{\gamma-\beta-1} t^{\beta+k-1} dt$$

$$= \frac{1}{B(\beta, \gamma-\beta)} \int_0^1 (1-t)^{\gamma-\beta-1} t^{\beta-1} \left\{ \sum_{k=0}^{\infty} \frac{(\alpha)_k}{k!} (xt)^k \right\} dt$$

(on interchanging the order of intergration and summation)

$$= \frac{1}{B(\beta, \gamma-\beta)} \int_0^1 (1-t)^{\gamma-\beta-1} t^{\beta-1} (1-xt)^{-\alpha} dt$$

$$\therefore \sum_{k=0}^{\infty} \frac{(\alpha)_k}{k!} (xt)^k = 1 + \frac{\alpha(xt)}{1} + \frac{\alpha(\alpha+1)}{2} (xt)^2 + \dots = (1-xt)^{-\alpha}$$

We thus get the integral formula for hypergeometric series as

$$F(\alpha, \beta, \gamma, x) = \frac{1}{B(\beta, \gamma-\beta)} \int_0^1 (1-t)^{\gamma-\beta-1} t^{\beta-1} (1-xt)^{-\alpha} dt \quad \dots (28)$$

which is valid for $|x| < 1$ and $\gamma > \beta > 0$.

COROLLARY 3. Gauss Theorem or Gauss Formula.

If we put $x=1$ in (25), we have

$$\begin{aligned} F(\alpha, \beta, \gamma, 1) &= \frac{1}{B(\beta, \gamma-\beta)} \int_0^1 (1-t)^{\gamma-\beta-1} t^{\beta-1} (1-t)^{-\alpha} dt \\ &= \frac{1}{B(\beta, \gamma-\beta)} \int_0^1 (1-t)^{\gamma-\beta-\alpha-1} t^{\beta-1} dt = \frac{B(\beta, \gamma-\alpha-\beta)}{B(\beta, \gamma-\beta)} \quad \dots (29) \end{aligned}$$

$$\text{Using } B(P, Q) = \frac{\Gamma P \Gamma Q}{\Gamma(P+Q)}, (27) \text{ yields}$$

$$F(\alpha, \beta, \gamma, 1) = \frac{\Gamma\gamma \Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)} \quad \dots (30)$$

which is known as Gauss's theorem

COROLLARY 4. Vandermond's theorem.

In (28) if we put $\alpha = -n$, we get

$$F(-n, \beta, \gamma, -1) = \frac{\Gamma\gamma \Gamma(\gamma-\beta+n)}{\Gamma(\gamma+n) \Gamma(\gamma-\beta)}$$

$$\begin{aligned}
 &= \frac{1.2 \dots (\gamma-1) \cdot 1.2 \dots (\gamma-\beta+n-1)}{1.2 \dots (\gamma+n-1) \cdot 1.2 \dots (\gamma-\beta-1)} \\
 &= \frac{(\gamma-\beta)(\gamma-\beta+1) \dots (\gamma-\beta+n-1)}{\gamma(\gamma+1) \dots (\gamma+n-1)} = \frac{(\gamma-\beta)_n}{(\gamma)_n} \dots (3)
 \end{aligned}$$

This is known as Vandermond's theorem

COROLLARY 5. Kummer's theorem.

In (27) if we put $x = -1$ and $\gamma = \beta - \alpha + 1$, we get

$$\begin{aligned}
 F(\alpha, \beta, \beta - \alpha + 1, -1) &= \frac{1}{B(\beta, \gamma - \beta)} \int_0^1 (1-t^2)^{-\alpha} t^{\beta-1} dt \\
 &\quad \text{Put } t^2 = y \text{ so that } 2t \, dt = dy \\
 &= \frac{\Gamma(\beta - \alpha + 1)}{\Gamma\beta \Gamma(\gamma - \beta)} \int_0^1 (1-y)^{-\alpha} y^{\beta-1/2} \frac{dy}{2\sqrt{y}} \\
 &= \frac{1}{2} \frac{\Gamma(\beta - \alpha + 1)}{\Gamma\beta \Gamma(\gamma - \beta)} \int_0^1 (1-y)^{-\alpha} y^{\beta/2-1} dy \\
 &= \frac{1}{2} \cdot \frac{\Gamma(\beta - \alpha + 1)}{\Gamma\beta \Gamma(\gamma - \beta)} \cdot B\left(\frac{\beta}{2}, 1 - \alpha\right) \\
 &= \frac{1}{2} \cdot \frac{\Gamma(\beta - \alpha + 1)}{\Gamma\beta \Gamma(1 - \alpha)} \cdot \frac{\Gamma\frac{\beta}{2} \Gamma(1 - \alpha)}{\Gamma\left(\frac{\beta}{2} + 1 - \alpha\right)} \\
 &\quad \because \gamma = \beta - \alpha + 1 \text{ gives, } \gamma - \beta = 1 - \alpha \\
 &= \frac{\Gamma(\beta - \alpha + 1)}{\Gamma(\beta + 1)} \cdot \frac{\Gamma\left(\frac{\beta}{2} + 1\right)}{\Gamma\left(1 - \alpha + \frac{\beta}{2}\right)} \dots (32)
 \end{aligned}$$

(on dividing and multiplying by β)

Which is known as Kummer's theorem.

[D] Linear Relationships of Hypergeometric Functions.

If we put $1-t=p$ in (29) we get

$$\begin{aligned}
 F(\alpha, \beta, \gamma, x) &= \frac{1}{B(\beta, \gamma - \beta)} \int_0^1 p^{\gamma-\beta-1} (1-p)^{\beta-1} \{1-x(1-p)\}^{-\alpha} dp \\
 &= \frac{(1-x)^{-\alpha}}{B(\beta, \gamma - \beta)} \int_0^1 (1-p)^{\beta-1} p^{\gamma-\beta-1} \left\{1 - \frac{xp}{x-1}\right\}^{-\alpha} dp \\
 \text{or } F(\alpha, \beta, \gamma, x) &= \frac{(1-x)^{-\alpha}}{B(\beta, \gamma - \beta)} B(\gamma - \beta, \beta) F\left(\alpha, \gamma - \beta, \gamma, \frac{x}{x-1}\right) \\
 &\quad \text{by using (28)} \\
 &= (1-x)^{-\alpha} F\left(\alpha, \gamma - \beta, \gamma, \frac{x}{x-1}\right) \dots (33)
 \end{aligned}$$

$\therefore$ by symmetry property

$$B(\beta, \gamma - \beta) = B(\gamma - \beta, \beta).$$

Similarly it may be shown that

$$F(\alpha, \beta, \gamma, x) = (1-x)^{-\beta} F\left(\gamma - \alpha, \beta, \gamma, \frac{x}{x-1}\right) \dots (34)$$

With $x = \frac{1}{2}$, (31) yields

$$F\left(\alpha, \beta, \gamma, \frac{1}{2}\right) = 2^\alpha F(\alpha, \gamma - \beta, \gamma, -1) \dots (35)$$

If $\beta = 1 - \alpha$, this gives

$$F\left(\alpha, 1 - \alpha, \gamma, \frac{1}{2}\right) = 2^\alpha F\left(\alpha, \gamma + \alpha - 1, \gamma, -1\right).$$

$$\begin{aligned}
 &= \frac{2^\alpha \Gamma\gamma \Gamma\left(\frac{\gamma + \alpha - 1}{2} + 1\right)}{\Gamma(\gamma + \alpha) \Gamma\left(1 - \alpha + \frac{\gamma + \alpha - 1}{2}\right)} \text{ using (30) i.e. Kummer's theorem} \\
 &= \frac{2^\alpha \Gamma\gamma \Gamma\left(\frac{\gamma}{2} + \frac{1}{2} + \frac{\alpha}{2}\right)}{\Gamma(\gamma + \alpha) \Gamma\left(\frac{1}{2} - \frac{\alpha}{2} + \frac{\gamma}{2}\right)} \\
 &= \frac{2^\alpha \Gamma\left(\frac{\gamma}{2}\right) \cdot \frac{\gamma}{2} \left(\frac{\gamma}{2} + 1\right) \left(\frac{\gamma}{2} + 2\right) \dots (\gamma - 1) \cdot \Gamma\left(\frac{\gamma}{2} + \frac{1}{2}\right) \left(\frac{\gamma}{2} + \frac{1}{2}\right) \dots \left(\frac{\gamma}{2} - \frac{1}{2} + \frac{\alpha}{2}\right)}{\Gamma\left(\frac{\gamma}{2} + \frac{\alpha}{2}\right) \left(\frac{\gamma}{2} + \frac{\alpha}{2}\right) \dots (\gamma + \alpha - 1) \Gamma\left(\frac{1}{2} - \frac{\alpha}{2} + \frac{\gamma}{2}\right)} \\
 &= \frac{2^\alpha \Gamma\left(\frac{\gamma}{2}\right) \cdot \frac{1}{2^{\gamma/2}} \gamma (\gamma + 2) \dots (2\gamma - 2) \cdot \frac{1}{2^{\alpha/2}} \Gamma\left(\frac{\gamma}{2} + \frac{1}{2}\right)}{(\gamma + 1) (\gamma + 3) \dots (\gamma - 1 + \alpha)} \\
 &= \frac{1}{2^{\gamma/2 - \alpha/2}} \Gamma\left(\frac{\gamma}{2} + \frac{\alpha}{2}\right) \cdot (\gamma + \alpha) (\gamma + \alpha + 2) \dots (2\gamma + 2\alpha - 2) \\
 &\quad \Gamma\left(\frac{1}{2} - \frac{\alpha}{2} + \frac{\gamma}{2}\right)
 \end{aligned}$$

$$= \frac{\Gamma\left(\frac{\gamma}{2}\right) \Gamma\left(\frac{\gamma + 1}{2}\right)}{\Gamma\left(\frac{\gamma + \alpha}{2}\right) \Gamma\left(\frac{1 - \alpha + \gamma}{2}\right)} \dots (36)$$

(on simplifying)

Further, we have shown the solution of the hypergeometric series for $x=0$ as

$$y = AF(\alpha, \beta, \gamma, x) + Bx^{1-\gamma} F(\alpha + 1 - \gamma, \beta + 1 - \gamma, 2 - \gamma, x) \text{ by (3)}$$

Which is convergent for $|x| \leq 1$ i.e., in the interval $(-1, 1)$, and the solution for $x=1$ by (6) is

$$y = AF(\alpha, \beta, 1+\alpha+\beta-\gamma, 1-x) + B(1-x)^{\gamma-\alpha-\beta} F(\gamma-\alpha, \gamma-\alpha-\beta+1, 1-x)$$

which is convergent for $|1-x| \leq 1$ i.e., in the interval $(0, 2)$. Clearly the solution (3) and (6) are convergent in the common interval $(0, 1)$ in which there exist a linear relationship between different hypergeometric functions.

Let the relation be $F(\alpha, \beta, \gamma, x) = AF(\alpha, \beta, 1+\alpha+\beta-\gamma, 1-x) + B(1-x)^{\gamma-\alpha-\beta} F(\gamma-\alpha, \gamma-\beta, \gamma-\alpha-\beta+1, 1-x)$... (37)

If we put $x=0$ in (37) we get by Cor. 1 of § 8.5 (C),

$$F(\alpha, \beta, \gamma, 0) = 1 = AF(\alpha, \beta, 1+\alpha+\beta-\gamma, 1) + BF(\gamma-\alpha, \gamma-\alpha-\beta+1, 1) \\ = A \frac{\Gamma(1+\alpha+\beta-\gamma) \Gamma(1-\gamma)}{\Gamma(2+\beta-\gamma) \Gamma(1+\alpha-\gamma)} + B \frac{\Gamma(\gamma-\alpha-\beta+1) \Gamma(1-\gamma)}{\Gamma(1-\beta) \Gamma(1-\alpha)} \dots (38)$$

by Gauss theorem (29)

Putting again, $x=1$ in (37) we get with the help of Gauss theorem,

$$F(\alpha, \beta, \gamma, 1) = AF(\alpha, \beta, 1+\alpha+\beta-\gamma, 0) = \frac{\Gamma\gamma \Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha) \Gamma(\gamma-\beta)}$$

i.e. $A = \frac{\Gamma\gamma \Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha) \Gamma(\gamma-\beta)}$ since by Cor. 1 of § 8.5(C) it may be shown that $F(\alpha, \beta, 1+\alpha+\beta-\gamma, 0) = 1$.

Substituting this value of A in (38), we find

$$1 = \frac{\Gamma\gamma \Gamma(\gamma-\alpha-\beta) \Gamma(1+\alpha+\beta-\gamma) \Gamma(1-\alpha)}{\Gamma(\gamma-\alpha) \Gamma(\gamma-\beta) \Gamma(2+\beta-\gamma) \Gamma(1+\alpha-\gamma)} \\ + B \frac{\Gamma(\gamma-\alpha-\beta+1) \Gamma(1-\gamma)}{\Gamma(1-\beta) \Gamma(1-\alpha)} \\ = \frac{\sin \pi(\gamma-\alpha) \sin \pi(\gamma-\beta)}{\sin \pi\gamma \sin \pi(\gamma-\alpha-\beta)} + B \frac{\Gamma(1-\gamma) \Gamma(1+\gamma-\alpha-\beta)}{\Gamma(1-\alpha) \Gamma(1-\beta)}$$

by using $\Gamma p \Gamma(1-p) = \frac{\pi}{\sin p\pi}$

$$\text{or } B = \frac{\Gamma(1-\alpha) \Gamma(1-\beta)}{\Gamma(1-\gamma) \Gamma(1+\gamma-\alpha-\beta)} \\ \left[\frac{\sin \pi\gamma \sin \pi(\gamma-\alpha-\beta) - \sin \pi(\gamma-\alpha) \sin \pi(\gamma-\beta)}{\sin \pi\gamma \sin \pi(\gamma-\alpha-\beta)} \right] \\ = \frac{\Gamma(1-\alpha) \Gamma(1-\beta) \Gamma(\alpha+\beta-\gamma)}{\Gamma(1-\gamma) \pi} \cdot \frac{-\sin \pi\alpha \sin \pi\beta}{\sin \pi\gamma \sin \pi(\gamma-\alpha-\beta)} \\ = \frac{\Gamma(1-\alpha) \Gamma(1-\beta) \Gamma(\alpha+\beta-\gamma)}{\Gamma(1-\gamma) \pi} \cdot \frac{\pi}{\Gamma\alpha \Gamma(1-\alpha)} \cdot \frac{\pi}{\Gamma\beta \Gamma(1-\beta)} \\ = \frac{\pi}{\Gamma\gamma \Gamma(1-\gamma)}$$

by using $\Gamma p \Gamma(1-p) = \frac{\pi}{\sin p\pi}$

$$= \frac{\Gamma\gamma \Gamma(\alpha+\beta-\gamma)}{\Gamma\alpha \Gamma\beta}$$

Now (37) yields with the substitutions for A and B

$$F(\alpha, \beta, \gamma, x) = \frac{\Gamma\gamma \Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha) \Gamma(\gamma-\beta)} F(\alpha, \beta, 1+\alpha+\beta-\gamma, 1-x) \\ + \frac{\Gamma\gamma \Gamma(\alpha+\beta-\gamma)}{\Gamma\alpha \Gamma\beta} x^{\gamma-\alpha-\beta} F(\gamma-\alpha, \gamma-\beta, \gamma-\alpha-\beta+1, 1-x) \dots (39)$$

which is the required relationship.

If we now replace x by $\frac{1}{x}$ in (39), we get

$$F\left(\alpha, \beta, \gamma, \frac{1}{x}\right) = \frac{\Gamma\gamma \Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha) \Gamma(\gamma-\beta)} F\left(\alpha, \beta, 1+\alpha+\beta-\gamma, 1-\frac{1}{x}\right) \\ + \frac{\Gamma\gamma \Gamma(\alpha+\beta-\gamma)}{\Gamma\alpha \Gamma\beta} \left(1-\frac{1}{x}\right)^{\gamma-\alpha-\beta} F\left(\gamma-\alpha, \gamma-\beta, \gamma-\alpha-\beta+1, 1-\frac{1}{x}\right)$$

but from (33) we have

$$F\left(\alpha, \beta, \gamma, 1-\frac{1}{x}\right) = \left(\frac{1}{x}\right)^{-\alpha} F(\alpha, \gamma-\beta, \gamma, 1-x) \\ = x^{\alpha} F(\alpha, \gamma-\beta, \gamma, 1-x),$$

$$F\left(\alpha, \beta, \gamma, \frac{1}{x}\right) = \frac{\Gamma\gamma \Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha) \Gamma(\gamma-\beta)} x^{\alpha} F(\alpha, \alpha-\gamma+1, \\ \alpha+\beta-\gamma+1, 1-x) + \frac{\Gamma\gamma \Gamma(\alpha+\beta-\gamma)}{\Gamma\alpha \Gamma\beta} x^{\beta} (x-1)^{\gamma-\alpha-\beta} \\ F(\gamma-\alpha, 1-\alpha, \gamma-\alpha-\beta+1, 1-x) \dots (40)$$

where $1 < x < 2$ and $1 > \gamma > \alpha + \beta$.

Various Representations in Terms of Hypergeometric Functions.

(i) If we put $x = (r_2 - r_1)u + r_1$ i.e. $u = \frac{x-r_1}{r_2-r_1}$... (41)

$$\text{i.e. } \frac{du}{dx} = \frac{1}{r_2-r_1}$$

so that $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \frac{1}{r_2-r_1} \frac{dy}{du}$ and $\frac{d^2y}{dx^2} = \frac{d}{dx} \left(\frac{1}{r_2-r_1} \frac{dy}{du} \right)$

$$= \frac{1}{(r_2-r_1)^2} \frac{d^2y}{du^2}, r_1 \neq r_2 \text{ in the equation } (x-r_1)(x-r_2) \frac{d^2y}{dx^2} + \frac{b}{a} \\ (x-r_3) \frac{dy}{dx} + \frac{c}{a} y = 0, \dots (42)$$

then we find the Gauss equation or hypergeometric equation,

$$u(1-u) \frac{d^2y}{du^2} + \left[\frac{b}{a} \frac{r_1-r_3}{r_1-r_2} - \frac{b}{a} u \right] \frac{dy}{du} - \frac{C}{a} y = 0 \dots (43)$$

its series solution by usual method or comparison with § 8.3 $E(e_3)$,

$$y = AF(\alpha, \beta, \gamma, u) + B u^{1-\gamma} F(\alpha-\gamma+1, \beta-\gamma+1, 2-\gamma, u) \dots (44)$$

where $\gamma = \frac{b(r_1 - r_3)}{a(r_1 - r_2)}$, $\alpha + \beta + 1 = \frac{b}{a}$ and $\alpha\beta = \frac{c}{a}$... (41)

$$\therefore y = AF\left(\alpha, \beta, \gamma; \frac{x-r_1}{r_2-r_1}\right) + B\left(\frac{x-r_1}{r_2-r_1}\right)^{1-\gamma} F(\alpha-\gamma+1, \beta-\gamma+1, 2-\gamma, \frac{x-r_1}{r_2-r_1}) \quad \dots (46)$$

COROLLARY. Tschebycheff's equation is

$$(x-1)(x+1) \frac{d^2y}{dx^2} + x \frac{dy}{dx} - n^2y = 0 \quad \dots (47)$$

Comparing it with (42), we get

$$r_1=1, r_2=-1, r_3=0, a=1, b=1, c=-n^2, x=-2u+1, \alpha=n, \beta=-n, \gamma=\frac{1}{2} \text{ and hence by (46), the solution is}$$

$$y = AF\left(n, -n; \frac{1}{2}, \frac{1-x}{2}\right) + B\left(\frac{1-x}{2}\right)^{1/2} F\left(n+\frac{1}{2}, -n+\frac{1}{2}, \frac{3}{2}; \frac{1-x}{2}\right) \quad \dots (48)$$

with the help of (48) and § 8.3 $E(e_3)$, the Tschebycheff's Polynomials may be found as

$$\left. \begin{aligned} T_0(x) &= 1, T_1(x) = x, T_2(x) = 2x^2 - 1, \\ T_3(x) &= 4x^3 - 3x, T_4(x) = 8x^4 - 8x^2 + 1 \end{aligned} \right\} \quad \dots (49)$$

[e_2] Legendre's Polynomials have already been discussed in problem on Legendre's Polynomials.

[e_3] Elliptical integrals of the first and second kind are

$$k(x) = \int_0^{\pi/2} \frac{d\phi}{(1-x^2 \sin^2 \phi)^{1/2}} \quad \dots (50)$$

$$\text{and } E(x) = \int_0^{\pi/2} (1-x^2 \sin^2 \phi)^{1/2} d\phi \quad \dots (51)$$

respectively.

$$\begin{aligned} \text{We have } k(x) &= \int_0^{\pi/2} (1-x^2 \sin^2 \phi)^{-1/2} d\phi \\ &= \int_0^{\pi/2} [1 + \frac{1}{2}x^2 \sin^2 \phi + \frac{\frac{1}{2}(\frac{1}{2}+1)}{2} x^4 \sin^4 \phi + \dots] d\phi \\ &= \int_0^{\pi/2} \sum_{k=0}^{\infty} \frac{\frac{1}{2}(\frac{1}{2}+1)\dots(\frac{1}{2}+k-1)}{|k|} x^{2k} \sin^{2k} \phi d\phi \\ &= \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k x^{2k}}{|k|} \int_0^{\pi/2} \sin^{2k} \phi d\phi \\ &= \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k x^{2k}}{|k|} \cdot \frac{\Gamma(2k+\frac{1}{2})}{\Gamma(k+1)} \cdot \frac{\sqrt{\pi}}{2} \text{ by Gamma integrals} \\ &= \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k x^{2k}}{|k|} \cdot \frac{(\frac{1}{2})_k}{|k|} \cdot \frac{\pi}{2} \end{aligned}$$

$$\begin{aligned} &= \frac{\pi}{2} \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k (\frac{1}{2})_k}{(1)_k |k|} (x^2)^k \text{ since } |k| = 1 \cdot 2 \dots (k-1) = (1)_k \\ &= \frac{\pi}{2} F\left(\frac{1}{2}, \frac{1}{2}, 1; x^2\right) \quad \dots (52) \end{aligned}$$

We can similarly show that

$$E(x) = \frac{\pi}{2} \left(-\frac{1}{2}, \frac{1}{2}, 1; x^2\right) \quad \dots (53)$$

Problem 33. Solve in series

$$x(I-x)y'' + 4(I-x)y' - 2y = 0.$$

Comparing it with the hypergeometric series we get

$$\alpha + \beta + 1 = 4, \gamma = 4, \alpha\beta = 2$$

$$\text{i.e. } \alpha = 1, \beta = 2, \gamma = 4 \text{ or } \alpha = 2, \beta = 1, \gamma = 4$$

For the first choice the solution is $y_1 = F(1, 2, 4, x)$
 $= F(2, 1, 4, x)$ by symmetry property

$$= 1 + \frac{x}{2} + \frac{3x^2}{10} + \frac{x^3}{5} + \frac{x^4}{7} + \dots$$

For the second choice the solution is $y_2 = x^{-3} F(-2, -1, -2, x)$
 $= x^{-3}(1-x)$ since the third term vanishes as one of $\alpha-\gamma+2$ or $\beta-\gamma+2$ is zero. Moreover fourth term has zero for its denominator as $\gamma=4$

Hence the complete solution is

$$y = AF(1, 2, 4, x) + B \frac{1-x}{x^3}.$$

Problem 34. Solve in Series

$$(x-x^2) \frac{d^2y}{dx^2} + \left(\frac{3}{2} - 2x\right) \frac{dy}{dx} - \frac{1}{4}y = 0.$$

Comparing with Gauss equation, we have

$$\alpha + \beta + 1 = 2, \alpha\beta = \frac{1}{4}, \gamma = \frac{3}{2} \text{ giving } \alpha = \beta = \frac{1}{2}, \gamma = \frac{3}{2}.$$

So the solution is

$$\begin{aligned} y &= AF(\alpha, \beta, \gamma, x) + Bx^{1-\gamma} F(\alpha-\gamma+1, \beta-\gamma+1, 2-\gamma, x) \\ &= AF\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, x\right) + Bx^{-\frac{1}{2}} F\left(0, 0, \frac{1}{2}, x\right) \\ &= AF\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, x\right) + Bx^{-\frac{1}{2}} \text{ since } F\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, x\right) \\ &= 1 + \frac{x}{6} + \frac{3x^2}{40} + \frac{5x^3}{112} + \dots \text{ and } F(0, 0, \frac{1}{2}, x) = 1. \end{aligned}$$

Problem 35. Transform $y'' + n^2y = 0$ to hypergeometric form by the substitution $u = \sin^2 x$ and prove that

$$(i) \cos nx = F\left(\frac{n}{2}, -\frac{n}{2}, \frac{1}{2}, \sin^2 x\right)$$

$$(ii) \sin nx = n \sin x F\left(\frac{1}{2} - \frac{1}{2}n, \frac{1}{2} + \frac{1}{2}n, \frac{3}{2}, \sin^2 x\right)$$

$$\text{for } -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}.$$

$$\text{Given equation is } y'' + n^2 y = 0 \quad \dots (1)$$

$$\text{Also given that } u = \sin^2 x \quad \dots (2)$$

$$\text{which gives } \frac{du}{dx} = 2 \sin x \cos x = \sin 2x$$

$$\text{So that } \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \sin 2x \frac{dy}{du} \text{ and } \frac{d^2 y}{dx^2} = 2 \cos 2x \frac{dy}{du} + \sin^2 2x \frac{d^2 y}{du^2}$$

Substituting these values in (1),

$$4 \sin^2 x \cos^2 x \frac{d^2 y}{du^2} + 2(1 - 2 \sin^2 x) \frac{dy}{du} + n^2 y = 0$$

$$\text{or } u(1-u) \frac{d^2 y}{du^2} + \left(\frac{1}{2} - u\right) \frac{dy}{du} + \frac{n^2}{4} y = 0 \quad \dots (3)$$

which is hypergeometric form, with $\gamma = \frac{1}{2}$, $\alpha + \beta + 1 = 1$, $\alpha\beta = -\frac{n^2}{4}$.

i.e. $\alpha = \frac{n}{2}$, $\beta = -\frac{n}{2}$, $r = \frac{1}{2}$ and hence the solution is

$$y = AF\left(\frac{n}{2}, -\frac{n}{2}, \frac{1}{2}, u\right) + Bu^{1/2} F\left(\frac{1+n}{2}, \frac{1-n}{2}, \frac{3}{2}, u\right)$$

$$= AF\left(\frac{n}{2}, -\frac{n}{2}, \frac{1}{2}, \sin^2 x\right) + B \sin x F\left(\frac{1+n}{2}, \frac{1-n}{2}, \frac{3}{2}, \sin^2 x\right) \quad \dots (4)$$

$$\text{where } |\sin x| < 1 \text{ i.e. } -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$$

But $\sin nx$ and $\cos nx$ are the solutions of (1) and so of (3). We can therefore take from (4).

$$\sin nx = AF\left(\frac{n}{2}, -\frac{n}{2}, \frac{1}{2}, \sin^2 x\right) + B \sin x F\left(\frac{1+n}{2}, \frac{1-n}{2}, \frac{3}{2}, \sin^2 x\right) \quad \dots (5)$$

$$\text{When } x=0, A=0 \text{ and so } \frac{\sin nx}{\sin x} = BF\left(\frac{1+n}{2}, \frac{1-n}{2}, \frac{3}{2}, \sin^2 x\right) \quad (\text{by } 5)$$

Which follows that when $x=0$, $B=n$ and hence

$$\sin nx = n \sin x F\left(\frac{1-n}{2}, \frac{1+n}{2}, \frac{3}{2}, \sin^2 x\right) \quad \dots (6)$$

$$\text{Also, } \cos nx = AF\left(\frac{n}{2}, -\frac{n}{2}, \frac{1}{2}, u\right) + B\sqrt{u} F\left(\frac{1+n}{2}, \frac{1-n}{2}, \frac{3}{2}, u\right) \text{ where } u = \sin^2 x \quad \dots (7)$$

when $x=0$, $A=1$

and differentiating (7) w.r.t. x , we find

$$-n \sin nx = A(-n^2) F\left(\frac{n+2}{2}, \frac{2-n}{2}, \frac{3}{2}, u\right) \frac{du}{dx} + B\sqrt{u} \frac{1-n^2}{6} F\left(\frac{3+n}{2}, \frac{3-n}{2}, \frac{5}{2}, u\right) \frac{du}{dx} + BF\left(\frac{1+n}{2}, \frac{1-n}{2}, \frac{3}{2}, u\right) \frac{1}{2\sqrt{u}} \frac{du}{dx} \quad \text{by §8.5 [C] (2).}$$

$$\text{Where } u = \sin^2 x \text{ and } \frac{du}{dx} = \sin 2x$$

When $x=0$, $B=0$

$$\text{So that } \cos nx = F\left(\frac{n}{2}, -\frac{n}{2}, \frac{1}{2}, \sin^2 x\right) \quad \dots (8)$$

8.6. CONFLUENT HYPERGEOMETRIC EQUATIONS AND FUNCTIONS

[A] Confluent Hypergeometric Differential Equation.

$$\text{We have } x(1-x) \frac{d^2 y}{dx^2} + \{\gamma - (\alpha + \beta + 1)x\} \frac{dy}{dx} - \alpha\beta y = 0. \quad \dots (1)$$

Replacing x by $\frac{x}{\beta}$ we have

$$x\left(1 - \frac{x}{\beta}\right) \frac{d^2 y}{dx^2} + \left\{\gamma - \left(1 + \frac{\alpha+1}{\beta}\right)x\right\} \frac{dy}{dx} - \alpha y = 0 \quad \dots (2)$$

Its solution as compared with the hypergeometric series, is

$$F\left(\alpha, \beta, \gamma, \frac{x}{\beta}\right)$$

Hence if we let $\beta \rightarrow \infty$; then (2) reduces to

$$x \frac{d^2 y}{dx^2} + (\gamma - x) \frac{dy}{dx} - \alpha y = 0 \quad \dots (3)$$

$$\text{Whose solution is } \lim_{\beta \rightarrow \infty} F\left(\alpha, \beta, \gamma, \frac{x}{\beta}\right) \quad \dots (4)$$

We call the equation (3) as the confluent hyper-geometric differential equation.

Now since $(\beta)_k = \beta(\beta+1)(\beta+2)\dots(\beta+k-1)$, therefore

$$\lim_{\beta \rightarrow \infty} \frac{(\beta)_k}{\beta^k} = \lim_{\beta \rightarrow \infty} \left(1 + \frac{1}{\beta}\right) \left(1 + \frac{2}{\beta}\right) \dots \left(1 + \frac{k-1}{\beta}\right) = 1$$

As such (4) may be represented as

$$\lim_{\beta \rightarrow \infty} F\left(\alpha, \beta, \gamma, \frac{x}{\beta}\right) = \sum_{r=0}^{\infty} \frac{(\alpha)_r}{(\gamma)_r} \cdot \frac{x^r}{r!} = F(\alpha, \gamma, x) \text{ (say)} \dots (5)$$

Here $F(\alpha, \gamma, x)$ or sometimes, ${}_1F_1(\alpha, \gamma, x)$ showing that there is one parameter of the type α and one of the type γ , is the solution of (3) and is said to be the *confluent hypergeometric function*.

Note. Since equation (3) has a removable (non-essential) singularity at $x=0$ so its solution may be developed directly by series method about $x=0$, choosing the series as $y = \sum_{r=0}^{\infty} a_r x^{k+r}$.

Substituting for $y, \frac{dy}{dx}, \frac{d^2y}{dx^2}$ in (3) and equating the coefficient of first and the general term (i.e. x^{k+r}) to zero, it is easy to find that indicial equation $k(k+\gamma-1)=0$ gives $k=0, 1-\gamma$ when $a_0 \neq 0$... (6)

and the recurrence relation is $a_{r+1} = \frac{k+r+\alpha}{(k+r+1)(k+r+\gamma)} a_r$... (7)

which for $k=0$ gives

$$y = a_0 \left[1 + \frac{\alpha}{\gamma} x + \frac{\alpha(\alpha+1)}{1.2 \cdot \gamma(\gamma+1)} x^2 + \dots \right] \\ = a_0 \sum_{k=0}^{\infty} \frac{(\alpha)_k}{k! (\gamma)_k} x^k = a_0 F(\alpha, \gamma, x) \dots (8)$$

and for $k=1-\gamma$ gives

$$y = a_0 x^{1-\gamma} \left[1 + \frac{\alpha'}{\gamma'} x + \frac{\alpha'(\alpha'+1)}{1.2 \gamma'(\gamma'+1)} x^2 + \dots \right]$$

where

$$\alpha' = \alpha - \gamma + 1 \text{ and } \gamma' = 2 - \gamma \\ = a_0 x^{1-\gamma} \sum_{k=0}^{\infty} \frac{(\alpha')_k}{k! (\gamma')_k} x^k = a_0 x^{1-\gamma} F(\alpha', \gamma', x) \\ = a_0 x^{1-\gamma} F(\alpha - \gamma + 1, 2 - \gamma, x) \dots (9)$$

where $F(\alpha - \gamma + 1, 2 - \gamma, x)$ is said to be the confluent hypergeometric function of the second kind.

The general solution is therefore,

$$y = AF(\alpha, \gamma, x) + Bx^{1-\gamma} F(\alpha - \gamma + 1, 2 - \gamma, x) \dots (10)$$

It is valid for $\gamma > 0$.

By ratio test

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{(\alpha)_{n+1} \frac{n}{n+1}}{(\alpha)_n \frac{n}{n+1}} \cdot \frac{(\gamma)_n}{(\gamma)_{n+1}} x \right| = \left| \frac{\alpha + n}{(\gamma + n)(n+1)} x \right|$$

which $\rightarrow 0$ as $n \rightarrow \infty$ i.e. $\left| \frac{u_{n+1}}{u_n} \right| < 1$ for all x , it is obvious that the series is convergent.

8) Simple Properties of Confluent Hypergeometric Functions.

[1] Differentiation of Confluent Hypergeometric Functions i.e.

$$\frac{d}{dx} F(\alpha, \gamma, x) = \frac{\alpha}{\gamma} F(\alpha+1, \gamma+1, x) \dots (11)$$

which may be similarly shown as in case of hypergeometric functions.

$$\frac{d^2}{dx^2} F(\alpha, \gamma, x) = \frac{\alpha}{\gamma} \left(\frac{\alpha+1}{\gamma+1} \right) F(\alpha+2, \gamma+2, x) \dots (12)$$

and in general

$$\frac{d^m}{dx^m} F(\alpha, \gamma, x) = \frac{(\alpha)_m}{(\gamma)_m} F(\alpha+m, \gamma+m, x) \dots (13)$$

$$\text{When } x=0, F(\alpha, \gamma, x)=1 \dots (14)$$

$$\left[\frac{d}{dx} F(\alpha, \beta, x) \right]_{x=0} = \frac{\alpha}{\gamma} \dots (15)$$

$$\text{and } \left[\frac{d^m}{dx^m} F(\alpha, \gamma, x) \right]_{x=0} = \frac{(\alpha)_m}{(\gamma)_m} \dots (16)$$

In case α is a negative integer, the series stops after a certain number of terms but when γ is also negative then the series restarts after a certain number of terms, as shown in case of hypergeometric functions.

[2] Integral Formulae for the Confluent Hypergeometric Function.

$$\text{We have } \frac{(\alpha)_k}{(\gamma)_k} = \frac{1}{B(\alpha, \gamma-\alpha)} \int_0^1 (1-t)^{\gamma-\alpha-1} t^{\alpha+k-1} dt \dots (17)$$

Just as in §8.5 [C-3]

$$\therefore F(\alpha, \gamma, x) = \sum_{k=0}^{\infty} \frac{(\alpha)_k}{k! (\gamma)_k} x^k \\ = \sum_{k=0}^{\infty} \frac{x^k}{k!} \cdot \frac{1}{B(\alpha, \gamma-\alpha)} \int_0^1 (1-t)^{\gamma-\alpha-1} t^{\alpha+k-1} dt \\ = \frac{1}{B(\alpha, \gamma-\alpha)} \int_0^1 (1-t)^{\gamma-\alpha-1} t^{\alpha-1} \sum_{k=0}^{\infty} \frac{(xt)^k}{k!} dt \\ = \frac{1}{B(\alpha, \gamma-\alpha)} \int_0^1 e^{xt} t^{\alpha-1} (1-t)^{\gamma-\alpha-1} dt \\ \because \sum_{k=0}^{\infty} \frac{(xt)^k}{k!} = 1 + \frac{xt}{1} + \frac{x^2 t^2}{2} + \dots = e^{xt} \\ = \frac{\Gamma \gamma}{\Gamma \alpha \Gamma(\gamma-\alpha)} \int_0^1 e^{xt} t^{\alpha-1} (1-t)^{\gamma-\alpha-1} dt \dots (18)$$

$$\text{using } B(l, m) = \frac{\Gamma l \Gamma m}{\Gamma(l+m)}$$

which is the integral formulae for the confluent hypergeometric function.

COROLLARY 1. In (18) if we put $t=1-p$, then

$$\begin{aligned} F(\alpha, \gamma, x) &= \frac{\Gamma\gamma}{\Gamma\alpha \Gamma(\gamma-\alpha)} \int_0^1 e^{x(1-p)} (1-p)^{\alpha-1} p^{\gamma-\alpha-1} dp \\ &= \frac{\Gamma\gamma}{\Gamma\alpha \Gamma(\gamma-\alpha)} e^x \int_0^1 e^{-xp} p^{\gamma-\alpha-1} (1-p)^{\alpha-1} dp \\ &= \frac{\Gamma\gamma}{\Gamma\alpha \Gamma(\gamma-\alpha)} e^x \frac{\Gamma(\gamma-\alpha) \Gamma\alpha}{\Gamma\gamma} F(\gamma-\alpha, \gamma, -x) \text{ using (18)} \\ &= e^x F(\gamma-\alpha, \gamma, -x) \end{aligned} \quad \dots (19)$$

COROLLARY 2. In (18) if we put $x=0$, then we find

$$\begin{aligned} F(\alpha, \gamma, 0) &= \frac{\Gamma\gamma}{\Gamma\alpha \Gamma(\gamma-\alpha)} \int_0^1 t^{\alpha-1} (1-t)^{\gamma-\alpha-1} dt \\ &= \frac{\Gamma\gamma}{\Gamma\alpha \Gamma(\gamma-\alpha)} B(\alpha, \gamma-\alpha) \\ &= \frac{\gamma}{\Gamma\alpha \Gamma(\gamma-\alpha)} \cdot \frac{\Gamma\alpha \Gamma(\gamma-\alpha)}{\Gamma\gamma} = 1 \\ \text{i.e. } F(\alpha, \gamma, 0) &= 1 \end{aligned} \quad \dots (20)$$

[C] Various Representations in terms of Confluent Hypergeometric Functions.

[c₁] Elementary Functions.

$$\text{We have } F(\alpha, \gamma, x) = \sum_{k=0}^{\infty} \frac{(\alpha)_k}{k! (\gamma)_k} x^k$$

$$\begin{aligned} \text{Putting } \gamma=\alpha, \text{ we have } F(\alpha, \alpha, x) &= \sum_{k=0}^{\infty} \frac{x^k}{k!} = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \dots \\ &= e^x \end{aligned} \quad \dots (21)$$

Similarly,

$$F(\alpha+1, \alpha, x) = \left(1 + \frac{x}{\alpha}\right) e^x \quad \dots (22)$$

$$\begin{aligned} F(1, 2, x) &= \sum_{k=0}^{\infty} \frac{(1)_k x^k}{(2)_k k!} = \sum_{k=0}^{\infty} \frac{x^k}{(k+1)!} \\ &= \frac{1}{x} \left[x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \right] \\ &= \frac{e^x - 1}{x} \end{aligned} \quad \dots (23)$$

$$F(-2, 1, x) = 1 - 2x + \frac{x^2}{2}$$

$$F(n, n+1, -x) = n \int_0^x t^{n-1} e^{-t} dt \quad \dots (24)$$

[c₂] Error Function.

$$\text{We have } \operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-n^2} dn \text{ (see Ch. VI)}$$

$$\begin{aligned} &= \frac{2}{\sqrt{\pi}} \int_0^x \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \frac{x^{2k}}{k!} dn \\ &= \frac{2}{\sqrt{\pi}} \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \int_0^x n^{2k} dn = \sum_{k=0}^{\infty} \frac{(-1)^k}{\sqrt{k!}} \frac{x^{2k+1}}{(2k+1)!} \\ &= x \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \frac{x^{2k}}{(2k+1)!} \end{aligned}$$

$$\begin{aligned} &= \frac{2x}{\sqrt{\pi}} \sum_{k=0}^{\infty} \frac{(-x^2)^k}{k! (2k+1)!} \\ &= \frac{2x}{\sqrt{\pi}} \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k (-x^2)^k}{k! \left(\frac{3}{2}\right)_k} \\ &\quad \because 2k+1 = \frac{(2k+1)(2k-1)(2k-3)\dots 1}{(2k-1)(2k-3)\dots 1} \\ &\quad \quad \frac{1}{2} \cdot \frac{3}{2} \dots \frac{2k-1}{2} \cdot \frac{2k+1}{2} \\ &= \frac{\left(\frac{1}{2} \cdot \frac{3}{2} \dots \frac{2k-1}{2} \cdot \frac{2k-3}{2} \dots \frac{1}{2}\right) \cdot \frac{1}{2}}{\frac{3}{2} \left(\frac{3}{2} + 1\right) \dots \frac{2k+1}{2} \left(\frac{3}{2}\right)_k} \\ &= \frac{1}{2} \left(\frac{1}{2} + 1\right) \dots \frac{2k-1}{2} = \left(\frac{1}{2}\right)_k \\ &= \frac{2x}{\sqrt{\pi}} F\left(\frac{1}{2}, \frac{3}{2}, -x^2\right) \end{aligned} \quad \dots (25)$$

[c₃] Fresnel's integrals. The first and second kinds are respectively,

$$C(x) = \frac{1}{2\sqrt{2}} \left[e^{\pi i/4} \phi\left(\frac{\sqrt{\pi}}{\sqrt{2}} x e^{-\pi i/4}\right) + e^{-\pi i/4} \phi\left(\frac{\sqrt{\pi}}{\sqrt{2}} x e^{\pi i/4}\right) \right] \quad \dots (26)$$

$$\text{and } S(x) = \frac{1}{2i\sqrt{2}} \left[e^{\pi i/4} \phi\left(\frac{\sqrt{\pi}}{\sqrt{2}} x e^{-\pi i/4}\right) - e^{-\pi i/4} \phi\left(\frac{\sqrt{\pi}}{\sqrt{2}} x e^{\pi i/4}\right) \right] \quad \dots (27)$$

$$\text{where } \phi(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-n^2} dx$$

$$\therefore \text{ by [c}_2\text{]}, \phi\left\{\sqrt{\frac{\pi}{2}} x e^{-\pi i/4}\right\} = \frac{2}{\sqrt{\pi}} \int_0^{\sqrt{\pi/2} x e^{-\pi i/4}} e^{-n^2} dn$$

$$= \sqrt{2} x e^{-\pi i/4} \sum_{k=4}^{\infty} \frac{\left(\frac{1}{2}\right)_k \left(\frac{1}{2} \pi i x^2\right)_k}{\left(\frac{3}{2}\right)_k |k|}$$

$$= \sqrt{2} x e^{-\pi i/4} F\left(\frac{1}{2}, \frac{3}{2}, -\frac{\pi i x^2}{2}\right)$$

Similarly $\phi \left\{ \sqrt{\frac{\pi}{2}} x e^{\pi i/4} \right\} = \sqrt{2} x e^{\pi i/4} F\left(\frac{1}{2}, \frac{3}{2}, \frac{\pi i x^2}{2}\right)$

Hence $C(x) = \frac{x}{2} \left[F\left(\frac{1}{2}, \frac{3}{2}, \frac{\pi i x^2}{2}\right) + F\left(\frac{1}{2}, \frac{3}{2}, -\frac{\pi i x^2}{2}\right) \right]$... (28)

and $S(x) = \frac{x}{2i} \left[F\left(\frac{1}{2}, \frac{3}{2}, \frac{\pi i x^2}{2}\right) - F\left(\frac{1}{2}, \frac{3}{2}, -\frac{\pi i x^2}{2}\right) \right]$... (29)

[c] Bessel (Cylindrical) Functions.

We have $J_n(x) = \frac{\left[\frac{x}{2}\right]^n}{\Gamma\left(\frac{1}{2}\right) \Gamma\left[n + \frac{1}{2}\right]} \int_{-1}^1 (1-t^2)^{n-\frac{1}{2}} e^{ixt} dt$

Put $2p = 1+t$, $J_n(x) = \frac{2^{2n} \left[\frac{x}{2}\right]^n e^{-in}}{\Gamma\left(\frac{1}{2}\right) \Gamma\left[n + \frac{1}{2}\right]} \int_0^1 e^{2xip} p^{n-\frac{1}{2}} (1-p)^{n-\frac{1}{2}} dp$

$$= \frac{\left[\frac{x}{2}\right]^n}{\Gamma(n+1)} e^{-ix} F\left[n + \frac{1}{2}, 2n+1, 2ix\right]$$
 ... (30)

Since by putting $\alpha = n + \frac{1}{2}$, $\gamma = 2n+1$ and x for $2ix$ in

$$F(\alpha, \gamma, x) = \frac{\Gamma\gamma}{\Gamma\alpha \Gamma(\gamma-\alpha)} \int_0^1 e^{xt} t^{\alpha-1} (1-t)^{\gamma-\alpha-1} dt,$$

we get

$$F\left[n + \frac{1}{2}, 2n+1, 2ix\right] = \frac{\Gamma(2n+1)}{\Gamma\left[n + \frac{1}{2}\right] \Gamma\left[n + \frac{1}{2}\right]} \int_0^1 e^{2ixt} t^{n-\frac{1}{2}} (1-t)^{n-\frac{1}{2}} dt$$

Problem 36. Solve in series the Whittakar Equation $\frac{d^2 W}{dx^2} + \left\{ -\frac{1}{4} - \frac{k}{x} + \frac{\frac{1}{4} - m^2}{x^2} \right\} W(x) = 0$ and represent its functions in terms of confluent hypergeometric functions.

Taking $W(x) = x^{\gamma/2} e^{-x/2} y(x)$, $k = \frac{\gamma}{2} - \alpha$, $m = -\frac{1}{2} + \frac{\gamma}{2}$, the equation reduces to

$$x \frac{d^2 y}{dx^2} + (\gamma - x) \frac{dy}{dx} - \alpha y = 0$$

Which is confluent hypergeometric equation and hence its functions by putting $\alpha = \frac{1}{2} - k + m$ and $\gamma = 1 + 2m$ in $F(\alpha, \gamma, x)$ and $x^{\gamma-1} F(1+\alpha-\gamma, 2-\gamma, x)$ are $F(\frac{1}{2}-k+m, 1+2m, x)$ and $x^{-2m} F(\frac{1}{2}-k-m, 1-2m, x)$ respectively.

Thus the solution of given equation is

$$y = AF\left[\frac{1}{2} - k + m, 1 + 2m, x\right] + Bx^{-2m} F\left[\frac{1}{2} - k - m, 1 - 2m, x\right]$$

where A and B are arbitrary constants.

Also Whittakar's functions are

$$W_{k, m}(x) = x^{\frac{1}{2}+m} e^{-x/2} F\left(\frac{1}{2} - k + m, 1 + 2m, x\right)$$

and $W_{k, -m}(x) = x^{\frac{1}{2}-m} e^{-x/2} F\left[\frac{1}{2} - k - m, 1 - 2m, x\right]$.

8.7. HERMITE EQUATION, FUNCTIONS AND POLYNOMIALS

[A] Hermite's Differential Equation.

This equation is of the form

$$\frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + 2\nu y = 0 \quad \dots (1)$$

where ν is a parameter.

Suppose its series solution is

$$y = \sum_{r=0}^{\infty} a_r x^{k+r}, \quad a_0 \neq 0 \quad \dots (2)$$

So that $\frac{dy}{dx} = \sum_{r=0}^{\infty} a_r (k+r) x^{k+r-1}$

and $\frac{d^2 y}{dx^2} = \sum_{r=0}^{\infty} a_r (k+r) (k+r-1) x^{k+r-2}$

Substituting the values of y , $\frac{dy}{dx}$ and $\frac{d^2 y}{dx^2}$ in (1), we get the identity

$$\sum_{r=0}^{\infty} [(k+r)(k+r-1) x^{k+r-2} - 2(k+r-\nu) x^{k+r-1}] a_r \equiv 0 \quad \dots (3)$$

Equating the coefficient of the first term (i.e. x^{k-2}) (by putting $r=0$), to zero, we get

$$a_0 k(k-1) = 0 \text{ giving } k=0, 1 \text{ as } a_0 \neq 0 \quad \dots (4)$$

Now-equating to zero the coefficient of second term (i.e. x^{k-1}) in (3) we get

$a_1 k(k+1) = 0$ i.e. $a_1 = 0$ when $k = -1$ and a_1 may or may not be zero when $k = 0$.

Also equating the coefficient of x^{k+r} to zero, we find

$$a_{r+2}(k+r+2)(k+r+1) - 2a_r(k+r-\nu) = 0$$

giving the recurrence relation

$$a_{r+2} = \frac{2(k+r-\nu)}{(k+r+2)(k+r+1)} a_r \quad \dots (5)$$

$$\text{when } k=0, (5) \text{ becomes } a_{r+2} = \frac{2(r-\nu)}{(r+2)(r+1)} a_r \quad \dots (6)$$

$$\text{and when } k=1, (5) \text{ becomes } a_{r+2} = \frac{2(1+r-\nu)}{(r+3)(r+2)} a_r \quad \dots (7)$$

Case I. When $k=0$, putting $r=0, 1, 2, 3, \dots$ in (6) we have

$$a_2 = -\frac{2}{2} \nu a_0; \quad a_4 = -\frac{2(\nu-1)}{3} a_2$$

$$a_4 = \frac{2^2 \nu(\nu-2)}{4} a_0; \quad a_6 = \frac{2^3(\nu-1)(\nu-3)}{5} a_4$$

$$\dots \dots \dots$$

$$a_{2r} = \frac{(-2)^r \nu(\nu-2) \dots (\nu-2r+2)}{2r} a_0;$$

$$a_{2r+1} = \frac{(-2)^r (\nu-1)(\nu-3) \dots (\nu-2r+1)}{2r+1} a_1$$

Now if $a_1 = 0$, then $a_3 = a_5 = a_7 = a_{2r+1} = \dots = 0$.

But if $a_1 \neq 0$, then (2) gives for $k=0$, $y = \sum_{r=0}^{\infty} a_r x^r$

i.e. $y = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots$

$$= a_0 + a_2 x^2 + a_4 x^4 + \dots + a_1 x + a_3 x^3 + a_5 x^5 + \dots$$

$$= a_0 \left[1 - \frac{2\nu}{2} x^2 + \frac{2^2 \nu(\nu-2)}{4} x^4 - \dots \right]$$

$$+ (-1)^r \frac{2^r}{2r} \nu(\nu-2) \dots (\nu-2r+2) x^{2r} + \dots]$$

$$+ a_1 x \left[1 - \frac{2(\nu-1)}{3} x^2 + \frac{2^3(\nu-1)(\nu-3)}{5} x^4 - \dots \right]$$

$$+ (-1)^r \frac{2^r}{2r+1} (\nu-1)(\nu-3) \dots (\nu-2r+1) x^{2r+1} + \dots] \quad \dots (8)$$

$$= a_0 \left[1 + \sum_{r=1}^{\infty} \frac{(-1)^r 2^r}{2r} \nu(\nu-2) \dots (\nu-2r+2) x^{2r} \right]$$

$$+ a_1 \left[x + \sum_{r=1}^{\infty} \frac{(-1)^r 2^r}{2r+1} (\nu-1)(\nu-3) \dots (\nu-2r+1) x^{2r+1} \right] \quad \dots (9)$$

Case II. When $k=1$, then $a_1 = 0$ and so by putting $r=0, 1, 2, 3$, in (7) we find

$$a_2 = -\frac{2(\nu-1)}{3} a_0$$

$$a_4 = \frac{2^2(\nu-1)(\nu-3)}{5} a_0$$

$$\dots \dots \dots$$

$$a_{2r} = (-1)^r \frac{2^r (2\nu-1)(\nu-3) \dots (\nu-2r+1)}{2r+1} a_0$$

Hence the solution is

$$y = a_0 x \left[1 - \frac{2(\nu-1)}{3} x^2 + \frac{2^2(\nu-1)(\nu-3)}{5} x^4 - \dots \right. \\ \left. + \frac{(-1)^r 2^r (\nu-1)(\nu-3) \dots (\nu-2r+1)}{2r+1} x^{2r} + \dots \right] \quad \dots (1)$$

Clearly the solution (10) is included in the second part of (8) except that a_0 is replaced by a_1 and hence in order that the Hermite equation may have two independent solutions, a_1 must be zero, even if $k=0$ and then (8) reduces to

$$y = a_0 \left[1 - \frac{2\nu}{2} x^2 + \frac{2^2 \nu(\nu-2)}{4} x^4 - \dots \right. \\ \left. + (-1)^r \frac{2^r}{2r} \nu(\nu-2) \dots (\nu-2r+2) x^{2r} + \dots \right] \quad \dots (11)$$

The complete integral of (1) is then given by

$$y = A \left[1 - \frac{2\nu}{2} x^2 + \frac{2^2 \nu(\nu-2)}{4} x^4 - \dots \right] + B \left[1 - \frac{2(\nu-1)}{3} x^2 \right. \\ \left. + \frac{2^2(\nu-1)(\nu-3)}{5} x^4 - \dots \right] \quad \dots (1)$$

where A and B are arbitrary constants.

(ii) Hermite Polynomials.

The Hermite polynomial $H_n(x)$ is defined as

$$f(x, t) = e^{2tx - t^2} = \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!} \quad \dots (13)$$

for all integral values of n and all real values of x .

(13) can be written as

$$f(x, t) = e^{x^2} e^{-(x-t)^2} = \frac{H_0(x)}{0!} + \frac{H_1(x)}{1!} t + \frac{H_2(x)}{2!} t^2 \\ + \dots + \frac{H_n(x)}{n!} t^n + \dots$$

So that $\left[\frac{\partial^n f(x,t)}{\partial t^n} \right]_{t=0} = \frac{H_n(x)}{|n|} |n| = H_n(x)$

$$= \left[\frac{\partial^n}{\partial t^n} e^{-(x-t)^2} \right]_{t=0} e^{x^2} \quad \dots (14)$$

If we put $x-t=p$ i.e. $t=x-p=0$ for $t=0$ gives $x=p$
 and $\frac{\partial}{\partial t} = -\frac{\partial}{\partial p}$ so that $\frac{\partial^n}{\partial t^n} \left\{ e^{-(x-t)^2} \right\} = (-1)^n \frac{\partial^n}{\partial p^n} e^{-p^2}$
 $\therefore \left[\frac{\partial^n}{\partial t^n} e^{-(x-t)^2} \right]_{t=0} = (-1)^n \frac{\partial^n}{\partial x^n} e^{-x^2} = (-1)^n \frac{d^n}{dx^n} e^{-x^2}$

From (14) and (15), we therefore have

$$H_n(x) = e^{x^2} (-1)^n \frac{d^n}{dx^n} (e^{-x^2}) \quad \dots (16)$$

From (16) we can calculate Hermite polynomials of various degrees such as

$$\left. \begin{aligned} H_0(x) &= 1 & H_4(x) &= 16x^4 - 48x^2 + 12 \\ H_1(x) &= 2x & H_5(x) &= 32x^5 - 160x^3 + 120x \\ H_2(x) &= 4x^2 - 2 & H_6(x) &= 64x^6 - 480x^4 + 720x^2 - 120 \\ H_3(x) &= 8x^3 - 12x & H_7(x) &= 128x^7 - 1344x^5 + 3360x^3 - 1680x \end{aligned} \right\} \quad \dots (17)$$

[C] Hermite Polynomial in Terms of Confluent Hypergeometric Functions.

We have $\sum_{n=0}^{\infty} H_n(x) \frac{t^n}{|n|} = e^{2tx-t^2} = e^{2tx} \cdot e^{-t^2}$

$$= \left[1 + 2tx + \frac{(2x)^2}{|2|} t^2 + \dots + \frac{(2x)^n}{|n|} t^n + \dots \right] \left[1 - \frac{t^2}{|1|} + \frac{(t^2)^2}{|2|} - \dots + \frac{(-1)^n t^{2n}}{|n|} + \dots \right]$$

Equating the coefficients of $\frac{t^n}{|n|}$ on either side we get

$$H_n(x) = (2x)^n - \frac{n(n-1)}{|1|} (2x)^{n-2} + \frac{n(n-1)(n-2)(n-3)}{|2|} (2x)^{n-4} - \dots$$

$$= (2x)^n F\left(-\frac{n}{2}, \frac{1}{2}, -\frac{n}{2}, -\frac{1}{x^2}\right) \quad \dots (18)$$

Aliter, we have $\sum_{n=0}^{\infty} H_n(x) \frac{t^n}{|n|} = e^{2tx-t^2} = e^{2tx} e^{-t^2}$

$$= \sum_{n=0}^{\infty} \frac{(2x)^n t^n}{|n|} \times \sum_{n=0}^{\infty} (-1)^n \frac{t^{2n}}{|n|}$$

$$\therefore H_n(x) = \sum_{k=0}^{\infty} \sum_{k=0}^{n/2} (-1)^k \frac{|n|}{|k|} \frac{(2x)^{n-2k}}{|n-2k|} \quad \dots (19)$$

Even Hermite polynomials are, therefore

$$H_{2n}(x) = \sum_{k=0}^n (-1)^k \frac{|2n|}{|k|} \frac{|2n-2k|}{|2n-2k|} \cdot (2x)^{2n-2k}$$

$$= |2n| (-1)^n \sum_{k=0}^n \frac{(-1)^k (2x)^{2k}}{|n-k| |2k|} = (-1)^n \frac{|2n|}{|n|} \sum_{k=0}^n (-1)^k \frac{(2x)^{2k}}{|2k|}$$

$$= (-1)^n \frac{|2n|}{|n|} \sum_{k=0}^{\infty} \frac{(-n)_k (x^2)^k}{(\frac{1}{2})_k |k|} \because |2k| = 2^{2k} \left(\frac{1}{2}\right)_k |k|$$

$$= (-1)^n \frac{|2n|}{|n|} F\left(-n, \frac{1}{2}; x^2\right) \quad \dots (20)$$

Similarly odd Hermite polynomials are

$$H_{2n+1}(x) = \sum_{k=0}^{(2n+1)/2} \frac{(-1)^k |2n+1| (2x)^{2n+1-2k}}{|k| |2n+1-2k|} = (-1)^n \frac{|2n+1|}{|n|} 2x \sum_{k=0}^n \frac{(-n)_k x^{2k}}{(\frac{3}{2})_k |k|}$$

$$= (-1)^n \frac{|2n+1|}{|n|} 2x F\left(-n, \frac{3}{2}; x^2\right) \quad \dots (21)$$

[D] Recurrence formulae for $H_n(x)$ and to show that $H_n(x)$ is a solution of Hermite Equation.

Hermite equation is $y'' - 2xy' + 2ny = 0$ for integral values taking $y = H_n$.

Also, $e^{2tx-t^2} = \sum_{n=0}^{\infty} \frac{H_n(x) t^n}{|n|} \quad \dots (22)$

I. Differentiating partially w.r.t. x , we have

$$2t e^{-2tx-t^2} = \sum_{n=0}^{\infty} \frac{H'_n(x) t^n}{|n|}$$

i.e. $2t \sum_{n=0}^{\infty} \frac{H_n(x) t^n}{|n|} = \sum_{n=0}^{\infty} \frac{H'_n(x) t^n}{|n|}$

which yields on equating the coefficients of $\frac{t^n}{|n|}$ on either side,

$$2 \frac{|n|}{|n-1|} H_{n-1}(x) = H'_n(x)$$

i.e. $2n H_{n-1}(x) = H'_n(x) \quad \dots (23)$

II. Differentiating partially w.r.t. t , both sides of (22) we get

$$2(x-t) e^{2tx-t^2} = \sum_{n=1}^{\infty} \frac{H_n(x) t^{n-1}}{|n-1|} \because n=0 \text{ corresponds to the vanishing of R.H.S.}$$

$$\text{or } 2x \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!} - 2 \sum_{n=0}^{\infty} H_n(x) \frac{t^{n+1}}{n!} = \sum_{n=1}^{\infty} H_n(x) \frac{t^{n-1}}{(n-1)!}$$

Equating the coefficients of t^n on either side we find

$$2x \frac{H_n(x)}{n!} - 2 \frac{H_{n-1}(x)}{(n-1)!} = \frac{H_{n+1}(x)}{n!}$$

$$\text{i.e. } 2x H_n(x) = 2n H_{n-1}(x) + H_{n+1}(x) \quad \dots (24)$$

III. Eliminating $H_{n-1}(x)$ from (23) and (24) we get

$$2x H_n(x) = H_n'(x) + H_{n+1}(x)$$

$$\text{or } H_n'(x) = 2x H_n(x) - H_{n+1}(x) \quad \dots (25)$$

IV. Differentiating (25) w.r.t. x we find

$$H_n''(x) = 2x H_n'(x) + 2H_n(x) - H_{n+1}'(x)$$

Putting $H_{n+1}'(x) = 2(n+1) H_n(x)$ obtained from (23) on replacing n by $n+1$; we have

$$H_n''(x) = 2x H_n'(x) + 2H_n(x) - 2(n+1) H_n(x)$$

$$\text{or } H_n''(x) - 2x H_n'(x) + 2n H_n(x) = 0 \quad \dots (26)$$

which clearly follows that $y = H_n(x)$ is a solution of Hermite equation.

Note. e^{2tx-t^2} is known as **Generating function of Hermite Polynomial**.

Problem 37. Prove that

$$(i) H_{2m}(0) = (-1)^m 2^{2m} \left(\frac{1}{2} \right)_m$$

$$(ii) H'_{2m+1}(0) = (-1)^m 2^{2m+1} \left(\frac{3}{2} \right)_m$$

$$(iii) H_{2m+1}(0) = 0$$

$$(iv) H'_{2m}(0) = 0$$

$$(v) \frac{d^m}{dx^m} \left\{ H_n(x) \right\} = \frac{2^m n!}{(n-m)!} H_{n-m}(x), \text{ for } m < n$$

(i) Even Hermite polynomials are

$$H_{2m}(x) = \sum_{k=0}^m \frac{(-1)^k}{k!} \frac{2m!}{(2m-2k)!} (2x)^{2m-2k}$$

$$\therefore H_{2m}(0) = \frac{(-1)^m}{m!} \frac{2m!}{m!} = (-1)^m \frac{2m(2m-1)\dots 3.2.1}{m(m-1)\dots 3.2.1}$$

$$= (-1)^m 2^{2m} \frac{m!}{m!} \frac{1}{2} \left(\frac{1}{2} + 1 \right) \dots \left(\frac{1}{2} + m - 1 \right) = (-1)^m 2^{2m} \left(\frac{1}{2} \right)_m$$

(ii) From recurrence relation I, we have on replacing n by $2m+1$,

$$H'_{2m+1}(x) = 2(2m+1) H_{2m}(x)$$

$$\therefore H'_{2m+1}(0) = 2(2m+1) H_{2m}(0) = 2(2m+1) \cdot (-1)^m 2^{2m} \left(\frac{1}{2} \right)_m$$

by part (i)

$$\begin{aligned} &= (2m+1) (-1)^m 2^{2m+1} \left[\frac{(2m-1)(2m-3)\dots 3.1}{2^m} \right] \\ &= (-1)^m (2m+1) \left[\frac{3}{2} \cdot \left(\frac{3}{2} + 1 \right) \dots \left(\frac{3}{2} + m - 1 \right) \right] \\ &= (-1)^m 2^{2m+1} \left(\frac{3}{2} \right)_m. \end{aligned}$$

(iii) Odd Hermite Polynomials are

$$H_{2m+1}(x) = \sum_{k=0}^{2m+1/2} \frac{(-1)^k}{k!} \frac{2m+1}{(2m+1-2k)!} (2x)^{2m+1-2k}$$

$\therefore H_{2m+1}(0) = 0$, since all terms containing x become zero.

(iv) From recurrence relation I, we have

$$H'_{2m}(x) = 2(2m) H_{2m-1}(x)$$

$$\therefore H'_{2m}(0) = 4m H_{2m-1}(0) = 0 \text{ by (iii)}$$

(v) From recurrence relation I, we have

$$H'_n(x) = 2n H_{n-1}(x) \quad \dots (i)$$

$$\text{i.e. } \frac{d}{dx} \left\{ H_n(x) \right\} = 2n H_{n-1}(x)$$

$$\therefore \frac{d^2}{dx^2} \left\{ H_n(x) \right\} = 2n \frac{d}{dx} \left\{ H_{n-1}(x) \right\} = 2n \cdot 2(n-1) H_{n-2}(x)$$

$$\text{by using (1)} \quad = 2^2 n(n-1) H_{n-2}(x).$$

$$\text{Similarly } \frac{d^3}{dx^3} \left\{ H_n(x) \right\} = 2^3 n(n-1)(n-2) H_{n-3}(x)$$

Proceeding similarly m times we find

$$\begin{aligned} \frac{d^m}{dx^m} \left\{ H_n(x) \right\} &= 2^m n(n-1)\dots(n-m+1) H_{n-m}(x) \text{ where } m < n \\ &= \frac{2^m n!}{(n-m)!} H_{n-m}(x) \end{aligned}$$

[E] Hermite Functions.

An equation closely related to Hermite equation is

$$\frac{d^2 \psi}{dx^2} + (\lambda - x^2) \psi = 0 \quad \dots (27)$$

If we change the dependent variable ψ to y by the substitution

$$\psi = e^{-x^2/2} y \quad \dots (28)$$

$$\text{So that } \frac{d\psi}{dx} = e^{-x^2/2} \frac{dy}{dx} - y e^{-x^2/2} \cdot x$$

$$\text{and } \frac{d^2\psi}{dx^2} = e^{-x^2/2} \frac{d^2y}{dx^2} - 2x e^{-x^2/2} \frac{dy}{dx} - (e^{-x^2/2} - x^2 e^{-x^2/2}) y$$

We get from (1)

$$y'' - 2xy' + (\lambda - 1)y = 0 \quad \dots (29)$$

If we put $\lambda - 1 = 2\nu$, then (29) reduces to Hermite equation *i.e.*

$$y'' - 2xy' + 2\nu y = 0$$

It therefore follows that the general solution of (27) is given by

$$\psi = e^{-x^2/2} y$$

where y is given by (12) of § 8.7.

Thus if the parameter λ be of the form $1 + 2n$, n being a positive integer, then the solution of (27) will be a constant multiple of the function ψ_n defined by

$$\psi_n(x) = e^{-x^2/2} H_n(x) \quad \dots (30)$$

where $H_n(x)$ is the Hermite polynomial of degree n .

Here the function $\psi_n(x)$ is said to be the *Hermite function of order n* .

Recurrence Relations for $\psi_n(x)$:

Differentiating (30) w.r.t. x , we have

$$\begin{aligned} \psi'_n(x) &= e^{-x^2/2} H'_n(x) - e^{-x^2/2} x H_n(x) \\ &= 2n e^{-x^2/2} H_{n-1}(x) - x e^{-x^2/2} H_n(x) \\ &\quad \therefore H'_n(x) = 2n H_{n-1}(x) \text{ by (21)} \\ &= 2n \psi_{n-1}(x) - x \psi_n(x) \text{ using (30)} \end{aligned}$$

$$\therefore 2n \psi_{n-1}(x) = x \psi_n(x) + \psi'_n(x) \quad \dots (31)$$

Also from (25), $2x H_n(x) = 2n H_{n-1}(x) + H_{n+1}(x)$

which may be expressed by using (30), as

$$2x e^{-x^2/2} H_n(x) = 2n e^{-x^2/2} H_{n-1}(x) + e^{-x^2/2} H_{n+1}(x) \quad \dots (32)$$

i.e.

Eliminating $2n \psi_{n-1}(x)$ from (31) and (32) we find

$$x \psi_n(x) + \psi'_n(x) = 2x \psi_n(x) - \psi_{n+1}(x) \quad \dots (33)$$

i.e.

[F] Orthogonal Properties of Hermite Polynomials.

Now since $H_n(x)$ is a solution of Hermite equation, we have

$$H''_n(x) - 2x H'_n(x) + 2n H'_n(x) = 0 \text{ by (26)}$$

If we put $y = e^{-x^2/2} H_n(x)$ *i.e.* $H_n(x) = y e^{x^2/2}$

So that $H'_n(x) = y' e^{x^2/2} + xy e^{x^2/2}$

$$\text{and } H''_n(x) = y'' e^{x^2/2} + 2xy' e^{x^2/2} + y(1+x^2) e^{x^2/2} \quad \dots (34)$$

then we get

$$y'' + (1 - x^2 + 2n)y = 0 \quad \dots (34)$$

Since $y = e^{-x^2/2} H_n(x) = \psi_n(x)$ by (30), it therefore follows that $\psi_n(x)$ satisfies (34) and hence

$$\psi''_n + (2n + 1 - x^2) \psi_n = 0 \quad \dots (35)$$

a function ψ_m , this relation is

$$\psi''_m + (2m + 1 - x^2) \psi_m = 0 \quad \dots (36)$$

Multiplying (35) by ψ_m ; (36) by ψ_n and subtracting we get

$$2(m - n) \psi_m \psi_n = \psi_m \psi''_n - \psi_n \psi''_m \quad \dots (37)$$

Integrating over $(-\infty, \infty)$, we have

$$\begin{aligned} 2(m - n) \int_{-\infty}^{\infty} \psi_m \psi_n dx &= \int_{-\infty}^{\infty} (\psi_m \psi''_n - \psi_n \psi''_m) dx \\ &= \left[\psi_m \psi'_n - \psi_n \psi'_m \right]_{-\infty}^{\infty} - \int_{-\infty}^{\infty} (\psi'_m \psi'_n - \psi'_n \psi'_m) dx \\ &\quad \text{(on integrating by parts)} \\ &= 0 \quad \therefore \psi_n(x) \rightarrow 0 \text{ as } |x| \rightarrow \infty \text{ for all positive integral values of } n. \end{aligned}$$

$$\text{or } \int_{-\infty}^{\infty} \psi_m \psi_n dx = 0 \text{ if } m \neq n$$

$$\text{Symbolically } I_{m, n} = \int_{-\infty}^{\infty} \psi_m \psi_n dx = \int_{-\infty}^{\infty} e^{-x^2} H_m(x) H_n(x) dx = 0 \quad \text{when } m \neq n \quad \dots (38)$$

$$\text{In particular } I_{n-1, n+1} = 0 \quad \dots (39)$$

Now from (32) we have $2x \psi_n(x) = 2n \psi_{n-1}(x) + \psi_{n+1}(x)$

$$\therefore \int_{-\infty}^{\infty} 2x \psi_n(x) \psi_{n-1} dx = 2n \int_{-\infty}^{\infty} \psi_{n-1}(x) \psi_{n-1}(x) dx$$

$$\therefore \int_{-\infty}^{\infty} \psi_{n-1} \psi_{n+1} dx = 0 \text{ by (39)} \quad \dots (40)$$

$$\text{Also } \psi_n(x) = e^{-x^2/2} H_n(x) = (-1)^n e^{x^2/2} \frac{d^n}{dx^n} (e^{-x^2}) \text{ by (16)}$$

Thus (40) gives

$$-\int_{-\infty}^{\infty} 2x e^{x^2/2} \frac{d^n}{dx^n} (e^{-x^2/2}) \frac{d^{n-1}}{dx^{n-1}} (e^{-x^2}) dx = 2n I_{n-1, n-1}$$

$$\begin{aligned} \text{or } 2n I_{n-1, n-1} &= - \int_{-\infty}^{\infty} d(e^{x^2}) \frac{d^n}{dx^n} (e^{-x^2}) \frac{d^{n-1}}{dx^{n-1}} (e^{-x^2}) dx \\ &= \left[e^{-x^2} \frac{d^n}{dx^n} (e^{-x^2}) \cdot \frac{d^{n-1}}{dx^{n-1}} (e^{-x^2}) \right]_{-\infty}^{\infty} + \int_{-\infty}^{\infty} e^{-x^2} \left\{ \frac{d^n}{dx^n} (e^{-x^2}) \frac{d^n}{dx^n} (e^{-x^2}) + \frac{d^{n+1}}{dx^{n+1}} (e^{-x^2}) \cdot \frac{d^{n-1}}{dx^{n-1}} (e^{-x^2}) \right\} dx \\ &\quad \text{(on integrating by parts)} \\ &= 0 + I_{n, n} + I_{n+1, n-1} \\ &= I_{n, n} \text{ by (39)} \end{aligned}$$

$$\therefore I_{n, n} = 2n I_{n-1, n-1} \quad \dots (41)$$

$$\begin{aligned}
&= n \int_{-\infty}^{\infty} e^{-x^2} H_{n-1}(x) H_n(x) dx + m \int_{-\infty}^{\infty} e^{-x^2} H_n(x) H_{m-1}(x) dx \\
&= n \sqrt{\pi} 2^{n-1} \left| \underline{n-1} \right| \delta_{m, n-1} + m \sqrt{\pi} 2^n \left| \underline{n} \right| \delta_{n, m-1} \\
&\quad \text{(by orthogonal properties)} \\
&= \sqrt{\pi} [2^{n-1} \left| \underline{n} \right| \delta_{m, n-1} + 2^n \left| \underline{n+1} \right| \delta_{n+1, m}] \\
&\quad \therefore \delta_{n, m-1} = \delta_{n+1, m}
\end{aligned}$$

Problem 40. Verify, $P_n(x) = \frac{2}{\sqrt{\pi} \left| \underline{n} \right|} \int_0^{\infty} t^n e^{-t^2} H_n(xt) dt$.

Problem 41. Show that if m is an integer, $\int_{-\infty}^{\infty} x^m e^{-x^2} H_n(x) dx = 0$.

8.8. LAGUERRE EQUATION AND POLYNOMIAL WITH PROPERTIES

[A] Laguerre's Differential Equation.

This equation is of the form

$$xy'' + (1-x)y' + \nu y = 0 \quad \dots (1)$$

Dividing by x , it is observed that $x=0$ is a regular singularity of (1) and hence it has a series solution. Let its series solution be

$$y = \sum_{r=0}^{\infty} a_r x^{k+r} \quad a_0 \neq 0 \quad \dots (2)$$

$$\therefore y' = \sum_{r=0}^{\infty} a_r (k+r) x^{k+r-1}$$

$$\text{and } y'' = \sum_{r=0}^{\infty} a_r (k+r)(k+r-1) x^{k+r-2}$$

Substituting these values in (1), we get the identity

$$\sum_{r=0}^{\infty} \left[(k+r)^2 x^{k+r-1} - (k+r-\nu) x^{k+r} \right] a_r \equiv 0 \quad \dots (3)$$

Equating to zero the coefficient of x^{k-1} (the first and the lowest term); we get

$$k^2 = 0 \text{ i.e. } k=0 \text{ as } a_0 \neq 0 \quad \dots (4)$$

Again equating to zero the coefficient of x^{k+r} in (3) we get

$$(k+r+1)^2 a_{r+1} - (k+r-\nu) a_r = 0$$

which gives the recurrence relation

$$a_{r+1} = \frac{k+r-\nu}{(k+r+1)^2} a_r \quad \dots (5)$$

For $k=0$, this yields, $a_{r+1} = \frac{r-\nu}{(r+1)^2} a_r$

$$\therefore a_1 = -\nu a_0 = (-1) \nu a_0$$

$$a_2 = \frac{1-\nu}{2^2} = (-1)^2 \frac{\nu(\nu-1)}{(\underline{2})^2} a_0$$

$$\text{Similarly } a_3 = (-1)^3 \frac{\nu(\nu-1)(\nu-2)}{(\underline{3})^2} a_0$$

$$\dots \dots \dots a_r = (-1)^r \frac{\nu(\nu-1)\dots(\nu-r+1)}{(\underline{r})^2} a_0$$

Hence the solution is

$$\begin{aligned}
y = a_0 \left[1 - \nu x + \frac{\nu(\nu-1)}{(\underline{2})^2} x^2 - \dots \right. \\
\left. + (-1)^r \frac{\nu(\nu-1)\dots(\nu-r+1)}{(\underline{r})^2} x^r + \dots \right] \dots (6)
\end{aligned}$$

$$= a_0 \sum_{k=0}^{\infty} \frac{(-1)^k \left| \underline{\nu} \right|}{(\underline{k})^2 \left| \underline{\nu-k} \right|} \quad \dots (7)$$

In case ν is a positive integer put $\nu=n$, so that Laguerre equation becomes

$$xy'' + (1-x)y' + ny = 0 \text{ for positive integral } n. \quad \dots (8)$$

When $\nu=n$ (a positive integer) and $a_0 = \left| \underline{n} \right|$ then solution for (8) is said to be the Laguerre polynomial of degree n and denoted by $L_n(x)$ i.e.

$$L_n(x) = (-1)^n \left[x^n - \frac{n^2}{\underline{1}} x^{n-1} + \frac{n^2(n-1)^2}{\underline{2}} x^{n-2} + \dots + (-1)^n \left| \underline{n} \right| \right] \quad \dots (9)$$

Then the solution of Laguerre equation for ν to be a positive integer is

$$y = AL_n(x) \quad \dots (10)$$

From (9), it is easy to show that

$$\left. \begin{aligned} L_n(0) &= \left| \underline{n} \right|, & L_2(x) &= x^2 - 4x + 2 \\ L_0(x) &= 1, & L_3(x) &= -x^3 + 9x^2 - 36x + 6 \\ L_1(x) &= 1-x, & L_4(x) &= x^4 - 16x^3 + 72x^2 - 96x + 48 \end{aligned} \right\} \quad \dots (11)$$

Also $L_n(x)$ being the solution of (8), we should have

$$xL_n''(x) + (1-x)L_n'(x) + nL_n(x) = 0 \quad \dots (12)$$

[B] Laguerre Polynomials with their Representation in Terms of Confluent Hypergeometric Series.

The Laguerre Polynomials $L_n(x)$ are given by the relation

$$(1-t) \sum_{n=0}^{\infty} \frac{L_n(x)}{\left| \underline{n} \right|} t^n = e^{\frac{-xt}{1-t}} \quad \dots (13)$$

where n is a positive integer and x is a positive real number.

(13) can be written as

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{L_n(x) t^n}{n!} &= \frac{1}{1-t} e^{-\frac{xt}{1-t}} \\
&= \frac{1}{1-t} \left[1 - \frac{xt}{1-t} + \frac{x^2 t^2}{2! (1-t)^2} - \dots + \frac{(-1)^k x^k t^k}{k! (1-t)^k} + \dots \right] \\
&= \sum_{k=0}^{\infty} \frac{(-1)^k x^k t^k}{k! (1-t)^{k+1}} \\
&= \sum_{k=0}^{\infty} \frac{(-1)^k x^k t^k}{k!} (1-t)^{-(k+1)} \\
&= \sum_{k=0}^{\infty} \frac{(-1)^k x^k}{k!} t^k \left[1 + (k+1)t + \frac{(k+1)(k+2)}{2!} t^2 + \dots \right. \\
&\quad \left. \dots + \frac{(k+1)(k+2)\dots(k+l)}{l!} t^l + \dots \right] \\
&= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{(-1)^k (k+1)_l}{k! l!} x^k t^{k+l} \text{ where } (k+1)_l = \frac{\Gamma(k+1+l)}{\Gamma(k+1)}
\end{aligned}$$

Equating the coefficients of t^n on either side (coefficient of t^n on R.H.S. being obtained by putting $l=n-k$), we get

$$\frac{L_n(x)}{n!} = \sum_{k=0}^{\infty} \frac{(-1)^k (k+1)_{n-k}}{k! (n-k)!} x^k \quad \dots (14)$$

$$\text{Here } (k+1)_{n-k} = \frac{\Gamma(k+1+n-k)}{\Gamma(k+1)} = \frac{\Gamma(n+1)}{\Gamma(k+1)} \frac{n!}{k!}$$

$$\begin{aligned}
\therefore \frac{(-1)^k}{(n-k)!} &= \frac{(-1)^k n(n-1)\dots(n-k+1)}{n!} \\
&= \frac{(-n)(-n+1)(-n+2)\dots(-n+k-1)}{n!} = \frac{(-n)_k}{n!}
\end{aligned}$$

$\therefore$ (14) yields

$$\begin{aligned}
L_n(x) &= \frac{n!}{n!} \sum_{k=0}^{\infty} \frac{(-n)_k}{k!} \cdot \frac{n!}{(k!)^2} x^k = \frac{n!}{n!} \sum_{k=0}^{\infty} \frac{(-n)_k}{(k!)^2} x^k \\
&= \frac{n!}{n!} \left[1 + \frac{(-n)}{1! 1!} x + \frac{(-n)(-n+1)}{2! 2!} x^2 \right. \\
&\quad \left. + \frac{(-n)(-n+1)(-n+2)}{3! 3!} x^3 + \dots \right] \\
&= \frac{n!}{n!} F(-n, 1; x) \quad \dots (15)
\end{aligned}$$

From which it follows that $L_n(x)$ is a polynomial of degree n in x and that the coefficient of x^n is $(-1)^n$.

[C] Recurrence formulae for Laguerre Polynomials. The generating function for Laguerre Polynomial is

$$e^{-\frac{xt}{1-t}} = (1-t) \sum_{n=0}^{\infty} \frac{L_n(x)}{n!} t^n \text{ by (13)} \quad \dots (16)$$

I. Differentiating w.r.t. ' t ' it gives

$$-\frac{x}{(1-t)^2} e^{-\frac{xt}{1-t}} = (1-t) \sum_{n=0}^{\infty} \frac{L_n(x) t^{n-1}}{(n-1)!} - \sum_{n=0}^{\infty} \frac{L_n(x) t^n}{n!}$$

Using (16), we have

$$-\frac{x}{(1-t)^2} (1-t) \sum_{n=0}^{\infty} \frac{L_n(x) t^n}{n!} = (1-t) \sum_{n=0}^{\infty} \frac{L_n(x) t^{n-1}}{(n-1)!} - \sum_{n=0}^{\infty} \frac{L_n(x) t^n}{n!}$$

$$\text{or } x \sum_{n=0}^{\infty} \frac{L_n(x) t^n}{n!} + (1-t)^2 \sum_{n=0}^{\infty} \frac{L_n(x) t^{n-1}}{(n-1)!} - (1-t) \sum_{n=0}^{\infty} \frac{L_n(x) t^n}{n!} = 0$$

Equating to zero the coefficient of t^n , we find

$$\begin{aligned}
x \frac{L_n(x)}{n!} + \frac{L_{n+1}(x)}{(n+1)!} - 2 \frac{L_n(x)}{(n-1)!} + \frac{L_{n-1}(x)}{(n-2)!} - \frac{L_n(x)}{n!} + \frac{L_{n-1}(x)}{(n-1)!} &= 0 \\
\text{i.e. } L_{n+1}(x) + (x-2n-1) L_n(x) + n^2 L_{n-1}(x) &= \dots (17)
\end{aligned}$$

II. Again differentiating (16) w.r.t. x we get

$$\begin{aligned}
-\left(\frac{t}{1-t}\right) e^{-\frac{xt}{1-t}} &= (1-t) \sum_{n=0}^{\infty} \frac{L'_n(x) t^n}{n!} \\
\text{or } -\left(\frac{t}{1-t}\right) (1-t) \sum_{n=0}^{\infty} \frac{L_n(x)}{n!} t^n &= (1-t) \sum_{n=0}^{\infty} \frac{L'_n(x)}{n!} t^n \text{ by (16)}
\end{aligned}$$

$$\text{or } t \sum_{n=0}^{\infty} \frac{L_n(x) t^n}{n!} + (1-t) \sum_{n=0}^{\infty} \frac{L'_n(x) t^n}{n!} = 0$$

Equating to zero the coefficient of t^n , we get

$$\begin{aligned}
\frac{L'_n(x)}{n!} - \frac{L'_{n-1}(x)}{(n-1)!} + \frac{L_{n-1}(x)}{(n-1)!} &= 0 \\
\text{i.e. } L'_n(x) - n L'_{n-1}(x) + n L_{n-1}(x) &= 0 \quad \dots (18)
\end{aligned}$$

III. Now differentiating (17) w.r.t. x , we find

$$L'_{n+1}(x) - (x-2n-1) L'_n(x) + L_n(x) + n^2 L'_{n-1}(x) = 0.$$

Differentiating it again w.r.t. x , we have

$$L''_{n+1}(x) + (x-2n-1) L''_n(x) + 2L'_n(x) + n^2 L''_{n-1}(x) = 0$$

Replacing n by $n+1$, this yields

$$L''_{n+2}(x) + (x-2n-3) L''_{n+1}(x) + (n+1)^2 L''_n(x) + 2L'_{n+1}(x) = 0 \quad (a)$$

Whence from (18),

$$L'_n(x) = n \{L'_{n-1}(x) - L_{n-1}(x)\}$$

$$\text{or } L'_{n+1}(x) = (n+1) \{L'_n(x) - L_n(x)\} \text{ (on replacing } n \text{ by } n+1) \quad (b)$$

$$\therefore L''_{n+1}(x) = (n+1) \{L''_n(x) - L'_n(x)\} \text{ (on differentiating)} \quad (c)$$

$$\text{or } L''_{n+2}(x) = (n+2) \{L''_{n+1}(x) - L'_{n+1}(x)\} \text{ on replacing } n \text{ by } n+1$$

Thus we have from (a)

$$(n+2) \{L''_{n+1}(x) - L'_{n+1}(x)\} + (x-2n-3) L'_{n+1}(x) + (n+1)^2 L_n''(x) + 2L'_{n+1}(x) = 0$$

$$\text{or } (x-n-1) L''_{n+1}(x) - nL'_{n+1}(x) + (n+1)^2 L_n''(x) = 0$$

Eliminating $L''_{n+1}(x)$ and $L'_{n+1}(x)$ by (b) and (c), we get

$$x L_n''(x) + (1-x) L'_n(x) + n L_n(x) = 0 \quad \dots (19)$$

which clearly shows that $y = A L_n(x)$ is a solution of Laguerre equation.

[D] Differentiation formula for Laguerre Polynomial (Analog of Rodrigues formula)

We have.

$$(1-t)^{-1} e^{\left(1 - \frac{1}{1-t}\right)x} = \sum_{n=0}^{\infty} \frac{L_n(x) t^n}{|n|} \text{ by (13)}$$

Differentiating it w.r.t. 't' n times by Leibnitz theorem for successive differentiation, we have

$$e^x \frac{d^n}{dt^n} \left[(1-t)^{-1} e^{-\frac{x}{1-t}} \right] = L_n(x) + L_{n+1}(x) t + \dots \quad \dots (20)$$

since all terms upto the term containing t^{n-1} vanish when differentiated n times.

$$\text{Now } \frac{d}{dt} \left[(1-t)^{-1} e^{-\frac{x}{1-t}} \right] = \frac{1-x-t}{(1-t)^2} e^{-\frac{x}{1-t}}$$

So that

$$\lim_{t \rightarrow 0} \frac{d}{dt} \left[(1-t)^{-1} e^{-\frac{x}{1-t}} \right] = (1-x) e^{-x} = \frac{d}{dx} (x e^{-x})$$

Similarly

$$\lim_{t \rightarrow 0} \frac{d^2}{dt^2} \left[(1-t)^{-1} e^{-\frac{x}{1-t}} \right] = \frac{d^2}{dx^2} (x^2 e^{-x})$$

And in general

$$\lim_{t \rightarrow 0} \frac{d^n}{dt^n} \left[(1-t)^{-1} e^{-\frac{x}{1-t}} \right] = \frac{d^n}{dx^n} (x^n e^{-x})$$

Hence proceeding to the limit as $t \rightarrow 0$, (20) yields

$$e^n \frac{d^n}{dx^n} (x^n e^{-x}) = L_n(x) \quad \dots (21)$$

[E] Integral Property of Laguerre Polynomials.

Laguerre equation is $xy'' + (1-x)y' + ny = 0$ for positive integral n.

$$\text{If we take } y_n = \frac{1}{2\pi i} \oint \frac{t^{-n-1}}{1-t} e^{-\frac{xt}{1-t}} dt$$

$$\text{So that } y_n' = \frac{1}{2\pi i} \oint \frac{t^{-n}}{(1-t)^2} e^{-\frac{xt}{1-t}} dt$$

$$\text{and } y_n'' = \frac{1}{2\pi i} \oint \frac{t^{-n+1}}{(1-t)^3} e^{-\frac{xt}{1-t}} dt$$

Then Laguerre Polynomials being the solution of (22), its L.H.S. becomes with the substitutions of (23),

$$\frac{1}{2\pi i} \oint \left[\frac{xt^2}{(1-t)^2} - \frac{(1-x)t}{1-t} + n \right] \frac{t^{-n-1}}{1-t} e^{-\frac{xt}{1-t}} dt$$

$$\text{i.e. } -\frac{1}{2\pi i} \oint \frac{d}{dt} \left[\frac{t^{-n}}{1-t} \right] e^{-\frac{xt}{1-t}} dt$$

which should be zero for the contour being closed and hence L.H.S. of (22) vanishes for

$$y = \frac{1}{2\pi i} \oint \frac{t^{-n-1}}{1-t} e^{-\frac{xt}{1-t}} dt \quad \dots (24)$$

It means, for this value y is a solution of (22) and therefore we may express $L_n(x)$ which is an already established solution of (22), as

$$L_n(x) = A y_n(x) \quad A \text{ being an arbitrary constant} \quad \dots (25)$$

Now for $x=0$, we get from (9) and (23), $L_n(0) = |n| \quad \dots (26)$

$$\begin{aligned} \text{and } y_n(0) &= \frac{1}{2\pi i} \oint \frac{t^{-n-1}}{1-t} dt \\ &= \frac{1}{2\pi i} \times 2\pi i \text{ by contour integration} \\ &= 1 \end{aligned} \quad \dots (27)$$

So (25) gives for $x=0$, with the help of (26) and (27)

$$L_n(0) = A y_n(0) \text{ i.e., } |n| = A.$$

Hence from (25),

$$\begin{aligned} L_n(x) &= |n| y_n(x) \\ &= \frac{|n|}{2\pi i} \oint \frac{t^{-n-1}}{1-t} e^{-\frac{xt}{1-t}} dt \text{ by (24)} \end{aligned} \quad \dots (28)$$

Now by contour integration, $y_n = \text{coefficient of } t^n \text{ in } (1-t)^{-1} e^{-\frac{xt}{1-t}}$

$$\text{Hence } (1-t)^{-1} e^{-\frac{xt}{1-t}} = \sum_{n=0}^{\infty} y_n t^n = \sum_{n=0}^{\infty} L_n(x) \frac{t^n}{|n|} \quad \dots (29)$$

which is the generating function of Laguerre Polynomials.

[F] Orthogonal Properties of Laguerre Polynomials.

The Laguerre polynomials themselves do not form an orthonormal set since the Laguerre equation is not self-adjoint. We, therefore, introduce a function

$$\phi_n(x) = \frac{1}{|n|} e^{-x/2} L_n(x) \quad \dots (30)$$

and then will show that ϕ 's form an orthonormal set i.e.

$$\int_0^{\infty} \phi_m(x) \phi_n(x) dx = \int_0^{\infty} e^{-x} \frac{L_m(x)}{|m|} \frac{L_n(x)}{|n|} dx = \delta_{m,n} \quad \dots (31)$$

Over the interval $0 \leq x < \infty$, when $\delta_{m,n} = 0$ for $m \neq n$
 $= 1$ for $m = n$.

Since $L_n(x) = e^{-x} \frac{d^n}{dx^n} (x^n e^{-x})$ by (21), therefore, we have

$$\int_0^\infty e^{-x} x^m L_n(x) dx = \int_0^\infty e^{-x} x^m e^x \frac{d^n}{dx^n} (x^n e^{-x}) dx \\ = \int_0^\infty x^m \frac{d^n}{dx^n} (x^n e^{-x}) dx$$

Integrating the R.H.S. by parts, we get

$$\int_0^\infty e^{-x} x^m L_n(x) dx = \left[x^m \frac{d^{n-1}}{dx^{n-1}} x^n e^{-x} \right]_0^\infty \\ - \int_0^\infty m x^{m-1} \frac{d^{n-1}}{dx^{n-1}} (x^n e^{-x}) dx \\ = (-1) \cdot m \int_0^\infty x^{m-1} \frac{d^{n-1}}{dx^{n-1}} (x^n e^{-x}) dx \\ = (-1)^2 m(m-1) \int_0^\infty x^{m-2} \frac{d^{n-2}}{dx^{n-2}} (x^n e^{-x}) dx \\ = \dots \\ = (-1)^m \frac{m!}{n!} \int_0^\infty \frac{d^{n-m}}{dx^{n-m}} (x^n e^{-x}) dx \\ = 0 \text{ if } n > m.$$

Similarly, $\int_0^\infty e^{-x} x^n L_m(x) dx = 0$ for $m > n$.

Now $L_m(x)$ being a polynomial of degree m in x and $L_n(x)$ that of degree n in x , we have

$$\int_0^\infty e^{-x} L_m(x) L_n(x) dx = 0 \text{ if } m \neq n. \\ \text{i.e. } \int_0^\infty e^{-x} \frac{L_m(x)}{m!} \frac{L_n(x)}{n!} dx = 0 \text{ if } m \neq n \quad \dots (32)$$

In case $m = n$, then the term of degree n in $L_n(x)$ is $(-1)^n x^n$,

$$\therefore \int_0^\infty e^{-x} \{L_n(x)\}^2 dx = (-1)^n \int_0^\infty e^{-x} x^n L_n(x) dx \\ = (-1)^n \int_0^\infty e^{-x} x^n e^x \frac{d^n}{dx^n} (x^n e^{-x}) dx \\ = (-1)^n \int_0^\infty x^n \frac{d^n}{dx^n} (x^n e^{-x}) dx \\ = (-1)^{2n} \frac{n!}{n!} \int_0^\infty x^n e^{-x} dx \text{ (on integrating by parts } n \text{ times)} \\ = \frac{n!}{n!}$$

$$\text{or } \int_0^\infty e^{-x/2} \frac{L_n(x)}{n!} e^{-x/2} \frac{L_n(x)}{n!} dx = 1 \quad \dots (33)$$

Combining (32) and (33) we have

$$\int_0^\infty \phi_m(x) \phi_n(x) dx = \int_0^\infty e^{-x/2} \frac{L_m(x)}{m!} e^{-x/2} \frac{L_n(x)}{n!} dx = \delta_{m,n}$$

Note. $\phi_n(x)$ satisfies the equation

$$x \phi_n''(x) + \phi_n'(x) + \left(n + \frac{1}{2} - \frac{x}{4}\right) \phi_n(x) = 0$$

[G] The associated Laguerre Polynomials and Functions.

For positive integral n , Laguerre differential equation is

$$xy_2 + (1-x)y_1 + xy = 0 \text{ where } y_1 = \frac{dy}{dx} \text{ and } y_2 = \frac{d^2y}{dx^2}.$$

Differentiating it m times by Leibnitz theorem, we get

$$x y_{m+2} + {}^m C_1 y_{m+1} + (1-x) y_{m+1} + {}^m C_1 y_m (-1) + n y_m = 0 \\ \text{or } x y_{m+2} + (m-1-x) y_{m+1} + (n-m) y_m = 0$$

Writing $y_m = D^m y = D^m L_n(x)$, for $n \geq m$, this can be expressed as

$$x D^2(y_m) + (m+1-x) D(y_m) + (n-m) y_m = 0 \quad \dots (34)$$

From (34) it follows that

$$y_m = D^m L_n(x) = \frac{d^m}{dx^m} L_n(x) \equiv L_n^m(x) \text{ (say)} \quad \dots (35)$$

is the solution of

$$x y'' + (m+1-x) y' + (n-m) y = 0 \quad \dots (36)$$

where m is an integer such that $m \geq 0$.

The polynomial $L_n^m(x)$ introduced here is said to be the *Associated Laguerre Polynomials* of degree $n-m$.

It may be shown that in terms of confluent hypergeometric functions,

$$L_n^m(x) = \frac{(-1)^m (n!)}{m! (n-m)!} F(-n+m, m+1, x), n \geq m \quad \dots (37)$$

$$\text{or } = (m+1)_k F(-n, m+1, -x) \quad \dots (38)$$

From $L_n(x) = e^x D^n (x^n e^{-x})$, $D \equiv \frac{d}{dx}$

we may infer that

$$L_n^m(x) = D^m \{e^x D^n (x^n e^{-x})\} = \frac{e^x x^{n-m} \frac{d^n}{dx^n} (e^{-x} x^{n+m})}{dx^n} \quad \dots (39)$$

From $e^{\frac{-xt}{1-t}} = (1-t) \sum_{n=0}^\infty \frac{L_n(x) t^n}{n!}$, we may conclude that the

generating function for Associated Laguerre polynomials is

$$(-1)^m t^m e^{\frac{-xt}{1-t}} = (1-t)^{m+1} \sum_{n=m}^\infty L_n^m(x) \frac{t^n}{n!} \quad \dots (40)$$

Note 1. The generalized Laguerre Polynomials are defined by

$$L_n^m(x) = \frac{m+n!}{m! n!} F(-n, m+1, x) \quad \dots (41)$$

where n is a positive integer or zero.

This is a solution of $xy'' + (m+1-x)y' + ny = 0$... (42)

Note 2. The associated Laguerre equation (36) is not self-adjoint and hence does not form orthonormal system, but it can be made so by multiplying the weighing function $e^{-x} x^m$. We thus define the integral

$$I_{p,n} = \int_0^\infty y_{n,m} y_{p,m} x^q dx \quad \dots (43)$$

For $q=1$, where $y_{n,m} = e^{-x/2} x^{(q-1)/2} L_n^m(x)$... (44)

known as associated Laguerre function is the solution of

$$xy'' + 2y' + \left[n - \frac{q-1}{2} - \frac{4}{x} \cdot \frac{q^2-1}{4x} \right] y = 0$$

which is obtained by putting

$$y = e^{-x/2} x^{(q-1)/2} v \text{ i.e. } v = y e^{x/2} e^{-(q-1)/2} \text{ in} \quad \dots (45)$$

$$xv'' + (q+1-x)v' + (n-q)v = 0 \text{ (analogous to (36), } v = L_n^m(x))$$

we have

$$y_{n,m} = e^{-x/2} x^{(m-1)/2} L_n^m(x) \text{ and } y_{p,m} = e^{-x/2} x^{(m-1)/2} L_p^m(x)$$

$$\text{So that } I_{p,n} = \int_0^\infty e^{-x} x^{m-1} L_n^m(x) L_p^m(x) \cdot x^q dx \quad \dots (47)$$

Problem 42. Show that $L_n(2x) = \left| n \sum_{m=0}^n \frac{2^{n-m} (-1)^m}{|m| |n-m|} L_{n-1}(x) \right|$, $n > m$

$$\text{where } L_n(x) = \frac{1}{|n|} e^x D^n (x^n e^{-x}).$$

We have $L_0(x) = 1$

$$L_1(x) = \frac{1}{|1|} (1-x) = 1-x$$

Replacing x by $2x$, $L_1(2x) = 1-2x$

$$= 2(1-x) - 1 = 2L_1(x) - 2^0 \frac{1}{|1|} L_0(x)$$

$$= \left| 1 \sum_{m=0}^1 \frac{2^{1-m} (-1)^m}{|m| |1-m|} L_{1-m}(x) \right|$$

$$\text{Also, } L_2(2x) = \frac{1}{|2|} (2-8x+4x^2) = 1-4x+2x^2 = 4 \left(1-2x + \frac{x^2}{2} \right) - 4(1-x) + 1$$

$$= 2^2 L_2(x) - 2 \frac{|2|}{|1| |1|} L_1(x) + \frac{|2|}{|0| |2|} L_0(x)$$

$$= \left| 2 \sum_{m=0}^2 \frac{2^{2-m} (-1)^m}{|m| |2-m|} L_{2-m}(x) \right|$$

$$\begin{aligned} \text{and } L_3(3x) &= \frac{1}{|3|} (6-36x+36x^2-8x^3) = 1-6x+6x^2-\frac{4}{3}x^3 \\ &= 2^3 \left(1-3x + \frac{3}{2}x^2 - \frac{x^3}{6} \right) - 12 \left(1-2x + \frac{x^2}{2} \right) + 2.3(1-x) - 1 \\ &= 2^3 L_3(x) - \frac{2^2 |3|}{|1| |2|} L_2(x) + 2 \frac{|3|}{|2| |1|} L_1(x) - 2^0 \frac{|3|}{|3|} L_0(x) \\ &= \left| 3 \sum_{m=0}^3 \frac{2^{3-m} (-1)^m}{|m| |3-m|} L_{3-m}(x) \right| \end{aligned}$$

$$\text{Similarly } L_4(2x) = \left| 4 \sum_{m=0}^4 \frac{2^{4-m} (-1)^m}{|m| |4-m|} L_{4-m}(x) \right|$$

We can thus generalize the result as

$$L_n(2x) = \left| n \sum_{m=0}^n \frac{2^{n-m} (-1)^m}{|m| |n-m|} L_{n-1}(x) \right|$$

Problem 43. Prove the following:

$$(i) \frac{d}{dx} L_n^m(x) = L_n^{m+1}(x)$$

$$(ii) L_n^m(x) - n L_{n-1}^m(x) + n L_{n-1}^{m-1}(x) = 0.$$

$$(i) \text{ We have } L_n^m(x) = D^m L_n(x)$$

$$\therefore D \left[L_n^m(x) \right] = D^{m+1} L_n(x) = L_n^{m+1}(x)$$

$$\begin{aligned} (ii) \text{ L.H.S.} &= D^m \{ e^x D^n (x^n e^{-x}) - n D^{m-1} \{ e^x D^{n-1} (x^{n-1} e^{-x}) \} \\ &\quad + n D^{m-1} \{ e^x D^{n-1} (x^{n-1} e^{-x}) \} \} \\ &= D^{m-1} \{ e^x D^n (x^n e^{-x}) + e^x D^{n+1} (x^n e^{-x}) \} - n D^{m-1} \{ e^x D^{n-1} (x^{n-1} e^{-x}) \\ &\quad + e^x D^n (x^{n-1} e^{-x}) \} + n D^{m-1} \{ e^x D^{n-1} (x^{n-1} e^{-x}) \} \\ &= D^{m-1} \{ e^x D^n (x^n e^{-x}) + e^x D^n (-x^n e^{-x} + n x^{n-1} e^{-x}) \} \\ &\quad - n D^{m-1} \{ e^x D^{n-1} (x^{n-1} e^{-x}) + e^x D^n (x^{n-1} e^{-x}) \} + n D^{m-1} \{ e^x D^{n-1} (x^{n-1} e^{-x}) \} \\ &= n D^{m-1} \{ e^x D^n (x^{n-1} e^{-x}) \} - n D^{m-1} \{ e^x D^n (x^{n-1} e^{-x}) \} \\ &= 0. \end{aligned}$$

Problem 44. Prove the following:

$$(i) L_3^1(x) = -18 + 18x - 3x^2$$

$$(ii) L_4^2(x) = 144 - 96x + 12x^2$$

$$(iii) L_4^4(x) = 24.$$

Problem 45. Show that $L_n^m(2x) = \left| m+n \sum_{k=0}^n \frac{2^{n-k} (-1)^k}{|(m+n-k)| |k|} L_{n-k}^m(x) \right|$

Problem 46. Prove the following:

$$(i) L_n^m(x) - L_{n-1}^m(x) = L_{n-1}^{m-1}(x)$$

$$(ii) \frac{d}{dx} L_n^m(x) = -L_{n-1}^{m+1}(x)$$

$$(iii) n L_n^m(x) = (2n+m-1) L_{n-1}^m(x) - (n+m-1) L_{n-2}^m(x)$$

Problem 47. Establish the completeness of Laguerre functions.

We have considered the sequence of functions $\phi_1(x), \phi_2(x), \dots, \phi_n(x)$ such that

$$\int_a^b \phi_m(x) \phi_n(x) dx = 0 \text{ for } m \neq n$$

Then we call the functions $\phi_k(x), k=1, 2, 3, \dots$ to form an *orthogonal sequence* for the interval (a, b) .

In addition if $\int_a^b \{\phi_n(x)\}^2 dx = 1$ for all values of n , then we call the functions as to form an *orthonormal set*.

In case there is no integrable function $\psi(x) (\neq 0)$ such that

$$\int_a^b \psi(x) \phi_n(x) dx = 0 \text{ for all values of } n,$$

then we say that the sequence is a *complete orthogonal sequence*.

Problem 48. Find a series solution of

$$(i) xy'' + (1+x)y' + y = 0, \quad \text{Ans. } y = e^{-x} L_0(x) = e^{-x}$$

$$(ii) xy'' + (1+x)y' + 2y = 0, \quad \text{Ans. } y = e^{-x} L_1(x) = e^{-x}(1-x)$$

$$(iii) xy'' + (1+x)y' + \frac{3}{2}y = 0, \quad \text{Ans. } y = e^{-x} L_{\frac{1}{2}}(x)$$

$$(iv) y'' + \left(\frac{1}{4x^2} + \frac{1}{2x} - \frac{1}{4}\right)y = 0, \quad \text{Ans. } y = e^{-x/2} x^{1/2} L_0(x) = e^{-x/2} x^{1/2}$$

Problem 49. If a function $f(x)$ defined in $(0, \infty)$ is expressed as

$$f(x) = C_0 L_0(x) + C_1 L_1(x) + C_2 L_2(x) + \dots \quad \dots (1)$$

$$\text{then show that } C_k = \frac{\int_0^\infty e^{-x} L_k(x) f(x) dx}{\int_0^\infty e^{-x} [L_k(x)]^2 dx} \text{ for } k=0, 1, 2, \dots$$

Hint. Integrating over 0 to ∞ after multiplying (1) by $e^{-x} L_k(x)$ and using orthogonal properties i.e. $\int_0^\infty e^{-x} L_m(x) L_n(x) dx = 0, m \neq n$ the result is obtained.

Problem 50. Show that the Laguerre and Hermite Polynomials are related by

$$H_{2m}(x) = (-1)^m 2^{2m} \left| \frac{m}{\sqrt{\pi}} \right| L_m^{-1/2}(x^2)$$

$$\text{and } H_{2m+1}(x) = (-1)^m 2^{2m+1} \left| \frac{m}{\sqrt{\pi}} \right| x L_m^{1/2}(x^2).$$

8.9. RICCATI'S DIFFERENTIAL EQUATION

$$\text{In general form it is } x \frac{dy}{dx} - ay + by^2 = cx^n \quad \dots (1)$$

$$\text{But the particular form } \frac{dy}{dx} + by^2 = cx^m \quad \dots (2)$$

is commonly known as Riccati's equation.

Considering first (1) and changing the independent variable x to z by the transformation $z = x^a$ and dependent variable y to u by the transformation $y = uz$, it becomes

$$\frac{du}{dz} + \frac{b}{a} u^2 = \frac{c}{a} z^{\frac{n}{a}-2} \quad \dots (3)$$

which is of the form (2).

Now considering (2) and making a substitution $y = \frac{u}{x}$, it becomes

$$x \frac{du}{dx} - u + bu^2 = cx^n \text{ where } m = n - 2 \quad \dots (4)$$

which is of the form (1).

There arise two cases:

I. If $n = 2a$, (1) can be integrated in finite series

Put $y = ux^a$ in (1) which reduces to

$$x^{a+1} \frac{du}{dx} + bx^{2a} u^2 = cx^n$$

$$\text{or } x^{1-a} \frac{du}{dx} + bu^2 = cx^{n-2a} = c \text{ for } n = 2a$$

$$\therefore x^{1-a} \frac{du}{dx} = c - bu^2$$

$$\text{or } \frac{du}{c - bu^2} = x^{a-1} dx$$

which is integrable as variables are separated.

II. If $\frac{n \pm 2a}{2n}$ is a positive integer, (1) is integrable in finite terms.

Put $y = A + \frac{x^n}{y_1}$ in (1) which reduces to

$$-aA + bA^2 + (n-a+2bA) \frac{x^n}{y_1} + b \frac{x^{2n}}{y_1^2} - \frac{x^{n+1}}{y_1^2} \frac{dy_1}{dx} = cx^n$$

Choosing A such that $-aA + bA^2 = 0$ i.e. $A = a/b$ so that the substitution becomes $y = \frac{a}{b} + \frac{x^n}{y_1}$ and the reduced equation is

$$x \frac{dy_1}{dx} - (a+n) y_1 + c y_1^2 = b x^n \quad \dots (5)$$

which retains the form (1) except that a has been replaced by $a+n$; b by c and c by b .

Making a second substitution $y_1 = \frac{a+n}{c} + \frac{x^n}{y_2}$, the last equation reduces to $x \frac{dy_2}{dx} - (a+2n)y_2 + by_2^2 = cx^n$... (6)

Hence, making such k transformations we find,

$$x \frac{dy_k}{dx} - (a+kn)y_k + cy_k^2 = bx^n \text{ when } k \text{ is odd} \quad \dots (7)$$

$$\text{and } x \frac{dy_k}{dx} - (a+kn)y_k + cy_k^2 = bx^n \text{ when } k \text{ is even} \quad \dots (8)$$

In either case the equation is integrable in finite terms by I when $n=2(a+kn)$ i.e. if $\frac{n-2a}{2n}$ is a positive integer.

Now with the choice $A=0$; the first substitution $y = \frac{x^n}{y_1}$ reduces (1) to $x \frac{dy_1}{dx} - (n-a)y_1 + cy_1^2 = bx^n$... (9)

which is the same as (5) except that the sign of a is reversed.

The second substitution $y = \frac{n-a}{c} + \frac{x^n}{y_2}$ yields

$$x \frac{dy_2}{dx} - (2n-a)y_2 + by_2^2 = cx^n \quad \dots (10)$$

Making such k transformations, we have

$$x \frac{dy_k}{dx} - (kn-a)y_k + cy_k^2 = bx^n \text{ when } k \text{ is odd} \quad \dots (11)$$

$$\text{and } x \frac{dy_k}{dx} - (kn-a)y_k + by_k^2 = cx^n \text{ when } k \text{ is even} \quad \dots (12)$$

In either case the equation is integrable in finite terms if $n=2(kn-a)$ i.e. if $\frac{n+2a}{2n}$ is a positive integer.

But since $m=n-2$ is the condition for integrability of (1), we have $m+2 \pm 2 = 2k(m+2)$

Taking negative sign, $m = -\frac{4k}{2k-1}$

and taking positive sign, $m = \frac{4(k-1)}{2k-1} = \frac{4k'}{2k'+1}$ when $k'=k-1$.

Hence Riccati's equation is integrable in finite terms if $m = -\frac{4k}{2k \pm 1}$ where k is zero or a positive integer. The integration is carried by the substitution $y = y_1 + \frac{1}{v}$, where y_1 is supposed to be a known particular

integral of the equation, whence the transformed equation becomes linear.

In fact Riccati's equation and its general form are particular cases of the equation $\frac{dy}{dx} = P + Qy + Ry^2$

where P, Q, R are functions of x , since the substitution

$$y = -\frac{1}{R} - \frac{1}{u} \frac{du}{dx} \text{ reduces it to the form}$$

$$\frac{d^2u}{dx^2} - \left(Q + \frac{1}{R} \frac{dR}{dx} \right) \frac{du}{dx} + PRu = 0$$

which is integrable.

COROLLARY. Relation between Riccati's and Bessel's Equations.

$$\text{Riccati's equation is } \frac{dy}{dx} + by^2 = cx^m \quad \dots (13)$$

which is non-linear first order equation.

$$\text{Put } by = \frac{1}{v} \frac{dv}{dx} \text{ so that } b \frac{dy}{dx} = \frac{1}{v} \frac{d^2v}{dx^2} - \frac{1}{v^2} \left(\frac{dv}{dx} \right)^2$$

then (13) becomes

$$\frac{d^2v}{dx^2} - bcvx^m = 0 \quad \dots (14)$$

Taking $bc = a^2$, this yields

$$\frac{d^2v}{dx^2} - a^2x^mv = 0 \quad \dots (15)$$

when b and c have same sign (in case exponential functions occur in y)

$$\text{and } \frac{d^2v}{dx^2} + a^2x^mv = 0 \quad \dots (16)$$

when b and c have opposite sign (in case circular functions occur in y)

Changing x to z by $qz = x^a$ when $q = \frac{1}{2} m+1 = \frac{1}{n}$ (say).

The reduced equation is

$$\frac{d^2v}{dz^2} - \frac{n-1}{z} \frac{dv}{dz} - bcv = 0 \quad \dots (17)$$

which is integrable in finite form if

$$\frac{1}{n} = \frac{1}{2} m+1 = 1 - \frac{2k}{2k \pm 1} = \frac{\pm 1}{2k \pm 1}$$

i.e. if n is an odd integer (17) can be written as by putting $n-1=2p$,

$$\frac{d^2v}{dz^2} - \frac{2p}{z} \frac{dv}{dz} - bcv = 0 \quad \dots (18)$$

which is integrable in a finite form when p is an integer.

Reducing it to normal form by the substitution $vz^{-p}=w$, we have

$$\frac{d^2w}{dz^2} - bcw = \frac{p(p+1)}{z^2} w \quad \dots (19)$$

If we further reduce it by the substitution $w=t\sqrt{z}$, we find

$$z^2 \frac{d^2t}{dz^2} + z \frac{dt}{dz} + \left[(-bc)z^2 - \left(p + \frac{1}{2} \right)^2 \right] t = 0$$

$$\text{or } \frac{d^2t}{dz^2} + \frac{1}{z} \frac{dt}{dz} + \left[(-bc) - \frac{\left(p + \frac{1}{2} \right)^2}{z^2} \right] t = 0 \quad \dots (20)$$

which is clearly Bessel's equation with solution

$$t = A J_{p+1/2} \{ z(-bc)^{1/2} \} + B J_{-(p+1/2)} \{ z(-bc)^{1/2} \}$$

Problem 51. Solve $\frac{dy}{dx} = \cos x - y \sin x + y^2$

Since $y = \sin x$ is a particular solution of the given equation, we therefore, have $y_1 = \sin x$.

$$\text{Put } y = \sin x + \frac{1}{v} \text{ i.e. } \frac{dy}{dx} = \cos x - \frac{1}{v^2} \frac{dv}{dx},$$

The reduced equation is

$$\frac{dv}{dx} + v \sin x = -1$$

which is linear in v and hence its integrating factor

$$= e^{\int \sin x dx} = e^{-\cos x}$$

Thus the solution is

$$v \cdot e^{-\cos x} = A - \int e^{-\cos x} dx$$

$$\text{or } v = A e^{\cos x} - e^{\cos x} \int e^{-\cos x} dx$$

$$\text{i.e. } y = \sin x + \left[A e^{\cos x} - e^{\cos x} \int e^{-\cos x} dx \right]^{-1}$$

$$\text{or } \frac{1}{y - \sin x} = A e^{\cos x} - e^{\cos x} \int e^{-\cos x} dx.$$

8.10. THE DIRAC-DELTA FUNCTION WITH ITS FORMAL PROPERTIES

Consider a function $\delta(x)$ which is zero everywhere except at $x=0$ and tends to ∞ in such a manner that $\int_{-\infty}^{\infty} \delta(x) dx = 1$... (1)

$$\text{with } \begin{cases} \delta(t) = 0 & \text{if } t \neq 0 \\ = \infty & \text{if } t = 0 \end{cases} \quad \dots (2)$$

This is known as *Dirac-delta function* and used in mathematical physics wherever exist functions with non-zero values in very short

interval e.g. an impulsive force acting for a short while is defined as $\delta(x-\xi)$ by $\lim_{a \rightarrow 0} c e^{-(x-\xi)^2/a}$

where the constant $C(a)$ is chosen such that $\int_{-\infty}^{\infty} \delta(x-\xi) dx = 1$

and hence using the mean value theorem of integral calculus, we have

$$\int_{-\infty}^{\infty} f(x) \delta(x-\xi) dx = f(\xi).$$

Let us again consider a function

$$\delta_a(x) = \begin{cases} \frac{1}{2a}, & -a < x < a \\ 0, & |x| > a \end{cases} \quad \dots (3)$$

$$\begin{aligned} \text{Then, } \int_{-\infty}^{\infty} \delta_a(x) dx &= \int_{-\infty}^{-a} \delta_a(x) dx + \int_{-a}^a \delta_a(x) dx + \int_a^{\infty} \delta_a(x) dx \\ &= 0 + \int_{-a}^a \frac{1}{2a} dx + 0 = \frac{1}{2a} [a - (-a)] \\ &= 1 \end{aligned} \quad \dots (4)$$

In case $f(x)$ is integrable in the interval $(-a, a)$, then from mean value theorem, $\int_{-\infty}^{\infty} f(x) \delta_a(x) dx = \frac{1}{2a} \int_{-a}^a f(x) dx = f(\theta a); |\theta| \leq 1$

Let us now define $\delta(x) = \lim_{a \rightarrow 0} \delta_a(x)$

As such (3) and (4) yield

$$\delta(x) = 0, \text{ when } x \neq 0 \quad \dots (5)$$

$$\text{and } \int_{-\infty}^{\infty} \delta(x) dx = 1 \quad \dots (6)$$

which define Dirac-delta function.

Further, since we have

$$\int_{-a}^a f(x) \delta_a(x) dx = \frac{1}{2a} \int_{-\infty}^{\infty} f(x) dx = f(\theta a), \quad |\theta| \leq 1$$

$$\therefore \int_{-\infty}^{\infty} f(x) dx = f(0) \quad \dots (7)$$

which by change of variable, reduces to

$$\int_{-\infty}^{\infty} f(x) \delta(x-a) dx = f(a) \quad \dots (8)$$

$$\text{or symbolically, } f(x) \delta(x-a) = f(a) \delta(x-a) \quad \dots (9)$$

$$\text{In case } f(x) = x, (9) \text{ yields, } x \delta x = 0 \quad \dots (10)$$

In a similar manner we can show

$$\delta(-x) = \delta x \quad \dots (11)$$

$$\delta(ax) = \frac{1}{a} \delta(x), \quad a > 0 \quad \dots (12)$$

$$\delta(a^2 - x^2) = \frac{1}{2a} \{ \delta(x-a) + \delta(x+a) \}, a > 0 \quad \dots (13)$$

Now assuming that $\delta'(x)$ i.e., differential of $\delta(x)$ exists and regarding $\delta'(x)$ and $\delta(x)$ both as ordinary functions in the rule for integrating by parts, we have

$$\begin{aligned} \int_{-\infty}^{\infty} f(x) \delta'(x) dx &= \left[f(x) \delta(x) \right]_{-\infty}^{\infty} - \int_{-\infty}^{\infty} f'(x) \delta(x) dx \\ &= 0 - f'(0) \text{ by (7)} \\ &= -f'(0) \end{aligned} \quad \dots (14)$$

If $\delta^{(n)}(x)$ be the n th derivative of $\delta(x)$, then similarly we find on repeating this process n times,

$$\int_{-\infty}^{\infty} f(x) \delta^{(n)}(x) dx = (-1)^n f^{(n)}(0) \quad \dots (15)$$

Problem 52. Show that the function $\delta(x) = \lim_{\epsilon \rightarrow 0} \frac{\sin(2\pi\epsilon x)}{\pi x}$ is a Dirac-delta function.

$$\text{We have } \delta(x) = \lim_{\epsilon \rightarrow 0} \frac{\sin(2\pi\epsilon x)}{\pi x}$$

$$\therefore \delta(x) = 0, \text{ when } x \neq 0. \quad \dots (1)$$

and $\int_{-\infty}^{\infty} \frac{\sin(2\pi\epsilon x)}{\pi x} dx = 2 \int_0^{\infty} \frac{\sin(2\pi\epsilon x)}{\pi x} dx$, the function being even.

$$= \frac{2}{\pi} \cdot \frac{\pi}{2} = 1 \quad \dots (2)$$

It follows from (1) and (2) that the given function is Dirac-delta function.

8.11. RIEMANN-ZETA FUNCTION

If $s = \sigma + it$ where σ and t are real and if $\epsilon > 0$, then the series

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad \dots (1)$$

is a uniformly convergent series of analytic functions in a domain in which $\sigma \geq 1 + \epsilon$ and hence the series is an analytic function of s in that domain. This function $\zeta(s)$ is called as *Riemann-zeta-function*.

The *generalized zeta function* for $\sigma \geq 1 + \epsilon$ is defined as

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(a+n)^s} \quad \dots (2)$$

where a is a constant.

If $0 < a \leq 1$ and $\arg(a+n) = 0$, then $\zeta(s, 1) = \zeta(s)$ (3)

The expression of $\zeta(s, a)$ as an infinite integral, when $\sigma > 1 + \epsilon$ and $\arg x = 0$, is

$$\zeta(s, a) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{x^{s-1} e^{-ax}}{1 - e^{-x}} dx \quad \dots (4)$$

The expression of $\zeta(s, a)$ as a contour integral is

$$\zeta(s, a) = - \frac{\Gamma(1-s)}{2\pi i} \int_{\infty}^{(0+)} \frac{(-z)^{s-1} e^{-az}}{1 - e^{-z}} dz \quad \dots (5)$$

Riemann's relation between $\zeta(s)$ and $\zeta(1-s)$ is

$$2^{1-s} \Gamma(s) \zeta(s) \cos\left(\frac{1}{2}s\pi\right) = \pi^s \zeta(1-s) \quad \dots (6)$$

Hermite's formula for $\zeta(s, a)$ is

$$\zeta(s, a) = \frac{1}{2} a^{-s} + \frac{a^{1-s}}{s-1} + 2 \int_0^{\infty} (a^2 + y^2)^{-\frac{1}{2}s} \left\{ \sin\left(s \arctan \frac{y}{a}\right) \right\} \frac{dy}{e^{2\pi y} - 1} \quad \dots (7)$$

$$\therefore \zeta(0, a) = \frac{1}{2} - a \quad \dots (8)$$

$$\text{and } \left\{ \frac{d}{ds} \zeta(s, a) \right\}_{s=0} = \log \Gamma(a) - \frac{1}{2} \log(2\pi) \quad \dots (9)$$

$$\therefore \zeta'(0) = -\frac{1}{2} \log(2\pi) \quad \dots (10)$$

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 53. Solve the equation $x^2 y'' + xy' + (x^2 - n^2) y = 0$, $n = \text{constant}$, in the neighbourhood of the point $x=0$. What are the singular points of the equation? (Bombay, 1965)

Problem 54. Prove that if the functions $P_l(x)$ are defined by the equation

$$\frac{1}{\sqrt{1-2xt+t^2}} = \sum_{l=0}^{\infty} P_l(x) t^l$$

then $P_l(x)$ is a polynomial of l^{th} degree, satisfying the following relations:

$$(a) \int_{-1}^1 P_l(x) P_l'(x) dx = \frac{2}{2l+1} \delta_{ll'}$$

$$(b) (l+1) P_{l+1}(x) - (2l+1)x P_l(x) + l P_{l-1}(x) = 0 \quad (\text{Bombay, 1965})$$

Problem 55. Define Bessel functions and obtain the series solution of the Bessel differential equation for $n=0$ either by the use of operators or otherwise.

$$\text{Show that } \left[J_{\frac{1}{2}}(x) \right]^2 + \left[J_{-\frac{1}{2}}(x) \right]^2 = \frac{2}{\pi x} \quad (\text{Nagpur, 1965})$$

Problem 56. What are Legendre Polynomials? Show that

$$(i) (m+1) P_{m+1}(x) = (2m+1)x P_m(x) - m P_{m-1}(x)$$

$$(ii) (x^2-1) P'_m(x) = mx P_m(x) - m P_{m-1}(x), \text{ where } P_m(x) = \frac{d}{dx} P_m(x)$$

$$(iii) P_{2m}(-x) = P_{2m}(x)$$

$$(iv) P_{2m+1}(-x) = -P_{2m+1}(x) \quad (\text{Agra, 1965})$$

Problem 57. Write the differential equation satisfied by Bessel's function of order n .

Express the following Bessel functions in terms of trigonometric functions:

$$(i) J_{1/2}(x), (ii) J_{-1/2}(x); (iii) J_{3/2}(x), (iv) J_{-3/2}(x) \quad (\text{Agra, 1966, Vikram '67})$$

Problem 58. The generating function of Legendre Polynomial is

$$T(w, s) = (1 - 2ws + s^2)^{-1/2} = \sum_{l=0}^{\infty} s^l P_l(w)$$

Using this prove the following properties of Legendre polynomials

$$(i) \int_{-1}^1 P_m(w) P_n(w) dw = \frac{2}{2m+1} \delta_{mn}$$

$$(ii) P_m(-w) = (-1)^m P_m(w)$$

$$(iii) \sum_{l=0}^{\infty} (2l+1) P_l(w) = 0 \text{ for } w \neq \pm 1 \quad (\text{Agra, 1967})$$

Problem 59. Show that $4x^2 - 2$ is a polynomial solution of the differential equation,

$$x^2 \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + 2\alpha y = 0$$

where α is a positive integer.

(Vikram, 1967)

Hint. Given equation is Hermite differential equation and show that $H_2(x) = 4x^2 - 2$.

Problem 60. Show that $\int_0^{\infty} (\cos xy) (\cos xy') dx = \frac{1}{2} \pi \delta(y-y')$ with $y > 0, y' > 0$ and $\delta(y-y')$ being the Dirac-deltas function. (Agra, 1968)

Problem 61. The generating function for Bessel functions of integral orders is

$$G(x, h) = e^{x/2(h-h^{-1})} = \sum_{n=-\infty}^{\infty} h^n J_n(x)$$

(a) Show by direct substitution that $G(x, h)$ satisfies the equation

$$x^2 \frac{\partial^2 G}{\partial x^2} + x \frac{\partial G}{\partial x} + x^2 G - \left(h \frac{\partial}{\partial h} \right)^2 G = 0$$

use this result to show that $J_n(x)$ satisfies Bessel's differential equation,

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - n^2) y = 0$$

(b) In the generating function, putting $h = e^{i\theta}$, show that

$$\cos(x \sin \theta) = J_0(x) + 2 \sum_{n=1}^{\infty} J_{2n}(x) \cos 2n\theta$$

$$\sin(x \sin \theta) = 2 \sum_{n=0}^{\infty} J_{2n+1}(x) \sin(2n+1)\theta. \quad (\text{Agra, 1968})$$

Problem 62. Solve by series integration, the equation

$$(1-x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + \lambda(\lambda+1) y = 0$$

where λ is an integer. Discuss the nature of the solution for $\lambda = 1, 2$. (Vikram, 1969)

Problem 63. Write short note on Bessel's functions. (Vikram, 1969)

Problem 64. Solve the equation

$$(1-x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + n(n+1) y = 0$$

where n is a positive integer. If $P_n(x)$ is the Polynomial solution, prove that these functions form an orthogonal system. (Agra, 1969)

Problem 65. (a) Solve the equation in series

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + \left(x^2 - \frac{1}{4} \right) y = 0$$

(b) Show that $\int_{-1}^1 P_m(x) P_n(x) dx = 0$ unless $m=n$

$$= \frac{2}{2n+1} \text{ if } m=n$$

m, n being positive integers.

(Agra, 1970)

Problem 66. Obtain the solution of the Bessel's differential equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - n^2) y = 0$$

in the form of a power series. Discuss this solution in the neighbourhood of $x=0$.

Show that $\frac{d}{dx} [x^n J_n(x)] = x^n J_{n-1}(x)$ ($n=1, 2, 3, \dots$) (Bombay, 1970)

Problem 67. (a) Show that Bessel functions defined by $e^{x/2(t-1/t)}$

$= \sum_{n=-\infty}^{\infty} J_n(x) t^n$ have the integral representation

$$J_n(x) = \frac{1}{2\pi} \int_0^{2\pi} \cos(x \sin \theta - n\theta) d\theta \quad (\text{Agra, 1973})$$

(b) If α and β are the roots of the equation $J_0(x) = 0$, show that

$$\int_0^1 x J_0(\alpha x) J_0(\beta x) dx = \frac{1}{2} \delta_{\alpha\beta} J_2(\alpha) \quad (\text{Agra, 1971})$$

Problem 68. Explain the significance of spherical harmonics and discuss some general properties of harmonic functions. (Agra, 1971)

Problem 69. Prove that the polynomial solution $P_n(x)$ of the Legendre equation

$$(1-x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + n(n+1) y = 0$$

satisfies the orthogonality condition $\int_{-1}^1 P_n(x) P_m(x) dx = \frac{2}{2n+1} \delta_{nm}$

Show that this property allows for the expansion of an arbitrary function $f(x)$ into a series of Legendre polynomials provided that $f(x)$ is sectionally continuous in $-1 < x < 1$ and $\frac{df(x)}{dx}$ is sectionally continuous in $-1 < x < 1$. (Agra, 1972)

Problem 70. Prove that $[J_0(x)]^2 + 2[J_1(x)]^2 + 2[J_2(x)]^2 + \dots = 1$. (Agra, 1973)

Problem 71. Prove that the function $y = \frac{d^n}{dx^n} (x^2 - 1)^n$

satisfies Legendre's differential equation

$$(1-x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + n(n+1) y = 0$$

Hence obtain Rodrigue's formula for Legendre-polynomials $P_n(x)$.

Using this formula prove that

$$\int_{-1}^1 x^m P_n(x) dx = 0 \text{ for } m < n. \quad (\text{Agra, 1973})$$

Problem 72. Show that $\int_{-1}^1 x P_n(x) P_{n-1}(x) dx = \frac{2n}{4n^2 - 1}$ (Agra, 1974)

Problem 73. Express the electrostatic potential at a point P in space, due to two equal but opposite point charges such that the distance between them is small compared to their distance from P , in terms of Legendre-polynomials. (Agra, 1975)

We know that the potential at a point $P(x, y, z)$ distant r from O (origin), and ρ from z -axis such that OP makes an angle θ with z -axis, due to a charge q at $z = \zeta$ is given by

$$\phi(r, \theta) = \frac{q}{4\pi\epsilon} \cdot \frac{1}{\rho}, \quad \epsilon \text{ being inductive capacity.}$$

Here ρ is given by

$$\rho = (r^2 + \zeta^2 - 2r\zeta \cos \theta)^{1/2}$$

$$\text{or } \frac{1}{\rho} = \frac{1}{\zeta} \left[1 + \left(\frac{r}{\zeta} \right)^2 - 2 \frac{r}{\zeta} \cos \theta \right]^{-1/2}$$

$= \frac{1}{\zeta} \left[1 + \frac{r}{\zeta} \cos \theta + \left(\frac{r}{\zeta} \right)^2 \left(\frac{3}{2} \cos^2 \theta - \frac{1}{2} \right) + \dots \right]$, on expanding by Binomial theorem under assumption $\left| \left(\frac{r}{\zeta} \right)^2 - 2 \frac{r}{\zeta} \cos \theta \right| < 1$

$$= \frac{1}{\zeta} \sum_{n=0}^{\infty} P_n(\cos \theta) \left(\frac{r}{\zeta} \right)^n, r < \zeta$$

where coefficients of $\frac{r}{\zeta}$ being polynomials in $\cos \theta$ are termed as Legendre polynomials, such that

$$P_0(\cos \theta) = 1, P_1(\cos \theta) = \cos \theta,$$

$$P_2(\cos \theta) = \frac{1}{2} (3 \cos^2 \theta - 1) = \frac{1}{4} (3 \cos 2\theta + 1)$$

$$P_3(\cos \theta) = \frac{1}{2} (5 \cos^3 \theta - 3 \cos \theta) = \frac{1}{8} (5 \cos 3\theta + 3 \cos \theta) \text{ etc...}$$

In case $r > \zeta$, we have on interchanging r and ζ ,

$$\frac{1}{\rho} = \frac{1}{r} \left[1 + \left(\frac{\zeta}{r} \right)^2 - 2 \frac{\zeta}{r} \cos \theta \right]^{-1/2}$$

$$= \frac{1}{r} \sum_{n=0}^{\infty} P_n(\cos \theta) \left(\frac{\zeta}{r} \right)^n$$

Let us now consider the potential at P due to a dipole consisting of a point charge $+q$ at a point $z=d$ on the x -axis and an equal but opposite charge $-q$ at O , the origin, such that the distance of P is very large as compared to d .

In view of the last relation, the potential at a distant point P is given by

$$\phi = \frac{q}{4\pi\epsilon} \left(\frac{1}{\rho} - \frac{1}{r} \right)$$

$$= \frac{q}{4\pi\epsilon} \cdot \frac{1}{r} \left[\sum_{n=0}^{\infty} P_n(\cos \theta) \left(\frac{d}{r} \right)^n - 1 \right]$$

$$= \frac{q}{4\pi\epsilon} \cdot \frac{1}{r} \left[1 + \frac{d}{r} \cos \theta + \left(\frac{d}{r} \right)^2 \left(\frac{3}{2} \cos^2 \theta - \frac{1}{2} \right) + \dots - 1 \right]$$

$$= \frac{qd}{4\pi\epsilon} \cdot \frac{\cos \theta}{r^2}, \text{ neglecting higher order terms as } d \ll r.$$

Problem 74. For Bessel function $J_n(x)$, find out a and b , where

$$\frac{d}{dx} J_n(x) = a J_{n-1}(x) + b J_{n+1}(x)$$

(Agra, 1975)

Hint. Compare with the recurrence relation

$$J_{n-1}(x) - J_{n+1}(x) = 2 \frac{d}{dx} J_n(x).$$

$$\text{Ans. } a = \frac{1}{2}, b = -\frac{1}{2}.$$

Problem 75. Solve the differential equation

$$(1-x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + l(l+1)y = 0$$

where l is a positive integer. Discuss the orthogonal properties of the functions obtained as solution. (Agra, 1975, 1976)

Problem 76. (a) For Bessel function $J_n(x)$ prove that

$$J_n(x) = \frac{1}{\pi} \int_0^\pi \cos(n\theta - x \sin \theta) d\theta$$

and hence show that $J_0(x) = \frac{2}{\pi} \int_0^1 \frac{\cos xt}{\sqrt{1-t^2}} dt$

(b) Show that the recurrence relation

$$J_n'(x) = \frac{1}{2} [J_{n-1}(x) - J_{n+1}(x)]$$

follows directly from differentiation of

$$J_n(x) = \frac{1}{\pi} \int_0^\pi \cos(n\theta - x \sin \theta) d\theta$$

(Rohilkhand, 1976)

Hint. For (a): In a section it has been proved that

$$J_n(x) = \frac{1}{\pi} \int_0^\pi \cos(n\theta - x \sin \theta) d\theta$$

$$\text{Put } \theta = 0, J_0(x) = \frac{1}{\pi} \int_0^\pi \cos(x \sin \theta) d\theta$$

$$= \frac{2}{\pi} \int_0^{\pi/2} \cos(x \sin \theta) d\theta \text{ (being even function of } \theta)$$

$$\text{Now put } \sin \theta = t \text{ so that } d\theta = \frac{dt}{\cos \theta} = \frac{dt}{\sqrt{1-t^2}}$$

$$\text{Hence } J_0(x) = \frac{2}{\pi} \int_0^1 \frac{\cos xt}{\sqrt{1-t^2}} dt.$$

Problem 77. (a) Derive the Rodrigue's formula for the Legendre polynomial.

(b) Using Rodrigue's formula prove that

$$\int_{-1}^1 x^m P_n(x) dx = 0 \text{ if } m < n.$$

(Rohilkhand, 1976)

Hint. For (b): Using Rodrigue's formula, we have

$$\int_{-1}^1 x^m P_n(x) dx = \frac{1}{2^n n!} \int_{-1}^1 x^m \left(\frac{d}{dx} \right)^n (x^2-1)^n dx$$

On integrating R.H.S. by parts repeatedly, the first term always vanishes and we are left with

$$\int_{-1}^1 x^m P_n(x) dx = \frac{(-1)^m}{2^n n!} \int_{-1}^1 \left(\frac{d}{dx} \right)^{n-m} (x^2-1)^n dx$$

$$= \frac{(-1)^m}{2^n n!} \left[\left(\frac{d}{dx} \right)^{n-m-1} (x^2-1)^n \right]_{-1}^1 = 0 \text{ for } m < n.$$

Problem 78. Show that the coefficient of t^n in the expansion of

$$e^{i(t-t^{-1})^{1/2}} \text{ equals } \frac{1}{\pi} \int_0^\pi \cos(n\phi - x \sin \phi) d\phi$$

(Agra, 1976)

Hint: The coefficient of t^n in $e^{i(t-t^{-1})^{1/2}}$ is $P_n(x)$ which is also equal to

$$\frac{1}{\pi} \int_0^\pi \cos(n\phi - x \sin \phi) d\phi \text{ and hence the proposition.}$$

Problem 79. Show that two independent solutions of Bessel's differential equation $x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - n^2)y = 0$; cannot be obtained by series integration method. Explain how the second solution is obtained in this case. If $J_0(\rho\lambda)$ is Bessel's function of zero order and $z > 0$, show that

$$\int_0^\infty e^{-z\lambda} J_0(\rho\lambda) d\lambda = \frac{1}{\sqrt{\rho^2 + z^2}}.$$

(Rohilkhand, 1977)

Problem 80. Define associated Legendre' polynomials and prove their Orthogonality condition.

If $P_n^m(x)$ are the Associated Legendre' polynomials, show that

$$P_n^{-m}(x) = (-1)^m \frac{n-m}{n+m} P_n^m(x).$$

(Rohilkhand, 1977)

CHAPTER 9

FOURIER'S SERIES, INTEGRALS AND TRANSFORMS

9.1. DEFINITION AND EXPANSION OF A FUNCTION OF x

A Fourier series is a representation employed to express a periodic function $f(x)$ defined in an interval say $(-\pi, \pi)$ a linear relation between the sines and cosines of the same period, viz.

$$\begin{aligned} f(x) &= a_0 + a_1 \cos x + a_2 \cos 2x + a_3 \cos 3x + \dots + a_n \cos nx + \dots \\ &\quad + b_1 \sin x + b_2 \sin 2x + b_3 \sin 3x + \dots + b_n \sin nx + \dots \\ &= a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx. \end{aligned} \quad \dots (1)$$

In order to determine the values of the coefficients a_0 , a_n and b_n , let us first integrate both the sides of (1) between the limits $-\pi$ and π , whence we get

$$\begin{aligned} \int_{-\pi}^{\pi} f(x) dx &= a_0 \int_{-\pi}^{\pi} dx \text{ other integrals vanishing,} \\ \text{i.e. } a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(v) dv, \end{aligned}$$

replacing x by v to distinguish from $f(x)$.

Again multiplying both sides of (1) by $\cos nx$ and integrating between the limits $-\pi$ and π , we get

$$\begin{aligned} \int_{-\pi}^{\pi} f(x) \cos nx dx &= a_n \int_{-\pi}^{\pi} \cos^2 nx dx \\ &\quad \text{other integrals vanishing} \\ &= \frac{a_n}{2} \int_{-\pi}^{\pi} (1 + \cos 2nx) dx = \frac{a_n}{2} \cdot 2\pi \end{aligned}$$

$$\therefore a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx = \frac{1}{\pi} \int_{-\pi}^{\pi} f(v) \cos nv dv.$$

Further multiplying both sides of (1) by $\sin nx$ and integrating between the limits $-\pi$ and π , we get as above

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(v) \sin nv dv.$$

Hence

$$\begin{aligned} f(x) &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(v) dv + \frac{1}{\pi} \sum_{n=1}^{\infty} \cos nx \int_{-\pi}^{\pi} f(v) \cos nv dv \\ &\quad + \frac{1}{\pi} \sum_{n=1}^{\infty} \sin nx \int_{-\pi}^{\pi} f(v) \sin nv dv \end{aligned} \quad \dots (2)$$

The expansion on R.H.S. of (2) is known as *Fourier's series for $f(x)$* and a_0 , a_n , b_n etc, are known as *Fourier's constants for $f(x)$* .
Now (2) may also be written as

$$f(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(v) dv + \frac{1}{\pi} \sum_{n=1}^{\infty} \int_{-\pi}^{\pi} f(v) \cos n(x-v) dv \quad \dots (3)$$

which is valid for $-\pi \leq x \leq \pi$

Deductions from (2)

(i) If $f(x)$ be an odd function of x , i.e. if $f(-x) = -f(x)$, then

$$\int_{-\pi}^{\pi} f(v) dv = 0.$$

$$\text{Also } \int_{-\pi}^{\pi} \cos nv f(v) dv = 0$$

$$\text{and } \int_{-\pi}^{\pi} \sin nv f(v) dv = 2 \int_0^{\pi} \sin nv f(v) dv. \quad \dots (4)$$

$$\text{As such } f(x) = \frac{2}{\pi} \sum_{n=1}^{\infty} \sin nx \int_0^{\pi} \sin nv f(v) dv.$$

(ii) If $f(x)$ be an even function of x , i.e., $f(-x) = f(x)$,

$$f(x) = \frac{1}{\pi} \int_0^{\pi} f(v) dv + \frac{2}{\pi} \sum_{n=1}^{\infty} \cos nx \int_0^{\pi} \cos nv f(v) dv. \quad \dots (5)$$

COROLLARY 1. To find a cosine series for $f(x)$ when $0 \leq x \leq \pi$, let us assume that

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx. \quad \dots (6)$$

Integrating both sides from 0 to π , we have

$$\begin{aligned} \int_0^{\pi} f(x) dx &= \int_0^{\pi} a_0 dx \text{ other integrals vanishing} \\ &= a_0 \pi. \end{aligned}$$

$$\therefore a_0 = \frac{1}{\pi} \int_0^{\pi} f(v) dv.$$

Again multiplying both sides of (6) by $\cos nx$ and integrating from 0 to π , we get

$$\begin{aligned} \int_0^{\pi} \cos nx f(x) dx &= a_n \int_0^{\pi} \cos^2 nx dx \\ &\quad \text{other integrals vanishing} \\ &= \frac{a_n}{2} \int_0^{\pi} (1 + \cos 2nx) dx \\ &= \frac{a_n}{2} \cdot \pi. \end{aligned}$$

$$\therefore a_n = \frac{2}{\pi} \int_0^{\pi} \cos nv f(v) dv.$$

Hence

$$f(x) = \frac{1}{\pi} \int_0^\pi f(v) dv + \frac{2}{\pi} \sum_{n=1}^{\infty} \cos nx \int_0^\pi \cos nv f(v) dv \quad \dots (7)$$

COROLLARY 2. To find a sine series for $f(x)$ when $0 \leq x \leq \pi$.
(Agra, 1961)

Let us assume that

$$f(x) = \sum_{n=1}^{\infty} b_n \sin nx.$$

Multiplying both sides by $\sin nx$ and integrating from 0 to π , we have $\int_0^\pi f(x) \sin nx dx = \frac{1}{2} b_n \int_0^\pi dx$, other integrals vanishing

$$= \frac{\pi}{2} \cdot b_n,$$

$$i.e. \quad b_n = \frac{2}{\pi} \int_0^\pi \sin nv f(v) dv.$$

$$\text{Hence} \quad f(x) = \frac{2}{\pi} \sum_{n=1}^{\infty} \sin nx \int_0^\pi f(v) \sin nv dv. \quad \dots (8)$$

Important Remark

We generally take the Fourier series expansion in the interval $(-\pi, \pi)$ with period 2π , as

$$f(x) = \frac{a_0}{2} + (a_1 \cos x + b_1 \sin x) + \dots + (a_n \cos nx + b_n \sin nx) + \dots$$

$$= \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx) \quad \dots (9)$$

where

$$\left. \begin{aligned} a_0 &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx \end{aligned} \right\} \quad \dots (10)$$

and

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$

In case the interval is $(-l, l)$ with period $2l$, we have

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l} \right) \quad \dots (11)$$

$$\text{when } a_n = \frac{1}{l} \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx \text{ and } b_n = \frac{1}{l} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx \quad \dots (12)$$

[These are obtained by replacing x by $\frac{\pi x}{l}$ in (9) and (10)]

12. DIRICHLET'S CONDITIONS

A function $f(x)$ is said to satisfy the Dirichlet's condition in any interval (a, b) in which the function is defined, if it is subjected to either of the conditions.

(i) $f(x)$ is bounded in (a, b) i.e., there exists an upper bound M such that $|f(x)| \leq M$ for all the values of x in (a, b) and the interval (a, b) can be divided into a finite number of open sub-intervals in each of which the function $f(x)$ is monotonic.

(ii) $f(x)$ has a finite number of points of infinite discontinuity in the interval (a, b) , but when the arbitrary small neighbourhoods of these points are excluded then $f(x)$ remains bounded in the deleted interval and the interval can be divided into a finite number of open sub-intervals in each of which $f(x)$ is monotonic. Also the infinite integral

$$\int_a^b f(x) dx \text{ is to be absolutely convergent.}$$

Here below we clarify some terms used in these conditions:

Monotonic Functions. A function $f(x)$ defined in (a, b) is said to be monotonically increasing if $x_1, x_2 \in (a, b)$, $x_2 > x_1 \Rightarrow f(x_2) \geq f(x_1)$ strictly monotonically increasing if $x_1, x_2 \in (a, b)$, $x_2 > x_1 \Rightarrow f(x_2) > f(x_1)$ monotonically decreasing if $x_1, x_2 \in (a, b)$, $x_2 < x_1 \Rightarrow f(x_2) \leq f(x_1)$ and strictly monotonically decreasing if $x_1, x_2 \in (a, b)$, $x_2 < x_1 \Rightarrow f(x_2) < f(x_1)$.

If $f(x)$ be monotonically increasing in $[a, b]$, then for any point c such that $a < c < b$, $f(c-0)$ and $f(c+0)$ both exist.

Similarly, if $f(x)$ be monotonically decreasing in $[a, b]$ then for $a < c < b$, $f(c-0)$ and $f(c+0)$ both exist.

A function $f(x)$ tends to the limit l as $x \rightarrow c$ i.e., $\lim_{x \rightarrow c} f(x) = l$, if for every $\epsilon > 0$, there exists $\delta > 0$ such that $|f(x) - l| < \epsilon$. In case x is any point of (a, b) , then it satisfies the condition $0 < |x - c| < \delta$.

Right handed and left handed limits.

$$i.e. \quad \lim_{x \rightarrow c+0} f(x) \text{ and } \lim_{x \rightarrow c-0} f(x)$$

When a function $f(x)$ tends to l as x tends to c through values greater than c if for every $\epsilon > 0$ there exists $\delta > 0$ such that

$$|f(x) - l| < \epsilon \text{ and } x \text{ is any point of } (a, b) \text{ satisfying the condition } c < x < c + \delta, \text{ then we say that the right handed limit exists and write } f(c+0) = \lim_{x \rightarrow c+0} f(x) = l.$$

Similarly, when $f(x)$ tends to l as x tends to c through values smaller than c if to each $\epsilon > 0$ there corresponds $\delta > 0$ such that

$$x \in (a, b) \cap (c - \delta, c) \Rightarrow f(x) \in (l - \epsilon, l + \epsilon)$$

then we say that the left handed limit exists and write

$$f(c-0) = \lim_{x \rightarrow c-0} f(x) = l.$$

If $f(c+0)=f(c-0)=l$ i.e., $\lim_{x \rightarrow c+0} f(x) = \lim_{x \rightarrow c-0} f(x) = l$,

then we say that $\lim_{x \rightarrow c} f(x)$ exists at $x=c$.

If c is a limit point of (a, b) then $f(x)$ is said to be **continuous at** c if and only if $\lim_{x \rightarrow c} f(x)$ exists and equals $f(c)$ i.e., if

$$\lim_{x \rightarrow c+0} f(x) = \lim_{x \rightarrow c-0} f(x) = f(c).$$

A function $f(x)$ is said to be **continuous in an interval** (a, b) if it is continuous at every point of the interval.

Discontinuities. A function $f(x)$ is said to be **discontinuous at a point** $x=c$ of its domain if $f(x)$ is not continuous at c . The discontinuity of $f(x)$ at c arises in either of the ways:

- (i) $\lim_{x \rightarrow c} f(x)$ exists but $\lim_{x \rightarrow c} f(x) \neq f(c)$
- (ii) $\lim_{x \rightarrow c} f(x)$ does not exist.

In first case the discontinuity is said to be of the **first kind** or a **simple discontinuity** while in the second case it is of the **second kind**.

Actually we classify the discontinuities of five types as follows:

(1) **Removable discontinuity.** When $f(c+0)$ and $f(c-0)$ both exist and are equal but differ from $f(c)$ i.e., $f(c+0)=f(c-0) \neq f(c)$.

(2) **Discontinuity of first kind or ordinary discontinuity.** When $f(c+0)$ and $f(c-0)$ both exist and are finite but have different values while $f(c)$ may or may not be equal to either of them.

(3) **Discontinuity of the second kind.** When either $f(c+0)$ or $f(c-0)$ or both do not exist.

(4) **Mixed discontinuity.** When only one of $f(c+0)$ and $f(c-0)$ exists and the other does not exist.

(5) **Infinite discontinuity.** When either of $f(c+0)$ and $f(c-0)$ or $f(c)$ or both are infinite e.g. if $f(x) = \frac{1}{x}$

$$f(0+0) = \lim_{h \rightarrow 0} f(0+h) = \lim_{h \rightarrow 0} \frac{1}{h} = \infty$$

$$f(0-0) = \lim_{h \rightarrow 0} f(0-h) = \lim_{h \rightarrow 0} \left(\frac{1}{-h} \right) = -\infty$$

$$\text{and } f(0) = \frac{1}{0} = \infty$$

Hence $f(x)$ has an infinite discontinuity at $x=0$.

Note. We denote by $[a, b]$ closed interval, (a, b) open interval, $[a, b)$ semi-closed i.e., closed on the left and open on the right and similarly $[a, b]$ open on the left and closed on the right.

3. ASSUMPTIONS FOR THE VALIDITY OF FOURIER'S SERIES EXPANSION, WITH ALLIED THEOREMS

The Fourier's series expansion and the determination of Fourier's constants is valid under the following assumptions:

- (i) The expansion of $f(x)$ in a series of sines and cosines of integral multiples of x is possible in the given interval.
- (ii) The given function $f(x)$ is single-valued, continuous and integrable in the given range.

(iii) The series $a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$ is term by term integrable i.e., the series is uniformly convergent in the interval $(-\pi, \pi)$.

(iv) The given function $f(x)$ given in the interval $(-\pi, \pi)$ satisfies Dirichlet's conditions so that the sum on the R.H.S. of (2) of §9.1, has a limit as $n \rightarrow \infty$ and is equal to $f(x)$ at any point x $(-\pi < x < \pi)$, x being continuous and it is equal to $\frac{1}{2} [f(x+0) + f(x-0)]$ when there is an ordinary discontinuity at the point; also it is equal to $\frac{1}{2} [f(-\pi+0) + f(\pi-0)]$ at $x = \pm\pi$ when the limits $f(\pi-0)$ and $f(-\pi+0)$ exist.

THEOREM 1. If a function $f(x)$ is bounded and integrable in $[0, a]$, $a > 0$ and is monotonic in some interval $[0, c]$, c being positive and less than a i.e., $0 < c < a$, then

$$\lim_{n \rightarrow \infty} \int_0^a f(x) \frac{\sin nx}{x} dx = f(0+) \int_0^{\infty} \frac{\sin x}{x} dx \text{ where } f(0+) = f(0+0).$$

Suppose that $f(0+0)=0$ or without affecting the result, $f(0)=0$ and take a positive number $h < c$.

We have by second mean value theorem that if $f(x)$ is continuous in $[a, a+h]$, derivable in $(a, a+h)$ then there exists at least one number θ between 0 and 1 such that

$$f(a+h) - f(a) = hf'(\theta), \quad 0 < \theta < 1 \text{ i.e., } \theta \in (0, 1)$$

Thus there exists $h' \in [0, h]$, such that

$$\begin{aligned} \int_0^h f(x) \frac{\sin nx}{x} dx &= f(0) \int_0^{h'} \frac{\sin nx}{x} dx + f(h) \int_{h'}^h \frac{\sin nx}{x} dx \\ &= f(h) \int_{h'}^h \frac{\sin nx}{x} dx \text{ since } f(0)=0 \\ &= f(h) \int_{nh'}^{nh} \frac{\sin v}{v} dv \text{ on putting } nx=v \\ &= f(h) \int_{nh'}^{nh} \frac{\sin x}{x} dx \text{ (changing the variable } v \text{ to } x) \\ &\dots (1) \end{aligned}$$

Now $\int_0^{\infty} \frac{\sin x}{x} dx$ being convergent, there exists some $k > 0$ such

$$\text{that } \left| \int_0^t \frac{\sin x}{x} dx \right| \leq k, \text{ for every } t \geq 0.$$

$$\text{i.e. } \left| \int_{nh}^{nh'} \frac{\sin x}{x} dx \right| = \left| \int_0^{nh} \frac{\sin x}{x} dx - \int_0^{nh'} \frac{\sin x}{x} dx \right| \leq 2k$$

Also we have $f(h) \rightarrow 0$ as $h \rightarrow (0+0)$

from (1), (2) and (3) we conclude that given an $\epsilon > 0$, there exists a $\delta > 0$ such that

$$\left| \int_0^h f(x) \frac{\sin nx}{x} dx \right| \leq 2k |f(h)| < \frac{\epsilon}{2} \text{ for } 0 < h \leq \delta.$$

$$\text{But } \int_0^a f(x) \frac{\sin nx}{x} dx = \int_0^\delta f(x) \frac{\sin nx}{x} dx + \int_\delta^a f(x) \frac{\sin nx}{x} dx$$

where the second integral on the right tends to 0 as $n \rightarrow \infty$ and there exists a positive integer m such that for every $n \geq m$, we have

$$\left| \int_0^a f(x) \frac{\sin nx}{x} dx \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

$$\text{Hence } \lim_{n \rightarrow \infty} \int_0^a f(x) \frac{\sin nx}{x} dx = f(+0) \int_0^\infty \frac{\sin x}{x} dx$$

In a general case replacing $f(x)$ by $(f(x) - f(0))$ we have

$$\lim_{n \rightarrow \infty} \int_0^a [f(x) - f(+0)] \frac{\sin nx}{x} dx = 0$$

$$\text{But } \int_0^a \frac{\sin nx}{x} dx = \int_0^{na} \frac{\sin y}{y} dy \rightarrow \int_0^\infty \frac{\sin y}{y} dy \text{ as } n \rightarrow \infty$$

$$\text{Hence } \lim_{n \rightarrow \infty} \int_0^a f(x) \frac{\sin nx}{x} dx = f(+0) \int_0^\infty \frac{\sin x}{x} dx.$$

Note. The theorem is also true when $0 < a < \pi$.

THEOREM 2. If $f(x)$ is bounded and integrable in $(-\pi, \pi)$ and monotonic in $[-c, 0)$ and $(0, c)$ where c is some positive number less than π , then

$$\frac{1}{2} a_0 + \sum_{n=1}^{\infty} a_n = \frac{f(+0) + f(-0)}{2}$$

where $f(+0) = f(0+0)$ and $f(-0) = f(0-0)$

We have

$$\begin{aligned} \frac{1}{2} a_0 + \sum_{n=1}^m a_n &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx + \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sum_{n=1}^m \cos nx dx \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \left\{ 1 + 2 \sum_{n=1}^m \cos nx \right\} dx \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \cdot \frac{\sin \left(m + \frac{1}{2}\right) x}{\sin \frac{x}{2}} dx \text{ (by summation)} \end{aligned}$$

$$= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \frac{\sin \left(m + \frac{1}{2}\right) x}{\sin \frac{x}{2}} dx + \frac{1}{2\pi} \int_0^{\pi} f(x) \frac{\sin \left(m + \frac{1}{2}\right) x}{\sin \frac{x}{2}} dx$$

Replacing x by $-x$ in the first integral on the right we have

$$\begin{aligned} \frac{1}{2} a_0 + \sum_{n=1}^m a_n &= \frac{1}{2\pi} \int_0^{\pi} f(-x) \frac{\sin \left(m + \frac{1}{2}\right) x}{\sin \frac{x}{2}} dx + \frac{1}{2\pi} \int_0^{\pi} f(x) \frac{\sin \left(m + \frac{1}{2}\right) x}{\sin \frac{x}{2}} dx \\ &= \frac{1}{2\pi} \cdot 2 \int_0^{\pi/2} \left[f(-2x) \frac{\sin (2m+1)x}{\sin x} + f(2x) \frac{\sin (2m+1)x}{\sin x} \right] dx \text{ (on replacing } x \text{ by } 2x) \\ &= \frac{1}{\pi} [f(-0) + f(+0)] \int_0^\infty \frac{\sin x}{x} dx \text{ as } m \rightarrow \infty \text{ by theorem 1.} \end{aligned}$$

If we take $f(x) = 1$ for every value of x , then we find

$$\frac{1}{2} a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} 1 \cdot dx = 1 \text{ and } a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} \cos nx dx = 0.$$

Hence we get

$$1 = \frac{1+1}{\pi} \int_{-\infty}^{\infty} \frac{\sin x}{x} dx \quad \text{i.e. } \int_0^\infty \frac{\sin x}{x} dx = \frac{\pi}{2}$$

$$\text{So that } \frac{1}{2} a_0 + \sum_{n=1}^{\infty} a_n = \frac{f(+0) + f(-0)}{2}.$$

THEOREM 3. If $f(x)$ is bounded and integrable in $[-\pi, \pi]$ and if it is possible to divide $[-\pi, \pi]$ into finite number of open sub-intervals, in each of which $f(x)$ is monotonic, then

$$\begin{aligned} \frac{1}{2} a_0 + \sum_{n=1}^{\infty} (a_n \cos n\xi + b_n \sin n\xi) &= \frac{1}{2} [f(\xi-0) + f(\xi+0)] \\ &= \frac{1}{2} [f(\pi-0) + f(-\pi+0)] \text{ for } \xi = \pm\pi. \end{aligned}$$

In order to prove it, let us first prove a lemma.

Lemma. If $f(x)$ is bounded and integrable in every interval and is periodic with 2π as its period, then $\int_{-\pi}^{\pi} f(x) dx = \int_{-\pi}^{\pi} f(a+x) dx$, a being any number whatsoever.

Putting $a+x=y$, we have

$$\begin{aligned}\int_{-\pi}^{\pi} f(a+x) dx &= \int_{a-\pi}^{a+\pi} f(y) dy \\ &= \int_{a-\pi}^{-\pi} f(y) dy + \int_{-\pi}^{\pi} f(y) dy + \int_{\pi}^{a+\pi} f(y) dy\end{aligned}$$

Again putting $y = z - 2\pi$, we have

$$\int_{a-\pi}^{-\pi} f(y) dy = \int_{a+\pi}^{\pi} f(z-2\pi) dz = - \int_{\pi}^{a+\pi} f(z) dz = - \int_{\pi}^{a+\pi} f(y) dy$$

$$\therefore \int_{-\pi}^{\pi} f(a+x) dx = \int_{-\pi}^{\pi} f(y) dy \quad \dots (1)$$

Now we come to the main theorem. We have

$$\begin{aligned}\frac{1}{2} a_0 + \sum_{n=1}^m (a_n \cos n\xi + b_n \sin n\xi) &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx + \sum_{n=1}^m \\ &\quad \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \{ \cos nx \cos n\xi + \sin nx \sin n\xi \} dx \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \left[1 + 2 \sum_{n=1}^m \cos n(x-\xi) \right] dx \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x+\xi) \left[1 + 2 \sum_{n=1}^m \cos nx \right] dx \text{ by Lemma} \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x+\xi) \frac{\sin(m+\frac{1}{2})x}{\sin \frac{x}{2}} dx \text{ (by summation)} \\ &= \frac{1}{2\pi} \int_{-\pi}^0 f(x+\xi) \frac{\sin(m+\frac{1}{2})x}{\sin \frac{x}{2}} dx + \frac{1}{2\pi} \int_0^{\pi} f(x+\xi) \frac{\sin(m+\frac{1}{2})x}{\sin \frac{x}{2}} dx \\ &= \frac{1}{2\pi} \int_0^{\pi} f(-x+\xi) \frac{\sin(m+\frac{1}{2})x}{\sin \frac{x}{2}} dx + \frac{1}{2\pi} \int_0^{\pi} f(x+\xi) \frac{\sin(m+\frac{1}{2})x}{\sin \frac{x}{2}} dx \\ &\quad \text{(on replacing } x \text{ by } -x \text{ in the first integral)} \\ &= \frac{1}{\pi} \int_0^{\pi/2} f(-2x+\xi) \frac{\sin(m+\frac{1}{2})x}{\sin \frac{x}{2}} dx + \frac{1}{\pi} \int_0^{\pi/2} f(2x+\xi) \\ &\quad \frac{\sin(m+\frac{1}{2})x}{\sin \frac{x}{2}} dx \quad \text{(on replacing } x \text{ by } 2x)\end{aligned}$$

$$\begin{aligned}\therefore \frac{1}{2} a_0 + \sum_{n=1}^{\infty} (a_n \cos n\xi + b_n \sin n\xi) &= \frac{1}{\pi} \{ f(\xi-0) + f(\xi+0) \} \\ &\quad \int_0^{\infty} \frac{\sin x}{x} dx \text{ as } m \rightarrow \infty \text{ by theorem 1.}\end{aligned}$$

$$\begin{aligned}&= \frac{1}{\pi} \{ f(\xi-0) + f(\xi+0) \} \cdot \frac{\pi}{2} \text{ as in theorem 2.} \\ &= \frac{1}{2} [f(\xi+0) + f(\xi-0)] \text{ when } m \rightarrow \infty.\end{aligned}$$

we can thus restate this results as follows:

If $f(x)$ is bounded and integrable in $[-\pi, \pi]$ and if it is possible to divide $[-\pi, \pi]$ into a finite number of open subintervals in each of which $f(x)$ is monotonic, then the Fourier series corresponding to $f(x)$ converges for every x and if S be the sum function of the series, then

$$S(x) = \frac{1}{2} [f(x+0) + f(x-0)] \text{ for every } x \text{ in } (-\pi, \pi)$$

$$S(x) = \frac{1}{2} [f(\pi-0) + f(-\pi+0)] \text{ for every } x = \pm\pi$$

and

$$S(x+2\pi) = S(x).$$

Note 1. Here the relation $S(x+2\pi) = S(x)$ is very useful in determining the value of the sum function at a point which does not lie in the interval $[-\pi, \pi]$

Note 2. If $x = \xi$ be a point of continuity of $f(x)$, then

$$\begin{aligned}\frac{1}{2} a_0 + \sum_{n=1}^{\infty} (a_n \cos n\xi + b_n \sin n\xi) &= \frac{f(\xi+0) + f(\xi-0)}{2} \\ &= \frac{f(\xi) + f(\xi)}{2} = f(\xi)\end{aligned}$$

Conclusively if $f(x)$ satisfies the conditions of the theorem 3 in the interval $[-\pi, \pi]$, then the sum of the Fourier series corresponding to $f(x)$ is actually $f(x)$ at all such points x of $[-\pi, \pi]$, $f(x)$ being continuous and at points x , the points of discontinuity, the sum of the Fourier series is $\frac{1}{2} [f(x+0) + f(x-0)]$.

Note 3. Half range series.

If $f(x)$ satisfies the conditions of theorem 3 in $[0, \pi]$, then the sum of the sine series $\sum b_n \sin nx$ where $b_n = \frac{2}{\pi} \int_0^{\pi} f(x) \sin nx dx$ is equal to $\frac{1}{2} [f(x+0) + f(x-0)]$ at every point x s.t. $0 < x < \pi$ and is zero when $x=0$ or π .

Define an odd function $F(x)$ in $[-\pi, \pi]$ which is identical with $f(x)$ in $[0, \pi]$.

Thus, $F(x) = f(x)$ in $[0, \pi]$ and $F(x) = -F(-x) = -f(-x)$ in $[-\pi, 0]$ so that $F(x)$ satisfies the condition of theorem 3 in $[-\pi, \pi]$ provided $f(x)$ does so in $[0, \pi]$. Hence the sum of the series $\sum b_n \sin nx$ with

$$\begin{aligned}b_n &= \frac{2}{\pi} \int_0^{\pi} F(x) \sin nx dx = \frac{1}{\pi} [F(x+0) + F(x-0)] \\ &= \frac{1}{\pi} [f(x+0) + f(x-0)] \text{ at every point } x \text{ in } (0, \pi)\end{aligned}$$

and this sum is $\frac{1}{2} [F(+0) + F(-0)] = 0$ for F odd at $x=0$ or π .

Similarly if $f(x)$ satisfies the conditions of theorem 3 in $[0, \pi]$, then the sum of the series $\frac{1}{2} a_0 + \sum a_n \cos nx$ where $a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx$

dx is $\frac{1}{2} [f(x+0) + f(x-0)]$ at every point x in $(0, \pi)$ and is equal to $f(+0)$ for $x=0$ and $f(\pi-0)$ for $x=\pi$.

Note 4. Interval $[0, 2\pi]$.

If $f(x)$ satisfies the conditions of the theorem 3 in $[0, 2\pi]$ then the sum of the series $\frac{1}{2} a_0 + \sum (a_n \cos nx + b_n \sin nx)$ where a_n

$$= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx, \quad b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx,$$

is $\frac{1}{2} [f(x+0) + f(x-0)]$ at every point x in $(0, 2\pi)$ and is $\frac{1}{2} [f(2\pi-0) + f(+0)]$ at $x=0$ or 2π .

Also it is periodic with period 2π .

Writing $x=y+\pi$ so that y varies in $[-\pi, \pi]$ as x varies in $[0, 2\pi]$, let us assume that $f(x)=f(y+\pi)=F(y)$, whence F satisfies conditions of theorem 3 in $[-\pi, \pi]$. As such the sum of the series

$\frac{1}{2} a_0 + \sum (\alpha_n \cos ny + \beta_n \sin ny)$ where

$$\alpha_n = \frac{1}{\pi} \int_{-\pi}^{\pi} F(y) \cos ny \, dy \quad \text{and} \quad \beta_n = \frac{1}{\pi} \int_{-\pi}^{\pi} F(y) \sin ny \, dy,$$

is $\frac{1}{2} [F(y+0) + F(y-0)]$ for $-\pi < y < \pi$ and

is $\frac{1}{2} [F(\pi-0) + F(-\pi+0)]$ at $y=\pm\pi$, being periodic with period 2π .

Changing the variable by the substitution $x=y+\pi$, we have

$$\alpha_n = \frac{1}{\pi} \int_0^{2\pi} F(x-\pi) \cos n(x-\pi) \, dx = \frac{(-1)^n}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx$$

$$\beta_n = \frac{1}{\pi} \int_0^{2\pi} F(x-\pi) \sin n(x-\pi) \, dx = \frac{(-1)^n}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx$$

and $\cos ny = (-1)^n \cos nx$ and $\sin ny = (-1)^n \sin nx$

So that $\frac{1}{2} [F(y+0) + F(y-0)] = \frac{1}{2} [f(y+\pi+0) + f(y+\pi-0)]$

$$= \frac{1}{2} [f(x+0) + f(x-0)]$$

and $\frac{1}{2} [F(\pi-0) + F(-\pi+0)] = \frac{1}{2} [f(2\pi-0) + f(+0)]$

Note 5. Interval $[-l, l]$, l being a real number.

If $f(x)$ satisfies the conditions of theorem 3 in $[-l, l]$, then the sum function of the series $\frac{1}{2} a_0 + \sum (a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l})$ where $a_n = \frac{1}{l}$

$$\int_{-l}^l f(x) \cos \frac{n\pi x}{l} \, dx \quad \text{and} \quad b_n = \frac{1}{l} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} \, dx$$

is $\frac{1}{2} [f(x+0) + f(x-0)]$ for $-l < x < l$

and is $\frac{1}{2} [f(l-0) + f(-l+0)]$ for $x=\pm l$, being periodic with period $2l$.

Putting $y = \frac{\pi x}{l}$ and considering the function $F(y)$ such that

$$f(x) = f\left(\frac{ly}{\pi}\right) = F(y)$$

so that y varies in $[-\pi, \pi]$ as x varies in $[-l, l]$, we can prove the proposition.

Note 6. Interval $[a, b]$ in general.

If $f(x)$ satisfies the condition of theorem 3 in $[a, b]$, then the sum of the series

$$\frac{1}{2} a_0 + \sum \left(a_n \cos \frac{2n\pi x}{b-a} + b_n \sin \frac{2n\pi x}{b-a} \right) \quad \text{where} \quad a_n = \frac{2}{b-a} \int_a^b$$

$$f(x) \cos \frac{2n\pi x}{b-a} \, dx \quad b_n = \frac{2}{b-a} \int_a^b f(x) \sin \frac{2n\pi x}{b-a} \, dx$$

is $\frac{1}{2} [f(x+0) + f(x-0)]$ for $a < x < b$ and is $\frac{1}{2} [f(a+0) + f(b-0)]$ for $x=a$ or b , being periodic with period $b-a$.

The result is obvious by putting $y = \frac{2\pi x}{b-a} + \frac{b+a}{b-a} \pi$ so that y varies in $[-\pi, \pi]$ as x varies in $[a, b]$, (the transformation is obtained from $y=Lx+M$ such that $y=-\pi$ when $x=a$ and $y=\pi$ when $x=b$, by determining the constants L, M).

9.4. COMPLEX REPRESENTATION OF A FOURIER'S SERIES

We have $f(x) = \sum_{n=-\infty}^{\infty} a_n e^{in\pi x}$... (1)

where $a_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-in\pi x} \, dx$ and $a_{-n} = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{in\pi x} \, dx$... (2)

Here a_n and a_{-n} are said to be *conjugate imaginaries*.

In case we consider a function $f(t)$ which is periodic with a period

$T = \frac{2\pi}{\omega}$ then we can write

$$f(t) = \sum_{n=-\infty}^{\infty} a_n e^{in\omega t}, \quad f(t) \text{ being defined in } (-\infty, \infty) \quad \dots (3)$$

Where $\omega = \frac{2\pi}{T}$ and $f(t+T)=f(t)$... (4)

Now R.H.S. of (3) being real, the coefficient of the series on the R.H.S. of (3) must be such that no imaginary terms occur.

Integrating (3) over 0 to T we have

$$\int_0^{2\pi/\omega} f(t) \, dt = \int_0^{2\pi/\omega} \left(\sum_{n=-\infty}^{\infty} a_n e^{in\omega t} \right) dt = \sum_{n=-\infty}^{\infty} a_n$$

$$a_n \int_0^{2\pi/\omega} e^{in\omega t} \, dt, \quad \dots (5)$$

(under the assumption that term by term integration is permissible).

$$=0 \text{ when } n \neq 0 \quad \because \int_0^{2\pi/\omega} \frac{1}{in\omega} \left[e^{in\omega t} \right]_0^{2\pi/\omega} = \frac{1}{in\omega} \frac{1}{(e^{i2n\pi} - 1)} = 0$$

$$=T \text{ when } n=0 \quad \because \int_0^{2\pi/\omega} dt = \frac{2\pi}{\omega} = T$$

$$\therefore (5) \text{ reduces to } \int_0^T f(t) dt = a_0 T \text{ giving } a_0 = \frac{1}{T} \int_0^T f(t) dt = \overline{f(t)} \text{ (say)} \dots (6)$$

where $\overline{f(t)}$ denotes the mean value of $f(t)$.

Now multiplying (3) by $e^{-in\omega t}$ and integrating over 0 to T , we have

$$\int_0^T f(t) e^{-in\omega t} dt = a_n T, \text{ other terms being equal to zero.}$$

This gives
$$a_n = \frac{1}{T} \int_0^T f(t) e^{-in\omega t} dt \dots (7)$$

Replacing n by $-n$ in (7) we get $a_{-n} = \frac{1}{T} \int_0^T f(t) e^{in\omega t} dt \dots (8)$

From (7) and (8) we conclude that $a_{-n} = \overline{a_n}$. $\dots (9)$

In order to find the usual real form of the Fourier series, (3) can be expressed as

$$f(t) = \sum_{n=-\infty}^{-1} a_n e^{in\omega t} + a_0 + \sum_{n=1}^{\infty} a_n e^{in\omega t}$$

$$= \sum_{n=\infty}^1 a_{-n} e^{-in\omega t} + a_0 + \sum_{n=1}^{\infty} a_n e^{in\omega t}$$

(on replacing n by $-n$ in the first term)

$$\text{or } f(t) = a_0 + \sum_{n=1}^{\infty} (a_n e^{in\omega t} + a_{-n} e^{-in\omega t}) dt \dots (10)$$

$$= a_0 + \sum_{n=1}^{\infty} (a_n + a_{-n}) \cos n\omega t + \sum_{n=1}^{\infty} i(a_n - a_{-n}) \sin n\omega t$$

$$\because e^{in\omega t} = \cos n\omega t + i \sin n\omega t \text{ and } e^{-in\omega t} = \cos n\omega t - i \sin n\omega t$$

If we put $a_n + a_{-n} = \alpha_n$, $i(a_n - a_{-n}) = \beta_n$ and $\alpha_0 = 2a_0$ then we find

$$f(t) = \frac{\alpha_0}{2} + \sum_{n=1}^{\infty} \alpha_n \cos n\omega t + \sum_{n=1}^{\infty} \beta_n \sin n\omega t \dots (11)$$

which is the same form as (1) of §9.1 and we can thus determine the coefficients α_n and β_n as

$$\alpha_n = a_n + a_{-n} = \frac{1}{T} \int_0^T f(t) \{e^{-in\omega t} + e^{in\omega t}\} dt$$

$$= \frac{2}{T} \int_0^T f(t) \cos n\omega t dt \dots (12)$$

$$\text{and } \beta_n = i(a_n - a_{-n}) = \frac{1}{T} \int_0^T f(t) i \{e^{-in\omega t} - e^{in\omega t}\} dt$$

$$= \frac{1}{T} \int_0^T f(t) \frac{1}{i} \{e^{in\omega t} - e^{-in\omega t}\} dt \quad \because i^2 = -1$$

$$= \frac{2}{T} \int_0^T f(t) \sin n\omega t dt \dots (13)$$

The introduction of the term $\frac{\alpha_0}{2}$ term in (11) enables (12) to give general term α_n applicable for α_0 as well. In either of the forms real or complex the constant term of Fourier series is always equal to the mean value of the function.

Important Note. The students are advised to commit to memory the following two integrals:

$$(i) \int e^{ax} \sin bx dx = \frac{e^{ax}}{(a^2 + b^2)} \{a \sin bx - b \cos bx\}$$

$$= \frac{e^{ax}}{r} \sin (bx - \alpha), \text{ where } r = \sqrt{a^2 + b^2} \text{ and } \tan \alpha = \frac{b}{a}.$$

$$(ii) \int e^{ax} \cos bx dx = \frac{e^{ax}}{a^2 + b^2} \{a \cos bx + b \sin bx\}$$

$$= \frac{e^{ax}}{r} \cos (bx - \alpha), \text{ where } r = \sqrt{a^2 + b^2}, \tan \alpha = \frac{b}{a}.$$

Problem 1. Obtain Fourier's series for the expansion of $f(x) = x \sin x$ in the interval $-\pi < x < \pi$. Hence deduce that

$$\frac{\pi}{4} = \frac{1}{2} - \frac{1}{1.3} + \frac{1}{3.5} - \frac{1}{5.7} + \dots \quad (\text{Vikram, 1969})$$

Here $x \sin x$ is an even function of x and we have already shown that when $-\pi \leq x \leq \pi$ and $f(x)$ is an even function of x , the Fourier's series is

$$f(x) = \frac{1}{\pi} \int_0^{\pi} f(v) dv + \frac{2}{\pi} \sum_{n=1}^{\infty} \cos nx \int_0^{\pi} f(v) \cos nv dv,$$

where $f(x) = x \sin x$; $\therefore f(v) = v \sin v$.

Thus,

$$x \sin x = \frac{1}{\pi} \int_0^{\pi} v \sin v dv + \frac{2}{\pi} \sum_{n=1}^{\infty} \cos nx \int_0^{\pi} v \sin v \cos nv dv.$$

$$\text{Now } \int_0^{\pi} v \sin v dv = \left\{ -v \cos v \right\}_0^{\pi} + \left\{ \sin v \right\}_0^{\pi} = \pi$$

$$\begin{aligned}
 \text{and } \int_0^{\pi} v \sin v \cos nv \, dv &= \frac{1}{2} \int_0^{\pi} v \{ \sin(n+1)v - \sin(n-1)v \} \, dv \\
 &= \frac{1}{2} \left[\left\{ -v \frac{\cos(n+1)v}{n+1} \right\}_0^{\pi} + \left\{ \frac{\sin(n+1)v}{(n+1)^2} \right\}_0^{\pi} \right. \\
 &\quad \left. - \left\{ -v \frac{\cos(n-1)v}{n-1} \right\}_0^{\pi} - \left\{ \frac{\sin(n-1)v}{(n-1)^2} \right\}_0^{\pi} \right] \\
 &= \frac{1}{2} \left[-\pi \frac{\cos(n+1)\pi}{n+1} + \pi \frac{\cos(n-1)\pi}{n-1} \right] \\
 &= \frac{1}{2} \left[\frac{\pi \cos n\pi}{n+1} - \frac{\pi \cos n\pi}{n-1} \right], \quad n \text{ being an integer.}
 \end{aligned}$$

$$\text{Hence } x \sin x = 1 - \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{\cos n\pi \cos nx}{n^2 - 1}.$$

Corresponding to $n=1$, we have

$$\begin{aligned}
 \int_0^{\pi} v \sin v \cos v \, dv &= \frac{1}{2} \int_0^{\pi} v \sin 2v \, dv \\
 &= \frac{1}{2} \left[\left\{ -\frac{v \cos 2v}{2} \right\}_0^{\pi} + \left\{ \frac{\sin 2v}{4} \right\}_0^{\pi} \right] = -\frac{\pi}{4}.
 \end{aligned}$$

Hence

$$\begin{aligned}
 x \sin x &= 1 - \frac{2}{\pi} \left[-\frac{\pi}{4} \cos x + \sum_{n=1}^{\infty} \frac{\cos n\pi \cos nx}{n^2 - 1} \right] \\
 &= 1 - 2 \left[-\frac{\cos x}{4} + \frac{\cos 2x}{1.3} - \frac{\cos 3x}{2.4} + \frac{\cos 4x}{3.5} - \dots \right].
 \end{aligned}$$

Putting $x=\pi/2$, we get

$$\frac{\pi}{2} = 1 - 2 \left[-\frac{1}{1.3} + \frac{1}{3.5} - \frac{1}{5.7} + \dots \right]$$

or

$$\frac{\pi}{4} = \frac{1}{2} + \frac{1}{1.3} - \frac{1}{3.5} + \frac{1}{5.7} + \dots$$

Problem 2. Find a series of sines and cosines of multiples of x , which will represent $x+x^2$ in the interval $-\pi < x < \pi$. (Agra, 1969)

$$\text{Deduce that } \frac{\pi^2}{6} = \sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots \quad (\text{Agra, 1962})$$

We have

$$\begin{aligned}
 x+x^2 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} (v+v^2) \, dv + \frac{1}{\pi} \sum_{n=1}^{\infty} \cos nx \int_{-\pi}^{\pi} (v+v^2) \cos nv \, dv \\
 &\quad + \frac{1}{\pi} \sum_{n=1}^{\infty} \sin nx \int_{-\pi}^{\pi} (v+v^2) \sin nv \, dv.
 \end{aligned}$$

$$\text{Now } \int_{-\pi}^{\pi} (v+v^2) \, dv = \left[\frac{v^2}{2} + \frac{v^3}{3} \right]_{-\pi}^{\pi} = \frac{2\pi^2}{3}.$$

$$\begin{aligned}
 \int_{-\pi}^{\pi} (v+v^2) \cos nv \, dv &= \left\{ \frac{v \sin nv}{n} \right\}_{-\pi}^{\pi} + \left\{ \frac{\cos nv}{n^2} \right\}_{-\pi}^{\pi} \\
 &\quad + \left\{ \frac{v^2 \sin nv}{n} \right\}_{-\pi}^{\pi} - 2 \int_{-\pi}^{\pi} \frac{v \sin nv}{n} \, dv \\
 &= -2 \left[\left\{ -\frac{\cos nv}{n^2} \right\}_{-\pi}^{\pi} + \left\{ \frac{\sin nv}{n^3} \right\}_{-\pi}^{\pi} \right] \\
 &= \frac{4\pi \cos n\pi}{n^2}
 \end{aligned}$$

$$\begin{aligned}
 \text{and } \int_{-\pi}^{\pi} (v+v^2) \sin nv \, dv &= \left\{ -\frac{v \sin nv}{n} \right\}_{-\pi}^{\pi} + \left\{ \frac{\sin nv}{n^2} \right\}_{-\pi}^{\pi} \\
 &\quad - \left\{ \frac{v^2 \cos nv}{n} \right\}_{-\pi}^{\pi} + 2 \int_{-\pi}^{\pi} v \cos nv \, dv \\
 &= -\frac{2\pi \cos n\pi}{n}.
 \end{aligned}$$

Hence the series is

$$\begin{aligned}
 x+x^2 &= \frac{1}{2\pi} \cdot \frac{2\pi^3}{3} + 4 \sum_{n=1}^{\infty} \frac{\cos nx \cos n\pi}{n^2} - 2 \sum_{n=1}^{\infty} \frac{\sin nx \cos n\pi}{n} \\
 &= \frac{\pi^2}{3} + 4 \left[-\frac{\cos x}{1^2} + \frac{\cos 2x}{2^2} - \frac{\cos 3x}{3^2} + \dots \right] \\
 &\quad - 2 \left[-\frac{\sin x}{1} + \frac{\sin 2x}{2} - \frac{\sin 3x}{3} + \dots \right] \\
 &= \frac{\pi^2}{3} - 4 \left(\cos x - \frac{1}{2^2} \cos 2x + \frac{1}{3^2} \cos 3x + \dots \right) \\
 &\quad + 2 \left(\sin x - \frac{1}{2} \sin 2x + \frac{1}{3} \sin 3x + \dots \right).
 \end{aligned}$$

Second part. We have

$$\begin{aligned}
 x+x^2 &= \frac{\pi^2}{3} - 4 \left[\cos x - \frac{1}{2^2} \cos 2x + \frac{1}{3^2} \cos 3x + \dots \right] \\
 &\quad + 2 \left[\sin x - \frac{1}{2} \sin 2x + \frac{1}{3} \sin 3x + \dots \right].
 \end{aligned}$$

At extremum π and $-\pi$, the sum of series

$$\begin{aligned}
 = f(\pi) &= \frac{1}{2} [f(-\pi=0) + f(+\pi=0)] \\
 &= \frac{1}{2} [-\pi + \pi^2 + \pi + \pi^2] = \pi^2.
 \end{aligned}$$

Putting $x=\pi$ in the above series, we have

$$f(\pi) = \pi^2 = \frac{\pi^2}{3} + 4 \left[1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots \right]$$

$$\text{or } \frac{\pi^2}{6} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots$$

$$\text{or } \frac{\pi^2}{6} = \sum_{n=1}^{\infty} \frac{1}{n^2}.$$

Problem 3. Find the series of sines and cosines of multiples of x which represents $f(x)$ in the interval $-\pi < x < \pi$, where

$$f(x) = 0, \text{ when } -\pi < x \leq 0,$$

$$= \frac{\pi x}{4}, \text{ when } 0 < x < \pi$$

$$\text{and hence deduce } \frac{\pi^2}{8} = 1 + \frac{1}{3^2} + \frac{1}{5^2} + \dots$$

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx),$$

$$\text{where } a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{2\pi} \left[\int_0^{\pi} f(x) dx + \int_{-\pi}^0 f(x) dx \right]$$

$$= \frac{1}{2\pi} \left[\frac{\pi x^2}{8} \right]_0^{\pi} = \frac{\pi^2}{16},$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx \cos nx$$

$$= \frac{1}{\pi} \left[\int_0^{\pi} f(x) \cos nx dx + \int_{-\pi}^0 f(x) \cos nx dx \right]$$

$$= \frac{1}{4n^2} [1 - \cos n\pi] = 0 \text{ for } n \text{ even}$$

$$= \frac{1}{2n^2} \text{ for } n \text{ odd.}$$

$$\text{Similarly } b_n = -\frac{(-1)^n \pi}{4n}.$$

$$\text{So } f(x) = \frac{\pi^2}{16} + \sum_{n=1}^{\infty} \left[\frac{(-1)^{n-1}}{4n^2} \cos nx - \frac{(-1)^n \pi}{4n} \sin nx \right]$$

$$\therefore \frac{\pi x}{4} = \frac{\pi^2}{16} + \left(-\frac{1}{2} \cos x + \frac{\pi}{4} \sin x \right) - \frac{\pi}{4 \cdot 2} \sin 2x$$

$$\left(-\frac{1}{2 \cdot 3^2} \cos 3x + \frac{\pi}{4 \cdot 3} \sin 3x \dots \right)$$

$$\text{Now } f(\pi) = \frac{1}{2} [f(-\pi) + f(\pi)] = \frac{1}{2} \left(0 + \frac{\pi^2}{4} \right) = \frac{\pi^2}{8}.$$

$$\text{So } f(\pi) = \frac{\pi^2}{8} = \frac{\pi^2}{16} + \frac{1}{2} \left(1 + \frac{1}{5} + \frac{1}{5^2} + \dots \right)$$

$$\text{or } \frac{\pi^2}{8} = 1 + \frac{1}{3^2} + \frac{1}{5^2} + \dots$$

Problem 4. Find Fourier's series for $f(x)$ in the interval $(-\pi, \pi)$ where

$$f(x) = \pi + x \text{ when } -\pi < x < 0,$$

$$f(x) = \pi - x \text{ when } 0 < x < \pi.$$

Hint. Here $a_0 = \frac{\pi}{2}$, $a_n = 0$ for n even

$$= \frac{4}{\pi n^2} \text{ for } n \text{ odd,}$$

$$b_n = 0.$$

$$\text{So } f(x) = \frac{\pi}{2} + \frac{4}{\pi} \left(\cos x + \frac{1}{3^2} \cos 3x + \frac{1}{5^2} \cos 5x + \dots \right)$$

Problem 5. Obtain the Fourier series for a function $f(x)$, where

$$f(x) = \cos x \text{ for } 0 \leq x \leq \pi$$

$$f(x) = -\cos x \text{ for } -\pi \leq x \leq 0.$$

$$\text{Let } f(x) = a_0 + \sum (a_n \cos nx + b_n \sin nx).$$

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = 0,$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} -f(x) \cos nx dx = 0,$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx = \frac{4n}{\pi(n^2-1)} [1 + (-1)^n]$$

$$= 0 \text{ for } n \text{ odd}$$

$$= \frac{8n}{\pi(n^2-1)} \text{ for } n \text{ even.}$$

$$\therefore f(x) = \frac{8}{\pi} \left[\frac{2}{1 \cdot 3} \sin 2x + \frac{4}{3 \cdot 5} \sin 4x \dots \right].$$

Problem 6. Find the value of $\sum_{n=1}^{\infty} \frac{1}{n^2}$ using Fourier series.

$$\text{Let } x^2 = \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx),$$

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} x^2 dx = \frac{\pi^2}{3},$$

$$\text{and } a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 \cos nx dx = (-1)^n \frac{4}{n^2}$$

$$b_n = 0.$$

$$\text{Hence } x^2 = \frac{\pi^2}{3} - 4 \left[\frac{1}{1^2} \cos x - \frac{1}{2^2} \cos 2x + \frac{1}{3^2} \cos 3x + \dots \right]$$

$$\therefore 0 = \frac{\pi^2}{3} - 4 \left[1 - \frac{1}{2^2} + \frac{1}{3^2} + \dots \right]$$

$$= \frac{\pi^2}{3} - 4 \left[\left\{ \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots \right\} - \frac{2}{2^2} - \frac{2}{4^2} \dots \right]$$

$$= \frac{\pi^2}{3} - 4 \sum_{n=1}^{\infty} \frac{1}{n^2} + 4 \left(\frac{2}{2^2} \right) \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{3} - 2 \sum_{n=1}^{\infty} \frac{1}{n^2}.$$

Hence
$$2 \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{3}$$

or
$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

Problem 7. Prove that $\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$, using the Fourier's series.

L.H.S. = $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$

$$= \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots + \frac{1}{2^2} + \frac{1}{4^2} + \dots - \frac{1}{2^2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

$$= \sum_{n=1}^{\infty} \frac{1}{n^2} - \frac{1}{4} \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{3}{4} \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{3}{4} \cdot \frac{\pi^2}{6} = \frac{\pi^2}{8}$$

$$\left[\text{since } \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \text{ by Problem 6.} \right]$$

Problem 8. Using Fourier's series prove that

$$\sum_{n=1}^{\infty} \frac{1}{n^4} = \frac{\pi^4}{90}$$

Hint. Take the function $f(x) = x^4 - 2\pi^2 x^2$ and proceed similarly as in Problem 6.

Problem 9. Find a function of x which is equal to kx , when x lies between 0 and $l/2$ and is $k(l-x)$ when x lies between $l/2$ and l .

Given that $f(x) = kx$ when $0 \leq x \leq l/2$,
 $f(x) = k(l-x)$ when $l/2 \leq x \leq l$.

We know that, when $0 \leq y \leq \pi$,

$$f(y) = \frac{1}{\pi} \int_0^{\pi} f(u) du + \frac{2}{\pi} \sum_{n=1}^{\infty} \cos ny \int_0^{\pi} \cos nu f(u) du.$$

Putting $y = \frac{\pi x}{l}$ when $0 \leq x \leq l$, and $u = \frac{\pi v}{l}$, i.e., $du = \frac{\pi}{l} dv$,

we get

$$f\left(\frac{\pi x}{l}\right) = \frac{1}{l} \int_0^l f\left(\frac{\pi v}{l}\right) dv + \frac{2}{l} \sum_{n=1}^{\infty} \cos \frac{n\pi x}{l} \int_0^l f\left(\frac{\pi v}{l}\right) \cos \frac{n\pi v}{l} dv.$$

This may be expressed as

$$\phi(x) = \frac{1}{l} \int_0^l \phi(v) dv + 2 \sum_{n=1}^{\infty} \cos \frac{n\pi x}{l} \int_0^l \pi(v) \cos \frac{n\pi v}{l} dv.$$

Now
$$\int_0^l \phi(v) dv = \int_0^{l/2} \phi(v) dv + \int_{l/2}^l \phi(v) dv$$

$$= \int_0^{l/2} kv dv + \int_{l/2}^l k(l-v) dv$$

$$= \frac{k}{2} \frac{l^2}{2} + k \left[lv - \frac{v^2}{2} \right]_{l/2}^l$$

$$= \frac{kl^2}{8} + kl^2 - \frac{kl^2}{2} - \frac{kl^2}{2} + \frac{kl^2}{8} = \frac{kl^2}{4}$$

and
$$\int_0^l \phi(v) \cos \frac{n\pi v}{l} dv$$

$$= \int_0^{l/2} kv \cos \frac{n\pi v}{l} dv + \int_{l/2}^l k(l-v) \cos \frac{n\pi v}{l} dv$$

$$= \frac{kl}{n\pi} \left\{ v \sin \frac{n\pi v}{l} \right\}_0^{l/2} + \frac{kl^2}{n^2\pi^2} \left\{ \cos \frac{n\pi v}{l} \right\}_0^{l/2} + kl \cdot \frac{1}{n\pi} \left\{ \sin \frac{n\pi v}{l} \right\}_{l/2}^l$$

$$- \frac{kl}{n\pi} \left\{ v \sin \frac{n\pi v}{l} \right\}_{l/2}^l - \frac{kl^2}{n^2\pi^2} \left\{ \cos \frac{n\pi v}{l} \right\}_{l/2}^l$$

$$= \frac{kl^2}{2n\pi} \sin \frac{n\pi}{2} + \frac{kl^2}{n^2\pi^2} \left[\cos \frac{n\pi}{2} - 1 \right] - \frac{kl^2}{n\pi} \sin \frac{n\pi}{2} + \frac{kl^2}{2n\pi} \sin \frac{n\pi}{2}$$

$$- \frac{kl^2}{n^2\pi^2} \left[\cos n\pi - \cos \frac{n\pi}{2} \right]$$

$$= \frac{kl^2}{n^2\pi^2} \left[2 \cos \frac{n\pi}{2} - \cos n\pi - 1 \right] = -\frac{4kl^2}{n^2\pi^2} \text{ for } n=4m+2 \text{ and } 0$$

for other values of n

Hence if n is of the form $4m+2$.

$$\phi(x) = \frac{1}{l} \cdot \frac{kl^2}{4} + \frac{2}{l} \sum_{n=1}^{\infty} \cos \frac{\pi x}{l} \left(-\frac{4kl^2}{n^2\pi^2} \right)$$

$$= \frac{kl}{4} - \frac{8kl}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} \cos \frac{n\pi x}{l}$$

$$= \frac{kl}{4} - \frac{8kl}{\pi^2} \left[\frac{1}{2^2} \cos \frac{2\pi x}{l} + \frac{1}{6^2} \cos \frac{6\pi x}{l} + \frac{1}{10^2} \cos \frac{10\pi x}{l} + \dots \right]$$

when $n=4m+2$, where $m=0, 1, 2, \dots$; $\therefore n=2, 6, 10, \dots$.

Problem 10. If $f(x) = x$ for $0 \leq x \leq \frac{\pi}{2}$.

$$f(x) = \pi - x \text{ for } \frac{\pi}{2} \leq x \leq \pi,$$

Express this function by a sine series and also by a cosine series.

(i) To Express as a Sine Series. We know that

$$f(x) = \frac{2}{\pi} \sum_{n=1}^{\infty} \sin n\pi x \int_0^{\pi} f(v) \sin nv dv$$

Here

$$\int_0^{\pi} f(v) \sin nv dv$$

$$\begin{aligned}
&= \int_0^{\pi/2} v \sin nv \, dv + \int_{\pi/2}^{\pi} (\pi-v) \sin nv \, dv \\
&= \left[-v \frac{\cos nv}{n} + \frac{\sin nv}{n^2} \right]_0^{\pi/2} + \left[-(\pi-v) \frac{\cos nv}{n} - \frac{\sin nv}{n^2} \right]_{\pi/2}^{\pi} \\
&= -\frac{\pi}{2n} \cos \frac{n\pi}{2} + \frac{1}{n^2} \sin \frac{n\pi}{2} + \frac{\pi}{2n} \cos \frac{n\pi}{2} + \frac{1}{n^2} \sin \frac{n\pi}{2} \\
&= \frac{2}{n^2} \sin \frac{n\pi}{2}.
\end{aligned}$$

$$\begin{aligned}
\text{Hence } f(x) &= \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{1}{n^2} \sin \frac{n\pi}{2} \sin nx \\
&= \frac{4}{\pi} \left[\sin x - \frac{1}{3^2} \sin 3x + \frac{1}{5^2} \sin 5x - \dots \right].
\end{aligned}$$

(ii) To Express as a Cosine Series. We know that

$$f(x) = \frac{1}{\pi} \int_0^{\pi} f(v) \, dv + \frac{2}{\pi} \sum_{n=1}^{\infty} \cos nx \int_0^{\pi} f(v) \cos nv \, dv.$$

$$\begin{aligned}
\text{Now } \int_0^{\pi} f(v) \, dv &= \int_0^{\pi/2} v \, dv + \int_{\pi/2}^{\pi} (\pi-v) \, dv \\
&= \left[\frac{v^2}{2} \right]_0^{\pi/2} + \left[\pi v - \frac{v^2}{2} \right]_{\pi/2}^{\pi} = \frac{\pi^2}{8} + \frac{\pi^2}{2} - \frac{3\pi^2}{8} = \frac{\pi^2}{2} \\
\text{and } \int_0^{\pi} \cos nv f(v) \, dv &= \int_0^{\pi/2} v \cos nv \, dv + \int_{\pi/2}^{\pi} (\pi-v) \cos nv \, dv \\
&= \left\{ \frac{v \sin nv}{n} \right\}_0^{\pi/2} + \left\{ \frac{\cos nv}{n^2} \right\}_0^{\pi/2} + \frac{\pi}{n} \left\{ \sin nv \right\}_{\pi/2}^{\pi} \\
&\quad - \left[\frac{v \sin nv}{n} + \frac{\cos nv}{n^2} \right]_{\pi/2}^{\pi} \\
&= \frac{\pi}{2n} \sin \frac{n\pi}{2} + \frac{1}{n^2} \left(\frac{n\pi}{2} - 1 \right) - \frac{\pi}{n} \sin \frac{n\pi}{2} + \frac{\pi}{2n} \sin \frac{n\pi}{2} - \frac{\cos n\pi}{n^2} \\
&\quad + \frac{1}{n^2} \cos \frac{n\pi}{2} \\
&= \frac{1}{n^2} \left\{ 1 + \cos n\pi - 2 \cos \frac{n\pi}{2} \right\} \\
&= -\frac{4}{n^2}, \text{ if } n \text{ is of the form } 4m+2 \text{ and is zero for other}
\end{aligned}$$

values of n .

Hence the cosine series when n is of the form $4m+2$ is

$$\begin{aligned}
f(x) &= \frac{\pi}{4} - \frac{8}{\pi} \sum_{n=1}^{\infty} \frac{\cos nx}{n^2} \\
&= \frac{\pi}{4} - \frac{8}{\pi} \left\{ \frac{1}{2^2} \cos 2x + \frac{1}{6^2} \cos 6x + \dots \right\}
\end{aligned}$$

by putting $n=2, 6, 10, \dots$

Problem 11. Find a series of sines of multiples of x which represents $f(x)$ in the interval $(0, \pi)$. Hence deduce that

$$\frac{\pi^2}{8} = 1 + \frac{1}{3^2} + \frac{1}{5^2} + \dots$$

Draw graph of the function.

$$\text{Let } x = a_0 + \sum_{n=1}^{\infty} a_n \cos nx.$$

$$\begin{aligned}
a_0 &= \frac{1}{\pi} \int_0^{\pi} x \, dx = \frac{\pi}{2} \\
a_n &= \frac{2}{\pi} \int_0^{\pi} x \cos nx \, dx = 0 \text{ for } n \text{ even} \\
&= -4/(n\pi^2) \text{ for } n \text{ odd.}
\end{aligned}
\quad \left. \vphantom{\begin{aligned} a_0 \\ a_n \end{aligned}} \right\} \text{ for cosine series.}$$

Hence the coefficients of cosines of even multiples are zero.

$$\text{Now } x = \frac{\pi}{2} - \frac{4}{\pi} \left[\cos x + \frac{1}{3^2} \cos 3x + \dots \right].$$

Putting $x=0$, we have

$$0 = \frac{\pi}{2} - \frac{4}{\pi} \left[1 + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right]$$

$$\text{or } \frac{\pi^2}{8} = 1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots$$

Graph. In the interval $(0, \pi)$, the line $y=x$ gives the curves represented by the series. Hence $f(x)$ represented by the above series contains cosine terms only. So this is an even function and therefore the curve is symmetrical about the axis of y along which $f(x)$ is plotted. 2π is the period of the series, hence the portion between π to $-\pi$, repeats indefinitely on both the sides and the sum is continuous for all values of x .




Fig. 9.1.

In fact the graph of the sum of n terms of Fourier series for $f(x)$ approximates to the graph of $f(x)$ the greater the value of n

is, the closer is the approximation. Retaining upto three terms of the Fourier series i.e.

$$y = \frac{\pi}{2} - \frac{\pi}{4} \left\{ \frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right\}$$

the graph is as shown below :




Fig. 9.2.

Problem 12. Find the series of sines of multiples of x which represents x in the interval $\pi > x > 0$. Show by a graph the nature of the series.

Here

$$x = \sum_{n=1}^{\infty} b_n \sin nx,$$

where

$$b_n = \frac{2}{\pi} \int_0^{\pi} f(x) \sin nx \, dx \quad [\text{for sine series}]$$

$$= \frac{2}{\pi} \int_0^{\pi} x \sin nx \, dx$$

$$= \frac{2}{\pi} \left[-x \frac{\cos nx}{n} + \frac{\sin nx}{n^2} \right]_0^{\pi}$$

$$= -\frac{2}{\pi} \cdot \pi \frac{1}{n} \cos n\pi = -\frac{2}{n} (-1)^n$$

$\therefore$

$$x = 2 \left[\sin x - \frac{\sin 2x}{2} + \frac{1}{3} \sin 3x + \dots \right]$$

The sum is discontinuous at $x = \pi$.

Graph. The curve is symmetrical about the origin. For unrestricted




Fig. 9.3.

values of x , the series represented between $(-\pi, \pi)$ repeats indefinitely in both the directions. The points $\pm\pi, \pm3\pi, \dots$ are points of discontinuity.

Problem 13. Find the Fourier series for the periodic function $f(x)$ defined by

$$f(x) = -\pi \quad \text{if } -\pi < x < 0$$

$$f(x) = x \quad \text{if } 0 < x < \pi,$$

$$\text{Hence prove that } \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$$

Plot the graph for the series.

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

$$\begin{aligned} \text{Here } a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \, dx \\ &= \frac{1}{2\pi} \left[\int_{-\pi}^0 -\pi \, dx + \int_0^{\pi} x \, dx \right] \\ &= -\frac{\pi}{4}. \end{aligned}$$

$$\begin{aligned} \text{And } a_n &= \frac{1}{2\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\ &= \frac{1}{\pi n^2} [\cos n\pi - 1] = 0 \quad \text{if } n \text{ is even} \end{aligned}$$

$$\text{and } = -\frac{2}{\pi n^2} \quad \text{if } n \text{ is odd.}$$

$$\begin{aligned} b_n &= \frac{1}{2\pi} \left[\int_{-\pi}^0 -\pi \sin nx \, dx + \int_0^{\pi} x \sin nx \, dx \right] \\ &= \frac{1}{n} [1 - 2 \cos n\pi] = -\frac{1}{n} \quad \text{for } n \text{ even} \end{aligned}$$

$$\text{and } = \frac{3}{n} \quad \text{for } n \text{ odd.}$$

$$\begin{aligned} \therefore f(x) &= -\frac{\pi}{4} - \frac{2}{\pi} \cos x - \frac{2}{\pi} \frac{\cos 3x}{3^2} - \frac{2}{\pi} \frac{\cos 5x}{5^2} \dots \\ &\quad + 3 \sin x - \frac{\sin 2x}{2} + \frac{3 \sin 3x}{3} - \frac{\sin 4x}{4} + \dots \end{aligned}$$

One discontinuity occurs at $x = 0$.

$$\text{Hence } f(0) = \frac{f(0^+) + f(0^-)}{2} = -\frac{\pi}{2} \quad \text{since } f(0^+) = 0 \text{ and } f(0^-) = -\pi.$$

Putting $x = 0$ in the series, we have

$$f(0) = -\frac{\pi}{4} - \frac{2}{\pi} \left[\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right] = -\frac{\pi}{2}.$$

$$\text{Hence } \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$$

Graph. Discontinuities occur at $x = \pm\pi$ as follows :




Fig. 9.4.

Problem 14. Find a series of sines and cosines of multiples of x which will represent $\frac{\pi}{2 \sinh \pi} e^x$ in the interval $-\pi < x < \pi$. Find the sum of the series for $x = \pm\pi$.

$$\text{We have } f(x) = \frac{\pi}{2 \sinh \pi} e^x = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx) \quad \dots (1)$$

$$\begin{aligned} \text{Here } a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{\pi}{2 \sinh \pi} e^x dx \\ &= \frac{1}{2\pi} \cdot \frac{\pi}{2 \sinh \pi} [e^x]_{-\pi}^{\pi} \\ &= \frac{1}{4 \sinh \pi} [e^{\pi} - e^{-\pi}] = \frac{1}{4 \sinh \pi} \cdot 2 \sinh \pi = \frac{1}{2} \quad \dots (2) \end{aligned}$$

$$\begin{aligned} a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx, n \geq 1 \\ &= \frac{1}{2 \sinh \pi} \int_{-\pi}^{\pi} e^x \cos nx dx = \frac{1}{2 \sinh \pi} \end{aligned}$$

$$\left[\frac{e^x}{1+n^2} (\cos nx + n \sin nx) \right]_{-\pi}^{\pi} \\ = \frac{(e^{\pi} - e^{-\pi}) \cos n\pi}{(1+n^2) \cdot 2 \sinh \pi} = \frac{2 \sinh \pi \cos nx}{(1+n^2) \cdot 2 \sinh \pi} = \frac{\cos nx}{1+n^2} = \frac{(-1)^2}{1+n^2} \quad \dots (3)$$

$$\begin{aligned} \text{Similarly } b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx = \frac{1}{2 \sinh \pi} \int_{-\pi}^{\pi} e^x \sin nx dx \\ &= \frac{1}{2 \sinh \pi} \left[\frac{e^x}{1+n^2} (\sin nx - n \cos nx) \right]_{-\pi}^{\pi} = \frac{1}{2 \sinh \pi} \cdot \frac{1}{1+n^2} \\ &\quad (-n) (-1)^n (e^{\pi} - e^{-\pi}) \\ &= -(-1)^n \frac{n}{1+n^2} \quad \dots (4) \end{aligned}$$

Substituting the value of a_0, a_n, b_n from (2), (3), (4) in (1) we get

$$\begin{aligned} f(x) &= \frac{\pi e^x}{2 \sinh \pi} = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{(-1)^n}{1+n^2} (\cos nx - n \sin nx) \\ &= \frac{1}{2} + \sum_{n=1}^{\infty} \frac{(-1)^n \cos nx}{1+n^2} + \sum_{n=1}^{\infty} \frac{n(-1)^{n+1}}{1+n^2} \sin nx \\ &= \frac{1}{2} + \left(-\frac{1}{1+1^2} \cos x + \frac{1}{1+2^2} \cos 2x - \frac{1}{1+3^2} \cos 3x + \dots \right) \\ &\quad + \left(\frac{1}{1+1^2} \sin x - \frac{2}{1+2^2} \sin 2x + \frac{3}{1+3^2} \sin 3x - \dots \right) \end{aligned}$$

Now when $x = \pm\pi$, the sum of the series $= \frac{1}{2} [f(-\pi+0) + f(\pi-0)]$

$$\begin{aligned} &= \frac{1}{2} \left[\frac{\pi}{2 \sinh \pi} e^{-\pi} + \frac{\pi}{2 \sinh \pi} e^{\pi} \right] \\ &= \frac{1}{2} \cdot \frac{\pi}{2 \sinh \pi} \cdot (e^{-\pi} + e^{\pi}) \\ &= \frac{1}{2} \cdot \frac{\pi}{2 \sinh \pi} \cdot 2 \cosh \pi = \frac{\pi}{2} \coth \pi \end{aligned}$$

Problem 15. Find a Fourier's series of $f(x)$ in the interval $[-l, l]$ where

$$\begin{aligned} f(x) &= \frac{l}{4} \text{ for } -l \leq x \leq -\frac{l}{2} \\ &= \frac{x^2}{l} \text{ for } -\frac{l}{2} \leq x \leq \frac{l}{2} \\ &= \frac{l}{4} \text{ for } \frac{l}{2} \leq x \leq l \end{aligned}$$

and draw the graph of the function represented by the series.

Since the function $f(x)$ has the same value for positive and negative values of x in $[-l, l]$, it therefore follows that the function $f(x)$ is even and so the Fourier's series for $f(x)$ will be purely a cosine series i.e.

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l}$$

$$\text{where } a_0 = \frac{1}{l} \int_0^l f(x) dx = \frac{1}{l} \int_0^{l/2} \frac{x^2}{l} dx + \frac{1}{l} \int_{l/2}^l \frac{l}{4} dx = \frac{l^3}{24l} + \frac{l}{8} = \frac{l}{6}$$

$$\begin{aligned} \text{and } a_n &= \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx = \frac{2}{l} \int_0^{l/2} \frac{x^2}{l} \cos \frac{n\pi x}{l} dx + \frac{2}{l} \\ &\quad \int_{l/2}^l \frac{l}{4} \cos \frac{n\pi x}{l} dx = \frac{2l}{n^2 \pi^2} \left[\cos \frac{n\pi}{2} - \frac{2}{n\pi} \sin \frac{n\pi}{2} \right] \end{aligned}$$

$$\therefore f(x) = \frac{l}{6} + \frac{2l}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} \left(\cos \frac{n\pi}{2} - \frac{2}{n\pi} \sin \frac{n\pi}{2} \right) \cos \frac{n\pi x}{l}$$

Clearly the shape of the curve between $(-l, -\frac{l}{2})$ and $(\frac{l}{2}, l)$ is given by $y = \frac{l}{4}$ which represents a straight line parallel to the axis of x at a distance $\frac{l}{4}$ from x axis.

Also the shape of the curve between $(-\frac{l}{2}, \frac{l}{2})$ is given by $y = \frac{x^2}{l}$ or $x^2 = ly$ which represents a parabola whose latus rectum is l and the axis of the parabola is y -axis. Hence the shape of curve is as shown in Fig. 9.5.




Fig. 9.5.

Note. If the values of x are not restricted between $(-l, l)$ then the part of the curve from $-l$ to l will be repeated indefinitely in both directions.

Problem 16. If the function defined by $y = x^2$ from 0 to $\frac{\pi}{2}$ and by $y = 0$ from $\frac{\pi}{2}$ to π be represented by a series of sines of multiples of x , show that the coefficient of $\sin nx$ is

$$\left(\frac{4}{\pi n^3} - \frac{\pi}{2n} \right) \cos \frac{n\pi}{2} + \frac{2}{n^3} \sin \frac{n\pi}{2} - \frac{4}{\pi n^3}.$$

To what value does the series converge at the point $x = \frac{\pi}{2}$?

Sketch the graph of the function represented by the series for values of x not restricted to lie between 0 and π ; and also indicate the graph of the cosine series which represents the same function in the interval 0 to π .

$$\begin{aligned} \text{We have } f(x) &= x^2 \text{ for } 0 < x < \frac{\pi}{2} \\ &= 0 \text{ for } \frac{\pi}{2} < x < \pi. \end{aligned}$$

For a purely sine series in $(0, \pi)$, the function must be odd in $(-\pi, \pi)$ so that $f(x)$ in $(-\pi, 0)$ is defined such that $f(-x) = -f(x)$ in the form

$$\begin{aligned} y &= -x^2 \text{ for } -\frac{\pi}{2} < x < 0 \\ &= 0 \text{ for } -\pi < x < -\frac{\pi}{2} \end{aligned}$$

and then $f(x) = \sum_{n=1}^{\infty} b_n \sin nx$, where

$$\begin{aligned} b_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \sin nx \, dx = \frac{2}{\pi} \int_0^{\pi/2} x^2 \sin nx \, dx \\ &= \frac{2}{\pi} \left[\left(-\frac{\cos nx}{n} \cdot x^2 \right)_0^{\pi/2} + \int_0^{\pi/2} \frac{\cos nx}{n} \cdot 2x \, dx \right] \\ &= \frac{2}{\pi} \left[-\frac{\pi^2}{4n} \cos \frac{n\pi}{2} + \left(\frac{2x \sin nx}{n^2} \right)_0^{\pi/2} - \frac{2}{n^2} \int_0^{\pi/2} \sin nx \, dx \right] \\ &= \frac{2}{\pi} \left[-\frac{\pi^2}{4n} \cos \frac{n\pi}{2} + \frac{\pi \sin n\pi/2}{n^2} + \frac{2}{n^2} \left(\frac{\cos nx}{n} \right)_0^{\pi/2} \right] \\ &= \frac{2}{\pi} \left[-\frac{\pi^2}{4n} \cos \frac{n\pi}{2} + \frac{\pi}{n^2} \sin \frac{n\pi}{2} + \frac{2}{n^2} \cos \frac{n\pi}{2} - \frac{2}{n^3} \right] \\ &= \left(\frac{4}{\pi n^3} - \frac{\pi}{2n} \right) \cos \frac{n\pi}{2} + \frac{2}{n^2} \sin \frac{n\pi}{2} - \frac{4}{\pi n^3} \end{aligned}$$

which is the required coefficient of $\sin nx$.

Had we supposed $y = x^2$ in $(-\frac{\pi}{2}, 0)$ and $y = 0$ in $(-\pi, -\frac{\pi}{2})$, the form of the Fourier's series would have been purely a cosine series.

$$\begin{aligned} \text{Now, the sum of the series at } x = \frac{\pi}{2} &= \frac{1}{2} [f(x+0) + f(x-0)] \\ &= \frac{1}{2} \left[f\left(\frac{\pi}{2} + 0\right) + f\left(\frac{\pi}{2} - 0\right) \right] \\ &= \frac{1}{2} \left[0 + \frac{\pi^2}{4} \right] = \frac{\pi^2}{8}. \end{aligned}$$




Fig. 9.6

The shape of the curve between $(-\pi, -\frac{\pi}{2})$ and $(\frac{\pi}{2}, \pi)$ is given by $y = 0$ i.e., x -axis.

The shape of the curve between $(0, \frac{\pi}{2})$ is given by $y = x^2$ which

is a parabola whose latus rectum is unity and axis is the y -axis. Also the shape of the curve between $(-\frac{\pi}{2}, 0)$ is given by $y = -x^2$ which is a similar parabola as in $(0, \frac{\pi}{2})$ except that it has its concavity towards negative y -axis. The shape is as shown in Fig. 9.6.

The period of Fourier's series for $f(x)$ being 2π , the part of the curve lying between $(-\pi, \pi)$ is repeated indefinitely in both the directions.

Again for a purely cosine series in $(-\pi, 0)$ the function $f(x)$ must be such that $f(-x) = f(x)$ i.e., $f(x)$ in $(-\pi, 0)$ is defined such that,

$$y = x^2 \text{ for } -\frac{\pi}{2} < x < 0$$

$$= 0 \text{ for } -\pi < x < -\frac{\pi}{2}$$




Fig. 9.7

As such the only difference here will be between $(-\frac{\pi}{2}, 0)$ where the shape of the curve will be a parabola given by $y = x^2$ with its concavity towards the positive end of y -axis. The shape is as shown in Fig. 9.7.

9.5. PARSEVAL'S IDENTITY FOR FOURIER SERIES

If the Fourier series for $f(x)$ converges uniformly in $(-l, l)$, then

$$\frac{1}{l} \int_{-l}^l \{f(x)\}^2 dx = \frac{a_0^2}{2} + \sum (a_n^2 + b_n^2)$$

where a_0, a_n, b_n are Fourier's constants.

$$\text{We have } f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l} \right) \quad \dots (1)$$

Multiplying both sides of (1) by $f(x)$ and integrating term by term from $-l$ to l , we get

$$\begin{aligned} \int_{-l}^l \{f(x)\}^2 dx &= \frac{a_0}{2} \int_{-l}^l f(x) dx + \sum_{n=1}^{\infty} \left\{ a_n \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx \right. \\ &\quad \left. + b_n \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx \right\} \\ &= \frac{a_0^2}{2} l + l \sum_{n=1}^{\infty} (a_n^2 + b_n^2), \text{ by using Fourier coefficient so that} \\ &\quad \int_{-l}^l f(x) dx = l a_0, \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx = l a_n \text{ and} \\ &\quad \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx = l b_n. \\ \therefore \frac{1}{l} \int_{-l}^l \{f(x)\}^2 dx &= \frac{a_0^2}{2} + \sum (a_n^2 + b_n^2) \quad \dots (2) \end{aligned}$$

COROLLARY. Riemann's theorem. An important consequence of Parseval's identity is Riemann's theorem i.e.,

$$\lim_{n \rightarrow \infty} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx = 0, \quad \lim_{n \rightarrow \infty} \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx = 0 \quad \dots (3)$$

9.6. FOURIER'S INTEGRAL

If $f(x)$ satisfies the Dirichlet's conditions in $-\pi \leq x \leq \pi$, and $\int_{-\infty}^{\infty} f(x) dx$ converges i.e., is integrable in $-\infty < x < \infty$, then we have the Fourier series expansion for $f(x)$ as

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx, \quad \dots (1)$$

$$\text{where } \left. \begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx, \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx, \\ \text{and } b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx. \end{aligned} \right\} \quad \dots (2)$$

This function of x may be developed into a trigonometric series for all values of x between $x = -c$ and $x = c$ by putting

$$z = \frac{\pi}{c} x, \text{ where } z = -\pi \text{ when } x = -c$$

$$\text{and } z = \pi \text{ when } x = c,$$

$$\text{i.e., } f(x) = f\left(\frac{c}{\pi} z\right)$$

Then the series (1) may be developed in terms of z as

$$\begin{aligned} f\left(\frac{c}{\pi} z\right) &= a_0 + a_1 \cos z + a_2 \cos 2z + a_3 \cos 3z + \dots \\ &\quad + b_1 \sin z + b_2 \sin 2z + b_3 \sin 3z + \dots \quad \dots (3) \end{aligned}$$

$$\left. \begin{aligned} \text{where } a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f\left(\frac{c}{\pi} z\right) dz, \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f\left(\frac{c}{\pi} z\right) \cos nz \, dz, \\ \text{and } b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f\left(\frac{c}{\pi} z\right) \sin nz \, dz. \end{aligned} \right\} \dots (4)$$

Now if we replace z by $\frac{\pi}{c}x$, then (3) becomes

$$f(x) = a_0 + a_1 \cos \frac{\pi x}{c} + a_2 \cos \frac{2\pi x}{c} + a_3 \cos \frac{3\pi x}{c} + \dots \\ + b_1 \sin \frac{\pi x}{c} + b_2 \sin \frac{2\pi x}{c} + b_3 \sin \frac{3\pi x}{c} + \dots (5)$$

Its coefficients being the same as those of (3) is therefore valid from $x=c$ to $x=-c$, where

$$a_0 = \frac{1}{2\pi} \int_{-c}^c f(x) \cdot \frac{\pi}{c} dx = \frac{1}{2c} \int_{-c}^c f(x) dx$$

$$\therefore \text{ when } z = \frac{\pi}{c}x, dz = \frac{\pi}{c} dx$$

$$\left. \begin{aligned} a_0 &= \frac{1}{2c} \int_{-c}^c f(t) dt \text{ (say)} \\ a_n &= \frac{1}{\pi} \int_{-c}^c f(x) \cos \frac{n\pi x}{c} \cdot \frac{\pi}{c} dx \\ &= \frac{1}{c} \int_{-c}^c f(t) \cos \frac{n\pi t}{c} \cdot dt \\ \text{and } b_n &= \frac{1}{\pi} \int_{-c}^c f(x) \sin \frac{n\pi x}{c} \cdot \frac{\pi}{c} dx \\ &= \frac{1}{c} \int_{-c}^c f(t) \sin \frac{n\pi t}{c} dt. \end{aligned} \right\} \dots (6)$$

Now if we substitute the values of the coefficients $a_0, a_1, a_2, a_3, \dots, b_1, b_2, b_3, \dots$ given by (6) in (5), then we get

$$f(x) = \frac{1}{2c} \int_{-c}^c f(t) dt + \frac{1}{c} \int_{-c}^c f(t) \cos \frac{\pi x}{c} dt \\ + \frac{1}{c} \int_{-c}^c f(t) \cos \frac{2\pi t}{c} \cos \frac{2\pi x}{c} dx + \dots \\ + \frac{1}{c} \int_{-c}^c f(t) \sin \frac{\pi t}{c} \sin \frac{\pi x}{c} dt \\ + \frac{1}{c} \int_{-c}^c f(t) \sin \frac{2\pi t}{c} \sin \frac{2\pi x}{c} dt + \dots \\ = \frac{1}{c} \int_{-c}^c f(t) \left[\frac{1}{2} + \cos \frac{\pi t}{c} \cos \frac{\pi x}{c} + \cos \frac{2\pi t}{c} \cos \frac{2\pi x}{c} + \dots \right. \\ \left. + \sin \frac{\pi t}{c} \sin \frac{\pi x}{c} + \sin \frac{2\pi t}{c} \sin \frac{2\pi x}{c} + \dots \right] dt$$

$$= \frac{1}{c} \int_{-c}^c f(t) \left[\frac{1}{2} + \left\{ \cos \frac{\pi t}{c} \cos \frac{\pi x}{c} + \sin \frac{\pi t}{c} \sin \frac{\pi x}{c} \right\} \right. \\ \left. + \left\{ \cos \frac{2\pi t}{c} \cos \frac{2\pi x}{c} + \sin \frac{2\pi t}{c} \sin \frac{2\pi x}{c} \right\} + \dots \right] dt \\ = \frac{1}{c} \int_{-c}^c f(t) \left[\frac{1}{2} + \cos \frac{\pi}{c}(x-t) + \cos \frac{2\pi}{c}(x-t) + \dots \right] dt \\ = \frac{1}{2c} \int_{-c}^c f(x) \left[1 + 2 \cos \frac{\pi}{c}(x-t) + 2 \cos \frac{2\pi}{c}(x-t) + \dots \right] dt \\ = \frac{1}{2\pi} \int_{-c}^c f(t) \left[\frac{\pi}{c} \cos \frac{0\pi}{c}(x-t) + \frac{\pi}{c} \cos \frac{\pi}{c}(x-t) \right. \\ \left. + \frac{\pi}{c} \cos \left(\frac{-\pi}{c} \right)(x-t) + \frac{\pi}{c} \cos \frac{2\pi}{c}(x-t) \right. \\ \left. + \frac{\pi}{c} \cos \left(\frac{-2\pi}{c} \right)(x-t) + \dots \right] dt$$

$$\text{since } 2 \cos \phi = \cos \phi + \cos (-\phi)$$

$$= \frac{1}{2\pi} \int_{-c}^c f(t) \left[\dots \frac{\pi}{c} \cos \left(-\frac{n\pi}{c} \right)(x-t) + \dots \right. \\ \left. + \frac{\pi}{c} \cos \left(-\frac{2\pi}{c} \right)(x-t) + \frac{\pi}{c} \cos \left(-\frac{\pi}{c} \right)(x-t) \right. \\ \left. + \frac{\pi}{c} \cos \frac{0\pi}{c}(x-t) + \frac{\pi}{c} \cos \frac{\pi}{c}(x-t) + \frac{\pi}{c} \cos \frac{2\pi}{c}(x-t) \right. \\ \left. + \dots + \frac{\pi}{c} \cos \frac{n\pi}{c}(x-t) + \dots \right] dt \\ = \frac{1}{2\pi} \int_{-c}^c f(t) \left[\lim_{n \rightarrow \infty} \sum_{r=-n}^{r=n} \frac{\pi}{c} \cos \frac{r\pi}{c}(x-t) \right] dt \\ = \frac{1}{2\pi} \int_{-c}^c f(t) \left[\lim_{n \rightarrow \infty} \sum_{r=-n}^n \frac{1}{c/\pi} \cos \frac{r}{c/\pi}(x-t) \right] dt.$$

If c becomes indefinitely large i.e. as $c \rightarrow \infty$, $\frac{c}{\pi} \rightarrow \infty$, we have

$$\lim_{c \rightarrow \infty} \sum_{r=-\infty}^{\infty} \frac{1}{c/\pi} \cos \frac{r}{c/\pi}(x-t) = \int_{-\infty}^{\infty} \cos u(x-t) du$$

(by the definition of integral as the limit of a sum).

$$\text{Hence } f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(t) dt \int_{-\infty}^{\infty} \cos u(x-t) du. \dots (7)$$

This double integral is known as *Fourier's Integral* and holds if x is a point of continuity of $f(x)$.

$$\text{Aliter. We have } f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l} \right) \dots (8)$$

where $a_n = \frac{1}{l} \int_{-l}^l f(u) \cos \frac{n\pi u}{l} du$ and $b_n = \frac{1}{l} \int_{-l}^l f(u) \sin \frac{n\pi u}{l} du$... (9)

so that

$$a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l} = \frac{1}{\pi} \int_{-l}^l f(u) \cos \frac{n\pi}{l} (u-x) dx$$

$$\text{and } \frac{a_0}{2} = \frac{1}{2l} \int_{-l}^l f(u) du$$

$\therefore$ (8) gives

$$f(x) = \frac{1}{2l} \int_{-l}^l f(u) du + \frac{1}{l} \sum_{n=1}^{\infty} \int_{-l}^l f(u) \cos \frac{n\pi}{l} (u-x) du \dots (10)$$

Assuming that $\int_{-\infty}^{\infty} |f(u)| du$ converges, the first term on R.H.S. of (10) approaches zero as $l \rightarrow \infty$ and hence (10) yields

$$f(x) = \lim_{l \rightarrow \infty} \frac{1}{l} \sum_{n=1}^{\infty} \int_{-\infty}^{\infty} f(u) \cos \frac{n\pi}{l} (u-x) dx \dots (11)$$

Putting $\frac{\pi}{l} = \Delta t$, (11) can be written as

$$f(x) = \lim_{\Delta t \rightarrow 0} \sum_{n=1}^{\infty} \Delta t F(n \Delta t) \dots (12)$$

$$\text{where } F(t) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(x) \cos t(u-x) du \dots (13)$$

Thus (12) gives

$$f(x) = \int_0^{\infty} F(t) dt = \frac{1}{\pi} \int_0^{\infty} dt \int_{-\infty}^{\infty} f(u) \cos t(u-x) du \dots (14)$$

which is Fourier's Integral formula.

Note. The complex form of Fourier's integral is

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{itx} dt \int_{-\infty}^{\infty} f(u) e^{-itu} du$$

$$= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(u) e^{it(u-x)} du dt \dots (15)$$

9.7. DIFFERENT FORMS OF FOURIER'S INTEGRALS

Fourier's Integral is

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(t) dt \int_{-\infty}^{\infty} \cos u(x-t) du \dots (1)$$

$$\text{Here } \int_{-\infty}^{\infty} \cos u(x-t) du = \int_{-\infty}^0 \cos u(x-t) dt + \int_0^{\infty} \cos u(x-t) dt$$

Say $I = I_1 + I_2$

Replacing u by $-u$ in I_1 , we get

$$I_1 = - \int_{\infty}^0 \cos u(x-t) du = \int_0^{\infty} \cos u(x-t) du = I_2.$$

$$\text{Hence } I = \int_{-\infty}^{\infty} \cos u(x-t) du = 2 \int_0^{\infty} \cos u(x-t) du.$$

Therefore (1) becomes

$$f(x) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(t) dt \int_0^{\infty} \cos u(x-t) du \dots (2)$$

Now as the limits of integration in (2) do not involve the variables u or t , the order of integration may be changed, i.e.

$$f(x) = \frac{1}{\pi} \int_0^{\infty} du \int_{-\infty}^{\infty} f(t) \cos u(x-t) dt \dots (3)$$

(Agra, 1964)

If $f(x)$ be an odd function of x , i.e. $f(-x) = -f(x)$, then

$$\int_{-\infty}^{\infty} f(t) \cos u(x-t) dt = \int_{-\infty}^0 f(t) \cos u(x-t) dt + \int_0^{\infty} f(t) \cos u(x-t) dt.$$

Replacing t by $-t$, we have

$$\int_{-\infty}^0 f(t) \cos u(x-t) dt = - \int_0^{\infty} f(-t) \cos u(x+t) dt$$

$$= - \int_0^{\infty} f(t) \cos u(x+t) dt$$

$$\therefore f(-t) = -f(t).$$

$$\text{Thus } \int_{-\infty}^{\infty} f(t) \cos u(x-t) dt$$

$$= - \int_0^{\infty} f(t) \cos u(x+t) dt + \int_0^{\infty} f(t) \cos u(x-t) dt$$

$$= \int_0^{\infty} f(t) [\cos u(x-t) - \cos u(x+t)] dt$$

$$= \int_0^{\infty} f(t) \sin ux \sin ut. dt$$

Substituting it in (3), we get

$$f(x) = \frac{2}{\pi} \int_0^{\infty} du \int_0^{\infty} f(t) \sin ut \sin ux dt \dots (4)$$

Changing the order of integration this may be written as

$$f(x) = \frac{2}{\pi} \int_0^{\infty} f(t) dt \int_0^{\infty} \sin ut \sin ux du \dots (5)$$

Again if $f(x)$ be an even function of (x) , i.e., $f(-x) = f(x)$, we have

$$\int_{-\infty}^{\infty} f(t) \cos u(x-t) dt = \int_{-\infty}^0 f(t) \cos u(x-t) dt + \int_0^{\infty} f(t) \cos u(x-t) dt$$

Repeating t by $-t$ in the second integral, we get

$$\begin{aligned} \int_{-\infty}^{\infty} f(t) \cos u(x-t) dt &= - \int_{\infty}^0 f(-t) \cos u(x+t) dt + \int_0^{\infty} f(t) \cos u(x-t) dt \\ &= \int_0^{\infty} f(t) \cos u(x+t) dt + \int_0^{\infty} f(t) \cos u(x-t) dt \\ &= \int_0^{\infty} f(t) [\cos u(x+t) + \cos u(x-t)] dt \\ &= 2 \int_0^{\infty} f(t) \cos ut \cos ux dt \end{aligned}$$

Its substitution in (3) yields,

$$\begin{aligned} f(x) &= \frac{2}{\pi} \int_0^{\infty} du \int_0^{\infty} f(t) \cos ut \cos ux dt \quad \dots (6) \\ &= \frac{2}{\pi} \int_0^{\infty} f(t) dt \int_0^{\infty} \cos ut \cos ux du \quad (\text{On changing the order of integration}) \end{aligned}$$

Problem 17. Show that the sum function of the Integral formula is $\frac{1}{2} [f(x+0) + f(x-0)]$ corresponding to the function $f(x)$ in the interval $0 < x < l$.

By Weierstrass test, since $\left| \int_{-\infty}^{\infty} f(u) \cos t(x-u) du \right| \leq \int_{-\infty}^{\infty} |f(u)| du$ (which converges), the integral $\int_{-\infty}^{\infty} f(u) \cos t(x-u) du$ converges absolutely and uniformly for all values of t . Thus by reversing the order of integration, we have,

$$\begin{aligned} \frac{1}{\pi} \int_{t=0}^l dt \int_{u=-\infty}^{\infty} f(u) \cos t(x-u) du &= \frac{1}{\pi} \int_{u=-\infty}^{\infty} f(u) du \int_{t=0}^l \cos t(x-u) dt \\ &= \frac{1}{\pi} \int_{u=-\infty}^{\infty} f(u) \frac{\sin l(u-x)}{u-x} du \\ &= \frac{1}{\pi} \int_{p=-\infty}^{\infty} f(x+p) \frac{\sin lp}{p} dp \quad \text{when } u=x+p \\ &= \frac{1}{\pi} \int_{-\infty}^0 f(x+p) \frac{\sin lp}{p} dp + \frac{1}{\pi} \int_0^{\infty} f(x+p) \frac{\sin lp}{p} dp \\ &= \frac{1}{2} [f(x+0) + f(x-0)] \quad \text{when } l \rightarrow \infty \text{ by theorem 3 of § 9.3.} \end{aligned}$$

Problem 18. Show that $\int_0^{\infty} \frac{\cos ux}{u^2+1} du = \frac{\pi}{2} e^{-x}$, $x \geq 0$.

Putting $f(x) = e^{-x}$ in the Fourier's Integral form (6) of § 9.7, we have

$$\begin{aligned} (x) = e^{-x} &= \frac{2}{\pi} \int_0^{\infty} du \int_0^{\infty} e^{-t} \cos ut \cos ux dt = \frac{2}{\pi} \int_0^{\infty} \cos ux du \int_0^{\infty} e^{-t} \cos ut dt \\ &= \frac{2}{\pi} \int_0^{\infty} \cos ux du \left[\frac{e^{-t}}{1+u^2} (-\cos ut + u \sin ut) \right]_0^{\infty} \\ &= \frac{2}{\pi} \int_0^{\infty} \frac{\cos ux}{1+u^2} du \\ \therefore \int_0^{\infty} \frac{\cos ux}{1+u^2} du &= \frac{\pi}{2} e^{-x}. \end{aligned}$$

Note. If $x=0$, this result reduces to $\int_0^{\infty} \frac{du}{1+u^2} = \frac{\pi}{2}$.

9.8. A REMARK ON CONVERGENCE OF FOURIER'S SERIES

The Fourier constants in real form by (12) and (13) of § 9.4 are

$$\alpha_n = \frac{2}{T} \int_0^T f(t) \cos n\omega t dt \quad \text{and} \quad \beta_n = \frac{2}{T} \int_0^T f(t) \sin n\omega t dt \quad \dots (1)$$

It is plausible from (1) that α_n and β_n must diminish indefinitely as n increases since the more rapid is the fluctuation in sign of $\cos n\omega t$ and $\sin n\omega t$, the more complete is the canceling of the various elements of integrals (1) which are known as *Riemann-Lebesgue theorem*. Stokes has formulated rather definite results, according to which if $f(t)$ satisfies Dirichlet's conditions under the careful vigilance whether in particular cases discontinuities of $f(t)$ or its derivatives are introduced at the terminal points of the various segments, then

(i) if $f(t)$ has a finite number of isolated discontinuities in a period the coefficient converge to zero ultimately, like the sequence

$$\left\{ \frac{1}{m} \right\} = 1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, \dots \quad \dots (2)$$

(ii) if $f(t)$ is everywhere continuous while its first derivative $f'(t)$ possesses a finite number of isolated discontinuities then the coefficients converge like the sequence

$$\left\{ \frac{1}{m^2} \right\} = 1, \frac{1}{2^2}, \frac{1}{3^2}, \dots, \frac{1}{n^2}, \dots \quad \dots (3)$$

(iii) if $f(t)$ and $f'(t)$ are continuous but the second derivative $f''(t)$ is discontinuous at isolated points, the coefficient converge like the sequence

$$\left\{ \frac{1}{n^3} \right\} = 1, \frac{1}{2^3}, \frac{1}{3^3}, \dots, \frac{1}{n^3}, \dots \quad \dots (4)$$

(iv) if in general, $f(t)$ and its derivatives upto $(n-1)$ order are continuous but n th derivative in a period has a finite number of isolated discontinuities, the coefficients converge like the sequence,

$$\left\{ \frac{1}{n^{n+1}} \right\} = 1, \frac{1}{2^{n+1}}, \frac{1}{3^{n+1}}, \dots, \frac{1}{n^{n+1}} \dots (5)$$

The above statements can be demonstrated as follows:

Integrating by parts, we have from (1),

$$\beta_n = \frac{2}{T} \int_0^T f(t) \sin n\omega t dt = -\frac{1}{n} \left[\frac{f(t) \cos n\omega t}{\pi} \right]_0^T + \frac{1}{\pi n} \int_0^T f'(t) \cos n\omega t dt \quad \because T = \frac{2\pi}{\omega} \dots (6)$$

The first term on R.H.S. of (6) vanishes in either case (i) there are no points of discontinuity of $f(t)$ in the range from $t=0$ to $t=T$, (ii) there is no discontinuity of $f(t)$ at $t=0$ or T . In case there is a discontinuity,

there exist an upper limit M (say) to the coefficient of $\frac{1}{n}$ for all n .

The second term on R.H.S. of (6) vanishes as $n \rightarrow \infty$ due to fluctuations of $\cos n\omega t$. Hence β_n is comparable with M/n .

When there is no discontinuity in $f(t)$, we have

$$\beta_n = \frac{1}{\pi n} \int_0^T f'(t) \cos n\omega t dt \dots (7)$$

Integrating again by parts, we find,

$$\beta_n = \frac{1}{n^2} \left[\frac{f'(t) \sin n\omega t}{\pi \omega} \right]_0^T - \frac{1}{\pi n^2 \omega} \int_0^T f''(t) \sin n\omega t dt \dots (8)$$

If $f'(t)$ has discontinuities, take M as the upper limit of the coefficient of $\frac{1}{n^2}$ as discussed above, then β_n is comparable with M/n^2 , since second term on (8) vanishes due to fluctuations of $\sin n\omega t$.

As regards the convergence of a Fourier series, the above statements are very useful and so one should see initially how well or how poorly the series will converge.

Differentiation of a Fourier series makes the convergence poorer while Integration increases its rate of convergence. Conclusively if a Fourier series has been differentiated until it converges as $\frac{1}{n}$, it is no further differentiable.

9.9. PHYSICAL APPLICATIONS OF FOURIER'S SERIES

[1] Fourier Series Involving Phase Angles

By (11) of § 9.4 we have

$$f(t) = \frac{\alpha_0}{2} + \sum_{n=1}^{\infty} \alpha_n \cos n\omega t + \sum_{n=1}^{\infty} \beta_n \sin n\omega t \dots (1)$$

where $\alpha_n = \frac{2}{T} \int_0^T f(t) \cos n\omega t dt$, $\beta_n = \frac{2}{T} \int_0^T f(t) \sin n\omega t dt$ and $T = \frac{2\pi}{\omega}$ (period) $\dots (2)$

Let $\alpha_n \cos n\omega t + \beta_n \sin n\omega t = \gamma_n \cos (n\omega t - \phi_n)$, ϕ_n being the phase angles.

$$= \gamma_n \cos n\omega t \cos \phi_n + \gamma_n \sin n\omega t \sin \phi_n \dots (3)$$

Equating coefficients of $\cos n\omega t$ and $\sin n\omega t$ on either side of (3), we get

$$\alpha_n = \gamma_n \cos \phi_n \text{ and } \beta_n = \gamma_n \sin \phi_n \dots (4)$$

which give $\gamma_n = \sqrt{\alpha_n^2 + \beta_n^2}$ and $\phi_n = \tan^{-1} \frac{\beta_n}{\alpha_n} \dots (5)$

Hence the series (1) takes the form

$$f(t) = \frac{\alpha_0}{2} + \sum_{n=1}^{\infty} \gamma_n \cos (n\omega t - \phi_n) \dots (6)$$

or $f(t) = \frac{\alpha_0}{2} + \sum_{n=1}^{\infty} \gamma_n \sin \left(n\omega t + \frac{\pi}{2} - \phi_n \right) \dots (7)$

[2] Effective Values and the Average of a Product.

When dealing with the problems in electrical-circuit theory and in the theory of mechanical vibrations, we require to find the root mean square or effective value of a periodic function. In terms of complex Fourier-series expansion, a periodic function $f(t)$ is given by

$$f(t) = \sum_{n=-\infty}^{\infty} a_n e^{in\omega t} \text{ where } T = \frac{2\pi}{\omega} \text{ (period)} \dots (8)$$

The root-mean square or effective value of the function say f_e over a period T is given by

$$\begin{aligned} f_e^2 &= \frac{1}{T} \int_0^T f^2(t) dt \\ &= \frac{1}{T} \int_0^T \left[\sum_{n=-\infty}^{\infty} a_n e^{in\omega t} \sum_{m=-\infty}^{\infty} a_m e^{im\omega t} \right] dt \\ &= \frac{1}{T} \int_0^T \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} a_n a_m e^{i(n+m)\omega t} dt \\ &= \frac{1}{T} \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} a_n a_m \int_0^T e^{i(n+m)\omega t} dt \dots (9) \end{aligned}$$

(on carrying term by term integration)

$$\text{But } \int_0^{2\pi/\omega} e^{ip\omega t} dt = \left[\frac{e^{ip\omega t}}{ip\omega} \right]_0^{2\pi/\omega} = 0 \text{ for integral } m \neq 0 \left. \begin{aligned} &= \frac{2\pi}{\omega} = T \text{ for } m=0 \end{aligned} \right\} \dots (10)$$

which follows that all the integrals in (9) vanish except those for $m=-n$.

$$\therefore (9) \text{ reduces to } f^2_E = \frac{1}{T} \sum_{n=-\infty}^{\infty} a_n a_{-n} T = \sum_{n=-\infty}^{\infty} a_n a_{-n} \dots (11)$$

Regarding a_{-n} as the conjugate of a_n , the quantity in summation of R.H.S. of (11) is the square of the magnitude of a_n . Also the summation over negative values of n yields the same result as summation over positive values of n . Thus (11) reduces to

$$f^0_E = 2 \sum_{n=1}^{\infty} |a_n|^2 + a_0^2 \text{ where } a_0 \text{ corresponds } n=0 \dots (12)$$

Again, to find the average value over a product of two periodic functions with the same period T (say), let us assume two functions

$$f_1(t) = \sum_{n=-\infty}^{\infty} a_n e^{in\omega t}$$

$$\text{and } f_2(t) = \sum_{m=-\infty}^{\infty} b_m e^{im\omega t} \text{ with } T = \frac{2\pi}{\omega} \dots (13)$$

Then average of the product

$$\begin{aligned} &= \frac{1}{T} \int_0^T f_1(t) f_2(t) dt \\ &= \frac{1}{T} \int_0^T \left[\sum_{n=-\infty}^{\infty} a_n e^{in\omega t} \sum_{m=-\infty}^{\infty} b_m e^{im\omega t} \right] dt \\ &= \frac{1}{T} \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} a_n b_m \int_0^T e^{i(n+m)\omega t} dt \text{ (on interchanging} \\ &\quad \text{the order of integration, and summation)} \\ &= \sum_{n=-\infty}^{\infty} a_n b_{-n} \text{ [evaluated just as in (11)]} \dots (14) \end{aligned}$$

[3] **Thermal State.** Let us first consider the Fourier's problem of the permanent state of temperatures in a thin rectangular plate of breadth π and of infinite length, whose faces are impervious to heat. Suppose we have to find the temperature at any point of the plate assuming that the two long edges of the plate are kept at the constant temperature zero, that one of the short edges say the base of the plate is kept at the temperature unity and that the temperatures of the points in the plate decrease indefinitely as we recede from the base.

We know from the Analytical Theory of heat that the change of temperature at different points of a solid when heat flows within the solid, is given by

$$\frac{\partial u}{\partial t} = a^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} \right).$$

where t represents the time, u the temperature at any point (x, y, z) of the solid and a is a constant.

In case of permanent state of temperatures in a thin rectangular plate this reduces to

$$0 = a^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right),$$

i.e. taking the base of the plate as x -axis and one end of the base as origin, the temperature u of any point is given by

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0, \dots (15)$$

$$\text{provided, } u=0 \text{ when } x=0, \dots (16) \quad u=0 \text{ when } x=\pi, \dots (17)$$

$$u=0 \text{ when } y=\infty, \dots (18) \quad u=1 \text{ when } y=0 \dots (19)$$

Let us suppose that $u = A^{\alpha y + \beta x}$ is a solution of (15); then we have

$$\frac{\partial^2 u}{\partial x^2} = A\beta^2 e^{\alpha y + \beta x} \text{ and } \frac{\partial^2 u}{\partial y^2} = A\alpha^2 e^{\alpha y + \beta x}.$$

Substituting these values in (15), we get

$$\alpha^2 + \beta^2 = 0 \text{ or } \beta = \pm i\alpha.$$

Thus we get $u = Ae^{\alpha y \pm i\alpha x}$, where A and α are two constants,

$$\text{i.e. } u = Ae^{\alpha y} e^{i\alpha x} \text{ and } u = Ae^{\alpha y} e^{-i\alpha x}.$$

$$\text{Adding, } 2u = Ae^{\alpha y} (e^{i\alpha x} + e^{-i\alpha x})$$

$$= Ae^{\alpha y} 2 \cos \alpha x.$$

$$\therefore u = Ae^{\alpha y} \cos \alpha x. \dots (20)$$

This solution may also be adjusted in the form

$$u = Be^{\alpha y} \sin \alpha x. \dots (21)$$

The solution (21) satisfies the conditions

$$u=0 \text{ when } x=0, u=0 \text{ when } x=\pi, \alpha \text{ being an integer.}$$

Also $u=0$, when $y=\infty$, α being negative.

Hence u may be expressed as a sum of terms of the form $Ce^{-ny} \sin nx$, n being a positive integer, which satisfies the conditions (16), (17) and (18).

$$\therefore u = C_1 e^{-y} \sin x + C_2 e^{-2y} \sin 2x + C_3 e^{-3y} \sin 3x + \dots (22)$$

Again to satisfy the condition (19), if we put $y=0$ in (15), we get

$$u = C_1 \sin x + C_2 \sin 2x + C_3 \sin 3x + \dots (23)$$

If the series on R.H.S. of (23) be developed to represent unity the condition (19) is satisfied. For this we have to find the suitable values for the coefficients $C_1, C_2, C_3, \dots$ etc.

Consider the Fourier's series defined by $f(x)=1$ for all values of x lying between 0 and π .

$$\text{Here } f(x) \equiv C_1 \sin x + C_2 \sin 2x + C_3 \sin 3x + \dots,$$

where $C_n = \frac{2}{\pi} \int_0^\pi \sin nx \, dx$, $\therefore f(x)=1$; also $f(x)$ is an odd function of x as $f(-x) = -f(x)$.

$$\text{Now } \int_0^\pi \sin nx \, dx = \left[-\frac{\cos nx}{n} \right]_0^\pi = \frac{1}{n} [1 - (-1)^n]$$

= 0 if n is even

and

= $\frac{2}{\pi}$ if n is odd.

Hence

$$f(x) = \sum_{n=1}^{\infty} C_n \sin nx$$

$$= \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{2}{n} \sin nx, \text{ when } n \text{ is odd and zero when } n \text{ is even}$$

$$1 = \frac{4}{\pi} \left[\frac{\sin x}{1} + \frac{\sin 3x}{3} + \frac{\sin 5x}{5} + \dots \right], \text{ } n \text{ being odd which}$$

gives the values of the constants $C_1, C_2, C_3, \dots$ in order that the condition (19) is satisfied.

Substituting values of constants given by $C_n = \frac{4}{n\pi}$ (n being odd) in (22), the required solution to give the temperature at any point is

$$u = \frac{4}{\pi} [e^{-y} \sin x + \frac{1}{3} e^{-3y} \sin 3x + \frac{1}{5} e^{-5y} \sin 5x + \dots]. \quad \dots (24)$$

Note. Similarly we can find the potential function at any point of a long, thin rectangular conducting sheet of breadth π through which an electric current flows, supposing that the two long edges are kept at potential zero and one short edge at potential unity.

Deductions.

(i) If the temperature of the base of the plate be a given function of x , while the other conditions remain unaltered.

We have already shown that

$$f(x) = \sum_{n=1}^{\infty} C_n \sin nx,$$

$$\text{where } C_n = \frac{2}{\pi} \int_0^{\pi} f(y) \sin ny \, dy.$$

Thus we have from (22)

$$u = \frac{2}{\pi} \sum_{n=1}^{\infty} \left[e^{-ny} \sin nx \int_0^{\pi} f(y) \sin ny \, dy \right]. \quad \dots (25)$$

(ii) Assuming the temperature of the base as unity and the breadth of the plate as π , we have prove that

$$u = \frac{4}{\pi} [e^{-y} \sin x + \frac{1}{3} e^{-3y} \sin 3x + \frac{1}{5} e^{-5y} \sin 5y + \dots]. \quad \dots (26)$$

In order to sum the series on R.H.S., let us consider

$$\log(1+z) = z - \frac{1}{2}z^2 + \frac{1}{3}z^3 - \frac{1}{4}z^4 + \dots \text{ when } |z| < 1$$

$$\text{and } \log(1-z) = -z - \frac{1}{2}z^2 - \frac{1}{3}z^3 - \frac{1}{4}z^4 - \dots \text{ when } |z| < 1.$$

$$\therefore \frac{1}{2} [\log(1+z) - \log(1-z)] = z + \frac{z^3}{3} + \frac{z^5}{5} + \dots \text{ when } |z| < 1.$$

$\dots (27)$

Now z being a complex quantity, may be expressed as

$$z = r(\cos \theta + i \sin \theta).$$

Then,

$$\log(1+z) = \log[1+r(\cos \theta + i \sin \theta)]$$

$$= \frac{1}{2} \log[(1+r \cos \theta)^2 + r^2 \sin^2 \theta] + i \tan^{-1} \frac{r \sin \theta}{1+r \cos \theta}$$

$$\left[\because \log(X+iY) = \frac{1}{2} \log(X^2+Y^2) + i \tan^{-1} \frac{Y}{X} \right]$$

$$= \frac{1}{2} \log[1+2r \cos \theta + r^2] + i \tan^{-1} \frac{r \sin \theta}{1+r \cos \theta}.$$

Similarly,

$$\log(1-z) = \frac{1}{2} \log[1-2r \cos \theta + r^2] - i \tan^{-1} \frac{r \sin \theta}{1-r \cos \theta}.$$

$$\therefore \frac{1}{2} [\log(1+z) - \log(1-z)]$$

$$= \frac{1}{2} \left[\log \frac{1+2r \cos \theta + r^2}{1-2r \cos \theta + r^2} + i \left\{ \tan^{-1} \frac{r \sin \theta}{1+r \cos \theta} + \tan^{-1} \frac{r \sin \theta}{1-r \cos \theta} \right\} \right]$$

$$= \frac{1}{2} \left[\frac{1}{2} \log \frac{1+2r \cos \theta + r^2}{1-2r \cos \theta + r^2} + i \tan^{-1} \frac{2r \sin \theta}{1-r^2} \right]$$

$$\left[\because \tan^{-1} a + \tan^{-1} b = \tan^{-1} \frac{a+b}{1-ab} \right].$$

Hence from (27), we have

$$\frac{1}{2} \left[\frac{1}{2} \log \frac{1+2r \cos \theta + r^2}{1-2r \cos \theta + r^2} + i \tan^{-1} \frac{2r \sin \theta}{1-r^2} \right]$$

$$= r(\cos \theta + i \sin \theta) + \frac{r^3(\cos \theta + i \sin \theta)^3}{3} + \dots$$

$$= r(\cos \theta + i \sin \theta) + \frac{r^3(\cos 3\theta + i \sin 3\theta)}{3} + \dots$$

Equating real and imaginary parts, we get

$$\frac{1}{4} \log \frac{1+2r \cos \theta + r^2}{1-2r \cos \theta + r^2} = r \cos \theta + \frac{r^3 \cos 3\theta}{3} + \frac{r^5 \cos 5\theta}{5} + \dots \quad \dots (28)$$

$$\text{and } \frac{1}{2} \tan^{-1} \frac{2r \sin \theta}{1-r^2} = r \sin \theta + \frac{r^3 \sin 3\theta}{3} + \frac{r^5 \sin 5\theta}{5} + \dots \quad \dots (29)$$

(28) and (29) are valid for all values of θ provided $r < 1$.

Also e^{-y} is less than one if y is positive.

Hence replacing r by e^{-y} , (29) gives

$$\frac{e^{-y}}{1} \sin x + \frac{e^{-3y}}{3} \sin 3x + \frac{e^{-5y}}{5} \sin 5x + \dots - \frac{1}{2} \tan^{-1} \frac{2e^{-y} \sin x}{1-e^{-2y}}$$

$$= \frac{1}{2} \tan^{-1} \frac{2 \sin x}{e^y - e^{-y}}$$

$$= \frac{1}{2} \tan^{-1} \frac{\sin x}{\sinh y}.$$

Substituting the value of this series in (18), we get

$$u = \frac{2}{\pi} \tan^{-1} \frac{\sin x}{\sinh y} \quad \dots (30)$$

For Isothermal lines, $u = \text{constant}$

$$\text{i.e. } \frac{2}{\pi} \tan^{-1} \frac{\sin x}{\sinh y} = a \text{ (say)}$$

$$\text{or } \frac{\sin x}{\sinh y} = \tan \frac{a\pi}{2} \quad \dots (31)$$

Again

$$\frac{1}{4} \log \frac{1+2r \cos \theta + r^2}{1-2r \cos \theta + r^2} = \frac{1}{4} \log \frac{1+2e^{-v} \cos x + e^{-2v}}{1-2e^{-v} \cos x + e^{-2v}}$$

replacing r by e^{-v} and θ by x

$$= \frac{1}{4} \log \frac{\frac{e^v + e^{-v}}{2} + \cos x}{\frac{e^v + e^{-v}}{2} - \cos x} = \frac{1}{4} \log \frac{\cosh y + \cos x}{\cosh y - \cos x}$$

For the lines of flow,

$$\frac{1}{\pi} \log \frac{\cosh y + \cos x}{\cosh y - \cos x} = b \text{ (a constant)}$$

$$\text{or } \frac{\cosh y + \cos x}{\cosh y - \cos x} = e^{\pi b} \quad \dots (32)$$

[since $u_1 = \frac{1}{4} \log \frac{\cosh y + \cos x}{\cosh y - \cos x}$ is the solution for the problem if the isothermal lines are the lines of flow of the existing problem and conversely the lines of flow are the isothermal lines of the present problem.]

[4] Transverse Vibrations of a String

We now consider the problem of the transverse vibrations of stretched string fastened at the ends. Suppose that the string is initially distorted into some given curve and then allowed to swing. Let the length of the string be l and the equation of the curve to which the string is distorted initially be $y=f(x)$ with reference to the position of equilibrium of the string as x -axis and one of the ends of the origin.

In acoustics, such vibrations are given by

$$\frac{\partial^2 y}{\partial t^2} = a^2 \frac{\partial^2 y}{\partial x^2} \quad \dots (33)$$

We have to get an expression for y , which is the solution of the equation (33) subject to the conditions

$$y=0 \text{ when } x=0. \quad \dots (34)$$

$$y=0 \text{ when } x=l, \quad \dots (35)$$

$$y=f(x) \text{ when } t=0, \quad \dots (36)$$

$$\text{and } \frac{\partial y}{\partial t} = 0 \text{ when } t=0. \quad \dots (37)$$

Let $y = Ae^{\alpha x + \beta t}$ be the solution of (33),

$$\therefore \frac{\partial^2 y}{\partial t^2} = A^2 \beta e^{\alpha x + \beta t} \text{ and } \frac{\partial^2 y}{\partial x^2} = A \alpha^2 e^{\alpha x + \beta t}.$$

Substituting these values in (33), we get

$$\beta^2 = \alpha^2 a^2,$$

$$\text{giving } \beta = \pm \alpha a$$

$$\therefore y = Ae^{\alpha x \pm aat} \quad \dots (38)$$

is a solution of (33).

Here if we replace α by $+ai$ and $-ai$ in succession

$$y = Ae^{(x \pm ta) ai},$$

$$y = Ae^{-(x \pm ta) ai}.$$

Adding and then dividing by 2, we have

$$y = A \cos \alpha (x \pm at). \quad \dots (39)$$

This may further be expressed as

$$y = B \sin \alpha (x \pm at). \quad \dots (40)$$

From (38),

$$y = A \cos \alpha (x + at) = A \{ \cos \alpha x \cot \alpha a t - \sin \alpha x \sin \alpha a t \}$$

$$\text{and } y = A \cos \alpha (x - at) = A \{ \cos \alpha x \cos \alpha a t + \sin \alpha x \sin \alpha a t \}.$$

Writing y successively equal to half the sum and difference of these values, we can write

$$y = A \cos \alpha x \cos \alpha a t,$$

$$y = A \sin \alpha x \sin \alpha a t.$$

Similarly from (40),

$$y = B \sin \alpha x \cos \alpha a t,$$

$$y = B \cos \alpha x \sin \alpha a t.$$

Out of these four values of y , if we take

$$y = B \sin \alpha x \cos \alpha a t,$$

it is obvious that this satisfies the conditions (34) and (37) and this may be made to satisfy (35) by putting $\alpha = \frac{n\pi}{l}$ when n is any integer.

Hence we can express

$$y = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \cos \frac{n\pi at}{l},$$

$$\text{i.e., } y = b_1 \sin \frac{\pi x}{l} \cos \frac{\pi at}{l} + b_2 \sin \frac{2\pi x}{l} \cos \frac{2\pi at}{l} + b_3 \sin \frac{3\pi x}{l} \cos \frac{3\pi at}{l} + \dots \quad \dots (41)$$

The relation (41) satisfies the conditions (34), (35) and (37). In order that this may satisfy the condition (11) also, let us put $t=0$ in (41), whence we get

$$y = b_1 \sin \frac{\pi x}{l} + b_2 \sin \frac{2\pi x}{l} + b_3 \sin \frac{3\pi x}{l} + \dots \quad (42)$$

If we can however develop $f(x)$ into a series of the form (42), then a comparison will give the values of the coefficients b_1, b_2, b_3 etc., which when substituted in (41), will give a solution of the partial differential equation (33) under the conditions mentioned above.

Now consider the Fourier's series defined by

$$f(x) = b_1 \sin \frac{\pi x}{l} + b_2 \sin \frac{2\pi x}{l} + b_3 \sin \frac{3\pi x}{l} + \dots$$

for all values of x lying between 0 and l .

$$\text{Then } b_n = \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$$

$$= \frac{2}{l} \int_0^l f(v) \sin \frac{n\pi v}{l} dv$$

(replacing x by v in order to discriminate).

Substituting this value of b_n in

$$y = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \cos \frac{n\pi at}{l},$$

the complete solution of the differential equation (8) may be given as

$$y = \frac{2}{l} \sum_{n=1}^{\infty} \sin \frac{n\pi x}{l} \cos \frac{n\pi at}{l} \int_0^l f(v) \sin \frac{n\pi v}{l} dv. \quad (43)$$

Deduction If the shape of the curve into which the string is distorted be given by $y=f(x)=b \sin \frac{n\pi x}{l}$, then

$$f(v) = b \sin \frac{n\pi v}{l}.$$

$$\begin{aligned} \text{Thus } \int_0^l f(v) \sin \frac{n\pi v}{l} dv &= b \int_0^l \sin^2 \frac{n\pi v}{l} dv \\ &= \frac{b}{2} \int_0^l \left(1 - \cos \frac{2n\pi v}{l} \right) dv \\ &= \frac{b}{2} \left[v - \frac{l}{2n\pi} \sin \frac{2n\pi v}{l} \right]_0^l = b \frac{l}{2}. \end{aligned}$$

Hence the solution (43) reduces to

$$y = b \sin \frac{n\pi x}{l} \cos \frac{n\pi at}{l}. \quad (44)$$

Note. For more applications see the next chapter.

9.10. THE FOURIER TRANSFORMS

[A] **Fourier Sine Transforms.** They can be subdivided in two namely the infinite Fourier sine and transform the Finite Fourier sine transforms.

[a₁] **The Infinite Fourier sine Transform** of a function $F(x)$ of x such that $0 < x < \infty$ is denoted by $f_s(n)$, n being a positive integer and is defined as

$$f_s(n) = \int_0^{\infty} F(x) \sin nx \, dx \quad (1)$$

Here $F(x)$ is called as the *Inverse Fourier sine transform* of $f_s(n)$ and defined as

$$F(x) = \frac{2}{\pi} \int_0^{\infty} f_s(n) \sin nx \, dn \quad (2)$$

Thus if $f_s(n) = f_s[F(x)]$, then $F(x) = f_s^{-1}[f_s(n)]$ where f is the symbol for Fourier transform and f^{-1} for its inverse.

Problem 19. Find the sine transform of e^{-x} .

We have

$$f_s(n) = \int_0^{\infty} e^{-x} \sin nx \, dx = \left[\frac{e^{-x}}{1+n^2} (-\sin nx - n \cos nx) \right]_0^{\infty} = \frac{n}{1+n^2}.$$

Problem 20. Find the inverse sine transform of $e^{-\lambda n}$.

$$\begin{aligned} \text{We have } f_s^{-1}[e^{-\lambda n}] &= \frac{2}{\pi} \int_0^{\infty} e^{-\lambda n} \sin nx \, dn = \frac{2}{\pi} \left[\frac{e^{-\lambda n}}{\lambda^2 + x^2} \right. \\ &\quad \left. (-\lambda \sin nx - x \cos nx) \right]_0^{\infty} \\ &= \frac{2}{\pi} \cdot \frac{x}{\lambda^2 + x^2}. \end{aligned}$$

[a₂] **The Finite Fourier sine transform** of a function $F(x)$ of x such that $0 < x < l$ is denoted by $f_s(n)$, n being a positive integer and is defined as

$$f_s(n) = \int_0^l F(x) \sin \frac{n\pi x}{l} \, dx \quad (4)$$

In case $l=\pi$, this becomes

$$f_s(n) = \int_0^{\pi} F(x) \sin nx \, dx \quad (5)$$

and the inversion formula is

$$F(x) = \frac{2}{\pi} \sum_{n=1}^{\infty} f_s(n) \sin nx \quad (6)$$

whence a_n is the coefficient of $\sin nx$ in the expansion of $F(x)$ in a sine series and is given by

$$\begin{aligned} a_n &= \frac{2}{\pi} \int_0^{\pi} F(x) \sin nx \, dx \\ &= \frac{2}{\pi} f_s(n) \text{ by (5)} \end{aligned} \quad (7)$$

Problem 21. Find the Fourier sine transform of $F(x)=x$ such that $0 < x < 2$.

We have $f_s(n) = \int_0^2 F(x) \sin \frac{n\pi x}{2} dx$ $\because l=2$ in the existing case,

$$= \int_0^2 x \sin \frac{n\pi x}{2} dx$$

$$= \left[x \cdot \frac{-\cos \frac{n\pi x}{2}}{\frac{n\pi}{2}} \right]_0^2 + \int_0^2 \frac{2}{n\pi} \cos \frac{n\pi x}{2} dx \text{ (on integrating by parts)}$$

$$= \left[\frac{-2x}{n\pi} \cos \frac{n\pi x}{2} + \frac{4}{n^2\pi^2} \sin \frac{n\pi x}{2} \right]_0^2 = -\frac{4}{n\pi} \cos n\pi.$$

[B] **Fourier Cosine Transforms.** They can also be subdivided into two namely Infinite and finite cosine transforms.

[b₁] *The Infinite Fourier Cosine Transform* of $F(x)$ for $0 < x < \infty$, is defined as

$$f_c(n) = \int_0^\infty F(x) \cos nx \, dx, \quad n \text{ being a positive integer.} \quad \dots (8)$$

Here the function $F(x)$ is called as the *Inverse cosine transform* of $f_c(n)$ and is defined as

$$F(x) = \frac{2}{\pi} \int_0^\infty f_c(n) \cos nx \, dx \quad \dots (9)$$

Thus if $f_c(n) = f_c[F(x)]$, then $F(x) = f_c^{-1}[f_c(n)]$ $\dots (10)$

Problem 22. Find the cosine transform of $x^n e^{-ax}$.

$$\text{we have } \int_0^\infty e^{-ax} \cos nx \, dx = \frac{a}{a^2 + n^2} \text{ and } f_c(n) = \int_0^\infty x^n e^{-ax} \cos nx \, dx$$

Differentiating the first relation n times w.r.t. 'a' we find

$$\int_0^\infty x^n e^{-ax} \cos nx \, dx = (-1)^n \frac{d^n}{da^n} \left(\frac{a}{a^2 + n^2} \right) \\ = \frac{\left| \frac{n}{a^2 + n^2} \cos \left\{ (n+1) \tan^{-1} \frac{n}{a} \right\} \right|}{(a^2 + n^2)^{(n+1)/2}} \text{ by usual method.}$$

$$\text{Hence } f_c(n) = \frac{\left| \frac{n}{a^2 + n^2} \cos \left\{ (n+1) \tan^{-1} \frac{n}{a} \right\} \right|}{(a^2 + n^2)^{(n+1)/2}}$$

Problem 23. Find $f_c^{-1}\{e^{-\lambda n}\}$

$$\text{We have } f_c^{-1}\{e^{-\lambda n}\} = \frac{2}{\pi} \int_0^\infty e^{-\lambda n} \cos nx \, dn = \frac{2}{\pi} \left[\frac{e^{-\lambda n}}{\lambda^2 + x^2} \right. \\ \left. (-\lambda \cos nx + x \sin nx) \right]_0^\infty = \frac{2}{\lambda^2 + x^2}$$

[b₂] *The Finite Fourier cosine transform* of $F(x)$ for $0 < x < l$ is defined as

$$f_c(n) = \int_0^l F(x) \cos \frac{n\pi x}{l} \, dx \quad \dots (11)$$

when $l=\pi$, this becomes

$$f_c(n) = \int_0^\pi F(x) \cos nx \, dx \quad \dots (12)$$

and the inversion formula is

$$F(x) = \frac{1}{\pi} f_c(0) + \frac{2}{\pi} \sum_{n=1}^\infty f_c(n) \cos nx \quad \dots (13)$$

$$\text{when } f_c(0) = \int_0^\pi F(x) \, dx \quad \dots (14)$$

Also b_n the coefficient of $\cos nx$ in the expansion of $F(x)$ in a cosine series is given by

$$b_n = \frac{2}{\pi} \int_0^\pi F(x) \cos nx \, dx = \frac{2}{\pi} f_c(n) \text{ by (12)} \quad \dots (15)$$

Problem 24. Find the finite Fourier cosine transform of x .

$$\text{We have } f_c(n) = \int_0^\pi x \cos nx \, dx \\ = \left[\frac{x \sin nx}{n} \right]_0^\pi - \frac{1}{n} \int_0^\pi \sin nx \, dx \text{ (on integrating by parts)} \\ = 0 - \frac{1}{n} \left[\frac{\cos nx}{-n} \right]_0^\pi \\ = \frac{1}{n^2} \{(-1)^n - 1\}, \quad n=1, 2, 3, \dots$$

$$\text{But if } n=0, f_c(0) = \int_0^\pi x \, dx = \left[\frac{x^2}{2} \right]_0^\pi = \frac{\pi^2}{2}.$$

Note. On the next page are tabulated some useful Fourier sine and cosine transforms in a concise form.

[C] **The Complex Fourier Transforms.**

The Complex Fourier Transform of a function $F(x)$ for $-\infty < x < \infty$, is defined as

$$f(n) = \int_{-\infty}^\infty F(x) e^{inx} \, dx \quad \dots (16)$$

where e^{inx} is said to be the *Kernel* of the transform.

$$\text{The inversion formula is } F(x) = \frac{1}{2\pi} \int_{-\infty}^\infty f(n) e^{-inx} \, dn \quad \dots (17)$$

Problem 25. Find the Fourier Complex Transform of

$$F(x) = \begin{cases} 1-x^2, & |x| < 1 \\ 0, & |x| > 1 \end{cases}$$

Infinite Fourier Sine Transforms

$$f_s(n) = \int_0^\infty F(x) \sin nx \, dx, \quad F(x) = \frac{2}{\pi} \int_0^\infty f_s(n) \sin x \, dn$$

$F(x)$	$f_s(n)$
e^{-x}	$\frac{n}{(1+n^2)}$
$xe^{-x^2/2}$	$\sqrt{\frac{\pi}{2}} ne^{-n^2/2}$
$\frac{\sin x}{x}$	$\frac{1}{2} \log \left \frac{1+n}{1-n} \right $
$x(1-x^2)^2, 0 < x < 1, v > -1$ 0, $x > 1$	$\sqrt{\frac{\pi}{2}} 2^{v-1/2} \Gamma(v+1) n^{-1/2-v} J_{v+3/2}(n)$
$x^{\mu-1}, 0 < \mu < 1$	$\sqrt{\mu} \sin \left(\frac{\mu\pi}{2} \right) n^{-\mu}$
$x^\nu e^{-x}$	$2^\nu \frac{\mu^\nu}{(\Gamma^\nu)} \frac{n}{(\mu^2+n^2)^{\nu+1}}$
0, $0 < x < a$ $(x^2-a^2)^{-1/2}, x > a$	$J_0(an)$

Finite Fourier Sine Transforms

$$f_s(n) = \int_0^\pi F(x) \sin nx \, dx, \quad F(x) = \frac{2}{\pi} \sum_{n=1}^\infty f_s(n) \sin nx$$

$F(x)$	$f_s(n)$
1	$\frac{1}{n} \left[1 + (-1)^{n+1} \right]$
x	$(-1)^{n+1} \frac{\pi}{n}$
$x, 0 < x \leq \pi/2$ $\pi-x, \pi/2 < x < \pi$	$\frac{2}{n^2} \sin \frac{n\pi}{2}$
$1 - \frac{x}{\pi}$	$\frac{1}{n}$
x^2	$\frac{\pi^2}{n} (-1)^{n-1} - \frac{2}{n^3} \left[1 - (-1)^n \right]$
$x(\pi-x)$	$\frac{2}{n^3} [1 - (-1)^n]$
x^3	$(-1)^{n+1} \pi \left[\frac{6}{n^3} - \frac{\pi^2}{n} \right]$
$x(\pi^2-x^2)$	$\frac{n}{n^2+\lambda^2} \left[1 + (-1)^n e^{\lambda\pi} \right]$
$e^{\lambda x}$	$\frac{n}{n^2-\lambda^2} [1 - (-1)^n \cos \lambda\pi]$
$\cos \lambda x$	$\begin{cases} \frac{n}{n^2-m^2} [1 - (-1)^{n+m}], n \neq m \\ 0, n = m \end{cases}$
$\cos mx, m=1, 2, 3, \dots$	$\begin{cases} \frac{n}{n^2-m^2} [1 - (-1)^{n+m}], n \neq m \\ 0, n = m \end{cases}$
$\sin mx, m=1, 2, 3, \dots$	$\frac{\pi}{2}, n=m$
$\frac{\sin \lambda(\pi-x)}{\sin \lambda x}$	$\frac{n}{n^2-\lambda^2}, \lambda \neq 0, 1, 2, \dots$
$\frac{\sin \lambda(\pi-x)}{\sin \lambda x}$	$\frac{\pi}{\lambda^2+\pi^2}$

Infinite Fourier Cosine Transforms

$$f_c(n) = \int_0^\infty F(x) \cos nx \, dx, \quad F(x) = \frac{2}{\pi} \int_0^\infty f_c(n) \cos nx \, dx$$

$F(x)$	$f_c(n)$
$1, 0 < x < a$ $0, x > a$	$\frac{\sin na}{n}$
$x^{\mu-1}, 0 < \mu < 1$	$\sqrt{\frac{\pi}{2}} n^{-\mu} \sin \frac{\mu\pi}{2}$
e^{-x}	$\frac{1}{1+n^2}$
$\text{Sech } x^\pi$	$\sqrt{\frac{\pi}{2}} \frac{1}{1+n^4}$
e^{-x^2}	$\sqrt{\frac{\pi}{2}} e^{-n^2/4}$
$\cos x, 0 < x < a$ $0, x > a$	$\frac{1}{2} \left[\sin \frac{a(1-n)}{1-n} + \sin \frac{a(1+n)}{1+n} \right]$
$\sin \frac{x^2}{2}$	$\sqrt{\frac{\pi}{2}} \left[\cos \frac{n^2}{2} - \sin \frac{n^2}{2} \right]$
$\cos \frac{x^2}{2}$	$\sqrt{\frac{\pi}{2}} \left[\cos \frac{n^2}{2} + \sin \frac{n^2}{2} \right]$
$(1-x^2)^v, 0 < x < 1$ $0, x > 1$ and $v > -3/2$	$\sqrt{\pi} 2^{v-1/2} \Gamma(v+1) n^{-v-1/2} J_{v+1/2}(n)$

Finite Fourier Cosine Transforms

$$f_c(n) = \int_0^\pi F(x) \cos nx \, dx, \quad F(x) = \frac{1}{\pi} f_c(0) + \sum_{n=1}^\infty f_c(n) \cos nx$$

$F(x)$	$f_c(n)$
1	$\begin{cases} \pi, n=0 \\ 0, n=1, 2, 3, \dots \end{cases}$
$1, 0 < x < \pi/2$ $-1, \pi/2 < x < \pi$	$\begin{cases} 0, n=0 \\ 2/\pi \sin n\pi/2, n=1, 2, 3, \dots \end{cases}$
x	$\begin{cases} \pi^2/2, n=0 \\ 1/n^2 [(-1)^{n-1}], n=1, 2, 3, \dots \end{cases}$
x^2	$\begin{cases} \pi^3/3, n=0 \\ 2\pi/n^2 (-1)^n, n=1, 2, 3, \dots \end{cases}$
x^3	$\begin{cases} \pi^4/4, n=0 \\ 3\pi^2/n^2 (-1)^n + 6/n^4 [(-1)^{n-1}], n=1, 2, 3, \dots \end{cases}$
$\left(1 - \frac{x}{\pi}\right)^2$	$\begin{cases} \pi/3, n=0 \\ 2/\pi n^2, n=1, 2, 3, \dots \end{cases}$
$e^{\lambda x}$	$\frac{\lambda}{\lambda^2+n^2} [(-1)^n e^{\lambda\pi-1}]$
$\sin \lambda x$	$\frac{\lambda}{n^2-\lambda^2} [(-1)^n \cos \lambda\pi-1], n \neq \lambda$
$\sin mx, m=1, 2, 3, \dots$	$\begin{cases} 0, n=m \\ m/n^2-m^2 [(-1)^{n+m-1}], n \neq m \end{cases}$
$\frac{\cos \lambda(\pi-x)}{\sin \lambda \pi}$	$\frac{\lambda}{\lambda^2+n^2}$

We have $f(n) = \int_{-1}^1 (1-x^2) e^{inx} dx = \left[(1-x^2) \frac{e^{inx}}{in} + \frac{2}{in} \int_{-1}^1 x e^{inx} dx \right]$ (on integrating by parts)

$$= 0 + \frac{2}{in} \left[\frac{x e^{inx}}{in} \right]_{-1}^1 - \frac{2}{(in)^2} \int_{-1}^1 e^{inx} dx$$

$$= \frac{2}{-n^2} [e^{in} + e^{-in}] + \frac{2}{n^2} \left[\frac{e^{inx}}{in} \right]_{-1}^1 = -\frac{2}{n^2} (e^{in} + e^{-in}) + \frac{2}{in^3} (e^{in} - e^{-in})$$

$$= -\frac{4}{n^2} \cos n + \frac{4}{n^3} \sin n = -\frac{4}{n^3} (n \cos n - \sin n).$$

Problem 26. Find the Complex Fourier transform of $e^{-|x|}$ and then invert it.

We have $f(n) = \int_{-\infty}^{\infty} e^{-|x|} e^{inx} dx = \int_{-\infty}^0 e^{(1+in)x} dx + \int_0^{\infty} e^{-(1-in)x} dx$

$$= \frac{1}{1+in} + \frac{1}{1-in} = \frac{2}{1+n^2}$$

So that the inversion formula gives,

$$F(n) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{2}{1+n^2} e^{inx} dx = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{e^{-tnx}}{1+n^2} dx \text{ which may be integrated by contour Integration.}$$

Note. Several other Complex Fourier Transforms have been tabulated on page 843.

[D] Parseval's Identity for Fourier Integrals.

It is stated as $\int_{-\infty}^{\infty} |F(x)|^2 dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} |f(n)|^2 dn \dots (18)$

where $f(n)$ is the Fourier transform of $F(x)$.

Problem 27. Find the Fourier transform of

$$F(x) = \begin{cases} I, & |x| < a \\ 0, & |x| > a \end{cases}$$

Hence or otherwise evaluate $\int_0^{\infty} \frac{\sin^2 nx}{n^2} dx$.

We have $f(n) = \int_{-\infty}^{\infty} F(x) e^{-inx} dx = \int_{-a}^a 1 \cdot e^{-inx} dx$

$$\left[\frac{e^{-inx}}{-in} \right]_{-a}^a = \frac{e^{+ina} - e^{-ina}}{in} = \frac{2 \sin na}{n} \text{ for } n \neq 0$$

$f(n)$	$F(x)$	$f(n) = \int_{-\infty}^{\infty} F(x) e^{inx} dx$	$F(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(n) e^{-inx} dn$
$\frac{\sin a}{\cosh n + \cos a}$	$\frac{\sinh ax}{\sinh \pi x}, -\pi < a < \pi$	$\frac{1}{ n }$	$\frac{1}{ x }$
$2 \frac{\cos a/2 \cosh a/2}{\cosh n + \cos a}$	$\frac{\cosh ax}{\cosh \pi x}, -\pi < a < \pi$	$\frac{e^{ia(\omega+n)} - e^{ib(\omega+n)}}{n}$	$\frac{e^{i\omega x}, a < x < b}{0, x < a, x > b}$
$\begin{cases} 0, & n > b \\ \pi J_0 [a\sqrt{b^2 - n^2}], & n < b \end{cases}$	$\frac{\sin \{ b(a^2 + x^2)^{1/2} \}}{(a^2 + x^2)^{1/2}}, \dots$	$\frac{i}{\omega + n + i\lambda}$	$\frac{e^{-\lambda x + i\omega x}, x > 0}{0, x < 0}$
$\pi J_0 [a\sqrt{n^2 + b^2}]$	$\frac{\cosh \{ b(a^2 - x^2)^{1/2} \}}{(a^2 - x^2)^{1/2}}, x < a$	$\frac{[\lambda^2 + n^2]^{1/2} + \lambda]^{1/2}}{(\lambda^2 + n^2)^{1/2}}$	$\frac{e^{-\lambda} x }{ x ^{1/2}}, \dots$
$\pi J_0 [a\sqrt{n^2 - b^2}]$	$\frac{\cos \{ b(a^2 - x^2)^{1/2} \}}{\sqrt{a^2 - x^2}}, x < a$	$\frac{\sqrt{\pi}}{\lambda} e^{-n^2/4\lambda}$	$e^{-\lambda x^2}, R(\lambda) > 0$
$i^N \sqrt{2} J_{N+1/2}(n)$	$\frac{P_N(x), x < 1}{0, x > 1}$	$\frac{\sqrt{\pi}}{\lambda} \sin \left(\frac{n^2}{4\lambda} + \frac{\pi}{4} \right)$	$\sin \lambda x^2, \dots$
		$\frac{\sqrt{\pi}}{\lambda} \cos \left(\frac{n^2}{4\lambda} - \frac{\pi}{4} \right)$	$\cos \lambda x^2$
		$\pi J_0(an)$	$(a^2 - x^2)^{-1/2}, x < a$ $0, x > a$

$$\text{For } n=0, f(n) = 2 \lim_{n \rightarrow 0} \frac{\sin na}{n} = 2 \lim_{n \rightarrow 0} \frac{1}{n} \left[na - \frac{1}{3} n^3 a^3 + \dots \right] = 2a$$

Now using Parseval's identity, we find

$$\int_{-a}^a 1^2 dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{4 \sin^2 na}{n^2} dn \quad \text{when } n \neq 0$$

$$\text{i.e. } \frac{1}{2\pi} \cdot 2 \int_0^{\infty} \frac{4 \sin^2 na}{n^2} dn = \left[x \right]_{-a}^a = 2a$$

$$\therefore \int_0^{\infty} \frac{\sin^2 nx}{n^2} dx = \frac{\pi x}{2}$$

[E] Relation between the Fourier Transform of the Derivatives of a Function.

If $f(n)$ be the Fourier transform of $F(x)$, then we have to express the Fourier transform of the function $\frac{d^m F}{dx^m}$ in terms of $f(n)$.

We have by the definition of Fourier-transform,

$$f \left[\frac{d^m F}{dx^m} \right] = \int_{-\infty}^{\infty} \frac{d^m F}{dx^m} e^{inx} dx = f^m(n) \quad (\text{say}) \quad \dots (19)$$

$$\text{So that } f^m(n) = \int_{-\infty}^{\infty} \frac{d^m F}{dx^m} e^{inx} dx = \left[\frac{d^{m-1} F}{dx^{m-1}} e^{inx} \right]_{-\infty}^{\infty}$$

$$- \int_{-\infty}^{\infty} (in) e^{inx} \frac{d^{m-1} F}{dx^{m-1}} dx \quad (\text{on integrating by parts})$$

$$= -in \int_{-\infty}^{\infty} \frac{d^{m-1} F}{dx^{m-1}} e^{inx} dx, \text{ under the assumption } \frac{d^{m-1} F}{dx^{m-1}} \rightarrow 0$$

$$\text{as } |x| \rightarrow \infty \quad \dots (20)$$

$$= -in f^{m-1}(n) \text{ by (19)}$$

Repeating the same process under the assumption $\frac{d^r F}{dx^r} \rightarrow 0$

as $|x| \rightarrow \infty, r=1, 2, 3, \dots, (m-1)$ we get after $(m-1)$ operations, $f^m(n) = (-in)^m f(n) \quad \dots (21)$

which follows that the Fourier transform of $\frac{d^m F}{dx^m}$ is $(-in)^m$ times the

Fourier transform of $F(x)$ subject to the condition that $\frac{d^r F}{dx^r} \rightarrow 0$ when $|x| \rightarrow \infty$, for $r=1, 2, 3, \dots, (m-1)$.

By similar procedure we can find a relation between the sine and cosine Fourier transforms of the derivatives of a function, such as

$$f_c^m(n) = \int_0^{\infty} \frac{d^m F}{dx^m} \cos nx dx = \left[\frac{d^{m-1} F}{dx^{m-1}} \cos nx \right]_0^{\infty}$$

$$+ n \int_0^{\infty} \frac{d^{m-1} F}{dx^{m-1}} \sin nx dx, \text{ integrating by parts}$$

$$= -\alpha_{n-1} + n f_s^{m-1}(n) \quad \dots (22)$$

Under the assumptions,

$$\frac{d^{m-1} F}{dx^{m-1}} \rightarrow 0 \text{ as } x \rightarrow \infty \text{ and } \frac{d^{m-1} F}{dx^{m-1}} \rightarrow \alpha_{n-1} \text{ as } x \rightarrow 0.$$

$$\text{Similarly, integrating, } f_s^m(n) = \int_0^{\infty} \frac{d^m F}{dx^m} \sin nx dx$$

$$= -n f_c^{m-1}(n) \quad \dots (23)$$

(22) and (23) yield,

$$f_c^m(n) = -\alpha_{n-1} - n^2 f_s^{m-2}(n) \quad \dots (24)$$

Repeating the procedure $f_c^m(n)$ may be expressed as the sum of α 's and either $f_c(n)$ or $f_s(n)$ or $f_c'(n)$ or $f_s'(n)$ will occur when x is odd and in that case we can write $\alpha_0 + n f_s(n)$ in place of $f_c'(n)$. We thus have

$$f_c^{2m}(n) = - \sum_{r=0}^{m-1} (-1)^r \alpha_{2m-2r-1} n^{2r} + (-1)^m n^{2m} f_c(n) \quad \dots (25)$$

$$\text{and } f_c^{2m+1}(n) = - \sum_{r=0}^m (-1)^r \alpha_{2m-2r} n^{2r} + (-1)^m n^{2m+1} f_s(n) \quad \dots (26)$$

Similar procedure with the help of (22) and (23), will yield

$$f_s^m(n) = n \alpha_{n-2} - n^2 f_s^{m-2}(n) \quad \dots (27)$$

$$f_s^{2m}(n) = - \sum_{r=1}^m (-1)^r n^{2r-1} \alpha_{2n-2r} + (-1)^{m+1} n^{2m} f_s(n) \quad \dots (28)$$

$$\text{and } f_s^{2m+1}(n) = - \sum_{r=1}^m (-1)^r n^{2r-1} \alpha_{2n-2r+1} + (-1)^{m+1} n^{2m+1} f_c(n)$$

$$\quad \dots (29)$$

Note 1. The following results are easily deducible

$$\left. \begin{aligned} (i) \int_0^{\infty} \frac{d^2 F}{dx^2} \cos nx dx &= -n^2 f_c(n) \\ (ii) \int_0^{\infty} \frac{d^4 F}{dx^4} \cos nx dx &= n^4 f_c(n) \end{aligned} \right\} \text{when } x=0, \frac{dF}{dx} = \frac{d^3 F}{dx^3} = 0 \quad \dots (30)$$

$$\left. \begin{aligned} (iii) \int_0^{\infty} \frac{d^2 F}{dx^2} \sin nx dx &= n^2 f_s(n) \\ (iv) \int_0^{\infty} \frac{d^4 F}{dx^4} \sin nx dx &= n^4 f_s(n) \end{aligned} \right\} \text{when } x=0, F = \frac{d^2 F}{dx^2} = 0 \quad \dots (31)$$

$$(v) \int_0^{\infty} \frac{\partial F}{\partial t} \sin nx dx = \frac{\partial}{\partial t} \int_0^{\infty} F \sin nx dx = \frac{\partial f}{\partial t} \quad \dots (32)$$

Note 2. In case the transforms are finite, then consider

$$\int_0^{\pi} \frac{\partial F}{\partial x} \sin nx dx = \left[F(x) \sin nx \right]_0^{\pi} - n \int_0^{\pi} F(x) \cos nx dx, \text{ integrating by parts}$$

$$= -n f_c(n) \quad \dots (33)$$

under the assumption that $F(0)$ and $F(\pi)$ both are finite.

$$\begin{aligned}\text{Similarly, } \int_0^\pi \frac{\partial F}{\partial x} \cos nx \, dx &= \left[F(x) \cos nx \right]_0^\pi + n \int_0^\pi F(x) \sin nx \, dx \\ &= (-1)^n F(\pi) - F(0) + n f_s(n) \quad \dots (34)\end{aligned}$$

Assuming that $F(x) \rightarrow 0$ at $x=\pi$ and at $x=0$, (34) reduces to

$$\int_0^\pi \frac{\partial F}{\partial x} \cos nx \, dx = n f_s(n) \quad \dots (35)$$

and (33) reduces to

$$\begin{aligned}\int_0^\pi \frac{\partial^2 F}{\partial x^2} \sin nx \, dx &= -n \int_0^\pi \frac{\partial F}{\partial x} \cos nx \, dx \\ &= n[(-1)^{n+1} F(\pi) + F(0)] - n^2 f_s(n) \text{ by (34)} \dots (36)\end{aligned}$$

If $F(0)=F(\pi)=0$, then (36) yields,

$$\int_0^\pi \frac{\partial^2 F}{\partial x^2} \sin nx \, dx = -n^2 f_s(n) \quad \dots (37)$$

Similarly (34) yields

$$\int_0^\pi \frac{\partial^2 F}{\partial x^2} \cos nx \, dx = (-1)^n F'(\pi) - F'(0) - n^2 f_c(n) \quad \dots (38)$$

In case $\frac{\partial^2 F}{\partial x^2}$ vanishes at $x=0$ and at $x=\pi$, it is easy to see that

(31) gives

$$\int_0^\pi \frac{\partial^4 F}{\partial x^4} \sin nx \, dx = -n^2 \int_0^\pi \frac{\partial^2 F}{\partial x^2} \sin nx \, dx = n^4 f_s(n) \quad \dots (39)$$

and when $\frac{\partial F}{\partial x}, \frac{\partial^3 F}{\partial x^3}$ vanish at $x=0$ and at $x=\pi$, (38) gives

$$\int_0^\pi \frac{\partial^2 F}{\partial x^2} \cos nx \, dx = -n^2 f_c(n) \quad \dots (40)$$

$$\text{So that } \int_0^\pi \frac{\partial^4 F}{\partial x^4} \cos nx \, dx = n^4 f_c(n) \quad \dots (41)$$

Problem 28. Determine the function F such that

$$\frac{\partial^2 F}{\partial x^2} + \frac{\partial^2 F}{\partial y^2} = 0, \quad 0 < x < \pi,$$

with the boundary condition $F=0$ when $x=0$ and $x=\pi$

$$= 0 \text{ when } y=0$$

$$= F_0 \text{ (const.) when } y=\pi.$$

F being given to be zero when $x=0$ and $x=\pi$, we have to use the finite sine transform i.e. $f(n) = \int_0^\pi F(x) \sin nx \, dx$

Applying it to the given differential equation we have

$$\int_0^\pi \frac{\partial^2 F}{\partial x^2} \sin nx \, dx + \int_0^\pi \frac{\partial^2 F}{\partial y^2} \sin nx \, dx = 0$$

with the condition, $f=0$ when $y=0$ and $f = \int_0^\pi F_0 \sin nx \, dx$ when $y=\pi$

$$\text{By (37) we have, } \int_0^\pi \frac{\partial^2 F}{\partial x^2} \sin nx \, dx = -n^2 f$$

$$\therefore -n^2 f + \frac{\partial^2 f}{\partial y^2} = 0 \text{ where } \frac{\partial^2 f}{\partial y^2} = \int_0^\pi \frac{\partial^2 F}{\partial y^2} \sin nx \, dx$$

$$\text{or } \frac{\partial^2 f}{\partial y^2} - n^2 f = 0.$$

Its general solution is $f = A \sinh ny$

$$\text{But } f = F_0 \int_0^\pi \sin nx \, dx \text{ when } y=\pi$$

$$\begin{aligned}&= F_0 \left[-\frac{\cos nx}{n} \right]_0^\pi = 0 \text{ when } n \text{ is even} \\ &= -2 F_0/n \text{ when } n \text{ is odd.}\end{aligned}$$

So that considering the two solutions for f we conclude $f=0$ when n is even

and $f = \frac{2F_0}{n} \operatorname{cosech} n\pi \sinh ny$ when n is odd.

Hence the inversion formula will give on replacing n by $2m+1$,

$$F = \frac{4F_0}{\pi} \sum_{m=0}^{\infty} (2m+1)^{-1} \operatorname{cosech} (2m+1)\pi \sinh (2m+1)y \sin (2m+1)x$$

[F] Multiple Fourier Transforms.

If $F(x, y)$ be a function of two variables x and y , then assuming it to be the function of x only, its fourier transform $\phi(n, y)$ is given by

$$\phi(n, y) = \int_{-\infty}^{\infty} F(x, y) e^{inx} \, dx \quad \dots (42)$$

Now if $f(n, l)$ be the Fourier complex transform of $\phi(n, y)$ which is regarded as function of y only then

$$f(n, l) = \int_{-\infty}^{\infty} \phi(n, y) e^{ily} \, dy \quad \dots (43)$$

These two results when combined, give

$$f(n, l) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) e^{i(nx+ly)} \, dx \, dy \quad \dots (44)$$

and the inversion formula is

$$f(x, y) = \frac{1}{4\pi^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(n, l) e^{-i(nx+ly)} \, dn \, dl \quad \dots (45)$$

Note 1. The result may be generalized for any number of variables.

Note 2. In case the Fourier transforms are finite such that $F(x, y)$ is a function of two independent variables x, y where $0 \leq x \leq \pi$ and $0 \leq y \leq \pi$, then the sine transform of $F(x, y)$ is given by

$$f_s(n, l) = \int_0^\pi \int_0^\pi F(x, y) \sin nx \sin ly \, dx \, dy \quad \dots (46)$$

and the inversion formula is

$$F(x, y) = \frac{4}{\pi^2} \sum_{n=1}^{\infty} \sum_{l=1}^{\infty} f_s(n, l) \sin nx \sin ly \quad \dots (47)$$

[G] Convolution or Faltung Theorem for Fourier Transforms.

If $F(x)$ and $G(x)$ are two functions such that $-\infty < x < \infty$ then their Faltung or Convolution $F * G$ is defined as

$$H(x) = F * G = \int_{-\infty}^{\infty} F(n) G(x-n) \, dn \quad \dots (48)$$

It is worth noting that the Fourier Transform of the Convolution of $F(x)$ and $G(x)$ is the product of their Fourier transforms i.e.

$$f[F * G] = f[F] f[G] \quad \dots (49)$$

Since $f[F * G] = \int_{-\infty}^{\infty} H(x) e^{-inx} \, dx$ by definition

$$= \int_{-\infty}^{\infty} F(x) e^{-inx} \, dx \int_{-\infty}^{\infty} G(x) e^{-inx} \, dx$$

$$= f[F] \cdot f[G].$$

[H] Evaluation of Integrals with the help of Fourier Inversion Theorem.

Let $I_1 = \int_0^\infty e^{-ax} \cos nx \, dx$ and $I_2 = \int_0^\infty e^{-ax} \sin nx \, dx$.

Integrating by parts, we have

$$I_1 = \left[-\frac{1}{a} e^{-ax} \cos nx \right]_0^\infty - \frac{n}{a} \int_0^\infty e^{-ax} \sin nx \, dx = \frac{1}{a} - \frac{n}{a} I_2$$

Similarly $I_2 = \frac{n}{a} I_1$

These give on solving $I_1 = \frac{a}{a^2 + n^2}$ and $I_2 = \frac{n}{a^2 + n^2}$

Thus taking $F(x) = e^{-ax}$, its sine and cosine Fourier transforms are $\frac{a}{a^2 + n^2}$ and $\frac{n}{a^2 + n^2}$ respectively, so that the inversion formula gives

$$e^{-ax} = \frac{2}{\pi} \int_0^\infty \frac{a}{a^2 + n^2} \cos nx \, dn \quad \dots (50)$$

$$\text{and } e^{-ax} = \frac{2}{\pi} \int_0^\infty \frac{n}{a^2 + n^2} \sin nx \, dn \quad \dots (51)$$

$$\text{i.e. } \int_0^\infty \frac{\cos nx}{a^2 + n^2} \, dn = \frac{\pi}{2a} e^{-ax} \text{ and } \int_0^\infty \frac{n \sin nx}{a^2 + n^2} \, dn = \frac{\pi}{2} e^{-ax} \dots (52)$$

9.11. APPLICATIONS OF FOURIER TRANSFORMS TO BOUNDARY VALUE PROBLEMS

Problem 29. Find the finite Fourier sine and cosine transform of

$\frac{\partial^2 U}{\partial x^2}$ where U is a function of x and t for $0 < x < l$, $t > 0$.

We have $f_s \left\{ \frac{\partial U}{\partial x} \right\} = \int_0^l \frac{\partial U}{\partial x} \sin \frac{n\pi x}{l} \, dx$, by definition

$$= \left[U(x, t) \sin \frac{n\pi x}{l} \right]_0^l - \frac{n\pi}{l} \int_0^l U(x, t) \cos \frac{n\pi x}{l} \, dx,$$

integrating by parts

$$= 0 - \frac{n\pi}{l} f_c[U] = -\frac{n\pi}{l} f_c\{U\} \quad \dots (1)$$

$$\text{Also } f_c \left\{ \frac{\partial U}{\partial x} \right\} = \int_0^l \frac{\partial U}{\partial x} \cos \frac{n\pi x}{l} \, dx = \left[U(x, t) \cos \frac{n\pi x}{l} \right]_0^l$$

$$+ \frac{n\pi}{l} \int_0^l U(x, t) \sin \frac{n\pi x}{l} \, dx$$

$$= U(l, t) \cos n\pi - U(0, t) + \frac{n\pi}{l} f_s\{U\} \quad \dots (2)$$

Replacing U by $\frac{\partial U}{\partial x}$ in (1), we get

$$f_s \left\{ \frac{\partial^2 U}{\partial x^2} \right\} = -\frac{n\pi}{l} f_c \left\{ \frac{\partial U}{\partial x} \right\}$$

$$= -\frac{n\pi}{l} \left[U(l, t) \cos n\pi - U(0, t) + \frac{n\pi}{l} f_s\{U\} \right] \text{ by (2)}$$

$$= -\frac{n\pi}{l} U(l, t) \cos n\pi + \frac{n\pi}{l} U(0, t) - n^2 \frac{\pi^2}{l^2} f_s\{U\}$$

Again replacing U by $\frac{\partial U}{\partial x}$ in (2) we find

$$f_c \left\{ \frac{\partial^2 U}{\partial x^2} \right\} = \frac{\partial U(l, t)}{\partial x} \cos n\pi - \frac{\partial U(0, t)}{\partial x} + \frac{n\pi}{l} f_s \left\{ \frac{\partial U}{\partial x} \right\}$$

$$= \frac{\partial U(l, t)}{\partial x} \cos n\pi - \frac{\partial U(0, t)}{\partial x} - \frac{n^2 \pi^2}{l^2} f_c\{U\} \text{ by (1)}$$

Problem 30. Solve $\frac{\partial U}{\partial t} = \frac{\partial^2 U}{\partial x^2}$, $x > 0$, $t > 0$ subject to the conditions

$$U(0, t) = 0, U(x, 0) = \begin{cases} 1, & 0 < x < l \\ 0, & x > l \end{cases} \text{ and } U(x, t) \text{ is bounded.}$$

Taking the Fourier sine transform of both sides of the given equation, we get

$$\int_0^\infty \frac{\partial U}{\partial t} \sin nx \, dx = \int_0^\infty \frac{\partial^2 U}{\partial x^2} \sin nx \, dx \quad \dots (1)$$

Denoting $\int_0^\infty U(x, t) \sin nx \, dx$ by $u = u(n, t)$, we have

$$\frac{du}{dt} = \int_0^\infty \frac{\partial U(x, t)}{\partial t} \sin nx \, dx = \int_0^\infty \frac{\partial^2 U}{\partial x^2} \sin nx \, dx \text{ by (1)}$$

$$= \left[\sin nx \frac{\partial U}{\partial x} \right]_0^\infty + n \int_0^\infty \cos nx \frac{\partial U}{\partial x} \, dx, \text{ on integrating by parts.}$$

$$= 0 - n \left[\cos nx U \Big|_0^\infty + n \int_0^\infty \sin nx \cdot U dx \right], \text{ under the assumption, } \frac{\partial U}{\partial x} \rightarrow 0 \text{ as } x \rightarrow \infty$$

$$= n U(0, t) - n^2 u, \text{ under the assumption } U \rightarrow 0 \text{ as } x \rightarrow \infty \quad \dots (2)$$

Now $u(n, t) = \int_0^\infty U(x, t) \sin nx dx$ gives

$$u(n, 0) = \int_0^\infty U(x, 0) \sin nx dx = \int_0^1 1 \cdot \sin nx dx \text{ by the condition for } U(x, 0)$$

$$= \left[-\frac{\cos nx}{n} \right]_0^1 = \frac{1 - \cos n}{n} \quad \dots (3)$$

Using the condition $U(0, t) = 0$, (2) yields $\frac{du}{dt} = -n^2 u$ or $\frac{du}{u} = -n^2 dt$.

Integrating $\log u = -n^2 t + \log A$, A being constant of integration
 $\therefore u = Ae^{-n^2 t}$.

Initially when $t = 0$, $u(n, t) = u(n, 0) = \frac{1 - \cos n}{n}$ by (3); so that

$$A = \frac{1 - \cos n}{n}$$

Hence $u(n, t) = \frac{1 - \cos n}{n} e^{-n^2 t}$

Applying the inversion formula for Fourier sine transform, we have

$$U(x, t) = \frac{2}{\pi} \int_0^\infty u(n, t) \sin nx dx = \frac{2}{\pi} \int_0^\infty \frac{1 - \cos n}{n} e^{-n^2 t} \sin nx dx$$

which gives the required solution, physically interpreted as the temperature at any point x at any time t in a solid $x > 0$.

Problem 31. Use finite Fourier transforms to solve $\frac{\partial U}{\partial t} = \frac{\partial^2 U}{\partial x^2}$,

$U(0, t) = 0$, $U(\pi, t) = 0$, $U(x, 0) = 2x$ where $0 < x < \pi$, $t > 0$.

Give physical interpretation of the problem.

Taking the finite Fourier sine transform of both sides of the given differential equation, we have

$$\int_0^\pi \frac{\partial U}{\partial t} \sin nx dx = \int_0^\pi \frac{\partial^2 U}{\partial x^2} \sin nx dx \quad \dots (1)$$

Denoting by $u = u(n, t) = \int_0^\pi U(x, t) \sin nx dx$,

$$\text{find } \int \frac{du}{dt} = \int_0^\pi \frac{\partial U}{\partial t} \sin nx dx = \int_0^\pi \frac{\partial^2 U}{\partial x^2} \sin nx dx \text{ by (1)}$$

$$= \left[\sin nx \frac{\partial U}{\partial x} \right]_0^\pi - n \int_0^\pi \cos nx \frac{\partial U}{\partial x} dx, \text{ integrating by parts}$$

$$= 0 - n \int_0^\pi \cos nx \frac{\partial U}{\partial x} dx \quad \because U(0, t) = U(\pi, t) = 0$$

$$= -n \left[\cos nx \cdot U(x, t) \right]_0^\pi - n^2 \int_0^\pi \sin nx \cdot U(x, t) dx$$

$= -n^2 u$, since the first integral vanishes for $U(0, t) = U(\pi, t) = 0$.
 Its solution is, $u = Ae^{-n^2 t}$ A being constant of integration.

$$\text{Now from (2), } u(n, 0) = \int_0^\pi U(x, 0) \sin nx dx = \int_0^\pi 2x \sin nx dx$$

$$\therefore U(x, 0) = 2x$$

$$= 2 \left[-x \frac{\cos nx}{n} \right]_0^\pi + \frac{2}{n} \int_0^\pi \cos nx dx,$$

$$= -\frac{2\pi}{n} \cos n\pi + \frac{2}{n^2} \left[\sin nx \right]_0^\pi = -\frac{2\pi}{n} \cos n\pi$$

When $t = 0$, $u = -\frac{2\pi}{n} \cos n\pi$ applied to $u = Ae^{-n^2 t}$ gives

$$A = -\frac{2\pi}{n} \cos n\pi$$

$$\therefore u = -\frac{2\pi}{n} \cos n\pi e^{-n^2 t}$$

Applying the inversion formula for finite Fourier sine transform, we have

$$U(x, t) = \frac{2}{\pi} \sum_{n=1}^\infty \left(-\frac{2\pi}{n} \cos n\pi e^{-n^2 t} \right) \sin nx.$$

For physical interpretation, $U(x, t)$ may be regarded as the temperature at any point x at an instant of time t in a solid bounded by the planes $x = 0$ and $x = \pi$. The boundary conditions $U(0, t) = 0$ and $U(\pi, t) = 0$ give the zero temperature at the ends while $U(x, 0) = 2x$ represents that the initial temperature is a function of x .

Problem 32. Determine the displacements $Y(x, t)$ in a horizontal string stretched from the origin to the point $(\pi, 0)$ when the motion is due to the weight of the string alone. The string may be taken to be initially at rest in the position $Y = 0$.

Taking the axes of reference as shown in Fig. 9.8, and the string being released from the rest in position $Y = 0$, the boundary value problem is

$$\frac{\partial^2 Y}{\partial t^2} = a^2 \frac{\partial^2 Y}{\partial x^2} + g \quad \dots (1)$$

where $0 < x < \pi$, $t > 0$ and $a^2 = \frac{\text{tension}}{\text{linear density}}$ with the boundary conditions.




Fig. 9.8

$$Y=0=\frac{\partial Y}{\partial t} \text{ when } t=0 \text{ and } Y=0 \text{ when } x=0 \text{ or } \pi. \quad \dots (2)$$

Taking finite Fourier sine transform of both sides of (1), we get

$$\int_0^\pi \frac{\partial^2 Y}{\partial t^2} \sin nx \, dx = a^2 \int_0^\pi \frac{\partial^2 Y}{\partial x^2} \sin nx \, dx + 2 \int_0^\pi \sin nx \, dS \quad \dots (3)$$

Suppose that $y(n, t) = \int_0^\pi Y(x, t) \sin nx \, dx$, then

$$\begin{aligned} \frac{d^2 y}{dt^2} &= -a^2 n^2 y + g \left[\frac{1+(-1)^{n+1}}{n} \right] \text{ by (37) of § 9.10 [E]} \\ &= -a^2 n^2 y + g \left[\frac{1-(-1)^n}{n} \right] \quad \dots (4) \end{aligned}$$

With transformed conditions (2) as $y = \frac{dy}{dt} = 0$ when $t=0$.

(4) may be written as

$$(D^2 + a^2 n^2) y = \frac{g}{n} [1 - (-1)^n] \text{ where } D \equiv \frac{d}{dt}.$$

Its complementary function is $A \cos nat + B \sin nat$

and particular integral is $\frac{g}{n} [1 - (-1)^n] \frac{1}{a^2 n^2} \left[1 + \left(\frac{D}{an} \right)^2 \right], 1$

$$\text{i.e. } \frac{g}{a^2 n^3} [1 - (-1)^n]$$

So the complete integral of (4) is

$$y = A \cos nat + B \sin nat + \frac{g}{a^2 n^3} [1 - (-1)^n]$$

$$\therefore \frac{dy}{dt} = -Ana \sin nat + Bna \cos nat$$

Applying the conditions, $y=0=\frac{dy}{dt}$ when $t=0$, we get

$$A = -\frac{g}{a^2 n^3} [1 - (-1)^n] \text{ and } B=0$$

Hence the solution of (4) is

$$Y = \frac{g}{a^2 n^3} [1 - (-1)^n] (1 - \cos ant.)$$

Using the inversion formula, we find

$$y = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{g}{a^2 n^3} [1 - (-1)^n] (1 - \cos nat) \sin nx = \frac{2g}{\pi a^2} \sum_{n=1}^{\infty} \frac{1 - (-1)^n}{n^3} (1 - \cos nat) \sin nx$$

$$= \frac{2g}{\pi a^2} \sum_{n=1}^{\infty} \frac{x(\pi-x)}{2} (1 - \cos nat) \sin nx, \text{ if we apply}$$

$$f^{-1} \left\{ \frac{1 - (-1)^n}{n^3} \right\} = \frac{x(\pi-x)}{2}$$

Problem 33. Let $V(x, y)$ denote the electrostatic potential in a region bounded by the planes $x=0$, $x=\pi$ and $y=0$ in which there is a uniform distribution of space charge of density $\frac{h}{4\pi}$. If the planes $x=0$ and $y=0$ are kept at potential zero, the plane $x=\pi$ at an other fixed potential $V=1$ and V is finite as $y \rightarrow \infty$, then determine V .

We know that the electrostatic potential function $V(x, y)$ satisfies Poisson's equation which is

$$\begin{aligned} \nabla^2 V &= -4\pi \rho \text{ or } \frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} \\ &= -4\pi \rho \text{ in two dimensions.} \end{aligned}$$

Here $\rho = \frac{h}{4\pi}$, the volume density of charge.

$$\therefore \frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = -h \text{ where}$$

$$0 < x < \pi, y > 0 \quad \dots (1)$$

with the boundary conditions

$$V(0, y) = 0, V(\pi, y) = 1 \text{ for } y > 0$$

$$\text{and } V(x, 0) = 0 \text{ for } 0 < x < \pi$$

$$\text{also } V(x, y) \text{ is finite for } y > 0, 0 \leq x \leq \pi.$$




Fig. 9.9

Taking the finite Fourier sine transform of both sides of (1), we get

$$\int_0^\pi \frac{\partial^2 V}{\partial x^2} \sin nx \, dx + \int_0^\pi \frac{\partial^2 V}{\partial y^2} \sin nx \, dx = -h \int_0^\pi \sin nx \, dx \quad \dots (3)$$

Assuming that $v = \int_0^\pi V \sin nx \, dx$, (3) gives

$$-n^2 v + \frac{d^2 v}{dy^2} = n(-1)^n - h f_s \{1\} \quad \dots (4)$$

since $V=0$ at $x=0$ and $V=1$ at $x=\pi$ so that $v=0$ when $y=0$ and x is finite.

The complete integral of (4) is

$$v = A e^{-ny} + B e^{ny} + \frac{n(-1)^n - h f_s \{1\}}{-n^2}$$

But v is finite for $y \rightarrow \infty$, $\therefore B=0$

$$\text{Also when } y=0, v=0 \therefore A = \frac{n(-1)^n - h f_s \{1\}}{n^2}$$

Hence the solution of (4) is

$$v = \frac{h f_s \{1\} - n(-1)^n}{n^2} (1 - e^{ny})$$

Using the inversion formula, we find

$$V = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{h f_s \{1\} - n(-1)^n}{n^2} \sin nx.$$

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 34. Expand the function $f(t) = kt$ in the interval $-T/2 < t < T/2$ in a Complex Fourier series. Plot the function defined by the series outside this range.

Problem 35. Show that if $f(t) = \phi(-t)$ for $-T/2 < t < 0$ and $f(t) = \phi(t)$ for $0 < t < T/2$ then the real Fourier series for $f(t)$ contains no sine terms.

Problem 36. Show that if $f(t) = -\phi(-t)$ for $-T/2 < t < 0$ and $f(t) = \phi(t)$ for $0 < t < T/2$ then the real Fourier series for $f(t)$ contains no cosine terms.

Problem 37. Expand the function of period 12 defined as follows in the interval $-6 < t < 6$,

$$\begin{aligned} f(t) &= 0 \text{ for } -6 \leq t \leq -3 \\ &= t+3 \text{ for } -3 < t \leq 0 \\ &= 3-t \text{ for } 0 < t \leq 3 \\ &= 0 \text{ for } 3 < t \leq 6 \end{aligned}$$

Plot the function.

Problem 38. Using Fourier cosine integral, show that for $x > 0$,

$$e^{-bx} = \frac{2b}{\pi} \int_0^{\infty} \frac{\cos ux}{u^2 + b^2} dx.$$

Problem 39. (a) State the Dirichlet conditions for the expansion of a function $f(x)$ in a Fourier series. Obtain the Fourier expansion of the function

$$f(x) = \begin{cases} t, & 0 < t < 2 \\ 0, & 2 < t < 4 \end{cases}$$

(b) State the Fourier Integral theorem. Show that if

$$g(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(t) e^{-ixt} dt, \text{ then } f(t) = \sqrt{\frac{1}{2\pi}} \int_{-\infty}^{\infty} g(x) e^{ixt} dx$$

Obtain $f(t)$ for the function $g(x)$ defined by

$$\begin{aligned} g(x) &= 0, \quad x < 0 \\ &= 1, \quad x > 0 \end{aligned}$$

(Bombay, 1963)

Problem 40. Expand the periodic function

$$f(x) = \begin{cases} -1 & \text{for } -\pi < x < 0 \\ 1 & \text{for } 0 < x < \pi \end{cases}$$

in Fourier series.

Problem 41. Obtain a Fourier series expansion of the periodic function $f(t)$, the period of which is T and the form of which within the first period i.e., for $0 \leq t < T$ is given by $f(t) = t(T-t)$

(Agra, 1963)

Obtain the form of the Fourier coefficients for large n to order $\left(\frac{1}{n^2}\right)$.

(Agra, 1967)

Problem 42. Obtain the Fourier series expansions of the following periodic function, each with period T and having the following forms during the first period i.e. $0 \leq t < T$;

$$(i) f(t) = \sin \frac{\pi t}{T}$$

$$(ii) f(t) = \left(\frac{t^2}{T^2}\right)$$

which one has the more rapidly convergent expansion.

(Agra, 1968)

Problem 43. Discuss briefly the role of Fourier's series in mathematical Physics.

(Vikram, 1969)

Problem 44. (a) State the conditions under which a function can be expanded in the form of a Fourier series.

(b) Express the following function in a Fourier series

$$F(x) = x + x^2; \quad -\pi < x < \pi.$$

(Agra, 1969)

Problem 45. (a) State the Dirichlet conditions for the Fourier expansion of a periodic function. From the Fourier series obtain the Fourier Integral formula

$$f(x) = \frac{1}{\pi} \int_0^{\infty} \int_{-\infty}^{\infty} f(\xi) \cos [\alpha (\xi - x)] d\xi d\alpha \quad (-\infty < x < \infty)$$

(b) Let $f(\xi) = 1$ for $0 < \xi < 1$ and $f(\xi) = 0$ for $\xi > 1$

Use the above formula to show that

$$\int_0^{\infty} \frac{\sin \alpha \cos \alpha \xi}{\alpha} d\alpha = \frac{\pi}{4} \text{ when } \xi = 1. \quad (\text{Bombay, 1970})$$

Problem 46. (a) Find the Fourier-cosine-series for the function $f(x) = x$ in the range $0 < x < \pi$.

(b) Using the Fourier-sine-integral evaluate $\int_0^{\infty} \frac{\sin x}{x} dx$. (Agra, 1971)

Problem 47. If V is the Temperature at time t and k the diffusivity of the material, find V from the partial differential equation $\frac{\partial V}{\partial t} - k \frac{\partial^2 V}{\partial x^2}$, $x > 1, t > 0$

With the boundary condition $V = V_0$ for $x = 0, t > 0$

$V = 0$ for $t = 0, x > 0$

$$\left[\text{Ans. } V = V_0 \left[1 - \frac{2}{\pi} \int_0^{\infty} e^{-n^2 kt} \sin nx \cdot \frac{dn}{n} \right] \right]$$

Problem 48. Determine the solution of the equation $\frac{\partial^4 V}{\partial x^4} + \frac{\partial^2 V}{\partial y^2} = 0$, $-\infty < x < \infty$,

$y > 0$ satisfying the conditions

(i) V and its partial derivatives tend to zero as $x \rightarrow \pm \infty$

(ii) $V = F(x), \frac{\partial V}{\partial y} = 0$ when $y = 0$

$$\left[\text{Ans. } V(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(x) \cos(n^2 y) e^{-inx} dn \right]$$

Problem 49. Solve the wave equation $\frac{\partial^2 y}{\partial x^2} = \frac{1}{c^2} \frac{\partial^2 y}{\partial t^2}$ for a stretched string fixed

by a rigid supports at the ends $x = 0$ and $x = l$ if at $t = 0, \frac{\partial y}{\partial t} = 0$ and $y(x, t) = y_0(x)$.

(Agra, 1974)

Problem 50. Find the Fourier transform of the Gaussian Probability function:

$$f(x) = N e^{-\alpha x^2} \quad (N, \alpha \text{ are constants}).$$

(Agra, 1974)

THE LAPLACE TRANSFORMS

10.1. INTRODUCTION

The Laplace transform which is a part of the new-growing topic known as operational calculus is easily and effectively applicable to the boundary value problems of differential equations arising in physics, mathematics and engineering. The subject was mainly originated in the work of Heaviside who found it useful to solve the equation of electro-magnetic theory in the end of nineteenth century.

10.2. DEFINITION OF THE INTEGRAL TRANSFORM

All such transforms as Laplace transform, Fourier-transform and Hankel transform are included in the term Integral transform and we define it as follows:

If there is a known function $K(\alpha, x)$ of two variables α and x such that the integral

$$\int_0^{\infty} K(\alpha, x) F(x) dx \quad \dots (1)$$

is convergent, then the integral (1) is termed as the Integral transform of the function $F(x)$ and is denoted by $\tilde{F}(x)$ or $T\{F(x)\}$ i.e.

$$\tilde{F}(x) = T\{F(x)\} = \int_0^{\infty} K(\alpha, x) F(x) dx \quad \dots (2)$$

The function $K(\alpha, x)$ introduced here is sometimes known as the *Kernel* of the transformation and α is a parameter (real or complex) independent of x .

10.3. DEDUCTION OF THE DEFINITION OF THE LAPLACE TRANSFORM FROM THAT OF THE INTEGRAL TRANSFORM

In §10.2 we have defined the Integral transform of $F(x)$ as

$$T\{F(x)\} = \int_0^{\infty} K(\alpha, x) F(x) dx \quad \dots (1)$$

where $K(\alpha, x)$ is the Kernel of the transformation.

If we take the Kernel,

$$\left. \begin{aligned} K(\alpha, t) = K(s, t) &= 0 \text{ for } t < 0 \\ &= e^{-st} \text{ for } t \geq 0 \end{aligned} \right\} \quad \dots (2)$$

then the transform

$$T\{F(x)\} = \int_0^{\infty} e^{-st} F(t) dt \text{ for } t \geq 0 \quad \dots (3)$$

is known as the *Laplace transform*.

10.4. DEFINITION OF THE LAPLACE TRANSFORM

If $F(t)$ be a function of t defined for all positive values of t (i.e. $t > 0$), then the Laplace transform of $F(t)$ denoted by $L\{F(t)\}$ or $\tilde{F}(s)$ or $f(s)$ is defined by the expression

$$L\{F(t)\} = \tilde{F}(s) = f(s) = \int_0^{\infty} e^{-st} F(t) dt \quad \dots (1)$$

where s is a parameter (real or complex).

If the integral $\int_0^{\infty} e^{-st} F(t) dt$ converges for some value of s , then the Laplace transform of $F(t)$ is said to exist, otherwise it does not exist.

Problem 1. Find the Laplace transform of the following functions:

- (i) $F(t) = 1$
- (ii) $F(t) = t$
- (iii) $F(t) = t^n, n = 0, 1, 2, 3, \dots$

By definition of Laplace transform, we have

$$L\{F(t)\} = \int_0^{\infty} e^{-st} F(t) dt \quad \dots (1)$$

(i) when $F(t) = 1$, (1) becomes

$$L\{1\} = \int_0^{\infty} e^{-st} 1 dt = \left[\frac{e^{-st}}{-s} \right]_0^{\infty} = \frac{1}{s}, s > 0$$

(ii) when $F(t) = t$, (1) gives

$$\begin{aligned} L\{t\} &= \int_0^{\infty} e^{-st} \cdot t dt \\ &= \left[\frac{e^{-st}}{-s} \cdot t \right]_0^{\infty} - \int_0^{\infty} \frac{e^{-st}}{-s} dt, \text{ (integrating by parts).} \\ &= 0 + \frac{1}{s} \int_0^{\infty} e^{-st} dt \\ &= \frac{1}{s} \left[\frac{e^{-st}}{-s} \right]_0^{\infty} = \frac{1}{s^2}, s > 0 \end{aligned}$$

(iii) when $F(t) = t^n$, the transform (1) reduces to

$$\begin{aligned} L\{t^n\} &= \int_0^{\infty} e^{-st} t^n dt \\ &= \left[\frac{e^{-st}}{-s} \cdot t^n \right]_0^{\infty} + \frac{n}{s} \int_0^{\infty} e^{-st} \cdot t^{n-1} dt, \text{ (integrating by parts).} \\ &= 0 + \frac{n}{s} \left[\frac{e^{-st}}{-s} t^{n-1} \right]_0^{\infty} + \frac{n(n-1)}{s^2} \int_0^{\infty} e^{-st} \cdot t^{n-2} dt \\ &\quad \text{integrating by parts} \end{aligned}$$

$$\begin{aligned}
&= \frac{n(n-1)(n-2)}{s^3} \int_0^\infty e^{-st} \cdot t^{n-3} dt \text{ (repeating the process of integration by parts, whence first integral vanishes for both the limits)} \\
&= \dots\dots\dots \\
&= \frac{n(n-1)(n-2)\dots\dots 3 \cdot 2 \cdot 1}{s^n} \int_0^\infty e^{-st} \cdot t^0 dt \text{ where } t^0 = 1 \\
&= \frac{n}{s^n} \cdot \frac{1}{s} = \frac{n}{s^{n+1}}, s > 0 \text{ or } \frac{(n+1)}{s^{n+1}}, s > 0
\end{aligned}$$

Problem 2. Find the Laplace transform of e^{at} .

Here $F(t) = e^{at}$

Hence by definition of Laplace transform, we have

$$\begin{aligned}
L\{e^{at}\} &= \int_0^\infty e^{-st} \cdot e^{at} dt \\
&= \int_0^\infty e^{-(s-a)t} dt \\
&= \left[\frac{e^{-(s-a)t}}{-(s-a)} \right]_0^\infty \\
&= \frac{1}{s-a}, s > a \text{ for which the integral converges,} \\
&\quad \text{otherwise it diverges for } s \leq a.
\end{aligned}$$

Problem 3. Find the Laplace transform of $\sin at$ and $\cos at$.

We have, by definition of Laplace transform

$$L\{F(t)\} = \int_0^\infty e^{-st} F(t) dt$$

when $F(t) = \sin at$, then

$$\begin{aligned}
L\{\sin at\} &= \int_0^\infty e^{-st} \sin at dt \\
&= \left[\frac{e^{-st}}{s^2 + a^2} (-s \sin at - a \cos at) \right]_0^\infty \\
&\quad \left[\because \int e^{ax} \sin bx dx = \frac{e^{ax}}{a^2 + b^2} (a \sin bx - b \cos bx) \right] \\
&= \frac{a}{s^2 + a^2}, s > 0
\end{aligned}$$

Again when $F(t) = \cos at$, then

$$\begin{aligned}
L\{\cos at\} &= \int_0^\infty e^{-st} \cos at dt \\
&= \left[\frac{e^{-st}}{s^2 + a^2} (-s \cos at + a \sin at) \right]_0^\infty \\
&\quad \left[\because \int e^{ax} \cos bx dx = \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx) \right]
\end{aligned}$$

$$= \frac{s}{s^2 + a^2}, s > 0$$

Problem 4. Find the Laplace transform of $\sinh at$ and $\cosh at$.

We have, by definition of Laplace transform

$$\begin{aligned}
L\{\sinh at\} &= \int_0^\infty e^{-st} \sinh at dt \\
&= \frac{1}{2} \int_0^\infty e^{-st} (e^{at} - e^{-at}) dt \\
&= \frac{1}{2} \left[\int_0^\infty e^{-st} \cdot e^{at} dt - \int_0^\infty e^{-st} \cdot e^{-at} dt \right] \\
&= \frac{1}{2} \left[\frac{1}{s-a} - \frac{1}{s+a} \right] \text{ by Problem 2} \\
&= \frac{a}{s^2 - a^2}, s > |a| \\
\text{and } L\{\cosh at\} &= \int_0^\infty e^{-st} \cosh at dt \\
&= \frac{1}{2} \int_0^\infty e^{-st} (e^{at} + e^{-at}) dt \\
&= \frac{1}{2} \left[\frac{1}{s-a} + \frac{1}{s+a} \right] \text{ by Problem 2} \\
&= \frac{s}{s^2 - a^2}, s > |a|
\end{aligned}$$

Problem 5. Find the Laplace transform of the following functions:

(i) $F(t) = t \sin at$

(ii) $F(t) = t \cos at$

By definition, the Laplace transform of a function $F(t)$ is given by

$$L\{F(t)\} = \int_0^\infty e^{-st} F(t) dt$$

(i) when $F(t) = t \sin at$, we have

$$\begin{aligned}
L\{t \sin at\} &= \int_0^\infty e^{-st} \cdot t \sin at dt \\
&= \left[t \cdot \frac{e^{-st}}{s^2 + a^2} (-s \sin at - a \cos at) \right]_0^\infty \\
&\quad - \int_0^\infty \frac{e^{-st}}{s^2 + a^2} (-s \sin at - a \cos at) dt \\
&\quad [\text{on integrating by parts, treating } t \text{ as first function and } e^{-st} \sin at \text{ as the second function and using the result} \\
&\quad \int e^{ax} \sin bx dx = \frac{e^{ax}}{a^2 + b^2} (a \sin bx - b \cos bx), \text{ while} \\
&\quad \text{the first integral vanishes for both the limits}] \\
&= \frac{s}{s^2 + a^2} \int_0^\infty e^{-st} \sin at dt + \frac{a}{s^2 + a^2} \int_0^\infty e^{-st} \cos at dt
\end{aligned}$$

$$= \frac{s}{s^2+a^2} \cdot \frac{a}{s^2+a^2} + \frac{a}{s^2+a^2} \cdot \frac{s}{s^2+a^2} \text{ by Problem 3 for } s>0$$

$$= \frac{2as}{(s^2+a^2)^2}, s>0.$$

(ii) when $F(t)=t \cos at$, we have

$$L\{t \cos at\} = \int_0^\infty e^{-st} \cdot t \cos at \, dt$$

On R.H.S., integrating by parts treating t as first function and $e^{-st} \cos at$ as second function and using the result $\int e^{ax} \cos bx \, dx = \frac{e^{ax}}{a^2+b^2} (a \cos bx + b \sin bx)$, while the first integral vanishes for both the limits, we are left with

$$L\{t \cos at\} = \frac{s}{s^2+a^2} \int_0^\infty e^{-st} \cos at \, dt - \frac{a}{s^2+a^2} \int_0^\infty e^{-st} \sin at \, dt$$

(as in (i))

$$= \frac{s}{s^2+a^2} \cdot \frac{s}{s^2+a^2} - \frac{a}{s^2+a^2} \cdot \frac{a}{s^2+a^2} \text{ by Problem 3 for } s>0$$

$$= \frac{s^2-a^2}{(s^2+a^2)^2}, s>0.$$

Problem 6. Find the Laplace transform of t^a , where a is positive but not necessarily an integer.

Hint: Proceed just like in Problem 1 (iii) and get the result $\frac{\Gamma(a+1)}{s^{a+1}}$ $s>0$ since if a is not an integer, then $\Gamma(a)$ is not defined.

Problem 7. Find the Laplace transform of e^{-at}

Ans. $\frac{1}{s+a}$ (replace a by $-a$ in Problem 2)

Problem 8. Find the Laplace transform of the following functions.

(i) $F(t)=e^{at} \sin bt$ Ans. $b/\{(s-a)^2+b^2\}$

(ii) $F(t)=e^{at} \cos bt$ Ans. $(s-a)/\{(s-a)^2+b^2\}$

Hint. $L\{e^{at} \sin bt\} = \int_0^\infty e^{-st} \cdot e^{at} \sin bt \, dt$

$$= \int_0^\infty e^{-(s-a)t} \sin bt \, dt$$

$$= \left[\frac{e^{-(s-a)t}}{(s-a)^2+b^2} \left\{ -(s-a) \sin bt - b \cos bt \right\} \right]_0^\infty$$

$$= \frac{b}{(s-a)^2+b^2} \text{ etc.}$$

Note. The results so far derived along with a few more can be tabulated as following:

$F(t)$	$L\{F(t)\}=f(s)=\tilde{F}(s)$
1	$1/s, s>0$
t	$1/s^2, s>0$
$t^n, n=0, 1, 2, \dots$	$n!/s^{n+1} \text{ or } \Gamma(n+1)/s^{n+1}, s>0$
$t^a, a>0 \text{ but not necessarily an integer}$	$\Gamma(a+1)/s^{a+1}, a>0$
e^{at}, e^{-at}	$\frac{1}{s-a}; \frac{1}{s+a}, s>a$
$\sin at$	$a/(s^2+a^2), s>0$
$\cos at$	$s/(s^2+a^2), s>0$
$\sinh at$	$a/(s^2-a^2), s>0$
$\cosh at$	$s/(s^2-a^2), s>0$
$t \sin at$	$2as/(s^2+a^2)^2, s>0$
$t \cos at$	$(s^2-a^2)/(s^2+a^2)^2$
$e^{at} \sin bt$	$b/\{(s-a)^2+b^2\}$
$e^{at} \cos bt$	$(s-a)/\{(s-a)^2+b^2\}$
$t^{n-1} e^{at}, n>0$	$n!/(s-a)^n$
$J_0(at) \text{ and } t J_0(at)$	$1/(s^2+a^2)^{1/2} \text{ and } s/(s^2+a^2)^{3/2}$

10.5. FUNCTIONS OF EXPONENTIAL ORDER

A function $F(t)$ is said to be the function of exponential order m as $t \rightarrow \infty$, when for a given positive integer N , there exists real constants $M>0$ and m such that

$$|e^{-mt} F(t)| < M$$

$$\text{or } |F(t)| < M e^{mt} \text{ for } t > N$$

As an illustrative example the function $F(t) = t^2$ is of exponential order 3 since $|t^2| = t^2 < e^{3t}$ for all $t > 0$ because

$$\begin{aligned} \lim_{t \rightarrow \infty} e^{-mt} t^2 &= \lim_{t \rightarrow \infty} \frac{t^2}{e^{mt}} \\ &= \lim_{t \rightarrow \infty} \frac{t^2}{1 + mt + \frac{m^2 t^2}{2} + \frac{m^3 t^3}{6} + \dots} \\ &= \lim_{t \rightarrow \infty} \frac{1}{\frac{1}{t^2} + \frac{m}{t} + \frac{m^2}{2} + \frac{m^3 t}{6} + \dots} \\ &= 0. \end{aligned}$$

Problem 9. Show that function $t^3 [=F(t)]$ is of exponential as $t \rightarrow \infty$. Take m as some fixed positive value, and consider

$$\begin{aligned} \lim_{t \rightarrow \infty} (e^{-mt} t^3) &= \lim_{t \rightarrow \infty} \frac{t^3}{e^{mt}} \\ &= \lim_{t \rightarrow \infty} \frac{3t^2}{m e^{mt}} \quad \text{on differentiating numerator and denominator w.r.t. 't' by L' Hospital's rule} \\ &= \lim_{t \rightarrow \infty} \frac{6t}{m^2 e^{mt}} \quad (\text{by L' Hospitals' rule}) \\ &= \lim_{t \rightarrow \infty} \frac{6}{m^3 e^{mt}} \quad (\quad , \quad , \quad) \\ &= 0. \end{aligned}$$

This follows that $|t^3| < e^{mt}$ for all $t > 0$

$\therefore t^3$ is of exponential order given by

$$t^3 = O(e^{mt}), \quad t \rightarrow \infty \text{ for any fixed positive value of } m.$$

Note. Here the notation 'O' stands for 'of the order of'.

Problem 10. Show that the function $F(t) = e^{t^3}$ is not of exponential order as $t \rightarrow \infty$.

Taking m as some fixed positive value, we have

$$\begin{aligned} \lim_{t \rightarrow \infty} e^{-mt} F(t) &= \lim_{t \rightarrow \infty} e^{-mt} e^{t^3} \\ &= \lim_{t \rightarrow \infty} e^{t(t^2 - m)} \\ &= \infty \text{ for all values of } m. \end{aligned}$$

Hence it is not possible to find a number M such that $e^{t^3} < M e^{mt}$ for all values of m .

Thus the given function is not of exponential order as $t \rightarrow \infty$.

Problem 11. Show that the function $F(t) = t^n$ for $n = 1, 2, 3, \dots$ is of exponential order as $t \rightarrow \infty$.

Hint. Repeated application of L' Hospital's rule gives

$$\lim_{t \rightarrow \infty} e^{-mt} t^n = \lim_{t \rightarrow \infty} \frac{t^n}{e^{mt}} = \lim_{t \rightarrow \infty} \frac{n}{m^n e^{mt}} = 0$$

and so $t^n = O(e^{mt}), t \rightarrow \infty$ etc...

Problem 12. Show that the function $F(t) = e^{t^2}$ is not of exponential order as $t \rightarrow \infty$.

Note 2. Actually, functions of exponential order do not grow in absolute value more rapidly than $M e^{mt}$ as t increases but in practice there is no such restriction since M and m may be as large as we desire. For example the bounded functions like $\sin at$ or $\cos at$ are of exponential order.

10.6. PIECEWISE OR SECTIONAL CONTINUITY

Given a closed interval $[a, b]$ a function $F(t)$ is called **piecewise continuous** or **sectionally continuous** in $[a, b]$, i.e. $a \leq t \leq b$, if the interval can be divided into a finite number of sub-intervals such that in each of these sub intervals, the function remains continuous and possesses finite right and left hand limits.

In other words, the function $F(t)$ is sectionally or piecewise continuous in the closed interval $[a, b]$, if the closed interval $[a, b]$ can be divided into a finite number of sub-intervals $c \leq t \leq d$ such that

- $F(t)$ is continuous in the open interval $c < t < d$,
- $F(t)$ approaches a limit as t approaches each end-point within interval i.e.

$$\lim_{t \rightarrow c+0} F(t) \text{ and } \lim_{t \rightarrow d-0} F(t) \text{ both exist.}$$

For example, referring to the Fig. 10.1 it is evident that the func-




Fig. 10.1

tion $F(t)$ is continuous in the open intervals (a, t_1) , (t_1, t_2) , (t_2, t_3) etc., i.e. $a < t < t_1$; $t_1 < t < t_2$; $t_2 < t < t_3$ etc,

The right hand and left hand limits at t , exist and are given by

$$\lim_{\epsilon \rightarrow 0} F(t_1 + \epsilon) = F(t_1 + 0) = F(t_1 +)$$

$$\text{and } \lim_{\epsilon \rightarrow 0} F(t_1 - \epsilon) = F(t_1 - 0) = F(t_1 -)$$

where ϵ is positive.

Similar is the case for points t_2 , t_3 and t_4 .

10.7. A FUNCTION OF CLASS A

If a function $F(t)$ is sectionally continuous over every finite interval in the range $t \geq 0$ and is of exponential order as $t \rightarrow \infty$, then the function is termed as 'a function of class A'.

10.8. SUFFICIENT CONDITIONS FOR THE EXISTENCE OF LAPLACE TRANSFORMS

If a function $F(t)$ is piecewise or sectionally continuous in every finite interval $0 \leq t \leq N$ and is of exponential order m for $t > N$ then its Laplace transform $L\{F(t)\}$ i.e. $f(s)$ exists for all $s > m$.

or

If a function $F(t)$ is of class A, then the Laplace transform of $F(t)$ i.e., $L\{F(t)\}$ exists.

or

If a function $F(t)$ is sectionally continuous on every finite interval in the range $t \geq 0$ and satisfies the condition

$$|F(t)| \leq Me^{mt}$$

for all $t \geq 0$ and for some constants m and M , then $L\{F(t)\}$ i.e. $f(s)$ exists for all $s > m$.

We have, for any positive integer N ,

$$\begin{aligned} L\{F(t)\} &= \int_0^\infty e^{-st} F(t) dt \\ &= \int_0^N e^{-st} F(t) dt + \int_N^\infty e^{-st} F(t) dt \\ &= I_1 + I_2 \text{ (say)} \end{aligned} \quad \dots (1)$$

Since $F(t)$ is sectionally continuous in every finite interval $0 \leq t \leq N$, it therefore, follows that the integral I_1 exists.

Also $F(t)$ being of exponential order m for $t > N$, I_2 exists, since

$$\begin{aligned} |I_2| &= \left| \int_N^\infty e^{-st} F(t) dt \right| \leq \int_N^\infty |e^{-st} F(t)| dt \\ &\leq \int_0^\infty e^{-st} |F(t)| dt \\ &\leq \int_0^\infty e^{-st} M e^{mt} dt \text{ as } F(t) \text{ is of exponential order } m \text{ and so} \\ &\quad |F(t)| \leq M e^{mt} \end{aligned}$$

$$\leq \int_0^\infty e^{-(s-m)t} M dt$$

$$\therefore |I_2| \leq \frac{M}{s-m} \text{ for } s > m. \text{ (Here R.H.S.} \rightarrow 0 \text{ as } s \rightarrow \infty)$$

COROLLARY 1. From (1), it therefore follows that $L\{F(t)\}$ exists for $s > m$ (2)

If the function $F(t)$ is real valued in $L\{F(t)\} = \int_0^\infty e^{-st} F(t) dt = f(s)$ such that (i) $F(t)$ is R-integrable (i.e. has Riemann-integral) and (ii) $F(t)$ satisfies, $|F(t)| \leq M e^{m_0 t}$ for all $t > N$; M , m_0 , N being positive constants,

then $\int_0^\infty e^{-st} F(t) dt$ is convergent (absolutely) for $m > m_0$, s being a complex number.

Supposing, $s = m + in$, we have

$$\begin{aligned} |e^{-st} F(t)| &= |e^{-(m+in)t} F(t)| \\ &= e^{-mt} |F(t)| \because |e^{-nit}| = 1 \text{ for every } t \geq N \\ &\leq e^{-mt} M e^{m_0 t} \\ &\leq M e^{-(m-m_0)t} \end{aligned}$$

Hence, if the integral $\int_0^\infty e^{-(m-m_0)t} dt$ is convergent, then

$$\int_0^\infty e^{-st} F(t) dt \text{ for } m > m_0, \text{ is absolutely convergent} \quad \dots (3)$$

COROLLARY 2. $L\{F(t)\} = f(s) = \int_0^\infty e^{-st} F(t) dt$ converges, for all s greater than some fixed value s_0 of s .

In corollary 1, if we replace m_0 by s_0 and m by s then the proposition follows.

COROLLARY 3. If the function $F(t)$ is of class A and $L\{F(t)\} = f(s)$, then $\lim_{s \rightarrow \infty} f(s) = 0$

The function $F(t)$ being of class A is bounded over the range $0 \leq t \leq N$, so that $|F(t)| < M_1$ (say) for $m \leq t \leq N$.

Also $F(t)$ being of exponential order m , we have

$$|F(t)| < M_2 e^{mt} \text{ for } t \geq N$$

Let us assume that $M = \text{Max. } \{M_1, M_2\}$

and $p = \text{Max. } \{m, 0\}$

Thus, $|F(t)| < M e^{pt}$ for $t > 0$

$$\text{i.e. } \int_0^\infty e^{-st} F(t) dt < M \int_0^\infty e^{-st} e^{pt} dt = M \int_0^\infty e^{-(s-p)t} dt \\ = \frac{M}{s-p}, \quad s > p$$

Here $\frac{M}{s-p} \rightarrow 0$ as $s \rightarrow \infty$.

$$\text{Hence } \lim_{s \rightarrow \infty} \int_0^\infty e^{-st} F(t) dt = 0$$

$$\text{i.e. } \lim_{s \rightarrow \infty} f(s) = 0.$$

... (4)

Note. The conditions mentioned in this article for the existence of Laplace transforms are only sufficient but not necessary as is evident from the following example:

$$\text{Let } F(t) = \frac{1}{\sqrt{t}} = t^{-1/2}$$

Obviously the function $F(t) = t^{-1/2}$ is not sectionally continuous in every finite interval in the range $t \geq 0$, for

$$F(t) \rightarrow \infty \text{ as } t \rightarrow 0.$$

But $t^{-1/2}$ is integrable from 0 to any positive value N .

Also $t^{-1/2}$ is of exponential order, since

$$|F(t)| \text{ i.e. } |t^{-1/2}| < M e^{mt} \text{ for all } t > 0 \text{ with } M=1 \text{ and } m=0$$

As such the Laplace transform of $t^{-1/2}$ exists and may be evaluated as follows:

$$L\{F(t)\} = L\{t^{-1/2}\} = \int_0^\infty e^{-st} t^{-1/2} dt \text{ for } s > 0,$$

$$= 2 \int_0^\infty e^{-sx^2} dx, \quad s > 0$$

$$\therefore -\frac{1}{2} t^{-1/2} dt = dx$$

Again put $s^{1/2}x = y$ so that $s^{1/2} dx = dy$

$$\therefore L\{t^{-1/2}\} = 2s^{-1/2} \int_0^\infty e^{-y^2} dy, \quad s > 0$$

$$= 2s^{-1/2} \cdot \frac{1}{2} \sqrt{\pi}$$

$$\therefore \int_0^\infty e^{-y^2} dy = \frac{1}{2} \sqrt{\pi}$$

$$= \sqrt{\frac{\pi}{s}} \text{ for } s > 0, \text{ even if } t^{-1/2} \rightarrow \infty \text{ as } t \rightarrow 0$$

the Laplace transform of $t^{-1/2}$ exists even if the function is not sectionally continuous.

Alternatively, the integral $\int_0^\infty e^{-st} t^{-1/2} dt, s > 0$ can be evaluated by a

substitution $\sqrt{st} = x$ i.e. $\frac{dt}{\sqrt{t}} = \frac{2}{\sqrt{s}} dx$ whence we have

$$L\{t^{-1/2}\} = \frac{2}{\sqrt{s}} \cdot \int_0^\infty e^{-x^2} dx = \frac{2}{\sqrt{s}} \cdot \frac{\sqrt{\pi}}{2} = \sqrt{\frac{\pi}{s}} \quad \dots (5)$$

Problem 13. Show that the Laplace transform of the function

$$F(t) = t^n, \quad 0 > n > -1$$

exists, although it is not a function of the class A.

Here the function $F(t) = t^n$ is not sectionally continuous and so is not of the class A, in every finite interval in the range $t \geq 0$, since $t^n \rightarrow 0$ as $t \rightarrow 0$ for $0 > n > -1$

Hence the function has infinite discontinuity at $t=0$

$$\text{Also, } \lim_{t \rightarrow \infty} \{e^{-mt} F(t)\} = \lim_{t \rightarrow \infty} \left(\frac{t^n}{e^{mt}} \right)$$

$$= \lim_{t \rightarrow \infty} \left(\frac{nt^{n-1}}{me^{mt}} \right) \text{ by L' Hospital rule}$$

$$= \lim_{t \rightarrow \infty} \frac{n^2 t^{n-2}}{m^2 e^{mt}}$$

$$= \dots \dots \dots$$

$$= \lim_{t \rightarrow \infty} \frac{|n|}{m^n e^{mt}} \text{ by repeated application of L' Hospital rule}$$

$$= 0$$

which follows that $F(t) = t^n$ is of exponential order.

[Or this may be argued thus, $t^n \rightarrow 0$ as $t \rightarrow \infty$ so that t^n is of exponential order with $M=1$ and $m=0$, for, $t^n < M e^{mt}$].

Moreover, the function t^n ($0 > n > -1$) is integrable from 0 to any positive number N .

$$\text{Further, } L\{F(t)\} = \int_0^\infty e^{-st} t^n dt \quad \text{Put } st = x$$

$$= \int_0^\infty e^{-x} \left(\frac{x}{s} \right)^n \frac{dx}{s} \quad \text{i.e. } dt = \frac{dx}{s}$$

$$= \frac{1}{s^{n+1}} \int_0^\infty x^n e^{-x} dx$$

$$= \frac{\Gamma(n+1)}{s^{n+1}} \text{ by the definition of gamma function.}$$

Hence the Laplace transform of t^n exists even if it is not a function of the class A.

10.9. SOME PROPERTIES OF LAPLACE TRANSFORMS

[A] Linearity Property

A Laplace transform $L\{F(t)\}$ is said to be linear if for every pair of function $F_1(t)$ and $F_2(t)$ and for every pair of constants C_1 and C_2 , we have

$$L\{C_1 F_1(t) + C_2 F_2(t)\} = C_1 L\{F_1(t)\} + C_2 L\{F_2(t)\} \\ = C_1 f_1(s) + C_2 f_2(s)$$

where $f_1(s)$ and $f_2(s)$ are linear transforms of $F_1(t)$ and $F_2(t)$ respectively

$$\text{We have } L\{F_1(t)\} = f_1(s) = \int_0^{\infty} e^{-st} F_1(t) dt$$

$$\text{and } L\{F_2(t)\} = f_2(s) = \int_0^{\infty} e^{-st} F_2(t) dt$$

$$\text{So that } L\{C_1 F_1(t)\} = C_1 f_1(s) = \int_0^{\infty} e^{-st} C_1 F_1(t) dt = C_1 L\{F_1(t)\}$$

$$\text{and } L\{C_2 F_2(t)\} = C_2 f_2(s) = \int_0^{\infty} e^{-st} C_2 F_2(t) dt = C_2 L\{F_2(t)\}$$

$$\begin{aligned} \therefore L\{C_1 F_1(t) + C_2 F_2(t)\} &= \int_0^{\infty} e^{-st} \{C_1 F_1(t) + C_2 F_2(t)\} dt \text{ by definition} \\ &= \int_0^{\infty} e^{-st} C_1 F_1(t) dt + \int_0^{\infty} e^{-st} C_2 F_2(t) dt \\ &= C_1 L\{F_1(t)\} + C_2 L\{F_2(t)\} \\ &= C_1 f_1(s) + C_2 f_2(s) \end{aligned} \quad \dots (1)$$

The result may be generalized for any number of functions and for the same number of arbitrary constants i.e.,

$$L\left\{\sum_{r=1}^n C_r F_r(t)\right\} = \sum_{r=1}^n C_r L\{F_r(t)\} \quad \dots (2)$$

Problem 14. Find Laplace Transform of $4e^{5t} + 6t^3 - 4 \cos 3t + 3 \sin 4t$.

Applying the linearity property, we have

$$\begin{aligned} L\{4e^{5t} + 6t^3 - 4 \cos 3t + 3 \sin 4t\} &= 4L\{e^{5t}\} + 6L\{t^3\} - 4L\{\cos 3t\} + 3L\{\sin 4t\} \\ &= 4\left(\frac{1}{s-5}\right) + 6\left(\frac{3!}{s^4}\right) - 4\left(\frac{s}{s^2+9}\right) + 3\left(\frac{4}{s^2+16}\right) \\ &= \frac{4}{s-5} + \frac{36}{s^4} - \frac{4s}{s^2+9} + \frac{12}{s^2+16} \end{aligned}$$

[B] First Translation (or Shifting) Property

If $f(s)$ be the Laplace transform of $F(t)$, then the Laplace transform of $e^{at} F(t)$ is $f(s-a)$, where a is any real or complex number i.e., if $L\{F(t)\} = f(s)$, then $L\{e^{at} F(t)\} = f(s-a)$.

$$\text{Given, } L\{F(t)\} = \int_0^{\infty} e^{-st} F(t) dt = f(s)$$

$$\begin{aligned} \therefore L\{e^{at} F(t)\} &= \int_0^{\infty} e^{-st} e^{at} F(t) dt \\ &= \int_0^{\infty} e^{-(s-a)t} F(t) dt \\ &= \int_0^{\infty} e^{-ut} F(t) dt \text{ by putting } u = s-a \end{aligned}$$

$$= f(u)$$

$$= f(s-a)$$

$\dots (3)$

Problem 15. Find the Laplace transform of $e^{-2t} \sin 3t$.

$$\text{We have } L\{\sin 3t\} = \frac{3}{s^2+9}$$

$$\therefore L\{e^{-2t} \sin 3t\} = \frac{3}{(s+2)^2+9} = \frac{3}{s^2+4s+13}$$

[C] Second Translation (or Shifting) Property

$$\text{If } L\{F(t)\} = f(s) \text{ and } G(t) = \begin{cases} F(t-a), & t > a \\ 0, & t < a \end{cases}$$

$$\text{Then } L\{G(t)\} = e^{-as} f(s).$$

$$\begin{aligned} \text{We have } L\{G(t)\} &= \int_0^{\infty} e^{-st} G(t) dt \\ &= \int_0^a e^{-st} G(t) dt + \int_a^{\infty} e^{-st} G(t) dt \\ &= \int_0^a e^{-st} \cdot 0 dt + \int_a^{\infty} e^{-st} F(t-a) dt \\ &= \int_a^{\infty} e^{-st} F(t-a) dt \\ &= \int_0^{\infty} e^{-s(u+a)} F(u) du, \text{ by taking } u = t-a \text{ i.e. } du = dt. \\ &\quad \text{when } t = a, u = 0 \text{ and when } t = \infty, u = \infty. \\ &= e^{-sa} \int_0^{\infty} e^{-su} F(u) du \\ &= e^{-sa} f(s) \end{aligned} \quad \dots (4)$$

Problem 16. Find the Laplace transform of $F(t)$, where

$$F(t) = \begin{cases} \cos\left(t - \frac{2\pi}{3}\right), & t > \frac{2\pi}{3} \\ 0, & t < \frac{2\pi}{3} \end{cases}$$

$$\begin{aligned} \text{We have, } L\{F(t)\} &= \int_0^{\infty} e^{-st} F(t) dt \\ &= \int_0^{2\pi/3} e^{-st} \cdot 0 dt + \int_{2\pi/3}^{\infty} e^{-st} \cos\left(t - \frac{2\pi}{3}\right) dt \\ &= \int_0^{\infty} e^{-s\left(u + \frac{2\pi}{3}\right)} \cos u du \quad \text{by taking } u = t - \frac{2\pi}{3} \\ &= e^{-\frac{2\pi s}{3}} L\{\cos u\} \\ &= e^{-\frac{2\pi s}{3}} \frac{s}{s^2+1} \end{aligned}$$

[D] The Change of Scalar Property

If $L\{F(t)\}=f(s)$, then $L\{F(at)\}=\frac{1}{a}f\left(\frac{s}{a}\right)$... (5)

we have $L\{F(t)\}=\int_0^\infty e^{-st}F(t) dt=f(s)$

$$\begin{aligned}\therefore L\{F(at)\} &= \int_0^\infty e^{-st}F(at) dt && \text{(on replacing } t \text{ by } at) \\ &= \int_0^\infty e^{-su/a}F(u)\frac{du}{a} && \text{by taking } at=u \\ &= \frac{1}{a}\int_0^\infty e^{-pu}F(u) du && \text{where } p=\frac{s}{a}. \\ &= \frac{1}{a}\int_0^\infty e^{-pu}F(u) du && \text{(replacing } u \text{ by } t) \\ &= \frac{1}{a}f(p) \\ &= \frac{1}{a}f\left(\frac{s}{a}\right) \quad \therefore p=\frac{s}{a}.\end{aligned}$$

Problem 17. Find the Laplace transform of $\cos 5t$.

We have, $L\{\cos t\}=\frac{s}{s^2+1}$, $s>0$

$$\therefore L\{\cos 5t\}=\frac{1}{5}\frac{s/5}{(s/5)^2+1}=\frac{s}{s^2+25}.$$

[E] Laplace Transform of Derivatives

If $F(t)$ is continuous for $t\geq 0$ and of exponential order as $t\rightarrow\infty$ while $F'(t)$ is sectionally continuous i.e., $F'(t)$ is of class A for $t\geq 0$, and if $L\{F(t)\}=f(s)$, then $L\{F'(t)\}=sf(s)-F(0)$.

In general if $L\{F(t)\}=f(s)$ and $F(t)$, $F'(t)$, $F''(t)$, ..., $F^{(n-1)}(t)$ are sectionally continuous for $t\geq 0$ and of exponential order as $t\rightarrow\infty$ while $F^{(n)}(t)$ is

$$\begin{aligned}L\{F^{(n)}(t)\} &= s^n f(s) - \sum_{r=0}^{n-1} s^{n-1-r} F^{(r)}(0) \\ &= s^n f(s) - s^{n-1}F(0) - s^{n-2}F'(0) \dots - sF^{(n-2)}(0) - F^{(n-1)}(0)\end{aligned}$$

$$\text{Since } L\{F(t)\}=f(s)=\int_0^\infty e^{-st}F(t) dt$$

$$\begin{aligned}\therefore L\{F'(t)\} &= \int_0^\infty e^{-st}F'(t) dt = \left[e^{-st}F(t) \right]_0^\infty + s \int_0^\infty e^{-st}F(t) dt, \text{ integrating by parts} \\ &= 0 - F(0) + s f(s) \\ &= s f(s) - F(0)\end{aligned}$$

Applying the result (6), we have

$$L\{F''(t)\}=sL\{F'(t)\}-F'(0)$$

... (6)

$$\begin{aligned}&= s\{s f(s) - F(0)\} - F'(0) \text{ by (6)} \\ &= s^2 f(s) - s F(0) - F'(0) \quad \dots (7)\end{aligned}$$

$$\text{Similarly } L\{F''(t)\}=s^2 f(s) - s^2 F(0) - s F'(0) - F''(0) \quad \dots (8)$$

Generalizing it, we find

$$\begin{aligned}L\{F^{(n)}(t)\} &= s^n f(s) - s^{n-1}F(0) - s^{n-2}F'(0) \dots - sF^{(n-2)}(0) - F^{(n-1)}(0) \\ &= s^n f(s) - \sum_{r=0}^{n-1} s^{n-1-r} F^{(r)}(0) \quad \dots (9)\end{aligned}$$

Problem 18. Find the Laplace transform of $F'(t)$ when $F(t)=e^{3t}$.

Given $F(t)=e^{3t}$, $\therefore F(0)=1$ and $F'(t)=3e^{3t}$

As such $L\{e^{3t}\}=sL\{e^{3t}\}-1$ by (6) above

$$= \frac{s}{s-3} - 1 = \frac{3}{s-3}.$$

$$\text{Aliter. } L\{F'(t)\}=L\{3e^{3t}\}=3L\{e^{3t}\}=\frac{3}{s-3}.$$

[F] Derivatives of Laplace Transforms

If the function $F(t)$ is sectionally continuous for $t\geq 0$ and if $L\{F(t)\}=f(s)$, then $f'(s)=L\{-tF(t)\}$

$$\text{We have } f(s)=\int_0^\infty e^{-st}F(t) dt.$$

Differentiating either side w.r.t. 's' we get

$$\begin{aligned}f'(s) &= \int_0^\infty (-t) e^{-st} F(t) dt = \int_0^\infty e^{-st} \{-tF(t)\} dt \\ &= L\{-tF(t)\} \quad \dots (10)\end{aligned}$$

In general if $F(t)$ is sectionally continuous for $t\geq 0$ and if $L\{F(t)\}=f(s)$, then

$$f^{(n)}(s)=L\{(-t)^n F(t)\} \quad \dots (11)$$

where $f^{(n)}(s)=\frac{d^n}{ds^n}f(s)$ for all integral values of n .

We may state it as

$$L\{t^n F(t)\}=(-1)^n f^{(n)}(s)=(-1)^n \frac{d^n}{ds^n}f(s) \quad \dots (12)$$

Problem 19. Find the Laplace transform of $t^3 e^t$.

$$\text{Since } L\{e^t\}=f(s)=\frac{1}{s-a}$$

$$\begin{aligned}\therefore L\{t^3 e^t\} &= (-1)^3 \frac{d^3}{ds^3} \left(\frac{1}{s-a} \right) = (-1)^3 = \frac{(-1)(-2)(-3)}{(s-1)^4} \\ &= \frac{6}{(s-1)^4}\end{aligned}$$

[G] Laplace Transform of Integrals

$$(i) \text{ If } L\{F(t)\}=f(s), \text{ then } L\left\{\int_0^t F(u) du\right\}=\frac{f(s)}{s} \quad \dots (13)$$

$$\text{Let } G(t) = \int_0^t F(u) du$$

$$\text{Then } G'(t) = \frac{d}{dt} \left[\int_0^t F(u) du \right] = F(t) \text{ and } G(0) = \int_0^0 F(u) du = 0.$$

Applying the property [E], we have

$$L\{G'(t)\} = sL\{G(t)\} - G(0)$$

$$\text{i.e. } L\{F(t)\} = sL\{G(t)\} - 0 \text{ or } f(s) = sL\left\{\int_0^t F(u) du\right\}$$

$$\text{i.e. } L\left\{\int_0^t F(u) du\right\} = \frac{f(s)}{s}$$

$$\text{(ii) If } L\{F(t)\} = f(s) \text{ then } L\left\{\frac{F(t)}{t}\right\} = \int_s^\infty f(u) du \quad \dots (14)$$

$$\text{Let } G(t) = \frac{F(t)}{t}, \text{ so that } F(t) = tG(t)$$

$$\therefore L\{F(t)\} = L\{tG(t)\} \text{ (on taking Laplace transform)}$$

$$= (-1) \frac{d}{ds} L\{G(t)\} \text{ by property [F]}$$

$$\text{i.e. } -f(s) = \frac{d}{ds} L\{G(t)\}$$

Integrating both sides with regard to s , we get

$$-\int_\infty^s f(s) ds = L\{G(t)\}$$

$$\text{i.e., } L\{G(t)\} = \int_s^\infty f(u) du, \text{ on the assumption that } \lim_{s \rightarrow \infty} L\{G(s)\} \rightarrow 0.$$

$$\text{Problem 20. Find the Laplace transform of } \int_0^\infty \frac{\sin t}{t} dt.$$

$$\text{We have } L\{\sin t\} = \frac{1}{s^2 + 1} = f(s) \text{ (say)}$$

$$\begin{aligned} \text{and } L\left\{\frac{\sin t}{t}\right\} &= \int_s^\infty \frac{du}{u^2 + 1} \text{ by (14), } \therefore f(u) = \frac{1}{u^2 + 1} \\ &= \left[\tan^{-1} u \right]_s^\infty = \frac{\pi}{2} - \tan^{-1} s \\ &= \cot^{-1} s \therefore \tan^{-1} s + \cot^{-1} s = \frac{\pi}{2} \\ &= \tan^{-1} \frac{1}{s}. \end{aligned}$$

Hence by (13) of property [G],

$$L\left\{\int_0^\infty \frac{\sin t}{t} dt\right\} = \frac{1}{s} \tan^{-1} \frac{1}{s}.$$

[H] Periodic Functions

If $F(t)$ is a periodic function with period $T > 0$, so that

$$F(t+T) = F(t), \text{ then } L\{F(t)\} = \frac{\int_0^T e^{-st} F(t) dt}{1 - e^{-sT}} \quad \dots (15)$$

$$\text{We have } L\{F(t)\} = \int_0^\infty e^{-st} F(t) dt$$

$$= \int_0^T e^{-st} F(t) dt + \int_T^{2T} e^{-st} F(t) dt + \int_{2T}^{3T} e^{-st} F(t) dt + \dots$$

$$= \sum_{n=0}^\infty \int_{nT}^{(n+1)T} e^{-st} F(t) dt$$

If we put $t = u + nT$, then $F(u + nT) = F(u) \therefore F(t + T) = F(t)$ (given)
Thus,

$$L\{F(t)\} = \sum_{n=0}^\infty \int_0^T e^{-s(u+nT)} F(u) du$$

$$\begin{aligned} \text{or } L\{F(t)\} &= \sum_{n=0}^\infty e^{-snT} \int_0^T e^{-su} F(u) du \\ &= \left(1 + e^{-sT} + e^{-2sT} + \dots \right) \int_0^T e^{-su} F(u) du \\ &= \frac{1}{1 - e^{-sT}} \int_0^T e^{-su} F(u) du \therefore \frac{1}{1 - e^{-sT}} \\ &= (1 - e^{-sT})^{-1} = 1 + e^{-sT} + e^{-2sT} + \dots \text{ when } |e^{-sT}| < 1 \\ &= \frac{\int_0^T e^{-st} F(t) dt}{1 - e^{-sT}} \text{ (replacing } u \text{ by } t) \end{aligned}$$

Problem 21. Find the Laplace transform of $F(t)$ when $F(t)$ is a periodic function with period 2π , such that

$$F(t) = \begin{cases} \sin t, & 0 < t < \pi \\ 0, & \pi < t < 2\pi \end{cases}$$

$$\begin{aligned} \text{We have } L\{F(t)\} &= \frac{1}{1 - e^{-2\pi s}} \int_0^{2\pi} e^{-st} F(t) dt \\ &= \frac{1}{1 - e^{-2\pi s}} \left[\int_0^\pi e^{-st} \sin t dt + \int_\pi^{2\pi} e^{-st} \cdot 0 \cdot dt \right] \\ &= \frac{1}{1 - e^{-2\pi s}} \left[\frac{e^{-st}}{s^2 + 1} (-s \sin t - \cos t) \right]_0^\pi + 0 \\ &\therefore \int e^{ax} \sin bx dx = \frac{e^{ax}}{a^2 + b^2} (a \sin bx - b \cos bx) \\ &= \frac{1}{1 - e^{-2\pi s}} \cdot \frac{e^{-\pi s} + 1}{s^2 + 1} \\ &= \frac{1}{(1 - e^{-\pi s})(1 + e^{-\pi s})} \cdot \frac{1 + e^{-\pi s}}{1 + s^2} \\ &= \frac{1}{(1 - e^{-\pi s})(1 + s^2)} \end{aligned}$$

[I] Initial Value Theorem

If $L\{F(t)\} = f(s)$ then $\lim_{t \rightarrow 0} F(t) = \lim_{s \rightarrow \infty} sf(s)$... (16)

We have $L\{F'(t)\} = sf(s) - F(0)$ by property [E]

$$\text{i.e., } \int_0^{\infty} e^{-st} F'(t) dt = sf(s) - F(0)$$

Taking the limit as $s \rightarrow \infty$,

$$\lim_{s \rightarrow \infty} \int_0^{\infty} e^{-st} F'(t) dt = \lim_{s \rightarrow \infty} sf(s) - F(0)$$

$$\begin{aligned} \text{or } \lim_{s \rightarrow \infty} sf(s) &= F(0) + \int_0^{\infty} \left(\lim_{s \rightarrow \infty} e^{-st} \right) F'(t) dt \\ &= F(0) + 0 \quad \because \lim_{s \rightarrow \infty} e^{-st} = 0 \\ &= \lim_{t \rightarrow 0} F(t) \end{aligned}$$

Problem 22. Verify the initial value theorem for the function

$$F(t) = e^{-3t}$$

$$\therefore F(t) = e^{-3t}$$

$$\therefore f(s) = L\{F(t)\} = L\{e^{-3t}\} = \frac{1}{s+3}$$

$$\text{Now } \lim_{t \rightarrow 0} F(t) = \lim_{t \rightarrow 0} e^{-3t} = 1$$

$$\text{and } \lim_{s \rightarrow \infty} sf(s) = \lim_{s \rightarrow \infty} \frac{s}{s+3} = 1.$$

$$\text{Hence } \lim_{t \rightarrow 0} F(t) = \lim_{s \rightarrow \infty} sf(s).$$

[J] Final-Value Theorem

If $L\{F(t)\} = f(s)$, then $\lim_{t \rightarrow \infty} F(t) = \lim_{s \rightarrow 0} sf(s)$... (17)

We have $L\{F'(t)\} = sf(s) - F(0)$ by Prop. [E]

$$\text{i.e., } \int_0^{\infty} e^{-st} F'(t) dt = sf(s) - F(0)$$

Taking the limit as $s \rightarrow 0$,

$$\lim_{s \rightarrow 0} \int_0^{\infty} e^{-st} F'(t) dt = \lim_{s \rightarrow 0} sf(s) - F(0)$$

$$\text{or } \lim_{s \rightarrow 0} sf(s) = F(0) + \lim_{s \rightarrow 0} \int_0^{\infty} e^{-st} F'(t) dt$$

$$= F(0) + \int_0^{\infty} \left(\lim_{s \rightarrow 0} e^{-st} \right) F'(t) dt$$

$$= F(0) + \int_0^{\infty} 1 \cdot F'(t) dt$$

$$= F(0) + \int_0^{\infty} \frac{d}{dt} F(t) dt$$

$$= F(0) + \left[F(t) \right]_0^{\infty}$$

$$= F(0) + \lim_{t \rightarrow \infty} F(t) - F(0)$$

$$= \lim_{t \rightarrow \infty} F(t).$$

Problem 23. Verify the Final-value theorem for the function $F(t) = e^{-2t}$.

We have $F(t) = e^{-2t}$ so that $f(s) = L\{F(t)\} = L\{e^{-2t}\} = \frac{1}{s+2}$

$$\therefore \lim_{t \rightarrow \infty} F(t) = \lim_{t \rightarrow \infty} e^{-2t} = 0$$

$$\text{and } \lim_{s \rightarrow 0} sf(s) = \lim_{s \rightarrow 0} \frac{s}{s+2} = 0$$

$$\text{Hence } \lim_{t \rightarrow \infty} F(t) = \lim_{s \rightarrow 0} sf(s).$$

[K] Behaviour of $f(s)$ as $s \rightarrow 0$ and $s \rightarrow \infty$

$$\text{We have } L\{F(t)\} = f(s) = \int_0^{\infty} e^{-st} F(t) dt$$

$$\text{when } s \rightarrow 0, f(0) = \int_0^{\infty} F(t) dt \quad \dots (18)$$

$$\text{and when } s \rightarrow \infty, \lim_{s \rightarrow \infty} f(s) = \int_0^{\infty} 0 \cdot F(t) dt = 0 \quad \dots (19)$$

[L] Relationship with Fourier Transform

Let us define a function $F(t)$ such that

$$F(t) = \begin{cases} e^{-st} f(t), & t > 0 \\ 0, & t < 0 \end{cases}$$

Then $f[F(t)] = \int_{-\infty}^{\infty} e^{int} F(t) dt$ by definition of complex Fourier transform

$$= \int_{-\infty}^0 e^{int} \cdot 0 dt + \int_0^{\infty} e^{int} e^{-st} f(t) dt$$

$$= \int_0^{\infty} e^{-(s-in)t} f(t) dt$$

$$= \int_0^{\infty} e^{-st} f(t) dt \quad \text{by taking } x - in = s$$

$$= L\{f(t)\} \quad \text{by definition of Laplace transform}$$

$$\text{Hence } f[F(t)] = L\{f(t)\} \quad \dots (20)$$

10.10 SOME METHODS FOR FINDING LAPLACE TRANSFORMS

[1] **Direct method.** This is based on the definition of Laplace transforms given in § 10.4, e.g.

$$\begin{aligned} L\{(t^2+1)^2\} &= L\{t^4+2t^2+1\} = L\{t^4\} + 2L\{t^2\} + L\{1\} \\ &= \int_0^\infty e^{-st} t^4 \cdot dt + 2 \int_0^\infty e^{-st} t^2 dt + \int_0^\infty e^{-st} \cdot 1 \cdot dt \\ &= \frac{4}{s^5} + 2 \cdot \frac{2}{s^3} + \frac{1}{s} = \frac{24+4s^2+s^4}{s^5} \quad \dots (1) \end{aligned}$$

[2] **Series-expansion method.** If the function $F(t)$ is expressible as a Power series e.g.

$$F(t) = a_0 + a_1 t + a_2 t^2 + \dots = \sum_{n=0}^{\infty} a_n t^n$$

then the Laplace transform of $F(t)$ is obtained by taking Laplace transforms of each term in the series e.g.

$$\begin{aligned} L\{\sin \sqrt{t}\} &= L\left\{\sqrt{t} - \frac{(\sqrt{t})^3}{|3|} + \frac{(\sqrt{t})^5}{|5|} - \frac{(\sqrt{t})^7}{|7|} + \dots\right\} \\ &= L\{t^{1/2}\} - \frac{1}{|3|} L\{t^{3/2}\} + \frac{1}{|5|} L\{t^{5/2}\} - \frac{1}{|7|} L\{t^{7/2}\} + \dots \\ &= \frac{|3/2|}{s^{3/2}} - \frac{|5/2|}{|3| s^{5/2}} + \frac{|7/2|}{|5| s^{7/2}} - \frac{|9/2|}{|7| s^{9/2}} + \dots \\ &= \frac{\sqrt{\pi}}{2s^{3/2}} \left[1 - \frac{1}{2^2 s} + \frac{1}{2} \left(\frac{1}{2^2 s}\right)^2 - \frac{1}{3} \left(\frac{1}{2^2 s}\right)^3 + \dots \right] \\ &= \frac{\sqrt{\pi}}{2s^{3/2}} \left[e^{-2\frac{1}{2}s} \right] = \frac{e^{-\frac{1}{2}s} \sqrt{\pi}}{2s^{3/2}} \quad \dots (2) \end{aligned}$$

[3] **Method of differential equations.** If a differential equation satisfied by the function $F(t)$ can be determined, then its Laplace transform may be evaluated by using the properties of Laplace transforms e.g.

$$\begin{aligned} \text{if } F(t) &= \sin \sqrt{t} \text{ then } F'(t) = \frac{1}{2\sqrt{t}} \cos \sqrt{t} \\ \text{and } F''(t) &= \frac{1}{2} \left[-\frac{1}{2t^{3/2}} \cos \sqrt{t} - \frac{1}{2t} \sin \sqrt{t} \right] = \frac{1}{2t} [-F'(t) - \frac{1}{2}F(t)] \\ \text{i.e. } 4tF''(t) + 2F'(t) + F(t) &= 0 \quad \dots (3) \\ \text{which is clearly satisfied by } F(t) &= \sin \sqrt{t} \end{aligned}$$

Assuming that $L\{F(t)\} = L\{\sin \sqrt{t}\} = f(s)$, we have on using the properties [E] and [F] of § 10.9,

$$\begin{aligned} L\{4tF''(t)\} &= 4L\{tF''(t)\} \\ &= -4 \frac{d}{ds} [s^2 f(s) - sF(0) - F(0)] \end{aligned}$$

$$\begin{aligned} &= -8s f(s) - 4s^2 f'(s) - 4F(0) \\ L\{2F'(t)\} &= 2L\{F'(t)\} = 2[sf(s) - F(0)] \\ &= 2sf(s) - 2F(0) \end{aligned}$$

and

$$L\{F(t)\} = f(s)$$

∴ Taking Laplace transform of (3), we get

$$\begin{aligned} L\{4tF''(t)\} + L\{2F'(t)\} + L\{F(t)\} &= 0 \\ \text{i.e. } -8s f(s) - 4s^2 f'(s) - 4F(0) + 2s f(s) - 2F(0) + f(s) &= 0 \\ \text{or } 4s^2 f'(s) + (6s-1) f(s) &= 0 \text{ since } F(0) = \sin \sqrt{0} = 0 \\ \text{or } \frac{f'(s)}{f(s)} &= \frac{1-6s}{4s^2} = \frac{1}{4s^2} - \frac{3}{2s} \end{aligned}$$

Integrating with regard to 's', we find

$$\begin{aligned} \log f(s) &= \log C - \frac{1}{4s} - \frac{3}{2} \log s, \quad C \text{ being constant of integration} \\ &= \log \frac{C}{s^{3/2}} e^{-\frac{1}{4s}} \end{aligned}$$

$$\text{i.e. } f(s) = \frac{C e^{-1/4s}}{s^{3/2}} \quad \dots (4)$$

Now to determine C , apply the limits of initial-value theorem i.e. when $t \rightarrow 0, s \rightarrow \infty$.

$$\text{For } t \text{ small, } \sin \sqrt{t} = \sqrt{t} \text{ so that } L\{\sqrt{t}\} = \frac{|3/2|}{s^{3/2}} = \frac{\sqrt{\pi}}{2s^{3/2}}$$

and for s large, $f(s) = \frac{C}{s^{3/2}}$ as $\lim_{s \rightarrow \infty} e^{-1/4s} = 1$

∴ $L\{F(t)\} = f(s)$ gives for $t \rightarrow 0, s \rightarrow \infty$,

$$\frac{\sqrt{\pi}}{2s^{3/2}} = \frac{C}{s^{3/2}} \text{ i.e. } C = \frac{\sqrt{\pi}}{2}.$$

$$\text{Hence } f(s) = \frac{\sqrt{\pi} e^{-1/4s}}{2s^{3/2}} \quad \dots (5)$$

[4] **Method of differentiation with respect to a parameter.** This method is based on differentiation of Laplace transform of a known function w.r.t. a parameter, e.g. if

$$F(t) = t \sin at \text{ then we have to find } L\{t \sin at\}$$

$$\text{Consider } L\{\cos at\} = \int_0^\infty e^{-st} \cos at \, dt = \frac{s}{s^2 + a^2}$$

Differentiating w.r.t. a , we get

$$\frac{d}{da} \int_0^\infty e^{-st} \cos at \, dt = \frac{d}{da} \left(\frac{s}{s^2 + a^2} \right)$$

$$\text{i.e. } \int_0^\infty e^{-st} (-t \sin at) \, dt = \frac{-2as}{(s^2 + a^2)^2}$$

$$\text{or } +L\{t \sin at\} = \frac{+2as}{(s^2 + a^2)^2} \quad (6)$$

10.11. THE LAPLACE TRANSFORMS OF SOME SPECIAL FUNCTIONS

[1] The Gamma function. Euler's Gamma function is defined as

$$\Gamma n = \int_0^{\infty} e^{-x} x^{n-1} dx, n > 0.$$

Its important properties have already been discussed in chapter on Beta and Gamma functions, but a few of them are mentioned here.

$$\Gamma(n+1) = n\Gamma n \text{ for } n > 0 \text{ and } \Gamma n = \frac{(n-1)!}{n} \text{ for } n < 0$$

$$\Gamma(n+1) = |n| \text{ for positive integral values of } n.$$

$$\Gamma n \Gamma(1-n) = \frac{\pi}{\sin n\pi}, 0 < n < 1, \Gamma \frac{1}{2} = \sqrt{\pi}$$

For large n , $\Gamma(n+1) \approx \sqrt{2\pi n} n^n e^{-n}$ (Stirling formula)

$$\text{Now we have } L\{t^n\} = \int_0^{\infty} e^{-st} t^n dt$$

$$\text{Put } st = u \text{ i.e. } t = \frac{u}{s} \text{ and } dt = \frac{du}{s}$$

$$\therefore L\{t^n\} = \frac{1}{s^{n+1}} \int_0^{\infty} e^{-u} u^n du = \frac{\Gamma(n+1)}{s^{n+1}} \quad \dots (1)$$

(by the definition of Gamma function)

If we now put $n = -\frac{1}{2}$,

$$L\{t^{-1/2}\} = \frac{\Gamma \frac{1}{2}}{s^{1/2}} = \frac{\sqrt{\pi}}{s} \quad \dots (2)$$

[2] Bessel Functions. Bessel function of order n is defined as

$$J_n(t) = \frac{t^n}{2^n \Gamma(n+1)} \left\{ 1 - \frac{t^2}{2(2n+2)} + \frac{t^4}{2 \cdot 4 (2n+2)(2n+4)} \dots \right\}$$

which satisfies Bessel's differential equation

$$y''(t) + \frac{1}{t} y'(t) + \left(1 - \frac{n^2}{t^2}\right) y(t) = 0$$

$$\text{or } t^2 J_n''(t) + t J_n'(t) + (t^2 - n^2) J_n(t) = 0$$

Some important properties are:

$$J_{-n}(t) = (-1)^n J_n(t), n \text{ being positive integral}$$

$$J_n(t) = i^{-n} J_n(t), J_n \text{ being modified Bessel function of order } n.$$

$$J_{n+1}(t) = \frac{2n}{t} J_n(t) - J_{n-1}(t)$$

$$\frac{d}{dt} \{t^n J_n(t)\} = t^n J_{n-1}(t) \text{ which becomes } J'_0(t) = -J_1(t) \text{ for } n=0$$

$$e^{\frac{1}{2}t(x^2/x)} = \sum_{n=-\infty}^{\infty} J_n(t) x^n$$

known as generating function for the Bessel functions.

$J_0(t)$ is called Bessel function of order zero and has for its expansion

$$J_0(t) = 1 - \frac{t^2}{2^2} + \frac{t^4}{2^2 \cdot 4^2} - \frac{t^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots$$

$$\therefore L\{J_0(t)\} = L\{1\} - L\left\{\frac{t^2}{2^2}\right\} + L\left\{\frac{t^4}{2^2 \cdot 4^2}\right\} - \dots$$

$$= \frac{1}{s} - \frac{1}{2^2 s^3} + \frac{1}{2^2 \cdot 4^2 s^5} - \dots$$

$$= \frac{1}{s} \left[1 - \frac{1}{2} \left(\frac{1}{s^2}\right) + \frac{1 \cdot 3}{2 \cdot 4} \left(\frac{1}{s^4}\right) - \dots \right]$$

$$= \frac{1}{s} \left(1 + \frac{1}{s} \right)^{-1/2} = \frac{1}{\sqrt{s^2 + 1}} \quad \dots (3)$$

$$\text{Similarly } L\{J_1(t)\} = 1 - s / \sqrt{s^2 + 1} \quad \dots (4)$$

Aliter. $J_0(t)$ satisfies the equation

$$t J_0''(t) + J_0'(t) + t J_0(t) = 0$$

$$\therefore L\{t J_0''(t)\} + L\{J_0'(t)\} + L\{t J_0(t)\} = 0$$

Taking $L\{J_0(t)\} = f(s)$ and using properties [E] and [F] of § 10.9.

$$- \frac{d}{ds} \{s^2 f(s) - s F'(s) - F'(s)\} + \{s f(s) - F(s)\} - \frac{d}{ds} f(s) = 0$$

where $F(t) = J_0(t)$ gives $F(0) = 1$ and $F'(0) = 0$

$$\therefore -2s f(s) - s^2 f'(s) + 1 + s f(s) - 1 - f'(s) = 0$$

$$\text{or } s f(s) + (s^2 + 1) f'(s) = 0$$

$$\text{i.e. } \frac{f'(s)}{f(s)} = \frac{-s}{s^2 + 1} = -\frac{1}{2} \cdot \frac{2s}{s^2 + 1}$$

Integrating with regard to 's'

$$\log f(s) = -\frac{1}{2} \log(s^2 + 1) + \log C, C \text{ being constant of integration}$$

$$\therefore f(s) = \frac{C}{\sqrt{s^2 + 1}}$$

Applying initial value theorem, we have

$$\lim_{s \rightarrow \infty} s f(s) = \lim_{t \rightarrow 0} F(t)$$

$$\text{i.e. } \lim_{s \rightarrow \infty} \frac{sC}{\sqrt{s^2 + 1}} = \lim_{t \rightarrow 0} J_0(t) \text{ which gives } C = 1$$

$$\text{Hence } f(s) = \frac{1}{\sqrt{s^2 + 1}} \text{ i.e. } L\{J_0(t)\} = \frac{1}{\sqrt{s^2 + 1}}$$

Now using the change of scale property, we have

$$L\{J_0(at)\} = \frac{1}{a} f\left(\frac{s}{a}\right) \text{ where } f(s) = L\{J_0(t)\} = \frac{1}{\sqrt{s^2 + 1}}$$

$$\text{so that } f\left(\frac{s}{a}\right) = \frac{1}{\sqrt{\left(\frac{s}{a}\right)^2 + 1}} = \frac{1}{\sqrt{s^2 + a^2}}$$

Hence
$$L\{J_0(at)\} = \frac{1}{\sqrt{s^2 + a^2}} \quad \dots (5)$$

Further to deduce $L\{t J_0(at)\}$, using the property [F] of §10.9, we get

$$L\{t J_0(at)\} = -\frac{d}{ds} [L\{J_0(at)\}] = -\frac{d}{ds} \left(\frac{1}{\sqrt{s^2 + a^2}} \right) = \frac{s}{(s^2 + a^2)^{3/2}} \quad \dots (6)$$

Similarly $L\{t J_1(t)\} = \frac{1}{(s^2 + 1)^{3/2}} \quad \dots (7)$

[3] The error function and its complement

The error function of a variable t denoted by $\text{erf}(t)$ or $E_r(t)$ is defined as

$$\text{erf}(t) = \frac{2}{\sqrt{\pi}} \int_0^t e^{-x^2} dx = E_r(t)$$

and the complement of the error function denoted by $\text{erf } C(t)$ is defined by

$$\text{erf } C(t) = 1 - \text{erf}(t) = 1 - \frac{2}{\sqrt{\pi}} \int_0^t e^{-x^2} dx = \frac{2}{\sqrt{\pi}} \int_t^\infty e^{-x^2} dx.$$

It is notable that

$$\lim_{t \rightarrow 0} \text{erf}(t) = 0 \text{ and } \lim_{t \rightarrow \infty} \text{erf}(t) = 1.$$

Thus $\text{erf } \sqrt{t} = \frac{2}{\sqrt{\pi}} \int_0^{\sqrt{t}} e^{-x^2} dx = \frac{2}{\sqrt{\pi}} \int_0^{\sqrt{t}} \left\{ 1 - x^2 + \frac{x^4}{2!} - \frac{x^6}{3!} + \dots \right\} dx$

$$= \frac{2}{\sqrt{\pi}} \left[t^{1/2} - \frac{t^{3/2}}{3} + \frac{t^{5/2}}{5 \cdot 2} - \frac{t^{7/2}}{7 \cdot 3} + \dots \right]$$

$$\therefore L\{\text{erf } \sqrt{t}\} = \frac{2}{\sqrt{\pi}} L\left\{ t^{1/2} - \frac{t^{3/2}}{3} + \frac{t^{5/2}}{5 \cdot 2} - \frac{t^{7/2}}{7 \cdot 3} + \dots \right\}$$

$$= \frac{2}{\sqrt{\pi}} \left[\frac{\Gamma(3/2)}{s^{3/2}} - \frac{\Gamma(5/2)}{s^{5/2}} + \frac{\Gamma(7/2)}{s^{7/2}} - \frac{\Gamma(9/2)}{s^{9/2}} + \dots \right]$$

$$= \frac{1}{s^{3/2}} - \frac{1}{2} \cdot \frac{1}{s^{5/2}} + \frac{1.3}{2.4} \cdot \frac{1}{s^{7/2}} - \frac{1.3.5}{2.4.6} \cdot \frac{1}{s^{9/2}} + \dots$$

$$= \frac{1}{s^{3/2}} \left(1 + \frac{1}{s} \right)^{-1/2} = \frac{1}{s \sqrt{s+1}} \quad \dots (8)$$

[4] The sine, cosine and exponential integrals

The sine integral is defined as $S_i(t) = \int_0^t \frac{\sin x}{x} dx$

and the cosine integral is defined as $C_i(t) = \int_0^t \frac{\cos x}{x} dx$

Also the exponential integral is defined as $E_i(t) = \int_t^\infty \frac{e^{-x}}{x} dx$

we have $S_i(t) = \int_0^t \frac{\sin x}{x} dx = \int_0^t \frac{1}{x} \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \right) dx$

$$= \int_0^t \left\{ 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \frac{x^6}{7!} + \dots \right\} dx$$

$$= t - \frac{t^3}{3 \cdot 3!} + \frac{t^5}{5 \cdot 5!} - \frac{t^7}{7 \cdot 7!} + \dots$$

$$\therefore L\{S_i(t)\} = L\left\{ t - \frac{t^3}{3 \cdot 3!} + \frac{t^5}{5 \cdot 5!} - \frac{t^7}{7 \cdot 7!} + \dots \right\}$$

$$= \frac{1}{s^2} - \frac{3}{3!} \cdot \frac{1}{s^4} + \frac{1}{5!} \cdot \frac{1}{s^6} - \frac{1}{7!} \cdot \frac{1}{s^8} + \dots$$

$$= \frac{1}{s} \left[\frac{1/s}{1} - \frac{(1/s)^3}{3} + \frac{(1/s)^5}{5} - \frac{(1/s)^7}{7} + \dots \right]$$

$$= \frac{1}{s} \tan^{-1} \frac{1}{s} \quad \dots (9)$$

Similarly $L\{C_i(t)\} = \frac{1}{2s} \log(s^2 + 1) \quad \dots (10)$

and $L\{E_i(t)\} = L\left\{ \int_t^\infty \frac{e^{-x}}{x} dx \right\}$; Put $x = ty$ i.e. $\frac{dx}{x} = \frac{dy}{y}$

(on logarithmic differentiation)

$$= L \int_1^\infty \frac{e^{-ty}}{y} dy$$

$$= \int_0^\infty e^{-st} \left\{ \int_1^\infty \frac{e^{-ty}}{y} dy \right\} dt \text{ by definition of Laplace transform}$$

$$= \int_1^\infty \frac{1}{y} \left\{ \int_0^\infty e^{-(s+ty)t} dt \right\} dy, \text{ by changing the order of integration}$$

$$= \int_1^\infty \frac{1}{y} \cdot \frac{1}{s+ty} dy$$

$$= \int_1^\infty \frac{1}{s} \left[\frac{1}{y} - \frac{1}{s+ty} \right] dy = \frac{1}{s} \left[\log y - \log(s+ty) \right]_1^\infty$$

$$= \frac{1}{s} \left[-\log\left(\frac{s}{y} + 1\right) \right]_1^\infty$$

$$= \frac{1}{s} \log(s+1) \quad \dots (11)$$

[5] Heaviside unit function or unit step function

Unit step function is defined as

$$U(t-a) = \begin{cases} 0, & t < a \\ 1, & t > a. \end{cases}$$

$$\therefore L\{U(t-a)\} = \int_0^\infty e^{-st} U(t-a) dt$$

$$= \int_0^a e^{-st} \cdot 0 \, dt + \int_a^\infty e^{-st} \cdot 1 \, dt = \left[\frac{e^{-st}}{-s} \right]_a^\infty$$

$$= \frac{1}{s} e^{-as} \quad \dots (12)$$

[6] Dirac-delta function or unit impulse function

A function $\delta(t)$ such that

$$\lim_{\epsilon \rightarrow 0} F_\epsilon(t) = \delta(t) \text{ where } F_\epsilon(t) = \begin{cases} 1/\epsilon, & 0 \leq t \leq \epsilon \\ 0, & t > \epsilon \end{cases}$$

and $\int_0^\infty F_\epsilon(t) \, dt = 1$

With the properties:

(i) $\int_0^\infty \delta(t) \, dt = 1$

(ii) $\int_0^\infty \delta(t) G(t) \, dt = G(0)$ for any continuous function $G(t)$

(iii) $\int_0^\infty \delta(t-a) G(t) \, dt = G(a)$ for any continuous function $G(t)$
is called as unit impulse function or Dirac-delta function.

We have $L\{\delta(t)\} = \int_0^\infty e^{-st} \lim_{\epsilon \rightarrow 0} F_\epsilon(t) \, dt$ which is not defined since

$\lim_{\epsilon \rightarrow 0} F_\epsilon(t)$ does not exist and hence it is useful to consider $\delta(t) = \lim_{\epsilon \rightarrow 0} F_\epsilon(t)$ to be such that $L\{\delta(t)\} = 1$ (13)

Also $L\{F_\epsilon(t)\} = \int_0^\infty e^{-st} F_\epsilon(t) \, dt = \int_0^\epsilon e^{-st} \cdot \frac{1}{\epsilon} \, dt + \int_\epsilon^\infty e^{-st} \cdot 0 \, dt$
 $= \frac{1}{\epsilon} \int_0^\epsilon e^{-st} \, dt = \frac{1 - e^{-\epsilon s}}{\epsilon s}$... (14)

[7] Null functions. A null function $N(t)$ for all $t > 0$ is defined as

$$\int_0^t N(x) \, dx = 0$$

e.g. if $F(t) = \begin{cases} 1, & t=1 \\ 0, & \text{otherwise,} \end{cases}$ then it is a null function, since

$\int_0^t F(x) \, dx = 0$ for all $t > 0$. As such the Laplace transform of a null function is zero i.e. $L\{N(t)\} = 0$... (15)

10.12. EVALUATION OF INTEGRALS WITH THE HELP OF LAPLACE TRANSFORMS

By definition, we have

$$L\{F(t)\} = \int_0^\infty e^{-st} F(t) \, dt = f(s) \quad \dots (1)$$

Assuming that the integral is convergent and proceeding to the limit

$s \rightarrow 0$, this reduces to

$$\int_0^\infty F(t) \, dt = f(0) \quad \dots (2)$$

(1) and (2) are sometimes used to evaluate integrals.

Problem 24. Evaluate the following integrals

(a) $\int_0^\infty t^2 e^{-t} \sin t \, dt$

(b) $\int_0^\infty J_0(t) \, dt$

(c) $\int_0^\infty e^{-t} \operatorname{erf} \sqrt{t} \, dt$.

(a) We have $L\{\sin t\} = \frac{1}{s^2 + 1}$

$\therefore L\{t^2 \sin t\} = (-1)^2 \frac{d^2}{ds^2} L\{\sin t\}$ by Prop. [F] of §10.9

$$= \frac{d^2}{ds^2} \left(\frac{1}{s^2 + 1} \right) = \frac{d}{ds} \left(\frac{-2s}{(s^2 + 1)^2} \right) = -\frac{2(1 - 3s^2)}{(1 + s^2)^3}$$

So that $\int_0^\infty e^{-st} \cdot t^2 \sin t \, dt = -\frac{2(1 - 3s^2)}{(1 + s^2)^3}$.

Putting $s=1$, we find $\int_0^\infty e^{-t} t^2 \sin t \, dt = \frac{4}{4} = 1$

(b) We have $L\{J_0(t)\} = \frac{1}{\sqrt{s^2 + 1}}$ by (3) of §10.11

i.e. $\int_0^\infty e^{-st} J_0(t) \, dt = \frac{1}{\sqrt{s^2 + 1}}$

Proceeding to the limit as $s \rightarrow 0$, we get

$$\int_0^\infty J_0(t) \, dt = 1.$$

(c) We have, $L\{\operatorname{erf} \sqrt{t}\} = \frac{1}{s \sqrt{s+1}}$ by (8) of §10.11

i.e. $\int_0^\infty e^{-st} \operatorname{erf} \sqrt{t} \, dt = \frac{1}{s \sqrt{s+1}}$.

Proceeding to the limit $s \rightarrow 1$, we find

$$\int_0^\infty e^{-t} \operatorname{erf} \sqrt{t} \, dt = \frac{1}{\sqrt{2}}.$$

10.13. LAPLACE TRANSFORM OF THE LAPLACE TRANSFORM

We have $L\{F(t)\} = \int_0^\infty e^{-st} F(t) \, dt = f(s)$

$$\therefore L[L\{F(t)\}] = L \left\{ \int_0^\infty e^{-st} F(t) dt \right\}$$

$$\text{i.e. } L\{f(s)\} = \int_0^\infty e^{-ps} ds \int_0^\infty e^{-st} F(t) dt \quad \dots (1)$$

Here the area of integration being the whole positive quadrant, we can change the order of integration and hence we get

$$L[L\{F(t)\}] = \int_0^\infty F(t) dt \int_0^\infty e^{-s(t+p)} ds$$

$$= \int_0^\infty F(t) dt \cdot \left[\frac{e^{-s(t+p)}}{t+p} \right]_0^\infty$$

$$= \int_0^\infty \frac{F(t)}{t+p} dt \quad \dots (2)$$

10.14. THE INVERSE LAPLACE TRANSFORM

If $f(s)$ be the Laplace transform of a function $F(t)$ i.e. $L\{F(t)\} = f(s)$ then $F(t)$ is said to be an *Inverse Laplace transform* of $f(s)$ and written as

$$F(t) = L^{-1}\{f(s)\} \quad \dots (1)$$

Here L^{-1} is known as the *Inverse Laplace transformation operator*

e.g. if $L\{e^{2t}\} = \frac{1}{s-2}$ then $e^{2t} = L^{-1}\left\{\frac{1}{s-2}\right\}$.

Uniqueness of Inverse Laplace transform.

If $N(t)$ be a null function then $L\{N(t)\} = 0$

and $L\{F(t)\} = f(s)$

$$\therefore L\{F(t) + N(t)\} = L\{F(t)\} + L\{N(t)\}$$

$$= f(s)$$

So that $L^{-1}\{f(s)\} = F(t)$

and also $L^{-1}\{f(s)\} = F(t) + N(t)$.

Showing that we can have two different functions with the same Laplace transform and hence the inverse Laplace transform of a function is not unique. It is, however, unique if we do not allow null function which in general, does not arise in cases of physical interest. This result is known as *Lerch's theorem* which states:

If $F(t)$ is sectionally continuous function in every finite interval $0 \leq t \leq N$ and is of exponential order for $t < N$ such that $L\{F(t)\} = f(s)$ then the inverse Laplace transform of $f(s)$ i.e. $L^{-1}\{f(s)\} = F(t)$ is unique.

Conclusively, a given function $f(s)$ cannot have more than one inverse transform $F(t)$ that is continuous for each positive t . But a function $f(s)$ may not have a continuous inverse transform e.g. the Laplace transform of $s^{-1} e^{-kt}$ is the step function which is discontinuous.

Note. Some inverse transforms can be tabulated as follows:

$f(s)$	$L^{-1}\{f(s)\} = F(t)$	$f(s)$	$L^{-1}\{f(s)\} = F(t)$
$\frac{1}{s}, s > 0$	1	$2s/(s^2 + a^2)^2$	$t \sin at/a$
$\frac{1}{s^2}, s > 0, \dots$	t	$(s^2 - a^2)/(s^2 + a^2)$	$t \cos at$
$\frac{1}{s^{n+1}}, n = 0, 1, 2, 3, \dots$	$t^n/n!$	$1/\{(s-a)^2 + b^2\}$	$e^{at} \sin bt/b$
$1/(s-a), \dots$	e^{at}	$(s-a)/\{(s-a)^2 + b^2\}$	$e^{at} \cos bt$
$1/(s^2 + a^2), s > 0$	$\frac{\sin at}{a}$	$1/\sqrt{s^2 + a^2}$	$J_0(at)$
$s/(s^2 + a^2), s > 0, \dots$	$\cos at$	$s/(s^2 + a^2)^{3/2}$	$t J_0(at)$
$1/(s^2 - a^2), s > a $	$\frac{\sinh at}{a}$	$1/(s-a)^n$	$\frac{t^{n-1} e^{at}}{(n-1)!}, n > 0$
$s/(s^2 - a^2), s > a $	$\cosh at$		

10.15. SOME PROPERTIES OF INVERSE LAPLACE TRANSFORM

[A] Linearity Property

If the Laplace transform of $F_1(t)$ and $F_2(t)$ be $f_1(s)$ and $f_2(s)$ respectively and C_1, C_2 are constants then

$$L^{-1}\{C_1 f_1(s) + C_2 f_2(s)\} = C_1 L^{-1}\{f_1(s)\} + C_2 L^{-1}\{f_2(s)\}$$

$$= C_1 F_1(t) + C_2 F_2(t) \quad \dots (1)$$

By Prop. [A] of § 10.9, we have

$$L\{C_1 F_1(t) + C_2 F_2(t)\} = C_1 L\{F_1(t)\} + C_2 L\{F_2(t)\} = C_1 f_1(s) + C_2 f_2(s)$$

$$\therefore C_1 F_1(t) + C_2 F_2(t) = L^{-1}\{C_1 f_1(s) + C_2 f_2(s)\}$$

$$\text{i.e. } L^{-1}\{C_1 f_1(s) + C_2 f_2(s)\} = C_1 F_1(t) + C_2 F_2(t) = C_1 L^{-1}\{f_1(s)\} + C_2 L^{-1}\{f_2(s)\}$$

Problem 25. Find $L^{-1}\left\{\frac{2(s-a)}{(s-a)^2 + b^2} + \frac{8-6s}{16s^2+9} + \frac{24-30\sqrt{s}}{s^4}\right\}$

$$L^{-1}\left\{\frac{2(s-a)}{(s-a)^2 + b^2} + \frac{8-6s}{16s^2+9} + \frac{24-30\sqrt{s}}{s^4}\right\} = L^{-1}\left\{\frac{2(s-a)}{(s-a)^2 + b^2}\right\}$$

$$+ L^{-1}\left\{\frac{8-6s}{16s^2+9}\right\} + L^{-1}\left\{\frac{24-30\sqrt{s}}{s^4}\right\}$$

$$= 2L^{-1}\left\{\frac{s-a}{(s-a)^2 + b^2}\right\} + \frac{1}{2} L^{-1}\left\{\frac{1}{s^2+9/16}\right\} - \frac{3}{8} L^{-1}\left\{\frac{s}{s^2+9/16}\right\}$$

$$+ 24L^{-1}\left\{\frac{1}{s^4}\right\} - 30L^{-1}\left\{\frac{1}{s^{7/2}}\right\}$$

$$= 2e^{at} \cos bt + \frac{1}{2} \sin \frac{3t}{4} - \frac{3}{8} \cos \frac{3t}{4} + 24 \frac{t^3}{3} - 30 \frac{t^{5/2}}{\Gamma 7/2}$$

$$= 2e^{at} \cos bt + \frac{1}{2} \sin \frac{3t}{4} - \frac{3}{8} \cos \frac{3t}{4} + 4t^3 - \frac{16}{\sqrt{\pi}} t^{5/2}$$

[B] First Translation (or Shifting) Property

If $L^{-1}\{f(s)\} = F(t)$, then $L^{-1}\{f(s-a)\} = e^{at}F(t)$.

We have, $f(s) = \int_0^\infty e^{-st} F(t) dt = L\{F(t)\}$

$$\therefore f(s-a) = \int_0^\infty e^{-(s-a)t} F(t) dt = \int_0^\infty e^{-st} \cdot e^{at} F(t) dt$$

$$= L\{e^{at} F(t)\}$$

$$\text{i.e. } L^{-1}\{f(s-a)\} = e^{at} F(t) \quad \dots (2)$$

(2) may also be expressed as

$$L^{-1}\{f(s)\} = e^{-at} L^{-1}\{f(s-a)\} \quad \dots (3)$$

Which follows that the substitution of $s-a$ for s in the transform corresponds to the multiplication of the object function $F(t)$ by the function e^{at} .

We may illustrate it as follows:

$$\therefore \frac{|m|}{s^{m+1}} = L\{t^m\}, \text{ where } m=1, 2, \dots \text{ and } s > 0$$

$$\therefore \frac{|m|}{(s-a)^{m+1}} = L\{t^m e^{at}\}, s > a$$

$$\text{Also, } \therefore L\{\cos bt\} = \frac{s}{s^2 + b^2}$$

$$\therefore L\{e^{-at} \cos bt\} = \frac{s+a}{(s+a)^2 + b^2}, s > -a.$$

$$\text{Problem 26. Find (a) } L^{-1}\left\{\frac{1}{\sqrt{2s+5}}\right\} \text{ (b) } L^{-1}\left\{\frac{s+1}{s^2+6s+25}\right\}$$

$$\text{(a) We have } L^{-1}\left\{\frac{1}{\sqrt{2s+5}}\right\} = \frac{2}{\sqrt{2}} L^{-1}\left\{\frac{1}{(s+5/2)^{1/2}}\right\}$$

$$= \frac{1}{\sqrt{2}} e^{-5t/2} \cdot \frac{t^{-1/2}}{\Gamma^{1/2}} = \frac{1}{\sqrt{2\pi}} t^{-1/2} e^{-5t/2}$$

$$\text{(b) We have } L^{-1}\left\{\frac{s+1}{s^2+6s+25}\right\} = L^{-1}\left\{\frac{s+3-2}{(s+3)^2+4^2}\right\}$$

$$= L^{-1}\left\{\frac{s+3}{(s+3)^2+4^2}\right\} - 2L^{-1}\left\{\frac{1}{(s+3)^2+4^2}\right\}$$

$$= e^{-3t} \cos 4t - 2e^{-3t} \frac{\sin 4t}{4}$$

$$= e^{-3t} \left[\cos 4t - \frac{1}{2} \sin 4t\right].$$

[C] Second Translation (or Shifting) Property

$$\text{If } L^{-1}\{f(s)\} = F(t), \text{ then } L^{-1}\{e^{-as} f(s)\} = \begin{cases} F(t-a), & t > a \\ 0, & t < a \end{cases} \dots (4)$$

$$\text{Let } e^{-as} f(s) = L\{G(t)\}, \text{ then } L^{-1}\{e^{-as} f(s)\} = G(t) = \begin{cases} F(t-a), & t > a \\ 0, & t < a \end{cases}$$

$$\text{So that } L\{G(t)\} = \int_0^\infty e^{-st} G(t) dt = \int_0^a e^{-st} \cdot 0 \cdot dt + \int_a^\infty e^{-st} F(t-a) dt$$

$$= \int_a^\infty e^{-st} F(t-a) dt \quad \text{Put } t-a=u \text{ so that } dt=du$$

$$= \int_0^\infty e^{-s(u+a)} F(u) du = e^{-as} \int_0^\infty e^{-su} F(u) du$$

$$= e^{-as} \int_0^\infty e^{-st} F(t) dt, \text{ on replacing } u \text{ by } t.$$

$$= e^{-as} L\{F(t)\} = e^{-as} f(s)$$

$$\therefore G(t) = L^{-1}\{e^{-as} f(s)\}$$

Note. In terms of Heaviside unit step function $G(t)$ can be expressed as

$$F(t-a) \cdot U(t-a).$$

$$\text{Problem 27. Find } L^{-1}\left\{\frac{s e^{-2\pi s/3}}{s^2+9}\right\}$$

$$\therefore L^{-1}\left\{\frac{s}{s^2+9}\right\} = \cos 3t$$

$$\therefore L^{-1}\left\{\frac{s e^{-2\pi s/3}}{s^2+9}\right\} = \begin{cases} \cos 3\left(t - \frac{2\pi}{3}\right), & t > \frac{2\pi}{3} \\ 0, & t < \frac{2\pi}{3} \end{cases}$$

$$= \begin{cases} \cos 3t, & t > \frac{2\pi}{3} \\ 0, & t < \frac{2\pi}{3} \end{cases}$$

$$= \cos 3t \cdot U\left(t - \frac{2\pi}{3}\right)$$

[D] Change of Scale Property

$$\text{If } L^{-1}\{f(s)\} = F(t), \text{ then } L^{-1}\{f(as)\} = \frac{1}{a} F\left(\frac{t}{a}\right), a > 0 \quad \dots (5)$$

$$\therefore f(s) = \int_0^\infty e^{-st} F(t) dt, \quad \therefore f(as) = \int_0^\infty e^{-ast} F(t) dt$$

$$\text{Put } at=u \text{ so that } dt = \frac{du}{a}$$

$$\begin{aligned}\therefore f(as) &= \int_0^\infty e^{-us} F\left(\frac{u}{a}\right) \cdot \frac{du}{a} = \frac{1}{a} \int_0^\infty e^{-su} F\left(\frac{u}{a}\right) du \\ &= \frac{1}{a} \int_0^\infty e^{-st} F\left(\frac{t}{a}\right) dt \quad (\text{on replacing } u \text{ by } t) \\ &= \frac{1}{a} L\left\{F\left(\frac{t}{a}\right)\right\} = L\left\{\frac{1}{a} F\left(\frac{t}{a}\right)\right\} \\ \therefore L^{-1}\{f(as)\} &= \frac{1}{a} F\left(\frac{t}{a}\right),\end{aligned}$$

Problem 28. If $L^{-1}\left\{\frac{e^{-1/s}}{s^{1/2}}\right\} = \frac{\cos 2\sqrt{t}}{\sqrt{\pi t}}$, find $L^{-1}\left\{\frac{e^{-a/s}}{s^{1/2}}\right\}$, $a > 0$

$$\therefore L^{-1}\left\{\frac{e^{-1/s}}{s^{1/2}}\right\} = \frac{\cos 2\sqrt{t}}{\sqrt{\pi t}}, \therefore L^{-1}\left\{\frac{e^{-a/s}}{(cs)^{1/2}}\right\} = \frac{1}{c} \cdot \frac{\cos 2\sqrt{t/c}}{\sqrt{\pi t/c}}$$

or $\frac{1}{\sqrt{c}} L^{-1}\left\{\frac{e^{-1/cs}}{s^{1/2}}\right\} = \frac{1}{\sqrt{c}} \cdot \frac{\cos 2\sqrt{t/c}}{\sqrt{\pi t/c}}$

Putting $c = \frac{1}{a}$, we find

$$L^{-1}\left\{\frac{e^{-a/s}}{s^{1/2}}\right\} = \frac{\cos 2\sqrt{at}}{\sqrt{\pi t}}.$$

[E] Inverse Laplace Transform of Derivatives

If $L^{-1}\{f(s)\} = F(t)$, then $L^{-1}\{f^{(n)}(s)\} = (-1)^n \frac{d^n}{dt^n} F(t)$... (6)

By Prop. [F] of § 10.9, we have

$$L\{t^n F(t)\} = (-1)^n f^{(n)}(s), \therefore t^n F(t) = L^{-1}\{(-1)^n f^{(n)}(s)\} = (-1)^n L^{-1}\{f^{(n)}(s)\}$$

or $L^{-1}\{f^{(n)}(s)\} = (-1)^n t^n F(t) = (-1)^n \cdot \frac{(-1)^{2n}}{(-1)^{2n}} F(t) \therefore (-1)^{2n} = 1$
 $= (-1)^n t^n F(t) = (-1)^n \left\{ \frac{d^n}{ds^n} f(s) \right\}$

Problem 29. Find $L^{-1}\left\{\frac{s}{(s^2+a^2)^2}\right\}$

Consider $\frac{d}{ds} \left\{ \frac{1}{s^2+a^2} \right\} = \frac{-2s}{(s^2+a^2)^2}$ i.e. $\frac{s}{(s^2+a^2)^2} = -\frac{1}{2} \frac{d}{ds} \left(\frac{1}{s^2+a^2} \right)$

$$\therefore L^{-1}\left\{\frac{s}{(s^2+a^2)^2}\right\} = -\frac{1}{2} L^{-1}\left\{\frac{d}{ds} \left(\frac{1}{s^2+a^2} \right)\right\} \\ = \frac{t \sin at}{2a} \text{ since } L^{-1}\left(\frac{1}{s^2+a^2}\right) = \frac{\sin at}{a}$$

[F] Multiplication by s^n

If $L^{-1}\{f(s)\} = F(t)$ and $F(0) = 0$, then $L^{-1}\{sf(s)\} = F'(t)$... (7)

By Prop. [E] of § 10.9, $L\{F'(t)\} = sf(s) - F(0) = sf(s) \therefore F(0) = 0$

$$\therefore F'(t) = L^{-1}\{sf(s)\}.$$

In case $F(0) \neq 0$, then $L^{-1}\{sf(s) - F(0)\} = F'(t)$

$$\text{or } L^{-1}\{sf(s)\} = F'(t) + L^{-1}\{F(0)\} = F'(t) + F(0)\delta(t)$$

where $\delta(t)$ is the Dirac-delta function.

Problem 30. Find $L^{-1}\left\{\frac{I}{(s^2+I)^2}\right\}$ when $L^{-1}\left\{\frac{s}{(s^2+I)^2}\right\} = \frac{I}{2} t \sin t$.

We have, $L^{-1}\left\{s \cdot \frac{s}{(s^2+I)^2}\right\} = \frac{d}{dt} \left(\frac{I}{2} t \sin t \right)$

or $L^{-1}\left\{\frac{s^2+I-I}{(s^2+I)^2}\right\} = \frac{1}{2} (t \cos t + \sin t)$

or $L^{-1}\left\{\frac{1}{s^2+I}\right\} - L^{-1}\left\{\frac{1}{(s^2+I)^2}\right\} = \frac{1}{2} (t \cos t + \sin t)$

or $L^{-1}\left\{\frac{1}{(s^2+I)^2}\right\} = L^{-1}\left\{\frac{1}{(s^2+I)}\right\} - \frac{1}{2} (t \cos t + \sin t)$
 $= \sin t - \frac{1}{2} (t \cos t + \sin t) = \frac{1}{2} (\sin t - t \cos t)$

[G] Division by Powers of s

If $L^{-1}\{f(s)\} = F(t)$, then (i) $L^{-1}\left\{\frac{f(s)}{s}\right\} = \int_0^t F(u) du$... (8)

(ii) $L^{-1}\left\{\frac{f(s)}{s^2}\right\} = \int_0^t \int_0^v F(u) du dv$... (9)

(i) Let $G(t) = \int_0^t F(u) du = L^{-1}\left\{\frac{f(s)}{s}\right\}$, then $G'(t) = F(t)$, $G(0) = 0$

$\therefore L\{G'(t)\} = sL\{G(t)\} - G(0) = sL\{G(t)\}$
 $= f(s) \therefore L^{-1}\left\{\frac{f(s)}{s}\right\} = G(t) \text{ yields } sL\{G(t)\} = f(s).$

Hence using the Prop. [G] of § 10.9,

$L\{G(t)\} = \frac{f(s)}{s}$ or $L^{-1}\left\{\frac{f(s)}{s}\right\} = G(t) = \int_0^t F(u) du.$

(ii) Let $G(t) = \int_0^t \int_0^v F(u) du dv = L^{-1}\left\{\frac{f(s)}{s^2}\right\}$, then $s^2 L\{G(t)\} = f(s)$

Also $G'(t) = \int_0^t F(u) du$, $G''(t) = F(t)$ and $G(0) = G'(0) = 0$

so that $L\{G''(t)\} = s^2 L\{G(t)\} - sG(0) - G'(0) = s^2 L\{G(t)\} = f(s)$

$\therefore L\{G(t)\} = \frac{f(s)}{s^2}$ or $L^{-1}\left\{\frac{f(s)}{s^2}\right\} = G(t) = \int_0^t \int_0^v F(u) du dv$

This may be written as $L^{-1}\left\{\frac{f(s)}{s^2}\right\} = \int_0^t \int_0^t F(t) (dt)^2$... (10)

or in general $L^{-1}\left\{\frac{f(s)}{s^n}\right\} = \int_0^t \int_0^t \dots \int_0^t F(t) (dt)^n$... (11)

Problem 31. Find $L^{-1} \left\{ \frac{1}{s^2(s^2+1)} \right\}$

$$\therefore L^{-1} \left\{ \frac{1}{s^2+1} \right\} = \sin t, \therefore L^{-1} \left\{ \frac{1}{s(s^2+1)} \right\} = \int_0^t \sin u du = 1 - \cos t$$

$$\text{and hence } L^{-1} \left\{ \frac{1}{s^2(s^2+1)} \right\} = \int_0^t (1 - \cos u) du = t - \sin t.$$

[H] **The Convolution Theorem or the Convolution Property**

If $L^{-1}\{f(s)\} = F(t)$ and $L^{-1}\{g(s)\} = G(t)$, then

$$L^{-1}\{f(s)g(s)\} = \int_0^t F(u)G(t-u)du = F * G \quad \dots (12)$$

where $F * G$ is known as **Convolution or Faltung** of F and G and this convolution is commutative i.e. $F * G = G * F$.

$$\text{The result (12) follows from } f(s)g(s) = L \left\{ \int_0^t F(u)G(t-u)du \right\} \dots (13)$$

which may be proved as below:

$$\therefore f(s) = L\{F(t)\} \text{ and } g(s) = L\{G(t)\}$$

$$\therefore f(s) \cdot g(s) = L\{F(t)\} \cdot L\{G(t)\}$$

$$\begin{aligned} &= \int_0^\infty e^{-sw} F(w) dw \int_0^\infty e^{-sv} G(v) dv \\ &= \int_0^\infty \int_0^\infty e^{-s(w+v)} F(w) G(v) dw dv \\ &= \int_R e^{-s(w+v)} F(w) G(v) dw dv \quad \dots (14) \end{aligned}$$

where R represents the positive quadrant of the plane over which the double integration extends.

In order to transform the double integration from the region R (w, v) to another region $S(x, y)$, let us make the substitutions

$$w = y, v = x - y$$




Fig. 10.2

whose mappings are shown in Fig. 10.2.

When $w=0, y=0, w=\infty, y=\infty$

and when $v=0, x-y=0, v=\infty, (x-y)=\infty$

The Jacobian of the transformation is

$$\frac{\partial(w, v)}{\partial(x, y)} = \begin{vmatrix} \frac{\partial w}{\partial x} & \frac{\partial w}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{vmatrix} = \begin{vmatrix} 0 & 1 \\ 1 & -1 \end{vmatrix} = -1 \text{ giving } dw dv = dx dy.$$

Thus (14) transforms to

$$\begin{aligned} f(s)g(s) &= \iint_S e^{-sx} F(y) G(x-y) dx dy \\ &= \int_0^\infty e^{-sy} \left[\int_0^x F(y) G(x-y) dy \right] dx \\ &= \int_0^\infty e^{-st} \left[\int_0^t F(y) G(t-y) dy \right] dt \text{ (on replacing the variables } y \text{ by } u \text{ and } x \text{ by } t) \\ &= L \left\{ \int_0^t F(u) G(t-u) du \right\} = L\{F * G\} \end{aligned}$$

$$\text{Hence } L^{-1}\{f(s)g(s)\} = \int_0^t F(u)G(t-u)du = F * G$$

Now to show that the operation $F * G$ is commutative, we have

$$\begin{aligned} F * G &= \int_0^t F(u)G(t-u)du = \int_0^t F(t-z)G(z)dz, \text{ when } t-u=z \\ &= \int_0^t G(z)F(t-z)dz = G * F. \quad \dots (15) \end{aligned}$$

Thus this theorem may be restated as

If $f(s)$ and $g(s)$ are the transforms of two functions $F(t)$ and $G(t)$ that are sectionally continuous on each interval $0 \leq t \leq u$, then the transform of the convolution $F(t) * G(t)$ exists and it is $f(s)g(s)$.

e.g. if $F(t)=t$ and $G(t)=e^{at}$ satisfy these conditions, then

$$\begin{aligned} L^{-1} \left\{ \frac{1}{s^2} \cdot \frac{1}{s-a} \right\} &= t * e^{at} = \int_0^t u e^{a(t-u)} du = e^{at} \int_0^t u e^{-au} du \\ &= \frac{1}{a^2} (e^{at} - at - 1). \end{aligned}$$

$$\text{COROLLARY. If } f(s) = L\{F(t)\}, \text{ then } \int_0^t \int_0^v F(u) du dv = \int_0^t (t-u) F(u) du \quad \dots (16)$$

By convolution theorem,

$$L \left\{ \int_0^t (t-u) F(u) du \right\} = L\{t\} L\{F(t)\} = \frac{f(s)}{s^2}$$

$$t * F(t)$$

$$\therefore L\{t\} = \frac{1}{s^2} \text{ and } L\{F(t)\} = f(s)$$

Hence by (9),

$$\int_0^t (t-u) F(u) du = L^{-1} \left\{ \frac{f(s)}{s^2} \right\} = \int_0^t \int_0^v F(v) du dv$$

which may also be expressed as

$$\int_0^t \int_0^t F(t) (dt)^2 = \int_0^t (t-u) F(u) du \quad \dots (17)$$

or in general $\int_0^t \int_0^t \dots \int_0^t F(t) (dt)^n = \int_0^t \frac{(t-u)^{n-1}}{(n-1)!} F(u) du \quad \dots (18)$

Problem 32. Using the convolution theorem, evaluate

$$L^{-1} \left\{ \frac{1}{s^2(s+1)^2} \right\}$$

$$\therefore L^{-1} \left\{ \frac{1}{s^2} \right\} = t \text{ and } L^{-1} \left\{ \frac{1}{(s+1)^2} \right\} = te^{-t}$$

$$\begin{aligned} \therefore L^{-1} \left\{ \frac{1}{s^2(s+1)^2} \right\} &= \int_0^t (ue^{-u}) (t-u) du = \int_0^t (ut-u^2) e^{-u} du \\ &= \left[-(ut-u^2) e^{-u} \right]_0^t + \int_0^t (t-2u) e^{-u} du, \text{ integrating by parts} \\ &= 0 + \int_0^t (t-2u) e^{-u} du = \left[-(t-2u) e^{-u} \right]_0^t - 2 \int_0^t e^{-u} du \\ &= te^{-t} + t - 2 \left[-e^{-u} \right]_0^t = te^{-t} + t + 2e^{-t} - 2 \\ &= (t+2) e^{-t} + t - 2. \end{aligned}$$

10.16. EVALUATION OF SOME INTEGRALS BY INVERSE LAPLACE TRANSFORMS

[1] **The Beta Function.** Which is defined as

$$\beta(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx \text{ for } m > 0, n > 0.$$

The result $\beta(m, n) = \frac{\Gamma m \Gamma n}{\Gamma(m+n)} = \int_0^1 x^{m-1} (1-x)^{n-1} dx$ can be exhibited by the help of inverse Laplace transform as shown below.

$$\text{Let } G(t) = \int_0^t x^{m-1} (t-x)^{n-1} dx$$

By convolution theorem,

$$L\{G(t)\} = L\{t^{m-1}\} \cdot L\{t^{n-1}\} = \frac{\Gamma m}{s^m} \cdot \frac{\Gamma n}{s^n} = \frac{\Gamma m \Gamma n}{s^{m+n}}$$

$$\therefore G(t) = L^{-1} \left\{ \frac{\Gamma m \Gamma n}{s^{m+n}} \right\} = \Gamma m \Gamma n L^{-1} \left\{ \frac{1}{s^{m+n}} \right\} = \Gamma m \Gamma n \frac{t^{m+n-1}}{\Gamma(m+n)}$$

$$\text{i.e. } \int_0^t x^{m-1} (t-x)^{n-1} dx = \frac{\Gamma m \Gamma n}{\Gamma(m+n)} t^{m+n-1} \quad \dots (1)$$

Putting $t=1$, this reduces to

$$\int_0^1 x^{m-1} (1-x)^{n-1} dx = \frac{\Gamma m \Gamma n}{\Gamma(m+n)} = \beta(m, n) \quad \dots (2)$$

Problem 33. Show that $\int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta = \frac{\Gamma m \Gamma n}{2\Gamma(m+n)} = \frac{1}{2} \beta(m, n).$

$$\text{We have } \beta(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$$

Putting $x = \sin^2 \theta$, this gives

$$\beta(m, n) = 2 \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta = \frac{\Gamma m \Gamma n}{\Gamma(m+n)}$$

$$\text{i.e. } \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta = \frac{\Gamma m \Gamma n}{2\Gamma(m+n)} = \frac{1}{2} \beta(m, n)$$

[2] **Evaluation of** $\int_0^t J_0(u) J_0(t-u) du$

$$\text{Let } G(t) = \int_0^t J_0(u) J_0(t-u) du$$

By convolution theorem,

$$L\{G(t)\} = L\{J_0(t)\} \cdot L\{J_0(t)\} = \frac{1}{\sqrt{s^2+1}} \cdot \frac{1}{\sqrt{s^2+1}} = \frac{1}{s^2+1}$$

$$\text{So that } G(t) = L^{-1} \left(\frac{1}{s^2+1} \right)$$

$$\text{i.e. } \int_0^t J_0(u) J_0(t-u) du = \sin t \quad \dots (3)$$

[3] **Evaluation of** $\frac{1}{\pi} \int_{-1}^1 e^{itw} (1-w^2)^{-1/2} dw$

$$\therefore \frac{1}{\sqrt{s^2+1}} = \frac{1}{\sqrt{s+i}} \cdot \frac{1}{\sqrt{s-i}} \text{ and } L\{J_0(t)\} = \frac{1}{\sqrt{s^2+1}}$$

$\therefore$ by convolution theorem

$$\begin{aligned} J_0(t) &= L^{-1} \left\{ \frac{1}{\sqrt{s^2+1}} \right\} = L^{-1} \left\{ \frac{1}{\sqrt{s+i}} \cdot \frac{1}{\sqrt{s-i}} \right\} \\ &= \int_0^t \frac{u^{-1/2} e^{-iu}}{\sqrt{\pi}} \cdot \frac{(t-u)^{-1/2} e^{i(t-u)}}{\sqrt{\pi}} du \quad \because L^{-1} \left\{ \frac{1}{\sqrt{s+a}} \right\} \\ &= \frac{t^{-1/2} e^{-at}}{\sqrt{\pi}} \end{aligned}$$

$$= \frac{1}{\pi} \int_0^t e^{i(t-u)} u^{-1/2} (t-u)^{-1/2} du$$

$$= \frac{1}{\pi} \int_0^t e^{it(1-v)} v^{-1/2} (1-v)^{-1/2} dv \text{ by putting } u=vt$$

$$= \frac{1}{\pi} \int_{-1}^1 e^{itw} (1-w^2)^{-1/2} dw \text{ by putting } 1-2v=w$$

$$\text{Hence } \frac{1}{\pi} \int_{-1}^1 e^{itw} (1-w^2)^{-1/2} dw = J_0(t) \quad \dots (4)$$

[4] Evaluation of $\frac{2}{\sqrt{\pi}} \int_0^{\sqrt{t}} e^{-x^2} dx$.

$$\therefore L^{-1} \left\{ \frac{1}{\sqrt{s}} \right\} = \frac{1}{\sqrt{\pi t}}, \therefore L^{-1} \left\{ \frac{1}{\sqrt{s+1}} \right\} = \frac{e^{-t}}{\sqrt{\pi t}}$$

then by convolution theorem,

$$\begin{aligned} L^{-1} \left\{ \frac{1}{s\sqrt{s+1}} \right\} &= \int_0^t 1 \cdot \frac{e^{-u}}{\sqrt{\pi u}} du \\ &= \frac{2}{\sqrt{\pi}} \int_0^{\sqrt{t}} e^{-x^2} du \text{ by putting } \sqrt{u} = x \text{ i.e. } \frac{1}{\sqrt{u}} du = 2dx \\ &= \text{erf } \sqrt{t} \end{aligned}$$

$$\text{Hence } \frac{2}{\sqrt{\pi}} \int_0^{\sqrt{t}} e^{-x^2} dx = L^{-1} \left\{ \frac{1}{s\sqrt{s+1}} \right\} = \text{erf } \sqrt{t} \quad \dots (5)$$

Problem 34. Show that $\int_0^\infty \cos tx^2 dx = \frac{1}{2} \sqrt{\frac{\pi}{t}}$.

Let $G(t) = \int_0^\infty \cos tx^2 dx$, then taking Laplace transform,

$$\begin{aligned} L\{G(t)\} &= \int_0^\infty e^{-st} dt \int_0^\infty \cos tx^2 dx = \int_0^\infty dx \int_0^\infty e^{-st} \cos tx^2 dt \\ &= \int_0^\infty L\{\cos tx^2\} dx = \int_0^\infty \frac{s}{s^2+x^4} dx \end{aligned}$$

If we put $x^2 = s \tan \theta$ or $x = \sqrt{s} \sqrt{\tan \theta}$, then we get

$$\begin{aligned} L\{G(t)\} &= \frac{1}{2\sqrt{s}} \int_0^{\pi/2} \frac{d\theta}{\sqrt{\tan \theta}} = \frac{1}{2\sqrt{s}} \int_0^{\pi/2} \sin^{-1/2} \theta \cos^{+1/2} \theta d\theta \\ &= \frac{1}{2\sqrt{s}} \frac{\Gamma \frac{1}{2} \Gamma \frac{3}{2}}{2\Gamma 1} \text{ by Problem 33.} \\ &= \frac{1}{2\sqrt{s}} \frac{\pi}{2 \sin \frac{\pi}{4}} \text{ using } \Gamma n \Gamma(1-n) = \frac{\pi}{\sin n\pi}, 0 < n < 1 \\ &= \frac{1}{2\sqrt{s}} \cdot \frac{\pi \sqrt{2}}{2} = \frac{\pi \sqrt{2}}{4\sqrt{s}} \end{aligned}$$

Using inverse transform,

$$\begin{aligned} G(t) &= \int_0^\infty \cos tx^2 dx = \frac{\pi \sqrt{2}}{4} L^{-2} \left\{ \frac{1}{\sqrt{s}} \right\} = \frac{\pi \sqrt{2}}{4} \cdot \frac{t^{-1/2}}{\sqrt{\pi}} \\ &= \frac{\sqrt{2\pi}}{4} t^{-1/2} \end{aligned}$$

Putting $t=1$, we get

$$\int_0^\infty \cos x^2 dx = \frac{\sqrt{2\pi}}{4} = \frac{1}{2} \sqrt{\frac{\pi}{2}}$$

Problem 35. Show that $\int_0^\infty e^{-x^2} dx = \frac{1}{2} \sqrt{\pi}$

Take $G(t) = \int_0^\infty e^{-tx^2} dx$ so that

$$L\{G(t)\} = \int_0^\infty \frac{dx}{s+x^2} = \left[\frac{1}{\sqrt{s}} \tan^{-1} \frac{x}{\sqrt{s}} \right]_0^\infty = \frac{\pi}{2\sqrt{s}}$$

Taking inverse,

$$G(t) = \int_0^\infty e^{-tx^2} dx = \frac{\pi}{2} \frac{t^{-1/2}}{\sqrt{\pi}} = \frac{1}{2} \sqrt{\frac{\pi}{t}} t^{-1/2}$$

$$\text{Putting } t=1, \int_0^\infty e^{-x^2} dx = \frac{\sqrt{\pi}}{2}$$

10.17. SOME METHODS FOR FINDING INVERSE LAPLACE TRANSFORM

[1] Partial Fraction-method

Consider a rational fraction $\frac{N(s)}{D(s)}$ where $N(s)$ and $D(s)$ are polynomials in s with no common factor and the degree of $N(s)$ is lower than degree of $D(s)$.

$$\text{Let } f(s) = \frac{N(s)}{D(s)} \quad \dots (1)$$

Its inverse Laplace transform can be determined whenever the elementary factors of $D(s)$ are known.

(i) Taking first the case when $D(s)$ has no repeated factors, we can write

$$D(s) = (s-\alpha_1)(s-\alpha_2)\dots(s-\alpha_n)$$

where all α_r 's are distinct and coefficient of s^n being assumed unity without any loss of generality. Here $D(s)$ has been taken as a polynomial of degree n so that $r=1, 2, 3, \dots, n$.

By theory of partial fractions, we have

$$\begin{aligned} f(s) &= \frac{N(s)}{D(s)} = \frac{N(s)}{(s-\alpha_1)(s-\alpha_2)\dots(s-\alpha_n)} \\ &= \frac{A_1}{s-\alpha_1} + \frac{A_2}{s-\alpha_2} + \dots + \frac{A_r}{s-\alpha_r} + \dots + \frac{A_n}{s-\alpha_n} \end{aligned}$$

Multiplying both sides by $(s-\alpha_r)$ and taking the limit as $s \rightarrow \alpha_r$, we have

$$A_r = \lim_{s \rightarrow \alpha_r} \frac{N(s) \cdot (s-\alpha_r)}{D(s)} = N(\alpha_r) \cdot \lim_{s \rightarrow \alpha_r} \frac{s-\alpha_r}{D(s)}$$

$$= N(\alpha_r) \cdot \lim_{s \rightarrow \alpha_r} \frac{1}{D'(s)} \text{ by L' Hospital's rule}$$

$$= \frac{N(\alpha_r)}{D'(\alpha_r)}$$

$$\begin{aligned} \text{Thus } f(s) &= \frac{N(s)}{D(s)} = \frac{N(\alpha_1)}{D'(\alpha_1)(s-\alpha_1)} + \dots + \frac{N(\alpha_r)}{D'(\alpha_r)(s-\alpha_r)} \\ &\quad + \dots + \frac{N(\alpha_n)}{D'(\alpha_n)(s-\alpha_n)} \\ &= \sum_{r=1}^n \frac{N(\alpha_r)}{D'(\alpha_r)} \cdot \frac{1}{s-\alpha_r} \quad \dots (2) \end{aligned}$$

where $D'(\alpha_r) = (s-\alpha_1)(s-\alpha_2)\dots(s-\alpha_{r-1})(s-\alpha_{r+1})\dots(s-\alpha_n)$ at $s=\alpha_r$.

If $f(t)$ be the inverse Laplace transform of $f(s)$, then

$$\begin{aligned} F(t) &= L^{-1}\{f(s)\} = L^{-1}\left\{\sum_{r=1}^n \frac{N(\alpha_r)}{D'(\alpha_r)} \cdot \frac{1}{s-\alpha_r}\right\} \\ &= \sum_{r=1}^n \frac{N(\alpha_r)}{D'(\alpha_r)} e^{\alpha_r t} \quad \dots (3) \end{aligned}$$

which is known as *Heaviside Expansion formula*.

(ii) Now taking the case when $D(s)$ has repeated linear factors, we can write

$$f(s) = \frac{N(s)}{D(s)} = \frac{\phi(s)}{(s-\alpha)^{n+1}} \quad \dots (4)$$

where $\phi(s)$ is the quotient of polynomials $N(s)$ and the one obtained by removing the factor $(s-\alpha)^{n+1}$ from $D(s)$. Then, we have

$$\begin{aligned} \frac{\phi(s)}{(s-\alpha)^{n+1}} &= \frac{A_0}{s-\alpha} + \frac{A_1}{(s-\alpha)^2} + \dots + \frac{A_r}{(s-\alpha)^{r+1}} + \dots + \\ &\quad \frac{A_n}{(s-\alpha)^{n+1}} + g(s) \end{aligned}$$

where $g(s)$ denotes the sum of partial fractions corresponding to the other factors of $D(s)$.

Multiplying both sides by $(s-\alpha)^{n+1}$ we get

$$\phi(s) = A_0(s-\alpha)^n + A_1(s-\alpha)^{n-1} + \dots + A_r(s-\alpha)^{r-n} + \dots + A_n + g(s)(s-\alpha)^{n+1}$$

Taking the limit as $s \rightarrow \alpha$, $A_n = \phi(\alpha)$ and so we have on differentiating both sides $(n-r)$ times w.r.t. s ,

$$\phi^{(n-r)}(\alpha) = \frac{(n-r)!}{(n-r)!} A_r \text{ i.e. } A_r = \frac{\phi^{(n-r)}(\alpha)}{(n-r)!}$$

Hence from (4)

$$f(s) = \sum_{r=0}^n \frac{\phi^{(n-r)}(\alpha)}{(n-r)!} \cdot \frac{1}{(s-\alpha)^{r+1}} + g(s) \quad \dots (5)$$

$$\therefore \phi^{(0)}(\alpha) = \phi(\alpha)$$

$\therefore$ Taking the inverse transform, we have

$$F(t) = L^{-1}\{f(s)\} = \sum_{r=0}^n \frac{\phi^{(n-r)}(\alpha)}{(n-r)!} \cdot \frac{t^r e^{\alpha t}}{r!} + G(t) \quad \dots (6)$$

by using the relations $L\{e^{\lambda t} F(t)\} = f(s-\lambda)$ and $L\{t^{r+1}\} = \Gamma(r+1) s^{-r-1}$

Also here, $L^{-1}\{g(s)\} = G(t)$

Applying Leibnitz theorem for differentiation of product of two functions, (6) can be written as

$$F(t) = \frac{1}{n!} \left\{ \frac{\partial^n}{\partial s^n} \phi(s) e^{st} \right\}_{s=\alpha} + G(t) \quad \dots (7)$$

(iii) Taking the case when $D(s)$ has non-repeated quadratic factors of the form $(s-\alpha)^2 + \beta^2$, $s > 0$ and α, β being real numbers, then we can write

$$f(s) = \frac{N(s)}{D(s)} = \frac{\phi(s)}{(s-\alpha)^2 + \beta^2} = \frac{A_1 s + A_2}{(s-\alpha)^2 + \beta^2} + g(s) \quad \dots (8)$$

where $g(s)$ denotes the sum of partial fractions corresponding to the other factors of $D(s)$ and A_1, A_2 are real constants. Then we have

$$\phi(\alpha + i\beta) = \lim_{s \rightarrow \alpha + i\beta} \phi(s) = (\alpha + i\beta) A_1 + A_2 = \phi_1 + i\phi_2 \text{ (say)}$$

i.e., $(\alpha A_1 + A_2) + i\beta A_1 = \phi_1 + i\phi_2$, ϕ_1, ϕ_2 being real and imaginary parts of ϕ .

Equating real and imaginary parts,

$$\phi_1 = \alpha A_1 + A_2 \text{ and } \phi_2 = \beta A_1 \text{ giving } A_1 = \frac{\phi_2}{\beta} \text{ and } A_2 = \phi_1 - \frac{\alpha}{\beta} \phi_2.$$

Thus (8) reduces to

$$f(s) = \frac{1}{\beta} \cdot \frac{(s-\alpha)\phi_2 + \beta\phi_1}{(s-\alpha)^2 + \beta^2} + g(s) \quad \dots (9)$$

Taking the inverse transform, we have

$$F(t) = L^{-1}\{f(s)\} = \frac{1}{\beta} (\phi_2 \cos \beta t + \phi_1 \sin \beta t) e^{\alpha t} + G(t) \quad \dots (10)$$

where $G(t) = L^{-1}\{g(s)\}$.

(iv) Lastly taking the case when $D(s)$ has the square of a quadratic function of the form $\{(s-\alpha)^2 + \beta^2\}^2$, then we can write,

$$\begin{aligned} f(s) &= \frac{N(s)}{D(s)} = \frac{\phi(s)}{\{(s-\alpha)^2 + \beta^2\}^2} = \frac{A_1 s + A_2}{(s-\alpha)^2 + \beta^2} \\ &\quad + \frac{A_3 s + A_4}{\{(s-\alpha)^2 + \beta^2\}^2} + g(s) \quad \dots (11) \end{aligned}$$

where $G(s)$ denotes the sum of partial fractions corresponding to the other factors of $D(s)$ and A_1, A_2, A_3, A_4 are real constants. Then we have

$$\phi(s) = (A_1 s + A_2) \{(s-\alpha)^2 + \beta^2\} + (A_3 s + A_4) + g(s) \{(s-\alpha)^2 + \beta^2\}^2$$

$$\text{So that } \phi(\alpha + i\beta) = \lim_{s \rightarrow \alpha + i\beta} \phi(s) = A_3(\alpha + i\beta) + A_4 = \phi_1 + i\phi_2 \text{ (say)}$$

Giving $A_3\alpha + A_4 = \phi_1$ and $\beta A_3 = \phi_2$ which yield

$$A_3 = \frac{\phi_2}{\beta} \text{ and } A_4 = \frac{1}{\beta}(\beta\phi_1 - \alpha\phi_2)$$

$$\text{Also } \phi'(s) = A_1[(s-\alpha)^2 + \beta^2] + 2(A_1s + A_2)(s-\alpha) + \frac{d}{ds}[g(s)\{(s-\alpha)^2 + \beta^2\}]$$

so that $\phi'(\alpha + i\beta) = 2i\beta[A_1(\alpha + i\beta) + A_2] = \phi_1' + i\phi_2'$ (say)

giving $-2\beta^2 A_1 = \phi_1'$ and $2\alpha\beta A_1 + 2\beta A_2 = \phi_2'$

which yield $A_1 = -\frac{1}{2\beta^2}\phi_1'$ and $A_2 = \frac{\beta\phi_2' + \alpha\phi_1'}{2\beta^2}$.

Hence (11) reduce to

$$f(s) = \frac{1}{2\beta^2} \frac{(-\phi_1's + \beta\phi_2' + \alpha\phi_1')}{(s-\alpha)^2 + \beta^2} + \frac{1}{\beta} \frac{(\phi_2s + \beta\phi_1 - \alpha\phi_2)}{\beta[(s-\alpha)^2 + \beta^2]} + g(s) \quad \dots (12)$$

Taking inverse Laplace transform, we get

$$F(t) = L^{-1}\{f(s)\} = \frac{1}{2\beta^2} e^{\alpha t} [|\phi(\alpha + i\beta)| \{\sin(\beta t + \psi) - \beta t \cos(\beta t + \psi)\} - \beta\{|\phi'(\alpha + i\beta)| \cos(\beta t + \psi')\}] \quad \dots (13)$$

where $|\phi(\alpha + i\beta)| = \sqrt{\phi_1'^2 + \phi_2'^2}$ and $|\phi'(\alpha + i\beta)| = \sqrt{\phi_1'^2 + \phi_2'^2}$, giving

$$\psi = \tan^{-1} \frac{\phi_2'}{\phi_1'} \text{ and } \psi' = \tan^{-1} \frac{\phi_2'}{\phi_1'}$$

As a generalization if $D(s)$ has $(n+1)$ power of the quadratic factors $\{(s-\alpha)^2 + \beta^2\}$ then corresponding term in $F(t)$ will be of the form $t^r e^{\alpha t} \sin \beta t$ and $t^r e^{\alpha t} \cos \beta t$ where $r=0, 1, 2, \dots, n$.

Problem 36. Find the inverse Laplace transform of the following functions

$$(i) \frac{1}{(s+1)(s-2)}, \quad (ii) \frac{3s+1}{(s-1)(s^2+1)},$$

$$(iii) \frac{1}{(s-2)(s+2)^2}, \quad (iv) \frac{4s}{(s-2)(s^2+4)}.$$

$$(i) \text{ Here } f(s) = \frac{1}{(s+1)(s-2)} = \frac{1}{3(s+1)} - \frac{1}{3(s-2)}$$

$$\therefore L^{-1}\{f(s)\} = \frac{1}{3}[e^{-t} - e^{2t}].$$

$$(ii) \text{ Here } f(s) = \frac{3s+1}{(s-1)(s^2+1)} = \frac{3s+1}{(s-1)(s+i)(s-i)} = \frac{N(s)}{D(s)} \text{ (say)}$$

Then $N(s) = 3s+1$, $D(s) = s^3 - s^2 + s - 1$ so that $D'(s) = 3s^2 - 2s + 1$ and zeros $\alpha_1 = 1$, $\alpha_2 = i$, $\alpha_3 = -i$

Using Heaviside's expansion formula, we have

$$L^{-1}\{f(s)\} = \frac{N(1)}{D'(1)} e^t + \frac{N(i)}{D'(i)} e^{it} + \frac{N(-i)}{D'(-i)} e^{-it}$$

$$\begin{aligned} &= \frac{4e^t}{2} + \frac{3i+1}{-(2+2i)} e^{it} + \frac{-3i+1}{-2+2i} e^{-it} \\ &= 2e^t - \frac{(3i+1)(1-i)}{2(1+i)(1-i)} e^{it} + \frac{(3i-1)(1+i)}{2(1-i)(1+i)} e^{-it} \\ &= 2e^t - \frac{1}{2}(i+2)e^{it} + \frac{1}{2}(i-2)e^{-it} \\ &= 2e^t - \frac{1}{2}(e^{it} - e^{-it}) - (e^{it} + e^{-it}) \\ &= 2e^t - \sin t - 2 \cos t. \end{aligned}$$

$$(iii) \text{ Here } f(s) = \frac{1}{(s-2)(s+2)^2}.$$

The term in the inverse transform of $f(s)$ corresponding to $(s-2)$ is $\frac{1}{16} e^{2t}$ and corresponding to $(s+2)^2$ we have $\phi(s) = (s-2)^{-1}$ which yields, $\phi(-2) = -\frac{1}{4}$ and $\phi'(s) = -\frac{1}{(s-2)^2}$ yields $\phi'(-2) = -\frac{1}{16}$

so that by (5) of § 10.17,

$$\begin{aligned} L^{-1}\{f(s)\} &= \frac{1}{16} e^{2t} + \frac{\phi'(-2)}{1} \frac{e^{-2t}}{1} + \frac{\phi(-2)te^{-2t}}{0} \\ &= \frac{1}{16} e^{2t} - \frac{1}{16} e^{-2t} - \frac{1}{4} te^{-2t} \\ &= \frac{1}{16} [e^{2t} - e^{-2t} - 4te^{-2t}] \end{aligned}$$

$$(iv) \text{ Here } f(s) = \frac{4s}{(s-2)(s^2+4)}.$$

The term in the inverse transform corresponding to $(s-2)$ is e^{2t} and corresponding to s^2+4 we have $\phi(s) = \frac{4s}{s-2}$, which yields $\phi(2i) = 2(1-i)$ so that its real and imaginary parts are $\phi_1 = 2$, $\phi_2 = -2$.

Also comparing with (9) of § 10.17, we have $\alpha = 0$ and $\beta = 2$

$$\begin{aligned} \text{Hence } L^{-1}\{f(s)\} &= e^{2t} + \frac{1}{2} L^{-1}\left\{\frac{-2s}{s^2+4} + \frac{4}{s^2+4}\right\} \\ &= e^{2t} + \frac{1}{2}[-2 \cos 2t + 2 \sin 2t] \\ &= e^{2t} + (\sin 2t - \cos 2t) = e^{2t} + \sqrt{2} \sin\left(2t - \frac{\pi}{4}\right). \end{aligned}$$

[2] Series Expansion Method

If $f(s)$ has a series expansion in inverse powers of s such as

$$f(s) = \frac{a_0}{s} + \frac{a_1}{s^2} + \frac{a_2}{s^3} + \dots \quad \dots (14)$$

then under suitable conditions, we can invert term by term to find

$$F(t) = a_0 + a_1 t + \frac{a_2 t^2}{2} + \dots \quad \dots (15)$$

Problem 37. Find $L^{-1} \left\{ \frac{e^{-1/s}}{s} \right\}$

We have $L^{-1} \left\{ \frac{e^{-1/s}}{s} \right\} = L^{-1} \left\{ \frac{1}{s} \left(1 - \frac{1}{s} + \frac{1}{2 \cdot s^2} - \frac{1}{3 \cdot s^3} + \dots \right) \right\}$

$$= L^{-1} \left\{ \frac{1}{s} \right\} - L^{-1} \left\{ \frac{1}{s^2} \right\} - \frac{1}{2} L^{-1} \left\{ \frac{1}{s^3} \right\} - \frac{1}{3} L^{-1} \left\{ \frac{1}{s^4} \right\} + \dots$$

$$= 1 - t + \frac{t^2}{(2)^2} - \frac{t^3}{(3)^2} + \dots$$

$$= 1 - \frac{(2\sqrt{t})^2}{2^2} + \frac{(2\sqrt{t})^4}{2^2 \cdot 4^2} - \frac{(2\sqrt{t})^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots$$

$$= J_0(2\sqrt{t}).$$

[3] Method of Differential Equations

By solving differential equations, the inverse Laplace transforms of some functions can be found as is evident from the following Problem.

Problem 38. Find $L^{-1} \left\{ e^{-\sqrt{s}} \right\}$

Let $y(s) = e^{-\sqrt{s}}$ and $L \{ Y(t) \} = y(s)$.

So that $L^{-1} \{ y(s) \} = L^{-1} \{ e^{-\sqrt{s}} \} = Y(t)$

Also $\frac{dy}{ds} = y' = -\frac{e^{-\sqrt{s}}}{2\sqrt{s}}$ and $\frac{d^2y}{ds^2} = y'' = \frac{e^{-\sqrt{s}}}{4s} + \frac{e^{-\sqrt{s}}}{4s^{3/2}}$

$$\therefore 4sy'' = e^{-\sqrt{s}} + \frac{e^{-\sqrt{s}}}{\sqrt{s}} = y - 2y'$$

$$\therefore 4sy'' + 2y' - y = 0 \quad \dots (1)$$

which is the differential equation satisfied by $y = e^{-\sqrt{s}}$.

Now $y = L \{ Y \}$ gives $y' = L \{ -tY \}$ and $y'' = L \{ (-1)^2 t^2 Y \} = L \{ t^2 Y \}$

So that $sy'' = L \left\{ \frac{d}{dt} (t^2 Y) \right\} = L \{ t^2 Y' + 2tY \}$

Thus (1) reduces to

$$4L \{ t^2 Y' + 2tY \} + 2L \{ -tY \} - L \{ Y \} = 0$$

or $L \{ 4t^2 Y' + (6t-1)Y \} = 0$ or $4t^2 Y' + (6t-1)Y = L^{-1} \{ 0 \} = 0$

i.e. $4t^2 \frac{dY}{dt} + (6t-1)Y = 0$ which gives on separating the variables,

$$\frac{dY}{Y} + \frac{6t-1}{4t^2} dt = 0 \quad \text{or} \quad \frac{dY}{Y} + \left(\frac{3}{2t} - \frac{1}{4t^2} \right) dt = 0$$

or $\frac{dY}{Y} = - \left(\frac{3}{2t} - \frac{1}{4t^2} \right) dt$

Integrating, $\log Y = -\frac{3}{2} \log t - \frac{1}{4t} + \log C$, C being a constant of integration

$$\therefore Y = C \frac{e^{-1/4t}}{t^{3/2}} \quad \text{or} \quad tY = \frac{C}{\sqrt{t}} e^{-1/4t} \quad \dots (2)$$

Now $L \{ tY \} = -\frac{d}{ds} \{ y \} = -\frac{d}{ds} (e^{-\sqrt{s}}) = \frac{e^{-\sqrt{s}}}{2\sqrt{s}}$

Applying the final value Theorem, i.e. when $t \rightarrow \infty$, $s \rightarrow 0$, we have from (2)

$$tY = \frac{C}{\sqrt{t}} e^{-1/4t} \rightarrow \frac{C}{\sqrt{t}} \quad \text{so that} \quad L \{ tY \} \rightarrow L \left\{ \frac{C}{\sqrt{t}} \right\} = C \sqrt{\pi} / \sqrt{s}$$

and $L \{ tY \} = \frac{e^{-\sqrt{s}}}{2\sqrt{s}} \rightarrow \frac{1}{2\sqrt{s}}$, whence $\frac{C \sqrt{\pi}}{\sqrt{s}} = \frac{2}{2\sqrt{s}}$ gives

$$C = \frac{1}{2\sqrt{\pi}}.$$

Hence the solution of (1) is $Y = \frac{1}{2\sqrt{\pi} t^{3/2}} \cdot e^{-1/4t}$

i.e. $L^{-1} \{ e^{-\sqrt{s}} \} = e^{-1/4t} / 2\sqrt{\pi} t^{3/2}.$

[4] Method of Differentiation with Respect to a Parameter

This method is illustrated with the help of following Problem.

Problem 39. Find $L^{-1} \left\{ \frac{1}{(s^2 + a^2)^{3/2}} \right\}$

We have $L \{ J_0(at) \} = \frac{1}{\sqrt{s^2 + a^2}}$.

Differentiating w.r.t. the parameter 'a' we get

$$\frac{d}{da} L \{ J_0(at) \} = \frac{d}{da} \left[\frac{1}{\sqrt{s^2 + a^2}} \right] \quad \text{or} \quad L \left\{ \frac{d}{da} J_0(at) \right\} = \frac{-a}{(s^2 + a^2)^{3/2}}$$

or $L \{ t J_0'(at) \} = \frac{-a}{(s^2 + a^2)^{3/2}}$ i.e. $L^{-1} \left\{ \frac{1}{(s^2 + a^2)^{3/2}} \right\} = -\frac{t}{a} J_0'(at)$

$$= \frac{t}{a} J_1(at) \quad \therefore J_0'(x) = -J_1(x)$$

10.18. APPLICATIONS OF LAPLACE TRANSFORMS TO DIFFERENTIAL EQUATIONS

The Laplace transforms can be successfully applied to solve

(i) ordinary differential equations with constant coefficients such as

$$\frac{d^2y}{dt^2} + a \frac{dy}{dt} + by = F(t), \quad a, b \text{ being constants.}$$

(ii) ordinary differential equations with variable coefficients such as

$$t^2 \frac{d^2y}{dt^2} + t \frac{dy}{dt} + y = F(t)$$

(iii) simultaneous ordinary differential equations such as

$$\frac{d^2y}{dt^2} + a \frac{dy}{dx} + b \frac{dx}{dt} = F_1(t) \quad \text{and} \quad C \frac{dy}{dt} + \frac{d^2x}{dt^2} + d \frac{dx}{dt} = F_2(t);$$

a, b, c, d being constants.

(iv) partial differential equations such as

$$\frac{\partial^2 y}{\partial t^2} + a \frac{\partial^2 y}{\partial x^2} + y = F(x, t)$$

(v) integral equations i.e. the equation containing a dependent variable under an integral sign, such as

$$Y(t) = F(t) + \int_a^b K(u, t) Y(u) du, \quad a, b \text{ being constants or functions of } t.$$

Here $F(t)$ and the Kernel $K(u, t)$ are known and $Y(t)$ is to be determined.

We explain these applications with the help of following problems.

Problem 40. Solve $x''(t) + 4x'(t) + 4x(t) = 4e^{-2t}$; $x(0) = -1$, $x'(0) = 4$ verify that your solution satisfies the differential equation and the boundary conditions.

$$\text{Let } L\{x(t)\} = f(s) \quad \dots (1)$$

$$\text{Then } L\{x'(t)\} = sf(s) - F(0) \text{ where } F(0) = x(0) = -1$$

$$= sf(s) + 1 \quad \dots (2)$$

$$\text{and } L\{x''(t)\} = s^2 f(s) - sF(0) - F'(0), \text{ where } F(0) = x(0) = -1$$

$$\text{and } F'(0) = x'(0) = 4$$

$$= s^2 f(s) + s - 4 \quad \dots (3)$$

Taking Laplace transform of the given equation, we get

$$L\{x''(t)\} + 4L\{x'(t)\} + 4L\{x(t)\} = 4L\{e^{-2t}\}$$

$$\text{or } s^2 f(s) + s - 4 + 4sf(s) + 4 + 4f(s) = \frac{4}{s+2} \text{ by (1), (2) and (3)}$$

$$\text{or } (s^2 + 4s + 4)f(s) = \frac{4}{s+2} - s \text{ i.e. } f(s) = \frac{4}{(s+2)^3} - \frac{s}{(s+2)^2}$$

$$= \frac{4}{(s+2)^3} + \frac{2}{(s+2)^2} - \frac{1}{s+2} \quad \because s = (s+2) - 2$$

Taking inverse Laplace transform of both sides, we find

$$L^{-1}\{f(s)\} = 4L^{-1}\left\{\frac{1}{(s+2)^3}\right\} + 2L^{-1}\left\{\frac{1}{(s+2)^2}\right\} - L^{-1}\left\{\frac{1}{s+2}\right\}$$

$$\text{i.e. } x(t) = \frac{4(-1)^2}{2} L^{-1}\left\{\frac{d^2}{ds^2}\left(\frac{1}{s+2}\right)\right\} + 2(-1) L^{-1}\left\{\frac{d}{ds}\left(\frac{1}{s+2}\right)\right\} - e^{-2t}$$

$$= 2(-1)^2 \cdot (-1)^2 t^2 e^{-2t} + 2(-1)(-1) t e^{-2t} - e^{-2t} = e^{-2t}(2t^2 + 2t - 1)$$

Hence the required solution is

$$x(t) = e^{-2t}(2t^2 + 2t - 1) \quad \dots (4)$$

Now to verify the result, we have from (4)

$$x'(t) = -2e^{-2t}(2t^2 + 2t - 1) + e^{-2t}(4t + 2) = e^{-2t}(-4t^2 - 4t + 2 + 4t + 2)$$

$$= 4e^{-2t}(-t^2 + 1) \quad \dots (5)$$

$$\text{and } x''(t) = -8e^{-2t}[-t^2 + 1] + 4e^{-2t}(-2t) = 8e^{-2t}(t^2 - t - 1)$$

$$\text{So that } x''(t) + 4x'(t) + 4x(t) = 8e^{-2t}(t^2 - t - 1) + 16e^{-2t}(-t^2 + 1)$$

$$+ 4e^{-2t}(2t^2 + 2t - 1) = 4e^{-2t} \dots (6)$$

which is the same as given and hence the result is verified.

Again from (4), we have $x(0) = -1$ and from (5), $x'(0) = 4$ which clearly satisfy the boundary conditions.

Problem 41. Solve the differential equation $ty''(t) + y'(t) + ty(t) = 0$.

Under the condition that $y(0) = 1$ and $y(t)$ is bounded.

$$\text{Let } L\{y(t)\} = f(s) \text{ so that } L^{-1}\{f(s)\} = y(t) \quad \dots (1)$$

$$\text{Now } L\{y'(t)\} = sf(s) - F(0), \text{ where } F(0) = y(0) = 1$$

$$= sf(s) - 1 \quad \dots (2)$$

$$\text{and } L\{y''(t)\} = s^2 f(s) - sF(0) - F'(0) \text{ where } F(0) = y(0) = 1$$

$$\text{and } F'(0) = y'(0) = k \quad (\text{const.}) \text{ say}$$

$$= s^2 f(s) - s - k.$$

Taking Laplace transform of the given equation,

$$L\{ty''(t)\} + L\{y'(t)\} + L\{ty(t)\} = 0$$

$$\text{or } -\frac{d}{ds} L\{y''(t)\} + L\{y'(t)\} - \frac{d}{ds} L\{y(t)\} = 0$$

$$\text{or } -\frac{d}{ds} \{s^2 f(s) - s - k\} + sf(s) - 1 - \frac{d}{ds} \{f(s)\} = 0 \text{ by (1) and (2)}$$

$$\text{or } -2sf(s) - s^2 f'(s) + 1 + sf(s) - 1 - f'(s) = 0$$

$$\text{or } -sf(s) - (s^2 + 1)f'(s) = 0$$

$$\text{or } \frac{f'(s)}{f(s)} = -\frac{s}{s^2 + 1} = -\frac{1}{2} \cdot \frac{2s}{s^2 + 1}$$

Integrating, $\log f(s) = -\frac{1}{2} \log(s^2 + 1) + \log C$, C being a constant of integration

$$\therefore f(s) = \frac{C}{\sqrt{s^2 + 1}} \quad \dots (3)$$

Taking inverse Laplace transform, we have

$$L^{-1}\{f(s)\} = C L^{-1}\left\{\frac{1}{\sqrt{s^2 + 1}}\right\} \text{ or } y(t) = C J_0(t)$$

$$\text{Initially when } t=0, y(t) = y(0) = 1, \text{ so that } 1 = C J_0(0) = C$$

$$\therefore J_0(0) = 1.$$

Hence the required solution is $y(t) = J_0(t)$.

Problem 42. Solve $X'' + Y' + 3X = 15e^{-t}$

$$Y'' - 4X' + 3Y = 15 \sin 2t.$$

Subject to $X(0)=35$, $X'(0)=-48$, $Y(0)=27$, $Y'(0)=-55$.

Let $L\{X\}=x$ and $L\{Y\}=y$ so that $L^{-1}\{x\}=X$ and $L^{-1}\{y\}=Y$.

Also $L\{X'\}=sL\{X\}-X(0)=sx-35$; $L\{X''\}=s^2L\{X\}-sX(0)-X'(0)$
 $=s^2x-35s+48$

and $L\{Y'\}=sL\{Y\}-Y(0)=sy-27$; $L\{Y''\}=s^2L\{Y\}-sY(0)-Y'(0)$
 $=s^2y-27s+55$

$\therefore$ Taking Laplace transforms of the given equations, we have

$$s^2x-35s+48+sy-27+3x=\frac{15}{s+1}$$

$$\text{and } s^2y-27s+55-4(sx-35)+3y=\frac{30}{s^2+4}$$

$$\text{or } (s^2+3)x+sy-\left(35s-21+\frac{15}{s+1}\right)=0$$

$$\text{and } -4sx+(s^2+3)y-\left(27s-195+\frac{30}{s^2+4}\right)=0$$

$$\begin{aligned} \text{Solving } & \frac{x}{-s\left(27s-195+\frac{30}{s^2+4}\right)+(s^2+3)\left(35s-21+\frac{15}{s+1}\right)} \\ & = \frac{y}{4s\left(35s-21+\frac{15}{s+1}\right)+(s^2+3)\left(27s-195+\frac{30}{s^2+4}\right)} \\ & = \frac{1}{(s^2+3)^2+4s^2} \end{aligned}$$

$$\text{So that } x = \frac{1}{s^4+10s^2+9} \left[-27s^2+195s-\frac{30s}{s^2+4}+35s^3-21s^2 \right. \\ \left. + \frac{15s^2}{s+1} + 105s-63 + \frac{45}{s+1} \right]$$

$$\begin{aligned} & = \frac{1}{(s^2+1)(s^2+9)(s+1)(s^2+4)} [(35s^3-48s^2+300s-63) \\ & \quad (s+1)(s^2+4)-30s(s+1)+15(s^2+3)(s^2+4)] \\ & = \frac{35s^3-48s^2-300s-63}{(s^2+1)(s^2+9)} + \frac{15(s^2+3)}{(s+1)(s^2+1)(s^2+9)} \end{aligned}$$

$$\begin{aligned} & = \frac{265s-15}{8(s^2+1)} + \frac{15s-369}{8(s^2+9)} + \frac{3}{s+1} - \frac{15}{8} \cdot \frac{s-1}{s^2+1} - \frac{9}{8} \cdot \frac{s-1}{s^2+9} \\ & \quad - \frac{10s}{(s^2+1)} + \frac{2s}{s^2+4} - \frac{6s}{8(s^2+9)} \end{aligned}$$

(by resolving into partial fractions)

$$= \frac{30s}{s^2+1} - \frac{45}{s^2+9} + \frac{3}{s+1} + \frac{2s}{s^2+4} \quad \dots (1)$$

$$\text{Similarly } y = \frac{30s}{s^2+9} - \frac{60}{s^2+1} - \frac{3}{s+1} + \frac{2}{s^2+4} \quad \dots (2)$$

Taking inverse Laplace transforms of (1) and (2) we get

$$\begin{aligned} L^{-1}\{x\} &= X = 30 \cos t - 45 \cdot \frac{\sin 3t}{3} + 3e^{-t} + 2 \cos 2t \\ &= 30 \cos t - 15 \sin 3t + 3e^{-t} + 2 \cos 2t \end{aligned}$$

$$\begin{aligned} \text{and } L^{-1}\{y\} &= Y = 30 \cos 3t - 60 \sin t - 3e^{-t} + \frac{2 \sin 2t}{2} \\ &= 30 \cos 3t - 60 \sin t - 3e^{-t} + \sin 2t \end{aligned}$$

which are the required solutions.

Problem 43. Find the solution of $\frac{\partial^2 V}{\partial x^2} - \frac{\partial^2 V}{\partial t^2} = xt$ satisfying the conditions $V = \frac{\partial V}{\partial t} = 0$ when $t=0$.

Let $L\{V\}=v$ so that $L^{-1}\{v\}=V$, then

$L\left\{\frac{\partial^2 V}{\partial t^2}\right\} = s^2L\{V\} - sV(0) - V'(0) = s^2v - 0 - 0 = s^2v$, taking t as principal variable

and $L\left\{\frac{\partial^2 V}{\partial x^2}\right\} = \frac{d^2}{dx^2} L\{V\} = \frac{d^2v}{dx^2}$, taking x as the secondary variable.

$$\text{Also } L\{xt\} = \frac{x}{s^2}.$$

$\therefore$ Taking Laplace transform of the given equation, we have

$$\frac{d^2v}{dx^2} - s^2v = \frac{x}{s^2} \quad \text{i.e. } (D^2 - s^2)v = \frac{x}{s^2} \quad \text{where } D \equiv \frac{d}{dx} \quad \dots (1)$$

Its complementary function is $C_1 e^{sx} + C_2 e^{-sx}$

$$\therefore D^2 - s^2 = 0 \text{ gives } D = \pm s$$

$$\begin{aligned} \text{and particular integral} &= \frac{x/s^2}{D^2 - s^2} = \frac{1}{-s^2} \left[1 - \left(\frac{D}{s^2}\right)^2 \right]^{-1} \frac{x}{s^2} \\ &= -\frac{1}{s^2} \left[1 + \frac{D^2}{s^2} + \dots \right] \cdot \frac{x}{s^2} = -\frac{x}{s^4} \end{aligned}$$

$\therefore$ The complete integral of (1) is

$$v = C_1 e^{sx} + C_2 e^{-sx} - \frac{x}{s^4} \quad \dots (2)$$

Now $V=0$ gives $L\{V\}=v=0$ for all values of x , therefore $C_1=0$, otherwise $v \rightarrow \infty$ as $x \rightarrow \infty$.

$$\text{As such (2) reduces to } v = C_2 e^{-sx} - \frac{x}{s^4} \quad \dots (3)$$

But when $x=0$, $v=0 \therefore C_2=0$

$$\text{Hence (3) becomes } v = -\frac{x}{s^4}$$

Taking inverse Laplace transform of both sides, we find
 $L^{-1}\{v\} = L^{-1}\left\{-\frac{x}{s^4}\right\}$ i.e. $V = -\frac{xt^3}{3} = -\frac{xt^3}{6}$ which is the required solution.

Problem 44. Solve the following integral equation and verify your solution.

$$F(t) = 1 + 2 \int_0^t F(t-x) e^{-2x} dx.$$

Taking Laplace transform of both sides, we get

$$L\{F(t)\} = L\{1\} + 2L\left\{\int_0^t F(t-x) e^{-2x} dx\right\}$$

Assuming $L\{F(t)\} = f(s)$ so that $L\{1\} = \frac{1}{s}$

and $L\left\{\int_0^t F(t-x) e^{-2x} dx\right\} = \frac{f(s)}{s+2}$, the last equation becomes,

$$f(s) = \frac{1}{s} + \frac{2f(s)}{s+2}$$

or $\left(1 - \frac{2}{s+2}\right) f(s) = \frac{1}{s}$ or $\frac{1}{s+2} f(s) = \frac{1}{s}$ or $f(s) = \frac{1}{s} \cdot \frac{s+2}{s} = \frac{1}{s} + \frac{2}{s^2}$

Taking inverse Laplace transform of both sides, we find

$$L^{-1}\{f(s)\} = L^{-1}\left\{\frac{1}{s}\right\} + 2L^{-1}\left\{\frac{1}{s^2}\right\} \text{ i.e. } F(t) = 1 + 2t \text{ which is the required solution.}$$

In order to verify the solution, putting

$$F(t) = 1 + 2t \text{ i.e. } F(t-x) = 1 + 2(t-x),$$

we have, R.H.S. of the given equation $= 1 + 2 \int_0^t \{1 + 2(t-x)\} e^{-2x} dx$

$$= 1 + 2 \left[\int_0^t e^{-2x} dx + 2t \int_0^t e^{-2x} dx - 2 \int_0^t x e^{-2x} dx \right]$$

$$= 1 - e^{-2t} + 1 - 2te^{-2t} + 2t + 2te^{-2t} + e^{-2t} - 1 = 1 + 2t = F(t) = \text{L.H.S.}$$

Hence the solution is verified.

10.19. APPLICATIONS OF LAPLACE TRANSFORMS TO BOUNDARY VALUE PROBLEMS

In view of §10.9 (E), it can be concluded that Laplace transforms are capable of transforming the differential equations with constant coefficients into algebraic equations in the form of transformed function. Since various problems used in physics are mathematically formulated in the shape of ordinary and partial differential equations involving one or more unknown functions together with initial or boundary conditions fitting the physical situation, therefore Laplace transform can be successfully applied to such boundary value problems.

Problem 45. A particle of mass 3 gms moves on the x -axis and is attracted towards a fixed point in its path with a force whose numerical value is $12x$. Assuming that the particle is initially at rest at $x=5$, determine the position of the particle at any time t , (a) when there is no other force, (b) when there is a damping force whose numerical value is 6 times the instantaneous velocity.

(a) Let P be the particle and O the fixed point. Since the particle is attracted towards O , therefore when the particle is on the right of




Fig. 10.3

O i.e. in the direction $\vec{OP}$ then the net force is $-12x$ (being in the direction of $\vec{PO}$) and similarly when the particle is moving along $\vec{PO}$, the force being in the direction of OP i.e. again $-12x$. Hence applying Newton's second law of motion i.e. mass $\times$ acceleration $=$ acting force, the equation of motion is $3 \frac{d^2x}{dt^2} = -12x$

$$\text{or} \quad \frac{d^2x}{dt^2} + 4x = 0 \quad \dots (1)$$

Let $L\{x\} = \xi$ so that $L^{-1}\{\xi\} = x$

$$\therefore L\left\{\frac{d^2x}{dt^2}\right\} = s^2 L\{x\} - sx(0) - x'(0) \text{ where } x(0)=5 \text{ and } x'(0)=0 \text{ (boundary conditions)}$$

$$= s^2 \xi - 5s$$

Taking Laplace transform of (1), we therefore get

$$s^2 \xi - 5s + 4\xi = 0 \text{ or } (s^2 + 4)\xi = 5s \text{ or } \xi = \frac{5s}{s^2 + 4}$$

$\therefore L^{-1}\{\xi\} = 5L^{-1}\left\{\frac{s}{s^2 + 4}\right\}$ i.e. $x = 5 \cos 2t$, which gives the position of the particle.

(b) In this case the equation of motion becomes

$$3 \frac{d^2x}{dt^2} = -12x - 6 \frac{dx}{dt} \text{ i.e., } \frac{d^2x}{dt^2} + 2 \frac{dx}{dt} + 4x = 0 \quad \dots (2)$$

with the initial conditions $x(0)=5$, $x'(0)=0$

Taking Laplace transform of (2), $L\left\{\frac{d^2x}{dt^2}\right\} + 2L\left\{\frac{dx}{dt}\right\} + 4L\{x\} = 0$

or $s^2 \xi - 5s + 2(s\xi - 0) + 4\xi = 0$ when $L\{x\} = \xi$

$$\text{or } (s^2 + 2s + 4)\xi = 5s \text{ or } \xi = \frac{5s + 10}{(s+2)^2} = \frac{5}{s+2}$$

$\therefore L^{-1}\{\xi\} = x = 5e^{-2t}$ which is the required solution.

Problem 46. An inductor of H henrys and a capacitor of C farads are in series with a generator of E volts. At $t=0$ the charge on the capacitor and current in the circuit are zero. Find the charge on the capacitor at any time $t > 0$ if $E = E_0$, a constant.

If Q be the instantaneous charge and I the current at any time t , then $I = \frac{dQ}{dt}$... (1)

Now voltage drop across the inductor $= H \frac{dI}{dt} = H \frac{d^2Q}{dt^2}$ by (1)

Voltage drop across the capacitor $= \frac{Q}{C}$ and the voltage drop across

the generator $= -$ voltage rise $= -E$.

Thus, according to Kirchoff's law, the algebraic sum of voltage drops (potential drops) around any closed circuit being zero, the equation determining Q is

$$H \frac{d^2Q}{dt^2} + \frac{Q}{C} - E = 0 \text{ or } \frac{d^2Q}{dt^2} + \frac{Q}{HC} = \frac{E}{H} \quad \dots (2)$$

with the initial (boundary) conditions $Q(0)=0$, $I(0)=0$, $Q'(0)=0$ and $E=E_0$ for $t>0$

Assuming that $L\{Q\}=q$, we have

$$L\left\{\frac{d^2Q}{dt^2}\right\} = s^2 L\{Q\} - sQ(0) - Q'(0) = s^2q - 0 - 0 = s^2q$$

$\therefore$ Laplace transform of (2) gives $L\left\{\frac{d^2Q}{dt^2}\right\} + \frac{1}{HC} L\{Q\} = \frac{E_0}{H} L\{1\}$

$$\text{i.e. } s^2q + \frac{q}{HC} = \frac{E_0}{Hs} \text{ or } \left(s^2 + \frac{1}{HC}\right)q = \frac{E_0}{Hs} \text{ or } q = \frac{E_0/H}{s\left(s^2 + \frac{1}{HC}\right)}$$

$$\text{or } q = \frac{E_0C}{s} - \frac{CE_0s}{s^2 + \frac{1}{HC}} = CE_0 \left\{ \frac{1}{s} - \frac{s}{s^2 + \frac{1}{HC}} \right\}$$

Taking inverse Laplace transform, we have

$$Q = L^{-1}\{q\} = CE_0 \left[L^{-1}\left\{\frac{1}{s}\right\} - L^{-1}\left\{\frac{s}{s^2 + \frac{1}{HC}}\right\} \right] \\ = CE_0 \left[1 - \cos \frac{t}{\sqrt{HC}} \right]$$

which gives the required charge on the capacitor.

Problem 47. A beam which is clamped at its ends $x=0$ and $x=l$ carries a uniform load w per unit length. Show that deflection at any point is $Y(x) = \frac{wx^2(l-x)^2}{24EI}$, where E is Young's modulus of elasticity for the beam and I is the moment of Inertia of a cross-section of the beam about the axis.

The transverse deflection $Y(x)$ at any point x is given by the differential equation

THE LAPLACE TRANSFORMS

$$\frac{d^4Y}{dx^4} = \frac{w}{EI}, \quad 0 < x < l \quad \dots (1)$$

with the initial condition, $y(0)=0$, $y'(0)=0$ and $y(l)=0$, $y'(l)=0$...

Assuming that

$$L\{Y\} = y \text{ i.e. } L^{-1}\{Y\} = Y, \text{ we have} \quad \text{Fig. 10.4} \\ L\left\{\frac{d^4Y}{dx^4}\right\} = s^4 L\{Y\} - s^3Y(0) - s^2Y'(0) - sY''(0) - Y'''(0). \\ = s^4y - 0 - 0 - sC_1 - C_2 \text{ under the assumption that } Y''(0)=C_1 \\ \text{and } Y'''(0)=C_2.$$

$\therefore$ Taking Laplace transform of (1), we have

$$L\left\{\frac{d^4Y}{dx^4}\right\} = \frac{w}{EI} L\{1\} \text{ i.e., } s^4y - sC_1 - C_2 = \frac{w}{EIs} \text{ or } y = \frac{C_1}{s^3} + \frac{C_2}{s^4} + \frac{w}{EIs^5}$$

Taking inverse Laplace transform, we find

$$Y = L^{-1}\{y\} = C_1 \frac{x^2}{2} + C_2 \frac{x^3}{3} + \frac{w}{EI} \frac{x^4}{4} = \frac{C_1x^2}{2} + \frac{C_2x^3}{6} + \frac{wx^4}{24EI} \quad \dots (2)$$

Applying the initial condition that $Y(x)=0$ when $x=l$ and $Y'(x)=0$ when $x=l$, (2) gives

$$0 = \frac{C_1l^2}{2} + \frac{C_2l^3}{6} + \frac{wl^4}{24EI} \quad \dots (3)$$

$$\text{and } Y' = C_1x + \frac{C_2x^2}{2} + \frac{wx^3}{6EI} \text{ gives } 0 = C_1l + \frac{C_2l^2}{2} + \frac{wl^3}{6EI} \quad \dots (4)$$

Solving (3) and (4); $C_1 = \frac{wl^2}{12EI}$ and $C_2 = -\frac{wl}{2EI}$ which when substituted in (2) give

$$Y = \frac{wl^2x^2}{24EI} - \frac{wlx^3}{12EI} + \frac{wx^4}{24EI} = \frac{wx^2}{24EI} [l^2 - 2lx + x^2] = \frac{wx^2(l-x)^2}{24EI}$$

Problem 48. A semi-infinite solid $x>0$ is initially at temperature zero. At time $t=0$, a constant temperature $U_0>0$ is applied and maintained at the face $x=0$. Find the temperature at any point of the solid at any time $t>0$.

Let $U(x, t)$ be the temperature of the solid at any point x and at any time t . Then one-dimensional heat conduction equation gives

$$\frac{\partial^2 U}{\partial x^2} = \frac{1}{k} \frac{\partial U}{\partial t}, \quad x>0, t>0 \quad \dots (1)$$

where k is the diffusivity defined by $\frac{k}{C\rho}$ where k is the thermal conductivity, C the specific heat and ρ the volume density. The boundary conditions are

(i) $U(x, 0)=0$, (ii) $U(0, t)=U_0$ (given) and (iii) U is bounded i.e.,

$$|U(x, t)| < M.$$




Fig. 10.5

Assuming that $L\{U(x, t)\} = u(x, s)$ i.e. $L^{-1}\{u(x, s)\} = U(x, t)$, we have

$$L\left\{\frac{\partial U}{\partial t}\right\} = sL\{U\} - U(x, 0) = su \text{ and } L\left\{\frac{\partial^2 U}{\partial x^2}\right\} = \frac{d^2}{dx^2} [L\{U\}] = \frac{d^2 u}{dx^2}$$

$\therefore$ Taking Laplace transform of (1)

$$\frac{d^2 u}{dx^2} = \frac{1}{k} su \text{ or } \frac{d^2 u}{dx^2} - \frac{s}{k} u = 0 \quad \dots (2)$$

$$\text{with boundary condition } u(0, s) = L\{U(0, t)\} = U_0 L\{1\} = \frac{U_0}{s} \quad \dots (3)$$

and $u = u(x, s)$ is required to be bounded.

$$\text{The solution of (2) is } u(x, s) = C_1 e^{x\sqrt{s/k}} + C_2 e^{-x\sqrt{s/k}} \quad \dots (4)$$

In order that u be bounded (finite) as $x \rightarrow \infty$, we must choose $C_1 = 0$ otherwise the temperature will be infinite when $x \rightarrow \infty$.

$$\text{Thus (4) reduces to } u(x, s) = C_2 e^{-x\sqrt{s/k}}$$

$$\text{But from (3) } u(x, s) = \frac{U_0}{s} \text{ when } x=0, \text{ therefore } C_2 = \frac{U_0}{s}$$

$$\text{Hence the solution of (2) is } u(x, s) = \frac{U_0}{s} e^{-x\sqrt{s/k}}$$

Taking inverse Laplace transform, we find

$$U(x, t) = U_0 \operatorname{erf} C \left(\frac{x}{2\sqrt{kt}} \right) = U_0 \left\{ 1 - \frac{2}{\sqrt{\pi}} \int_0^{x/2\sqrt{kt}} e^{-u^2} du \right\}$$

which gives the required temperature.

Problem 49. An infinitely long string having one end at $x=0$ is initially at rest on the x -axis. The end $x=0$ undergoes a periodic transverse displacement given by $A_0 \sin \omega t$, $t > 0$. Find the displacement of any point on the string at any time t .

Let $Y(x, t)$ be the transverse displacement of the string at any point x and at any time t . Then the transverse displacement is given by

$$\frac{\partial^2 Y}{\partial t^2} = a^2 \frac{\partial^2 Y}{\partial x^2}, \quad x > 0, t > 0 \quad \dots (1)$$

where $a^2 = \frac{T}{\rho}$, T being the constant tension in the string and ρ the constant linear density. The boundary conditions are

(i) $Y(x, 0) = 0$, (ii) $\frac{\partial Y}{\partial t}(x, 0) = 0$, (iii) $Y(0, t) = A_0 \sin \omega t$, and (iv) the




Fig. 10.6

displacement is bounded i.e., $|Y(x, t)| < M$.

Assuming that $L\{Y(x, t)\} = y(x, s)$ i.e., $L^{-1}\{Y(x, s)\} = Y(x, t)$ we have

$$L\left\{\frac{\partial^2 Y}{\partial t^2}\right\} = s^2 L\{Y\} - sY(x, 0) - \frac{\partial Y}{\partial t}(x, 0) = s^2 y - 0 - 0$$

$$\text{and } L\left\{\frac{\partial^2 Y}{\partial x^2}\right\} = \frac{d^2}{dx^2} L\{Y\} = \frac{d^2 y}{dx^2}$$

$\therefore$ Taking Laplace transform of (1), we get

$$s^2 y = a^2 \frac{d^2 y}{dx^2} \text{ i.e., } \frac{d^2 y}{dx^2} - \frac{s^2}{a^2} y = 0 \quad \dots (2)$$

with the boundary condition $Y(0, t) = L\{Y(0, t)\} = L\{A_0 \sin \omega t\}$

$$= \frac{A_0 \omega}{s^2 + \omega^2} \text{ and } y(x, s) \text{ is bounded.} \quad \dots (3)$$

$$\text{The solution of (2) is } y(x, s) = C_1 e^{sx/a} + C_2 e^{-sx/a} \quad \dots (4)$$

But $y(x, s)$ being bounded when $x \rightarrow \infty$, $C_1 = 0$ otherwise the displacement will be infinite when $x \rightarrow \infty$.

$$\therefore (4) \text{ reduces to } y(x, s) = C_2 e^{-sx/a}$$

Applying (3), we find $C_2 = \frac{A_0 \omega}{s^2 + \omega^2}$, therefore the solution of (2) is

$$y(x, s) = \frac{A_0 \omega}{s^2 + \omega^2} e^{-sx/a}$$

Taking inverse Laplace transform, we get

$$y(x, t) = \begin{cases} A_0 \sin \omega(t - x/a), & t > x/a \\ 0, & t < x/a \end{cases}$$

which shows that a point of the string is at rest until $t = x/a$ and there after it undergoes motion identical with that of the end $x=0$ but lags behind it in time by the amount x/a .

10.20. THE COMPLEX INVERSION FORMULA OR INVERSION THEOREM

If $L\{F(t)\} = f(s)$, where $F(t)$ has a continuous derivative and is of exponential order, then

$$F(t) = \int_{\gamma - i\infty}^{\gamma + i\infty} e^{st} f(s) ds, \quad t > 0. \quad \dots (1)$$

In the previous chapter we have introduced that if $\phi(t)$ be a function of t having continuous derivative such that $\int_{-\infty}^{\infty} \phi(t) dt$ is absolutely convergent, then the Fourier's integral is

$$\phi(t) = \frac{1}{\pi} \int_0^{\infty} dx \int_{-\infty}^{\infty} \phi(u) \cos x(t-u) du \quad \dots (2)$$

But $\cos x(t-u)$ being an even function of x and $\sin x(t-u)$ an odd function of x , we can express (2) as

$$\phi(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} \phi(u) \cos x(t-u) du$$

and $0 = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} \phi(u) \sin x(t-u) du$

or $\phi(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} \phi(u) [\cos xt \cos xu + \sin xt \sin xu] du$... (3)

and $0 = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} \phi(u) [\sin xt \cos xu - \cos xt \sin xu] du$... (4)

Multiplying (4) by i and adding to (3) we get

$$\begin{aligned} \phi(t) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} \phi(u) [(\cos xt \cos xu + \sin xt \sin xu) + i(\sin xt \cos xu - \cos xt \sin xu)] du \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} \phi(u) [(\cos xt + i \sin xt)(\cos xu - i \sin xu)] du \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} \phi(u) \cdot e^{ixt} e^{-ixu} du \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ixt} dx \int_{-\infty}^{\infty} \phi(u) e^{-ixu} du \end{aligned}$$

Assuming that $\phi(t) = e^{-\gamma t} F(t)$ for $t > 0$ and $\phi(t) = 0$ for $t < 0$ so that the Integral theorem is also satisfied, we have for $t > 0$

$$\begin{aligned} e^{-\gamma t} F(t) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ixt} dx \int_{-\infty}^{\infty} e^{-\gamma u} F(u) e^{-ixu} du, \gamma > t \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ixt} dx \int_{-\infty}^{\infty} e^{-(\gamma+ix)u} F(u) du \\ \therefore e^{-\gamma t} F(t) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ixt} dx f(\gamma+ix), \text{ put } \gamma+ix=s \text{ so that } dx = i ds \end{aligned}$$

$$\begin{aligned} &= \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{(s-\gamma)t} f(s) ds = \frac{e^{-\gamma t}}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{st} f(s) ds \\ \text{i.e. } F(t) &= \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{st} f(s) ds. \end{aligned}$$

Aliter. By Cauchy's integral formula, mentioned in chapter on complex variables, we have

$$f(s) = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z-s}$$

Assuming that $f(z)$ is regular on the right of $x=\gamma$ and distorting the contour C and integrating from $\gamma+i\infty$ to $\gamma-i\infty$, then to the right of $\gamma-i\infty$ to $\gamma+i\infty$ and back to $\gamma+i\infty$, we get,

$$\begin{aligned} f(s) &= \frac{1}{2\pi i} \int_{\gamma+i\infty}^{\gamma-i\infty} \frac{f(z) dz}{z-s} \\ &= \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{f(z) dz}{s-z} \dots (5) \end{aligned}$$

Now since γ must be smaller than the real part of s i.e. $\gamma < R(s)$ and assuming that $L^{-1}\{f(s)\} = F(t)$, we get on taking Laplace transform of (5),

$$L^{-1}\{f(s)\} = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} f(z) dz \cdot L^{-1}\left\{\frac{1}{s-z}\right\}$$

$$\begin{aligned} \text{i.e., } F(t) &= \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} f(z) e^{zt} dz \quad \because L\{e^{zt}\} = \int_0^\infty e^{-st} e^{zt} dt \\ &= \frac{1}{s-z} \text{ if } R(s) > R(z) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} f(s) e^{st} ds \text{ (on replacing } z \text{ by } s) \\ &= \text{sum of residues of } e^{st} f(s) \text{ at the poles of } f(s) \\ &= \Sigma (\text{residues of } e^{st} f(s) \text{ at poles of } f(s)) \dots (6) \end{aligned}$$

This gives the use of *residue theorem* in finding Inverse Laplace Transform.




Fig. 10.8

Note. Referred to Fig. 10.8, if arc BDEFA = Γ and

$$T = \sqrt{R^2 - r^2}, \text{ then}$$

$$F(t) = \lim_{R \rightarrow \infty} \frac{1}{2\pi i} \int_{\gamma+iT}^{\gamma+i\infty} e^{st} f(s) ds$$

$$\text{or } F(t) = \lim_{R \rightarrow \infty} \left\{ \frac{1}{2\pi i} \oint_C e^{st} f(s) ds - \frac{1}{2\pi i} \int_{\Gamma} e^{st} f(s) ds \right\}$$

where C is the contour of Fig. 10.8

and known as *Bromwich Contour*.

Problem. 50. Evaluate $L^{-1}\left\{\frac{1}{(s+1)(s-2)^2}\right\}$ by residue-method.

We have $L^{-1} \left\{ \frac{1}{(s+1)(s-2)^2} \right\} = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{e^{st}}{(s+1)(s-2)^2} ds$
 $= \frac{1}{2\pi i} \oint_C \frac{e^{st}}{(s+1)(s-2)^2} ds = \text{sum of residues of } \frac{e^{st}}{(s+1)(s-2)^2}$
 at poles $s = -1$ (simple pole) and $s = 2$ (pole of order 2).

But residue of $\frac{e^{st}}{(s+1)(s-2)^2}$ at $s = -1$ is $\lim_{s \rightarrow -1} (s+1)f(s)$ where

$$f(s) = \frac{e^{st}}{(s+1)(s-2)^2}; \quad = \lim_{s \rightarrow -1} \frac{e^{st}}{(s-2)^2} = \frac{1}{9} e^{-t}$$

and residue of $\frac{e^{st}}{(s+1)(s-2)^2}$ at $s = 2$ (order two)

$$= \lim_{s \rightarrow 2} \frac{1}{1!} \frac{d}{ds} \left[(s-2)^2 \cdot \frac{e^{st}}{(s+1)(s-2)^2} \right] = \lim_{s \rightarrow 2} \frac{d}{ds} \left[\frac{e^{st}}{(s+1)} \right]$$

$$= \lim_{s \rightarrow 2} \left\{ \frac{(s+1)t - 1}{(s+1)^2} \right\} e^{st} = \frac{1}{3} te^{2t} - \frac{1}{9} e^{2t}$$

$$\text{Hence } L^{-1} \left\{ \frac{1}{(s+1)(s-2)^2} \right\} = \frac{1}{9} e^{-t} + \frac{1}{3} te^{2t} - \frac{1}{9} e^{2t}.$$

Problem 51. Solve the following differential equation $\frac{d^2 X}{dt^2} + \lambda^2 X = 0$.

$$\text{Given equation is } \frac{d^2 X}{dt^2} + \lambda^2 X = 0 \quad \dots (1)$$

Assuming $L\{X(t)\} = x(s)$, $X(0) = c_1$ and $X'(0) = c_2$, we have

$$L \left\{ \frac{d^2 X}{dt^2} \right\} = s^2 L\{X\} - sX(0) - X'(0) = s^2 x - sc_1 - c_2.$$

$\therefore$ taking Laplace transform of (1), we find

$$L \left\{ \frac{d^2 X}{dt^2} \right\} + \lambda^2 L\{X\} = 0 \text{ i.e. } s^2 x - sc_1 - c_2 + \lambda^2 x = 0 \text{ or } (s^2 + \lambda^2)x = sc_1 + c_2$$

$$\text{or } x = \frac{sc_1 + c_2}{s^2 + \lambda^2} = \frac{sc_1 + c_2}{(s+i\lambda)(s-i\lambda)} = \frac{c_1}{s+i\lambda} + \frac{ic_2}{s-i\lambda}$$

$$= \frac{A_1}{s+i\lambda} + \frac{A_2}{s-i\lambda} \text{ where } A_1 = \frac{c_1}{2} + \frac{ic_2}{\lambda} \text{ and } A_2 = \frac{c_1}{2} - \frac{ic_2}{\lambda}$$

Applying Laplace inversion theorem, we get

$$X = \frac{A_2}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{e^{st}}{s+i\lambda} ds + \frac{A_1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{e^{st}}{s-i\lambda} ds = A_1 I_1 + A_2 I_2 \text{ (say)}$$

$$\text{Here } I_1 = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{e^{st}}{s+i\lambda} ds. \text{ Put } s+i\lambda = S \text{ i.e. } ds = dS$$

$$= \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{e^{(S-i\lambda)t}}{S} dS = e^{-\lambda t} \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{e^{St}}{S} dS$$

$$= e^{-\lambda t} (\text{Residue of } \frac{e^{St}}{S} \text{ at } S=0) = e^{-\lambda t} \lim_{S \rightarrow 0} S \cdot \frac{e^{St}}{S} = e^{-\lambda t}$$

Similarly $I_2 = e^{i\lambda t}$.

Hence $X = A_1 e^{-i\lambda t} + A_2 e^{i\lambda t}$ which is the required solution.

Problem 52. An infinitely long circular cylinder of unit radius has a constant initial temperature U_0 . At $t=0$, a temperature of 0°C is applied to the surface and is maintained. Find the temperature at any point of the cylinder at any time t .

Take any point P of the cylinder. Taking z -axis along the axis of the cylinder and x, y axes as shown in Fig. 10.9, let (r, ϕ, z) be the cylindrical coordinates of P . Now the axis of z being coincident with the axis of the cylinder, the temperature is independent of ϕ and z and as such it may be taken as $U(r, t)$.

The temperature at any point of the cylinder is given by

$$\frac{\partial U}{\partial t} = k \left(\frac{\partial^2 U}{\partial r^2} + \frac{1}{r} \frac{\partial U}{\partial r} \right), \quad 0 < r < 1 \quad \dots (1)$$

with the boundary conditions (i) $U(1, t) = 0$, (ii) $U(r, 0) = U_0$ and (iii) the temperature is finite (bounded) i.e. $|U(r, t)| < M$.




Fig. 10.9

$$\text{The equation (1) is equivalent to } \frac{\partial U}{\partial t} = \frac{\partial^2 U}{\partial r^2} + \frac{1}{r} \frac{\partial U}{\partial r} \quad \dots (2)$$

where t is to be replaced by kt since $\frac{1}{k} \frac{\partial U}{\partial t} = \frac{\partial U}{\partial (kt)}$

Assuming that $L\{U(r, t)\} = u(r, s)$, we have

$$L \left\{ \frac{\partial U}{\partial r} \right\} = \frac{d}{dr} L\{U\} = \frac{du}{dr}, \quad L \left\{ \frac{\partial^2 U}{\partial r^2} \right\} = \frac{d^2 u}{dr^2} L\{U\} = \frac{d^2 u}{dr^2}$$

$$\text{and } L \left\{ \frac{\partial U}{\partial t} \right\} = s L\{U\} - U(r, 0) = su - U_0$$

$\therefore$ taking Laplace transform of (2), we get

$$su(r, s) - U_0 = \frac{d^2 u}{dr^2} + \frac{1}{r} \frac{du}{dr} \text{ i.e. } \frac{d^2 u}{dr^2} + \frac{1}{r} \frac{du}{dr} - su = -U_0 \quad \dots (3)$$

with the boundary condition, $u(1, s) = 0$ and $u(r, s)$ is bounded.

In order to find the general solution of (3), let us take two steps.

First step. The complimentary function of (3) is given by

$$\frac{d^2 u}{dr^2} + \frac{1}{r} \frac{du}{dr} - su = 0 \text{ or } \frac{d^2 u}{dr^2} + \frac{1}{r} \frac{du}{dr} + (s^2)u = 0 \because i^2 = -1$$

or $\frac{d^2 u}{dr^2} + \frac{1}{r} \frac{du}{dr} + (i\sqrt{s})^2 u = 0$, which is modified Bessel's equation of order zero and may also be expressed as

$$\frac{1}{(i\sqrt{s})^2} \frac{d^2u}{dr^2} + \frac{1}{(i\sqrt{s})^2} \cdot \frac{1}{r} \frac{du}{dr} + u = 0 \text{ or } \frac{d^2u}{d(i\sqrt{sr})^2} + \frac{1}{i\sqrt{sr}} \frac{du}{d(i\sqrt{sr})} + u = 0,$$

$$\text{or } \frac{d^2u}{dp^2} + \frac{1}{p} \frac{du}{dp} + u = 0 \text{ when } i\sqrt{sr} = p.$$

Being Bessel equation of zero order, its solution is

$$u = a_0 J_0(p) \text{ i.e. } u(r, s) = a_0 J_0(i\sqrt{sr}), a_0 \text{ being arbitrary constant} \dots (4)$$

and this is the complementary function of (3).

Second step. The particular integral of (3) is given by

$$\frac{d^2u}{dr^2} + \frac{1}{r} \frac{du}{dr} - su = -U_0.$$

As discussed in the Chapter of differential equations, its reduced equation on changing the independent variable r to z by the substitution,

$$z = \int e^{-\int \frac{1}{r} ds} dr = \int \frac{1}{\sqrt{r}} dr = 2\sqrt{r}, \text{ becomes}$$

$$\frac{d^2u}{dz^2} + Q_1 u = X_1 \text{ where } Q_1 = \frac{Q}{\left(\frac{dz}{dx}\right)^2} = \frac{-s}{\frac{1}{r}} = -rs \text{ and } X_1 = \frac{X}{\left(\frac{dz}{dx}\right)^2} = -U_0 r$$

$$\text{or } \frac{d^2u}{dz^2} - sru = -U_0 r \text{ or } (D^2 - sr)u = -U_0 s \text{ where } D \equiv \frac{d}{dz}$$

$$\therefore \text{Particular integral} = \frac{-U_0 r}{D^2 - rs} = \frac{U_0}{s} \left\{ 1 - \frac{D^2}{rs} \right\}^{-1} = \frac{U_0}{s} \left\{ 1 + \frac{D^2}{rs} + \dots \right\} \cdot 1 = \frac{U_0}{s}$$

Hence the complete integral of (3) is

$$u(r, s) = a_0 J_0(i\sqrt{sr}) + \frac{U_0}{s}.$$

$$\text{Initially when } r=1, u(r, s) = u(1, s) = 0 \text{ gives } a_0 = \frac{-U_0}{s J_0(i\sqrt{s})}$$

$$\text{Thus } u(r, s) = \frac{U_0}{s} - \frac{U_0 J_0(i\sqrt{sr})}{s J_0(i\sqrt{s})}.$$

Now using complex inversion formula, we find

$$U(r, t) = U_0 - \frac{U_0}{2\pi i} \int_{\gamma-i\infty}^{+\infty} \frac{e^{st} J_0(i\sqrt{sr})}{s J_0(i\sqrt{s})} ds \quad \because L^{-1}\left\{\frac{1}{s}\right\} = 1 \dots (5)$$

$$\text{Here } \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{e^{st} J_0(i\sqrt{sr})}{s J_0(i\sqrt{s})} ds = \text{sum of residues of } \frac{e^{st} J_0(i\sqrt{sr})}{s J_0(i\sqrt{s})}$$

at the poles given by $J_0(i\sqrt{s}) = 0$

Now $J_0(i\sqrt{s})$ has simple poles at the points where

$$i\sqrt{s} = \lambda_1, \lambda_2, \lambda_3, \dots, \lambda_n, \dots \text{ and at } s=0$$

$$= \lambda_n \text{ where } n=1, 2, 3, \dots, n, \dots \text{ and at } s=0$$

$$\text{i.e. } s = -\lambda_n^2 \text{ where } n=1, 2, 3, \dots, n \text{ and at } s=0$$

$$\text{Residue of } \frac{e^{st} J_0(i\sqrt{sr})}{s J_0(i\sqrt{s})} \text{ at } s=0 \text{ is } = \lim_{s \rightarrow 0} (s-o) \cdot \frac{e^{st} J_0(i\sqrt{sr})}{s J_0(i\sqrt{s})} = \frac{e^0 J_0(o)}{J_0(o)} = 1$$

$$\text{and Residue of } \frac{e^{st} J_0(i\sqrt{sr})}{s J_0(i\sqrt{s})} \text{ at } s = -\lambda_n^2 \text{ is}$$

$$= \lim_{s \rightarrow -\lambda_n^2} (s + \lambda_n^2) \cdot \frac{e^{st} J_0(i\sqrt{sr})}{s J_0(i\sqrt{s})}$$

$$= \lim_{s \rightarrow -\lambda_n^2} \frac{e^{st} J_0(i\sqrt{sr})}{s} \lim_{s \rightarrow -\lambda_n^2} \frac{s + \lambda_n^2}{J_0(i\sqrt{s})} = e^{-\lambda_n^2 t} \frac{J_0(\lambda_n r)}{-\lambda_n^2} \lim_{s \rightarrow -\lambda_n^2} \frac{s + \lambda_n^2}{J_0(i\sqrt{s})} \left(\text{form } \frac{0}{0} \right)$$

$$= \frac{e^{-\lambda_n^2 t} J_0(\lambda_n r)}{-\lambda_n^2} \lim_{s \rightarrow -\lambda_n^2} \frac{1}{J_0'(i\sqrt{s}) (i/2\sqrt{s})} \quad (\text{differentiating num and denom. w.r.t.s})$$

$$= \frac{e^{-\lambda_n^2 t} J_0(\lambda_n r)}{-\lambda_n^2} \cdot \frac{-2\lambda_n}{J_0'(\lambda_n)} \quad \because \frac{i}{2\sqrt{s}} = \frac{i^2}{2i\sqrt{s}} = \frac{-1}{2\lambda_n}$$

$$= \frac{2e^{-\lambda_n^2 t} J_0(\lambda_n r)}{\lambda_n J_1(\lambda_n)} \quad \because J_0'(x) = -J_1(x)$$

Hence (5) yields,

$$U(r, t) = U_0 - U_0 \left\{ 1 - \sum_{n=1}^{\infty} \frac{2e^{-\lambda_n^2 t} J_0(\lambda_n r)}{\lambda_n J_1(\lambda_n)} \right\} = 2U_0 \sum_{n=1}^{\infty} \frac{e^{-\lambda_n^2 t} J_0(\lambda_n r)}{\lambda_n J_1(\lambda_n)}$$

Replacing t by kt , the required solution is

$$U(r, t) = 2U_0 \sum_{n=1}^{\infty} \frac{e^{-kt\lambda_n^2} J_0(\lambda_n r)}{\lambda_n J_1(\lambda_n)}$$

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 53. Solve $\frac{d^2x}{dt^2} + \omega^2 x = \cos \omega t$, $t > 0$, subject to the initial conditions

$$x = x_0 \text{ and } \frac{dx}{dt} = x_1 \text{ when } t = 0.$$

$$\text{Ans. } x = x_0 \cos \omega t + x_1 \frac{\sin \omega t}{\omega} + \frac{t}{2\omega} \sin \omega t.$$

Problem 54. Solve $3 \frac{dx_1}{dt} + 2x_1 + \frac{dx_2}{dt} = 1$, $t \geq 0$ and $\frac{dx_1}{dt} + 4 \frac{dx_2}{dt} + 3x_2 = 0$, subject to the initial conditions $x_1 = 0 = x_2$ when $t = 0$.

Ans. $x_1 = \frac{1}{2} - \frac{1}{5}e^{-t} - \frac{3}{10}e^{-6t/11}$ and $x_2 = \frac{1}{5}(e^{-t} - e^{-6t/11})$.

Problem 55. A constant electromotive force E is applied at $t=0$ to an electrical circuit consisting of an inductance L , resistance R and capacitance C in series. The initial values of the current i and the charge (q) on the capacitor are 0. Find the current.

Hint: The current is given by $L \frac{di}{dt} + Ri + \frac{q}{c} = E$ where $\frac{dq}{dt} = i$.

Ans. $i = \frac{E}{\omega L} e^{-at} \sin \omega t$ if $\omega^2 > 0$ } where $a = \frac{R}{2L}$ and $\omega^2 = \frac{1}{LC} - \frac{R^2}{4L^2}$
 $= \frac{E}{L} t e^{-at}$ if $\omega^2 = 0$
 $= \frac{E}{kL} e^{-at} \sinh kt$ if $\omega^2 < 0$ where $k^2 = -\omega^2$

Problem 56. Solve Bessel's equation $\frac{d^2 z}{du^2} + \frac{1}{u} \frac{dz}{du} + \left(1 - \frac{n^2}{u^2}\right) z = 0$.

Ans. $z = J_n(u)$.

Problem 57. Show that the solution of $\frac{d^2 y}{dt^2} + a^2 y = F(t)$ subject to the initial conditions $y=0=y'$ when $t=0$, is $y = \frac{1}{a} \int_0^t F(u) \sin a(t-u) du$.

Problem 58. A particle of mass m can perform small oscillations about a position of equilibrium under a restoring force mn^2 times the displacement. It is started from rest by a constant force F_0 which acts for a time T and then ceases. Show that the amplitude of the subsequent oscillation is $(2 F_0 / mn^2) \sin(nT/2)$.

Problem 59. If L_n be the Laguerre Polynomial given by $L_n(x) = \frac{e^x}{n!} \frac{d^n}{dx^n} (e^{-x} x^n)$ then show $L\{L_n(x)\} = \frac{(s-1)^n}{s^{n+1}}$.

Problem 60. Prove the following:

(i) $L\left\{\frac{2}{t} \sinh t\right\} = \log \frac{s+1}{s-1}$, (ii) $L\left\{\int_0^t J_0(\mu) J_0(t-\mu) d\mu\right\} = \frac{1}{s^2+1}$.

(iii) $L^{-1}\left\{\frac{f(s)}{\sinh(\mu s)}\right\} = 2 \sum_{k=0}^{\infty} F(t-2k, \mu-\mu) \psi(t-2k\mu-\mu)$.

Problem 61. (a) Find the Laplace transforms of the functions e^{-at} , $\cos(bt+a)$, (t^2+1)

(b) Solve by means of Laplace transforms

$z'' + 2z' + \int_0^t y dt = t$; $z'' + 2z' + y = \sin 2t$; $z(0) = -1$, $z'(0) = 1$ (Bombay, 1965)

Problem. 62. (a) Define Laplace transform of a function $f(t)$, obtain the Laplace transforms for (i) t^n , (ii) e^{-at} , (iii) $\sin \omega t$, (iv) te^{-at} , (v) $\int_0^t f(t) dt$.

(b) Solve the following differential equation: $\frac{d^2 x}{dt^2} + 4 \frac{dx}{dt} + 13x = 0$

using Laplace transforms, given that $x=3$ and $\frac{dx}{dt}=6$ when $t=0$.

(Bombay, 1970)

CHAPTER 11

HANKEL TRANSFORMS

11.1. INTRODUCTION

In §7.5 [B], we have mentioned the form of Laplace's equation as

$$\nabla^2 V \equiv \frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial z^2} = 0 \quad \dots (1)$$

This can be transformed to cylindrical coordinates (u, ϕ, z) by using the transformation $x = u \cos \phi$, $y = u \sin \phi$, $z = z$ so that $u^2 = x^2 + y^2$ and $\tan \phi = \frac{y}{x}$

Giving $\frac{\partial u}{\partial y} = \frac{y}{u} = \sin \phi$, $\frac{\partial \phi}{\partial x} = -\frac{\sin \phi}{u}$ and $\frac{\partial \phi}{\partial y} = \frac{\cos \phi}{u}$

$$\therefore \frac{\partial V}{\partial x} = \frac{\partial V}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial V}{\partial \phi} \frac{\partial \phi}{\partial x} + \frac{\partial V}{\partial z} \frac{\partial z}{\partial x} = \frac{\partial V}{\partial u} \cos \phi + \frac{\partial V}{\partial \phi} \left(-\frac{\sin \phi}{u}\right)$$

i.e. $\frac{\partial}{\partial x} \equiv \cos \phi \frac{\partial}{\partial u} - \frac{\sin \phi}{u} \frac{\partial}{\partial \phi}$

and $\frac{\partial^2 V}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial V}{\partial x} \right) = \left(\cos \phi \frac{\partial}{\partial u} - \frac{\sin \phi}{u} \frac{\partial}{\partial \phi} \right) \left(\cos \phi \frac{\partial V}{\partial u} - \frac{\sin \phi}{u} \frac{\partial V}{\partial \phi} \right)$

$$= \cos^2 \phi \frac{\partial^2 V}{\partial u^2} - \frac{2 \sin \phi \cos \phi}{u} \frac{\partial^2 V}{\partial u \partial \phi} + \frac{2 \sin \phi \cos \phi}{u^2} \frac{\partial V}{\partial \phi} + \frac{\sin^2 \phi}{u} \frac{\partial V}{\partial u} + \frac{\sin^2 \phi}{u^2} \frac{\partial^2 V}{\partial \phi^2}$$

similarly $\frac{\partial^2 V}{\partial y^2} = \sin^2 \phi \frac{\partial^2 V}{\partial u^2} + \frac{2 \sin \phi \cos \phi}{u} \frac{\partial^2 V}{\partial u \partial \phi} - \frac{2 \sin \phi \cos \phi}{u^2} \frac{\partial V}{\partial \phi} + \frac{\cos^2 \phi}{u} \frac{\partial V}{\partial u} + \frac{\cos^2 \phi}{u^2} \frac{\partial^2 V}{\partial \phi^2}$

so that $\nabla^2 V = \frac{\partial^2 V}{\partial u^2} + \frac{1}{u} \frac{\partial V}{\partial u} + \frac{1}{u^2} \frac{\partial^2 V}{\partial \phi^2} + \frac{\partial^2 V}{\partial z^2} = 0 \quad \dots (2)$

is the cylindrical form of (1).

In order to solve it by the method of separation of variables discussed in §7.5, we put $V = U\Phi Z$... (3)

where $U = U(u)$, $\Phi = \Phi(\phi)$, $Z = Z(z)$

we have,

$$\frac{\partial V}{\partial u} = \Phi Z \frac{\partial U}{\partial u}; \frac{\partial^2 V}{\partial u^2} = \Phi Z \frac{\partial^2 U}{\partial u^2}, \frac{\partial^2 V}{\partial \phi^2} = UZ \frac{\partial^2 \Phi}{\partial \phi^2} \text{ and } \frac{\partial^2 V}{\partial z^2} = U\Phi \frac{\partial^2 Z}{\partial z^2}$$

Their substitution in (2) yields,

$$\Phi Z \frac{\partial^2 U}{\partial u^2} + \frac{1}{u} \frac{\partial U}{\partial u} \Phi Z + \frac{1}{u^2} UZ \frac{\partial^2 \Phi}{\partial \phi^2} + U\Phi \frac{\partial^2 Z}{\partial z^2} = 0 \quad \dots (4)$$

Dividing throughout by $U\Phi Z$, we find

$$\frac{1}{U} \left(\frac{\partial^2 U}{\partial u^2} + \frac{1}{u} \frac{\partial U}{\partial u} \right) + \frac{1}{u^2} \frac{1}{\Phi} \frac{\partial^2 \Phi}{\partial \phi^2} + \frac{1}{Z} \frac{\partial^2 Z}{\partial z^2} = 0 \quad \dots (5)$$

The first three terms being independent of z , so must also be the last

$$\text{i.e. } \frac{\partial^2 Z}{\partial z^2} = CZ, \text{ } C \text{ being a constant.} \quad \dots (6)$$

Again the first two and the last terms in (5) being independent of ϕ ,

so must also be the third one and hence $\frac{\partial^2 \Phi}{\partial \phi^2} = D\Phi$, D being a constant

$\dots (7)$

As such (5) becomes

$$u^2 \frac{\partial^2 U}{\partial u^2} + u \frac{\partial U}{\partial u} + (D + Cu^2)U = 0 \quad \dots (8)$$

If we put $C = k^2$, $D = -n^2$, (6) and (7) become $\frac{\partial^2 Z}{\partial z^2} = k^2 Z$ and $\frac{\partial^2 \Phi}{\partial \phi^2}$

$= -n^2 \Phi$ whose solutions are $Z = A_1 e^{kz} + B_1 e^{-kz}$ and $\Phi = A_2 \cos n\phi + B_2 \sin n\phi$ respectively.

and (8) reduces to $u^2 \frac{\partial^2 U}{\partial u^2} + u \frac{\partial U}{\partial u} + (k^2 u^2 - n^2) U = 0$

With $v = ku$, this becomes, $v^2 \frac{\partial^2 U}{\partial v^2} + v \frac{\partial U}{\partial v} + (v^2 - n^2) U = 0$

$\dots (9)$

which is Bessel's differential equation of order n and its solutions are known as *cylindrical functions* or Bessel's functions of order n .

Hence if $J_n(v)$ be a Bessel function of order n , then the solution of Laplace's equation is $V = U\Phi Z = J_n(ku) e^{\pm in\phi} e^{\pm kz}$

$\dots (10)$

which is known as a *Cylindrical Harmonic* and this harmonic is symmetrical about z -axis if $n=0$.

In §8.4 we have already discussed the Bessel's equation in the form

$$\frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} + \left(1 - \frac{n^2}{x^2}\right)y = 0 \quad \dots (11)$$

which is the same form as (9).

Its solution is $y = AJ_n(x) + BJ_{-n}(x)$

$\dots (12)$

A, B being arbitrary constants; and $J_n(x)$ is known as *Bessel's function of first kind of order n* given by

$$J_n(x) = \sum_{r=0}^{\infty} \frac{(-1)^r \left(\frac{x}{2}\right)^{n+2r}}{r! \Gamma(n+r-1)} \quad \dots (13)$$

$$\text{and } J_{-n}(x) = \sum_{r=0}^{\infty} \frac{(-1)^r \left(\frac{x}{2}\right)^{-n+2r}}{r! \Gamma(-n+r-1)} \quad \dots (14)$$

such that $J_n(x) = (-1)^n J_{-n}(x)$

The series on the right of (12) is absolutely convergent for all values of x . In case n is an integer, the second independent solution of (11) may be shown to be

$$G_n(x) = \frac{\pi}{2} \operatorname{cosec} [J_{-n}(x) - e^{-in\pi} J_n(x)] \quad \dots (15)$$

where $G_n(x)$ is known as *Bessel function of the second kind of order n* .

In this case the general solution of (11) is

$$y = AJ_n(x) + BG_n(x) \quad \dots (16)$$

Also, we have the recurrence relations for Bessel's functions of the first kind

$$xJ_n' = nJ_n - xJ_{n+1} = xJ_{n-1} - nJ_n \quad \dots (17)$$

$$2J_n' = J_{n-1} - J_{n+1} \quad \dots (18)$$

$$2nJ_n = x\{J_{n-1} + J_{n+1}\} \quad \dots (19)$$

$$\frac{d}{dx} (x^{-n} J_n) = -x^{-n} J_{n+1} \quad \dots (20)$$

$$\frac{d}{dx} (x^n J_n) = x^n J_{n-1} \quad \dots (21)$$

Here (21) provides the definite integral

$$\int_0^\mu x^n J_{n-1}(x) dx = \left[x^n J_n(x) \right]_0^\mu = \mu^n J_n(\mu), \quad n > 0 \quad \dots (22)$$

If we put $n=1$ and make the substitutions $x = \xi r$, $\mu = \xi a$ in the integral on the left of (22), we find

$$\int_0^a r J_0(\xi r) dr = \frac{a}{\xi} J_1(a\xi) \quad \dots (23)$$

which is frequently used in the application of the theory of *Hankel transforms* to special functions.

Also we may write

$$\int_0^a r^3 J_0(\xi r) dr = \int_0^a r^2 \frac{1}{\xi} \frac{\partial}{\partial r} [r J_1(\xi r)] dr = \left[\frac{r^3}{\xi} J_1(\xi r) \right]_0^a - \frac{2}{\xi} \int_0^a r^2 J_1(\xi r) dr$$

Now putting on the right of (22), $n=2$, $x = \xi r$, $\mu = \xi a$ and using (19)

$$\text{with } n=1, \text{ we may find } \int_0^a r^3 J_0(\xi r) dr = \frac{a^2}{\xi^2} \left[2J_0(\xi a) + \left(a\xi - \frac{4}{a\xi}\right) J_1(\xi a) \right] \quad \dots (24)$$

Subtracting (24) from (23) multiplied by a^2 , we find

$$\int_0^a r(a^2 - r^2)J_0(\xi r)dr = \frac{4a}{\xi^3}J_1(\xi a) - \frac{2a^2}{\xi^2}J_0(\xi a) \quad \dots (25)$$

A few more integrals involving Bessel functions may be referred as

$$\int_0^\infty e^{-ax}J_0(\xi x)dx = (a^2 + \xi^2)^{-1/2} \quad \dots (26)$$

$$\int_0^\infty e^{-ax}J_1(\xi x)dx = \frac{1}{\xi} - \frac{a}{\xi\sqrt{(a^2 + \xi^2)}} \quad \dots (27)$$

$$\int_0^\infty x e^{-ax}J_0(\xi x)dx = a(a^2 + \xi^2)^{-3/2} \quad \dots (28)$$

$$\int_0^\infty x e^{-ax}J_1(\xi x)dx = \xi(a^2 + \xi^2)^{-3/2} \quad \dots (29)$$

$$\int_0^\infty e^{-ax}J_1(\xi x) \frac{dx}{x} = \frac{(a^2 + \xi^2)^{1/2} - a}{\xi} \quad \dots (30)$$

Further if n is half an odd integer we have

$$J_n(x) = \frac{x^n}{2^n \Gamma(n+1)} \left[1 - \frac{x^2}{1 \cdot 2^2(n+1)} + \frac{x^4}{2 \cdot 2^2(n+1)(n+2)} \right] \quad \dots (31)$$

which gives for $n = \frac{1}{2}$ and $n = -\frac{1}{2}$

$$J_{1/2}(x) = \sqrt{\frac{2}{x\pi}} \sin x \text{ and } J_{-1/2}(x) = \sqrt{\frac{2}{\pi x}} \cos x \quad \dots (32)$$

11.2. DEFINITION OF INFINITE HANKEL TRANSFORM

If $J_n(px)$ be the Bessel function of the first kind of order n , then the Hankel transform of a function $f(x)$, ($0 < x < \infty$) denoted by $\tilde{f}(p)$ is defined as

$$\tilde{f}(p) = \int_0^\infty f(x) \cdot x J_n(px) dx \quad \dots (1)$$

Here $xJ_n(px)$ is the *Kernel* of the transformation as referred in § 10.2.

We sometimes write (1) as

$$H\{f(x)\} = \tilde{f}(p) = \int_0^\infty f(x) \cdot x J_n(px) dx \quad \dots (2)$$

Problem 1. Taking $xJ_0(px)$ as the Kernel of the transformation, find the Hankel transform of the following functions:

$$(i) f(x) = \frac{e^{-ax}}{x}, (ii) f(x) = e^{-ax}, (iii) f(x) = \begin{cases} 1, & 0 < x < a, n=0 \\ 0, & x > a, n=0 \end{cases}$$

$$(iv) f(x) = \begin{cases} a^2 - x^2, & 0 < x < a, n=0 \\ 0, & x > a, n=0 \end{cases}$$

$$(i) \text{ we have } H\{f(x)\} = \tilde{f}(p) = \int_0^\infty \frac{e^{-ax}}{x} \cdot x J_0(px) dx =$$

$$\int_0^\infty e^{-ax} J_0(px) dx = (a^2 + p^2)^{-1/2} \text{ by (26) of §11.1.}$$

$$(ii) \tilde{f}(p) = \int_0^\infty e^{-ax} x J_0(px) dx = \frac{a}{(a^2 + p^2)^{3/2}} \text{ by (28) of §11.1.}$$

$$\text{Aliter. } \tilde{f}(p) = \int_0^\infty e^{-ax} x J_0(px) dx = \left[-\frac{1}{a} e^{-ax} x J_0(px) \right]_0^\infty + \frac{1}{a} \int_0^\infty e^{-ax} \frac{d}{dx} \{x J_0(px)\} dx \text{ (or integrating by parts)}$$

$$= 0 + \frac{1}{a} \int_0^\infty e^{-ax} \{J_0(px) + x J_0'(px)\} dx$$

$$= \frac{1}{a} \int_0^\infty e^{-ax} J_0(px) dx + \frac{1}{a} \int_0^\infty e^{-ax} x J_0'(px) dx$$

But we have $xJ_n'(x) = nJ_n(x) - xJ_{n+1}(x)$ by (17) of §11.1 writing $x=0$ and replacing x by px , we get $J_0'(xp) = -J_1(xp)$,

$$\therefore \tilde{f}(p) = \frac{1}{a} \int_0^\infty e^{-ax} J_0(px) dx - \frac{1}{a} \int_0^\infty e^{-ax} x J_1(px) dx$$

$$= \frac{1}{a(a^2 + p^2)^{1/2}} - \frac{1}{a} \cdot \frac{p^2}{(a^2 + p^2)^{3/2}} \text{ by (26) and (27) of §11.1}$$

$$(iii) H\{f(x)\} = \tilde{f}(p) = \int_0^\infty f(x) \cdot x J_0(px) dx = \int_0^a 1 \cdot x J_0(px) dx + \int_a^\infty 0 \cdot x J_0(px) dx = \frac{a}{p} J_1(ap) \text{ by (23) of §11.1}$$

$$(iv) H\{f(x)\} = \tilde{f}(p) = \int_0^\infty f(x) x \cdot J_0(px) dx = \int_0^a (a^2 - x^2) x J_0(px) dx + \int_a^\infty 0 \cdot x J_0(px) dx$$

$$= \int \frac{4a}{p^2} J_1(pa) - \frac{2a^2}{p^2} J_0(pa) \text{ by (25) of §11.1.}$$

Note. On the next page we have tabulated some useful Hankel transforms.

11.3. INVERSE FORMULA FOR HANKEL TRANSFORM

If $\tilde{f}(p)$ be the Hankel transform of the function $f(r)$ for $-\infty < x < \infty$,

$$\text{i.e., } \tilde{f}(p) = \int_{-\infty}^\infty f(r) \cdot r J_n(pr) dr \quad \dots (1)$$

$$\text{Then } f(r) = \int_0^\infty \tilde{f}(p) \cdot p J_n(pr) dp \quad \dots (2)$$

is said to be the inversion formula for the Hankel transform $\tilde{f}(p)$ and we may write $f(r) = H^{-1}\{\tilde{f}(p)\}$ $\dots (3)$

$f(x)$	n	$\tilde{f}(p)$	$f(x)$	n	$\tilde{f}(p)$
$x^n, 0 < x < a$ $1, x > a$	> -1	$\frac{a^{n+1}}{p} J_{n+1}(pa)$	$\frac{e^{-ax}}{x}$	1	$\frac{1}{p} - \frac{p}{p(p^2+a^2)^{1/2}}$
$1, 0 < x < a$ $0, x > a$	0	$\frac{a}{p} J_1(ap)$	e^{-ax}	1	$p(p^2+a^2)^{-3/2}$
$a^2-x^2, 0 < x < a$ $0, x > a$	0	$\frac{4a}{p^2} J_1(pa) - \frac{2a^2}{p^2} J_0(pa)$	$\frac{\sin ax}{a}$	0	$\begin{cases} 0, p > a \\ (a^2-p^2)^{-1/2}, 0 < p < a \end{cases}$
x^{m-1}	> -1	$\frac{2m\Gamma(\frac{1}{2} + \frac{1}{2}(m+\frac{1}{2}n))}{p^{m+1}\Gamma(\frac{1}{2} - \frac{1}{2}m + \frac{1}{2}n)}$	$\frac{\sin ax}{a}$	1	$\begin{cases} \frac{a}{(p^2-a^2)^{1/2}}, p > a \\ 0, p < a \end{cases}$
$x^{-2}e^{-ax}$	1	$\frac{\sqrt{p^2+a^2}-a}{p}$	$\frac{\sin x^2}{x^2}$	0	$\begin{cases} \sin^{-1} \frac{1}{p}, p > 1 \\ \frac{\pi}{2}, p < 1 \end{cases}$
e^{-ax}	0	$a(p^2+a^2)^{-3/2}$	$\frac{a}{(a^2+x^2)^{3/2}}$	0	e^{-ap}
$x^n e^{-ax^2}$	> -1	$\frac{p}{(2a)^{n+1}} e^{-p^2/4a}$	$x^{m-2}e^{-ax^2}$	> -1	$\frac{p^n \Gamma(\frac{n}{2} + \frac{m}{2})}{2^{n+1} p^{m/2} \Gamma(\frac{1}{2}) \Gamma(1+n)} \times$
$\frac{e^{-ax}}{x}$	0	$(p^2+a^2)^{-1/2}$			$F\left\{\frac{n}{2} + \frac{m}{2}, n+1, -\frac{p^2}{4a}\right\}$

in terms of confluent hypergeometric function.

In chapter 9, we have introduced that if $\tilde{f}(s)$ be the complex Fourier transform of $f(x)$ in $-\infty < x < \infty$, i.e., $\tilde{f}(s) = \int_{-\infty}^{\infty} f(x) e^{isx} dx$

Then we have $f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \tilde{f}(s) e^{-isx} ds$.

In §9.10 [F] dealing with multiple Fourier transform, we have considered the functions of two variables such as

$$\tilde{f}(s, t) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) e^{i(sx+ty)} dx dy \quad \dots (4)$$

$$\text{then } f(x, y) = \frac{1}{4\pi^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \tilde{f}(s, t) e^{-i(sm+tn)} ds dt \quad \dots (5)$$

$$\text{If we put } x=r \cos \theta, y=r \sin \theta, s=p \cos \alpha, t=p \sin \alpha, \text{ then (4) and (5) transform to } \tilde{f}(p, \alpha) = \int_0^{\infty} r dr \int_0^{2\pi} f(r, \theta) e^{ipr \cos(\theta-\alpha)} d\theta \quad \dots (6)$$

$$\text{and } f(r, \theta) = \frac{1}{4\pi^2} \int_0^{\infty} p dp \int_0^{2\pi} \tilde{f}(p, \alpha) e^{-ipr \cos(\theta-\alpha)} d\alpha \quad \dots (7)$$

Taking $f(r, \theta) = f(r) e^{-in\theta}$, (6) reduces to

$$\tilde{f}(p, \alpha) = \int_0^{\infty} f(r) r dr \int_0^{2\pi} e^{i\{-n\theta + pr \cos(\theta-\alpha)\}} d\theta$$

If we now make a substitution, $\phi = \alpha - \theta - \frac{\pi}{2}$, then it becomes

$$\begin{aligned} \tilde{f}(p, \alpha) &= \int_0^{\infty} f(r) r dr \int_0^{2\pi} e^{i\{n(\phi+\pi/2-\alpha)+pr \cos(\phi+\pi/2)\}} d\phi \\ &= \int_0^{\infty} f(r) r dr e^{in(\pi/2-\alpha)} \int_0^{2\pi} e^{i(n\phi-pr \sin \phi)} d\phi \\ &= \int_0^{\infty} f(r) r dr \cdot 2\pi e^{in(\pi/2-\alpha)} J_n(pr) dr \\ &\quad \therefore J_n(pr) = \frac{1}{2\pi} \int_0^{2\pi} e^{i(n\phi-pr \sin \phi)} d\phi \\ &= 2\pi e^{in(\pi/2-\alpha)} \int_0^{\infty} f(r) \cdot r J_n(pr) dr \\ &= 2\pi e^{in(\pi/2-\alpha)} \tilde{f}(b) \quad \dots (8) \end{aligned}$$

Again taking $f(r, \theta) = f(r) e^{-in\theta}$, and using (8), (7) yields

$$\begin{aligned} f(r) e^{-in\theta} &= \frac{1}{2\pi} \int_0^{\infty} p dp \int_0^{2\pi} 2\pi e^{in(\pi/2-\alpha)} \tilde{f}(p) \\ &\quad e^{-ipr \cos(\theta-\alpha)} d\alpha \end{aligned}$$

$$= \frac{1}{2\pi} \int_0^\infty p \tilde{f}(p) dp \int_0^{2\pi} e^{i\{x(\pi/2-\alpha)-pr \cos(\theta-\alpha)\}} d\alpha$$

Putting $\psi = \theta - \alpha + \frac{\pi}{2}$, this becomes

$$\begin{aligned} f(r) e^{-in\theta} &= \int_0^\infty p \tilde{f}(p) dp \int_0^{2\pi} e^{i\{n(\psi-\theta)-pr \cos(\pi/2-\psi)\}} d\psi \\ &= \frac{1}{2\pi} \int_0^\infty p \tilde{f}(p) dp e^{-in\theta} \int_0^{2\pi} e^{i(n\psi - pr \sin \psi)} d\psi \\ \text{i.e. } f(r) &= \frac{1}{2\pi} \int_0^\infty p \tilde{f}(p) dp \cdot 2\pi J_n(pr) dp \\ &= \int_0^\infty \tilde{f}(p) p J_n(pr) dp \end{aligned}$$

which is the required inversion formula.

Problem 2. Find $H^{-1}\{p^2 e^{-ap}\}$ when $n=1$.

$$\begin{aligned} \text{We have } H^{-1}\{p^2 e^{-ap}\} &= \int_0^\infty p^2 e^{-ap} \cdot p J_1(px) dp \\ &= \int_0^\infty e^{-ap} J_1(px) \frac{dp}{p} = \frac{(a^2+x^2)^{1/2}-a}{x} \end{aligned}$$

11.4. PARSEVAL'S THEOREM FOR HANKEL TRANSFORMS

If $\tilde{f}(p)$ and $\tilde{g}(p)$ be respectively the Hankel transform of $f(x)$ and $g(x)$

$$\text{then } \int_0^\infty x f(x) g(x) dx = \int_0^\infty p \tilde{f}(p) \tilde{g}(p) dp. \quad \dots (1)$$

$$\text{We have } \int_0^\infty p \tilde{f}(p) \tilde{g}(p) dp = \int_0^\infty p \tilde{f}(p) dp \int_0^\infty x g(x) J_n(px) dx$$

$$(\text{on substituting } \tilde{g}(p) = \int_0^\infty g(x) x J_n(px) dx, n \geq -\frac{1}{2})$$

$$= \int_0^\infty x g(x) dx \int_0^\infty p \tilde{f}(p) J_n(px) dx$$

(changing the order of integration)

$$= \int_0^\infty x g(x) f(x) dx \text{ by §11.3.}$$

11.5. LINEARITY PROPERTY

If $f(x)$ and $g(x)$ are two functions and C_1, C_2 two constants, then

$$H\{C_1 f(x) + C_2 g(x)\} = C_1 H\{f(x)\} + C_2 H\{g(x)\} \quad \dots (1)$$

$$\text{We have } H\{C_1 f(x) + C_2 g(x)\} = \int_0^\infty \{C_1 f(x) + C_2 g(x)\} \cdot x J_n(px) dx$$

$$= C_1 \int_0^\infty f(x) \cdot x J_n(px) dx + C_2 \int_0^\infty g(x) \cdot x J_n(px) dx$$

$$= C_1 H\{f(x)\} + C_2 H\{g(x)\}.$$

This result can be extended to any number of functions.

11.6. HANKEL TRANSFORM OF THE DERIVATIVES OF A FUNCTION

If $\tilde{f}_n(p)$ be the Hankel transform of order n of the function $f(x)$ i.e.

$$\tilde{f}_n(p) = \int_0^\infty x f(x) J_n(px) dx, \text{ then the Hankel transform of } \frac{df}{dx} \text{ is}$$

$$\tilde{f}_n'(p) = \int_0^\infty x \frac{df}{dx} J_n(px) dx \quad \dots (1)$$

$$= \left[x f(x) J_n(px) \right]_0^\infty - \int_0^\infty f(x) \frac{d}{dx} \{x J_n(px)\} dx \quad \dots (2)$$

(on integrating by parts)

$$= - \int_0^\infty f(x) \{J_n(px) + x J_n'(px)\} dx \quad \dots (3)$$

under the assumption that $x f(x) \rightarrow 0$ as $x \rightarrow 0$ or $x \rightarrow \infty$

$$= - \int_0^\infty f(x) \{(1-n) J_n(px)\} dx - \int_0^\infty px f(x) J_{n-1}(px) dx$$

($\because x J_n(px) = px J_n(px) - n J_n(px)$ by (17) of §11.1)

$$= (n-1) \int_0^\infty f(x) J_n(px) dx - p \tilde{f}_{n-1}(p)$$

$$= (n-1) I - p \tilde{f}_{n-1}(p) \text{ (say). } \dots \text{ where } I \equiv \int_0^\infty f(x) J_n(px) dx$$

The recurrence relation,

$$J_{n-1}(x) - \frac{2n}{x} J_n(x) + J_{n+1}(x) = 0 \text{ gives}$$

$$J_{n-1}(px) - \frac{2n}{px} J_n(px) + J_{n+1}(px) = 0 \quad \text{on replacing } x \text{ by } px$$

$$\text{i.e. } 2n J_n(x) = px J_{n-1}(x) + px J_{n+1}(x)$$

$$\text{so that } 2n I = 2n \int_0^\infty f(x) J_n(px) dx = p \left[\int_0^\infty x f(x) J_{n-1}(px) dx + \int_0^\infty x f(x) J_{n+1}(px) dx \right]$$

$$= p \tilde{f}_{n-1}(p) + p \tilde{f}_{n+1}(p) \quad \dots (5)$$

Hence (5) reduces to

$$\begin{aligned} \tilde{f}_n(p) &= \frac{n-1}{2n} p \tilde{f}_{n-1}(p) + \frac{n-1}{2n} p \tilde{f}_{n+1}(p) - p \tilde{f}_{n-1}(p) \\ &= -p \left[\frac{n+1}{2n} \tilde{f}_{n-1}(p) - \frac{n-1}{2n} \tilde{f}_{n+1}(p) \right] \quad \dots (6) \end{aligned}$$

which gives the required Hankel transform of $\frac{df}{dx}$

Replacing n by $(n-1)$ and $(n+1)$ in succession (6) yields,

$$\begin{aligned}\tilde{f}'_{n-1}(p) &= -p \left[\frac{n}{2(n-1)} \tilde{f}_{n-2}(p) - \frac{n-2}{2(n-1)} \tilde{f}_n(p) \right] \\ \text{and } \tilde{f}'_{n+1}(p) &= -p \left[\frac{n+2}{2(n+1)} \tilde{f}_n(p) - \frac{n}{2(n+1)} \tilde{f}_{n+2}(p) \right]\end{aligned}$$

Using these results and replacing f by f' in (6) we find

$$\begin{aligned}\tilde{f}''_n(p) &= -p \left[\frac{n+1}{2n} \tilde{f}'_{n-1}(p) - \frac{n-1}{2n} \tilde{f}'_{n+2}(p) \right] \\ &= \frac{p^2}{4} \left[\frac{n+1}{n-1} \tilde{f}_{n-2}(p) - 2 \frac{n^2-3}{n^2-1} \tilde{f}_n(p) + \frac{n-1}{n+1} \tilde{f}_{n+2}(p) \right] \\ &\dots (7)\end{aligned}$$

COROLLARY. Putting $n=1, 2, 3$ successively in (6) we have

$$\tilde{f}'_1(p) = -p \tilde{f}_0(p) \dots (8)$$

$$\tilde{f}'_2(p) = -p \left[\frac{3}{4} \tilde{f}_1(p) - \frac{1}{4} \tilde{f}_3(p) \right] \dots (9)$$

$$\tilde{f}'_3(p) = -p \left[\frac{2}{3} \tilde{f}_2(p) - \frac{1}{3} \tilde{f}_4(p) \right] \dots (10)$$

11.7. HANKEL TRANSFORMS OF $\frac{d^2f}{dx^2}$, $\frac{df}{dx} + \frac{1}{x} \frac{df}{dx}$ and

$\frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} f$ UNDER CERTAIN CONDITIONS

$$\begin{aligned}\text{We have } H \left\{ \frac{d^2f}{dx^2} \right\} &= \int_0^\infty \frac{d^2f}{dx^2} \cdot x J_n(px) dx = \left[\frac{df}{dx} x J_n(px) \right]_0^\infty \\ &\quad - \int_0^\infty \frac{df}{dx} \cdot \frac{d}{dx} \{ x J_n(px) \} dx\end{aligned}$$

Assuming that $xf(x) \rightarrow 0$ as $x \rightarrow 0$, $x \rightarrow \infty$, we have

$$\begin{aligned}H \left\{ \frac{d^2f}{dx^2} \right\} &= - \int_0^\infty \frac{df}{dx} \left[J_n(px) + p x J'_n(px) \right] dx \\ &= \int_0^\infty x \frac{d^2f}{dx^2} \cdot J_n(px) dx \\ \text{or } \int_0^\infty x \left(\frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right) J_n(px) dx &= -p \int_0^\infty \frac{df}{dx} \cdot x J'_n(px) dx \\ &= -p \left[f(x) \cdot x J'_n(px) \right]_0^\infty - \int_0^\infty f(x) \frac{d}{dx} \{ x J'_n(px) \} dx \\ &= p \int_0^\infty f(x) \frac{d}{dx} \{ x J'_n(px) \} dx \\ &\dots (1)\end{aligned}$$

$\therefore xf(x) \rightarrow 0$ as $x \rightarrow 0$ or ∞

But $J_n(px)$ satisfies Bessel's differential equation i.e.

$$\frac{d}{dx} \left(x \frac{dy}{dx} \right) + \left(1 - \frac{n^2}{x^2} \right) xy = 0,$$

$$\therefore \frac{d}{dx} \left(x \frac{d}{dx} J_n(x) \right) + \left(1 - \frac{n^2}{x^2} \right) x J_n(x) = 0$$

$$\text{or } \frac{1}{p} \frac{d}{dx} [px \cdot J'_n(px)] = - \left(p^2 - \frac{n^2}{x^2} \right) \frac{x}{p^2} J_n(px),$$

on replacing x by px

$$\text{or } \frac{d}{dx} [x J'_n(px)] = - \left(p^2 - \frac{n^2}{x^2} \right) \frac{x}{p} J_n(px)$$

As such (1) reduces to

$$\begin{aligned}\int_0^\infty \left(\frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right) \cdot x J_n(px) dx \\ = -p \int_0^\infty f(x) \left(p^2 - \frac{n^2}{x^2} \right) \frac{x}{p} J_n(px) dx\end{aligned}$$

$$\begin{aligned}\text{or } \int_0^\infty \left(\frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} f \right) \cdot x J_n(px) dx \\ = -p^2 \int_0^\infty f(x) \cdot x J_n(px) dx \\ = -p^2 \tilde{f}_n(p) \quad \text{by § 11.6} \dots (2)\end{aligned}$$

where $\tilde{f}_n(p)$ is the Hankel transform of order n of $f(x)$.

COROLLARY 1. If we put $n=0$ in (2), we get

$$\int_0^\infty x \left(\frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right) J_0(px) dx = -p^2 \tilde{f}_0(p) \dots (3)$$

where $\tilde{f}_0(p)$ is the Hankel Transform of Zeroeth order.

COROLLARY 2. If we put $n=1$ in (2), we find

$$\int_0^\infty \left(\frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{1}{x^2} f \right) x J_1(px) dx = -p^2 \tilde{f}_1(p) \dots (4)$$

$$\text{or } \int_0^\infty x \frac{df}{dx} J_1(px) dx = -p \tilde{f}(p) \dots (5)$$

where $\tilde{f}(p) = \int_0^\infty x f(x) J_0(px) dx$.

Problem 3. Find $H \left\{ \frac{\partial f}{\partial x} \right\}$ when $f = \frac{e^{-ax}}{x}$ and $n=1$.

We have

$$H \left\{ \frac{df}{dx} \right\} = \int_0^\infty x \frac{df}{dx} J_1(px) dx = -p \tilde{f}_0(p) \text{ by (5) of § 11.6.}$$

$$= -p \int_0^\infty x f(x) J_0(px) dx = -p \int_0^\infty e^{-ax} J_0(px) dx = \frac{-ap}{(a^2 + p^2)^{3/2}}$$

Problem 4. Find $H\left\{\frac{d^2 f}{dt^2}\right\}$ when $f=f(x, t)$.

$$\text{We have } H\left\{\frac{d^2 f}{dt^2}\right\} = \int_0^\infty x \frac{\partial^2 f}{\partial t^2} J_n(px) dn = \frac{\partial^2}{\partial t^2} \int_0^\infty x f(x, t) J_n(px) dx \\ = \frac{d^2}{dt^2} \tilde{f}(p, t).$$

Problem 5. Evaluate $\int_0^\infty \left(\frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx}\right) \cdot x J_0(px) dx$, when $f(x) = e^{-ax}$.

We have by (3) of §11.7,

$$\int_0^\infty x \left(\frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx}\right) J_0(px) dx = -p^2 \tilde{f}_0(p) \\ = -p^2 \int_0^\infty e^{-ax} J_0(px) dx = \frac{-ap^2}{(a^2 + p^2)^{3/2}}.$$

11.8. APPLICATIONS TO BOUNDARY VALUE PROBLEMS

Whenever there is symmetry about an axis in a problem, then the use of polar coordinates is advised. Using Hankel transform a variable ranging from 0 to ∞ can be excluded.

Problem 6. Find the potential $V(r, z)$ of a field due to a flat circular disc of unit radius with centre at origin and axis along z -axis, satisfying the differential equation $\frac{\partial^2 V}{\partial r^2} + \frac{1}{r} \frac{\partial V}{\partial r} + \frac{\partial^2 V}{\partial z^2} = 0$, $0 \leq r \leq \infty$, $z \geq 0$ and the boundary conditions:

$$(i) V = V_0 \text{ when } z=0, 0 \leq r < 1$$

$$(ii) \frac{\partial V}{\partial z} = 0 \text{ when } z=0, r > 1.$$

Taking the Hankel transform of the given equations for $n=0$, we have

$$\int_0^\infty \left(\frac{\partial^2 V}{\partial r^2} + \frac{1}{r} \frac{\partial V}{\partial r}\right) r J_0(pr) dr = - \int_0^\infty \frac{\partial^2 V}{\partial z^2} r J_0(pr) dr$$

$$\text{or } -p^2 \tilde{V} = - \frac{d^2 \tilde{V}}{dz^2} \text{ where } \tilde{V} = \int_0^\infty V \cdot r J_0(pr) dr \quad \dots (1)$$

$$\text{i.e., } \frac{d^2 \tilde{V}}{dz^2} - p^2 \tilde{V} = 0 \text{ whose solution is } \tilde{V}(p, z) = A e^{+pz} + B e^{-pz} \quad \dots (2)$$

Now since $V \rightarrow 0$ as $z \rightarrow \infty$ and so $\tilde{V} \rightarrow 0$ as $z \rightarrow \infty$, therefore $A=0$.

As such (2) reduces to $\tilde{V}(p, z) = B e^{-pz}$.

But B being independent of z may be taken as $B(p)$ i.e. function of p alone.

$$\therefore \tilde{V}(p, z) = B(p) e^{-pz}$$

Applying inversion formula,

$$V(r, z) = \int_0^\infty B(p) e^{-pz} \cdot p J_0(pr) dp \quad \dots (3)$$

Applying the conditions when $z=0$,

$$V = \int_0^\infty p \cdot B(p) J_0(pr) dp = V_0, 0 \leq r < 1 \quad \dots (4)$$

$$\text{and } \left(\frac{\partial V}{\partial z}\right)_{z=0} = \int_0^\infty -p^2 B(p) J_0(pr) dp = 0, r > 1$$

$$\text{i.e. } \int_0^\infty p^2 B(p) J_0(pr) dp = 0, r > 1 \quad \dots (5)$$

Comparing (4) and (5) with the integrals

$$\int_0^\infty J_0(pr) \frac{\sin p}{p} dp = \frac{\pi}{2}, 0 \leq r < 1 \text{ and}$$

$$\int_0^\infty J_0(pr) \sin p dp = 0, r > 0 \text{ respectively, then we find}$$

$$B(p) = \frac{2}{\pi} V_0 \frac{\sin p}{p^2}.$$

Hence the required solution is

$$V(r, z) = \frac{2V_0}{\pi} \int_0^\infty e^{-pz} \frac{\sin p}{p} J_0(pr) dp.$$

Problem 7. The magnetic potential V for a circular disc of radius a and strength ω , magnetised parallel to its axis, satisfying Laplace's equation is equal to $2\pi\omega$ on the disc itself and vanishes at exterior points in the plane of the disc. Show that at the point (r, z) , $z > 0$, $V = 2\pi\omega \int_0^\infty e^{-pz} J_0(pr) J_1(pa) dp$.

The magnetic potential V satisfies the Laplace's equation

$$\frac{\partial^2 V}{\partial r^2} + \frac{1}{r} \frac{\partial V}{\partial r} + \frac{\partial^2 V}{\partial z^2} = 0, 0 < r < \infty \quad \dots$$

With the boundary condition

$$V = 2\pi\omega, 0 \leq r < a, z=0 \text{ and } V=0, r > a, z=0.$$

Hankel transform of (1) for $n=0$ gives

$$\int_0^\infty \left(\frac{\partial^2 V}{\partial r^2} + \frac{1}{r} \frac{\partial V}{\partial r}\right) r J_0(pr) dr + \int_0^\infty \frac{\partial^2 V}{\partial z^2} \cdot r J_0(pr) dr = 0.$$

$$\text{or } -p^2 \tilde{V} + \frac{d^2 \tilde{V}}{dz^2} = 0 \text{ where } \tilde{V} = \int_0^\infty V \cdot r J_0(pr) dr \quad \dots (2)$$

$$\begin{aligned} \text{and } \left[\tilde{V} \right]_{z=0} &= \int_0^a \left[V \right]_{z=0} \cdot r J_0(pr) dr + \int_0^\infty \left[V \right]_{z=0} r J_0(pr) dr \\ &= \int_0^a 2\pi\omega \cdot r J_0(pr) dr + 0 = 2\pi\omega \int_0^a \frac{1}{p} \frac{d}{dr} \{r J_1(pr)\} dr \\ &= \frac{2\pi\omega a}{p} J_1(pa) \end{aligned} \quad \dots (3)$$

Solution of (2) is $\tilde{V} = Ae^{pz} + Be^{-pz}$ $\dots (4)$

But $\tilde{V} \rightarrow 0$ as $z \rightarrow \infty$ $\therefore A=0$ and so (4) reduces to $\tilde{V} = Be^{-pz}$ $\dots (5)$

From (3) and (5), we have $\left[\tilde{V} \right]_{z=0} = B = \frac{2\pi\omega a}{p} J_1(pa)$

Hence $\tilde{V} = \frac{2\pi\omega a}{p} J_1(pa) e^{-pz}$

Applying inversion formula we find

$$V(r, z) = 2\pi\omega a \int_0^\infty e^{-pz} J_0(pr) J_1(pr) dp.$$

11.9. FINITE HANKEL TRANSFORMS

If $f(r)$ be a function satisfying Dirichlet's conditions (given in §9.2) in the interval $(0, a)$, then its finite Hankel transform is defined as

$$\tilde{f}(p_i) = \int_0^a r f(r) J_n(p_i r) dr \quad \dots (1)$$

where p_i is a positive root of the transcendental equation $J_n(p_i a) = 0$ $\dots (2)$

Also at any point in $(0, a)$ where $f(x)$ is continuous, we have

$$f(r) = \frac{2}{a^2} \sum_i \tilde{f}(p_i) \frac{J_n(p_i r)}{\{J'_n(p_i a)\}^2} \quad \dots (3)$$

Note 1. For the sake of convenience the upper limit a is generally converted to unity (1) by suitable transformation.

Note 2. All the roots of $J_n(\lambda_i) = 0$ are real and unrepeatd (except $\lambda_i = 0$) It is more convenient to take the sum over all the positive roots λ_i . As such when $n=0$, (3) can be easily transformed by substitution $p_i = \frac{\lambda_i}{a}$. In case $a=1$, the equation is automatically reduced to $J_0(p_i) = 0$.

Note 3. When $n=0$ and $a=1$, then $J_0'(x) = -J_1(x)$ $\dots (4)$
so that the inversion formula (3) becomes

$$f(r) = 2 \sum_i \tilde{f}(p_i) \frac{J_0(p_i r)}{\{J_1(p_i)\}^2} \quad \dots (5)$$

where p_i are roots of $J_0(p_i) = 0$ $\dots (6)$

Note 4. If $f(r)$ is represented by generalized Fourier—Bessel series

$$f(r) = \sum_i C_i J_n(p_i r), \quad 0 \leq r \leq a \quad \dots (7)$$

then the coefficient C_i is given by

$$\begin{aligned} C_i &= \frac{2}{a^2 J_{n+1}^2(p_i a)} \int_0^a r f(r) J_n(p_i r) dr \\ &= \frac{2 \tilde{f}(p_i)}{a^2 \left[J_{n+1}(p_i a) \right]^2} = \frac{2 \tilde{f}(p_i)}{a^2 \left[J'_n(p_i a) \right]^2} \text{ using (4)} \end{aligned} \quad \dots (8)$$

$$\text{Hence } f(r) = \frac{2}{a^2} \sum_i \tilde{f}(p_i) \frac{J_n(p_i r)}{\left[J'_n(p_i a) \right]^2} \quad \dots (9)$$

In case $a=1$, this becomes

$$f(r) = 2 \sum_i \tilde{f}(p_i) \frac{J_n(p_i r)}{J_n^2(p_i)} \quad \dots (10)$$

Summations being taken over $J_n(p_i) = 0$.

Note 5. $f(r)$ is the inverse Hankel transform of $\tilde{f}(p_i)$.

Note 6. $r J_n(p_i r)$ is the Kernel of the transform.

Another forms of Hankel Transform

[1] When, the range of variation does not include the origin i.e. $f(r)$ satisfies Dirichlet's condition in $0 < b \leq r \leq a$, then the finite Hankel transform is defined as

$$\tilde{f}(p_i) = \int_b^a r f(r) \left[J_n(p_i r) Y_n(p_i a) - Y_n(p_i r) J_n(p_i a) \right] dr \quad \dots (11)$$

where Y_n is the Bessel function of order n of second kind and p_i is a root of the equation

$$J_n(p_i a) Y_n(p_i b) - J_n(p_i b) Y_n(p_i a) = 0 \quad \dots (12)$$

Also then at each point of (a, b) where $f(r)$ is continuous we have the inversion formula

$$f(r) = \sum_i \frac{2 p_i^2 J_n^2(p_i a)}{J_n^2(p_i b) - J_n^2(p_i a)} \tilde{f}(p_i) \left[J_n(p_i r) Y_n(p_i b) - J_n(p_i b) Y_n(p_i r) \right] \quad \dots (13)$$

where the summation extends over all the positive roots of (12).

[2] If $f(r)$ satisfies Dirichlet's conditions in the closed interval $[0, 1]$ i.e. $0 \leq r \leq 1$, and its finite Hankel transform is defined as

$$\tilde{f}(p_i) = \int_0^1 r f(r) J_n(p_i r) dr \quad \dots (14)$$

in which p_i is a root of the transcendental equation

$$p J'_n(p) + h J_n(p) = 0 \quad \dots (15)$$

then at each point of $[0, 1]$ where $f(r)$ is continuous, we have the inversion formula

$$f(r) = 2 \sum_p \frac{p^2 \tilde{f}(p)}{h^2 + p^2 - n^2} \cdot \frac{J_n(pr)}{J_n^2(p)} \quad \dots (16)$$

where the summation extends over all the positive roots of (15).

If $n=0$, (16) reduces to

$$\tilde{f}(r) = 2 \sum_p \frac{p^2 \tilde{f}(p)}{h^2 + p^2} \cdot \frac{J_0(pr)}{J_0^2(p)} \quad \dots (17)$$

and (15) reduces to $h J_0(p) = p J_1(p)$ $\therefore J_0'(p) = -J_1(p)$ $\dots (18)$

Also using (15), (16) may be written as

$$f(r) = 2 \sum_p \frac{\tilde{f}(p)}{1 + \frac{p^2 - n^2}{h^2}} \cdot \frac{J_n(pr)}{h^2 J_n^2(p)} \quad \dots (19)$$

$$\text{or} \quad f(r) = 2 \sum_p \frac{\tilde{f}(p)}{1 + \frac{p^2 - n^2}{h^2}} \cdot \frac{J_n(pr)}{J_n'(p)} \quad \dots (20)$$

where the summation extends over all the roots of the equation

$$J_n(p) = -\frac{p J_n'(p)}{h} \quad \dots (21)$$

Evidently if $h \rightarrow \infty$, then (20) and (21) reduce to (3) and (2) where $a=1$ and hence we may conclude that (3) and (2) are limiting case of (20) and (21) respectively.

Problem 8. Find $H_n \{r^n\}$ taking $r J_n(pr)$ as the Kernel of transform.

$$\therefore \frac{d}{dx} \left[x^n J_n(x) \right] = x^n J_{n-1}(x)$$

$$\therefore \int_0^b \lambda^n J_{n-1}(x) dx = \left[x^n J_n(x) \right]_0^b = b^n J_n(b), \quad n > 0$$

$$\therefore x^n J_n(x) \rightarrow 0 \text{ as } x \rightarrow 0$$

$$\text{Replacing } n \text{ by } (n+1), \int_0^b \lambda^{n+1} J_n(x) dx = b^{n+1} J_{n+1}(b)$$

$$\text{Writing } pr \text{ for } x \text{ and } pa \text{ for } b; \int_0^a r^n \cdot r J_n(pr) dr = \frac{a^{n+1}}{p} J_{n+1}(pa)$$

$$\therefore 1 + n \left\{ r^n \right\} = \frac{a^{n+1}}{p} J_{n+1}(ap) = \frac{1}{p} J_{n+1}(p) \text{ for } a=1.$$

Problem 9. Find the Hankel transform of $\frac{J_n(ax)}{J_n(a)}$, taking $x J_n(px)$ as the Kernel of the transform.

$J_n(ax)$ and $J_n(px)$ being the solutions of Bessel's equation, we have

$$x^2 \frac{d^2}{dx^2} J_n(ax) + x \frac{d}{dx} J_n(ax) + (\alpha^2 x^2 - n^2) J_n(ax) = 0 \quad \dots (1)$$

$$x^2 \frac{d^2}{dx^2} J_n(px) + x \frac{d}{dx} J_n(px) + (p^2 x^2 - n^2) J_n(px) = 0 \quad \dots (2)$$

Multiplying (1) by $J_n(px)$, (2) by $J_n(ax)$ and subtracting, we get

$$(\alpha^2 - p^2) x J_n(ax) J_n(px) = \frac{d}{dx} \left[x \left\{ J_n(ax) \frac{d}{dx} J_n(px) - J_n(px) \frac{d}{dx} J_n(ax) \right\} \right]$$

Integrating with regard to x from 0 to 1 and using $J_n(p)=0$, we find

$$(\alpha^2 - p^2) \int_0^1 x J_n(ax) J_n(px) dx = p J_n(\alpha) J_n'(p)$$

$$\therefore \int_0^1 \frac{J_n(ax)}{J_n(a)} x J_n(px) dx = \frac{p}{\alpha^2 - p^2} J_n'(p)$$

$$\text{i.e. } H_n \left\{ \frac{J_n(ax)}{J_n(a)} \right\} = \frac{p}{\alpha^2 - p^2} J_n'(p)$$

Note. For $n=0$, using $J_1(p) = -J_0'(p)$, this yields

$$H_0 \left\{ \frac{J_0(ax)}{J_0(a)} \right\} = -\frac{p}{\alpha^2 - p^2} J_1(p).$$

Here below we tabulate a few of finite Hankel transforms.

$f(x) = \frac{2}{a^2} \sum_i \tilde{f}_i(p_i) \frac{J_n(p_i x)}{ J'_n(p_i a) }$	n	$\tilde{f}_i(p_i) = \int_0^a f(x) J_n(px) dx$
x^n	> -1	$\frac{a^{n+1}}{p_i} J_{n+1}(p_i a)$
c	0	$\frac{ac}{p_i} J_1(pa)$
$a^2 - x^2$	0	$\frac{4a}{p_i^2} J_1(p_i a)$
$\frac{J_r(xa)}{J_r(ia)}$	> -1	$\frac{p_i \alpha}{a^2 - p_i^2} J_n'(p_i a)$
$\frac{J_0(ax)}{J_0(a)}$	0	$\frac{p_i \alpha}{a^2 - p_i^2} J_1(p_i a)$

Problem 10. Find $f(x)$ if $H\{f(x)\} = \frac{c}{p} J_1(pa)$, p being positive root of $J_0(pa)$.

$$\text{Inversion formula gives } f(x) = \frac{2}{a^2} \sum_i \tilde{f}(p) \frac{J_n(px)}{[J_n'(pa)]^2}$$

where summation extends over all positive roots of $J_n(pa)$.

$$\begin{aligned} \therefore f(x) &= \frac{2}{a} \sum_i \frac{c}{p} \frac{J_1(pa) J_0(px)}{\{J_0'(pa)\}^2} = \frac{2}{a} \sum_i \frac{c}{p} \cdot \frac{J_1(pa) J_0(px)}{\{J_1(pa)\}^2} \\ &\because J_0'(x) = -J_1(x) \\ &= \frac{2}{a^2} \sum_i \frac{c}{p} \frac{J_0(px)}{J_1(pa)} \end{aligned}$$

11.10. FINITE HANKEL TRANSFORM OF $\frac{df}{dx}$, p being root of $J_n(p) = 0$.

$$\begin{aligned} H_n \left\{ \frac{df}{dx} \right\} &= \int_0^1 \frac{df}{dx} \cdot x J_n(px) dx = \left[f(x) \cdot x J_n(px) \right]_0^1 \\ &\quad - \int_0^1 f(x) \frac{d}{dx} \{x J_n(px)\} dx \\ &= - \int_0^1 f(x) \frac{d}{dx} \{x J_n(px)\} dx \quad \dots (1) \\ &\because J_n(p) = 0 \text{ for } x=1 \text{ and } x J_n(px) = 0 \text{ for } x=0. \end{aligned}$$

Now we have the recurrence relations

$$2 J_n'(x) = J_{n-1}(x) - J_{n+1}(x) \text{ and } 2n J_n(x) = x[J_{n-1}(x) + J_{n+1}(x)]$$

which on replacing x by px , become

$$J_n'(px) = \frac{1}{2} [J_{n-1}(px) - J_{n+1}(px)] \text{ and } J_n(px) = \frac{px}{2n} [J_{n-1}(px) + J_{n+1}(px)]$$

Substituting them in $\frac{d}{dx} \{x J_n(px)\} = xp J_n'(px) + J_n(px)$ we get

$$\begin{aligned} \frac{d}{dx} \{x J_n(px)\} &= \frac{px}{2} [J_{n-1}(px) - J_{n+1}(px)] + \frac{px}{2n} [J_{n-1}(px) + J_{n+1}(px)] \\ &= \frac{p}{2n} [(n+1)x J_{n-1}(px) - (n-1)x J_{n+1}(px)] \end{aligned}$$

So that (1) yields

$$\begin{aligned} H_n \left\{ \frac{df}{dx} \right\} &= \frac{p}{2n} \int_0^1 [(n-1)x f(x) J_{n+1} - (n+1)x f(x) J_{n-1}] dx \\ &= \frac{p}{2n} [(n-1) H_{n+1} \{f\} - (n+1) H_{n-1} \{f\}] \quad \dots (2) \end{aligned}$$

where $H_n \{f\}$ denotes the Hankel transform of order n .

$$\text{Note. For } n=1, (2) \text{ yields } H_1 \left\{ \frac{df}{dx} \right\} = -p H_0 \{f(x)\} \quad \dots (3)$$

11.11. FINITE HANKEL TRANSFORM OF $\frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx}$.

Case I. When p is a root of $J_n(p) = 0$.

$$\begin{aligned} \text{We have } H_n \left\{ \frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right\} &= \int_0^1 x \left\{ \frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right\} J_n(px) dx \\ &= \int_0^1 x \frac{d^2f}{dx^2} J_n(px) dx + \int_0^1 \frac{df}{dx} J_n(px) dx \\ &= H_n \left\{ \frac{d^2f}{dx^2} \right\} + \int_0^1 \frac{df}{dx} J_n(px) dx \quad \dots (1) \end{aligned}$$

In result (2) of § 11.11, replacing f by $\frac{df}{dx}$, we get

$$H_n \left\{ \frac{d^2f}{dx^2} \right\} = \frac{p}{2n} \left[+ (n-1) H_{n+1} \left\{ \frac{df}{dx} \right\} - (n+1) H_{n-1} \left\{ \frac{df}{dx} \right\} \right] \dots (2)$$

Now replacing x by px in the recurrence relation

$$2n J_n(x) = x [J_{n-1}(x) + J_{n+1}(x)]$$

$$\text{we have } J_n(px) = \frac{px}{2n} [J_{n-1}(x) + J_{n+1}(px)].$$

$$\begin{aligned} \text{So that } \int_0^1 \frac{df}{dx} J_n(px) dx &= \int_0^1 \frac{df}{dx} \cdot \frac{px}{2n} [J_{n-1}(x) + J_{n+1}(px)] dx \\ &= \frac{p}{2n} \int_0^1 x \frac{df}{dx} J_{n-1}(px) dx + \frac{p}{2n} \int_0^1 x \frac{df}{dx} J_{n+1}(px) dx \\ &= \frac{p}{2n} \left[H_{n-1} \left\{ \frac{df}{dx} \right\} + H_{n+1} \left\{ \frac{df}{dx} \right\} \right] \quad \dots (3) \end{aligned}$$

With the help of (2) and (3); (1) yields,

$$H_n \left\{ \frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right\} = \frac{p}{2} \left[-H_{n-1} \left\{ \frac{df}{dx} \right\} + H_{n+1} \left\{ \frac{df}{dx} \right\} \right] \dots (4)$$

Case II. When p is a root of $p J_n'(p) + h J_n(p) = 0$... (5)

Taking $n=0$, (5) becomes $p J_0'(p) + h J_0(p) = 0$... (6)

$$\begin{aligned} \text{we have } H_0 \left\{ \frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right\} &= \int_0^1 \left(\frac{d^2f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right) x J_0(px) dx \\ &= \int_0^1 \frac{d^2f}{dx^2} x J_0(px) dx + \int_0^1 \frac{df}{dx} \cdot J_0(px) dx \\ &= \left[\frac{df}{dx} \cdot x J_0(px) \right]_0^1 - \int_0^1 \frac{df}{dx} \cdot \frac{d}{dx} \{x J_0(px)\} dx + \int_0^1 \frac{df}{dx} \cdot J_0(px) dx \\ &= \left[\frac{df}{dx} \right]_{x=1} J_0(p) - \int_0^1 \frac{df}{dx} \{xp J_0'(px) + J_0(px)\} dx + \int_0^1 \frac{df}{dx} J_0(px) dx \\ &= \left[\frac{df}{dx} \right]_{x=1} J_0(p) - p \int_0^1 \frac{df}{dx} x J_0'(px) dx \end{aligned}$$

$$= \left[\frac{df}{dx} \right]_{x=1} J_0(p) - p \left[f(x) \cdot x J_0'(px) \right]_0^1 + p \int_0^1 f(x) \cdot \frac{d}{dx} \{x J_0'(px)\} dx$$

$$= \left[\frac{df}{dx} \right]_{x=1} J_0(p) - p f(1) J_0'(p) + p \int_0^1 f(x) \cdot \{x p J_0''(px) + J_0'(px)\} dx$$

But $J_0(x)$ being the solution of Bessel's equation

$$x^2 \frac{d^2 y}{dx^2} + \frac{dy}{dx} + xy = 0, \text{ we have}$$

$$x J_0''(x) + J_0'(x) + x J_0(x) = 0.$$

Replacing x by px , $px J_0''(px) + J_0'(px) = -px J_0(px)$

$$\begin{aligned} \therefore H_0 \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right\} &= \left[\frac{df}{dx} \right]_{x=1} J_0(p) - p f(1) \left\{ -\frac{h J_0(p)}{p} \right\} \\ &\quad - p^2 \int_0^1 x f(x) J_0(px) dx \text{ by (6)} \\ &= \left\{ \left[\frac{df}{dx} \right]_{x=1} + h f(1) \right\} J_0(p) - p^2 H_0 \{f(x)\} \\ &= \left[\frac{df}{dx} + h f(x) \right]_{x=1} J_0(p) - p^2 H_0 \{f(x)\} \end{aligned} \quad \dots (7)$$

If boundary condition be such that $\frac{df}{dx} + h f(x) = 0$ when $x=1$, then

$$\text{we get } H_0 \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right\} = -p^2 H_0 \{f(x)\} \quad \dots (8)$$

Case III. When p is a root of $J_n(pa)$ $Y_n(pb) - J_n(pb) Y_n(pa) = 0$ $\dots (9)$

Integrating by parts, the following, we have

$$\begin{aligned} &\int_a^b x [J_n(px) Y_n(bp) - J_n(bp) Y_n(xp)] \left(\frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right) dx \\ &= -p \left\{ x f(x) [J_n'(xp) Y_n(bp) - J_n(bp) Y_n'(xp)] + p \int_a^b x f(x) \frac{d}{dx} \right. \\ &\quad \left. [J_n'(xp) Y_n(bp) - J_n(bp) Y_n'(xp)] + f(x) [J_n'(xp) Y_n(bp) - J_n(bp) \right. \\ &\quad \left. Y_n'(xp)] \right\} dx \quad \dots (10) \end{aligned}$$

It is easy to show that

$$J_n'(pa) Y_n(bp) - Y_n'(ap) J_n(pb) = \frac{1}{pa} \frac{J_n(pb)}{J_n(pa)} \quad \dots (11)$$

With the help of (11), (10) yields,

$$H_n \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right\} = \int_0^1 \left(\frac{n^2}{x^2} - p^2 \right) x f(x) [J_n(px) Y_n(bp) - J_n(bp) Y_n(xp)] dx$$

$$\text{or } H_n \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} \right\} = \frac{J_n(pb)}{J_n(pa)} f(a) - f(b) - p^2 H_n \{f(x)\} \quad \dots (12)$$

Note 1. If boundary condition is such that $f(a)=0=f(b)$, then (12) yields

$$H_n \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} \right\} = -p^2 H_n \{f(x)\} \quad \dots (13)$$

Note 2. If $n=0$ then (13) reduces to

$$H_0 \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right\} = -p^2 H_0 \{f(x)\} \quad \dots (14)$$

which is the same as (8).

11.12. FINITE HANKEL TRANSFORM OF $\frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} f(x)$, p being a root of $J_n(pa) = 0$.

We have,

$$\begin{aligned} H_n \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} f(x) \right\} &= \int_0^a \left[\frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} f(x) \right] x J_n(px) dx \\ &= \int_0^a \frac{d^2 f}{dx^2} \cdot x J_n(px) dx + \int_0^a \frac{df}{dx} J_n(px) dx - n^2 \int_0^a \frac{1}{x} f(x) J_n(px) dx \\ &= \left[\frac{df}{dx} \cdot x J_n(px) \right]_0^a - \int_0^a \frac{df}{dx} \cdot \frac{d}{dx} \{x J_n(px)\} dx + \int_0^a \frac{df}{dx} J_n(px) dx - x^2 \int_0^a \frac{1}{x} f(x) J_n(px) dx \\ &= - \int_0^a \frac{df}{dx} \{px J_n'(px) + J_n(px)\} dx + \int_0^a \frac{df}{dx} J_n(px) dx - n^2 \int_0^a \frac{1}{x} f(x) J_n(px) dx \quad \because J_n(pa) = 0 \\ &= -p \int_0^a \frac{df}{dx} \cdot x J_n'(px) dx - n^2 \int_0^a \frac{1}{x} f(x) J_n(px) dx \\ &= -p \left[f(x) \cdot x J_n'(px) \right]_0^a + p \int_0^a f(x) \cdot \frac{d}{dx} \{x J_n'(px)\} dx - n^2 \int_0^a \frac{1}{x} f(x) J_n(px) dx \\ &= -pa f(a) J_n'(ap) + p \int_0^a f(x) \cdot \{px J_n''(px) + J_n'(px)\} dx - n^2 \int_0^a \frac{1}{x} f(x) J_n(px) dx \end{aligned}$$

But $J_n(x)$ being the solution of Bessel's equation,

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - n^2) y = 0,$$

we have $x^2 J_n''(x) + x J_n'(x) + (x^2 - n^2) J_n(x) = 0$

Replacing x by px , this gives

$$p^2 x^2 J_n''(px) + px J_n'(px) + (p^2 x^2 - n^2) J_n(px) = 0$$

$$\text{i.e., } px J_n''(px) + J_n'(px) = px \left(\frac{n^2}{p^2 x^2} - 1 \right) J_n(px)$$

$$\begin{aligned} \therefore H_n \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} f(x) \right\} &= -pa f(a) J_n'(ap) + p \int_0^a f(x) \\ &\quad \left\{ px \left(\frac{n^2}{p^2 x^2} - 1 \right) J_n(px) \right\} dx - n^2 \int_0^a \frac{1}{x} f(x) J_n(px) dx \\ &= -pa f(a) J_n'(ap) - p^2 \int_0^a f(x) x J_n(px) dx \\ &= -pa f(a) J_n'(ap) - p^2 H_n \{f(x)\} \end{aligned} \quad \dots (1)$$

Note 1. If $a=1$, (1) becomes

$$H_n \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} - \frac{n^2}{x^2} f(x) \right\} = -p f(1) J_n'(p) - p^2 H_n \{f(x)\} \quad \dots (2)$$

Note 2. When $n=0$, (2) reduces to

$$\begin{aligned} H_0 \left\{ \frac{d^2 f}{dx^2} + \frac{1}{x} \frac{df}{dx} \right\} &= -p f(1) J_0'(p) - p^2 H_0 \{f(x)\} \\ &= -p f(1) J_1(p) - p^2 H_0 \{f(x)\} \quad \therefore J_0'(p) = -J_1(p) \quad \dots (3) \end{aligned}$$

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 11. Find the Finite Hankel transform $\phi(p)$ where

$$\phi(p) = \frac{2^{l+n-m}}{\Gamma(m-n)} \rho^n (1-\rho^2)^{m-n-1}$$

$$\text{We have } H_n \{\phi(p)\} = \frac{2^{l+n-m}}{\Gamma(m-n)} \int_0^1 \rho^{n+1} (1-\rho^2)^{m-n-1} J_n(p\rho) d\rho$$

If we expand Bessel function $J_n(p\rho)$ in ascending powers of $p\rho$ and integrate term by term, we get

$$\begin{aligned} H_n \{\phi(p)\} &= \frac{2^{l+n-m}}{\Gamma(m-n)} \sum_{s=0}^{\infty} \int_0^1 \rho^{n+1} (1-\rho^2)^{m-n-1} \frac{(-1)^s (p\rho)^{n+2s}}{2^{n+2s} s! \Gamma(n+s+1)} d\rho \\ &= \frac{1}{\Gamma(m-n)} \sum_{s=0}^{\infty} \frac{p^{n+2s}}{2^{m+2s}} \int_0^1 t^{n+s} (1-t)^{m-n-1} \frac{(-1)^s dt}{s! \Gamma(n+s+1)}, \text{ where } \rho^2 = t \\ &= \sum_{s=0}^{\infty} \frac{(-1)^s p^{m+2s}}{2^{m+2s} \Gamma(m+s+1)} \frac{p^{n-m}}{s!} = p^{n-m} J_m(p). \end{aligned}$$

Problem 12. Find the finite Hankel transform of $\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial V}{\partial r} \right) - \frac{n^2 V}{r}$, where $V=0$ when $r=0$ and $V=V_1$ when $r=1$.

$$\text{We have } \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial V}{\partial r} \right) - \frac{n^2 V}{r} = \frac{\partial^2 V}{\partial r^2} + \frac{1}{r} \frac{\partial V}{\partial r} - \frac{n^2 V}{r^2} = f(r) \text{ (say)}$$

Hence using (2) of § 11.12, we have

$$H\{f(r)\} = -p f(1) J_n'(p) - p^2 H_n \{f(r)\} \text{ when } f(r)=V \text{ and } f(0)=0, f(1)=V_1$$

$$= -p V_1 J_n'(p) - p^2 H_n V$$

Problem 13. Solve $\frac{\partial^2 V}{\partial r^2} + \frac{1}{r} \frac{\partial V}{\partial r} = \frac{1}{k} \frac{\partial V}{\partial t}$, $0 \leq r < 1$, $t > 0$

where $\frac{\partial V}{\partial r} + hV=0$, when $r=1$, $t > 0$ and $V=1$ when $t=0$, $0 \leq r < 1$; h, k being constants

$$\text{Assuming that } \tilde{V} = \int_0^1 V r J_0(pr) dr \quad \dots (1)$$

$$\text{where } p \text{ is a root of } p J_0'(p) + h J_0(p) = 0 \quad \dots (2)$$

and multiplying the given equation by $r J_0(pr)$ and then integrating with regard to r from 0 to 1, we have

$$\int_0^1 \left(\frac{\partial^2 V}{\partial r^2} + \frac{1}{r} \frac{\partial V}{\partial r} \right) r J_0(pr) dr = \frac{1}{k} \int_0^1 \frac{\partial V}{\partial t} r J_0(pr) dr = \frac{1}{k} \frac{d\tilde{V}}{dt}$$

$$\frac{1}{k} \frac{d\tilde{V}}{dt} = \left\{ \frac{\partial V}{\partial r} r J_0(pr) \right\}_0^1 - \int_0^1 \frac{\partial V}{\partial r} \{J_0(pr) + r p J_0'(pr)\} r p + \int_0^1 \frac{\partial V}{\partial r} J_0'(pr) dr$$

$$= \left[\frac{\partial V}{\partial r} r J_0(pr) \right]_0^1 - p \int_0^1 r \frac{\partial V}{\partial r} J_0'(pr) dr$$

$$= J_0(p) \left[\frac{\partial V}{\partial r} \right]_{r=1} - p \left[V J_0'(pr) \right]_0^1 - \int_0^1 V \{J_0'(pr) + p r J_0''(pr)\} dr$$

$$= J_0(p) \left[\frac{\partial V}{\partial r} \right]_{r=1} - p [V]_{r=1} J_0'(p) - p^2 \int_0^1 V r J_0(pr) dr \quad \therefore J_0(pr) \text{ satisfies}$$

Bessel's equation.

$$= J_0(p) \left[\frac{\partial V}{\partial r} + hV \right]_{r=1} - p^2 \tilde{V} \quad \therefore p J_0'(p) = -h J_0(p)$$

$$= -p^2 \tilde{V}$$

$$\text{i.e., } \frac{d\tilde{V}}{dt} = -k p^2 \tilde{V} \text{ where } \tilde{V} = \int_0^1 r J_0(pr) dr = \frac{J_1(p)}{p} \text{ when } t=0, \therefore V=1 \text{ when } t=0, 0 \leq r < 1$$

$$\therefore \text{Its solution is } \tilde{V} = \frac{J_1(p)}{p} e^{-k p^2 t}$$

Applying inversion formula, we find

$$V = 2 \sum_p e^{-k p^2 t} \frac{p J_1(p) J_0(pr)}{h^2 + p^2 J_0^2(p)}$$

where summation extends over all positive roots of (2).

$$\text{Problem 14. Solve } \frac{\partial^2 f}{\partial r^2} + \frac{1}{r} \frac{\partial f}{\partial r} = \frac{1}{k} \frac{\partial f}{\partial t}, 0 \leq r < 1, t > 0$$

where $f=f_0$ when $r=1$, $t > 0$ and $f=0$ when $t=0$, $0 \leq r < 1$.

Hint. Take $\tilde{f}(p) = \int_0^1 f(r) J_0(pr) dr$, p being root of $J_0(p)=0$ and proceed like in Problem 13.

$$\text{Ans. } f = 2 f_0 \sum_p (1 - e^{-k p^2 t}) \frac{J_0(pr)}{p J_1(p)}$$

Problem 15. Viscous fluid is contained between two infinitely long concentric circular cylinders of radii a and b . The inner cylinder is kept at rest and outer cylinder suddenly starts rotating with uniform angular velocity ω . Find the velocity v of the fluid if the equation of motion is

$$\frac{\partial^2 v}{\partial r^2} + \frac{1}{r} \frac{\partial v}{\partial r} - \frac{v}{r^2} = \frac{1}{\nu} \frac{\partial v}{\partial t}, a < r < b, t > 0$$

ν being Kinematic viscosity.

$$\text{Hint. Take } f^2(p) = \int_a^b f(r) \cdot r B_1(pr) dr, a < b > a \text{ where}$$

$B_1(pr) = J_1(pr) Y_1(pa) - Y_1(pr) J_1(pa)$, $Y_1(pr)$ being Bessel's function of second kind of order one, and p is a positive root of

$$J_1(pb) Y_1(pa) = Y_1(pb) J_1(pa).$$

Multiplying the given equation by $rB_1(pr)$ and integrate w.r.t. ' r ' from a to b with boundary conditions $v=b\omega$ when $r=b$

$$v=0 \quad \text{when } r=a$$

$$v=0 \quad \text{when } t=0.$$

$$\text{Ans. } v = \pi b \omega \sum_p \frac{1 - e^{-v p^2 t}}{J_1^2(pa) - J_1^2(pb)} J_1(pa) J_1(pb) B_1(pr).$$

Problem 16. If α and β are the roots of the equation $J_0(x) = 0$, show that

$$\int_0^1 x J_0(\alpha x) J_0(\beta x) dx = \frac{1}{2} \delta_{\alpha\beta} J_1^2(\alpha) \quad (\text{Agra, 1971})$$

CHAPTER 12

DIFFUSION, WAVE AND LAPLACE'S EQUATIONS

12.1. INTRODUCTION

In 7.5, dealing with special types of differential equations we have already mentioned diffusion equation (or Fourier equation of heat flow), Laplace's equations as particular cases of steady-state heat flow and wave equations in one, two and three dimensions along with the two methods of solutions namely (i) method of separation of variables and (ii) D'Alembert's method. Now in subsequent chapters namely, 8th, 9th, 10th and 11th chapters we have discussed various methods of solution which may be successfully applied to solve different types of equations encountered in physics, mechanics and applied mathematics. In the present chapter we shall try to make a systematic study of one, two and three dimensional diffusion, Laplace's and wave equations with boundary value problems while a few of them have been already solved by the methods of transforms.

12.2. DIFFUSION EQUATION OR FOURIER EQUATION OF HEAT FLOW (Agra, 1974)

Assuming that the temperature at any point (x, y, z) of a solid at time t is $u(x, y, z, t)$, the thermal conductivity of the solid is K , the density of the solid is ρ and specific heat is σ , the heat equation

$$\frac{\partial u}{\partial t} = h^2 \nabla^2 u \quad \dots (1)$$

where $h^2 = \frac{K}{\rho\sigma} = k$ (say), k being known as diffusivity, is said to be the equation of diffusion or the Fourier equation of heat flow.

We know that heat flows from points at higher temperature to the points at lower temperature and the rate of decrease of temperature at any point varies with the direction. In other words the amount of heat say ΔH crossing an element of surface ΔS in Δt seconds is proportional to the greatest rate of decrease of the temperature u i.e.

$$\Delta H = K \Delta S \Delta t \left| \frac{du}{dt} \right| \quad \dots (2)$$

If $\bar{v}$ be the velocity of heat flow given by

$$\bar{v} = -K \text{ grad } u = -K \cdot \nabla u \quad \dots (3)$$




Fig. 12.1

where $\hat{n}$ is the positive outward drawn normal vector to the element dS and the negative sign shows the increase of temperature with the increase of x so that $\frac{\partial u}{\partial x}$ is positive and heat flows towards negative x from points of higher temperature to those of lower temperature, thereby rendering the flux to be negative.

Now applying Gauss's divergence theorem according to which if V be the volume bounded by a closed surface S and A be a vector function of position with continuous derivative, we have the quantity of heat entering S per unit time as

$$\iint_S (K \nabla u) \cdot \hat{n} dS = \iiint_V \nabla \cdot (K \nabla u) dV \quad \dots (5)$$

$$\text{i.e.,} \quad \iiint_V \nabla \cdot A dV = \iint_S A \cdot \hat{n} dS = \oiint_S A \cdot dS \quad \dots (6)$$

Taking volume element $dv = dx dy dz$, the heat contained in

$$V = \iiint_V \sigma \rho u dV.$$

$\therefore$ The time rate of increase of heat is given by

$$\frac{\partial}{\partial t} \iiint_V \sigma \rho u dV = \iiint_V \sigma \rho \frac{\partial u}{\partial t} dV \quad \dots (7)$$

Equating R.H.S's of (5) and (7), we find

$$\iiint_V \left[\sigma \rho \frac{\partial u}{\partial t} - \nabla \cdot (K \nabla u) \right] dV = 0 \quad \dots (8)$$

But V being arbitrary and the integrand being assumed to be continuous the relation (8) will be identically zero for every point if

$$\sigma \rho \frac{\partial u}{\partial t} = \nabla \cdot (K \nabla u)$$

$$\text{or} \quad \frac{\partial u}{\partial t} = \frac{K}{\sigma \rho} \nabla \cdot \nabla u = h^2 \nabla^2 u = k \nabla^2 u \text{ where } h^2 = k = \frac{K}{\sigma \rho}.$$

$$\text{or} \quad \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = \frac{1}{k} \frac{\partial u}{\partial t} = h^2 \frac{\partial u}{\partial t} \quad \dots (9)$$

This is three dimensional diffusion equation.

COROLLARY 1. If the temperature within a substance be assumed to be independent of z i.e., there being no heat flow in direction of z , then (9) reduces to $\frac{\partial u}{\partial t} = h^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$ $\dots (10)$

which is known as two dimensional diffusion equation or the equation for two dimensional flow parallel to x - y plane.

$$\text{COROLLARY 2. Putting } \frac{\partial u}{\partial y} = \frac{\partial u}{\partial z} = 0 \text{ in (9) we get } \frac{\partial u}{\partial t} = h^2 \frac{\partial^2 u}{\partial x^2} \quad \dots (11)$$

which is the equation for the one-dimensional flow of heat along a bar.

COROLLARY 3. For steady-state heat flow, u is independent of time i.e., $\frac{\partial u}{\partial t} = 0$ and hence (9) reduces to

$$\nabla^2 u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0 \quad \dots (12)$$

which is known as three-dimensional Laplace's equation.

12.3. ONE-DIMENSIONAL DIFFUSION EQUATION

$$\frac{\partial u}{\partial t} = h^2 \frac{\partial^2 u}{\partial x^2} = k \frac{\partial^2 u}{\partial x^2}$$

$$[A] \text{ Independent derivation of } \frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}.$$

Consider one-dimensional flow of electricity in a long insulated cable and specify the current i and voltage E at any time in the cable by x -coordinate and time-variable t .

The potential drop E in a line-element δx of length at any point x is given by $-\delta E = iR\delta x + L\delta x \frac{\partial i}{\partial t}$ $\dots (1)$

where R and L are respectively resistance and inductance per unit length.

If C and G be respectively capacitance to earth and conductance per unit length, then we have

$$-\delta i = GE\delta x + C\delta x \frac{\partial E}{\partial t} \quad \dots (2)$$

$$\text{Rewriting (1) and (2), } \frac{\partial E}{\partial x} + Ri + L \frac{\partial i}{\partial t} = 0 \quad \dots (3)$$

and
$$\frac{\partial i}{\partial x} + GE + C \frac{\partial E}{\partial t} = 0 \quad \dots (4)$$

Differentiating (3) w.r.t. x and (4) w.r.t. t we have

$$\frac{\partial^2 E}{\partial x^2} + R \frac{\partial i}{\partial x} + L \frac{\partial^2 i}{\partial x \partial t} = 0 \quad \dots (5)$$

and
$$\frac{\partial^2 i}{\partial x \partial t} + G \frac{\partial E}{\partial t} + C \frac{\partial^2 E}{\partial t^2} = 0 \quad \dots (6)$$

Eliminating $\frac{\partial^2 i}{\partial x \partial t}$ from (5) and (6) we get

$$\frac{\partial^2 E}{\partial x^2} = CL \frac{\partial^2 E}{\partial t^2} + LG \frac{\partial E}{\partial t} - R \frac{\partial i}{\partial x} \quad \dots (7)$$

Again eliminating $\frac{\partial i}{\partial x}$ from (4) and (7) we find

$$\frac{\partial^2 E}{\partial x^2} = CL \frac{\partial^2 E}{\partial t^2} + (CR + GL) \frac{\partial E}{\partial t} + RGE \quad \dots (8)$$

Differentiation of (3) w.r.t. ' t ' and (4) w.r.t. ' x ' yields

$$\frac{\partial^2 E}{\partial x \partial t} + R \frac{\partial i}{\partial t} + L \frac{\partial^2 i}{\partial t^2} = 0 \quad \dots (9)$$

and
$$\frac{\partial^2 i}{\partial x^2} + G \frac{\partial E}{\partial x} + C \frac{\partial^2 E}{\partial x \partial t} = 0 \quad \dots (10)$$

Elimination of $\frac{\partial E}{\partial x}$ and $\frac{\partial^2 E}{\partial x \partial t}$ from (3), (9) and (10) gives

$$\frac{\partial^2 i}{\partial x^2} = CL \frac{\partial^2 i}{\partial t^2} + (CR + GL) \frac{\partial i}{\partial t} + RGi \quad \dots (11)$$

(7) and (11) follow that E and i satisfy a second order partial differential equation

$$\frac{\partial^2 u}{\partial x^2} = CL \frac{\partial^2 u}{\partial t^2} + (CR + LG) \frac{\partial u}{\partial t} + RGu \quad \dots (12)$$

which is known as *telegraphy equation*.

If the leakage to the ground is small then $G=0=L$ and hence (12) reduces to $\frac{\partial^2 u}{\partial x^2} = CR \frac{\partial u}{\partial t} = \frac{1}{k} \frac{\partial u}{\partial t}$ where $k = \frac{1}{CR}$.

which is one-dimensional diffusion equation.

[B] Solution of $\frac{\partial u}{\partial t} = h^2 \frac{\partial^2 u}{\partial x^2}$ or $u_t = h^2 u_{xx}$.

The solution of this equation by the method of separation of variables has already been discussed in §7.5. Here below we discuss the solution in different conditions.

[b₁] (Both the ends of a bar at temperature zero).

If both the ends of a bar of length l are at temperature zero and the initial temperature is to be prescribed function $F(x)$ in the bar, then find the temperature at a subsequent time t .

One dimensional heat equation is $\frac{\partial u}{\partial t} = h^2 \frac{\partial^2 u}{\partial x^2} \quad \dots (1)$

we have to find a function $u(x, t)$ satisfying (1) with the boundary conditions $u(0, t) = u(l, t) = 0, t \geq 0, l$ being the length of bar. $\dots (2)$

and $u(x, 0) = F(x), 0 < x < l \quad \dots (3)$

In order to apply the method of separation of variables, let us assume that

$$u(x, t) = X(x) T(t), X \text{ and } T \text{ being respectively the functions of } x \text{ and } t \text{ alone.}$$

So that $\frac{\partial u}{\partial t} = X \frac{dT}{dt}$ and $\frac{\partial^2 u}{\partial x^2} = T \frac{d^2 X}{dx^2}$.

Their substitution in (1) gives

$$\frac{1}{X} \frac{d^2 X}{dx^2} = \frac{1}{h^2 T} \frac{dT}{dt} \quad \dots (4)$$

The L.H.S. and R.H.S. of (4) are constants because of variables being separated and hence we can write $\frac{1}{X} \frac{d^2 X}{dx^2} = \frac{1}{h^2 T} \frac{dT}{dt} = -\lambda^2$ (constant of separation).

Here $\frac{1}{X} \frac{d^2 X}{dx^2} = -\lambda^2$ i.e., $\frac{d^2 X}{dx^2} + \lambda^2 X = 0$ gives $X = A \cos \lambda x + B \sin \lambda x \quad \dots (5)$

and $\frac{1}{h^2 T} \frac{dT}{dt} = -\lambda^2$ i.e., $\frac{dT}{dt} + h^2 \lambda^2 T = 0$ gives $T = Ce^{-\lambda^2 h^2 t} \quad \dots (6)$

In view of condition (2) i.e., $u=0$ at $x=0$, or (5) gives $A=0$ and λ be chosen such that $\sin \lambda l = 0$ i.e., $\lambda = \frac{n\pi}{l}$, n being an integer.

Hence the solution (1) i.e., $u=XT$ takes the form

$$u = B \sin \frac{n\pi}{l} x e^{-\frac{n^2 \pi^2 h^2}{l^2} t} \quad \dots (7)$$

Summing over for all values of n , this becomes

$$u(x, t) = \sum_{n=1}^{\infty} B_n \sin \left(\frac{n\pi}{l} x \right) e^{-\frac{n^2 \pi^2 h^2}{l^2} t} \quad \dots (8)$$

Applying condition (3) i.e., $u(x, 0) = F(x)$ at $t=0$ we have

$$F(x) = \sum_{n=1}^{\infty} B_n \sin \frac{n\pi}{l} x \text{ for } 0 < x < l \quad \dots (9)$$

So that $B_n = \frac{2}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx \quad \dots (10)$

which is obtained by multiplying (9) by $\sin \frac{n\pi x}{l}$ and then integrating from $x=0$ to $x=l$.

Hence the required solution is

$$u(x, t) = \frac{2}{l} \sum_{n=1}^{\infty} e^{-\frac{n^2 \pi^2 h^2}{l^2} x} \sin \frac{n\pi}{l} x \int_0^l F(u) \sin \frac{n\pi}{l} u du \dots (11)$$

Deduction: (Insulated faces)

If instead of the ends of a bar of length l having kept at temperature zero, they are impervious to heat and the initial temperature is the prescribed function $F(x)$ in the bar, then to find the temperature at a subsequent time t , we have the boundary conditions

$$\frac{\partial u}{\partial x} = 0 \text{ at } x=0 \text{ or } l \text{ for all } t \dots (12)$$

$$u(x, 0) = F(x), 0 < x < l \dots (13)$$

Then the solution follows from (5) as

$$u = A \cos \lambda x + B \sin \lambda x$$

which in view of (12) requires $B=0$ and $\sin \lambda x = 0$ i.e., $\lambda = \frac{n\pi}{l}$, $n=0, 1, 2, 3 \dots$

So that the general solution of the one-dimensional diffusion equation will be of the form

$$u(x, t) = B_0 + \sum_{n=1}^{\infty} A_n e^{-\frac{n^2 \pi^2 h^2}{l^2} t} \cos \frac{n\pi x}{l} \dots (14)$$

where B_0 corresponds to $n=0$.

$$\text{By (13), this yields, } F(x) = u(x, 0) = B_0 + \sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{l} \dots (15)$$

from which we can easily find the coefficients

$$A_n = \frac{2}{l} \int_0^l F(x) \cos \frac{n\pi x}{l} dx \dots (16)$$

$$\text{and } B_0 = \frac{1}{2} A_0 \dots (17)$$

Note 1. The temperature in a slab having initial temperature $F(x)$ and the faces $x=0, x=l$ thermally insulated is given by

$$u(x, t) = B_0 + \sum_{n=1}^{\infty} B_n e^{-n^2 h^2 t} \cos nx \dots (18)$$

$$\text{where } A_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx dx \dots (19)$$

$$\text{and } B_0 = \frac{1}{2} A_0 = \frac{1}{\pi} \int_0^{\pi} F(x) dx \dots (20)$$

Note 2. The temperature in a slab having initial temperature $F(x)$ and the faces $x=0, x=l$ thermally insulated is given by

$$u(x, t) = \frac{1}{l} \int_0^l F(x) dx + \frac{2}{l} \sum_{n=1}^{\infty} e^{-\frac{n^2 \pi^2 h^2}{l^2} t} \cos \frac{n\pi x}{l} \int_0^l F(x) \frac{n\pi x}{l} dx \dots (21)$$

[b₂] (One end of a bar at temperature u_0 and other at zero temperature).

If a bar of length l is at temperature v_0 such that one of its ends $x=0$ is kept at zero temperature and the other end $x=l$ is kept at temperature u_0 , then find the temperature at any point x of the bar at an instant of time $t > 0$.

or

A rod of length l and thermal conductivity h^2 is maintained at a uniform temperature v_0 . At $t=0$ the end $x=0$ is suddenly cooled to 0°C by application of ice and the end $x=l$ is heated to the temperature u_0 by applying steam, the rod being insulated along its length so that no heat can transfer from the sides. Find the temperature of the rod at any point at any time.

$$\text{The equation is } \frac{\partial u}{\partial t} = h^2 \frac{\partial^2 u}{\partial x^2}, 0 < x < l, t > 0 \dots (1)$$

$$\text{With boundary conditions } u(0, t) = 0, u(l, t) = u_0 \text{ for all } t \dots (2)$$

$$\text{and } u(x, 0) = v_0 \dots (3)$$

$$\text{Let the solution of (1) be } u(x, t) = X(x)T(t) \dots (4)$$

where X is a function of x alone and T is a function of t alone.

$$\text{Substituting from (4) } \frac{\partial u}{\partial t} = X \frac{dT}{dt} \text{ and } \frac{\partial^2 u}{\partial x^2} = T \frac{d^2 X}{dx^2}, \text{ in (1) we}$$

get $\frac{1}{X} \frac{d^2 X}{dx^2} = \frac{1}{h^2 T} \frac{dT}{dt}$ where variables are separated and hence terms as either side are constants.

Now there arise three possibilities :

$$[1] \frac{d^2 X}{dx^2} = 0, \frac{dT}{dt} = 0 \text{ whence the solution is } X = Ax + B, T = C \dots (5)$$

$$[2] \frac{d^2 X}{dx^2} = \lambda^2 x, \frac{dT}{dt} = h^2 \lambda^2 T, \text{ the solution being } X = Ae^{\lambda x} + Be^{-\lambda x}, T = Ce^{h^2 \lambda^2 t} \dots (6)$$

$$[3] \frac{d^2 X}{dx^2} = -\lambda^2 x, \frac{dT}{dt} = -h^2 \lambda^2 T, \text{ the solution being } X = A \cos \lambda x + B \sin \lambda x, C = e^{-h^2 \lambda^2 t} \dots (7)$$

The combined solution in any of the three cases is $u = XT$. But $u = XT$ increases indefinitely with time t so possibility [2] is ruled out since then $u \rightarrow \infty$ as $t \rightarrow \infty$. Conclusively the possibilities [1] and [3] determine the solution of (1) in the form

$$u(x, t) = u_s(x) + u_T(x, t) \dots (8)$$

where $u_s(x)$ is the temperature distribution after a long interval of time when there exists steady state of temperature and $u_T(x, t)$ is the transient effects which die down when the time passes. Consequently there exists uniform temperature after one end $x=0$ being kept at zero temperature and the end $x=l$ at $u=u_0$ so that

$$u_S(x) = \frac{u_0}{l} x, \text{ whence (8) yields } u(x, t) = \frac{u_0}{l} x + u_T(x, t) \quad \dots (9)$$

$$\text{with boundary conditions } u_T(0, t) = 0 = u_T(l, t) \text{ by (2)} \quad \dots (10)$$

$$\text{and } u_T(x, 0) = v_0 - \frac{u_0}{l} x \text{ by (3)} \quad \dots (11)$$

Hence the possibility [3] i.e., the solution (7) reduces to

$$u_T(x, t) = (A \cos \lambda x + B \sin \lambda x) e^{-\lambda^2 h^2 t^2} \quad \dots (12)$$

whence in view of (2), this requires $A=0$ and $\sin \lambda l = 0$ i.e., $\lambda = \frac{n\pi}{l}$, n being an integer.

We thus obtain a solution

$$u_T(x, t) = \sum_{n=1}^{\infty} B_n e^{-\frac{n^2 \pi^2 h^2}{l^2} t} \sin \frac{n\pi}{l} x \quad \dots (13)$$

In view of (11), this gives $u_T(x, 0) = v_0 - \frac{u_0}{l} x = \sum_{n=1}^{\infty} B_n \sin \frac{n\pi}{l} x$.

$$\Delta \quad B_n = \frac{2}{l} \int_0^l \left(v_0 - \frac{u_0}{l} x \right) \sin \frac{n\pi}{l} x \, dx$$

$$= \frac{2}{n\pi} [v_0 - (-1)^n (v_0 - u_0)] \text{ (on integrating by parts)}$$

Hence the general solution of (1) with the help of (9) and (13) is

$$u(x, t) = \frac{u_0}{l} x + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} [v_0 - (-1)^n (v_0 - u_0)] e^{-\frac{n^2 \pi^2 h^2}{l^2} t} \sin \frac{n\pi}{l} x \quad \dots (14)$$

which gives temperature at any point x of the bar at any time $t > 0$.

Note. If we set $v_0 = 0$, then (14) takes the form

$$u(x, t) = \frac{u_0}{l} \left[x + \frac{2l}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} (-1)^n e^{-\frac{n^2 \pi^2 h^2}{l^2} t} \sin \frac{n\pi}{l} x \right] \quad \dots (15)$$

[b₃] (Temperature in an infinite bar)

If an infinite bar of small cross-section is insulated such that there is no transfer of heat at the surface and the temperature of the bar at $t=0$ is given by an arbitrary function $F(x)$ of x (taking the bar along x -axis), then find the temperature of the rod at any point of the bar at any time t .

$$\text{The boundary value problem is } \frac{\partial u}{\partial t} = h^2 \frac{\partial^2 u}{\partial x^2} \quad \dots (1)$$

$$\text{With initial condition, } u(x, 0) = F(x), \quad -\infty < x < \infty \quad \dots (2)$$

$$\text{Let the solution be } u(x, t) = X(x)T(t) \quad \dots (3)$$

$$\text{whence (1) gives } \frac{1}{X} \frac{d^2 X}{dx^2} = \frac{1}{h^2 T} \frac{dT}{dt} = -\lambda^2 \text{ (say)} \quad \dots (4)$$

Then the solution of (1) is

$$u(x, t) = XT = (A \cos \lambda x + B \sin \lambda x) e^{-h^2 \lambda^2 t} \quad \dots (5)$$

Here the arbitrary constants A and B being periodic may be taken as $A = A(\lambda)$, $B = B(\lambda)$ and due to the linearity and homogeneity of the heat equation we may write

$$u(x, t) = \int_0^{\infty} u(x, t, \lambda) \, d\lambda = \int_0^{\infty} e^{-h^2 \lambda^2 t} [A(\lambda) \cos \lambda x + B(\lambda) \sin (\lambda x)] \, d\lambda \quad \dots (6)$$

The condition (2) claims that

$$u(x, 0) = F(x) = \int_0^{\infty} [A(\lambda) \cos \lambda x + B(\lambda) \sin \lambda x] \, d\lambda$$

In view of Fourier's integrals we have

$$A(\lambda) = \frac{1}{\pi} \int_{-\infty}^{\infty} F(\mu) \cos (\mu \lambda) \, d\mu \text{ and } B(\lambda) = \frac{1}{\pi} \int_0^{\infty} F(\mu) \sin (\lambda \mu) \, d\mu$$

$$\text{so that } u(x, 0) = \frac{1}{\pi} \int_0^{\infty} \left[\int_{-\infty}^{\infty} F(\mu) \cos \lambda (x - \mu) \, d\mu \right] \, d\lambda$$

As such (6) takes the form

$$u(x, t) = \frac{1}{\pi} \int_0^{\infty} \left[\int_{-\infty}^{\infty} F(\mu) \cos \lambda (x - \mu) e^{-h^2 \lambda^2 t} \, d\mu \right] \, d\lambda$$

$$= \frac{1}{\pi} \int_{-\infty}^{\infty} F(\mu) \left[\int_0^{\infty} e^{-h^2 \lambda^2 t} \cos \lambda (x - \mu) \, d\lambda \right] \, d\mu \quad \dots (7)$$

$$\text{But we know that } \int_0^{\infty} e^{-x^2} \cos 2bx \, dx = \sqrt{\pi} \frac{e^{-b^2}}{2}$$

$$\text{So that } \int_0^{\infty} e^{-h^2 \lambda^2 t} \cos \lambda (x - \mu) \, d\lambda = \frac{\sqrt{\pi}}{2h\sqrt{t}} e^{-\left(\frac{x-\mu}{2h\sqrt{t}}\right)^2}$$

$$= \frac{\sqrt{\pi}}{2h\sqrt{t}} e^{-\frac{(x-\mu)^2}{4h^2 t}}$$

Hence (7) gives

$$\text{as } u(x, t) = \frac{1}{2h\sqrt{\pi t}} \int_{-\infty}^{\infty} F(\mu) e^{-\frac{(x-\mu)^2}{4h^2 t}} \, d\mu \quad \dots (8)$$

which gives the required temperature at any point at any time.

Problem 1. Solve $\frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$ when $u=0$ for $t=\infty$ and $x=0$ or l .

Taking $u(x, t) = X(x)T(t)$, the solution of the given equation is

$$X = A \cos \lambda x + B \sin \lambda x, \quad T = C e^{-\lambda^2 t} \text{ by §12.3 [b]}_1$$

with boundary conditions, $u(0, t) = u(l, t) = 0$ and $u(x, \infty) = 0$.

Hence putting $h=1$ in (8) of §12.3 [b₁] the required solution is

$$u(x, t) = \sum_{n=1}^{\infty} B_n \sin \frac{x\pi}{l} e^{-\frac{n^2 \pi^2}{l^2} t}$$

Problem 2. Solve $\frac{\partial \theta}{\partial t} = h^2 \frac{\partial^2 \theta}{\partial x^2}$ under the boundary conditions

$$\theta = (0, t) = \theta(l, t) = 0, t > 0 \quad \dots (1)$$

$$\text{and } \theta(x, 0) = x, 0 < x < l, \quad \dots (2)$$

l being the length of the bar.

Proceeding just like in §12.3 [b_1], we get the required solution on putting $\theta(x, 0) = F(x) = x$ in (8) of §12.3 [b_1],

$$\theta(x, t) = \sum_{n=1}^{\infty} B_n \sin \frac{n\pi x}{l} e^{-n^2\pi^2 h^2/l^2 t}$$

$$\text{where } B_n = \frac{2}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx = \frac{2}{l} \int_0^l x \sin \frac{n\pi x}{l} dx$$

$$= -\frac{2l}{n\pi} \cos n\pi$$

$$= \begin{cases} \frac{2l}{n\pi} & \text{when } n \text{ is odd} \\ -\frac{2l}{n\pi} & \text{when } n \text{ is even} \end{cases}$$

$$\text{Hence } \theta(x, t) = \frac{2l}{\pi} \left[e^{-\pi^2 h^2 t/l^2} \sin \frac{\pi x}{l} - \frac{1}{2} e^{-2^2\pi^2 h^2 t/l^2} \right.$$

$$\left. \sin \frac{2\pi x}{l} + \frac{1}{4} e^{-3^2\pi^2 h^2 t/l^2} \sin \frac{3\pi x}{l} \dots \right]$$

Problem 3. Find the temperature $u(x, t)$ in a bar of length l , perfectly insulated, and whose ends are kept at temperature zero while the initial temperature is given by

$$F(x) = \begin{cases} x, & 0 < x < l/2 \\ l-x, & l/2 < x < l. \end{cases}$$

The boundary value problem is $\frac{\partial u}{\partial t} = h^2 \frac{\partial^2 u}{\partial x^2}$.

With conditions $u(0, t) = u(l, t) = 0$ and $u(x, 0) =$

$$F(x) = \begin{cases} x, & 0 < x < l/2 \\ l-x, & l/2 < x < l. \end{cases}$$

Hence by (8) of §12.3 [b_1] the required solution is

$$u(x, t) = \sum_{n=1}^{\infty} B_n F(x) \sin \frac{n\pi x}{l} e^{-n^2\pi^2 h^2 t/l^2}$$

$$\text{where } B_n = \frac{2}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx = \frac{2}{l} \int_0^{l/2} x \sin \frac{n\pi x}{l} dx$$

$$+ \int_{l/2}^l (l-x) \sin \frac{n\pi x}{l} dx$$

$$= \frac{4l}{n^2\pi^2} \sin \frac{n\pi}{2} = \begin{cases} \frac{4l}{n^2\pi^2} & \text{for } n=1, 5, 9, \dots \\ 0 & \text{for } n=2, 4, 6, \dots \\ -\frac{4l}{n^2\pi^2} & \text{for } n=3, 7, 11, \dots \end{cases}$$

Hence the solution is

$$u(x, t) = \frac{4l}{\pi^2} \left[\frac{1}{1^2} \sin \frac{\pi x}{l} e^{-h^2\pi^2 t/l^2} - \frac{1}{3^2} \sin \frac{3\pi x}{l} e^{-3^2 h^2\pi^2 t/l^2} + \dots \right]$$

Note. Had we considered the case of slab with its ends $x=0$ and $x=l$ maintained at temperature zero and initial temperature being

$$F(x) = \begin{cases} T_0, & 0 < x < l/2 \\ 0, & l/2 < x < l/2 \end{cases}$$

then we should have

$$B_n = \frac{2}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx = \frac{2T_0}{l} \int_0^{l/2} \sin \frac{n\pi x}{l} dx$$

$$= \frac{4T_0}{n\pi} \sin^2 \frac{n\pi}{4}$$

and the solution would be

$$u(x, t) = \frac{4T_0}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \sin^2 \frac{n\pi}{4} e^{-n^2\pi^2 h^2 t/l^2} \sin \frac{n\pi x}{l}$$

Problem 4. Solve $\frac{\partial u}{\partial t} = h^2 \frac{\partial^2 u}{\partial x^2}$, $0 < x < \pi$, $t < 0$, under the boundary conditions

$$u_x(0, t) = 0 = u_x(\pi, t) \text{ and } u(x, 0) = \sin x$$

By (14) of §12.3 [b_1], we have

$$u(x, t) = B_0 + \sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{l} e^{-n^2\pi^2 h^2 t/l^2} \text{ where}$$

$$B_0 = \frac{A_0}{2} \text{ and } l = \pi$$

$$= \frac{A_0}{2} = \sum_{n=1}^{\infty} A_n^* \cos nx e^{-n^2 h^2 t}$$

$$\text{where } A_n = \frac{2}{\pi} \int_0^{\pi} \sin x \cos nx dx = \begin{cases} 0, & \text{when } n \text{ is odd} \\ \frac{4}{\pi(4m^2-1)}, & \text{when } n=2m \end{cases}$$

$$\text{and } A_0 = \frac{2}{\pi} \int_0^{\pi} \sin x dx = \frac{4}{\pi}$$

Hence the required solution is

$$u(x, t) = \frac{2}{\pi} - \frac{4}{\pi} \sum_{m=1}^{\infty} (4m^2 - 1)^{-1} e^{-4m^2 h^2 t} \cos 2mx.$$

Problem 5. The face $x=0$ of a slab is maintained at temperature zero and heat is supplied at constant rate at the face $x=\pi$, so that

$\frac{\partial u}{\partial x} = \mu$ when $x=\pi$. If the initial temperature is zero, show that

$$u(x, t) = \mu x + \sum_{j=1}^{\infty} \frac{(-1)^j}{(j-\frac{1}{2})^2} \sin(j-\frac{1}{2})x e^{-(j-\frac{1}{2})^2 t}$$

where the unit of time is so chosen that $k=1$.

Taking $u(x, t)$ as the temperature of the slab, the boundary value problem is $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$, $0 < x < \pi$, $t > 0$ (1)

with condition $u(0, t) = 0$... (2)

$u(x, 0) = 0$... (3)

and $\frac{\partial}{\partial x} u(\pi, t) = \mu$... (4)

Applying the method of separation of variables, the solutions of the given equation are (i) $u = (Ae^{\lambda x} + Be^{-\lambda x})e^{\lambda^2 t}$

(ii) $u = A_1 + B_1 x$

(iii) $u = (A_2 \cos \pi x + B_2 \sin \pi x)e^{-\lambda^2 t}$

according as the constant of variation is λ^2 or $-\lambda^2$.

Here (i) is inadmissible as $u \rightarrow \infty$ when $t \rightarrow \infty$.

(ii) alone is inadequate to give the complete solution and hence the complete solution is given by (ii) and (iii) jointly

i.e. $u(x, t) = u_s(x) + u_T(x, t)$... (5)

where $u_s(x)$ is the temperature distribution after a long period of time when the slab has reached the steady state of the temperature distribution and $u_T(x, t)$ denotes the transient effects which die down with the passage of time.

From (ii) $u_s(x) = A_1 + B_1 x$... (6)

and from (iii) $u_T(x, t) = (A_2 \cos \lambda x + B_2 \sin \lambda x) e^{-\lambda^2 t}$... (7)

Applying (2), (6) gives $A_1 = 0$ and by (4), (6) gives $\mu = B_1$ so that $u_s = \mu x$... (8)

Thus with the help of (7) and (8), (5) reduces to

$u(x, t) = \mu x + (A_2 \cos \lambda x + B_2 \sin \lambda x) e^{-\lambda^2 t}$... (9)

Applying (2), i.e. $u(0, t) = 0$, we get $A_2 = 0$... (10)

Applying (4), i.e. $\mu = \frac{\partial}{\partial x} u(\pi, t)$,

we have $(\mu + \lambda B_2 \cos \lambda \pi) e^{-\lambda^2 t} = \mu$

i.e. $\cos \lambda \pi = 0$ giving $\lambda \pi = (2j-1)\frac{\pi}{2}$ i.e. $\lambda = j - \frac{1}{2}$... (11)

As such $u_T(x, t) = B_j \sin(j-\frac{1}{2})x e^{-(j-\frac{1}{2})^2 t}$ where we have set $B_j = B_2$. Summing over all j , the general solution is

$$u_T(x, t) = \sum_{j=1}^{\infty} B_j \sin(j-\frac{1}{2})x e^{-(j-\frac{1}{2})^2 t} \dots (12)$$

Hence from (5)

$$u(x, t) = \mu x + \sum_{j=1}^{\infty} B_j \sin(j-\frac{1}{2})x e^{-(j-\frac{1}{2})^2 t} \dots (13)$$

Applying the condition (3), $0 = \mu x + \sum_{j=1}^{\infty} B_j \sin(j-\frac{1}{2})x$

$$\begin{aligned} \text{i.e. } -\mu x &= \sum_{j=1}^{\infty} B_j \sin(j-\frac{1}{2})x \text{ so that } B_j = \frac{2}{\pi} \int_0^{\pi} (-\mu x) \sin(j-\frac{1}{2})x dx \\ &= \frac{2\mu}{\pi} \frac{(-1)^j}{(j-\frac{1}{2})^2} \end{aligned}$$

Hence (13) reduces to

$$u(x, t) = \mu x + \frac{2\mu}{\pi} \sum_{j=1}^{\infty} \frac{(-1)^j}{(j-\frac{1}{2})^2} \sin(j-\frac{1}{2})x e^{-(j-\frac{1}{2})^2 t}$$

which is the required relation.

12.4. TWO-DIMENSIONAL DIFFUSION EQUATION

$$\text{i.e. } \frac{\partial u}{\partial t} = h^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$$

Consider a thin rectangular plate whose surface is impervious to heat flow and which has an arbitrary function of temperature $F(x, y)$ at $t=0$. Its four edges say $x=0$, $x=a$, $y=0$, $y=b$ are kept at zero temperature. We have to determine the subsequent temperature at a point of the plate as t increases

$$\text{The boundary value problem is } \frac{\partial u}{\partial t} = h^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) \dots (1)$$

Subject to the conditions for all t , (i) $u(0, y, t) = 0$, (ii) $u(a, y, t) = 0$, (iii) $u(x, 0, t) = 0$, (iv) $u(x, b, t) = 0$ and the initial condition (v) $u(x, y, 0) = F(x, y)$

In order to apply the method of separation of variables, let us assume that

$$u(x, y, t) = X(x) Y(y) T(t) \dots (2)$$

where X is a function of x alone, Y is a function of y alone and T is a function of t alone. From (2) we have,

$$\frac{\partial u}{\partial t} = XY \frac{dT}{dt}, \quad \frac{\partial^2 u}{\partial x^2} = YT \frac{d^2 X}{dx^2} \text{ and } \frac{\partial^2 u}{\partial y^2} = XT \frac{d^2 Y}{dy^2}$$

Substituting them in (1), we find

$$\frac{1}{h^2 T} \frac{dT}{dt} = \frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2}, \text{ after dividing by } XYT \dots (3)$$

In (3), the variables being separated, we can assume

$$\frac{1}{X} \frac{d^2 X}{dx^2} = -\lambda_1^2; \quad \frac{1}{Y} \frac{d^2 Y}{dy^2} = -\lambda_2^2 \quad \text{and} \quad \frac{1}{h^2 T} \frac{dT}{dt} = -\lambda^2$$

$$\text{so that} \quad \lambda^2 = \lambda_1^2 + \lambda_2^2 \dots (4)$$

The general solutions of (4) are

$$X = A \cos \lambda_1 x + B \sin \lambda_1 x; \quad Y = C \cos \lambda_2 y + D \sin \lambda_2 y; \\ T = E e^{-\lambda^2 h^2 t} \dots (5)$$

So that with the help of (5), (2) gives the solution of (1) in the form $u(x, y, t) = (A \cos \lambda_1 x + B \sin \lambda_1 x) (C \cos \lambda_2 y + D \sin \lambda_2 y) e^{-\lambda^2 h^2 t}$... (6)

In view of condition (i), $0 = u(0, y, t) = A(C \cos \lambda_2 y + D \sin \lambda_2 y) e^{-\lambda^2 h^2 t}$, giving $A = 0$.

In view of condition (ii), we claim $\sin \lambda_1 x = 0$ i.e. $\lambda_1 = \frac{m\pi}{a}$, m being an integer.

Similarly applying conditions (iii) and (iv) to (6), we get $C = 0$ and $\lambda_2 = \frac{n\pi}{b}$, n being an integer. As such (5) takes the form

$$u_{mn}(x, y, t) = B_{mn} e^{-\lambda_{mn}^2 h^2 t} \sin \frac{m\pi}{a} x \sin \frac{n\pi}{b} y, \text{ where}$$

$$\lambda^2 = \lambda_{mn}^2 = \pi^2 \left(\frac{m^2}{a^2} + \frac{n^2}{b^2} \right).$$

Summing over all the possible values of m and n , the general solution is

$$u(x, y, t) = \sum_{m, n=1}^{\infty} B_{mn} e^{-\lambda_{mn}^2 h^2 t} \sin \frac{m\pi}{a} x \sin \frac{n\pi}{b} y \dots (7)$$

where B_{mn} are arbitrary constants to be determined by the condition (v)

$$\text{i.e. } F(x, y) = u(x, y, 0) = \sum_{m, n=1}^{\infty} B_{mn} \sin \frac{m\pi}{a} x \sin \frac{n\pi}{b} y \dots (8)$$

Multiplying both sides of (8) by $\sin \frac{m\pi}{a} x \sin \frac{n\pi}{b} y dx dy$ and integrating with regard to x from 0 to a and with regard to y from 0 to b we get on using orthogonality properties of the sines,

$$B_{mn} = \frac{4}{ab} \int_0^a \int_0^b F(x, y) \sin \frac{m\pi}{a} x \sin \frac{n\pi}{b} y dx dy \dots (9)$$

which gives the arbitrary constants of (7).

Problem 6. A rectangular plate bounded by the lines $x=0, y=0, x=a, y=b$ has an initial distribution of temperature given by $F(x, y) = B \sin \frac{\pi x}{a} \sin \frac{\pi y}{b}$. The edges are maintained at zero temperature and the plane faces are impervious to heat. Find the temperature at any point at any time.

By (7) of §12.4, the general solution is

$$u(x, y, t) = \sum_{m, n=1}^{\infty} B_{mn} e^{-\lambda_{mn}^2 h^2 t} \left\{ \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} \right\}$$

where $B_{mn} = \frac{4}{ab} \int_0^a \int_0^b F(x, y) \sin \frac{m\pi}{a} x \sin \frac{n\pi}{b} y dx dy$ by

(9) of § 12.4.

$$= \frac{4B}{ab} \int_0^a \int_0^b \sin \frac{\pi x}{a} \sin \frac{\pi y}{b} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} dx dy \\ \therefore F(x, y) = B \sin \frac{\pi x}{a} \sin \frac{\pi y}{b}$$

$$\therefore B_{m1} = \frac{4B}{ab} \int_0^a \frac{b}{2} \sin \frac{\pi x}{a} \sin \frac{m\pi x}{a} dx \\ \therefore \int_0^b \sin \frac{\pi y}{b} \sin \frac{n\pi y}{b} dy = \begin{cases} 0, & \text{for } n=2, 3, 4, \dots \\ b/2, & \text{for } n=1. \end{cases}$$

So that $B_{11} = B$.

$$\text{Also } \lambda_{11}^2 = \pi^2 \left(\frac{1}{a^2} + \frac{1}{b^2} \right).$$

Hence the solution is

$$u(x, y, t) = B_{11} e^{-\lambda_{11}^2 h^2 t} \sin \frac{\pi x}{a} \sin \frac{\pi y}{b} \\ = B e^{-h^2 \pi^2 \left(\frac{1}{a^2} + \frac{1}{b^2} \right) t} \sin \frac{\pi x}{a} \sin \frac{\pi y}{b}.$$

Problem 7. A semi-infinite plate having width π has its faces insulated. The semi-infinite edges are kept at 0°C while the infinite edge is maintained at 100°C . Assuming that the initial temperature is 0°C , find the temperature at any point at any time.

Taking the diffusivity i.e. $h^2 = 1$, the boundary value problem is

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \dots (1)$$

- with conditions (i) $u(0, y, t) = 0$,
(ii) $u(\pi, y, t) = 0$, (iii) $u(x, y, 0) = 0$;
(iv) $u(x, 0, t) = 100$ and
(v) $|u(x, y, t)| < M$

where $0 < x < \pi, y > 0, t > 0$.

Taking Laplace transform of (1) and assuming $L\{u(x, y, t)\} = U(x, y, t)$,




Fig. 12.2.

We have $\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} = sU$ by using condition (iii) ... (2)

But the finite Fourier sine transform of a function $F(x)$, $0 < x < l$ is defined as

$$f_s(x) = \int_0^l F(x) \sin \frac{n\pi x}{l} dx, n \text{ being an integer} \quad \dots (3)$$

Multiplying (2) by $\sin nx$ and integrating from 0 to π , we get

$$\int_0^\pi \frac{\partial^2 U}{\partial x^2} \sin nx dx + \int_0^\pi \frac{\partial^2 U}{\partial y^2} \sin nx dx = \int_0^\pi sU \sin nx dx$$

Setting $\tilde{U} = \int_0^\pi U \sin nx dx$, this becomes

$$-n^2 \tilde{U} + nU(\pi, y, s) \cos n\pi + nU(0, y, s) + \frac{d^2 \tilde{U}}{dy^2} = s\tilde{U} \quad \dots (4)$$

But from the Laplace transforms of conditions (i) and (ii) we have $U(0, y, s) = 0$, $U(\pi, y, s) = 0$

$$\therefore (4) \text{ reduces to } \frac{d^2 \tilde{U}}{dy^2} - (n^2 + s) \tilde{U} = 0 \quad \dots (5)$$

$$\text{Its solution is } \tilde{U} = A e^{y\sqrt{n^2+s}} + B e^{-y\sqrt{n^2+s}} \quad \dots (6)$$

By condition (v), $\tilde{U}$ being bounded, as $y \rightarrow \infty$, we have $A = 0$, so that (6) yields, $\tilde{U} = B e^{-y\sqrt{n^2+s}} \quad \dots (7)$

Applying condition (iv),

$$\tilde{U}(n, 0, s) = \int_0^\pi \frac{100}{s} \sin nx dx = \frac{100}{s} \left(\frac{1 - \cos n\pi}{n} \right) \quad \dots (8)$$

In (7) if we put $y = 0$, we get with the help of (8),

$$B = \tilde{U} = \frac{100}{s} \left(1 - \frac{\cos n\pi}{n} \right)$$

$$\text{Hence } \tilde{U} = \frac{100}{s} \frac{(1 - \cos n\pi)}{n} e^{-y\sqrt{n^2+s}}$$

Applying Fourier sine inversion formula, we find

$$U = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{100}{s} \left(\frac{1 - \cos n\pi}{n} \right) e^{-y\sqrt{n^2+s}} \sin nx \quad \dots (9)$$

$$\text{Now we have } L^{-1} \left\{ e^{-y\sqrt{s+n^2}} \right\} = \frac{y}{2\sqrt{\pi t^3}} e^{-\frac{y^2}{4t}}$$

$$\text{so that } L^{-1} \left\{ e^{-y\sqrt{s+n^2}} \right\} = \frac{y}{2\sqrt{\pi t^3}} e^{-\frac{y^2}{4t}} e^{-n^2 t}$$

$$\text{Thus } L^{-1} \left\{ \frac{e^{-y\sqrt{s+n^2}}}{s} \right\} = \int_0^t \frac{y}{2\sqrt{\pi v^3}} e^{-y^2/4v} e^{-n^2 v} dv$$

$$= \frac{2}{\sqrt{\pi}} \int_0^\infty \frac{y}{2\sqrt{t}} e^{-(p^2+n^2 y^2/4p^2)} dp \text{ where } p^2 = \frac{y^2}{4v}$$

Hence taking the inverse Laplace transform of (9) term by term, we get

$$u(x, y, t) = \frac{400}{\pi^{3/2}} \sum_{n=1}^{\infty} \left(\frac{1 - \cos n\pi}{n} \right) \sin nx \int_0^\infty \frac{y}{2\sqrt{t}} e^{-(p^2+n^2 y^2/4p^2)} dp$$

12.5. THREE-DIMENSIONAL DIFFUSION EQUATION

$$\text{We have } \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = \frac{1}{t^2} \frac{\partial u}{\partial t} \quad \dots (1)$$

where $u = u(x, y, z, t)$.

$$\text{Let, } u(x, y, z, t) = X(x)Y(y)Z(z)T(t) \quad \dots (2)$$

where X, Y, Z, T being respectively the functions of x, y, z, t alone.

$$\text{From (2) we have } \frac{\partial u}{\partial t} = XYZ \frac{dT}{dt}, \quad \frac{\partial^2 u}{\partial x^2} = YZT \frac{d^2 X}{dx^2}, \quad \frac{\partial^2 u}{\partial y^2} = XZT \frac{d^2 Y}{dy^2} \text{ and } \frac{\partial^2 u}{\partial z^2} = XYT \frac{d^2 Z}{dz^2}$$

Their substitution in (1) yields

$$\frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2} + \frac{1}{Z} \frac{d^2 Z}{dz^2} = \frac{1}{h^2 T} \frac{dT}{dt} = -\lambda^2 \text{ (say) as variables are separable.}$$

$$\text{Now taking } \frac{1}{X} \frac{d^2 X}{dx^2} = -\lambda_1^2, \quad \frac{1}{Y} \frac{d^2 Y}{dy^2} = -\lambda_2^2, \quad \frac{d^2 Z}{dz^2} = -\lambda_3^2$$

so that $\lambda_1^2 + \lambda_2^2 + \lambda_3^2 = \lambda^2$, we get the solutions

$$X = A_2 \cos \lambda_1 x + B_1 \sin \lambda_1 x = a \cos (\lambda_1 x + \alpha_{\lambda_1})$$

Similarly $Y = b \cos (\lambda_2 x + \alpha_{\lambda_2})$, $Z = c \cos (\lambda_3 x + \alpha_{\lambda_3})$ and

$$T = de^{-\lambda^2 h^2 t} = de^{-(\lambda_1^2 + \lambda_2^2 + \lambda_3^2) \lambda^2 t}$$

Hence for all values of t , the general solution of (11) is

$$u(x, y, z, t) = \sum_{\lambda_1=0}^{\infty} \sum_{\lambda_2=0}^{\infty} \sum_{\lambda_3=0}^{\infty} A_{\lambda_1 \lambda_2 \lambda_3} \cos (\lambda_1 x + \alpha_{\lambda_1}) \cos (\lambda_2 x + \alpha_{\lambda_2}) \cos (\lambda_3 x + \alpha_{\lambda_3}) e^{-h^2 (\lambda_1^2 + \lambda_2^2 + \lambda_3^2) t}$$

12.6. LAPLACE'S EQUATION

In §12.2 (cor 3) we have already derived the Cartesian form of three dimensional Laplace's equation as a particular case of steady heat flow in the form.

$$\nabla^2 u \equiv \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0 \quad \dots (1)$$

In cylindrical coordinates, (r, θ, z) , it is as shown in §11.1,

$$\nabla^2 u \equiv \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial z^2} = 0 \quad \dots (2)$$

and in Polar spherical coordinates (r, θ, ϕ) , it is as shown in §8.1,

$$\nabla^2 u \equiv \frac{\partial^2 u}{\partial r^2} + \frac{2}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial u}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 u}{\partial \phi^2} = 0 \quad \dots (3)$$

Two-dimensional Cartesian form of Laplace's equation is

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \dots (4)$$

Taking u as independent of z , the two-dimensional Laplace-equation in cylindrical coordinates is given by

$$\frac{\partial^2 u}{\partial x^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0 \quad \dots (5)$$

and in Polar coordinates (r, θ) it resumes the same form as (5).

One-dimensional Laplace's equation is $\frac{d^2 u}{dx^2} = 0$. $\dots (6)$

Its solution being easy and straight has no points of worth consideration and hence we shall consider only two and three dimensional Laplace equations.

12.7. TWO-DIMENSIONAL LAPLACE'S EQUATION (STEADY FLOW OF HEAT)

[A] Solution of two-dimensional Laplace-equation in Cartesian coordinates

$$\text{We have } \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0, \quad \dots (1)$$

(taking temperature as independent of time).

This can be solved either by the method of separation of variables or by the application of integral transforms as is evident from the following problems.

Problem 8. Determine the steady state temperature distribution in a thin plate bounded by the lines $x=0$, $x=l$, $y=0$ and $y=\infty$, assuming that heat cannot escape from either surface of the plate, the edges $x=0$ and $x=l$ being kept at a temperature zero and also the lower edge $y=0$ is kept at temperature $F(x)$ and the edge $y=\infty$ at temperature zero.

The boundary value problem is

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \dots (1)$$

With conditions

- (i) $u(0, y)=0$, (ii) $u(l, y)=0$
- (iii) $u(x, 0)=F(x)$, and
- (iv) $u(x, \infty)=0$.

In order to apply the method of separation of variables, assume that $u(x, y)=X(x)Y(y)$ $\dots (2)$

$$\text{So that } \frac{\partial^2 u}{\partial x^2} = Y \frac{d^2 X}{dx^2} \text{ and } \frac{\partial^2 u}{\partial y^2} = X \frac{d^2 Y}{dy^2}$$

$$\therefore (1) \text{ gives } \frac{1}{X} \frac{d^2 X}{dx^2} = -\frac{1}{Y} \frac{d^2 Y}{dy^2} = -\lambda^2 \text{ (say) as}$$

variables are separated

$$\text{Here } \frac{1}{X} \frac{d^2 X}{dx^2} = -\lambda^2 \text{ i.e. } \frac{d^2 X}{dx^2} + \lambda^2 X = 0 \text{ gives}$$

$$X = A \cos \lambda x + B \sin \lambda x \quad \dots (3)$$

$$\text{and } \frac{1}{Y} \frac{d^2 Y}{dy^2} = \lambda^2 \text{ i.e. } \frac{d^2 Y}{dy^2} - \lambda^2 Y = 0 \text{ gives } Y = C e^{\lambda y} + D e^{-\lambda y} \quad \dots (4)$$

As such a solution of (1) is

$$u(x, y) = XY = (A \cos \lambda x + B \sin \lambda x) (C e^{\lambda y} + D e^{-\lambda y}) \quad \dots (5)$$

Applying condition (iv) we have $C=0$ and applying (i) $A=0$, so that (5) takes the form

$$u(x, y) = B \sin \lambda x \cdot e^{-\lambda y} \quad \dots (6)$$

But condition (ii) yields, $\sin \lambda l = 0$

$$\text{i.e. } \lambda = \frac{n\pi}{l}, n \text{ being an integer.}$$

Hence for all distinct n , the general solution of (1) is

$$u(x, y) = \sum_{n=0}^{\infty} B_n \sin \frac{n\pi x}{l} e^{-n\pi y/l} \quad \dots (7)$$

which gives the required temperature in the thin plate, where

$$B_n = \frac{2}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx \text{ and } F(x) = u(x, \infty) = \sum_{n=0}^{\infty} \sin \frac{n\pi x}{l}$$

Problem 9. Temperature distribution in a finite plate.

Find the steady state temperature distribution of a thin rectangular plate bounded by the lines $x=0$, $x=l$, $y=0$, $y=b$ assuming that the edges $x=0$, $x=l$ and $y=0$ are maintained at temperature zero while the edge $y=b$ is maintained at temperature $F(x)$.




Fig. 12.3

The boundary value Problem is $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \dots (1)$

with the conditions (i) $u(0, y) = 0$, (ii) $u(l, y) = 0$, (iii) $u(x, 0) = F(x)$ and (iv) $u(x, b) = 0$.

Proceeding just like in Problem 8, we get the general solution of (1) as

$$u(x, y) = \sum_{n=1}^{\infty} \left[B_n e^{n\pi y/l} + C_n e^{-n\pi y/l} \right] \sin \frac{n\pi x}{l} \quad \dots (2)$$

In view of condition (i), $C_n = -B_n$ so that (2) reduces to

$$u(x, y) = \sum_{n=1}^{\infty} B_n \left[e^{n\pi y/l} - e^{-n\pi y/l} \right] \sin \frac{n\pi x}{l} \\ = \sum_{n=1}^{\infty} D_n \sinh \frac{n\pi y}{l} \sin \frac{n\pi x}{l} \text{ on setting } D_n = 2B_n$$

By condition (iv), $F(x) = \sum_{n=1}^{\infty} D_n \sinh \frac{n\pi b}{l} \sin \frac{n\pi x}{l}$ so that

$$D_n \sinh \frac{n\pi b}{l} = \frac{2}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx.$$

Hence the solution is

$$u(x, y) = \frac{2}{l} \sum_{n=1}^{\infty} \sinh \frac{n\pi y}{l} \operatorname{cosech} \frac{n\pi b}{l} \sin \frac{n\pi x}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx.$$

Problem 10. (Insulated at one side). Determine the steady state temperature in a rectangular plate of length a and width b with sides maintained at temperature zero while the lower end is kept at temperature $F(x)$ and upper one insulated.

The boundary value problem is $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \dots (1)$

with conditions (i) $u(0, y) = 0$, (ii) $u(a, y) = 0$, (iii) $u(x, 0) = F(x)$ and (iv) $u(x, b) = 0$

Proceeding just like in Problem 8, we have

$$u(x, y) = \sum_{n=1}^{\infty} \left[C_n \cosh \frac{n\pi y}{a} + D_n \sinh \frac{n\pi y}{a} \right] \sin \frac{n\pi x}{a} \quad \dots (2)$$

In view of (iii) we have

$$F(x) = \sum_{n=1}^{\infty} C_n \sin \frac{n\pi x}{a} \text{ so that } C_n = \frac{2}{a} \int_0^a F(x) \sin \frac{n\pi x}{a} dx \text{ and}$$

in view of (iv), $0 = u(x, b) = \sum_{n=1}^{\infty} \frac{n\pi}{a} \left(C_n \cosh \frac{n\pi b}{a} + D_n \sinh \frac{n\pi b}{a} \right) \sin \frac{n\pi x}{a}$, giving $D_n = -C_n \tanh \frac{n\pi b}{a}$

Hence $u(x, y) = \frac{2}{a} \sum_{n=1}^{\infty} \left(\cosh \frac{n\pi y}{a} - \tanh \frac{n\pi y}{a} \sin \frac{n\pi y}{a} \right) \sin \frac{n\pi x}{a} \int_0^a F(x) \sin \frac{n\pi x}{a} dx.$

Problem 11. Heat flows in a semi-infinite rectangular plate, the end $x=0$ being kept at temperature $T^\circ\text{C}$ and the edges $y=0$ and $y=a$ at temperature zero, then show that the temperature at any point (x, y) is given by

$$u(x, y) = \frac{4T}{\pi} \sum_{r=1}^{\infty} \frac{1}{2r+1} \sin \frac{(2r+1)\pi y}{a} e^{-(2r+1)\pi x/a}$$

The boundary value problem is $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \dots (1)$

with conditions (i) $u=0$ when $y=0$, (ii) $u=0$ when $y=a$, (iii) $u=T$ when $x=0$.

The solution by usual method is

$$u(x, y) = (A \cos ny + B \sin ny) e^{-nx}$$

In view of (i), $0 = Ae^{-nx}$ i.e. $A=0$.

In view of (ii), $0 = B \sin na e^{-nx}$ giving $\sin na = 0$ i.e., $na = (2r+1)\pi$.

Hence the general solution is

$$u(x, y) = \sum_{r=0}^{\infty} B_r \sin (2r+1) \frac{\pi y}{a} e^{-(2r+1)\pi x/a} \quad \dots (2)$$

In view of (iii), $T = \sum_{r=0}^{\infty} B_r \sin (2r+1) \frac{\pi y}{a}$ so that

$$B_r = \frac{2}{a} \int_0^a T \sin (2r+1) \frac{\pi y}{a} dy = \frac{4T}{(2r+1)\pi}$$

Hence $u(x, y) = \frac{4T}{\pi} \sum_{r=0}^{\infty} \frac{1}{2r+1} \sin \frac{(2r+1)\pi y}{a} e^{-(2r+1)\pi x/a}$

Problem 12. A square plate has its faces and its edges $x=0$ and $x=\pi$ ($0 < y < \pi$) insulated. Its edges $y=0$ and $y=\pi$ are kept at temperatures zero and $f(x)$ respectively. Show that the formula for its steady temperature is

$$u(x, y) = \frac{1}{2\pi} a_0 y + \sum_{n=1}^{\infty} a_n \frac{\sinh ny}{\sinh n\pi} \cos nx, \text{ where}$$

$$a_n = \frac{2}{\pi} \int_0^\pi f(x) \cos nx dx, n=0, 1, 2, \dots$$

Hint. The boundary value problem is $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$ with conditions

(i) $u_x(0, y) = 0$, (ii) $u_x(\pi, y) = 0$, (iii) $u(x, 0) = 0$, (iv) $u(x, \pi) = f(x)$.
Apply method of separation of variables.

Problem 13. If $u(x, y)$ denotes the electrostatic potential in a region bounded by the planes $x=0$, $x=\pi$ and $y=0$ in which there is a uniform distribution of space charge of density $\frac{h}{4\pi}$. If the planes $x=0$ and $y=0$ are kept at potential zero, the plane $x=\pi$ at an other fixed potential $u=1$ and u is finite as $y \rightarrow \infty$, then find u .

The function $u(x, y)$ satisfies Poisson's equation $\nabla^2 u = -4\pi\rho$ in two dimensions, where $\rho = \frac{h}{4\pi}$ and hence the boundary value problem is

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = -h \quad (0 < x < \pi, y > 0) \quad \dots (1)$$

with conditions $u=0$ when $x=0$, $u=1$ when $x=\pi$ $\dots (2)$

and $u=0$ when $y=0$, ($0 < x < \pi$) and $u < M$ ($0 \leq x \leq \pi, y > 0$) $\dots (3)$
where M is some constant.

Using finite Fourier transform, (1) gives

$$\int_0^\pi \frac{\partial^2 u}{\partial x^2} \sin \lambda x \, dx + \int_0^\pi \frac{\partial^2 u}{\partial y^2} \sin \lambda x \, dx = -h \int_0^\pi \sin \lambda x \, dx$$

$$\text{or} \quad \frac{d^2 \tilde{u}}{dy^2} - p^2 \tilde{u} = p(-1)^p - h F_s\{1\}, \quad \dots (4)$$

$$\text{where} \quad \tilde{u} = \int_0^\pi u \sin \lambda x \, dx$$

(under the conditions $u=0$ at $x=0$ and $u=1$ at $x=\pi$).

Finite Fourier transform of (3) gives $\tilde{u}=0$ when $y=0$ and $|\tilde{u}| < M\pi$ $\dots (5)$

$$\therefore \text{Solution of (4) is } \tilde{u} = Ae^{-py} + Be^{py} + \frac{p(-1)^p - h F_s\{1\}}{-p^2}$$

Since $\tilde{u}$ is finite when $y \rightarrow \infty$ $\therefore B=0$.

$$\text{Also } y=0, \tilde{u}=0 \text{ gives } A = \frac{p(-1)^p - h F_s\{1\}}{p^2}$$

$$\text{Hence } \tilde{u} = \frac{h F_s\{1\} - p(-1)^p}{p^2} (1 - e^{py})$$

Applying the inversion formula for finite Fourier sine transform, we find

$$u(x, y) = \frac{2}{\pi} \sum \tilde{u} \sin nx.$$

Problem 14. Solve $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$ for $0 < x < \pi$, $0 < y < \pi$ under the boundary conditions

$$u(x, 0) = x^2, u(x, \pi) = 0, u_x(0, y) = \frac{\partial}{\partial x} u_x(0, y) = 0 = u_x(\pi, y).$$

$$\text{Ans. } u(x, y) = \frac{\pi}{3}(\pi - y) + 4 \sum_{n=1}^{\infty} (-1)^n \frac{\sinh n(\pi - y) \cos nx}{n^2 \sinh n\pi}$$

[B] Solution of two-dimensional Laplace's equation in cylindrical (or Polar) coordinates.

$$\text{The Laplace's equation in this case is } \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0 \quad \dots (1)$$

$$\text{Assume } u(r, \theta) = R(r) \Theta(\theta)$$

$$\text{so that } \frac{\partial^2 u}{\partial r^2} = \Theta \frac{d^2 R}{dr^2}, \frac{\partial u}{\partial r} = \Theta \frac{dR}{dr} \text{ and } \frac{\partial^2 u}{\partial \theta^2} = R \frac{d^2 \Theta}{d\theta^2}.$$

$$\therefore (1) \text{ gives } \Theta \frac{d^2 R}{dr^2} + \frac{1}{r} \Theta \frac{dR}{dr} + \frac{1}{r^2} R \frac{d^2 \Theta}{d\theta^2} = 0 \quad \dots (2)$$

$$\text{or } \frac{1}{R} \left(r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} \right) = -\frac{1}{\Theta} \frac{d^2 \Theta}{d\theta^2} = n^2 \text{ (say)} \quad \dots (3)$$

as variables are separated.

$$\text{Here } -\frac{1}{\Theta} \frac{d^2 \Theta}{d\theta^2} = n^2 \text{ i.e., } \frac{d^2 \Theta}{d\theta^2} + n^2 \Theta = 0 \text{ gives } \Theta = A \cos n\theta + B \sin n\theta \quad \dots (4)$$

$$\text{and } \frac{1}{R} \left(r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} \right) = n^2 \text{ i.e., } r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} - n^2 R = 0$$

being homogeneous, takes the form $\frac{d^2 R}{ds^2} - n^2 R = 0$ on putting $r = e^s$

and then its solution is

$$R = Ce^{ns} + De^{-ns} \text{ i.e., } R = C r^n + D r^{-n} \quad \dots (5)$$

Taking $n=0$ we have from (3)

$$\frac{d^2 \Theta}{d\theta^2} = 0 \text{ giving } \Theta = A_0 \theta + B_0 \quad \dots (6)$$

$$\text{and } r^2 \frac{d^2 R}{dr^2} + r \frac{dR}{dr} = 0 \text{ or } \frac{d^2 R}{ds^2} = 0 \text{ (when } r = e^s) \text{ giving } R = C_0 s + D_0 = C_0 \log r + D_0 \quad \dots (7)$$

The solution of Laplace's equation in cylindrical coordinates when u is independent of z is known as *Circular Harmonics* and n is the degree of the harmonic. Hence the Circular Harmonics of degree zero are given by

$$u_0 = (A_0 + B_0)(C \log r + D) \text{ by (6) and (7)}$$

and those of degree n are given by

$$u_n = (A_n \cos n\theta + B_n \sin n\theta)(C_n r^n + D_n r^{-n}) \text{ by (4) and (5).}$$

The general single-valued solution of (1) for all possible n may be written as

$$u = A_0 \log r + \sum_{n=1}^{\infty} (A_n \cos n\theta + B_n \sin n\theta) (C_n r^n + D_n r^{-n}) + C_0 \quad \dots (8)$$

where A_0, A_n, B_n, C_n, D_n and C_0 all are arbitrary constants.

Problem. 15. For a semi-circular plate of radius a with boundary diameter at 0°C and surface at 100°C , show that the temperature distribution is given by

$$u = \frac{400}{\pi} \sum_{n=1}^{\infty} \frac{r^{2n-1} \sin(2n-1)\theta}{2n-1 a^{2n-1}}$$

$$\text{The boundary value problem. } \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0 \quad \dots (1)$$

Its solution by (8) of § 12.7 [B] is

$$u = A_0 \log r + \sum_{n=1}^{\infty} (A_n \cos n\theta + B_n \sin n\theta) (C_n r^n + D_n r^{-n}) + C_0 \quad \dots (2)$$

But temperature being finite at $r=0$, (2) should not contain terms of $\log r$ and r^{-n} and this will be so if $A_0=0=D_n$.

Moreover at $r=0$, u being zero, we should have $C_0=0$

$$\text{Hence (2) reduces to } u = \sum_{n=1}^{\infty} (A_n \cos n\theta + B_n \sin n\theta) C_n r^n \\ = \sum_{n=1}^{\infty} (a_n \cos n\theta + b_n \sin n\theta) r^n \text{ taking } a_n = A_n C_n \text{ etc.}$$

Now a being the radius of the sphere and assuming $u=u(a)$ at $r=a$, we have

$$u(a) = \sum_{n=1}^{\infty} (a_n \cos n\theta + b_n \sin n\theta) a^n \quad \dots (3)$$

$$\text{where } a_n = \frac{2}{\pi} \int_0^\pi \frac{u(a)}{a^n} \cos n\theta d\theta = \frac{200}{\pi a^n} \int_0^\pi \cos n\theta d\theta = 0 \text{ and} \\ b_n = \frac{2}{\pi} \int_0^\pi \frac{u(a)}{a^n} \sin n\theta d\theta$$

$$\text{i.e., } b_n = \frac{200}{\pi a^n} \int_0^\pi \sin n\theta d\theta = 0 \text{ when } n \text{ is even and } \frac{400}{n\pi a^n} \text{ when } n \text{ is odd.}$$

Hence (3) reduces to

$$u = \frac{400}{\pi} \sum_{n=1}^{\infty} \frac{r^{2n-1} \sin(2n-1)\theta}{2n-1 a^{2n-1}} \text{ for } n \text{ odd.}$$

Problem 16. Determine the steady state temperature at any point of a semi-circular metal plate of radius a whose circumference is maintained to a given temperature of T° whereas the base is kept at zero temperature

$\therefore u=u(r, \theta)$, the boundary value problem is

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0 \quad \dots (1)$$

with conditions (i) $u=0$ when $\theta=0$ for $0 \leq r < a$

(ii) u is finite when $r \rightarrow 0$

(iii) $u=T$ when $r=a$ for $0 < \theta < \pi$

Solution of (1) by (8) of § 12.7 [B] is

$$u = A_0 \log r + \sum_{n=1}^{\infty} (A_n \cos n\theta + B_n \sin n\theta) (C_n r^n + D_n r^{-n}) + C_0 \quad \dots (2)$$

In view of condition (ii), u being finite, (2) must not contain terms of $\log r$ and r^{-n} and this will be so when $A_0=0=D_n$. Thus (2) reduces to

$$u = \sum_{n=1}^{\infty} (a_n \cos n\theta + b_n \sin n\theta) r^n + C_0 \text{ when } a_n = A_n C_n, b_n = B_n C_n \quad \dots (3)$$

In view of condition (i) $C_0=0$ and hence (3) yields

$$u = \sum_{n=1}^{\infty} (a_n \cos n\theta + b_n \sin n\theta) r^n \quad \dots (4)$$

By condition (iii), this gives, $T = \sum_{n=1}^{\infty} (a_n \cos n\theta + b_n \sin n\theta) a^n$ from which we find

$$a_n = \frac{2}{\pi} \int_0^\pi \frac{T}{a^n} \cos n\theta d\theta = \frac{2T}{\pi a^n} \int_0^\pi \cos n\theta d\theta = 0 \text{ and} \\ b_n = \frac{2}{\pi} \int_0^\pi \frac{T}{a^n} \sin n\theta d\theta = \frac{2T}{\pi a^n} \int_0^\pi \sin n\theta d\theta = \frac{2T}{n\pi a^n} (1 - \cos n\pi)$$

Hence (4) gives the required solution as

$$u = \frac{2T}{n\pi a^n} \sum_{n=1}^{\infty} (1 - \cos n\pi) r^n \sin n\theta.$$

Problem 17. A long cylinder is made of two halves, the upper half is at the temperature T_1 and the lower half at the temperature T_2 . Find the distribution of temperature inside the cylinder.

Taking the axis of cylinder along z -axis, there is symmetry along z -axis and hence z -axis has no effect on the distribution of temperature. At the centre where $r=0$, we have $u=\text{finite}$. The boundary value problem is

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0 \quad \dots (1)$$

Its solution is (by (8) of § 12.7[B])

$$u = A_0 \log r + \sum_{n=1}^{\infty} (A_n \cos n\theta + B_n \sin n\theta) (C_n r^n + D_n r^{-n}) + C_0 \quad \dots (2)$$

u being finite, we have to eliminate the terms of $\log r$ and r^{-n} so that $A_0 = 0 = D_n$

$$\therefore (2) \text{ becomes } u = \sum_{n=1}^{\infty} (a_n \cos n\theta + b_n \sin n\theta) + C_0 \quad \dots (3)$$

where $a_n = A_n$, C_n etc.

Suppose that $u = F(\theta)$ at $r = R$ (say), then (3) gives

$$f(\theta) = C_0 + \sum_{n=1}^{\infty} (a_n \cos n\theta + b_n \sin n\theta) R^n \quad \dots (4)$$

$$\text{This gives } C_0 = \frac{1}{2\pi} \int_0^{2\pi} f(\theta) d\theta$$

$$a_n = \frac{1}{R^n \pi} \int_0^{2\pi} f(\theta) \cos n\theta d\theta \text{ and } b_n = \frac{1}{R^n \pi} \int_0^{2\pi} f(\theta) \sin n\theta d\theta$$

But we are given that

$$f(\theta) = T_1 \text{ for } \pi > \theta > 0 \text{ (upper half)}$$

$$\text{and } f(\theta) = T_2 \text{ for } 2\pi > \theta > \pi \text{ (lower half)}$$

$$\therefore C_0 = \frac{1}{2\pi} \left[\int_0^{\pi} T_1 d\theta + \int_{\pi}^{2\pi} T_2 d\theta \right] = \frac{T_1 + T_2}{2}$$

$$a_n = \frac{1}{\pi R^n} \left[\int_0^{\pi} T_1 \cos n\theta d\theta + \int_{\pi}^{2\pi} T_2 \cos n\theta d\theta \right] = 0$$

$$\text{and } b_n = \frac{1}{\pi R^n} \left[\int_0^{\pi} T_1 \sin n\theta d\theta + \int_{\pi}^{2\pi} T_2 \sin n\theta d\theta \right]$$

$$= \frac{1}{\pi R^n} \left[\frac{T_1 + T_2}{n} (1 - \cos n\pi) \right] = \begin{cases} \frac{2}{n\pi R^n} (T_1 - T_2) & \text{for } n \text{ odd} \\ 0, & \text{for } n \text{ even} \end{cases}$$

Hence the solution (3) reduces to

$$u = \frac{T_1 + T_2}{2} + \sum_{n=1}^{\infty} \frac{2}{n\pi R^n} (T_1 - T_2) \sin n\theta \cdot r^n$$

where $n = 1, 3, 5, \dots$

$$= \frac{T_1 + T_2}{2} + \frac{2}{\pi R} (T_1 - T_2) \sin \theta - r + \frac{2}{3\pi R^3} (T_1 - T_2) \sin 3\theta \cdot r^3 + \dots$$

which gives the required distribution of temperature.

Problem 18. If u is a function of r and θ satisfying

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0 \quad \dots (1)$$

within the region of the plane bounded by $r = a$, $r = 0$, $\theta = 0$, $\theta = \frac{\pi}{2}$ and also satisfying the boundary conditions $u = 0$ when $\theta = 0$, $u = 0$ when $\theta = \frac{\pi}{2}$, $u = 0$ when $r = b$ and $u = \theta \left(\frac{\pi}{2} - \theta \right)$ when $r = 0$ then show that

$$u = \frac{16}{\pi} \sum_{n=1}^{\infty} \frac{(r/b)^{4n-2} - (b/r)^{4n-2} \cdot \sin(4n-2)\theta}{(a/b)^{4n-2} - (b/a)^{4n-2} (4n-2)^3}.$$

The solution of (1) with the help of (4) and (5) of §12.7 [B] is $u(r, \theta) = (A \cos m\theta + B \sin m\theta) (C r^m + D r^{-m})$ $\dots (2)$

where we have taken m^2 as constant of separation. Applying the boundary condition $u = 0$, when $\theta = 0$, (2) gives $0 = A (C r^m + D r^{-m})$ i.e. $A = 0$.

$$\therefore (2) \text{ becomes } u(r, \theta) = (C' r^m + D' r^{-m}) \sin m\theta \quad \dots (3)$$

where $BC = C'$ and $BD = D'$.

$$\text{The condition } u = 0 \text{ when } \theta = 0, \text{ gives } 0 = (C' r^m + D' r^{-m}) \sin \frac{m\pi}{2}$$

$$\text{i.e. } \sin \frac{m\pi}{2} = 0 \text{ or } \frac{m\pi}{2} = (2n-1)\pi \text{ giving } m = 4n-2.$$

Also the condition $u = 0$ when $r = b$ gives $0 = (C' r^m + D' r^{-m}) \sin m\theta$ i.e. $C' b^m + D' b^{-m} = 0$ which yields with $m = 4n-2$, $D' = -C' b^{2m} = -C' b^{8n-4}$.

As such (3) reduces to $u = C' [r^{4n-2} - b^{8n-4} r^{-(4n-2)}] \sin(4n-2)\theta$

Considering all possible n , the general solution becomes

$$u = \sum_{n=1}^{\infty} C'_n [r^{4n-2} - b^{8n-4} r^{-(4n-2)}] \sin(4n-2)\theta \quad \dots (4)$$

Applying the condition $u = \theta \left(\frac{\pi}{2} - \theta \right)$ when $r = a$, (4) yields

$$\theta \left(\frac{\pi}{2} - \theta \right) = \sum_{n=1}^{\infty} C'_n [a^{4n-2} - b^{8n-4} a^{-(4n-2)}] \sin(4n-2)\theta$$

$$\text{So that } C'_n \left(a^{4n-2} - \frac{b^{8n-4}}{a^{4n-2}} \right) = \frac{4}{\pi} \int_0^{\pi/2} \theta \left(\frac{\pi}{2} - \theta \right) \sin(4n-2)\theta d\theta = \frac{4}{\pi} \cdot \frac{4}{(4n-2)^3}.$$

$$\text{Giving } C'_n = \frac{16}{\pi(4n-2)^3} \cdot \frac{a^{4n-2}}{a^{8n-4} - b^{8n-4}}.$$

Hence (4) reduces to

$$u = \sum_{n=1}^{\infty} \frac{16}{\pi} \sum_{n=1}^{\infty} \frac{\{r^{4n-2} - b^{8n-4} r^{-(4n-2)}\} a^{4n-2}}{a^{8n-4} - b^{8n-4}} \cdot \frac{\sin(4n-2)\theta}{(4n-2)^3}$$

$$= \frac{16}{\pi} \sum_{n=1}^{\infty} \left(\frac{r}{b} \right)^{4n-2} - \left(\frac{b}{r} \right)^{4n-2} \cdot \frac{\sin(4n-2)\theta}{(4n-2)^3}.$$




Fig. 12.4

12.8. THREE-DIMENSIONAL LAPLACE'S EQUATION

[A] Solution of three dimensional Laplace-equation in Cartesian coordinates.

$$\text{The equation is } \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0 \quad \dots (1)$$

$$\text{Suppose } u = X(x) Y(y) Z(z) \quad \dots (2)$$

$$\text{Then (1) yields, } \frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2} + \frac{1}{Z} \frac{d^2 Z}{dz^2} = 0 \quad \dots (3)$$

This relation being of the form $F_1(X) + F_2(Y) + F_3(Z) = 0$ will be true only if F_1, F_2, F_3 are constant functions since x, y, z and so X, Y, Z are independent functions. We therefore take constants $-n^2, -m^2, +p^2$ such that $p^2 = m^2 + n^2$ and

$$\frac{1}{X} \frac{d^2 X}{dx^2} = -n^2 \text{ i.e. } \frac{d^2 X}{dx^2} + n^2 X = 0 \text{ giving } X = A \cos nx + B \sin nx \quad \dots (4)$$

$$\frac{1}{Y} \frac{d^2 Y}{dy^2} = -m^2 \text{ i.e. } \frac{d^2 Y}{dy^2} + m^2 Y = 0 \text{ giving } Y = C \cos my + D \sin ny \quad \dots (5)$$

$$\frac{1}{Z} \frac{d^2 Z}{dz^2} = p^2 \text{ i.e. } \frac{d^2 Z}{dz^2} - p^2 Z = 0 \text{ giving } Z = E e^{pz} + F e^{-pz} \quad \dots (6)$$

As such the combined solution of (1) is

$$u = (A \cos nx + B \sin nx) (C \cos my + D \sin ny) (E e^{pz} + F e^{-pz}) \quad \dots (7)$$

As an alternative this may be taken as

$$u = (A e^{inx} + B e^{-inx}) (C e^{my} + D e^{-my}) [E \cos pz + F \sin pz] \quad \dots (8)$$

Note. We can easily verify that Laplace's equation $\nabla^2 u = 0$ is satisfied by

$$u = \frac{l}{\sqrt{(x-a)^2 + (y-b)^2 + (z-c)^2}} \quad \dots (9)$$

where l is a constant and (a, b, c) are coordinates of a fixed point.

[B] Solution of three-dimensional Laplace-equation in cylindrical coordinates

$$\text{We have } \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial z^2} = 0 \quad \dots (10)$$

$$\text{Suppose that } u(r, \theta, z) = R(r) \Theta(\theta) Z(z). \quad \dots (11)$$

$$\text{Then (10) yields } \frac{1}{R} \frac{d^2 R}{dr^2} + \frac{r}{R} \frac{dR}{dr} + \frac{1}{r^2 \Theta} \frac{d^2 \Theta}{d\theta^2} + \frac{1}{Z} \frac{d^2 Z}{dz^2} = 0. \quad \dots (12)$$

As variables are separated, we may take

$$\frac{1}{Z} \frac{d^2 Z}{dz^2} = \lambda^2 \text{ and } \frac{1}{\Theta} \frac{d^2 \Theta}{d\theta^2} = -\mu^2 \text{ i.e. } \frac{d^2 Z}{dz^2} - \lambda^2 Z = 0$$

$$\text{and } \frac{d^2 \Theta}{d\theta^2} + \mu^2 \Theta = 0 \quad \dots (13)$$

$$\text{These yield } Z = e^{\pm \lambda z} \text{ and } \Theta = e^{\pm i\mu\theta} \quad \dots (14)$$

or in other words the solutions of (13) are

$$Z = A_1 e^{\lambda z} + B_1 e^{-\lambda z}, \Theta = A_2 \cos \mu\theta + B_2 \sin \mu\theta \quad \dots (15)$$

Also then (12) reduces to

$$\frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + \left(\lambda^2 - \frac{\mu^2}{r^2} \right) R = 0 \quad \dots (16)$$

which is Bessel's equation and takes the form

$$\frac{d^2 R}{dx^2} + \frac{1}{x} \frac{dR}{dx} + \left(1 - \frac{\mu^2}{x^2} \right) R = 0, \text{ on putting } \lambda r = x. \quad \dots (17)$$

Its general solutions are

$$R = A_3 J_\mu(\lambda r) + B_3 J_{-\mu}(\lambda r), \text{ for fractional } \mu \quad \dots (18)$$

$$\text{and } R = A_3 J_\mu(\lambda r) + B_3 Y_\mu(\lambda r), \text{ for integral } \mu \quad \dots (19)$$

As such the solutions for (10), with the help of (15), (18) and (19) are

$$u(r, \theta, z) = (A_1 e^{\lambda z} + B_1 e^{-\lambda z}) (A_2 \cos \mu\theta + B_2 \sin \mu\theta) [A_3 J_\mu(\lambda r) + B_3 J_{-\mu}(\lambda r)] \quad \dots (20)$$

$$\text{and } u(r, \theta, z) = (A_1 e^{\lambda z} + B_1 e^{-\lambda z}) (A_2 \cos \mu\theta + B_2 \sin \mu\theta) [A_3 J_\mu(\lambda r) + B_3 Y_\mu(\lambda r)] \quad \dots (21)$$

Note 1. The general solution of (16) may be written as

$$R = A_{\lambda\mu} J_\mu(\lambda r) + B_{\lambda\mu} Y_\mu(\lambda r) \quad \dots (22)$$

where $A_{\lambda\mu}$ and $B_{\lambda\mu}$ are constants.

Since $Y_\mu(\lambda r) \rightarrow \infty$ as $r \rightarrow 0$, therefore in a physical problem if μ is finite along the line $r=0$, then we must have $B_{\lambda\mu}=0$ and hence the solution of (10) may be written as

$$u = \sum_{\lambda, \mu} A_{\lambda\mu} J_\mu(\lambda r) e^{\pm \lambda z \pm i\mu\theta} \quad \dots (23)$$

Note 2. Trying the superposition, the general solution of (10) may be written as

$$u(r, \theta, z) = \sum_{\mu=0}^{\infty} J_\mu(\lambda r) [e^{\lambda z} (A_\mu \cos \mu\theta + B_\mu \sin \mu\theta) + e^{-\lambda z} (C_\mu \cos \mu\theta + D_\mu \sin \mu\theta)] \quad \dots (24)$$

Note 3. In a problem if there is symmetry about z -axis, then we may take $\mu=0$ and the solution will be

$$u(r, \theta, z) = \sum_{\lambda} A_{\lambda} J_0(\lambda r) e^{\pm \lambda z} \quad \dots (25)$$

Note 4. If in a problem of symmetry about z -axis, $u \rightarrow 0$ as $r \rightarrow 0$ and $z \rightarrow \infty$, then the solution is of the form $u(r, \theta, z) = \sum_{\lambda} A_{\lambda} J_0(\lambda r) e^{-\lambda z}$

$\dots (26)$

[C] **Solution of three-dimensional Laplace-equation in spherical Polar coordinates**

We have
$$\frac{\partial^2 u}{\partial r^2} + \frac{2}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial u}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 u}{\partial \phi^2} = 0 \quad \dots (27)$$

or equivalently,
$$r^2 \frac{\partial^2 u}{\partial r^2} + 2r \frac{\partial u}{\partial r} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial u}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2 u}{\partial \phi^2} = 0 \quad \dots (28)$$

Suppose $u(r, \theta, \phi) = R(r) \Theta(\theta) \Phi(\phi) \quad \dots (29)$

Then (27) and (28) yield on dividing by $R\Theta\Phi$

$$\frac{1}{R} \frac{d^2 R}{dr^2} + \frac{2}{rR} \frac{dR}{dr} + \frac{1}{r^2 \Theta} \frac{d^2 \Theta}{d\theta^2} + \frac{\cot \theta}{r^2 \Theta} \frac{d\Theta}{d\theta} + \frac{1}{r^2 \sin^2 \theta} \cdot \frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2} = 0 \quad \dots (30)$$

and
$$\left[\frac{r^2}{R} \frac{d^2 R}{dr^2} + \frac{2r}{R} \frac{dR}{dr} + \frac{1}{\Theta \sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) \right] \sin^2 \theta = - \frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2} = \lambda^2 \text{ (say)} \quad \dots (31)$$

Considering (31), $-\frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2} = \lambda^2$ i.e. $\frac{d^2 \Phi}{d\phi^2} + A^2 \Phi = 0$ gives

$$\Phi = C e^{\pm i\lambda \phi} \quad \dots (32)$$

and
$$\frac{1}{R} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) = - \frac{1}{\Theta \sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) + \frac{\lambda^2}{\sin^2 \theta} = n(n+1) \text{ (say), gives } \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) - n(n+1)R = 0 \quad \dots (33)$$

and
$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) + \left[n(n+1) - \frac{\lambda^2}{\sin^2 \theta} \right] \Theta = 0 \quad \dots (34)$$

Considering (30), if we write $\frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2} = -\lambda^2$ and $\frac{r^2}{R} \frac{d^2 R}{dr^2} + \frac{2r}{R} \frac{dR}{dr} = n(n+1)$, then we have

$\Phi = C e^{\pm i\lambda \phi}$ which is (32); $r^2 \frac{dR}{dr^2} + \frac{2r}{dr} R - n(n+1)R = 0$ which is

(33) and
$$\frac{d^2 \Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} + \left\{ n(n+1) - \frac{\lambda^2}{\sin^2 \theta} \right\} \Theta = 0 \text{ which is (34).}$$

Now the equation (33) being homogeneous if we put $r=e^s$, then it reduces to $\{D(D-1)+2D-n(n+1)\}R=0$ where $D \equiv \frac{d}{ds}$ or $(D-n)(D+n+1)R=0$ giving $R=Ae^{ns}+Be^{-(n+1)s}=Ar^n+Br^{-n-1} \quad \dots (35)$

Again if we put $\cos \theta = \mu$ in (3) then since

$$\frac{d\Theta}{d\theta} = \frac{d\Theta}{d\mu} \frac{d\mu}{d\theta} = -\sin \theta \frac{d\Theta}{d\mu} \text{ i.e. } \frac{1}{\sin \theta} \frac{d}{d\theta} \equiv -\frac{d}{d\mu},$$

we have
$$\frac{d}{d\mu} \left\{ (1-\mu^2) \frac{d\Theta}{d\mu} \right\} + \left\{ n(n+1) - \frac{\lambda^2}{1-\mu^2} \right\} \Theta = 0$$

i.e. $(1-\mu^2) \frac{d^2 \Theta}{d\mu^2} - 2\mu \frac{d\Theta}{d\mu} + \left\{ n(n+1) - \frac{\lambda^2}{1-\mu^2} \right\} \Theta = 0 \quad \dots (36)$

which is Legendre's associated equation and hence has the solution

$$\Theta = A P_n^\lambda(\mu) + B Q_n^\lambda(\mu) = A P_n^\lambda(\cos \theta) + B Q_n^\lambda(\cos \theta) \quad \dots (37)$$

In other words if we take $\Theta = \Theta(\cos \theta)$ from associated Legendre equation, then the solution of (27) is of the form $(Ar^n + Br^{-n-1})$

$$\Theta(\cos \theta) e^{\pm i\lambda \phi}$$

So that summing over for all n and trying superposition, the general solution of (27) may be written as

$$u(r, \theta, \phi) = \sum_{n=0}^{\infty} \left(A_n r^n + \frac{B_n}{r^{n+1}} \right) \Theta(\cos \theta) e^{\pm i\lambda \phi} \quad \dots (38)$$

Note. If $\lambda=0$, then (36) reduces to $(1-\mu^2) \frac{d^2 \Theta}{d\mu^2} - 2\mu \frac{d\Theta}{d\mu} + n(n+1)\Theta = 0$

which is Legendre's equation and hence we have for integral n ,

$$\Theta = P_n(\mu) = P_n(\cos \theta) = \frac{1}{2^n n!} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n$$

and also $\Theta = Q_n(\mu) = Q_n(\cos \theta) = \frac{1}{2} P_n(\mu) \log \frac{\mu+1}{\mu-1} - \sum_{j=0}^p \frac{2n-4j-1}{(2j+1)(n-j)} P_{n-2j-1}(\mu)$

where $p = \frac{1}{2}(n-1)$ or $\frac{1}{2}n-1$ according as n is odd or even:

Thus $\Theta = C_n P_n(\mu) + D_n Q_n(\mu)$ so that $u = \sum_{n=1}^{\infty} \left(A_n r^n + \frac{B_n}{r^{n+1}} \right)$

$$\{ C_n P_n(\cos \theta) + D_n Q_n(\cos \theta) \} e^{-i\lambda \phi} \quad \dots (39)$$

In case $D_n=0$ under specified boundary conditions, then

$$\Theta = C_n P_n(\cos \theta)$$

Hence the solution is $u = \sum_{n=0}^{\infty} \left(A_n r^n + \frac{B_n}{r^{n+1}} \right) P_n(\cos \theta) \cdot e^{-i\lambda\phi} \dots (40)$

Note 2. If there is axial symmetry about z-axis, then u depends only r and θ and so (27) reduces to

$$\frac{\partial^2 u}{\partial r^2} + \frac{2}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial u}{\partial \theta} = 0 \dots (41)$$

Its solution by putting $\phi=0$ in (40), is

$$u(r, \theta) = \sum_{n=0}^{\infty} \left(A_n r^n + \frac{B_n}{r^{n+1}} \right) \{ C_n P_n(\cos \theta) + D_n Q_n(\cos \theta) \} \dots (42)$$

In case $D_n=0$ under specified conditions, then

$$u(r, \theta) = \sum_{n=0}^{\infty} \left(A_n r^n + \frac{B_n}{r^{n+1}} \right) P_n(\cos \theta) \dots (43)$$

Problem 19. If the surface S of a sphere of radius a is kept at a fixed distribution of electric potential $u=F(\theta)$, then find the potential u at all points in space which is assumed to be free of further charge.

In this case $\frac{\partial^2 u}{\partial \phi^2}$ being zero, we have the constant of separation i.e.

$\lambda=0$ and hence equation (36) of § 12.8[C], reduces to

$$(1-\mu^2) \frac{d^2 \Theta}{d\mu^2} - 2\mu \frac{d\Theta}{d\mu} + n(n+1)\Theta = 0 \dots (1)$$

Being Legendre's, equation the Legendre polynomial $P_n(u) = P_n(\cos \theta)$ is the solution of (1) i.e. we have

$$u = \sum_{n=0}^{\infty} \left(A_n r^n + \frac{B_n}{r^{n+1}} \right) P_n(\cos \theta) \dots (2)$$

Now to determine the potential u , we consider the problem in two cases:

Case I. Outside the sphere. Since the potential at infinity vanishes i.e. $\lim_{r \rightarrow \infty} u = 0$, the boundary condition requires that any positive power of r should not be present in the solution (2), thereby giving $A_n=0$ so that (2) reduces to

$$u = \sum_{n=0}^{\infty} \frac{B_n}{r^{n+1}} P_n(\cos \theta) \dots (3)$$

Assuming that $u=F(\theta)$ when $r=a$, (3) gives

$$F(\theta) = f(\cos \theta) \text{ (say)} = \sum_{n=0}^{\infty} \frac{B_n}{a^{n+1}} P_n(\cos \theta) \dots (4)$$

If we replace $\cos \theta$ by u in (4), we get $a^{n+1} f(u) = \sum_{n=0}^{\infty} B_n P_n(u)$

$$\text{so that } a^{n+1} \int_{-1}^1 f(u) P_n(u) du = \int_{-1}^1 \sum_n B_n P_n(u) \cdot P_n(u) du = B_n \int_{-1}^1$$

$$P_n^2(u) du = \frac{2B_n}{2n+1} \text{ giving } B_n = \frac{2n+1}{2} a^{n+1} \int_0^\pi F(\theta) P_n(\cos \theta) \sin \theta d\theta \text{ on setting } u = \cos \theta \dots (5)$$

Putting this value of B_n in (3), we get the required potential outside the sphere i.e.

$$u = \sum_{n=0}^{\infty} \frac{2n+1}{2} \left(\frac{a}{r} \right)^{n+1} P_n(\cos \theta) \int_0^\pi F(\theta) P_n(\cos \theta) \sin \theta d\theta \dots (6)$$

Case II. Inside the sphere. Since the potential inside the sphere can not be infinite, therefore the general solution must not contain any negative power of r , thereby giving $B_n=0$, so that (2) reduces to

$$u = \sum_{n=0}^{\infty} A_n r^n P_n(\cos \theta), r < a \dots (7)$$

when $r=a$, $u=F(\theta)$, therefore (7) gives $F(\theta) = \sum_{n=0}^{\infty} A_n a^n P_n(\cos \theta)$ from which we have as in case I,

$$A_n = \frac{2n+1}{a^n} \int_0^\pi F(\theta) P_n(\cos \theta) \sin \theta d\theta \dots (8)$$

Substituting (8) in (7) we get the required potential inside the sphere.

Problem 20. Find a solution of the equation

$$\frac{\partial}{\partial r} \left(r^2 \frac{\partial \phi}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \phi}{\partial \theta} \right) = 0 \dots (1)$$

in the form $\phi = f(r) \cos \theta$, given that (i) $-\frac{\partial \phi}{\partial r} = u \cos \theta$ when $r=a$ and (ii) $-\frac{\partial \phi}{\partial r} = 0$ when $r=\infty$.

We have $\phi = f(r) \cos \theta$

$$\therefore \frac{\partial \phi}{\partial r} = f'(r) \cos \theta, \frac{\partial \phi}{\partial \theta} = -f(r) \sin \theta.$$

Their substitution in (1) gives

$$\frac{\partial}{\partial r} \{ r^2 f'(r) \cos \theta \} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \{ -\sin \theta f(r) \sin \theta \} = 0$$

$$\text{i.e., } r^2 f''(r) + 2r f'(r) - 2f(r) = 0 \dots (2)$$

This equation being homogeneous, can be reduced to the form

$$\{ D(D-1) + 2D - 2 \} f(r) = 0 \text{ by putting } r = e^s \text{ and } D \equiv \frac{d}{ds}$$

$$\text{or } (D^2 + D - 2) f(r) = 0 \text{ or } (D-1)(D+2) f(r) = 0$$

$$\therefore f(r) = A e^s + B e^{-2s} = Ar + \frac{B}{r^2} \text{ so that } \phi = Ar \cos \theta + \frac{B}{r^2} \cos \theta \quad \dots (3)$$

$$(3) \text{ gives } \frac{\partial \phi}{\partial r} = A \cos \theta - \frac{2B}{r^3} \cos \theta.$$

$$\text{Applying the condition (i) } -u \cos \theta = A \cos \theta - \frac{2B}{a^3} \cos \theta \quad \dots (4)$$

$$\text{and applying (ii) } 0 = A \cos \theta \text{ i.e., } A = 0 \text{ and then from (4) } B = \frac{a^3 u}{2}$$

$$\text{Hence (3) yields } \phi = \frac{1}{2} \frac{a^3 u}{r^2} \cos \theta.$$

Problem 21. Find the permanent temperature within a solid sphere of radius unit when one half of the surface of the sphere is kept at constant temperature 0°C and the other half of the surface at 1°C .

(Rohilkhand, 1977)




Fig. 12.5

The distribution being symmetrical about z-axis, we have

$$\begin{aligned} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \phi}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \phi}{\partial \theta} \right) &= 0 \\ \text{i.e., } r^2 \frac{\partial^2 \phi}{\partial r^2} + 2r \frac{\partial \phi}{\partial r} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \phi}{\partial \theta} \right) &= 0 \quad \dots (1) \end{aligned}$$

with boundary conditions

$$(i) \phi = 1 \text{ for } 0 < \theta < \frac{\pi}{2} \text{ and } (ii) \phi = 0 \text{ for } \frac{\pi}{2} < \theta < \pi$$

under the consideration of distribution for upper half of the sphere i.e. for $0 < \theta < \pi$.

Assuming that $\phi = R(r) \Theta(\theta)$, (1) yields on dividing throughout by $R\Theta$,

$$\frac{r^2}{R} \frac{d^2 R}{dr^2} + \frac{2r}{R} \frac{dR}{dr} = -\frac{1}{\Theta} \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) = 0 \quad \dots (2)$$

or separating the variables and taking λ^2 as constant of separation,

$$\frac{r^2}{R} \frac{d^2 R}{dr^2} + \frac{2r}{R} \frac{dR}{dr} + \frac{1}{\Theta} \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) = \lambda^2 \text{ (say)}$$

$$\text{so that } r^2 \frac{d^2 R}{dr^2} + 2r \frac{dR}{dr} - \lambda^2 R = 0 \quad \dots (3)$$

$$\text{and } \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) + \lambda^2 \Theta = 0 \quad \dots (4)$$

Taking $\lambda^2 = n(n+1)$, (3) yields $r^2 \frac{d^2 R}{dr^2} + 2r \frac{dR}{dr} - n(n+1) R = 0$ which being a homogeneous equation can be solved by substitution $R = r^n$, to give the solution $R = Ar^n + \frac{B}{r^{n+1}}$.

Also taking $\cos \theta = \mu$, (4) yields, $\frac{\partial}{\partial \mu} \left\{ (1-\mu^2) \frac{d\Theta}{d\mu} \right\} + n(n+1) \Theta = 0$ which is Legendre's equation and hence $P_n(\mu)$ is a solution of it i.e., $\Theta = P_n(\mu) = P_n(\cos \theta)$

Combining the two solutions we have for all n

$$\phi = \sum_{n=0}^{\infty} \left(A_n r^n + \frac{B_n}{r^{n+1}} \right) P_n(\cos \theta) \quad \dots (5)$$

Now the temperature at the centre being finite it is required that $B_n = 0$

$$\therefore (5) \text{ reduces to } \phi = \sum_{n=0}^{\infty} A_n r^n P_n(\cos \theta) = \sum_{n=0}^{\infty} A_n r^n P_n(\mu) \quad \dots (6)$$

But by orthogonal properties of Legendre's polynomials, we have

$$\int_{-1}^1 P_n^2(\mu) d\mu = \frac{2}{2n+1}$$

$$\text{Also } r=1 \text{ gives } \phi = \sum A_n P_n(\mu) \quad \dots (7)$$

Multiplying (7) by $P_n(\mu)$ and integrating with regard to μ from -1 to 1, we have

$$\begin{aligned} A_n &= \frac{2n+1}{2} \int_{-1}^1 \phi P_n(\mu) d\mu = \frac{2n+1}{2} \int_0^\pi \phi P_n(\cos \theta) \sin \theta d\theta \\ &\quad \text{when } \mu = \cos \theta \\ &= \frac{2n+1}{2} \left[\int_0^{\pi/2} \phi P_n(\cos \theta) \sin \theta d\theta + \int_{\pi/2}^\pi \phi P_n(\cos \theta) \sin \theta d\theta \right] \end{aligned}$$

$$= \frac{2n+1}{2} \int_0^{\pi/2} P_n(\cos \theta) \sin \theta d\theta$$

$$\because \phi = 1 \text{ for } 0 < \theta < \pi/2 \text{ and } \phi = 0 \text{ for } \pi/2 < \theta < \pi.$$

$$= \frac{2n+1}{2} \int_0^{\pi/2} P_n(\mu) d\mu \text{ giving } A_0 = \frac{1}{2} \int_0^{\pi/2} 1 \cdot d\mu = \frac{1}{2}$$

$$\text{and } A_1 = \frac{3}{2} \int_0^{\pi/2} P_1(\mu) d\mu = \frac{3}{2} \int_0^{\pi/2} \mu d\mu = \frac{3}{4}$$

$$A_2 = \frac{5}{2} \int_0^{\pi/2} P_2(\mu) d\mu = \frac{5}{2} \int_0^{\pi/2} \frac{3\mu^2-1}{2} d\mu = \frac{5}{4} \left[\mu^3 - \mu \right]_0^{\pi/2} = 0$$

$$\begin{aligned} A_3 &= \frac{7}{2} \int_0^{\pi/2} P_3(\mu) d\mu = \frac{7}{4} \int_0^{\pi/2} (5\mu^3 - \mu) d\mu \\ &= \frac{7}{4} \left[\frac{5}{4} \mu^4 - \frac{3}{2} \mu^2 \right]_0^{\pi/2} = -\frac{7}{16} \text{ etc.} \end{aligned}$$

Hence $\phi = \frac{1}{2} + \frac{3}{4} \cdot r P_1(\cos \theta) - \frac{7}{16} r^3 P_3(\cos \theta) + \dots$

12.9. GENERAL PROPERTIES OF HARMONIC FUNCTIONS

(Agra, 1977)

We know that functions satisfying Laplace's differential equation are said to be the *Harmonic functions*. Now to discuss general properties of such functions, let us consider a vector point function $\mathbf{A}$ and a scalar point function u satisfying Laplace's equation i.e., $\nabla^2 u = 0$

such that $\mathbf{A} = \nabla u$... (1)

$\therefore \nabla \cdot \mathbf{A} = \nabla \cdot (\nabla u) = \nabla^2 u = 0$... (2)

But Gauss' divergence theorem gives $\iint_S \mathbf{A} \cdot d\mathbf{s} = \iiint_V (\nabla \cdot \mathbf{A}) du$... (3)

which with the help of (3) yields $\iint_S \mathbf{A} \cdot d\mathbf{s} = \iint_S (\nabla u) \cdot d\mathbf{s} = 0$... (4)

If we take curl of both sides of (2), we get $\nabla \times \mathbf{A} = \nabla \times \nabla u = 0$... (5)

But Stoke's theorem for a vector field $\mathbf{A}$ is

$\oint_C \mathbf{A} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{A}) \cdot d\mathbf{s} = 0$... (6)

where integral being taken over the closed curve C bounding the open surface S .

(7) with the help of (4) reduces to $\oint_C (\nabla u) \cdot d\mathbf{r} = 0$... (7)

From (5) and (8) certain important properties of harmonic functions can be deduced.

Applying Green's theorem i.e.,

$\iiint_V (v \nabla^2 w - w \nabla^2 v) du = \iint_S (v \nabla w - w \nabla v) \cdot d\mathbf{s}$... (8)

we can easily exhibit that if $\nabla^2 u = 0$ in a region bounded by a sphere of radius r then the value of u say u_0 at the centre of the sphere is given by

$u_0 = \frac{1}{4\pi r^2} \iint_S u ds$... (9)

where integral is taken over the surface of the sphere.

These results may be categorically stated as:

(i) From (10), the average value of a harmonic function on the

surface of a sphere in which it has no singularity i.e., the points where the function becomes infinite, is equal to its value at the centre of the sphere.

(ii) From (5), it follows that a harmonic function having no singularity in a given region cannot have a maximum or minimum value in the region.

(iii) From (ii) we conclude that a harmonic function without singularity within a region and being constant everywhere on the bounding surface of the region, has the same constant value everywhere inside the region.

(iv) Two harmonic functions having identical values on a closed contour and having no singularity within the contour, are identical throughout the region bounded by the contour.

(v) From Green's theorem it follows that if the normal derivative of a harmonic function is zero on a closed surface within which there is no singularity, the function is constant.

(vi) It follows from (v) that if two harmonic functions have the same normal derivative on a closed surface within which there are no singularities, they differ at most by an additive constant.

12.10. THE WAVE EQUATIONS

(A) Derivation of one-dimensional wave equation

Consider a flexible string of length l tightly stretched between two points $x=0$ and $x=l$ on x -axis, with its ends at these ends. If the string is set into small transverse vibration, the displacement say $u(x, t)$ from the x -axis of any point x of the string at any time t is given by

$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$ where $c^2 = \frac{T}{\rho}$ T being tension and ρ the linear density.

The equation $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$... (1)

is known as *one-dimensional wave equation*.

Let the string (assumed to be perfectly flexible) of length l tightly stretched between the points $x=0$ and $x=l$ on x -axis be distorted and then at a certain instant of time say $t=0$, it is released and allowed to vibrate. To determine its deflection (displacement from x -axis) at any point x at any time t , let us take the following assumptions:

- The string is uniform i.e. its mass m per unit length is constant.
- The string is perfectly elastic and so offers no resistance to any bending.




Fig. 12.6

(iii) The tension T is so large that the action of gravitational force on the string is negligible.

(iv) The motion of the string is a small transverse vibration in vertical plane i.e. each particle of the string moves strictly in the vertical plane so that the deflection and slope (gradient) at any point of the string are very small in absolute value.

Consider the motion of an element PQ of length δs of the string. The string being perfectly elastic the tensions T_1 at P and T_2 at Q are tangential to the curve of the string. Let T_1 and T_2 make angles α and β respectively with the horizontal.

There being no motion in the horizontal direction, we have

$$T_1 \cos \alpha = T_2 \cos \beta = T \text{ (say) = constant} \quad \dots (1)$$

Mass of the element PQ is $\rho \delta s$. By Newton's second law of motion we therefore have

$$T_2 \sin \beta - T_1 \sin \alpha = (\rho \delta s) \cdot \frac{\partial^2 u}{\partial t^2} \quad \dots (2)$$

$$\frac{\partial^2 u}{\partial t^2} \text{ being upward acceleration of } PQ$$

$$\text{Using (1), (2) yields } \frac{T_2 \sin \beta}{T_2 \cos \beta} - \frac{T_1 \sin \alpha}{T_1 \cos \alpha} = \frac{\rho \delta s}{T} \cdot \frac{\partial^2 u}{\partial t^2}$$

$$\text{i.e. } \tan \beta - \tan \alpha = \frac{\rho \delta s}{T} \cdot \frac{\partial^2 u}{\partial t^2} \quad \dots (3)$$

Replacing δs by δx since the gradient of the curve is very small, (4)

$$\text{gives } \left(\frac{\partial u}{\partial x} \right)_{x+\delta x} - \left(\frac{\partial u}{\partial x} \right)_x = \frac{\rho \delta x}{T} \cdot \frac{\partial^2 u}{\partial t^2} \quad \dots (5)$$

since $\tan \alpha$ and $\tan \beta$ are slopes at x and $x + \delta x$ respectively,

$$\left(\frac{\partial u}{\partial x} \right)_{x+\delta x} - \left(\frac{\partial u}{\partial x} \right)_x = \frac{\rho}{T} \cdot \frac{\partial^2 u}{\partial t^2} \cdot \delta x$$

or

$$\text{i.e. } \frac{u_x(x+\delta x, t) - u_x(x, t)}{\delta x} = \frac{\rho}{T} \cdot \frac{\partial^2 u}{\partial t^2}$$

Proceeding to the limit as $\delta x \rightarrow 0$, we get

$$\frac{\partial^2 u}{\partial x^2} = \frac{\rho}{T} \cdot \frac{\partial^2 u}{\partial t^2} = \frac{1}{c^2} \cdot \frac{\partial^2 u}{\partial t^2} \text{ where } \frac{1}{c^2} = \frac{\rho}{T}$$

Note 1 $c^2 = \frac{T}{\rho}$ reveals that the constant $\frac{T}{\rho}$ is positive.

Note 2. Since u is dependent of x and t both, therefore we have used the partial derivatives $\frac{\partial^2 u}{\partial x^2}$ and $\frac{\partial^2 u}{\partial t^2}$.

Note 3. If a force $F(x, t)$ per unit of mass acts in the u -direction along the string, in addition to the tension of the string, then

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} + F.$$

[B] Derivation of two-dimensional wave equation

In case of a rectangular membrane, the two dimensional wave equation is

DIFFUSION, WAVE AND LAPLACE'S EQUATIONS

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) \quad \dots (6)$$

Consider the motion of a stretched membrane supposed to be stretched and fixed along its entire boundary in the x - y plane. Let us take the following assumptions:

(i) The membrane is homogeneous i.e. its mass (say) ρ per unit area is constant.

(ii) The membrane is perfectly flexible and so thin that it offers no resistance to any bending.

(iii) The tension T per unit length caused by the stretching of the membrane is invariant during the motion and retains the same value at each of its points and in all the directions.




Fig. 12.7

(iv) The deflection $u(x, y, t)$ of the membrane during the motion is negligible as compared to the size of the membrane. Also all the angles of inclination are small.

Consider the motion of an element $ABCD$ of the membrane. Let its area be $\delta x \delta y$. T being the tension per unit length, the force acting on the edges are $T \delta x$ and $T \delta y$ approximately. Also the membrane being perfectly flexible, the tensions $T \delta x$ and $T \delta y$ are tangential to the membrane. Let α, β be the inclinations of these tensions with the horizontal. Then the horizontal components of the forces at one pair of opposite edges are $T \delta y \cos \alpha$ and $T \delta y \cos \beta$. When α and β are very small $\cos \alpha \rightarrow 1$ and $\cos \beta \rightarrow 1$ so that $T \delta y \cos \alpha \rightarrow T \delta y$ and $T \delta y \cos \beta \rightarrow T \delta y$ i.e. the horizontal components of the forces at opposite edges are nearly equal and hence the motion of the particles of the

membrane in horizontal direction is negligibly small. As such we assume that every particle of the membrane moves vertically.

$$\begin{aligned} \text{The resultant vertical force} &= T\delta y \sin \beta - T\delta y \sin \alpha \\ &= T\delta y (\tan \beta - \tan \alpha) \end{aligned}$$

$$\therefore \alpha, \beta \text{ being small } \sin \alpha = \alpha = \tan \alpha \text{ and } \sin \beta = \beta = \tan \beta$$

$$= T\delta y [u_x(x+\delta x, y_1) - u_x(x, y_2)] \quad \dots (7)$$

where u_x denotes the partial derivative of u w.r.t. x and y_1, y_2 are the values of y between y and $y+\delta y$.

Similarly, the resultant vertical force acting on the other two edges

$$= T\delta x [u_y(x_1, y+\delta y) - u_y(x_2, y)] \quad \dots (8)$$

where u_y denotes the partial derivative of u w.r.t. y and x_1, x_2 are the values of x between x and $x+\delta x$.

By Newton's second law of motion, we have

$$\begin{aligned} \text{Total vertical force on the element} &= \rho \delta x \delta y \frac{\partial^2 u}{\partial t^2} \\ \text{i.e. } T\delta y [u_x(x+\delta x, y_1) - u_x(x, y_2)] &+ T\delta x [u_y(x_1, y+\delta y) - u_y(x_2, y)] \\ &= \rho \delta x \delta y \frac{\partial^2 u}{\partial t^2} \end{aligned}$$

where $\frac{\partial^2 u}{\partial t^2}$ is the acceleration of the element.

$$\begin{aligned} \text{Thus } \frac{\partial^2 u}{\partial t^2} &= \frac{T}{\rho} \left[\frac{u_x(x+\delta x, y_1) - u_x(x, y_2)}{\delta x} \right. \\ &\quad \left. + \frac{T}{\rho} \left[\frac{u_y(x_1, y+\delta y) - u_y(x_2, y)}{\delta y} \right] \right] \end{aligned}$$

Proceeding to the limit as $\delta x \rightarrow 0$ and $\delta y \rightarrow 0$, we have

$$\frac{\partial^2 u}{\partial t^2} = \frac{T}{\rho} [u_{xx} + u_{yy}] = c^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) = c^2 \nabla^2 u \quad \dots (9)$$

$$\text{where } \nabla^2 \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$$

Note 1. If $u = v(x, y) e^{i\omega t}$, (9) yields $\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} + k^2 v = 0 \quad \dots (10)$

$$\text{where } k^2 = \left(\frac{\omega}{c} \right)^2$$

Note 2. The three-dimensional wave equation is

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} \right) = c^2 \nabla^2 u \quad \dots (11)$$

$$\text{where } \nabla^2 \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$$

12.11. GREEN'S FUNCTIONS FOR THE WAVE EQUATION

The wave equation is

$$\nabla^2 \psi \equiv \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \psi = \frac{1}{c^2} \frac{\partial^2 \psi}{\partial t^2} \quad \dots (1)$$

Also written as,

$$\square^2 \psi \equiv \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \psi = 0 \quad \dots (2)$$

If its solution be of the form

$$\psi(x, y, z, t) = \Psi(x, y, z) e^{\pm i c \lambda t} \quad \dots (3)$$

$$\text{Then (1) gives, } \nabla^2 \Psi + \lambda^2 \Psi = 0 \quad \dots (4)$$

which is known as *Space form of the wave equation* or *Helmholtz's equation*.

Taking $\mathbf{r} = x \mathbf{i} + y \mathbf{j} + z \mathbf{k}$ as the position vector of a point (x, y, z) and $\mathbf{r}' = x' \mathbf{i} + y' \mathbf{j} + z' \mathbf{k}$ as the position vector of an isolated point (x', y', z') , the *Green's function* $G(\mathbf{r}, \mathbf{r}')$ is defined as

$$G(\mathbf{r}, \mathbf{r}') = H(\mathbf{r}, \mathbf{r}') + \frac{1}{|\mathbf{r}' - \mathbf{r}|} \quad \dots (5)$$

$$\text{where } H(\mathbf{r}, \mathbf{r}') \text{ satisfies } \left(\frac{\partial^2}{\partial x'^2} + \frac{\partial^2}{\partial y'^2} + \frac{\partial^2}{\partial z'^2} \right) H(\mathbf{r}, \mathbf{r}') = 0 \quad \dots (6)$$

$$\begin{aligned} \text{Using Green's formula i.e. } \psi(\mathbf{r}) &= \frac{1}{4\pi} \int_S \left\{ \frac{1}{|\mathbf{r}' - \mathbf{r}|} \frac{\partial \psi(\mathbf{r}')}{\partial n} \right. \\ &\quad \left. - \psi(\mathbf{r}') \frac{\partial}{\partial n} \left(\frac{1}{|\mathbf{r}' - \mathbf{r}|} \right) \right\} dS' \quad \dots (7) \end{aligned}$$

it may be shown that

$$\psi(\mathbf{r}) = \frac{1}{4\pi} \int_S \left\{ G(\mathbf{r}, \mathbf{r}') \frac{\partial \psi(\mathbf{r}')}{\partial n} - \psi(\mathbf{r}') \frac{\partial G(\mathbf{r}, \mathbf{r}')}{\partial n} \right\} dS' \quad \dots (8)$$

where $\mathbf{n}$ is the unit outward drawn normal to the surface S .

Now we claim that the solution of space form of the wave equation under certain boundary conditions can be made to depend on the determination of the appropriate Green's function. Let us assume that $G(\mathbf{r}, \mathbf{r}')$ satisfies the equation

$$\left(\frac{\partial^2}{\partial x'^2} + \frac{\partial^2}{\partial y'^2} + \frac{\partial^2}{\partial z'^2} \right) G(\mathbf{r}, \mathbf{r}') + \lambda^2 G(\mathbf{r}, \mathbf{r}') = 0 \quad \dots (9)$$

under the assumption that $G(\mathbf{r}, \mathbf{r}')$ is finite and continuous w.r.t. either their variables x, y, z or x', y', z' for the points $\mathbf{r}, \mathbf{r}'$ belonging to a region V bounded by a closed surface S except in $\mathbf{r}$ -neighbourhood where there is a singularity of the type

$$\frac{e^{-i\lambda |\mathbf{r} - \mathbf{r}'|}}{|\mathbf{r} - \mathbf{r}'|} \text{ as } |\mathbf{r}' - \mathbf{r}| \rightarrow 0 \quad \dots (10)$$

Now $\Psi(\mathbf{r})$ being the solution of (4) and its partial derivatives of the first and second orders being continuous within the volume V on the closed surface S we have

$$\frac{1}{4\pi} \int_S \frac{e^{i\lambda|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|} \frac{\partial \Psi(\mathbf{r}')}{\partial n} - \Psi(\mathbf{r}') \frac{\partial}{\partial n} e^{i\lambda|\mathbf{r}-\mathbf{r}'|} \cdot \frac{1}{|\mathbf{r}-\mathbf{r}'|} dS' = \begin{cases} \Psi(\mathbf{r}), & \text{if } \mathbf{r} \text{ lies inside } V \\ 0, & \text{if } \mathbf{r}' \text{ does not lie inside } V \end{cases} \quad \dots (11)$$

Using (10), we therefore have

$$\Psi(\mathbf{r}) = \frac{1}{4\pi} \int_S \left\{ G(\mathbf{r}, \mathbf{r}') \frac{\partial \Psi(\mathbf{r}')}{\partial n} - \Psi(\mathbf{r}') \frac{\partial G(\mathbf{r}, \mathbf{r}')}{\partial n} \right\} dS' \quad \dots (12)$$

Taking $G(\mathbf{r}, \mathbf{r}')$ such that it satisfies the boundary condition

$$G_1(\mathbf{r}, \mathbf{r}') = 0 \quad \dots (13)$$

whereas the point $\mathbf{r}'$ lies on the surface S , then (12) reduces to

$$\Psi(\mathbf{r}) = -\frac{1}{4\pi} \int_S \Psi(\mathbf{r}') \frac{\partial G_1(\mathbf{r}, \mathbf{r}')}{\partial n} dS' \quad \dots (14)$$

which gives Ψ at any point $\mathbf{r}$ within S .

Again if $G_2(\mathbf{r}, \mathbf{r}')$ is such a function satisfying $\frac{G_2(\mathbf{r}, \mathbf{r}')}{\partial n} = 0$ for $\mathbf{r}'$ lying inside S ... (15)

we have $\Psi(\mathbf{r}) = \frac{1}{4\pi} \int_S \frac{\partial \Psi(\mathbf{r}')}{\partial n} G_2(\mathbf{r}, \mathbf{r}') dS'$... (16)

which gives Ψ at any point within S provided $\frac{\partial \Psi}{\partial n}$ is known at every point of S .

COROLLARY. *Green's function for Diffusion equation:*

The diffusion equation is $\frac{\partial u}{\partial t} = h^2 \nabla^2 u$... (17)

Let $u(\mathbf{r}, t)$ be a solution of it. Then for a volume V enclosed by a surface S , the boundary condition is $u(\mathbf{r}, t) = \phi(\mathbf{r}, t)$... (18)

when $\mathbf{r}$ lies inside S .

The Initial condition is $u(\mathbf{r}, 0) = f(\mathbf{r})$ when $\mathbf{r}$ lies inside V ... (19)

If we define Green's function $G(\mathbf{r}, \mathbf{r}', t-t')$, $t > t'$

such that $\frac{\partial G}{\partial t} = h^2 \nabla^2 G$... (20)

With boundary condition $G(\mathbf{r}, \mathbf{r}', t-t') = 0$ when $\mathbf{r}'$ lies inside S ... (21)

and initial condition $\lim_{t \rightarrow t'} G \rightarrow 0$... (22)

at all the points of V except at the point $\mathbf{r}$ where G takes the form

$$\frac{e^{-\frac{|\mathbf{r}-\mathbf{r}'|^2}{4h^2(t-t')}}}{8[\pi h^2(t-t')]^{3/2}} \quad \dots (23)$$

Now G being a function of t and hence of $(t-t')$ only, (20) is equivalent to

$$\frac{\partial G}{\partial t'} + h^2 \nabla^2 G = 0 \quad \dots (24)$$

Physically interpreted $G(\mathbf{r}, \mathbf{r}', t-t')$ in the temperature at any point $\mathbf{r}'$ at time t due to an instantaneous point source of unit strength generated at time t' of the point $\mathbf{r}$. Initially, the temperature of the solid is zero and the surface is kept at zero temperature.

Equations (17) and (18) being valid for $t' < t$, can be rewritten as

$$\frac{\partial u}{\partial t'} = h^2 \nabla^2 u, \quad t' < t \quad \dots (25)$$

and $u(\mathbf{r}', t) = \phi(\mathbf{r}', t)$ when $\mathbf{r}'$ lies inside S ... (26)

Equations (24) and (25) yield,

$$\frac{\partial}{\partial t'} (uG) = u \frac{\partial G}{\partial t'} + G \frac{\partial u}{\partial t'} = h^2 [G \nabla^2 u - u \nabla^2 G]$$

so that for an arbitrary small $\epsilon > 0$, we find

$$\int_0^{t-\epsilon} \left\{ \int_V \frac{\partial}{\partial t'} (uG) dv' \right\} dt' = h^2 \int_0^{t-\epsilon} \left\{ \int_V [G \nabla^2 u - u \nabla^2 G] dv' \right\} dt' \quad \dots (27)$$

or, changing the order of integration,

$$\begin{aligned} & \int_V (uG)_{t'=t-\epsilon} dv' - \int_V (uG)_{t'=0} dv' \\ &= u(\mathbf{r}, t) \int_V [G(\mathbf{r}, \mathbf{r}', t-t')]_{t'=t-\epsilon} dv' - \int_V G(\mathbf{r}, \mathbf{r}', t) f(\mathbf{r}') dv' \end{aligned}$$

By (23), for $G(\mathbf{r}, \mathbf{r}', t-t')$ we have $\int_V [G(\mathbf{r}, \mathbf{r}', t-t')]_{t'=t-0} dv' = 1$

So that when $\epsilon \rightarrow 0$, L.H.S. of (27)

$$= u(\mathbf{r}, t) - \int_V f(\mathbf{r}') G(\mathbf{r}, \mathbf{r}', t) dv'$$

Hence applying Green's theorem to the R.H.S. of (27) and using (18) and (21) we may find

$$-h^2 \int_0^t dt' \int_S \phi(\mathbf{r}', t) \frac{\partial G}{\partial n} dS' \text{ in limit when } \epsilon \rightarrow 0 \text{ and } \frac{\partial G}{\partial n} \text{ denoting}$$

the derivative of G along outward drawn normal to the surface S .

We shall ultimately find,

$$u(\mathbf{r}, t) = \int_V f(\mathbf{r}') G(\mathbf{r}, \mathbf{r}', t) dv' - h^2 \int_0^t dt' \int_S \phi(\mathbf{r}', t) \frac{\partial G}{\partial n} dS' \dots (28)$$

which gives the solution of (17) with boundary conditions (18) and (19).

12.12. HOMOGENEOUS AND INHOMOGENEOUS WAVE EQUATIONS

In the next Chapter we shall discuss Maxwell's electromagnetic field equations in the form $\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$... (1)

$$\nabla \times \mathbf{H} = \mathbf{J} + \frac{\partial \mathbf{D}}{\partial t} \dots (2)$$

$$\nabla \cdot \mathbf{B} = 0 \dots (3)$$

$$\text{and } \nabla \cdot \mathbf{D} = \rho \dots (4)$$

In addition to these equations, we have few more relations in a homogeneous isotropic medium,

$$\mathbf{D} = k\mathbf{E} \dots (5)$$

$$\mathbf{B} = \mu\mathbf{H} \dots (6)$$

$$\text{and } \mathbf{J} = \sigma\mathbf{E} \dots (7)$$

The method of integration to be used here for electrodynamical equations actually leads us to homogeneous wave equation as shown below. For the purpose of their integration, introduce a vector $\mathbf{A}$ known as *magnetic vector potential* such that

$$\mathbf{B} = \nabla \times \mathbf{A} \dots (8)$$

$$(1) \text{ and } (8) \text{ yield } \nabla \times \mathbf{E} = -\frac{\partial}{\partial t} (\nabla \times \mathbf{A}) = -\nabla \times \frac{\partial \mathbf{A}}{\partial t} \dots (9)$$

(on changing the order of time and space derivatives).

$$\text{We can write (9) as } \nabla \times \left(\mathbf{E} + \frac{\partial \mathbf{A}}{\partial t} \right) = 0 \dots (10)$$

which follows that $\left(\mathbf{E} + \frac{\partial \mathbf{A}}{\partial t} \right)$ is an irrotational vector and hence

it is expressible as the gradient of a scalar point function such that

$$\mathbf{E} + \frac{\partial \mathbf{A}}{\partial t} = -\nabla \phi, \phi \text{ being a scalar potential} \dots (11)$$

$$\text{or } \mathbf{E} = -\frac{\partial \mathbf{A}}{\partial t} - \nabla \phi \dots (12)$$

Multiply (2) by μ and using (6), we have

$$\nabla \times \mathbf{B} = \mu \mathbf{J} + \mu \frac{\partial \mathbf{D}}{\partial t} \dots (13)$$

$$\text{But } \nabla \times \mathbf{B} = \nabla \times (\nabla \times \mathbf{A}) = \nabla (\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A} \dots (14)$$

$$\therefore (13) \text{ gives } \nabla (\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A} = \mu \mathbf{J} + \mu \frac{\partial \mathbf{D}}{\partial t} \dots (15)$$

Differentiation of (12) w.r.t. 't' yields

$$\frac{\partial \mathbf{E}}{\partial t} = -\frac{\partial^2 \mathbf{A}}{\partial t^2} - \nabla \frac{\partial \phi}{\partial t} \dots (16)$$

Elimination of $\frac{\partial \mathbf{E}}{\partial t}$ from (15) and (16) with the help of (5) gives

$$\nabla (\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A} = \mu \mathbf{J} + \mu k \left(-\frac{\partial^2 \mathbf{A}}{\partial t^2} - \nabla \frac{\partial \phi}{\partial t} \right) \dots (17)$$

$$\text{or } -\nabla^2 \mathbf{A} = \mu \mathbf{J} - \mu k \frac{\partial^2 \mathbf{A}}{\partial t^2} - \nabla \left(\nabla \cdot \mathbf{A} \mu k \frac{\partial \phi}{\partial t} \right) \dots (18)$$

It follows from (18) that curl of $\mathbf{A}$ is specified by its divergence but $\text{div } \mathbf{A}$ is not specified. But to find $\mathbf{A}$ uniquely, curl $\mathbf{A}$ and $\text{div } \mathbf{A}$ both should be specified and hence let us assume that

$$\nabla \cdot \mathbf{A} = -\mu k \frac{\partial \phi}{\partial t} \dots (19)$$

$$\text{So that (18) yields } \nabla^2 \mathbf{A} - \mu k \frac{\partial^2 \mathbf{A}}{\partial t^2} = -\mu \mathbf{J} \dots (20)$$

$$\text{Also (4) with the help of (5) gives } \nabla \cdot \mathbf{E} = \frac{\rho}{k} \dots (21)$$

$$\text{which with the help of (12) becomes } \nabla \cdot \left(-\frac{\partial \mathbf{A}}{\partial t} - \nabla \phi \right) = \frac{\rho}{k} \dots (22)$$

$$\text{or } -\frac{\partial}{\partial t} (\nabla \cdot \mathbf{A}) - \nabla^2 \phi = \frac{\rho}{k} \dots (23)$$

Elimination of $\nabla \cdot \mathbf{A}$ from (19) and (23), yields

$$\nabla^2 \phi - \mu \kappa \frac{\partial^2 \phi}{\partial t^2} = -\frac{\rho}{\kappa} \quad \dots (24)$$

If we put $c = \frac{1}{\sqrt{\mu \kappa}}$, (20) and (24) reduce to

$$\nabla^2 A - \frac{1}{c^2} \frac{\partial^2 A}{\partial t^2} = -\mu J \quad \dots (25)$$

$$\text{and } \nabla^2 \phi - \frac{1}{c^2} \frac{\partial^2 \phi}{\partial t^2} = -\frac{\rho}{\kappa} \quad \dots (26)$$

which have got the same form and known as *Inhomogeneous wave equations* or *Lorentz's equations* and they lead to the conclusion that magnetic vector potential A and scalar potential ϕ are propagated in accordance with a equation of the form

$$\nabla^2 u - \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} = -f(x, y, z, t) \text{ which is claimed to solve} \quad \dots (27)$$

$$\text{with initial conditions } u=0 \text{ and } \frac{\partial u}{\partial t}=0 \text{ at } t=0 \quad \dots (28)$$

In order to use the method of Laplace transform, assume that $L\{u(x, y, z, t)\} = U(x, y, z, s)$ and $L\{f(x, y, z, t)\} = F(x, y, z, s)$ $\dots (29)$

$$\text{Taking Laplace transform of (27), we get } \nabla^2 U - \frac{s^2}{c^2} U = -F \quad \dots (30)$$

$$\text{It we put } \kappa^2 = -\frac{s^2}{c^2} \text{ } \kappa = \frac{s}{c} i, \text{ } i = \sqrt{-1}, \text{ then it becomes} \quad \dots (31)$$

$$\nabla^2 U + \kappa^2 U + F = 0$$

which is *Helmholtz's equation*.

In particular case (31) can be taken as $\nabla^2 U_0 + \kappa^2 U_0 = 0$ $\dots (32)$ which is the standard form of Helmholtz's equation and its particular solution is

$$U_0 = \frac{e^{\pm i\kappa r}}{r} \quad \dots (33)$$

where r is the distance from a point and U_0 is determined at another point.

Using this particular solution, we can find the general solution of (31) as

$$U(x, y, z, s) = \frac{1}{4\pi} \iiint F(x_1, y_1, z_1) e^{\pm i\kappa r} \cdot \frac{1}{r} dv \quad \dots (34)$$

$$\text{where } r = \sqrt{(x-x_1)^2 + (y-y_1)^2 + (z-z_1)^2} \text{ and } dv = dx dy dz \quad \dots (35)$$

It may be verified that (34) satisfies (31).

Now substituting $\kappa = \frac{s}{c} i$ (35) becomes

$$U(x, y, z, s) = \frac{1}{4\pi} \iiint \frac{F(x_1, y_1, z_1)}{r} e^{\pm s/c r} dv \quad \dots (36)$$

Taking inverse Laplace transform of (36) we find the solution of inhomogeneous wave equation (27) as

$$u(x, y, z, t) = \frac{1}{4\pi} \iiint \frac{f\left(x, y, z, t - \frac{r}{c}\right)}{r} dv \quad \dots (37)$$

[Since we define the Laplace transform of $F(t)$ as $f(s) = L\{F(t)\}$

$= \int_0^\infty e^{-st} F(t) dt$. Under the condition that definite integral of $F(t)$ exists and $F(t) = 0$ for $t < 0$. Also we define the inverse transform $L^{-1}\{f(s)\} = F(t)$ and $L\left\{\frac{d^2 F}{dt^2}\right\} = s^2 f - sF'(0) - s^2 F(0)$ where

$F'(0)$ is $\frac{dF}{dt}$ evaluated at $t=0$ and

$$L^{-1}\{e^{as} f(s)\} = \begin{cases} 0, & t < a \\ F(t-a), & t > a. \end{cases} \text{ Also } L\left\{\frac{\partial^2 u}{\partial t^2}\right\} = s^2 U$$

The equation (37) shows that the effects in variation of $F(x_1, y_1, z_1, t)$ do not approach the point (x, y, z) unless the time t is retarded by r/c .

As such we can write the solutions of (25) and (26) as

$$A = \frac{\mu}{4\pi} \iiint \frac{J(x_1, y_1, z_1, t - \frac{r}{c})}{r} dv \quad \dots (38)$$

and

$$\phi = \frac{1}{4\pi \kappa} \iiint \frac{\rho(x_1, y_1, z_1, t - \frac{r}{c})}{r} dv \quad \dots (39)$$

These give *retarded potentials* of electro-dynamics.

12.13. THEORY OF WAVE GUIDES

Here we have to discuss the propagation of electromagnetic waves travelling in the longitudinal direction in a homogeneous isotropic medium filling the interior of a metal tube of infinite length, under the assumptions

- (i) The tube has a uniform cross-section.
- (ii) The tube is placed straight along x -axis.
- (iii) The conductivity of the tube is infinite.
- (iv) The medium is devoid of free charges.
- (v) x - y plane is the plane of cross-section of the tube.
- (vi) x -axis is along the wave guide.

Taking E_0 , H_0 , σ , κ and μ as electric intensity, magnetic intensity, conductivity, electric inductive capacity and magnetic inductive capacity respectively, we can write the fundamental Maxwell's equations in the forms

$$\nabla \times \mathbf{E}_0 = -\mu \frac{\partial \mathbf{H}_0}{\partial t}, \quad \dots (1)$$

$$\nabla \times \mathbf{H}_0 = \sigma \mathbf{E}_0 + \kappa \frac{\partial \mathbf{E}_0}{\partial t}. \quad \dots (2)$$

$$\nabla \cdot \mathbf{E}_0 = 0 \quad \dots (3)$$

$$\text{and} \quad \nabla \cdot \mathbf{H}_0 = 0 \quad \dots (4)$$

In order to discuss the possible oscillations propagating inside the wave guide, we can take $\mathbf{E}_0$ and $\mathbf{H}_0$ of the form $e^{i\omega t}$ such that

$$\mathbf{E}_0 = \mathbf{E} e^{i(\omega t - ax)} \quad \dots (5)$$

$$\text{and} \quad \mathbf{H}_0 = \mathbf{H} e^{i(\omega t - ax)} \quad \dots (6)$$

The frequency of oscillation being given by

$$= \frac{\omega}{2\pi} \text{ is known as the propagation constant.}$$

$$\text{Now we have } \mathbf{E} = i \mathbf{E}_x + j \mathbf{E}_y + k \mathbf{E}_z \quad \dots (7)$$

$$\text{and} \quad \mathbf{H} = i \mathbf{H}_x + j \mathbf{H}_y + k \mathbf{H}_z \quad \dots (8)$$

If we substitute for $\mathbf{E}_0$ and $\mathbf{H}_0$ from (5) and (6) into (1) and (2) we get the Cartesian components as

$$\begin{aligned} \frac{\partial E_z}{\partial y} - \frac{\partial E_y}{\partial z} &= -i\omega\mu H_x; \quad \frac{\partial E_x}{\partial z} + aE_z = -i\omega\mu H_y; \\ aE_y + \frac{\partial E_x}{\partial y} &= i\omega\mu H_z \end{aligned} \quad \dots (9)$$

$$\begin{aligned} \text{and} \quad \frac{\partial H_z}{\partial y} - \frac{\partial H_y}{\partial z} &= (\sigma + i\omega\kappa)E_x; \quad \frac{\partial H_x}{\partial z} + aH_z = (\sigma + i\omega\kappa)E_y; \\ -aH_y - \frac{\partial H_x}{\partial y} &= (\sigma + i\omega\kappa)E_z \end{aligned} \quad \dots (10)$$

It is observed that there are two types of waves namely (i) *TE* (Transverse electric) or *H* waves and (ii) *TM* (transverse magnetic) or *E* waves, which exist independently and satisfy equations (9) and (10).

Case I. *TE or H waves* are characterized by

$$E_x = 0 \text{ and } H_x \neq 0 \quad \dots (11)$$

which follows that in the direction of propagation, the electric field has no component while the magnetic field has a component.

If we put $E_x = 0$, (9) and (10) yield

$$\frac{\partial E_z}{\partial y} - \frac{\partial E_y}{\partial z} = i\omega\mu H_x; \quad aE_z = -i\omega\mu H_y; \quad aE_y = i\omega\mu H_z \quad \dots (12)$$

$$\begin{aligned} \text{and} \quad \frac{\partial H_z}{\partial y} - \frac{\partial H_y}{\partial z} &= 0; \quad \frac{\partial H_x}{\partial z} + aH_z = (\sigma + i\omega\kappa)E_y; \\ -aH_y - \frac{\partial H_x}{\partial y} &= (\sigma + i\omega\kappa)E_z \end{aligned} \quad \dots (13)$$

Eliminations of H_y , H_z , E_y and E_z yields

$$\frac{\partial^2 H_x}{\partial z^2} + \frac{\partial^2 H_x}{\partial y^2} = -[a^2 - (\sigma + i\omega\kappa)i\omega\mu]H_x \quad \dots (14)$$

But in an electric region inside the wave guide $\sigma < \omega\kappa$ so that (14) reduces to

$$\frac{\partial^2 H_x}{\partial z^2} + \frac{\partial^2 H_x}{\partial y^2} = -(a^2 + \omega^2\mu\kappa)H_x = -K^2 H_x \quad \dots (15)$$

where $K^2 = a^2 + \omega^2\mu\kappa$

Hence magnetic intensity H_x can be determined under given boundary conditions.

Now from (12) and (13) we can derive

$$\begin{aligned} H_y &= -\frac{a}{\kappa^2} \frac{\partial H_x}{\partial y}; \quad H_z = -\frac{a}{\kappa^2} \frac{\partial H_x}{\partial z}; \quad E_y = \frac{i\omega\mu}{a} H_z; \\ E_z &= -\frac{i\omega\mu}{a} H_y \end{aligned} \quad \dots (16)$$

Thus $\mathbf{E}$ and $\mathbf{H}$ can be determined if H_x is known.

In case the surface of the metallic wave guide is a perfect conductor, then the tangential component of $\mathbf{E}$ vanishes and for a *rectangular wave guide* with its sides parallel to y and z axes, $E_x = 0 = E_y$ at the surface of the wave guide. As such it follows from (16) that

$$\frac{\partial H_x}{\partial y} = 0 = \frac{\partial H_x}{\partial z}.$$

If n be the normal to the surface then at the surface of a wave guide of any cross-section, we have

$$\frac{\partial H_x}{\partial n} = 0.$$

Taking general coordinate system, we can write (15) as

$$\nabla^2_{y,z} H_x + K^2 H_x = 0 \quad \dots (17)$$

$$\text{where} \quad \nabla^2_{y,z} \equiv \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \text{ subject to } E_z = -\frac{i\omega\mu}{a} H_y.$$

Its solution therefore gives the possible value of K and hence the value of ' a ' the constant of propagation such as

$$a = \sqrt{K^2 - \omega^2\mu\kappa} \quad \dots (18)$$

These are imaginary values of 'a' which lead to possible wave propagation along wave guide while for real 'a', the wave is rapidly attenuated as it proceeds along x-axis of the wave guide.

Case II. *TM or E waves* are characterised by $H_x=0$ and $E_x \neq 0$... (19)

Here (9) and (10) for $H_x=0$ yield

$$\frac{\partial E_z}{\partial y} - \frac{\partial E_y}{\partial z} = 0; \quad \frac{\partial E_x}{\partial z} + aE_z = -i\omega\mu H_y; \\ aE_y + \frac{\partial E_x}{\partial y} = i\omega\mu H_z \quad \dots (20)$$

$$\text{and } \frac{\partial H_z}{\partial y} - \frac{\partial H_y}{\partial z} = (\sigma + i\omega\kappa)E_x; \quad aH_z = (\sigma + i\omega\kappa)E_y; \\ -aH_y = (\sigma + i\omega\kappa)E_z \quad \dots (21)$$

Elimination of E_z , E_y , H_y and H_z yields

$$\frac{\partial^2 E_x}{\partial y^2} + \frac{\partial^2 E_x}{\partial z^2} = -(a^2 + \omega^2\mu\kappa)E_x = -\kappa^2 E_x \quad \dots (22)$$

For general coordinate system, this can be written as

$$\nabla^2_{y,z} E_x + \kappa^2 E_x = 0$$

where

$$\nabla^2_{y,z} = \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \quad \dots (23)$$

In case of the surface of a perfectly conducting wave guide E has no tangential component at the surface, thereby giving $E_x=0$ at the surface of the wave guide.

The determination of the possible values of K leads to the possible value of 'a' the propagation constant.

From (19) and (20) E_z , E_y , H_y and H_z can be determined.

12.14. SOLUTION OF ONE-DIMENSIONAL WAVE EQUATION

The equation is $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$... (1)

Its solution by D' Alembert's method has already been given in §7.5. Here we solve it by the method of separation of variables.

Assume $u(x, t) = X(x) T(t)$... (2)

where X is a function of x alone and T that of t alone.

$$\therefore \frac{\partial^2 u}{\partial x^2} = T \frac{\partial^2 X}{\partial x^2} \text{ and } \frac{\partial^2 u}{\partial t^2} = X \frac{\partial^2 T}{\partial t^2}$$

which when substituted in (1) give

$$X \frac{d^2 T}{dt^2} = c^2 T \frac{d^2 X}{dx^2} \text{ i.e., } \frac{1}{X} \frac{d^2 X}{dx^2} = \frac{1}{c^2 T} \frac{d^2 T}{dt^2}$$

on dividing throughout by XTc^2 .

As variables are separated, taking λ as constant of separation, we have

$$\frac{1}{X} \frac{d^2 X}{dx^2} = \frac{1}{c^2 T} \frac{d^2 T}{dt^2} = \lambda \text{ giving } \frac{d^2 X}{dx^2} = \lambda X \text{ and } \frac{d^2 T}{dt^2} = c^2 \lambda T \quad \dots (3)$$

There arise three possibilities:

$$(i) \lambda = 0, \text{ so that by (3) } \frac{d^2 X}{dx^2} = 0, \frac{d^2 T}{dt^2} = 0 \\ \text{giving } X = Ax + B, T = Ct + D \quad \dots (4)$$

$$(ii) \lambda = \mu^2, \quad \dots \frac{d^2 X}{dx^2} - \mu^2 X = 0, \frac{d^2 T}{dt^2} - \mu^2 c^2 T = 0 \text{ giving} \\ X = Ae^{\mu x} + Be^{-\mu x}, T = Ce^{\mu ct} + De^{-\mu ct} \quad \dots (5)$$

$$(iii) \lambda = -\mu^2, \quad \dots \frac{d^2 X}{dx^2} + \mu^2 X = 0, \frac{d^2 T}{dt^2} + \mu^2 c^2 T = 0 \text{ giving} \\ X = A \cos \mu x + B \sin \mu x; \\ T = C \cos \mu ct + D \sin \mu ct \quad \dots (6)$$

If we impose the boundary conditions

$$u(0, t) = 0, u(l, t) = 0 \text{ for all } t \quad \dots (7)$$

and the initial condition

$$u(x, 0) = F(x); \left(\frac{\partial u}{\partial t} \right)_{t=0} = g(x) \quad \dots (8)$$

then (7) asserts that $u(0, t) = X(0) T(t) = 0$ and $u(l, t) = X(l) T(t) = 0$

which imply that either $T(t) = 0$ or $X(0) = 0$ and $X(l) = 0$

Thus from (4) when $x=0$, we have $B=0$ and $X(l)=0=A l+B$ then gives $A=0$.

Also from (5) when $x=0$, we have

$$X(0)=0=A+B \text{ and } X(l)=0=Ae^{\mu l} + Be^{-\mu l} \text{ giving } A=B=0.$$

In either case $A=B=0$ give $X(x)=0$, $u(x, t)=0$, so that the solutions (4) and (5) fail to give the solution of (1) and it is the solution (6) which is periodic in time and is capable of giving a solution of (1).

Combining the two solutions of (6) we have a general solution of (1) as

$$u(x, t) = (A \cos \mu x + B \sin \mu x) (C \cos \mu ct + D \sin \mu ct) \quad \dots (9)$$

Now to determine the constants A , B and μ , we adjust them so as (9) satisfies (7) i.e.,

$$u(0, t) = 0 = A \{(\cos \mu ct + D \sin \mu ct)\} = 0 \text{ giving } A = 0$$

and $u(l, t) = 0 = (0 + B \sin \mu l) (C \cos \mu ct + D \sin \mu ct)$ holds for

$$\mu l = n\pi \text{ i.e., } \mu = \frac{n\pi}{l}, n \text{ being a positive integer.}$$

Hence the solution of (1) satisfying the boundary conditions (7), may be written as

$$u_n(x, t) = \left(C_n \cos \frac{n\pi ct}{l} + D_n \sin \frac{n\pi ct}{l} \right) \sin \frac{n\pi x}{l} \quad \dots (10)$$

Now applying the initial condition (8), (10) yields

$$u(x, 0) = C_n \sin \frac{n\pi x}{l} = F(x)$$

$$\text{and } \left(\frac{\partial u}{\partial t}\right)_{t=0} = \left[-\frac{n\pi c}{l} C_n \sin \frac{n\pi ct}{l} + \frac{n\pi c}{l} D_n \cos \frac{n\pi ct}{l}\right]_{t=0} \sin \frac{n\pi x}{l} \\ = \frac{n\pi c}{l} D_n \sin \frac{n\pi x}{l} = g(x)$$

It is notable that a mere single term as solution will not satisfy $u(x, 0)$ and $\left(\frac{\partial u}{\partial t}\right)_{t=0}$.

In fact the solution (2) is linear and homogeneous and hence it indicates that the sum of any number of distinct solutions of (1) is also a solution of (1). As such the required solution of (1) in place of (10) may be taken as

$$u(x, t) = \sum_{n=1}^{\infty} \left(C_n \cos \frac{n\pi ct}{l} + D_n \sin \frac{n\pi ct}{l} \right) \sin \frac{n\pi x}{l} \quad \dots (11)$$

where $C_n \sin \frac{n\pi x}{l} = F(x)$ and $\frac{n\pi c}{l} D_n \sin \frac{n\pi x}{l} = g(x)$.

Of course, the solution (11) satisfies (7) and hence together with (8), it provides

$$u(x, 0) = \sum_{n=1}^{\infty} C_n \sin \frac{n\pi x}{l} = F(x) \text{ and } \left(\frac{\partial u}{\partial t}\right)_{t=0} \\ = \sum_{n=1}^{\infty} \frac{n\pi c}{l} D_n \sin \frac{n\pi x}{l} = g(x) \quad \dots (12)$$

The R.H.S.'s of (12) being Fourier expansion, we have

$$C_n = \frac{2}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx \text{ and } \frac{n\pi c}{l} D_n = \frac{2}{l} \int_0^l g(x) \sin \frac{n\pi x}{l} dx \quad \dots (13)$$

Hence (11) gives the required solution of (1), for all values of C_n and D_n given by (13) satisfying (7) and (8).

COROLLARY 1. In (8) if we assume $g(x)=0$, the initial velocity, then (13) yields $D_n=0$ and hence (12) reduces to

$$u(x, t) = \sum_{n=1}^{\infty} C_n \cos \frac{n\pi ct}{l} \sin \frac{n\pi x}{l} = \frac{1}{2} \sum_{n=1}^{\infty} C_n \left[\sin \frac{n\pi}{l} (x-ct) \right. \\ \left. + \sin \frac{n\pi}{l} (x+ct) \right] \\ = \frac{1}{2} \sum_{n=1}^{\infty} C_n \sin \frac{n\pi}{l} (x-ct) + \frac{1}{2} \sum_{n=1}^{\infty} C_n \sin \frac{n\pi}{l} (x+ct) \quad \dots (14)$$

Thus replacing x by $x-ct$ and $x+ct$ successively in (11) we find two series

$$\sum_{n=1}^{\infty} C_n \sin \frac{n\pi}{l} (x-ct) \text{ and } \sum_{n=1}^{\infty} C_n \sin \frac{n\pi}{l} (x+ct)$$

we may therefore conclude that

$$u(x, t) = \frac{1}{2} [f(x-ct) + f(x+ct)] \quad \dots (15)$$

Which is the solution of wave equation (1), where f is the odd periodic extension of F with period $2l$.

COROLLARY 2. If we put $\lambda_n = \frac{n\pi c}{l}$ then the functions given by (10) are termed as the *Eigen functions* or *characteristic functions* and the values $\lambda_n = \frac{n\pi c}{l}$ are known as *Eigen Values* or *Characteristic Values* of the vibrating string and the set

$\Lambda = (\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_n)$ is known as the *Spectrum*.

We also observe that u_n represents a harmonic motion with frequency $\frac{\lambda_n}{2\pi} = \frac{nc}{2l}$ cycles per unit time. We call this motion as the n th normal mode of the string. In case $n=1$, the normal mode is called as the *fundamental mode* while the normal modes for $n=2, 3, 4, \dots$ are called as *Overtone*.

Note. In §7.5 while discussing the D'Alemberts' method for solving one-dimensional wave equation of the type

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \quad \dots (1)$$

We have found a solution of it in the form

$$u(x, t) = \phi(x+ct) + \psi(x-ct) \quad \dots (2)$$

We require the verification of the boundary condition

$$u(0, t) = 0, u(l, t) = 0 \quad \dots (3)$$

and the initial conditions $u(x, 0) = f(x)$... (4)

$$\text{and } \left(\frac{\partial u}{\partial t}\right)_{t=0} = g(x) \quad \dots (5)$$

$$\text{Obviously } u'(x, t) = \frac{\partial u}{\partial t} = c\phi'(x+ct) - c\psi'(x-ct) \quad \dots (6)$$

Applying (4) and (5) to (2) and (6) we get

$$u(x, 0) = \phi(x) + \psi(x) = f(x) \quad \dots (7)$$

$$\text{and } \left(\frac{\partial u}{\partial t}\right)_{t=0} = c\{\phi'(x) - \psi'(x)\} = 0 \quad \dots (8)$$

Assuming $g(x)=0$ in particular (8) yields

$$\phi'(x) = \psi'(x) \text{ giving on integration } \phi(x) = \psi(x) + \lambda \quad \dots (9)$$

So that (7) and (9) render

$$\phi(x) = \frac{1}{2}[F(x) - \lambda] \quad \dots (10)$$

$$\text{and } \psi(x) = \frac{1}{2}[F(x) + \lambda] \quad \dots (11)$$

whence with the help of (10) and (11), (2) yields

$$u(x, t) = \frac{1}{2}[F(x+ct) + F(x-ct)] \quad \dots (12)$$

which reduces to

$$u(0, t) = \frac{1}{2}[F(ct) + F(-ct)] = 0$$

and $u(l, t) = \frac{1}{2}[F(l+ct) + F(l-ct)] = 0 \quad \dots (13)$
by the use of (3) and (4).

It follows from (13) that the function F is odd and periodic with period $(2l)$ and hence (12) is the solution of (1). Physically interpreted (2) represents two plane waves travelling in opposite directions with the same period.

Problem 22. A string is stretched between two fixed points $(0, 0)$ and $(l, 0)$ and released at rest from the positions $u = \lambda \sin \pi x$. Show that the formula for its subsequent displacement $u(x, t)$ is given by $u(x, t) = \lambda \cos(c\pi t) \sin(\pi x)$, c^2 being diffusivity.

The boundary value problem is $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \quad \dots (1)$

with boundary conditions $u(0, t) = 0$ and $u(l, t) = 0 \quad \dots (2)$

and initial conditions $u(x, 0) = \lambda \sin \pi x = 0$ and $\left(\frac{\partial u}{\partial t}\right)_{t=0} = 0 \quad \dots (3)$

By (11) of §12.14, we therefore have

$$u(x, t) = \sum_{n=1}^{\infty} C_n \cos(n\pi ct) \sin(n\pi x) \text{ where } C_n = 2 \int_0^l \lambda \sin \pi x \sin \frac{n\pi x}{l} dx$$

It is obvious that $C_n = 0$ for $n = 2, 3, \dots$ but $C_1 = \lambda \int_0^l \sin^2 \pi x dx = \lambda$.

Hence $u(x, t) = \lambda \cos(\pi ct) \sin(\pi x)$.

Problem 23. Show that the deflection of vibrating string of length π (its ends being fixed and $c^2 = 1$), corresponding to zero initial velocity and initial deflection $F(x) = \lambda (\sin x - \sin 2x)$ is given by $u(x, t) = \lambda (\cos t \sin x - \cos 2t \sin 2x)$.

The boundary value problem is $\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2}$ (as $c^2 = 1$) $\dots (1)$

with conditions $g(x) = 0$ and $F(x) = \lambda (\sin x - \sin 2x)$.

Hence by (11) of §12.14 we have (as $D_n = 0$)

$$u(x, t) = \sum_{n=1}^{\infty} C_n \cos \frac{n\pi ct}{l} \sin \frac{n\pi x}{l} = \sum_{n=1}^{\infty} C_n \cos nt \sin nx$$

$\because c=1 \text{ and } \pi=l$

$$\text{where } C_n = \frac{2}{\pi} \int_0^{\pi} F(x) \sin \frac{n\pi x}{l} dx = \frac{2}{\pi} \int_0^{\pi} \lambda (\sin x - \sin 2x) \sin nx dx$$

$$= \frac{2\lambda}{\pi} \int_0^{\pi} \sin x \sin nx dx - \frac{2\lambda}{\pi} \int_0^{\pi} \sin 2x \sin nx dx$$

Clearly $C_n = 0$ for $n = 3, 4, 5, \dots$ and $C_1 = \lambda$, $C_2 = \lambda$.

Hence the required deflection of the vibrating string is given by

$$u(x, t) = C_1 \cos t \sin x + C_2 \cos 2t \sin 2x = \lambda (\sin x - \sin 2x)$$

which verifies the assertion.

Problem 24. Solve the wave equation $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$ if the string of length $2a$ is originally plucked at the middle point by giving it an initial displacement d from the mean position.

The boundary value problem is $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$ with initial conditions.

$$F(x) = u(x, 0) = \begin{cases} \frac{d}{a}, & 0 \leq x < a \\ \frac{d}{a} (2a - x), & a \leq x \leq 2a. \end{cases}$$

Also initial velocity being zero i.e., $g(x) = 0$, we have $D_n = 0$

$$\therefore \text{By (11) of §12.14, } u(x, t) = \sum_{n=1}^{\infty} C_n \cos \frac{n\pi ct}{2a} \sin \frac{n\pi x}{2a}.$$

$$\text{where } C_n = \frac{2}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx = \frac{2}{2a} \int_0^{2a} F(u) \sin \frac{n\pi u}{l} du$$

$$= \frac{1}{a} \int_0^a \frac{d}{a} u \sin \frac{n\pi u}{2a} du + \frac{1}{a} \int_a^{2a} \frac{d}{a} (2a - u) \sin \frac{n\pi u}{2a} du$$

$$= \frac{8d}{\pi^2} \frac{1}{n^2} \sin \frac{n\pi}{2}$$

which vanishes for $n = 2, 4, 6, \dots$

$$\text{Hence } u(x, t) = \frac{8d}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} \sin \frac{n\pi}{2} \cos \frac{n\pi ct}{2a} \sin \frac{n\pi x}{2a}$$

$$= \frac{8d}{\pi^2} \sum_{r=1}^{\infty} \frac{(-1)^{r+1}}{(2r-1)^2} \cos \frac{(2r-1)\pi ct}{2a} \sin \frac{(2r-1)\pi x}{2a}.$$

Problem 25. A string is stretched between two fixed points $(0, 0)$ and $(l, 0)$ and released at rest from the deflection given by

$$F(x) = \begin{cases} \frac{2\lambda}{l} x, & 0 < x < \frac{l}{2} \\ \frac{2\lambda}{l} (l - x), & \frac{l}{2} < x < l \end{cases}$$

Show that the deflection of the string at any time t is given by

$$u(x, t) = \frac{8\lambda}{\pi^2} \sum_{n=1}^{\infty} \sin \frac{n\pi}{2} \cos \frac{n\pi ct}{l} \sin \frac{n\pi x}{l}.$$

Put $a = \frac{l}{2}$ in the previous problem.

Problem 26. The points of trisection of a string are pulled aside through a distance d on opposite sides of the equilibrium-position and the string is released from rest. Show that the displacement of the string at any subsequent time is given by

$$u(x, t) = \frac{9d}{\pi^2} \sum_{r=1}^{\infty} \frac{1}{r^2} \sin \frac{2r\pi}{3} \sin \frac{2r\pi}{3a} x \cos \frac{2r\pi ct}{3a}.$$

Also show that the mid-point of the string always remains at rest.




Fig. 12.8

Consider OB as equilibrium position of the string of length $3a$ (say), and C, D are points of trisection, which are pulled through a distance d as opposite sides and released.

The boundary value problem is

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \quad \dots (1)$$

Using the conceptions of coordinate geometry, the equation of line OP is

$$y = \frac{d-0}{a-0}(x-0) \text{ i.e., } y = \frac{d}{a}x$$

the equation of PQ is $y-d = \frac{-d-d}{2a-a}(x-a)$ i.e. $y = \frac{d(3a-2x)}{a}$ and

the equation of QB is $y-(-d) = \frac{0-(-d)}{3a-2a}(x-2a)$ i.e., $y = \frac{d(x-3a)}{a}$.

Hence the initial deflection is stated as

$$F(x) = \begin{cases} \frac{d}{a}x, & 0 \leq x \leq a \\ \frac{d}{a}(3a-2x), & a \leq x \leq 2a \\ -\frac{d}{a}(x-3a), & 2a \leq x \leq 3a \end{cases}$$

and the initial velocity $g(x=0)$ so that $D_n=0$ in (11) of §12.14

$\therefore$ we have from (B) of §12.14

$$C_n = \frac{2}{l} \int_0^l F(x) \sin \frac{n\pi x}{l} dx = \frac{2}{3a} \frac{d}{a} \left[\int_0^a x \sin \frac{n\pi x}{3a} dx + \int_a^{2a} (3a-2x) \sin \frac{n\pi x}{3a} dx + \int_{2a}^{3a} (x-3a) \sin \frac{n\pi x}{3a} dx \right]$$

$$= \frac{18d}{n^2\pi^2} \{1 + (-1)^n\} \sin \frac{n\pi}{3} \text{ on integrating by parts and simplifying}$$

Obviously $C_n=0$ for $x=1, 3, 5, 7, \dots$ i.e. n being odd

$$\text{and for even } n, C_n = \frac{18d}{n^2\pi^2} \cdot 2 \sin \frac{n\pi}{3} = \frac{36d}{n^2\pi^2} \sin \frac{n\pi}{3}.$$

Hence by (11) of §12.4, the solution is

$$u(x, t) = \frac{36}{\pi^2} \sum_{r=1}^{\infty} \frac{1}{(2r)^2} \sin \frac{2r\pi}{3} \sin \frac{2r\pi}{3a} x \cos \frac{2r\pi ct}{3a}$$

$$= \frac{9d}{\pi^2} \sum_{r=1}^{\infty} \frac{1}{r^2} \sin \frac{2r\pi}{3} \sin \frac{2r\pi}{3a} x \cos \frac{2r\pi ct}{a^3}$$

If we set $x = \frac{3a}{2}$, this result reduces to

$$u(x, t) = 0 \text{ since } \sin \frac{2r\pi}{3a} \cdot x = \sin r\pi = 0 \text{ for each } r.$$

This follows that the mid-point of the string always remains at rest.

12.15. SQUARE WAVE

(Agra, 1961)

In case of a square wave the displacement x is constant $=c$ (say) for certain interval of time and zero for the next same interval and so on. This wave is as shown in Fig. 12.9.




Fig. 12.9

Thus for OA , $x=f(x)=c$ from $t=0$ to $t=T/2$ }
and for AB , $x=f(x)=0$ from $t=T/2$ to T } $\dots (1)$

Taking time-axis of co-ordinates through the lowest point of the displacement curve, we have by Fourier theorem,

$$f(x) = a_0 + \sum_{r=1}^{\infty} a_r \cos rnt + \sum_{r=1}^{\infty} b_r \sin rnt. \quad \dots (2)$$

Here a_0 shows the distance between the axis of coordinates and the axis of the wave and $a_0 = \frac{1}{T} \int_0^T f(x) dt$, $\dots (3)$

$$a_r = \frac{2}{T} \int_0^T f(x) \cos(rnt) dt, \quad \dots (4)$$

$$b_r = \frac{2}{T} \int_0^T f(x) \sin(rnt) dt. \quad \dots (5)$$

Thus we have

$$a_0 = \frac{1}{T} \int_0^T f(x) dt = \frac{1}{T} \left[\int_0^{T/2} f(x) dt + \int_{T/2}^T f(x) dt \right]$$

$$= \frac{1}{T} \int_0^{T/2} c dt = \frac{c}{2}.$$

But the axis of displacement curve being the line $x=c/2$ shown by dotted line in Fig. 12.9, we have

$$a_r = \frac{2}{T} \left[\int_0^{T/2} f(x) \cos rnt dt + \int_{T/2}^T f(x) \cos rnt dt \right]$$

$$= \frac{2}{T} \int_0^{T/2} c \cos rnt dt = 0.$$

$$\text{and } b_r = \frac{2}{T} \left[\int_0^{T/2} c \sin rnt dt \right] = \frac{c}{\pi r} [1 - \cos r\pi] = 0 \text{ if } r \text{ is even}$$

$$= \frac{2c}{\pi} \text{ if } r \text{ is odd.}$$

As such the even terms of the sine series disappear and we are left with odd terms only, so that

$$x=f(x)=\frac{c}{2}+\frac{2c}{\pi}\left[\sin nt+\frac{\sin 3nt}{2}+\frac{1}{5}\sin 5nt+\dots+\frac{1}{r}\sin rnt+\dots\right]$$




Fig. 12.10




Fig. 12.11

This series represents a square wave. If we take first three terms, then curve is shown by thick lines in Fig. 12.10.

In case we take large number of terms, then the curve is as shown in Fig. 12.11.

12.16. SOLUTION OF TWO-DIMENSIONAL WAVE EQUATION

[A] Vibrations of a Rectangular Membrane

Consider the oscillation of a uniform rectangular membrane for which due to uniformity T and ρ are constants, T being tension in dynes per cm. length of the edge and ρ the density in gm. per cm² of the membrane. The free oscillations of it are given by

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) \quad \dots (1)$$

where $c = \sqrt{\frac{T}{\rho}}$ = constant.

With boundary conditions

$$u=0 \text{ for } x=0, x=a, y=0, y=b$$

i.e. for $u=u(x, y, t)$, the conditions are




Fig. 12.12

$$u(0, y, t)=0, u(a, y, t)=0, u(x, 0, t)=0, u(x, b, t)=0 \quad \dots (2)$$

Also the initial displacement and initial velocity are given by

$$u(x, y, 0)=F(x, y) \text{ and } u_t(x, y, 0) = \left[\frac{\partial}{\partial t} u(x, y, t) \right]_{t=0} = g(x, y) \quad \dots (3)$$

$$\text{Assume that } u(x, y, t)=X(x) Y(y) T(t) \quad \dots (4)$$

where X is the function of x only, Y is that of y only and T is that of t only.

$$\text{Then (1) yields } \frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2} = \frac{1}{c^2 T} \frac{d^2 T}{dt^2} \quad \dots (5)$$

Here variables have been separated and hence choice of the constant of separation gives way to three possibilities:

$$(i) \quad \frac{1}{X} \frac{d^2 X}{dx^2} = 0, \frac{1}{Y} \frac{d^2 Y}{dy^2} = 0, \frac{1}{c^2 T} \frac{d^2 T}{dt^2} = 0 \text{ giving } X=A_1 x+B_1, \\ Y=A_2 y+B_2, T=A_3 t+B_3 \quad \dots (6)$$

$$(ii) \quad \frac{1}{X} \frac{d^2 X}{dx^2} = \lambda_1^2, \frac{1}{Y} \frac{d^2 Y}{dy^2} = \lambda_2^2, \frac{1}{c^2 T} \frac{d^2 T}{dt^2} = \lambda^2 \text{ so that} \\ \lambda^2 = (\lambda_1^2 + \lambda_2^2) c^2, \text{ giving}$$

$$X=A_1 e^{\lambda_1 x} + B_1 e^{-\lambda_1 x}, Y=A_2 e^{\lambda_2 y} + B_2 e^{-\lambda_2 y}, T=A_3 e^{c\lambda t} + B_3 e^{-c\lambda t} \quad \dots (7)$$

$$(iii) \quad \frac{1}{X} \frac{d^2 X}{dx^2} = -\lambda_1^2, \frac{1}{Y} \frac{d^2 Y}{dy^2} = -\lambda_2^2, \frac{1}{c^2 T} \frac{d^2 T}{dt^2} = -\lambda^2$$

so that $\lambda^2 = (\lambda_1^2 + \lambda_2^2) c^2$, giving

$$X=A_1 \cos \lambda_1 x + B_1 \sin \lambda_1 x, Y=A_2 \cos \lambda_2 y + B_2 \sin \lambda_2 y, \\ T=A_3 \cos \lambda c t + B_3 \sin \lambda c t \quad \dots (8)$$

In view of 2nd and 4th conditions of (2), we conclude that

$$X(0)=X(a)=0, Y(0)=0=Y(b),$$

thereby rendering the solutions (5) and (6) unable to give the solution of (1).

As regards (7), the function $X(x)$ is a linear combination of $\sin \lambda_1 x$ and $\cos \lambda_1 x$ in which cosine is to be rejected in view of the condition $u=0$ at $x=0$ i.e. $u(0, y, t)=0$ and hence $X(0)=0$. We are left with $X(x)=B_1 \sin \lambda_1 x=0$ as $u(a, y, t)=0$ gives $X(a)=0$ so that $\sin \lambda_1 a=0=\sin m\pi$, m being a positive integer. Thus we get

$$\lambda_1 = \frac{m\pi}{a} \text{ and similarly } Y=\sin \lambda_2 y=0 \text{ yields}$$

$$\lambda_2 = \frac{n\pi}{b} \text{ for integral } n.$$

Consequently the required solution of (1) takes the form

$$(A \cos \lambda_{mn} t + B \sin \lambda_{mn} t) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} \quad \dots (9)$$

where A and B are constants and $\lambda_{mn} = \lambda$ is given by

$$\lambda_{mn}^2 = \left(\frac{m\pi c}{a} \right)^2 + \left(\frac{n\pi c}{b} \right)^2 \quad \dots (10)$$

Such quantities are termed as *eigen values* or *characteristic values* of vibrating membranes.

Trying superposition and accounting for different constants A and B for each choice of m and n , we can write the solution as

$$u(x, y, t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} (A_{mn} \cos \lambda_{mn} t + B_{mn} \sin \lambda_{mn} t) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} \quad \dots (11)$$

Now applying the condition (3), i.e. for $t=0$, $u(x, y, 0) = F(x, y)$, we get

$$F(x, y) = u(x, y, 0) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} \quad \dots (12)$$

which is known as the *Double Fourier Sine Series* of $F(x, y)$.

Multiplying (11) by $\sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b}$ and integrating over the rectangle

$$x=0, x=a, y=0, y=b \text{ we find the value of } A_{mn} \text{ i.e.,} \\ A_{mn} = \frac{4}{ab} \int_0^a \int_0^b F(x, y) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} dx dy \quad \dots (13)$$

Also differentiating (10) w.r.t. ' t ' and applying the condition (3)

i.e. for $t=0$ $\frac{\partial}{\partial t} u(x, y, t) = g(x, y)$, we get

$$B_{mn} = \frac{4}{ab\lambda_{mn}} \int_0^a \int_0^b g(x, y) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} dx dy \quad \dots (14)$$

Obviously the general term of (10) is a periodic function of time with period $\frac{2\pi}{\lambda_{mn}}$ and having the frequency

$$\frac{\lambda_{mn}}{2\pi} = \frac{c}{2} \left[\left(\frac{m}{a} \right)^2 + \left(\frac{n}{b} \right)^2 \right]^{1/2} \quad \dots (15)$$

which are termed as *characteristic frequencies* or *eigen frequencies* and the associated oscillations given by (8) are termed as *Modes*.

The *fundamental mode* is the mode of the lowest frequency obtained by putting $m=n=1$ in (8).

COROLLARY. If we assume that the initial velocity $g(x, y) = 0$ so that by (3), $u_t(x, y, 0) = 0$ then clearly $B_{mn} = 0$ and hence the solution of (1) reduces to the form

$$u(x, y, t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} \cos \lambda_{mn} t \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} \quad \dots (16)$$

$$\text{where } A_{mn} = \frac{4}{ab} \int_0^a \int_0^b F(x, y) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} dx dy \quad \dots (17)$$

$$\text{and } \lambda_{mn} = \pi^2 c^2 \left[\frac{m^2}{a^2} + \frac{n^2}{b^2} \right] \quad \dots (18)$$

Note. *Distinction between the behaviour of vibrating strings and membranes.*

The main difference is that for every eigen frequency of vibration of a string there exists a corresponding mode such that the string is divided into equal parts by fixed posited nodes, whereas for an oscillating membrane with a given characteristic property, there exists points on the membrane which are at rest and they constitute *nodal lines*. Since the shape of the nodal lines for a given frequency is not the same, therefore for a given eigen frequency there may be more than one mode e.g. for a rectangular membrane with $a=b$, by (14), the frequency is given by

$$\lambda_{mn} = \frac{c\pi}{a} \sqrt{m^2 + n^2} = \kappa \sqrt{m^2 + n^2} \text{ where } \kappa = \pi c/a \quad \dots (19)$$

and by (10) the fundamental mode is $u_{11} = (A_{11} \cos \lambda_{11} t + B_{11} \sin \lambda_{11} t)$

$$\sin \frac{\pi x}{a} \sin \frac{\pi y}{a} \text{ where } \lambda_{11} = a\sqrt{2}.$$

But u_{11} being zero for all t only when $x=0, y=0, x=a, y=a$, therefore in the interior of the membrane for this frequency, there does not exist a nodal line.

In case $m=1, n=2$ and $m=2, n=1$, we find two modes

$$u_{12} = (A_{12} \cos \lambda_{12} t + B_{12} \sin \lambda_{12} t) \sin \frac{\pi x}{a} \sin \frac{2\pi y}{a} \text{ and } u_{21} \\ = (A_{21} \cos \lambda_{21} t + B_{21} \sin \lambda_{21} t) \sin \frac{2\pi x}{a} \sin \frac{\pi y}{a} \quad \dots (20)$$

having the same frequency since $\lambda_{12} = \lambda_{21} = a\sqrt{5}$ for $y = \frac{a}{2}$,

$$u_{12} = 0 \text{ and for } x = \frac{a}{2}, u_{21} = 0$$

We can thus show the existence of oscillations with the same frequency but different nodal lines.

[B] Vibrations of a circular membrane. (Bessel's functions)

If we transform $\frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$ by the substitutions

$$x = r \cos \theta, y = r \sin \theta$$

$$\text{we get the polar form } \frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} \right) \quad \dots (1)$$

where $u(r, \theta, t)$ is the deflection of the membrane.

The equation (1) is known as the *wave equation for a circular membrane*, if the boundary is the circle $r=a$ so that the boundary condition is

$$u(a, \theta, t) = 0, \quad -\pi \leq \theta \leq 0, \quad t \geq 0 \quad \dots (2)$$

Let us take the initial conditions for $0 \leq r \leq a, -\pi \leq \theta \leq 0$, as

$$u(r, \theta, 0) = F(r, \theta) \quad \dots (3)$$

and

$$u_t(r, \theta, 0) = \left[\frac{\partial u}{\partial t}(r, \theta, t) \right]_{t=0} = g(r, \theta) \quad \dots (4)$$

Assume that $u(r, \theta, t) = R(r)\Theta(\theta)T(t)$ $\dots (5)$
where R is the function of r alone, Θ that of θ alone and T that of t alone.

Then (1) yields

$$\frac{1}{c^2 T} \frac{d^2 T}{dt^2} = \frac{1}{R} \frac{d^2 R}{dr^2} + \frac{1}{rR} \frac{dR}{dr} + \frac{1}{r^2 \Theta} \frac{d^2 \Theta}{d\theta^2} = -\lambda^2 \text{ (say) as variables are separated.}$$

The terms in it being constant we can take

$$\frac{1}{\Theta} \frac{d^2 \Theta}{d\theta^2} = -m^2.$$

As such we have the ordinary differential equations for R , Θ and T as

$$\frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + \left(\lambda^2 - \frac{m^2}{r^2} \right) R = 0 \quad \dots (6)$$

$$\frac{d^2 \Theta}{d\theta^2} + m^2 \Theta = 0 \quad \dots (7)$$

$$\frac{d^2 T}{dt^2} + c^2 \lambda^2 T = 0 \quad \dots (8)$$

The solution of (7) is of the form $\Theta = D e^{\pm im\theta}$ $\dots (9)$
 $m = 0, 1, 2, \dots$

D being a constant.

Also putting $s = \lambda r$, (6) becomes

$$\frac{d^2 R}{ds^2} + \frac{1}{s} \frac{dR}{ds} + \left(1 - \frac{m^2}{s^2} \right) R = 0 \quad \dots (10)$$

which is Bessel's equation and hence its general solution is

$$R(r) = K_1 J_m(s) + K_2 J_{-m}(s) = K_1 J_m(\lambda r) + K_2 J_{-m}(\lambda r) \quad \dots (11)$$

K_1, K_2 being constants.
But as $r \rightarrow 0, J_{-m} \rightarrow \infty$ thereby contradicting the hypothesis that deflection of the membrane is always finite. Thus to avoid the term containing J_{-m} choose $K_2 = 0$, so that (11) reduces to $R(r) = K_1 J_m(\lambda r)$ $\dots (12)$

The boundary condition (2) requires that

$$R(a) = K_1 J_m(\lambda a) = 0 \text{ i.e. } J_m(\lambda a) = 0 \quad \dots (13)$$

Assuming $\lambda_1, \lambda_2, \lambda_3, \dots$ as positive roots of (13), the general solution of (8) is

$$T = A \cos c\lambda t + B \sin c\lambda t \quad \dots (14)$$

With the help of (9), (12) and (14), the required oscillation has the form

$$\{A \cos c\lambda t + B \sin c\lambda t\} e^{\pm im\theta} J_m(\lambda r) \quad \dots (15)$$

Trying superposition and using distinct values of constants A and B for each choice of m and n , the general solution of (1) may be taken as

$$u(r, \theta, t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \{A_{mn} \cos(c\lambda_{mn}t) + B_{mn} \sin(c\lambda_{mn}t)\} e^{\pm im\theta} J_m(\lambda_{mn}r) \quad \dots (16)$$

which satisfies the conditions (2), (3) and (4).

In case the equation (1) is radially symmetrical i.e. the solution is independent of θ , we have from (16) by putting $m=0$,

$$u(r, t) = \sum_{n=1}^{\infty} \{A_n \cos c\lambda_n t + B_n \sin c\lambda_n t\} J_0(\lambda_n r) \quad \dots (17)$$

where $\lambda_1, \lambda_2, \lambda_3, \dots$ are the positive roots of $J_0(\lambda a) = 0$

Here (17) gives the general solution of (1).

In case $t=0$, the initial condition (3) gives

$$u(r, 0) = F(r) = \sum_{n=1}^{\infty} A_n J_0(\lambda_n r) \quad \dots (18)$$

since $F(r, \theta)$ reduces to $F(r)$ when independent of θ .

Here A_n is the coefficient of Fourier-Bessel series determined by

$$A_n = \frac{2}{a^2 J_1^2(\lambda_n a)} \int_0^a F(r) J_0(\lambda_n r) r dr \quad \dots (19)$$

Also the condition (4) requires,

$$u_t(r, 0) = \sum_{n=1}^{\infty} c\lambda_n B_n J_0(\lambda_n r) = g(r), \text{ } g \text{ being independent of } \theta \text{ and}$$

hence B_n are determined by

$$B_n = \frac{2}{c\lambda_n a^2 J_1^2(\lambda_n a)} \int_0^a g(r) J_0(\lambda_n r) r dr \quad \dots (20)$$

COROLLARY. If we take $g(r, \theta) = u_t(r, \theta, 0) = 0$ then (17) yields on putting $t=0$.

$$u(r, t) = \sum_{n=1}^{\infty} A_n J_0(\lambda_n r) \cos(\lambda_n ct) \quad \dots (21)$$

where $F(r) = \sum_{n=1}^{\infty} A_n J_0(\lambda_n r)$ and the coefficients A_n of Fourier-Bessel series are given by

$$A_n = \frac{2}{a^2 J_1^2(\lambda_n a)} \int_0^a F(r) J_0(\lambda_n r) r dr \quad \dots (22)$$

The solution (21) may be expressed by means of modes and eigen frequencies as

$$\frac{\lambda_n}{2\pi} = \frac{c\lambda_n}{2\pi} = \frac{c\lambda_n}{2\pi a}$$

and the fundamental mode is given by $J_0(\lambda_1 r) \cos(\lambda_1 ct)$.

Note. The generalized Fourier-Bessel series is

$$f(x) = \sum_{m=1}^{\infty} C_m J_n(\lambda_{mn})$$

where

$$\lambda_{mn} = \frac{\alpha_{mn}}{a}, n \text{ fixed and } m=1, 2, 3 \dots$$

$$\text{also } C_m = \frac{1}{a^2 J_{n+1}^2(\alpha_{mn})} \int_0^a x f(x) J_n(\lambda_{mn} x) dx, m=1, 2, 3 \dots$$

Problem 27. Find the deflection $u(x, y, t)$ of the square membrane with $a=b=1$ and $c=1$ under the conditions that the initial velocity is zero and the initial deflection is

$$F(x, y) = A \sin \pi x \sin 2\pi y.$$

The boundary value problem is

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left[\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right] \text{ where } c^2=1 \quad \dots (1)$$

$$\text{with boundary conditions } \left. \begin{array}{l} u=0 \text{ for } x=0, x=1, \\ y=0, y=1 \end{array} \right\} \quad \dots (2)$$

$$\text{and the initial conditions } u(x, y, 0) = F(x, y) = A \sin \pi x \sin 2\pi y \quad \dots (3)$$

$$\text{and } u_t(x, y, 0) = g(x, y) = 0 \quad \dots (4)$$

By (10) of §12.16, we therefore have the solution as

$$u(x, y, t) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} A_{mn} \lambda_{mn} t \sin m\pi x \sin n\pi y$$

$$\text{where } \lambda_{mn}^2 = \pi^2(m^2 + n^2) \text{ and } A_{mn} = 4A \int_0^1 \int_0^1 \sin \pi x \sin m\pi x \sin 2\pi y \sin n\pi y dx dy$$

Clearly $A_{mn}=0$ where n is odd.

$$\text{Also } A_{m2} = 4A \int_0^1 \int_0^1 \sin \pi x \sin m\pi x \sin^2 2\pi y dx dy$$

$$= 2A \int_0^1 \sin \pi x \sin m\pi x dx$$

$$= 0 \text{ when } n \text{ is even}$$

$$\text{and } A_{12} = 2A \int_0^1 \sin^2 \pi x dx = A.$$

Hence the required solution is

$$u(x, y, t) = A_{12} \cos \lambda_{12} t \sin \pi x \sin 2\pi y \\ = A \cos \sqrt{5} t \sin \pi x \sin 2\pi y$$

which gives the required deflection of the membrane.

12.17. SOLUTION OF THREE-DIMENSIONAL WAVE EQUATION

The equation is

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} \quad \dots (1)$$

Suppose that we have to solve it under the conditions

$$(i) \frac{\partial u}{\partial x} = 0 \text{ when } x=0, x=a$$

$$(ii) \frac{\partial u}{\partial y} = 0 \text{ when } y=0, y=a$$

$$(iii) \frac{\partial u}{\partial z} = 0 \text{ when } z=0, z=a$$

$$(iv) u \neq 0 \text{ at } t=0.$$

$$\text{Assume } u = X(x) Y(y) Z(z) T(t) \quad \dots (2)$$

Then (1) yields

$$\frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2} + \frac{1}{Z} \frac{d^2 Z}{dz^2} = \frac{1}{c^2 T} \frac{d^2 T}{dt^2} \quad \dots (3)$$

Here variables have been separated. Hence each of the four terms in this equation must be constant say

$$\frac{1}{X} \frac{d^2 X}{dx^2} = -\lambda_1^2; \quad \frac{1}{Y} \frac{d^2 Y}{dy^2} = -\lambda_2^2; \quad \frac{1}{Z} \frac{d^2 Z}{dz^2} = -\lambda_3^2$$

$$\text{and } \frac{1}{c^2 T} \frac{d^2 T}{dt^2} = -\lambda^2 \text{ so that } \lambda^2 = \lambda_1^2 + \lambda_2^2 + \lambda_3^2.$$

Then we get

$$X = A_1 \cos(\lambda_1 x + \alpha_1), Y = A_2 \cos(\lambda_2 y + \alpha_2), Z = A_3 \cos(\lambda_3 z + \alpha_3)$$

and

$$T = B \cos(\lambda c t + \alpha)$$

where $A_1, A_2, A_3, B, \alpha_1, \alpha_2, \alpha_3$ and α all are arbitrary constants.

Hence the solution of (1) is

$$u = \Sigma A \cos(\lambda_1 x + \alpha_1) \cos(\lambda_2 y + \alpha_2) \cos(\lambda_3 z + \alpha_3) \cos(\lambda c t + \alpha) \quad \dots (4)$$

However we can take the solution of (1) in the form

$$u = C e^{\pm i(\lambda_1 x + \lambda_2 y + \lambda_3 z + \lambda c t)}$$

where

$$\lambda^2 = \lambda_1^2 + \lambda_2^2 + \lambda_3^2 \quad \dots (5)$$

Actually, $e^{\pm i\lambda_1 x} = A_1 \cos \lambda_1 x \pm A_2 \sin \lambda_1 x$

and

$$\frac{\partial u}{\partial x} = 0 \text{ when } x=0 \text{ implies } A_2=0 \text{ etc.}$$

Then $u = C \cos \lambda_1 x \cos \lambda_2 y \cos \lambda_3 z \cos \lambda ct$

... (6)

The result (6) may be deduced from (4) since when

$$x=0, \sin \alpha_1 x = 0 \text{ etc.}$$

If we now define $\frac{\partial u}{\partial x}$ as the gradient in the direction of the normal of each face, we see that the boundary condition (iv) holds at $t=0$.

Also $\frac{\partial u}{\partial x} = 0$ when $x=0, y=0, z=0$ and the boundary condition (i) requires that $\sin \lambda_1 a \cos \lambda_2 y \cos \lambda_3 z \cos \lambda ct = 0$

i.e., $\sin \lambda_1 a = 0$ or $\lambda_1 a = N_1 \pi$ i.e., $\lambda_1 = \frac{N_1 \pi}{a}$, N_1 being an integer.

$$\text{Similarly } \lambda_2 = \frac{N_2 \pi}{a}, \lambda_3 = \frac{N_3 \pi}{a} \text{ and } \lambda = \sqrt{\lambda_1^2 + \lambda_2^2 + \lambda_3^2} \\ = \frac{\pi}{a} \sqrt{N_1^2 + N_2^2 + N_3^2}$$

As such (6) reduces to

$$u = C \cos \frac{N_1 \pi}{a} x \cos \frac{N_2 \pi}{a} y \cos \frac{N_3 \pi}{a} z \cos \left\{ (N_1^2 + N_2^2 + N_3^2)^{1/2} \frac{\pi ct}{a} \right\}$$

Hence the general solution of (1) is

$$u = \sum_{N_1=1}^{\infty} \sum_{N_2=1}^{\infty} \sum_{N_3=1}^{\infty} C_{N_1 N_2 N_3} \cos \frac{N_1 \pi}{a} x \cos \frac{N_2 \pi}{a} y \cos \frac{N_3 \pi}{a} z \cos \left\{ (N_1^2 + N_2^2 + N_3^2)^{1/2} \frac{\pi ct}{a} \right\} \quad \dots (6)$$

Note 1. Cylindrical form of three-dimensional wave equation is

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial z^2} = \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} \quad \dots (7)$$

Assuming $u = R(r) \cdot \Theta(\theta) \cdot Z(z) \cdot T(t)$, (7) yields

$$\frac{1}{R} \left(\frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} \right) + \frac{1}{r^2 \Theta} \frac{d^2 \Theta}{d\theta^2} + \frac{1}{Z} \frac{d^2 Z}{dz^2} = \frac{1}{c^2 T} \frac{d^2 T}{dt^2},$$

where variables are separated,

$$\text{Taking } \frac{1}{\Theta} \frac{d^2 \Theta}{d\theta^2} = -\mu^2, \frac{1}{Z} \frac{d^2 Z}{dz^2} = -\nu^2, \frac{1}{T} \frac{d^2 T}{dt^2} = -c^2 \lambda^2, \quad \dots (8)$$

λ^2, μ^2, ν^2 being constants, we have

$$\frac{1}{R} \left(\frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} \right) - \frac{\mu^2}{r^2} = \nu^2 - \lambda^2 = -\alpha^2 \text{ (say) giving}$$

$$\frac{d^2 R}{dr^2} + \frac{1}{r} \frac{dR}{dr} + \left(\alpha^2 - \frac{\mu^2}{r^2} \right) R = 0 \quad \dots (9)$$

which is Bessel's equation of order m and argument αr .

$\therefore$ Solutions of (8) and (9) are

$$\Theta = A_1 \cos \mu \theta + B_1 \sin \mu \theta, Z = A_2 \cos \nu z + B_2 \sin \nu z$$

$$T = A_3 \cos \lambda ct + B_3 \sin \lambda ct, R = A_4 J_\mu(\alpha r) + B_4 Y_\mu(\alpha r)$$

Hence the general solution of 7, is

$$u = \Sigma A J_\mu(\alpha r) \cos \mu \theta \cos \nu z \cos \lambda ct \quad \dots (10)$$

where $\alpha^2 = \lambda^2 - \nu^2$ and the term $Y_\mu(\alpha r)$ may be included if required by the conditions of the problem.

In case of axial symmetry, u is independent of θ i.e., $\mu=0$ and hence $u = \Sigma A J_0(\alpha r) \cos \nu z \cos \lambda ct \quad \dots (i1)$

Also if u is independent of z , then $\nu=0$ and $\alpha=\lambda$, hence

$$u = \Sigma A J_0(\lambda r) \cos \lambda ct. \quad \dots (12)$$

Note 2. Polar spherical form of three dimensional wave equation is

$$\frac{\partial^2 u}{\partial r^2} + \frac{2}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{\cot \theta}{r^2} \frac{\partial u}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 u}{\partial \phi^2} = \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} \quad \dots (13)$$

Assuming $u = R(r) \Theta(\theta) \Phi(\phi) T(t)$, (13) yields

$$\frac{1}{R} \frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} + \frac{1}{r^2 \Theta} \frac{d^2 \Theta}{d\theta^2} + \frac{\cot \theta}{r^2 \Theta} \frac{d\Theta}{d\theta} + \frac{1}{r^2 \sin^2 \theta} \frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2} \\ = \frac{1}{c^2 T} \frac{d^2 T}{dt^2} \quad \dots (14)$$

As variables are separated, we can take

$$\frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2} = -m^2, \frac{1}{c^2 T} \frac{d^2 T}{dt^2} = -p^2 \quad \dots (15)$$

$$\text{and } \frac{1}{\Theta} \frac{d^2 \Theta}{d\theta^2} + \frac{\cot \theta}{\Theta} \frac{d\Theta}{d\theta} = \frac{m^2}{\sin^2 \theta} - n(n+1) \quad \dots (16)$$

where m, n, p are constants.

(14) reduces with the help of (15) and (16), to

$$\frac{1}{R} \frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} = \frac{n(n+1)}{r^2} - p^2 \quad \dots (17)$$

Solutions of (15) are $\Phi = \cos m\phi$ or $\sin m\phi$ or $e^{\pm im\phi}$ and $T = \cos pct$ or $\sin pct$ or $e^{\pm icpt}$

Also equation (16) can be written as

$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) + \left\{ n(n+1) - \frac{m^2}{\sin^2 \theta} \right\} \Theta = 0 \quad \dots (18)$$

Putting $\cos \theta = \mu$, so that $\frac{1}{\sin \theta} \frac{d}{d\theta} = -\frac{d}{d\mu}$, (18) yields

$$\frac{d}{d\mu} \left\{ (1-\mu^2) \frac{d\Theta}{d\mu} \right\} + \left\{ n(n+1) - \frac{m^2}{1-\mu^2} \right\} \Theta = 0$$

$$(1-\mu^2) \frac{d^2\Theta}{d\mu^2} - 2\mu \frac{d\Theta}{d\mu} + \left\{ n(n+1) - \frac{m^2}{1-\mu^2} \right\} \Theta = 0 \quad \dots (19)$$

which is associated Legendre's equation, having the solution

$$\Theta = AP_n^m(\mu) + BQ_n^m(\mu) = AP_n^m(\cos \theta) + BQ_n^m(\cos \theta) \quad \dots (20)$$

$$\text{If we neglect } Q_n, \text{ we can write } \Theta = AP_n^m(\cos \theta) \quad \dots (21)$$

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 28. Solve $\frac{\partial^2 u}{\partial x^2} = 0$ subject to the conditions $u = \lambda^2$ when $x = 0$
 $u = 1$ when $x = 1$

$$\text{Ans. } u = x(1-\lambda^2) + \lambda^2 \text{ when } \frac{\partial u}{\partial x} = \phi(\lambda).$$

Problem 29. Transform the equation $\nabla^2 \Psi = \frac{1}{c^2} \frac{\partial^2 \Psi}{\partial t^2}$ to spherical polar coordinates. If Ψ depends on r and t only, show that the equation can be written in the form $\frac{\partial^2}{r^2} (r\Psi) = \frac{1}{c^2} \frac{\partial^2}{\partial t^2} (r\Psi)$. Hence show that the general solution is of the form

$$\Psi(r, t) = \frac{1}{r} \{ f(r-ct) + g(r+ct) \}$$

Explain the physical meaning of this solution. (Bombay, 1965)

Problem 30. Establish the equation for heat conduction $k \nabla^2 T = \frac{\partial T}{\partial t}$.

A slender rod whose curved surface is perfectly insulated stretches from $x=0$ to $x=\infty$. Find the temperature in the rod as a function of x and t if the left end of the rod is maintained at the constant temperature 0°C and if initially the temperature along the rod is given by $T(x, 0) = f(x)$. (Bombay, 1965)

Problem 31. Solve the partial differential equation for the vibrations of a square elastic membrane fixed at the edges, $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2}$

The length of each side of the square is L , c is the velocity of elastic waves and $u(x, y, z)$ is the displacement of the point (x, y) at time t normal to the plane of the membrane.

Obtain the lowest frequency of vibration if $c = 10,000$ metres per sec. and $L = \sqrt{2}$ metres. (Agra, 1967)

Problem 32. Solve the differential equation $\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} + \frac{\partial^2 \psi}{\partial z^2} + \lambda^2 \psi = 0$ in which λ is a constant and the function ψ exists only in the region defined by $0 \leq x \leq l_1$; $0 \leq y \leq l_2$; $0 \leq z \leq l_3$ (Vikram, 1967)

Problem 33. Solve the two-dimensional Laplace's equation $\nabla^2 \phi = 0$ in the first quadrant i.e., $0 \leq x < \infty$ and $0 \leq y < \infty$, with the boundary condition that on the x and y axes $\phi = \phi$. (Agra, 1968; Vikram, 1969)

Problem 34. Express ∇^2 in cylindrical and spherical polar coordinates. (Vikram, 1969)

Problem 35. Write down the general wave equation. Solve it for a rectangular elastic membrane under uniform tension and fixed at the boundaries. What are the different modes under which it can vibrate? (Agra, 1969)

Problem 36. Obtain the solution for a guided electromagnetic wave through an infinitely long cavity of rectangular cross-section and discuss the limitations on the propagation wave length. (Agra, 1969, 1970)

Problem 37. Obtain the partial differential equation for the heat conduction in a homogeneous medium.

i.e., four faces of an infinitely long rectangular prism bounded by the planes $x=0$, $x=a$, $y=0$ and $y=b$ are kept at temperature zero. If the initial temperature distribution is $\theta_0(x, y)$, find an expression for the temperature at a point in terms of x , y and the time t . (Bombay, 1970)

Problem 38. What is meant by 'circular membrane'? Find a general expression for the same. (Bombay, 1970)

Problem 39. Find an expression for the velocity of transverse waves along a stretched membrane.

A circular membrane is under a uniform tension. Considering the wave equation, discuss its vibrations. (Bombay, 1970)

Problem 40. Deduce the Fourier equation for one dimensional flow of heat along a solid bar. If one end of the bar is periodically heated and cooled and the bar is covered with heat insulating material, find the temperature changes along any point of the bar. (Agra, 1970)

Problem 41. Obtain the axially symmetrical solutions of the three dimensional Laplace's equation. (Agra, 1970; 1974)

Problem 42. Calculate the potential due to gravitating ring of uniform density at any point of its axis. (Agra, 1970)

Problem 43. (a) Explain the significance of spherical harmonics and discuss some general properties of harmonic functions.

(b) Determine the potential outside and inside a spherical surface which is kept at a fixed distribution of electrical potential of the form $V = F(\theta)$. It is assumed that the space inside and outside the surface is free of charges. (Agra, 1971)

Problem 44. Solve the one-dimensional wave equation $\frac{\partial^2 f}{\partial x^2} = \frac{1}{c^2} \frac{\partial^2 f}{\partial t^2}$ for a string of length l fixed at both ends. The boundary conditions are

$$\left\{ f(x, t) \right\}_{t=0} = \frac{2hx}{l}, \quad 0 \leq x < \frac{l}{2}$$

$$= \frac{2h}{l}(l-x), \quad \frac{l}{2} \leq x < l.$$

Discuss the nature of the solution. (Agra, 1972)

Problem 45. Write Laplace's equation in the spherical polar coordinates. Solve the equation by the method of separation of variables in the case of cylindrical symmetry. Use this solution to obtain the gravitational potential of a uniform ring of radius a and mass m . Assume the ring to be made of thin wire. (Agra, 1972; 74)

Problem 46. Solve the heat diffusion equation $h^2 \nabla^2 \theta = \frac{\partial \theta}{\partial t}$ for a semi-infinite plate bounded by the lines $x=0$, $x=\infty$; $y=0$ and $y=b$. θ is the temperature, t the time and h is constant. The boundary conditions are prescribed by

$$\left. \begin{aligned} \theta(x, y, t) &= \theta(y) \text{ for } x=0 \\ \theta(x, y, t) &= 0 \text{ for } y=0 \text{ and } y=b \end{aligned} \right\} 0 \leq t < \infty. \quad (\text{Agra, 1973})$$

Problem 47. Define the magnetic vector potential $\mathbf{A}$ and show that any electromagnetic field can be derived from this potential $\mathbf{A}$ and a scalar potential ϕ . Prove that the two potentials satisfy inhomogeneous wave equations of the same form. (Agra, 1973)

Problem 48. Write Laplace's equation in cartesian, cylindrical and spherical coordinates. Solve Laplace's equation for the spherically symmetric case. (Agra, 1975; 1976)

Hint. In addition to the solution of Laplace's equation discussed in this chapter, see relevant sections alongwith Problem 180 of Ch. 1.

Problem 49. Solve the wave equation for electromagnetic waves in a cylindrical wave guide. Discuss the solution, its significance and applications.

The electromagnetic wave equation is

$$\nabla^2 \psi - \mu \epsilon \frac{\partial^2 \psi}{\partial t^2} - \mu \sigma \frac{\partial \psi}{\partial t} = 0 \quad \dots (1)$$

where μ is permeability, σ is conductivity and ϵ is inductive capacity. By § 1.45, in cylindrical coordinates (r, θ, z) , we have

$$\nabla^2 \psi = \frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \psi}{\partial \theta^2} + \frac{\partial^2 \psi}{\partial z^2} = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial \psi}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 \psi}{\partial \theta^2} + \frac{\partial^2 \psi}{\partial z^2} \quad \dots (2)$$

$\therefore$ (1) can be written as

$$\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial \psi}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 \psi}{\partial \theta^2} + \frac{\partial^2 \psi}{\partial z^2} - \mu \epsilon \frac{\partial^2 \psi}{\partial t^2} - \mu \sigma \frac{\partial \psi}{\partial t} = 0 \quad \dots (3)$$

The elementary harmonic solutions of this equation may be expressed in the form

$$\psi = F(r, \theta) e^{\pm i h z - i \omega t} \quad \dots (4)$$

where $F(r, \theta)$ is a solution of

$$\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial F_1}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 F}{\partial \theta^2} + (k^2 - h^2) F = 0 \quad \dots (5)$$

which may be easily separated by writing

$$F(r, \theta) = F_1(r) F_2(\theta) \quad \dots (6)$$

where $F_1(r)$ and $F_2(\theta)$ are arbitrary solutions of the ordinary differential equations

$$r \frac{d}{dr} \left(r \frac{dF_1}{dr} \right) + \{ (k^2 - h^2) r^2 - n^2 \} F_1 = 0, \quad \dots (7)$$

and

$$\frac{d^2 F_2}{d\theta^2} + n^2 F_2 = 0 \quad \dots (8)$$

where n and h are separation constants and may be chosen in accordance with the physical requirements.

The solutions of equation (7) which is satisfied by radial function $F_1(r)$ being Bessel's functions which are generally reserved for that particular solution $J_n(r\sqrt{k^2-h^2})$ which is finite on the axis $r=0$, we use the name *circular cylinder function* for any particular solution of (7) and denote it by $F_1 = Z_n(r\sqrt{k^2-h^2})$ where n is the order of the function with argument $r\sqrt{k^2-h^2}$.

As such the particular solutions of (1), being periodic in t and θ , may be constructed from elementary waves of the form

$$\psi_m = e^{im\theta} Z_m(r\sqrt{k^2-h^2}) e^{\pm i h z - i \omega t}$$

where the propagation constant h is a complex number.

It follows from (9) that an explicit expression for h in terms of frequency ω and medium-constants can only be found by observing the behaviour of ψ over a cylinder $r=\text{constant}$ or on a plane $z=\text{constant}$.

Now to study the properties of the function $Z_n(r\sqrt{k^2-h^2})$, putting $\lambda = r\sqrt{k^2-h^2}$, we observe that $Z_n(\lambda)$ satisfies the equation

$$\frac{d^2 Z_n}{d\lambda^2} + \frac{1}{\lambda} \frac{dZ_n}{d\lambda} + \left(1 - \frac{n^2}{\lambda^2} \right) Z_n = 0 \quad \dots (10)$$

which is characterized by a regular singularity at $\lambda=0$ and an essential singularity at $\lambda=\infty$.

Clearly the cylinder function or the Bessel function of first kind i.e., $J_n(\lambda)$ is a particular solution of (10) and it is finite at $\lambda=0$. As such Bessel function may be

expanded in a series of ascending powers of λ and this series is convergent, for all finite values of the argument λ .

$$\text{Here } J_n(\lambda) = \sum_{s=0}^{\infty} \frac{(-1)^s}{s! \Gamma(n+s+1)} \left(\frac{\lambda}{2} \right)^{n+2s} \quad \dots (11)$$

Another solution in terms of Bessel function of the second kind defined by

$$N_n(\lambda) = \frac{1}{\sin n\pi} [J_n(\lambda) \cos n\pi - J_{-n}(\lambda)] \quad \dots (12)$$

may also be obtained.

Problem 50. Distinguish clearly between the phase and the group velocities. Find the relation between them. (Agra, 1975, 1976)

It is generally observed that the notion of phase velocity is only applicable to fields which are periodic in space i.e., the fields representing wave trains of infinite duration. Denoting by $\psi(z, t)$ the state of medium, such that

$$\phi(z, t) = A e^{ikz - i\omega t} \quad \dots (1)$$

the surfaces of constant phase or state may be defined by

$$kz - \omega t = \text{constant}$$

and the velocity of propagation of these surfaces is given by

$$v = \frac{dz}{dt} = \frac{\omega}{k} \text{ by (2)} \quad \dots (3)$$

where v is termed as *phase velocity*.

Now to discuss the concept of group velocity; consider first the superposition of two harmonic waves differing very slightly in frequency and wave number, such that

$$\psi_1 = \cos(kz - \omega t) \quad \dots (4)$$

$$\psi_2 = \cos[(k+\delta k)z - (\omega+\delta\omega)t] \quad \dots (5)$$

The resultant of (4) and (5) is

$$\begin{aligned} \psi &= \psi_1 + \psi_2 = \cos(kz - \omega t) + \cos[(k+\delta k)z - (\omega+\delta\omega)t] \\ &= 2 \cos \frac{1}{2}(z\delta k - t\delta\omega) \cos[(k+\frac{1}{2}\delta k)z - (\omega+\frac{1}{2}\delta\omega)t] \end{aligned} \quad \dots (6)$$

which is an expression for the phenomenon of 'beats'.

It follows from (6) that the field oscillates at a frequency negligibly different from ω , with its effective amplitude

$$a = 2 \cos \frac{1}{2}(z\delta k - t\delta\omega) \quad \dots (7)$$

varying slowly between the algebraic sum of amplitudes of the component-waves and zero.

Due to constructive and destructive interference, the field distribution along time and space axes forms periodically repeated 'beats' or 'groups', whence the surfaces over which the group amplitude 'a' is constant are given by

$$z\delta k - t\delta\omega = \text{constant}$$

yielding the *group velocity* as

$$u = \frac{z}{t} = \frac{\delta\omega}{\delta k} \quad \dots (8)$$

which follows that group velocity is the ratio of the difference of frequency to the difference of wave number.

In case the medium is non-dispersive, $\delta k = \frac{1}{v} \delta\omega$, so that the group velocity coincides with the phase velocity v . Of course, in a dispersive medium the group velocity and the phase velocity are quite different.

Conclusively the group velocity u differs from the phase velocity v in a dispersive medium, and if the dispersion is normal then $u < v$ while if the dispersion

is anomalous then $u > v$. Also in the neighbourhood of an absorption band the group velocity u may become infinite or even negative.

Problem 51. Find the solution of the equation

$$\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} + \frac{\partial^2 \psi}{\partial z^2} = -\phi(x, y, z)$$

where $\phi(x, y, z)$ is a known function. Discuss the uniqueness of the solution.

(Agra, 1971)

Hint. For its solution see § 1.59 where ρ has been regarded as a function of x, y, z .

Problem 52. Obtain the differential equation for the vibrations of a string under tension, solve the equation and give the expression for the velocity of the waves in the string.

(Agra, 1971)

Problem 53. (a) Discuss the modes of vibrations of a circular membrane.

(b) Solve the equation,

$$\frac{\partial V}{\partial t} = h^2 \frac{\partial^2 u}{\partial x^2}$$

subject to initial conditions $v = F(x)$ at $t = 0$.

(Rohilkhand, 1971)

Problem 54. Using the method of separation of variables, solve Laplace equation to find the temperature inside and outside a sphere when its boundary is at the finite temperature $\psi = f(\theta)$.

Find out the permanent temperature within a solid sphere of radius unity when one half of the surface of the sphere is kept at constant temperature 0°C and the other half of its surface is at 1°C .

(Rohilkhand, 1971)

See § 12.6, 12.7 and 12.8 (see Problem 21).

Problem 55. Derive the differential equation for the vibrations of a circular membrane and by solving it show that the allowed angular frequencies of vibrations

are determined by the equation $J_n \left[a \left(\frac{m\omega^2}{T} \right)^{1/2} \right] = 0$, $n = 0, 1, 2, \dots$

where a is the radius of the membrane, m is the mass per unit area of the membrane and T is the tension of unit length.

(Rohilkhand, 1971)

See § 12.16 [B].

Problem 56. Discuss the method of separation of variables for solving two dimensional Laplace equation in spherical coordinate system and hence find the distribution of the electro-static potential produced by conducting ring carrying a total electric charge.

(Rohilkhand, 1971)

See § 12.7 [B].

CHAPTER 13

MAXWELL'S ELECTRO-MAGNETIC FIELD EQUATIONS

13.1. INTRODUCTION

It was due to Clark Maxwell that a change in the electric displacement vector was supposed to be equivalent to an electric current known as *displacement current*. The total flow obtained by the resultant of *displacement current* and *conduction current* (which arises out of the actual motion of electrons or electric charges) was supposed to be such that the total flow into any surface would invariably be zero.

In fact the equation of electromagnetic field have not yet been finally established, but we have used the fact that when there is a flow of steady current, then

$$\text{Curl } \mathbf{H} = 4\pi \mathbf{j} \quad \dots (1)$$

where $\mathbf{H}$ is the negative field and $\mathbf{j}$ is the current vector. In case of those electromagnetic fields where the flow of current is unsteady, the equation (1) is insufficient and requires modification for unsteady flow of currents.

$$\text{Since we know that } \text{div curl } \mathbf{H} = 0 \quad \dots (2)$$

$$\therefore (1) \text{ yields, } \text{div } \mathbf{j} = 0 \quad \dots (3)$$

which is always true in case of steady flow of currents.

But in case of unsteady flow of currents when generally $\text{div } \mathbf{j} \neq 0$, the equation of continuity of charge taken in mixed units, is given by

$$\text{div } \mathbf{j} = -\frac{1}{c} \frac{\partial \rho}{\partial t} = -\frac{1}{4\pi c} \left(\frac{\partial}{\partial t} \text{div } \mathbf{D} \right) \text{ where } \text{div } \mathbf{D} = 4\pi \rho \quad \dots (4)$$

$$\text{or } \text{div } \left(\mathbf{j} + \frac{1}{4\pi c} \frac{\partial \mathbf{D}}{\partial t} \right) = 0 \quad \dots (5)$$

$$\text{or } \text{div } \mathbf{J} = 0 \text{ where } \mathbf{J} = \mathbf{j} + \frac{1}{4\pi c} \frac{\partial \mathbf{D}}{\partial t} \quad \dots (6)$$

Here ρ is the free charge density at a scalar, point, $\mathbf{D}$ is the Maxwell's electric displacement vector and $\mathbf{J}$ is called *Maxwell's total current*.

Since the operators div and $\frac{\partial}{\partial t}$ are commutative, it therefore follows that $\mathbf{J}$ is a quantity whose divergence is always zero and in case of steady current flow $\mathbf{J}$ reduces to $\mathbf{j}$. Conclusively in (1), $\mathbf{j}$ should be replaced by $\mathbf{J}$ in order that the result (1) will be valid for steady as well as unsteady current flows. We call $\mathbf{j}$ as *conduction current* and the

part $\frac{1}{4\pi c} \frac{\partial \mathbf{D}}{\partial t}$ of $\mathbf{J}$ as *displacement current*.

If $\mathbf{P}$ be the *polarisation* of the medium, then we have

$$\mathbf{D} = \mathbf{E} + 4\pi \mathbf{P}, \mathbf{E} \text{ being electric field} \quad \dots (7)$$

$$\text{As such (6) yields } \mathbf{J} = \mathbf{j} + \frac{1}{c} \frac{\partial \mathbf{P}}{\partial t} + \frac{1}{4\pi c} \frac{\partial \mathbf{E}}{\partial t} \quad \dots (8)$$

The part of $\frac{1}{4\pi c} \frac{\partial \mathbf{D}}{\partial t}$ coming from $\frac{\partial \mathbf{P}}{\partial t}$ is known as *Polarisation current* and that from $\frac{1}{4\pi c} \frac{\partial \mathbf{E}}{\partial t}$ the *ether displacement current* or *displacement current in vacuum*.

The modification of (1) is supported by experiments and the electromagnetic theory of light based on Maxwell's electromagnetic field equations, is given a sound theoretical basis by replacing $\mathbf{j}$ by $\mathbf{J}$ in (1) whence we get

$$\text{Curl } \mathbf{H} = 4\pi \mathbf{J} = 4\pi \mathbf{j} + \frac{1}{c} \frac{\partial \mathbf{D}}{\partial t} \quad \dots (9)$$

where $\mathbf{j}$ and $\mathbf{H}$ are measured in e.m.u and $\mathbf{D}$ in e.s.u.
and $c = 3 \times 10^{10} \text{ cm/sec.}$

Here the term $\frac{1}{c} \frac{\partial \mathbf{D}}{\partial t}$ assumes importance only if $\frac{\partial \mathbf{D}}{\partial t}$ is of the order of c , otherwise it may be neglected if $\frac{\partial \mathbf{D}}{\partial t}$ is small as compared to c . Those unsteady states in which the last term in R.H.S. of (9) may be neglected are known as *quasi-unsteady states*. Dealing with very high frequencies as compared to that of light, the displacement current has considerable significance.

If we now introduce $\mathbf{B}$ as magnetic induction of the medium, given by

$$\mathbf{B} = \mu \mathbf{H} \quad \dots (10)$$

$$\text{then (9) reduces to } \text{curl } \mathbf{B} = 4\pi \mu \mathbf{j} + \frac{\mu}{c} \frac{\partial \mathbf{D}}{\partial t} \quad \dots (11)$$

Using Gauss's, divergence theorem, it follow from (5),

$$\oint \left(\mathbf{j} + \frac{1}{4\pi c} \frac{\partial \mathbf{D}}{\partial t} \right) \cdot d\mathbf{S} = 0 \quad \dots (12)$$

13.2. MAXWELL'S EQUATIONS FOR ELECTROMAGNETIC FIELD

As already discussed in §13.1, taking Gaussian units or the mixed system of units such that the quantities $\mathbf{D}$, $\mathbf{E}$, ρ with allied quantities are measured in e.s.u. and the quantities $\mathbf{B}$, $\mathbf{H}$, $\mathbf{j}$ with allied quantities in e.m.u, the differential or local form of the electromagnetic field equations of Maxwell in accordance with the elementary laws of electricity and magnetism laid by Gauss, Faraday, Ampere and others can be summarised as follows:

$$\text{Divergence equations } \text{div } \mathbf{D} = \nabla \cdot \mathbf{D} = 4\pi \rho \quad \dots (1)$$

$$\text{div } \mathbf{B} = \nabla \cdot \mathbf{B} = 0 \quad \dots (2)$$

$$\text{Circuital equations } \text{curl } \mathbf{H} = \nabla \times \mathbf{H} = 4\pi \mathbf{j} + \frac{1}{c} \frac{\partial \mathbf{D}}{\partial t} \quad \dots (3)$$

$$\text{curl } \mathbf{E} = \nabla \times \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} \quad \dots (4)$$

where the equation (4) is the generalized law of induction.

These equations are considered in association of the following *subsidiary relations* or macroscopic constitutive equations

$$\mathbf{D} = K \mathbf{E} \quad \dots (5)$$

$$\mathbf{B} = \mu \mathbf{H} \quad \dots (6)$$

$$\mathbf{j} = \sigma \mathbf{E} \quad \dots (7)$$

where K is the dielectric constant, μ is the permeability of the medium and σ is the conductivity of the medium. Here $\mathbf{j}$, σ , $\mathbf{E}$ are measured in the mixed units, but if $\mathbf{j}$ is in e.m.u. and σ , $\mathbf{E}$ in e.s.u. then we have $\mathbf{j} = \frac{\sigma \mathbf{E}}{c}$, also if $\mathbf{j}$, σ are in e.m.u. and $\mathbf{E}$ in e.s.u., then $\mathbf{j} = c\sigma \mathbf{E}$.

In addition to the above, few other quantities such as the scalar potential ϕ and magnetic vector potential $\mathbf{A}$ defined by

$$\mathbf{B} = \text{curl } \mathbf{A} \quad \dots (8)$$

$$\mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{A}}{\partial t} - \nabla \phi \quad \dots (9)$$

are also required.

Since (8) does not provide the unique definition of $\mathbf{A}$, therefore we supplement it by $\text{div } \mathbf{A} = 0$ (in steady states) and

$$\text{div } \mathbf{A} + \frac{\mu}{c} K \frac{\partial \phi}{\partial t} = 0 \quad \dots (10)$$

Here ϕ is in e.s.u. and $\mathbf{A}$ in e.m.u.

13.3. TO SHOW $\nabla \cdot \mathbf{D} = 4\pi \rho$.

We know that the molecules of dielectric substance are formed of charged entities, such as atomic nuclei and electrons and hence when dielectric substances are inserted in the field of electromagnetic waves, the charged entities are displaced in opposite directions of their mean position *i.e.*, the dielectric is said to be polarised. Such polarised charges form an electric dipole. Let $\mathbf{r}$ be the separation between positive and negative charges in each dipole and δq the magnitude of each displaced charge; then the electric moment say $d\mathbf{p}$ of dipole being $\mathbf{r} \delta q$ is defined over the elementary volume δV as

$$d\mathbf{p} = \int_{\delta V} \mathbf{r} dq \quad \dots (1)$$

Let us introduce the polarisation vector $\mathbf{P}$ as the resultant of the electric dipole moment of molecules per unit volume *i.e.*,

$$\mathbf{P} = \lim_{\delta V \rightarrow 0} \frac{d\mathbf{p}}{\delta V} \quad \dots (2)$$

Now representing the charge density of the charge caused by polarisation of the dielectric, by ρ_P , the charge carried out of the surface S enclosing the volume V in the electric field of E , is given by

$-\int_V \rho_P dV$. Also the net charge diverging out of the elementary volume δV in this dielectric being $\text{div } \mathbf{P} \delta V$, the charge carried out of the volume V is $\int_V \text{div } \mathbf{P} dV = 0$. We thus have

$$\int_V \text{div } \mathbf{P} dV = - \int_V \rho_P dV \quad \dots (1)$$

Using Gauss's divergence theorem, this yields,

$$\int_S \mathbf{P} \cdot \mathbf{n} dS = - \int_V \rho_P dV \quad \dots (2)$$

where $\mathbf{n}$ is the unit outward drawn normal to the surface element δS .

But according to Gauss's theorem of electrostatics, the total normal outward electric field flux across any arbitrary closed surface is 4 times the charge enclosed by it i.e.,

$$\int_S \mathbf{E} \cdot \mathbf{n} dS = 4\pi \cdot (\text{total charge enclosed in volume } V) \quad \dots (3)$$

Now, there are two types of charges inside V .

(i) the free charge of density ρ and (ii) the polarised charge of density ρ_P .

$$\begin{aligned} \therefore \int_S \mathbf{E} \cdot \mathbf{n} dS &= 4\pi \left[\int_V \rho dV + \int_V \rho_P dV \right] \\ &= 4\pi \left[\int_V \rho dV - \int_S \mathbf{P} \cdot \mathbf{n} dS \right] \text{ by (2)} \end{aligned}$$

$$\text{or } \int_S (\mathbf{E} + 4\pi \mathbf{P}) \cdot \mathbf{n} dS = 4\pi \int_V \rho dV$$

$$\text{or } \int_S \mathbf{D} \cdot \mathbf{n} dS = 4\pi \int_V \rho dV \text{ by (7) of §13.1} \quad \dots (4)$$

$$\text{Hence by Gauss's divergence theorem, } \int_V \text{div } \mathbf{D} dV = \int_V 4\pi \rho dV$$

$$\text{which yields } \text{div } \mathbf{D} = 4\pi \rho \quad \dots (5)$$

Note 1. In case of an ordinary dielectric i.e. a dielectric free from charge, $\rho = 0$ and hence $\text{div } \mathbf{D} = \nabla \cdot \mathbf{D} = 0$

Note 2. By (7) of §13.1, we have $\mathbf{D} = \mathbf{E} + 4\pi \mathbf{P} = \mathbf{E} \left[1 + 4\pi \frac{\mathbf{P}}{\mathbf{E}} \right]$

$$\text{which follows that } \mathbf{D} = K\mathbf{E} \quad \dots (6)$$

where $K = 1 + 4\pi \frac{\mathbf{P}}{\mathbf{E}}$ and known as dielectric constant of the medium

13.4. TO SHOW $\text{DIV } \mathbf{B} = 0$

Since the lines of force are either closed or go off to infinity, therefore the surface integral of magnetic induction $\mathbf{B}$ over a closed surface is

zero, so that (6) of §13.3 reduces here

$$\int_S \mathbf{B} \cdot \mathbf{n} dS = 0 \text{ or } \text{div } \mathbf{B} = \nabla \cdot \mathbf{B} = 0 \quad \dots (1)$$

$$\text{Note. Since } \mathbf{B} = \mu \mathbf{H} \quad \therefore (11) \text{ yields } \nabla \cdot \mathbf{H} = 0 \quad \dots (2)$$

13.5. TO SHOW $\text{CURL } \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t}$

According to Faraday's law of electromagnetic induction the induced E.M.F. around a closed circuit is negative times the rate of change of total flux ϕ of the magnetic induction $\mathbf{B}$ through the circuit. Induced E.M.F.

$$= -\frac{1}{c} \frac{\partial \phi}{\partial t} \quad \dots (1)$$

Also if $\mathbf{E}$ be the electric field in the direction $d\mathbf{l}$, then

$$\text{E.M.F.} = \sum \mathbf{E} \cdot d\mathbf{l} \quad \dots (2)$$

If we consider an elementary rectangle $PQRS$ with sides δy and δz as shown in Fig 13.1, then around it we have

$$\begin{aligned} \text{E.M.F.} &= \frac{\text{work}}{\text{charge}} = \sum \mathbf{E} \cdot d\mathbf{l} \\ &= [E_z(y + dy) - E_z(y)] dz - [E_y(z + dz) - E_y(z)] dy \\ &= \frac{\partial E_z}{\partial y} dy dz - \frac{\partial E_y}{\partial z} dy dz \quad \dots (3) \end{aligned}$$

$$\therefore E_z(y + dy) = E_z(y) + \frac{\partial E_z}{\partial y} dy \text{ by Taylor's theorem}$$

$$\text{Also } \sum \mathbf{E} \cdot d\mathbf{l} = -\frac{1}{c} \frac{d\phi_z}{dt} = -\frac{1}{c} \frac{\partial B_z}{\partial t} dy dz \quad \dots (4)$$

Hence from (3) and (4) equating the two values of E.M.F., we have

$$-\frac{1}{c} \frac{\partial B_z}{\partial t} dy dz = \frac{\partial E_z}{\partial y} dy dz - \frac{\partial E_y}{\partial z} dy dz = \text{curl}_x \mathbf{E}$$

$$\text{Similarly } -\frac{1}{c} \frac{\partial B_y}{\partial t} = \text{curl}_z \mathbf{E} \text{ and } -\frac{1}{c} \frac{\partial B_x}{\partial t} = \text{curl}_y \mathbf{E}$$

where B_x, B_y, B_z are components of $\mathbf{B}$ along principal axes.

Combining the last three relations we can write

$$-\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} = \text{curl } \mathbf{E} \text{ i.e., } \nabla \times \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} \quad \dots (5)$$

Note. In case of free space when $\mu = 1$ and $\mathbf{B} = \mathbf{H}$, (5) yields

$$\text{Curl } \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{H}}{\partial t} \quad \dots (6)$$




Fig. 13.1

13.6. PHYSICAL INTERPRETATION OF MAXWELL'S EQUATIONS

Consider $\text{div } \mathbf{D} = 4\pi\rho$, which may be written as

$\int \text{div } \mathbf{D} dV = 4\pi \int \rho dV$ i.e., $\int_S \mathbf{D} \cdot d\mathbf{S} = 4\pi \int_V \rho dV$ by Gauss' divergence theorem.

It follows, Gauss's theorem i.e., the flux of the displacement vector $\mathbf{D}$ across any closed surface S is directly proportional to the total electric charge in V .

The equation $\text{div } \mathbf{B} = 0$, may be written as

$\int_S \mathbf{B} \cdot d\mathbf{S} = 0$, by Gauss's divergence theorem

which follows: Magnetic flux theorem i.e., the total normal magnetic induction across any closed surface S is zero

Also the equation $\text{curl } \mathbf{H} = 4\pi\mathbf{j} + \frac{1}{c} \frac{\partial \mathbf{D}}{\partial t}$ can be expressed by Stoke's theorem, as

$$\begin{aligned} \int_S \text{curl } \mathbf{H} \cdot d\mathbf{S} &= 4\pi \int_S \left(\mathbf{j} + \frac{1}{4\pi c} \frac{\partial \mathbf{D}}{\partial t} \right) \cdot d\mathbf{S} \text{ or } \int_C \mathbf{H} \cdot d\mathbf{l} \\ &= 4\pi \int_S \left(\mathbf{j} + \frac{1}{4\pi c} \frac{\partial \mathbf{D}}{\partial t} \right) \cdot d\mathbf{S} \end{aligned}$$

which is Ampere's generalized circuital relation i.e., the work done in carrying a unit pole round a closed circuit C is equal to 4π times the total current in the circuit.

Lastly the equation $\text{curl } \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t}$ can be written by the help of Stoke's theorem, as

$$\int \text{curl } \mathbf{E} \cdot d\mathbf{S} = \int \left(-\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} \right) \cdot d\mathbf{S} \text{ or } \int_C \mathbf{E} \cdot d\mathbf{l} = -\frac{1}{c} \int \frac{\partial \mathbf{B}}{\partial t} \cdot d\mathbf{S}$$

which is Generalized law of electromagnetic induction i.e., the total electromagnetic force around a closed circuit C is equal to $-\frac{1}{c}$ times the rate of change of magnetic induction through C

13.7. DECAYING OF FREE CHARGE

Consider $\text{curl } \mathbf{H} = 4\pi\mathbf{j} + \frac{\partial \mathbf{D}}{\partial t}$ and $\mathbf{j} = \sigma\mathbf{E}$ in e.s.u. or e.m.u. throughout.

Elimination of $\mathbf{j}$ yields, $\text{curl } \mathbf{H} = 4\pi\sigma\mathbf{E} + \frac{\partial \mathbf{D}}{\partial t} = \frac{4\pi\sigma\mathbf{D}}{K} + \frac{1}{c} \frac{\partial \mathbf{D}}{\partial t}$ $\therefore \mathbf{D} = K\mathbf{E}$

$$\therefore \text{div } \text{curl } \mathbf{H} = \text{div} \left(\frac{4\pi\sigma\mathbf{D}}{K} + \frac{1}{c} \frac{\partial \mathbf{D}}{\partial t} \right) = 0$$

which with the help of $\mathbf{D} = 4\pi\rho$, gives $\frac{4\pi\sigma\rho}{K} + \frac{\partial\rho}{\partial t} = 0$ or $\frac{d\rho}{\rho} = -\frac{4\pi\sigma}{K} dt$

It yields on integration, $\rho = \rho_0 e^{-t/T}$... (1)

where ρ_0 is the charge at any point $t=0$ and $T = \frac{K}{4\pi\sigma}$, T being known as time of relaxation.

It follows from (1) that charge dies out exponentially at a rate quite independent of any other electro-magnetic phenomena taking place simultaneously. In most ordinary case we take $\rho=0$.

13.8. POYNTING VECTOR

Since the electromagnetic effects in a field are completely attributed to charges at rest or in motion and a charge moving in a field experiences a force so that the work is done on the moving charge by the forces of the field, we therefore claim to determine the rate of change of energy on account of the interaction of the field and the charges in it.

Let $\mathbf{E}$ be the electric field and $\mathbf{B}$ the magnetic induction at a point in the field occupied by a charge moving with velocity $\mathbf{v}$. Also let ρ be the charge density and dV an element of volume at the point under consideration. Then the force on a charge ρdV (charge on the elementary volume), is $\left[\mathbf{E} + \frac{\mathbf{v} \times \mathbf{B}}{c} \right] \rho dV$.

$$\begin{aligned} \text{The rate of work done on this charge} &= \left[\mathbf{E} + \frac{\mathbf{v} \times \mathbf{B}}{c} \right] \cdot \mathbf{v} \rho dV \\ &= (\mathbf{E} \cdot \mathbf{v}) \rho dV \quad \because (\mathbf{v} \times \mathbf{B}) \cdot \mathbf{v} = 0. \end{aligned}$$

$\therefore$ the total rate of work done by the forces of the field in the region under consideration,

$$\begin{aligned} &= \int_V (\mathbf{E} \cdot \mathbf{v}) \rho dV = c \int_V (\mathbf{E} \cdot \mathbf{j}) dV \text{ in mixed units, since } \rho\mathbf{v} = \mathbf{j}c \\ &= c \int_V \mathbf{E} \cdot \left(\frac{1}{4\pi} \text{curl } \mathbf{H} - \frac{1}{4\pi c} \frac{\partial \mathbf{D}}{\partial t} \right) dV \\ &\quad \because \text{by Maxwell's equations, } \mathbf{j} = \frac{1}{4\pi} \text{curl } \mathbf{H} - \frac{1}{4\pi c} \frac{\partial \mathbf{D}}{\partial t} \\ &= \frac{c}{4\pi} \int_V (\mathbf{E} \cdot \text{curl } \mathbf{H}) dV - \frac{1}{4\pi} \int_V \left(\mathbf{E} \cdot \frac{\partial \mathbf{D}}{\partial t} \right) dV \\ &= \frac{c}{4\pi} \int_V \{ \text{div} (\mathbf{H} \times \mathbf{E}) + \mathbf{H} \cdot \text{curl } \mathbf{E} \} dV - \frac{1}{4\pi} \int_V \left(\mathbf{E} \cdot \frac{\partial \mathbf{D}}{\partial t} \right) dV \\ &\quad \because \mathbf{E} \cdot \text{curl } \mathbf{H} = \text{div} (\mathbf{H} \times \mathbf{E}) + \mathbf{H} \cdot \text{curl } \mathbf{E} \\ &= \frac{c}{4\pi} \int_V \text{div} (\mathbf{H} \times \mathbf{E}) dV + \frac{c}{4\pi} \int_V (\mathbf{H} \cdot \text{curl } \mathbf{E}) dV - \frac{1}{4\pi} \int_V \left(\mathbf{E} \cdot \frac{\partial \mathbf{D}}{\partial t} \right) dV \end{aligned}$$

Hence using the Gauss's divergence theorem and Maxwell's equation $\text{curl } \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t}$, we have the rate of change of energy

$$= \frac{c}{4\pi} \int_S (\mathbf{H} \times \mathbf{E}) \cdot d\mathbf{S} - \frac{1}{4\pi} \int_V \left(\mathbf{H} \cdot \frac{\partial \mathbf{B}}{\partial t} + \mathbf{E} \cdot \frac{\partial \mathbf{D}}{\partial t} \right) dV \quad \dots (1)$$

Regarding μ and K as independent of time, we can write (1) as

$$-\frac{\partial}{\partial t} \int_V \frac{1}{8\pi} (\mathbf{B} \cdot \mathbf{H} + \mathbf{E} \cdot \mathbf{D}) dV = \frac{c}{4\pi} \int_S (\mathbf{E} \times \mathbf{X}) \cdot d\mathbf{S} + \text{Rate at which field is doing work}$$

$$\therefore \frac{\partial}{\partial t} (\mathbf{B} \cdot \mathbf{H}) = 2\mathbf{H} \cdot \frac{\partial \mathbf{B}}{\partial t} \text{ and } \frac{\partial}{\partial t} (\mathbf{E} \cdot \mathbf{D}) = 2\mathbf{E} \cdot \frac{\partial \mathbf{D}}{\partial t} \quad \dots (2)$$

We conclude from (2) that the electromagnetic energy of the field per unit volume must be taken as $\frac{1}{8\pi} (\mathbf{B} \cdot \mathbf{H} + \mathbf{E} \cdot \mathbf{D}) \quad \dots (3)$

Then the equation (2) expresses the principle of conservation of energy if the first term on the right of (2) represents a rate of flow of radiation-energy outwards across S , the surface enclosing V .

$$\text{The vector (say) } \vec{\Pi} = \frac{c}{4\pi} (\mathbf{E} \times \mathbf{H}) \quad \dots (4)$$

is known as *Radiant vector* or *Poynting vector* and it gives the rate at which the energy is radiated across unit area.

It is notable that $\vec{\Pi}$ is normal to the plane of $\mathbf{E}$ and $\mathbf{H}$ and it is not unique since $\vec{\Pi} + \mathbf{A}$ also satisfies (2) because $\nabla \cdot \mathbf{A} = 0$. Also the flux of $\vec{\Pi}$ across a closed surface is only significant.

13.9. POYNTING THEOREM

The rate at which the electric energy in any given region is increasing is equal to the Integral $\frac{c}{4\pi} \iint [l(\gamma Y - \beta Z) + m(\alpha Z - \gamma X) + n(\beta X - \alpha Y)] dS$ taken over the boundary of the region, l, m, n being the direction cosines of the inward normal to the surface element dS ; α, β, γ , being the components of magnetic force and X, Y, Z being the components of electric force.

Assuming that the electro-magnetic energy is not confined to the regions occupied by electric charges, magnets and currents but spreads over the whole space, the magnetic kinetic energy T and electric potential energy W of an isotropic medium are given by

$$T = \frac{1}{8\pi} \iiint \mu (\alpha^2 + \beta^2 + \gamma^2) dx dy dz \quad \dots (1)$$

$$W = \frac{1}{8\pi} \iiint K (X^2 + Y^2 + Z^2) dx dy dz \quad \dots (2)$$

Assuming that the energy is localized in a medium, the total energy in any closed region is,

$$T + W = \iiint \left\{ \frac{K}{8\pi} (X^2 + Y^2 + Z^2) + \frac{\mu}{8\pi} (\alpha^2 + \beta^2 + \gamma^2) \right\} dx dy dz \quad \dots (3)$$

Differentiating it and replacing $\mu\alpha$, by α , KX by $4\pi f$, etc., we get

$$\frac{d}{dt} [T + W] = \iiint \left[X \frac{\partial f}{\partial t} + Y \frac{\partial g}{\partial t} + Z \frac{\partial h}{\partial t} + \frac{1}{4\pi} \left(\alpha \frac{d\alpha}{dt} + \beta \frac{d\beta}{dt} + \gamma \frac{d\gamma}{dt} \right) \right] dx dy dz$$

$$= \frac{c}{4\pi} \iiint \left[X \left(\frac{\partial \gamma}{\partial y} - \frac{\partial \beta}{\partial z} \right) + \dots - \alpha \left(\frac{\partial Z}{\partial y} - \frac{\partial Y}{\partial z} \right) - \dots \right] dx dy dz - c \iiint (uX + vY + wZ) dx dy dz \quad \dots (4)$$

where

$$-\frac{1}{c} \frac{d\alpha}{dt} = \frac{\partial Z}{\partial t} - \frac{\partial Y}{\partial z} \text{ etc.,}$$

and

$$4\pi \left(u + \frac{1}{c} \frac{\partial f}{\partial t} \right) = \frac{\partial \gamma}{\partial y} - \frac{\partial \beta}{\partial z} \text{ etc.}$$

Here u, v, w are components of ordinary current at any point, which is produced by the moving electric charges.

The last term on the right of (4) gives exactly the rate at which work is done or the energy is dissipated by the flow of currents so that the first part of this must represent the rate at which energy flows into the region from outside.

Hence using Green's theorem, the first part on the right of (4), yields

$$\frac{c}{4\pi} \iiint \left[\frac{\partial}{\partial x} (Z\beta - Y\gamma) + \frac{\partial}{\partial y} (Y\gamma - Z\alpha) + \frac{\partial}{\partial z} (Y\alpha - X\beta) \right] dx dy dz = -\frac{c}{4\pi} \iint [l(Z\beta - Y\gamma) + m(Y\gamma - Z\alpha) + n(Y\alpha - X\beta)] dS.$$

Problem 1. If $\mathbf{Z}$ be a vector satisfying the equation

$$\nabla^2 \mathbf{Z} = \frac{4\pi\sigma\mu}{c} \frac{\partial \mathbf{Z}}{\partial t} + \frac{K\mu}{c^2} \frac{\partial^2 \mathbf{Z}}{\partial t^2} \quad \dots (1)$$

then show that the field may be defined by the equations:

$$\mathbf{A} = 4\pi\sigma\mu\mathbf{Z} + \frac{K\mu}{c} \frac{\partial \mathbf{Z}}{\partial t} \quad \dots (2)$$

$$\phi = -\text{div } \mathbf{Z}, \quad \dots (3)$$

$$\mathbf{E} = \text{curl curl } \mathbf{Z}, \quad \dots (4)$$

$$\mathbf{B} = \text{curl } \mathbf{A} \quad \dots (5)$$

We have from (2) and (3) $\text{div } \mathbf{A} + 4\pi\sigma\mu\phi + \frac{K\mu}{c} \frac{\partial \phi}{\partial t} = 0$

which is one of the Maxwell's equations.

$$\begin{aligned} \text{Again from (4), we have } \mathbf{E} &= \text{grad}^2 \mathbf{Z} - \nabla^2 \mathbf{Z} \\ &= -\text{grad } \phi - \frac{\partial}{\partial t} \left(\frac{4\pi\sigma\mu}{c} \mathbf{Z} + \frac{K\mu}{c^2} \frac{\partial \mathbf{Z}}{\partial t} \right) \text{ by (1) and (3)} \\ &= -\text{grad } \phi - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t} \end{aligned}$$

which is another Maxwell's equation.

Also $\text{curl } \mathbf{B} = \text{curl}(\text{curl } \mathbf{A})$ by (5)

$$= \text{curl curl} \left(4\pi\sigma\mu\mathbf{Z} + \frac{K\mu}{c} \frac{\partial \mathbf{Z}}{\partial t} \right) \text{ by (2)}$$

$$= 4\pi\sigma\mu (\text{curl curl } \mathbf{Z}) + \left(\frac{K\mu}{c} \right) \frac{\partial}{\partial t} (\text{curl curl } \mathbf{Z})$$

$$= 4\pi\sigma\mu \mathbf{E} + \frac{K\mu}{c} \frac{\partial \mathbf{E}}{\partial t} \text{ by (4)}$$

$$= 4\pi\mu\mathbf{j} + \frac{\mu}{c} \frac{\partial \mathbf{D}}{\partial t}$$

$$\therefore \sigma\mathbf{E} = \mathbf{j} \text{ and } k\mathbf{E} = \mathbf{D}$$

This is also one of the Maxwell's equations.

Hence the given equations define the field under consideration.

Problem 2. Initially the electric and magnetic fields in free space are $E_y(x, 0) = \sin ax$, $H_z(x, 0) = \sin ax$. Find E_y as function of time.

We have discussed in the previous chapter that the most general solution of one-dimensional wave equation is

$$E_y(x, t) = f(x+ct) + g(x-ct) \quad \dots (1)$$

$$E_z(x, t) = -f(x+ct) + g(x-ct) \quad \dots (2)$$

The negative sign being taken since E and H the directions of propagation from a right handed system. This is also justified on applying Maxwell's equation $\text{curl } E = -\frac{1}{c} \frac{\partial H}{\partial t}$.

At time $t=0$, we have $E_y(0, t) = \sin ax = f(x) + g(x)$ by (1)

$E_z(0, t) = \sin ax = -f(x) + g(x)$ by (2)

solving $f(x)=0$, $g(x)=\sin ax$.

Hence $E_y(x, t) = \sin a(x-ct)$ is the required solution.

Problem 3. Show that a plane polarised electromagnetic wave with $H + ke^{pt+in.r}$, satisfies the equation $H=0$ only if n is perpendicular to k .

Take, $H = H_1 i + H_2 j + H_3 k$

$$K = K_1 i + K_2 j + K_3 k$$

$$n = n_1 i + n_2 j + n_3 k$$

and $i = x i + y j + z k$

$$\therefore n.r = n_1 x + n_2 y + n_3 z \text{ and } \frac{\partial H_1}{\partial x} = \frac{\partial}{\partial x} \left\{ K_1 e^{i(pt+in.r)} = n_1 K_1 e^{i(pt+n.r)} \right.$$

$$\text{and } \text{div } H = \left(i \cdot \frac{\partial}{\partial x} + j \cdot \frac{\partial}{\partial y} + k \cdot \frac{\partial}{\partial z} \right) (H_1 i + H_2 j + H_3 k)$$

$$= \frac{\partial H_1}{\partial x} + \frac{\partial H_2}{\partial y} + \frac{\partial H_3}{\partial z} = e^{i(pt+n.r)} \{ n_1 K_1 + n_2 K_2 + n_3 K_3 \}$$

$$= n.K e^{i(pt+n.r)}$$

$\text{div } H = 0 \Rightarrow n.K = 0$ i.e., vector n is $\perp$ to K .

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 4. Show that Maxwell's equations for free space are satisfied by

$$H = \frac{1}{c} \frac{\partial}{\partial t} (\text{grad } \phi \times k); E = -k \left(\frac{1}{c^2} \right) \frac{\partial^2 \phi}{\partial t^2} + \frac{\partial}{\partial z} (\text{grad } \phi)$$

where k is a unit vector along z -axis and ϕ satisfies $\nabla^2 \phi = \frac{1}{c} \frac{\partial^2 \phi}{\partial t^2}$.

Problem 5. If k is a constant vector and ϕ is a scalar function of position and time t , show that $c^2 E = c^2 (k \cdot \nabla) \nabla \phi - k \ddot{\phi}$; $cH = \frac{\partial}{\partial t} (\nabla \phi \times k)$ satisfy Maxwell's equations in free space provided that $\ddot{\phi} = c^2 \nabla^2 \phi$.

CHAPTER 14

SPECIAL THEORY OF RELATIVITY

14.1. INTRODUCTION

We know that all the physical laws deal with the characteristics of certain objects in space in the course of time, while the location of an event or the position of a body is described by a suitable frame of reference which constitutes a conceptual frame work rigidly connected with some material body or a well defined point. Since all the bodies cannot become suitable reference systems, therefore the choice of the reference body always played an important role in the development of science. The heliocentric frame of reference introduced by Galileo was not accepted until the time of Newton who gave a rather comprehensive presentation of it, with the help of his pail experiment. He filled a pail with water and twisted the rope supporting the pail, around its axis, so that the water gradually came into rotation with its plane surface shaped into a paraboloid. Ultimately when the water gained the same speed of rotation as the pail, he stopped the pail so that it gradually and eventually came to rest to resume the shape of a plane. Evidently the surface of water is not influenced by the state of motion of the pail, but the deviation from a plane increases with the deviation from the particular state of motion. This whole process is based on the frame of reference connected with the pail while the angular velocity of the pail is related to a more suitable frame of reference say earth.

There are frames of reference known as *inertial system* with respect to which the law of inertia takes its familiar form i.e. in absence of forces, the space coordinates of a mass point are linear functions of time, whereas there exist another class of frames of reference in which the space coordinates are not linear functions of time. According to laws of mechanics all the frames of reference which are inertial systems are equivalent for describing the nature and for formulating its laws. Such a notion leads to the 'Principle of Relativity'.

The development of Maxwell's electromagnetic field equations was found apparently incompatible with the principle of relativity, according to which electromagnetic waves in empty space should propagate with a constant velocity $c = 3 \times 10^{10}$ cm./sec. But it was not found true with respect to both of the two different inertial systems, moving relatively to each other. Consequently several experiments failed to find such a frame of reference with respect to

which the speed of electromagnetic radiation would be constant in all directions so that *absolute rest absolute motion* etc., might be defined and with respect to which Earth's motion could be determined. One such an experiment is due to Michelson and Morley.

Lorentz conceived the existence of one privileged frame of reference which could not be demonstrated experimentally and he had to introduce several assumptions in the support of his theory. Finally Albert Einstein recognised that the revised fundamental concepts about space and time would be capable of resolving the impasse between theory and experiment. This is that revised concept of Einstein, which is now known as **Special theory of relativity** and which establishes the fundamental equivalence of all inertial systems. According to Einstein there is nothing like *absolute motion*, but all motions are relative and hence the physical laws are independent of the motion of observer. Einstein also ruled out the concept of '*absolute time*' and time varies from one inertial system to another inertial system. He also ruled out the concept of '*absolute space*', since the time is not absolute and therefore the distance between two points measured in two inertial systems cannot be absolute as is evident from the **Galilean Transformation**.

Suppose S, S' are two frames of reference, one at rest and the other moving with uniform velocity v . O, O' are two observers at the origins of S and S' respectively, observing the same event at P whose coordinates are (x, y, z, t) w.r.t. O and (x', y', z', t') w.r.t. O' ; S, S' being supposed to be parallel to each other and their origins being coincident at $t=0, t'=0$. There arise two cases:

Case I. Let the frame S' have the velocity v w.r.t. S in the direction of X' only. Then O' will have velocity v along X' -axis only so that both X and X' axes coincide and consequently the two systems can be combined to each other by the equations

$$x' = x - vt, y' = y, z' = z, t' = t \quad \dots (1)$$

Case II. Let the frame S' have the velocity v along any straight line in any direction. Then if v_x, v_y, v_z be the components of v along




Fig. 14.1




Fig. 14.2

x, y, z , axes, the two systems can be related by the equations

$$x' = x - v_x t, y' = y - v_y t, z' = z - v_z t, t' = t \quad \dots (2)$$

(1) and (2) constitute the **Galilean Transformations**.

In either case the distance being expressed in terms of time, which is not absolute, it is clear that the space is not absolute.

The theory of relativity is studied under two heads:

(1) **Special theory of relativity**. Which deals with inertial systems i.e. systems moving with uniform velocity.

(2) **General theory of relativity**. Which deals with non-inertial systems i.e. systems moving with accelerated velocity.

In this chapter we have only to deal with special theory of relativity.

14.2. BASIC POSTULATES OF SPECIAL THEORY OF RELATIVITY

(Agra, 1969)

There are two basic postulates of special theory of relativity:

I. The principle of relativity. *The natural physical laws preserve or retain their form for all inertial frames i.e. they retain the same form relative to all observers in a state of relative uniform motion.*

This postulate is an extension of the conclusion drawn from Newtonian Mechanics i.e. the velocity is not absolute but it is relative as is evident from the failure of Michelson-Morley experiment performed to determine the velocity of earth through ether.

II. The principle of constancy of velocity of light. *The velocity of light in vacuum is independent of the velocity of observer or the velocity of the source i.e. it has the same value in all inertial frames.*

This postulate is not true according to Galilean transformations, but is verified experimentally that the velocity of light calculated by any means remains constant. It is this postulate, which draws a demarcation line between the classical theory and the theory of relativity given by Einstein. Of course, the constancy of velocity of light requires the following axioms as to introduce the transformation laws:

(i) The velocity of light c must have the same value in all inertial frames.

(ii) The transformations should be linear and approaching to Galilean transformations for low velocities i.e., $v \ll c$.

(iii) The transformation laws should be independent of '*absolute time*' and '*absolute space*' notions.

14.3. LORENTZ TRANSFORMATIONS

(Agra, 1962, 67, Kanpur, 69)

H.A. Lorentz introduced transformation equations relating the observations of position and time taken by two observers in two different inertial frames, in order to satisfy the above axioms.

Take S, S' two inertial frames of reference such that S' is moving with uniform velocity v along X' -axis relative to S . Let O, O' be two observers in two systems, situated at their origins. If we consider the two sets of axes X, Y, Z and X', Y', Z' parallel, then the choice of the origins of the two systems falls in taking their origins coincident at $t=0, t'=0$. It is also convenient to take X, X' axes of two systems coincident, so that the velocity of S' is permanently along X -axis. Let the two observers O, O' observe the same event at P whose coordinates are (x, y, z, t) w.r.t. S and (x', y', z', t') w.r.t. S' (see Fig. 14.1).

With these assumptions clearly, points which are at rest relative to S' will move with speed v relative to S in X -direction. Particularly the point at $x'=0$ will move with speed v in X -direction i.e. $x'=0$ will be identical with $x=vt$, so that the first of our transformation-equation is

$$x' = \alpha(x - vt) \quad \dots (1)$$

where α is a constant to be determined later and vt is the distance traversed by S' in time t along X -axis.

Now velocity of S' being along X -axis only, it follows from symmetry that

$$y' = y \text{ and } z' = z \quad \dots (2)$$

This set of equations is not complete, unless we formulate an equation connecting t' the time measure in S' , with the space and time coordinates i.e. x, y, z, t in S . Due to homogeneity of space and time, t' must linearly depend on t, x, y, z but for reasons of symmetry we assume that t' does not depend on y and z , otherwise two clocks in S' -system in $x'=0$ plane would appear to disagree as observed from S and hence we take

$$t' = \beta t + \gamma x \quad \dots (3)$$

where β and γ are not merely constants but are functions of v and are to be determined along with α .

Let us now assume that the event P is a light signal, produced when both t and t' are zero and when origins of two systems coincide. Also let the light pulse produced at $t=0$ spread out as a growing sphere such that the radius of wave front so produced grows with speed c . But the velocity of light in both the systems being the same i.e. c in all directions, the progress of the spherical wave front is described by either of the following equations:

$$x^2 + y^2 + z^2 = c^2 t^2 \quad \dots (4)$$

$$x'^2 + y'^2 + z'^2 = c^2 t'^2 \quad \dots (5)$$

Substituting values of x', y', z', t' from (1), (2) and (3) to (5), we get

$$\alpha^2 (x - vt)^2 + y^2 + z^2 = c^2 (\beta t + \gamma x)^2$$

$$\text{or } (\alpha^2 - c^2 \gamma^2)x^2 + y^2 + z^2 - 2\alpha\gamma x t (\alpha^2 v + c^2 \beta \gamma) = (c^2 \beta^2 - \alpha^2 v^2)t^2 \quad \dots (6)$$

The equations (4) and (6) representing the same motion are identical and hence comparison of various coefficients, yields

$$\alpha^2 - c^2 \gamma^2 = 1 \quad \dots (7)$$

$$\alpha^2 v + c^2 \beta \gamma = 0 \quad \dots (8)$$

$$c^2 \beta^2 - \alpha^2 v^2 = c^2 \quad \dots (9)$$

Multiplying (7) by v^2 and then subtracting (8) from it, we have

$$v(1 + c^2 \gamma^2) + c^2 \beta \gamma = 0 \quad \dots (10)$$

Again multiplying (7) by v^2 and adding it to (9), we get

$$-v^2(1 + c^2 \gamma^2) + c^2 \beta^2 = c^2 \quad \dots (11)$$

Now multiplying (10) by v and adding it to (11), we find

$$\beta^2 - 1 = -v\beta\gamma \text{ i.e. } \gamma = \frac{1 - \beta^2}{v\beta} \quad \dots (12)$$

Elimination of γ between (10) and (12) gives

$$v \left\{ 1 + c^2 \left(\frac{\beta^2 - 1}{v\beta} \right)^2 \right\} + c^2 \left(\frac{1 - \beta^2}{v} \right) = 0$$

$$\text{or } v^2 \beta^2 + c^2 (\beta^2 - 1)^2 + c^2 \beta^2 (1 - \beta^2) = 0$$

$$\text{or } \beta^2 \left[v^2 + c^2 - 2c^2 \right] + c^2 = 0 \text{ giving } \beta^2 = \frac{c^2}{c^2 - v^2} = \frac{1}{1 - \frac{v^2}{c^2}} \quad \dots (13)$$

With the help of (13), (9) yields (on taking positive roots only)

$$\alpha^2 = \frac{c^2}{c^2 - v^2} = \beta^2 \text{ so that } \alpha = \beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \quad \dots (14)$$

In view of (14), (8) gives

$$\gamma = \frac{-\alpha^2 v}{c^2 \beta} = \frac{-\alpha v}{c^2} = \frac{-\beta v}{c^2} \quad \dots (15)$$

From (12), (14) and (15), we get

$$\gamma = \frac{1 - \beta^2}{v\beta} = -\frac{\beta v}{c^2} = -\frac{v}{c^2 \sqrt{1 - \frac{v^2}{c^2}}} \quad \dots (16)$$

Also by (8),

$$\alpha^2 = \frac{-c\beta^2 \gamma}{v} = \beta^2 = \frac{1}{1 - \frac{v^2}{c^2}} \quad \dots (17)$$

In view of these results, the transformation equations (1), (2), (3) become

$$x' = \frac{x - vt}{\sqrt{1 - \frac{v^2}{c^2}}}, y' = y, z' = z \text{ and } t' = \frac{t - \frac{vx}{c^2}}{\sqrt{1 - \frac{v^2}{c^2}}} \quad \dots (18)$$

These are known as Lorentz transformation equations.

Inversely, we have

$$x = \frac{x' + vt'}{\sqrt{1 - \frac{v^2}{c^2}}}, y = y', z = z' \text{ and } t = \frac{t' + \frac{vx'}{c^2}}{\sqrt{1 - \frac{v^2}{c^2}}} \quad \dots (19)$$

These are known as Lorentz inverse transformation equations.

A look at the transformations (18) and (19) reveals that S has the relative velocity $-v$ w.r.t. S' . This conclusion is not trivial since neither the unit length nor the unit time is directly comparable in S and S' .

In case v is very small then $v/c \rightarrow 0$, so that by (14), $\beta \rightarrow 1$ and then Lorentz transformations reduce to the Galilean transformations i.e.

$$x' = x - vt, y' = y, z' = z, t' = t \quad \dots (20)$$

If $v > c$, then Lorentz transformations give imaginary values of x' and t' leading to an absurdity and hence v cannot be greater than c .

Again if $v < c$ so that $\frac{v^2}{c^2}$ is negligible, the Lorentz transformations take the form

$$x' = x - vt, y' = y, z' = z \text{ and } t' = t - \frac{vx}{c^2} \quad \dots (21)$$

In view of the transformations $y' = y$ and $z' = z$ so that

$$y^2 + z^2 = y'^2 + z'^2; \text{ (4) and (5) yield} \\ x^2 - c^2 t^2 = x'^2 - c^2 t'^2 \quad \dots (22)$$

We may also write

$$x^2 + (ict)^2 = x'^2 + (ict')^2, \text{ where } i = \sqrt{-1} \quad \dots (23)$$

Now using the conceptions of four-dimensional geometry and imaginary rotations, we can determine a necessary connection between two sets of coordinates (x_1, x_2, x_3, x_4) and (x'_1, x'_2, x'_3, x'_4) .

Setting $x = x_1, x' = x'_1$ and $ict = x_4, ict' = x'_4$, (23) can be written as

$$x_1^2 + x_4^2 = x_1'^2 + x_4'^2 \quad \dots (24)$$

Geometrically interpreted if x_1, x_4 and x'_1, x'_4 form two rectangular sets of coordinate axes, with the same origin, then L.H.S. and R.H.S. of (24) separately give the distances of the points (x_1, x_4) and (x'_1, x'_4) from the same origin. Consequently either the two sets of axes coincide or are inclined at an angle θ (say) with each other. In later case we can express

$$x_1' = x_1 \cos \theta + x_4 \sin \theta, x_4' = -x_1 \sin \theta + x_4 \cos \theta \quad \dots (25)$$

Of course, as is evident from (24) and (25) a circle in original set retains its shape in the rotated set and the coordinates in two sets are connected by circular functions $\sin \theta$ and $\cos \theta$.

If we conceive of an object at rest in S , moving with velocity v w.r.t. S' then we can write $\frac{dx_1}{dt'} = 0$ and $\frac{dx}{dt} = v$, so that

$$\frac{dx_1'}{dx_4'} = \frac{dx_1'}{dx_4} \bigg/ \frac{dx_4'}{dx_4} = \frac{\frac{dx_1}{dx_4} \cos \theta + \sin \theta}{-\frac{dx_1}{dx_4} \sin \theta + \cos \theta} = 0$$

$$\frac{dx_4}{dx_1} = -\tan \theta \quad \dots (26)$$

But when $x = x_1$ and $x_4 = ict$, $\frac{dx}{dt} = v$ gives

$$\frac{dx_1}{dx_4'} = \frac{dx_1}{dt} \cdot \frac{dt}{dx_4} = v \cdot \frac{1}{ic} = -\frac{iv}{c} \quad \dots (27)$$

$$\therefore (26) \text{ and } (27) \text{ give } \tan \theta = \frac{iv}{c} \quad \dots (28)$$

$$\text{Accordingly } \frac{\sin \theta}{iv} = \frac{\cos \theta}{c} = \frac{\sqrt{\cos^2 \theta + \sin^2 \theta}}{\sqrt{c^2 + (iv)^2}} = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\therefore \sin \theta = \frac{iv}{\sqrt{1 - \frac{v^2}{c^2}}} \text{ and } \cos \theta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

The substitution of these values in (25) yields Lorentz transformation equations.

The coordinates x and t in (22) may also be related with x', t' by hyperbolic functions such as

$$x' = x \cosh \zeta - ct \sinh \zeta; ct' = -x \sinh \zeta + ct \cosh \zeta \quad \dots (29)$$

COROLLARY. The above transformations have been derived in a particular case when the frame S' is moving with uniform velocity v along X -axis. We can generalize these results by taking the motion of S' w.r.t. S , along any straight line, so that its velocity

$$\mathbf{v} = v_x \mathbf{i} + v_y \mathbf{j} + v_z \mathbf{k}$$

where v_x, v_y, v_z are the components of $\mathbf{v}$ along X, Y, Z axes respectively.

Taking $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ as the position vector of an event its resolved parts along and perpendicular to $\mathbf{v}$ may be given by $\frac{(\mathbf{r} \cdot \mathbf{v})\mathbf{v}}{v^2}$ and

$$\mathbf{r} - \frac{(\mathbf{r} \cdot \mathbf{v})\mathbf{v}}{v^2}$$

Denoting them by $\mathbf{P}$ and $\mathbf{Q}$ respectively so that $\mathbf{r} = \mathbf{P} + \mathbf{Q}$ and applying the transformations of (18), we have

$$\mathbf{Q}' = \mathbf{Q} - \frac{(\mathbf{v} \cdot \mathbf{r})\mathbf{v}}{v^2} \quad \dots (30)$$

$$\text{and } \mathbf{P}' = \frac{\mathbf{P} - \mathbf{v}t}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{(\mathbf{r} \cdot \mathbf{v})\mathbf{v} - \mathbf{v}t}{\sqrt{1 - \frac{v^2}{c^2}}} \quad \dots (31)$$

$$\text{Thus } \mathbf{r}' = \mathbf{P}' + \mathbf{Q}' = \mathbf{r} + \frac{(\mathbf{r} \cdot \mathbf{v})\mathbf{v}}{v^2} (\gamma - 1) + \gamma \mathbf{v}t \quad \dots (32)$$

$$\text{where } \gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}.$$

Also we have the transformation

$$t' = \gamma \left(t - \frac{\mathbf{r} \cdot \mathbf{v}}{c^2} \right) \quad \dots (33)$$

(32) and (33) give the general Lorentz transformations without rotation. Their inverse transformations may be obtained by replacing $\mathbf{v}$ by $-\mathbf{v}$ whence we have

$$\mathbf{r} = \mathbf{r}' + \frac{\mathbf{r}' \cdot \mathbf{v}}{v^2} (\gamma - 1) + \gamma \mathbf{v}t' \text{ and } t = \gamma \left(t' + \frac{\mathbf{r}' \cdot \mathbf{v}}{c^2} \right) \quad \dots (34)$$

Problem 1. State the fundamental postulates of special theory of relativity and deduce the Lorentz transformations.

(Agra, 1954, 56, 62, 67)

§§14.2, 14.3 constitute the answer.

Problem 2. Show that Lorentz transformations $x' = \left(1 - \frac{v^2}{c^2}\right)^{-\frac{1}{2}} (x - vt)$, $y' = y$, $z' = z$ and $t = \left(1 - \frac{v^2}{c^2}\right)^{-\frac{1}{2}} \left(t - \frac{vx}{c^2}\right)$ may be obtained from the two basic postulates of the special theory of relativity.

(Agra, 1960)

This is the same as Problem 1.

Problem 3. Show that $x^2 + y^2 + z^2 - c^2 t^2$ is invariant.

If (x, y, z, t) and (x', y', z', t') be the coordinates of the same event observed by two observers in frames S and S' where S' is moving with velocity v relative to S , then we have to show that

$$x^2 + y^2 + z^2 - c^2 t^2 = x'^2 + y'^2 + z'^2 - c^2 t'^2$$

Lorentz transformations are

$$x' = \beta(x - vt), \quad y' = y, \quad z' = z \text{ and } t' = \beta \left(t - \frac{vx}{c^2} \right)$$

$$\text{where } \beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

We thus have

$$x'^2 + y'^2 + z'^2 - c^2 t'^2 = \beta^2 (x - vt)^2 + y^2 + z^2 - c^2 \beta^2 \left(t - \frac{vx}{c^2} \right)^2$$

$$\begin{aligned} &= \beta^2 \left[x^2 + v^2 t^2 - 2vxt - c^2 \left(t^2 + \frac{v^2 x^2}{c^4} - \frac{2vtx}{c^2} \right) \right] + y^2 + z^2 \\ &= \beta^2 \left(1 - \frac{v^2}{c^2} \right) x^2 - c^2 t^2 \left(1 - \frac{v^2}{c^2} \right) \beta^2 + y^2 + z^2 \\ &= x^2 - c^2 t^2 + y^2 + z^2 \quad \because \quad \beta^2 = \frac{1}{1 - \frac{v^2}{c^2}} \\ &= x^2 + y^2 + z^2 - c^2 t^2 \end{aligned}$$

which shows the invariance of $x^2 + y^2 + z^2 - c^2 t^2$.

Problem 4. Derive Lorentz inverse transformations.

Lorentz transformations are

$$x' = \beta(x - vt), \quad y' = y, \quad z' = z, \quad t' = \beta \left(t - \frac{vx}{c^2} \right)$$

$$\text{where } \beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}.$$

Solving them for x, y, z , we find

$$x = \frac{x'}{\beta} + v \left(t' + \frac{vx'}{c^2} \right) = \frac{1}{\beta} (x' + vt') + \frac{v^2 x'}{c^2}$$

$$\text{or } \left(1 - \frac{v^2}{c^2} \right) x = \frac{1}{\beta} (x' + vt') \text{ i.e. } x = \beta (x' + vt')$$

As such, $t' = \beta \left(t - \frac{vx}{c^2} \right)$ gives

$$\begin{aligned} t &= \frac{t'}{\beta} + \frac{v}{c^2} \beta (x' + vt') = \frac{t'}{\beta} \left(1 + \frac{v^2}{c^2} \beta^2 \right) + \frac{vx'}{c^2} \beta \\ &= \frac{t'}{\beta} \left[\frac{1 - \frac{v^2}{c^2} + \frac{v^2}{c^2}}{1 - \frac{v^2}{c^2}} \right] + \frac{vx'}{c^2} \beta = \frac{t'}{\beta \left(1 - \frac{v^2}{c^2} \right)} + \frac{vx'}{c^2} \beta \\ &= \frac{t'}{\beta} \beta^2 + \frac{vx'}{c^2} \beta = \beta \left(t' + \frac{vx'}{c^2} \right) \end{aligned}$$

Hence Lorentz inverse transformations are

$$x = \beta (x' + vt'), \quad y = y', \quad z = z', \quad t = \beta \left(t' + \frac{vx'}{c^2} \right)$$

14.4. THE KINEMATIC EFFECTS OF THE LORENTZ TRANSFORMATION

[A] Lorentz-Fitzgerald Contraction

(Agra, 1969)

Consider two frames of reference S, S' such that S' is moving with velocity v along x -axis relative to S . Then Lorentz inverse transformations are

$$x = \beta (x' + vt'), y = y', z = z', t = \beta \left(t' + \frac{vx'}{c^2} \right) \quad \dots (1)$$

$$\text{where } \beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

Suppose that there is a rod of length l placed parallel to Y -axis in system S , which is at rest. Then if y_1, y_2 be the Y -coordinates of the end points of the rod, we have

$$l = y_2 - y_1$$

$$\text{But from (1), } y_2' - y_1' = y_2 - y_1 = l$$

which follows that the length of the rod remains unaltered when measured by an observer in S' . Conclusively the lengths perpendicular to the direction of motion are unaltered.

Further suppose that the rod of length l is placed parallel to X -axis in system S , which is at rest. Let x_1, x_2 be the X -coordinates of the end points of rod, measured at the same instant t in S so that $t_1 = t_2 = t$ and $l = x_2 - x_1$. Also let x_1', x_2' be the X -coordinates of the two ends of the rod, measured by an observer in system S' at the same instant t' , so that $t_1' = t_2' = t'$. If l' be the length of the rod in the system S' , then $l' = x_2' - x_1'$.

$$\text{Now, } l = x_2 - x_1 = \beta (x_2' + vt_2') - \beta (x_1' + vt_1') \text{ by (1)}$$

$$= \beta (x_2' - x_1') + v\beta (t_2' - t_1')$$

$$= \beta l' = \frac{l'}{\sqrt{1 - \frac{v^2}{c^2}}} \quad \because t_1' = t_2'$$

$$\therefore l' = l \sqrt{1 - \frac{v^2}{c^2}} = l \left\{ 1 - \frac{v^2}{c^2} \right\}^{1/2} = l \left\{ 1 - \frac{1}{2} \cdot \frac{v^2}{c^2} + \dots \right\}$$

$$\text{which follows that } l' < l \text{ and } l' = l \left(1 - \frac{v^2}{c^2} \right)^{1/2} \quad \dots (2)$$

Here l is the length of the rod for an observer at rest w.r.t. the rod while to an observer in S' which is in relative motion w.r.t. the rod, the rod appears to be of length l' . Actually speaking to the first observer the rod appears to be at rest while to the second observer it appears to be in motion, the length of the rod being measured in the direction of motion. Clearly the length of the rod appears to be longest in the reference system w.r.t. which it is at rest. The length l of the rod measured by an observer which is at rest w.r.t. the rod is said to be the *proper length* of the rod.

Thus Lorentz-Fitzgerald contraction follows:

Every rigid body seems to be of maximum dimensions when at rest relative to the observer while its dimensions appear to be contracted in the direction of relative motion by the factor $\left(1 - \frac{v^2}{c^2} \right)^{1/2}$, when it moves with velocity v relative to the observer, whereas its dimensions perpendicular to the direction of motion are unaffected.

Consequently the length is not an absolute quantity. An interesting consequence of this fact is that a moving sphere seems to be a prolate ellipsoid. In case $v=c$, from (2) $l'=0$ i.e., the rod appears to be of length zero to an observer in system S' moving with velocity of light relative to S . Also if $v > c$, l' is imaginary, leading to an absurdity. Hence $v > c$ i.e., $v < c$ always.

Problem 5. Determine three dimensional volume element $dx dy dz$ under Lorentz transformations.

We have by Lorentz-Fitzgerald contraction

$$dx' = \left(1 - \frac{v^2}{c^2} \right)^{1/2} dx, dy' = dy, dz' = dz$$

$$\text{So that } dx' dy' dz' = \left(1 - \frac{v^2}{c^2} \right)^{1/2} dx dy dz.$$

Problem 6. A rod of length 100 cm. is placed in a satellite moving with velocity $0.8c$ relative to a laboratory. Determine its length as observed by an observer (a) in the satellite (b) in the laboratory.

(a) The observer being in the satellite is at rest relative to the rod and hence the length of the rod measured by him is 100 cm.

(b) The length of the rod as measured by an observer in the laboratory is given by

$$l' = l \sqrt{1 - \frac{v^2}{c^2}} \text{ where } l = 100 \text{ cm, } v = 0.8c$$

$$\therefore l' = 100 \sqrt{1 - \left(\frac{0.8c}{c} \right)^2} = 100 \sqrt{1 - 0.64} = 100 \times 0.6 = 60 \text{ cm.}$$

Problem 7. The length of a rocketship is 100 metres on the ground. When it is in flight its length appears to be 99 metres to an observer on the ground. Determine its speed.

$$\text{We have } l' = l \sqrt{1 - \frac{v^2}{c^2}}$$

$$\text{where } l = 100 \text{ metres} = 10^4 \text{ cm. ; } l' = 99 \text{ metres} = 99 \times 10^3 \text{ cm.}$$

$$\therefore \left(1 - \frac{v^2}{c^2} \right) = \left(\frac{l'}{l} \right)^2 = \left(\frac{99 \times 10^3}{10^4} \right)^2 = (.99)^2$$

$$\text{So that } \frac{v^2}{c^2} = 1 - (.99)^2 = (1 - 0.99)(1 + 0.99) = 0.01 \times 1.99$$

$$= 199 \times 10^{-4}$$

$$\therefore v = c (199 \times 10^{-4})^{1/2} = (\sqrt{199} \times 10^{-2}) \times 3 \times 10^{10} \\ = 3\sqrt{199} \times 10^8 = 3 \times 14.1 \times 10^8 = 4.23 \times 10^9 \text{ cm./sec.}$$

Problem 8. Determine the length of a rod (proper length 100 cm) moving with a velocity of $0.8c$ in a direction inclined at an angle of 60° to its own length.

Taking i, j as unit vectors along X and Y axes respectively, the proper length l of the rod moving with velocity $0.8c$ along X -axis is given by

$$l = l \cos 60^\circ \mathbf{i} + l \sin 60^\circ \mathbf{j} = \frac{l}{2} \mathbf{i} + l \sqrt{\frac{3}{2}} \mathbf{j}.$$

Now the motion being along X -axis, the contraction will take place along X -axis and the component of length along Y -axis will be unaffected.

Thus if l'_x and l'_y be the components of length of the rod when in motion along X -axis, we find

$$l'_x = l_x \sqrt{1 - \frac{v^2}{c^2}} = l \cos 60^\circ \sqrt{1 - \left(\frac{.8c}{c}\right)^2} = \frac{l}{2} \times 0.6 = 0.3l$$

$$\text{and } l'_y = l \sin 60^\circ = \frac{l \sqrt{3}}{2} = 0.866 l.$$

$$\begin{aligned} \therefore \text{Length of the moving rod} &= \sqrt{l'^2_x + l'^2_y} \\ &= l \sqrt{(0.3)^2 + (0.866)^2} = l \sqrt{.09 + 0.75} \text{ approx.} \\ &= l \sqrt{0.84} = 0.92 l \text{ approx.} \\ &= 100 \times .92 = 92 \text{ cm.} \end{aligned}$$

[B] Time-dilation or Apparent Retardation of Clocks

Consider two frames of reference S , S' such that S' is moving with velocity v along X -axis relative to S . Conceive of a clock placed in S at $x = x_1$, giving a signal at time t_1 , such that t_1 is the time measured by an observer in S' corresponding to it. Then Lorentz transformation gives

$$t_1' = \beta \left(t_1 - \frac{vx_1}{c^2} \right) \text{ where } \beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \quad \dots (3)$$

Suppose that the clock gives another signal at time t_2 in S , such that its corresponding time in S' is t_2' . Then

$$t_2' = \beta \left(t_2 - \frac{vx_1}{c^2} \right) \quad \dots (4)$$

Putting $t_2 - t_1 = t$ and $t_2' - t_1' = t'$ and then subtracting (3) from (4), we get,

$$\text{i.e. } t' = \beta t = \frac{t}{\sqrt{1 - \frac{v^2}{c^2}}} = t \left\{ 1 - \frac{v^2}{c^2} \right\}^{-1/2} = t \left\{ 1 + \frac{1}{2} \cdot \frac{v^2}{c^2} + \dots \right\}$$

$$\text{So that } t' > t \text{ and } t = t' \left(1 - \frac{v^2}{c^2} \right)^{1/2} \quad \dots (5)$$

which follows that the time interval t appears to be dilated or lengthened by the factor $\left(1 - \frac{v^2}{c^2} \right)^{-1/2}$ to the moving observer.

If we reverse the process i.e. put the clock at a point x_1 in S' giving signals at times t'_1, t'_2 while the corresponding times in S are t_1 and t_2 respectively then by Lorentz inverse transformation, we have

$$t_1 = \beta \left(t'_1 + \frac{vx'_1}{c^2} \right) \text{ and } t'_2 = \beta \left(t_2 + \frac{vx'_1}{c^2} \right).$$

So that $t' = t'_2 - t'_1 = \beta(t_2 - t_1) = \beta t$ implying $t' > t$ as above.

It follows that the time interval t' appears to be dilated by the factor $\left(1 - \frac{v^2}{c^2} \right)^{-1/2}$ to an observer moving with velocity v relative to S .

Conclusively a moving clock runs more slowly than a stationary one i.e. every clock seems to go at its fastest rate when it is at rest relative to the observer and if it moves w.r.t. the observer with velocity v , its rate appears to go at its slowest rate by the factor

$$\sqrt{1 - \frac{v^2}{c^2}}.$$

It should be noted that the time recorded by a clock moving with a given system is said to be the **proper time** for that system.

Problem 9. Show that four dimensional volume element $dx \, dy \, dz \, dt$ is invariant under Lorentz transformation. (Agra, 1967, 70)

Lorentz-Fitzgerald contraction gives

$$dx' = \left(1 - \frac{v^2}{c^2} \right)^{1/2} dx, \, dy' = dy, \, dz' = dz$$

$$\text{Time dilation gives } dt' = \frac{dt}{\sqrt{1 - \frac{v^2}{c^2}}}.$$

So that $dx' \, dy' \, dz' \, dt' = dx \, dy \, dz \, dt$.

Which shows the required invariance.

Problem 10. Write a short note on Clock Paradox.

We have already discussed in time-dilation that according to an observer in S , a clock in S' is going slow while from the point of view of S' , it is a clock in S which is moving fast and hence when the observer comes back from S' to S , he finds just reversed phenomena. We use to call it as a Clock Paradox.

To be more explanatory an observer posited in either frame of reference finds that the rate of clock in other system is slow. There is complete reciprocity between the two frames S and S' .

Problem 11. In the laboratory the life-time of a particle moving with speed 2.8×10^{10} cm./sec. is found to be 2.5×10^{-7} sec.. Calculate the proper life-time of the particle. (Agra, 1969)

$$\text{Time-dilation gives } t' = \frac{t}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\text{where } t' = 2.5 \times 10^{-7} \text{ sec., } v = 2.8 \times 10^{10} \text{ cm./sec.,} \\ c = 3 \times 10^{10} \text{ cm./sec.}$$

$$\begin{aligned}\therefore t &= t' \sqrt{1 - \frac{v^2}{c^2}} = 2.5 \times 10^{-7} \sqrt{1 - \left(\frac{2.8 \times 10^{10}}{3 \times 10^{10}}\right)^2} \\ &= 2.5 \times 10^{-7} \sqrt{1 - (0.93)^2} = 2.5 \times 10^{-7} \times \sqrt{1 - 0.87} \text{ approx.} \\ &= 2.5 \times 10^{-7} \times .36 = 9 \times 10^{-8} \text{ sec.}\end{aligned}$$

[C] Relativity of Simultaneity (Agra, 1964, 66, 67)

Consider two frames of reference S, S' such that S' is moving with velocity v along X -axis relative to S . Let the two events in S with space and time coordinates (x_1, y_1, z_1, t_1) and (x_2, y_2, z_2, t_2) respectively, occur simultaneously, so that

$$x_1 \neq x_2 \text{ but } t_1 = t_2$$

Let t'_1, t'_2 be the corresponding times in S' . Then by Lorentz transformations we have

$$t'_1 = \beta \left(t_1 - \frac{vx_1}{c^2} \right) \text{ and } t'_2 = \beta \left(t_2 - \frac{vx_2}{c^2} \right)$$

where

$$\beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\begin{aligned}\therefore t'_2 - t'_1 &= \beta(t_2 - t_1) + \beta \frac{v}{c^2} (x_1 - x_2) \\ &= \frac{\beta v}{c^2} (x_1 - x_2) \quad \because t_1 = t_2\end{aligned}$$

In view of $x_1 \neq x_2$, this follows that $t'_1 \neq t'_2$ i.e. the same two events are not simultaneous in S' .

Conclusively two simultaneous events at different points, for an observer at rest in S remain no longer simultaneous to an observer in S' which is moving with velocity v along X -axis relative to S .

In case $x_1 = x_2$, then however the events are simultaneous since $t'_2 = t'_1$.

[D] Relativistic Composition of Velocities (Agra 1964, 65, 68, 71)

Consider two frames of reference S, S' such that S' is moving with velocity v along X -axis relative to S which is at rest. Let the space and time coordinates of a moving particle P be (x, y, z, t) and (x', y', z', t') w.r.t. S and S' respectively. Also let u_x, u_y, u_z and u'_x, u'_y, u'_z be the velocity components of P w.r.t. S and S' respectively. Then

$$u_x = \frac{dx}{dt}, \quad u_y = \frac{dy}{dt}, \quad u_z = \frac{dz}{dt}$$

and

$$u'_x = \frac{dx'}{dt'}, \quad u'_y = \frac{dy'}{dt'}, \quad u'_z = \frac{dz'}{dt'}$$

Lorentz inverse transformations are

$$x = \beta(x' + vt'), \quad y = y', \quad z = z' \text{ and } t = \beta \left(t' + \frac{vx'}{c^2} \right)$$

where

$$\beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

Their differentiations give

$$dx = \beta(dx' + vdt'), \quad dy = dy', \quad dz = dz';$$

$$dt = \beta \left(dt' + \frac{v}{c^2} dx' \right), \text{ so that}$$

$$u_x = \frac{dx}{dt} = \frac{\beta(dx' + vdt')}{\beta \left(dt' + \frac{v}{c^2} dx' \right)} = \frac{\left(\frac{dx'}{dt'} + v \right)}{\left(1 + \frac{v}{c^2} \frac{dx'}{dt'} \right)} = \frac{v'_x + v}{1 + \frac{v}{c^2} u'_x} \dots (6)$$

$$\begin{aligned}u_y = \frac{dy}{dt} &= \frac{dy'}{\beta \left(dt' + \frac{v}{c^2} dx' \right)} = \frac{\frac{dy'}{dt'}}{\beta \left(1 + \frac{v}{c^2} \frac{dx'}{dt'} \right)} \\ &= \frac{u'_y}{1 + \frac{v}{c^2} u'_x} \dots (7)\end{aligned}$$

$$\begin{aligned}u_z = \frac{dz}{dt} &= \frac{dz'}{\beta \left(dt' + \frac{v}{c^2} dx' \right)} = \frac{\frac{dz'}{dt'}}{\beta \left(1 + \frac{v}{c^2} \frac{dx'}{dt'} \right)} \\ &= \frac{u'_z}{1 + \frac{v}{c^2} u'_x} \dots (8)\end{aligned}$$

(6), (7), (8) give the velocity components u_x, u_y, u_z as the composition of velocities (u'_x, u'_y, u'_z) and $(v, 0, 0)$.

COROLLARY 1. The relativistic law of addition of two velocities in the same direction say X -axis, is given by putting $u'_x = u', u'_y = 0, u'_z = 0$ in the above components whence we get

$$u_x = \frac{u' + v}{1 + \frac{u'v}{c^2}} \dots (9)$$

COROLLARY 2. The velocity of light is an absolute constant, quite independent of the motion of the reference system.

Putting $u' = c$ in (9), we get

$$u_x = \frac{c + v}{1 + \frac{cv}{c^2}} = \frac{(c + v)}{c + v} = c \dots (10)$$

which follows that the velocity of light c remains unchanged if compounded with a velocity smaller than c .

COROLLARY 3. The resultant of two velocities each being less than c , is also less than c .

Take $u' = c - p$, $v = c - q$ where $p, q > 0$ and p, q are constants. Then (9) yields

$$u_x = \frac{c - p + c - q}{1 + \frac{(c - p)(c - q)}{c^2}} = \frac{2c - p - q}{1 + \frac{c^2 - (p + q)c + pq}{c^2}}$$

$$= c \left[\frac{2c - p - q}{2c^2 - (p + q)c + pq} \right] = c \left[\frac{2c - (p + q)}{2c - (p + q) + \frac{pq}{c}} \right]$$

which follows that $u_x < c$ $\because 2c - (p + q) < 2c - (p + q) + \frac{pq}{c}$

for $p, q > 0$ and $\frac{pq}{c} > 0$.

COROLLARY 4. The addition of any velocity to c the velocity of light results in c .

Taking $u' = c$ and $v = c$, (9) gives

$$u_x = \frac{c + c}{1 + \frac{c \cdot c}{c^2}} = \frac{2c}{2} = c.$$

COROLLARY 5. The velocity of light i.e. c is known as **fundamental velocity** due to the following facts.

- (i) c is constant in all directions.
- (ii) c remains the same for all observers irrespective of the velocities of the source and the observer.
- (iii) c is invariant for any two systems related by Lorentz transformations.
- (iv) c remains unchanged by adding any velocity to it, which is less than c .
- (v) No velocity can be greater than c .

Problem 12. Deduce Fresnel's coefficient of drag from Lorentz transformations (or in particular from relativistic velocity composition law). (Agra, 1958, 66)

Assuming that the frame S' is water moving with velocity v along X -axis relative to another inertial frame S which is at rest, and taking μ as the refractive index of water, the velocity of a light particle (photon) relative to water (i.e. S')

is $\frac{c}{\mu} - u'$ (say) towards X -axis.

If u_x be the velocity of the light particle w.r.t. S , then we have by relativistic velocity addition law

$$u_x = \frac{\frac{c}{\mu} + v}{1 + \frac{c}{\mu} \cdot \frac{v}{c^2}} = \left(\frac{c}{\mu} + v \right) \left(1 + \frac{v}{c\mu} \right)^{-1}$$

$$= \left(\frac{c}{\mu} + v \right) \left\{ 1 - \frac{v}{c\mu} + \frac{v^2}{\mu^2 c^2} - \dots \right\}$$

$$= \left(\frac{c}{\mu} + v \right) \left(1 - \frac{v}{c\mu} \right), \text{ neglecting squares of } \frac{v}{c} \because v < c$$

$$= -\frac{c}{\mu} + v - \frac{v^2}{\mu^2 c}, \text{ neglecting } \frac{v^2}{\mu^2 c} \text{ as } v < c$$

$$= \frac{c}{\mu} + v \left(1 - \frac{1}{\mu^2} \right) = u' + v \left(1 - \frac{1}{\mu^2} \right)$$

which follows that the velocity u' is increased as observed from S due to drag of water or system S' moving with velocity v by an amount $v \left(1 - \frac{1}{\mu^2} \right)$

$$\therefore \text{Coefficient of drag} = \frac{v \left(1 - \frac{1}{\mu^2} \right)}{v} = 1 - \frac{1}{\mu^2}$$

which is known as *Fresnel's coefficient of drag*.

Problem 13. If a photon traverses the path in such a way that it moves in $X'-Y'$ plane making an angle ϕ with X -axis of system S' , then prove that for the frame S , $u_x^2 + u_y^2 = c^2$, where S' is moving with velocity v relative to S .

If in a system S' , $u_y' = c \sin \phi$ and $u_x' = c \cos \phi$ then show that in frame S , $u_x^2 + u_y^2 = c^2$ where S' is moving with velocity v relative to S .

We have

$$u_x = \frac{u_x' + v}{1 + \frac{v}{c^2} u_x'} = \frac{c \cos \phi + v}{1 + \frac{v}{c} \cos \phi}$$

and

$$u_y = \frac{u_y'}{\beta \left(1 + \frac{v}{c^2} u_x' \right)} = \frac{c \sin \phi \sqrt{1 - \frac{v^2}{c^2}}}{1 + \frac{v}{c} \cos \phi}$$

Squaring and adding these expressions, we find

$$u_x^2 + u_y^2 = \frac{1}{\left(1 + \frac{v}{c} \cos \phi \right)^2} \left[(c \cos \phi + v)^2 + c^2 \sin^2 \phi \left(1 - \frac{v^2}{c^2} \right) \right]$$

$$= \frac{1}{\left(1 + \frac{v}{c} \cos \phi \right)^2} [c^2 + v^2 + 2cv \cos \phi - v^2 \sin^2 \phi]$$

$$\because \sin^2 \phi + \cos^2 \phi = 1$$

$$= \frac{1}{\left(1 + \frac{v}{c} \cos \phi\right)^2} [c^2 + 2cv \cos \phi + v^2 \cos^2 \phi]$$

$$\therefore 1 - \sin^2 \phi = \cos^2 \phi$$

$$= \frac{c^2 \left(1 + \frac{v}{c} \cos \phi\right)^2}{\left(1 + \frac{v}{c} \cos \phi\right)^2} = c^2.$$

[E] Relativistic Composition of Accelerations

Consider two frames of reference S, S' such that S' is moving with velocity v along X -axis relative to S which is at rest. Let the space and time coordinates of a moving particle P be (x, y, z, t) and (x', y', z', t') w.r.t. S and S' respectively. Also let $u_x, u_y, u_z; a_x, a_y, a_z$ and $u'_x, u'_y, u'_z; a'_x, a'_y, a'_z$ be the velocity and acceleration components of P w.r.t. S and S' respectively. Then

$$a_x = \frac{du_x}{dt}, a_y = \frac{du_y}{dt}, a_z = \frac{du_z}{dt} \text{ and } a'_x = \frac{du'_x}{dt'}, a'_y = \frac{du'_y}{dt'}, a'_z = \frac{du'_z}{dt'}$$

Lorentz inverse transformations are

$$x = \beta(x' + vt'), y = y', z = z', t = \beta \left(t' + \frac{vx'}{c^2} \right) \text{ where } \beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}.$$

These give on differentiation,

$$dx = \beta(dx' + vdt'), dy = dy', dz = dz', dt = \beta \left(dt' + \frac{v}{c^2} dx' \right) \text{ giving}$$

$$\frac{dt}{dt'} = \beta \left(1 + \frac{v}{c^2} u'_x \right), \text{ so that by (6), (7) and (8), we have}$$

$$a_x = \frac{d}{dt} u_x = \frac{d}{dt} \left[\frac{u'_x + v}{1 + \frac{v}{c^2} u'_x} \right] = \frac{d}{dt'} \left[\frac{u'_x + v}{1 + \frac{v}{c^2} u'_x} \right] \frac{dt'}{dt}$$

$$= \frac{\frac{du'_x}{dt'} \left(1 + \frac{v}{c^2} u'_x \right) - (v + u'_x) \frac{v}{c^2} \frac{du'_x}{dt'}}{\left(1 + \frac{v}{c^2} u'_x \right)^2} \cdot \frac{1}{\beta \left(1 + \frac{v}{c^2} u'_x \right)}$$

$$= \frac{\frac{du'_x}{dt'} \left(1 - \frac{v^2}{c^2} \right)}{\beta \left(1 + \frac{v}{c^2} u'_x \right)^3} = \frac{\frac{du'_x}{dt'}}{\left(1 + \frac{v}{c^2} u'_x \right)^3} \beta^3 \quad \dots (10)$$

Similarly,

$$a_y = \frac{1}{\beta^2} \left[\frac{\frac{du'_y}{dt'}}{\left(1 + \frac{v}{c^2} u'_x \right)^2} - \frac{\frac{v}{c^2} \frac{du'_x}{dt'} u'_y}{\left(1 + \frac{v}{c^2} u'_x \right)^3} \right] \quad \dots (11)$$

$$a_z = \frac{1}{\beta^2} \left[\frac{\frac{du'_z}{dt'}}{\left(1 + \frac{v}{c^2} u'_x \right)^2} - \frac{\frac{v}{c^2} \frac{du'_x}{dt'} u'_z}{\left(1 + \frac{v}{c^2} u'_x \right)^3} \right] \quad \dots (12)$$

Equations (10), (11), (12) give the acceleration components a_x, a_y, a_z in terms of a'_x, a'_y, a'_z as observed in system S' . Evidently the components of acceleration in S' are constants, but the components of acceleration in S are not constant, in general owing to the fact that they contain velocity components along with acceleration components.

COROLLARY 1. In particular if the particle is instantaneously at rest relative to frame S' , then $u'_x = 0 = u'_y = u'_z$, so that (10) (11) and (12) reduce to

$$a_x = \left(1 - \frac{v^2}{c^2} \right)^{3/2} a'_x, a_y = \left(1 - \frac{v^2}{c^2} \right) a'_y, a_z = \left(1 - \frac{v^2}{c^2} \right) a'_z \quad \dots (13)$$

which follows that unlike Galilean transformations, the acceleration is different in two inertial frames. This is due to the relativity of space and time.

COROLLARY 2. Sets of instantaneous direction cosines of the motion of the particle P in frames S and S' .

Let l, m, n and l', m', n' be the instantaneous direction cosines of the motion of P in frames S and S' respectively. Then we have

$$l = \frac{u_x}{\sqrt{u_x^2 + u_y^2 + u_z^2}}, m = \frac{u_y}{\sqrt{u_x^2 + u_y^2 + u_z^2}}, n = \frac{u_z}{\sqrt{u_x^2 + u_y^2 + u_z^2}} \quad \dots (14)$$

$$\text{and } l' = \frac{u'_x}{\sqrt{u_x'^2 + u_y'^2 + u_z'^2}}, m' = \frac{u'_y}{\sqrt{u_x'^2 + u_y'^2 + u_z'^2}}, n' = \frac{u'_z}{\sqrt{u_x'^2 + u_y'^2 + u_z'^2}} \quad \dots (15)$$

Using Lorentz transformations, we have

$$l = \frac{\frac{u'_x + v}{1 + \frac{v}{c^2} u'_x}}{\frac{1}{1 + \frac{v}{c^2} u'_x} \sqrt{(u'_x + v)^2 + \frac{1}{\beta^2} (u_y'^2 + u_z'^2)}}$$

$$= \frac{u'_x + v}{\sqrt{u_x'^2 + 2u'_x v + v^2 + \left(1 - \frac{v^2}{c^2} \right) (u_y'^2 + u_z'^2)}}$$

$$= \frac{u'_x + v}{\sqrt{(u_x'^2 + u_y'^2 + u_z'^2) + 2u'_x v + \frac{v^2}{c^2} (c^2 - u_y'^2 - u_z'^2)}}$$

Dividing numerator and denominator of R.H.S. by

$\sqrt{u_x'^2 + u_y'^2 + u_z'^2}$ and using (15), we get

$$l = \frac{l' + \frac{v}{\sqrt{u_x'^2 + u_y'^2 + u_z'^2}}}{\left[1 + \frac{2lv}{\sqrt{u_x'^2 + u_y'^2 + u_z'^2}} + \frac{v^2}{c^2} \left(\frac{c^2}{u_x'^2 + u_y'^2 + u_z'^2} - m'^2 - n'^2 \right) \right]^{1/2}}$$

Similarly m and n can be expressed.

Problem 14. A particle P instantaneously at rest in frame S' experiences an acceleration $\mathbf{a}' = 3\mathbf{i} + 4\mathbf{j} + 12\mathbf{k}$. Determine the acceleration measured from an observer in frame S when S' is moving with velocity $.98c$ relative to S along X -axis.

Let a_x, a_y, a_z and a'_x, a'_y, a'_z be the components of $\mathbf{a}'$ in systems S and S' respectively. Then as we are given,

$$a'_x = 3, \quad a'_y = 4, \quad a'_z = 12$$

$$\text{By Cor. 1, } a_x = \left(1 - \frac{v^2}{c^2} \right) a'_x = \left\{ 1 - \left(\frac{.98c}{c} \right)^2 \right\} .3 = 0.024$$

(on simplification)

$$a_y = \left(1 - \frac{v^2}{c^2} \right) a'_y = \left\{ 1 - \left(\frac{.98c}{c} \right)^2 \right\} .4 = 0.1584 \quad ,$$

$$a_z = \left(1 - \frac{v^2}{c^2} \right) a'_z = \left\{ 1 - \left(\frac{.98c}{c} \right)^2 \right\} .12 = 0.4752 \quad ,$$

$$\text{Hence } \mathbf{u} = a_x \mathbf{i} + a_y \mathbf{j} + a_z \mathbf{k}$$

$$= 0.024 \mathbf{i} + 0.1584 \mathbf{j} + 0.4752 \mathbf{k}.$$

14.5. ENERGY—MOMENTUM RELATION

If a particle of mass m is moving with velocity v then its momentum p in classical Mechanics is given by $p = mv$... (1)

and the law of conservation of momentum states (i) the mass of a moving body is the same as that of the stationary body and (ii) the total momentum of a body remains unaltered unless an external force is applied. The first hypothesis does not stand true when examined by the help of Lorentz transformations, whereas the Lorentz invariance of law of conservation of momentum states that the mass of a moving body does not remain constant but it changes with velocity and given by

$$m = \frac{m_0}{\sqrt{1 - \frac{u^2}{c^2}}} \quad \dots (2)$$

(Agra, 1969, 70, 71)

where m_0 is the mass of the particle when at rest and m is its mass when moving with velocity u .

In order to prove (2), consider two frames S, S' such that S' is moving with velocity v along X -axis relative to S . Suppose that m_1 is

the mass of the particle moving with velocity u_1 in S along X -axis and its corresponding mass and velocity in S' are m_1' and u_1' respectively.

$$\text{Let } \beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}; \beta_1 = \frac{1}{\sqrt{1 - \frac{u_1^2}{c^2}}} \text{ and } \beta_1' = \frac{1}{\sqrt{1 - \frac{u_1'^2}{c^2}}} \quad \dots (3)$$

Velocity-composition law gives

$$u_1 = \frac{u_1' + v}{1 + \frac{v}{c^2} u_1'} \text{ giving } u_1' = \frac{u_1 - v}{1 - \frac{v}{c^2} u_1} \quad \dots (4)$$

$$\begin{aligned} \text{So that } \beta_1' u_1' &= \frac{u_1 - v}{\left(1 - \frac{u_1^2}{c^2} \right)^{1/2} \left(1 - \frac{v}{c^2} u_1 \right)} \\ &= \frac{u_1 - v}{\left\{ 1 - \frac{(u_1 - v)^2}{c^2 \left(1 - \frac{v}{c^2} u_1 \right)^2} \right\}^{1/2} \left(1 - \frac{v}{c^2} u_1 \right)} \text{ by (4)} \\ &= \frac{(u_1 - v) c}{\left[c^2 \left(1 - \frac{v}{c^2} u_1 \right)^2 - (u_1 - v)^2 \right]^{1/2}} \\ &= \frac{(u_1 - v) c}{\left[c^2 \left\{ 1 + \frac{v^2}{c^4} u_1^2 - \frac{2vu_1}{c^2} \right\} - (u_1^2 + v^2 - 2u_1v) \right]^{1/2}} \\ &= \frac{(u_1 - v)}{\frac{1}{c} \left[c^2 + \frac{v^2}{c^2} u_1^2 - 2vu_1 - u_1^2 - v^2 + 2vu_1 \right]^{1/2}} \\ &= \frac{u_1 - v}{\left[1 - \frac{u_1^2 + v^2}{c^2} + \frac{u_1^2 v^2}{c^4} \right]^{1/2}} = \frac{u_1 - v}{\left[\left(1 - \frac{v^2}{c^2} \right) \left(1 - \frac{u_1^2}{c^2} \right) \right]^{1/2}} \\ &= \beta \beta_1 (u_1 - v) \text{ by (3)} \\ \text{or } \frac{\beta_1' u_1'}{\beta_1} &= \beta (u_1 - v) \quad \dots (5) \end{aligned}$$

Assume that there are a number of such particles moving along X -axis and their masses and momentums being invariant in S , so that

$$\Sigma m_1 = \text{const.}, \quad \Sigma m_1 u_1 = \text{const.} \quad \dots (6)$$

But β and v being the same for every particle, (6) yield

$$\Sigma m_1 \beta v = \text{const.}, \quad \Sigma m_1 u_1 \beta = \text{const.}$$

which on subtraction give $\Sigma m_1 \beta (u_1 - v) = \text{const.}$

$$\text{or } \Sigma m_1 \frac{\beta_1' u_1'}{\beta_1} = \text{const. by (5)} \quad \dots (7)$$

In frame S' , applying law of conservation of momentum, we get
 $\Sigma m_1' u_1' = \text{const.} \quad \dots (8)$

Comparison of (7) and (8) renders $\frac{m_1 \beta_1}{\beta_1} = m_1'$
 or $\frac{m_1}{\beta_1} = \frac{m_1'}{\beta_1'} = m_0$ (say) as absolute constant giving
 $m_1 = \frac{m_0}{\sqrt{1 - \frac{u_1^2}{c^2}}}$ and $m_1' = \frac{m_0}{\sqrt{1 - \frac{u_1'^2}{c^2}}}$ with the help of (3).

We conclude from these results that for a body of mass m moving with velocity u relative to that system

$$m = \frac{m_0}{\sqrt{1 - \frac{u^2}{c^2}}}$$

which is the result (2) showing that the mass of a body increases with the increase of velocity.

When $u=0$, (2) gives $m=m_0$ i.e., mass of the particle at rest is m_0 and hence m_0 is known as the *rest mass*.

In case $u < c$ i.e., $\frac{u}{c} < 1$ then also $m=m_0$.

Now we establish the *mass-energy relation* i.e.,
 $E = mc^2 \quad \dots (9)$

(Agra, 1969, 71)

We know that the increase in energy of the particle by applying a force F is given in terms of work i.e., if dT be the increment in Kinetic energy T due to an increase dr in the displacement r , then

$$dT = F \cdot dr = F \cdot \frac{dr}{dt} \cdot dt = Fv dt \quad \dots (10)$$

$$\therefore v = \frac{dr}{dt} \text{ (velocity)}$$

But the force being defined as rate of change of momentum we have

$$F = \frac{d}{dt} (mv) \text{ or } F dt = d(mv) \quad \dots (11)$$

with the help of (11), (10) yields

$$\begin{aligned} dT &= v F dt = v \cdot d(mv) = v d \left(\frac{m_0 v}{\sqrt{1 - \frac{v^2}{c^2}}} \right) \text{ by (2)} \\ &= m_0 v \left[\frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} + \frac{v^2}{c^2 \sqrt{1 - \frac{v^2}{c^2}}} \right] \frac{dv}{\left(1 - \frac{v^2}{c^2} \right)} \text{ (on differentiation)} \end{aligned}$$

$$= \frac{m_0 v dv}{\left(1 - \frac{v^2}{c^2} \right)^{3/2}} \quad \dots (12)$$

Differentiation of (2) gives

$$dm = \frac{m_0 v dv}{c^2 \left(1 - \frac{v^2}{c^2} \right)^{3/2}} \text{ i.e., } c^2 dm = \frac{m_0 v dv}{\left(1 - \frac{v^2}{c^2} \right)^{3/2}} \quad \dots (13)$$

Comparing (12) and (13), we find

$$dT = c^2 dm \quad \dots (14)$$

If we consider the body initially at rest whence its rest mass is m_0 and on applying this force as it acquires velocity so its mass increases and becomes say $m = m_0 + dm$. Now the total Kinetic energy acquired by the body is given by

$$T = \int dT = \int_{m_0}^m c^2 dm \text{ by (4)} \\ = c^2 [m - m_0]$$

$$\text{or } T + m_0 c^2 = mc^2 \quad \dots (15)$$

Total energy $E =$ Kinetic energy of moving body + energy at rest

$$\text{i.e., } E = c^2 (m - m_0) + m_0 c^2 = mc^2$$

which establishes the relation (9).

In the last we prove the *energy-momentum relation* or *Einstein relation*:

$$E^2 = c^2 p^2 + m_0^2 c^4 \quad \dots (16)$$

where E is the energy and p is the momentum of a particle of rest mass m_0 .

Using (1) and (9), we have

$$\begin{aligned} E^2 - c^2 p^2 &= (mc^2)^2 - c^2 (mv)^2 \\ &= m^2 c^4 - m^2 c^2 v^2 \\ &= \left(\frac{m_0}{\sqrt{1 - \frac{v^2}{c^2}}} \right)^2 c^4 - \left(\frac{m_0 v}{\sqrt{1 - \frac{v^2}{c^2}}} \right)^2 c^2 v^2 \end{aligned}$$

by (2) on replacing v by u in the existing case.

$$= \frac{m_0^2 c^2}{1 - \frac{v^2}{c^2}} (c^2 - v^2) = m_0^2 c^4$$

$$\text{Hence } E^2 = c^2 p^2 + m_0^2 c^4$$

which establishes the relation (16).

Problem 15. Formulate the *energy-momentum transformation in the space time of special relativity*. (Agra, 1975)

Consider two systems S and S' such that S' is moving with velocity v along X -axis. If m and m' are the masses of a body in S and S' respectively, such that it is moving with velocities u (u_x, u_y, u_z) and u' (u_x', u_y', u_z') respectively. Then we have

$$m = \frac{m_0}{\sqrt{1 - \frac{u^2}{c^2}}}, \quad m' = \frac{m_0}{\sqrt{1 - \frac{u'^2}{c^2}}} \quad \dots (1)$$

where m_0 is the rest mass of the body.

Here $u^2 = u_x^2 + u_y^2 + u_z^2$ and $u'^2 = u_x'^2 + u_y'^2 + u_z'^2$

The velocity-composition law gives

$$u_x' = \frac{u_x - v}{1 - \frac{v}{c^2} u_x}, \quad u_y' = \frac{u_y \left(1 - \frac{v^2}{c^2}\right)^{1/2}}{1 - \frac{v}{c^2} u_x}, \quad u_z' = \frac{u_z \left(1 - \frac{v^2}{c^2}\right)^{1/2}}{1 - \frac{v}{c^2} u_x} \quad \dots (3)$$

Now from (1),

$$\frac{m}{m'} = \left(\frac{1 - \frac{u'^2}{c^2}}{1 - \frac{u^2}{c^2}} \right)^{1/2}$$

where $1 - \frac{u'^2}{c^2} = 1 - \frac{1}{c^2} (u_x'^2 + u_y'^2 + u_z'^2)$

$$= 1 - \left[(u_x - v)^2 + u_y^2 \left(1 - \frac{v^2}{c^2}\right) + u_z^2 \left(1 - \frac{v^2}{c^2}\right) \right] \left(\frac{1}{1 - \frac{v}{c^2} u_x} \right)^2 \text{ by (3)}$$

$$= \frac{c^2 \left(1 - \frac{u^2}{c^2}\right) \left(1 - \frac{v^2}{c^2}\right)}{c^2 \left(1 - \frac{v}{c^2} u_x\right)^2} \text{ (on simplification)}$$

$$\text{So that } \left(\frac{1 - \frac{v'^2}{c^2}}{1 - \frac{u^2}{c^2}} \right)^{1/2} = \frac{\left(1 - \frac{v^2}{c^2}\right)^{1/2}}{1 - \frac{v}{c^2} u_x}$$

$$\therefore \frac{m}{m'} = \frac{\left(1 - \frac{v^2}{c^2}\right)^{1/2}}{1 - \frac{v}{c^2} u_x} \text{ i.e., } m' = m \frac{\left(1 - \frac{v^2}{c^2}\right)^{1/2}}{\left(1 - \frac{v}{c^2} u_x\right)} = m \beta \left(1 - \frac{v}{c^2} u_x\right) \quad \dots (4)$$

$$\text{where } \beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

Also from (3),

$$u_x' = \frac{u_x - v}{1 - \frac{v}{c^2} u_x}, \quad u_y' = \frac{u_y}{\beta \left(1 - \frac{v}{c^2} u_x\right)}, \quad u_z' = \frac{u_z}{\beta \left(1 - \frac{v}{c^2} u_x\right)} \quad \dots (5)$$

Let p_x, p_y, p_z and p_x', p_y', p_z' be the components of momentum p of the particle, acquired due to velocity in frames S and S' respectively.

Then, $p_x = mu_x, p_y = mu_y, p_z = mu_z$ in frame S .

and $p_x' = m' u_x' = (mu_x - mv) \beta$ by (4) and (5)

$$= \left(p_x - \frac{v}{c^2} E \right) \beta \quad \because E = mc^2$$

Similarly, $p_y' = m' u_y' = mu_y = p_y$

and $p_z' = m' u_z' = mu_z = p_z$

Hence if E' corresponds to E , in frame S' , then

$E' = m' c^2 = \beta (mc^2 - mv u_x)$ by (4)

$$= \beta (E - v p_x)$$

Hence the required transformations are

$$p_x' = \beta \left(p_x - \frac{v E}{c^2} \right); \quad p_y' = p_y, \quad p_z' = p_z, \quad E' = \beta (E - v p_x)$$

$$\text{where } \beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

Problem 16. Show that $p^2 - \frac{E^2}{c^2}$ is Lorentz invariant.

Using notations of Problem 15, we have

$$\begin{aligned} p^2 &= p_x^2 + p_y^2 + p_z^2 \text{ and } p'^2 = p_{x1}'^2 + p_y'^2 + p_z'^2 \\ \therefore p'^2 - \frac{E'^2}{c^2} &= p_x'^2 + p_y'^2 + p_z'^2 - \frac{1}{c^2} \beta^2 (E - v p_x)^2 \quad \because E' = \beta (E - v p_x) \\ &= \beta^2 \left(p_x - \frac{v E}{c^2} \right)^2 + p_y^2 + p_z^2 - \beta^2 (E - v p_x)^2 \cdot \frac{1}{c^2} \\ &= \beta^2 \left[-\frac{1}{c^2} \left(1 - \frac{v^2}{c^2} \right) E^2 + p_x^2 \left(1 - \frac{v^2}{c^2} \right) \right] + p_y^2 + p_z^2 \\ &= p_x^2 + p_y^2 + p_z^2 - \frac{E^2}{c^2} \quad \because \beta^2 \left(1 - \frac{v^2}{c^2} \right) = 1 \\ &= p^2 - \frac{E^2}{c^2} \end{aligned}$$

which proves the required invariance.

Problem 17. Discuss space-like and time-like intervals.

(Agra 1966, 67, 70, 75)

If S, S' be two frames of reference such that S' is moving with velocity v along X -axis relative to S . Then Lorentz transformations are

$$x' = \beta(x - vt), \quad y' = y, \quad z' = z, \quad t' = \beta \left(t - \frac{vx}{c^2} \right)$$

where

$$\beta = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \quad \dots (1)$$

Take two events with coordinates (x_1, y_1, z_1, t_1) and (x_2, y_2, z_2, t_2) in S , then we have

$$s_{12}^2 = -(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2 + c^2(t_2 - t_1)^2 \quad \dots (2)$$

[Since in Minkowski's four dimensional space, the interval between two events (x, y, z, x_4) and $(x+dx, y+dy, z+dz, x_4+dx_4)$ is given by $ds^2 = dx^2 + dy^2 + dz^2 + dx_4^2$

On setting $x_4 = ict$ where we use to say the coordinates of an event (x, y, z, ict) as (x, y, z, t) , we have

$$ds^2 = dx^2 + dy^2 + dz^2 - c^2 dt^2$$

For the system S' , (2) becomes

$$s_{12}'^2 = -[(x_2' - x_1')^2 + (y_2' - y_1')^2 + (z_2' - z_1')^2] + c^2(t_2' - t_1')^2$$

$$= -[\beta^2\{(x_2 - x_1) - v(t_2 - t_1)\}^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2]$$

$$+ c^2\beta^2\left\{(t_2 - t_1) - \frac{v}{c^2}(x_2 - x_1)\right\}^2$$

$$= -[(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2] + c^2(t_2 - t_1)^2$$

$$\left[\text{on simplification and using } \beta^2\left(1 - \frac{v^2}{c^2}\right) = 1 \right]$$

$$= s_{12}^2 \text{ by (2)}$$

$$\therefore s_{12}' = s_{12} \quad \dots (3)$$

which follows that the interval s_{12} is Lorentz invariant.

Now if $s_{12} = 0$, the interval given by (2) is known as *singular interval* and then we have

$$-(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2 + c^2(t_2 - t_1)^2 = 0$$

which in the form of elements dx, dy, dz, dt , reduces to

$$-[dx^2 + dy^2 + dz^2] + c^2 dt^2 = 0$$

This is termed as the equation of a *null cone*.

Now if we assume that the two events occur at the same point in S' and the first event occurs after the second one, so that

$$x_2' = x_1', y_2' = y_1', z_2' = z_1'; t_2' > t_1'$$

then, we have $s_{12}' = c^2(t_2' - t_1')^2 > 0$

or $s_{12}' > 0$

But $s_{12} = s_{12}'$ by (3), therefore $s_{12} > 0$ i.e. the interval s_{12} is real. Such intervals which are real are known as *time like intervals*, because s_{12}' contains only time component.

Hence the condition for time-like interval is

$$c^2(t_2 - t_1)^2 > (x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2.$$

Conclusively if an interval is time like, then it corresponds to a frame of reference in which the interval between two events is real.

Again if two events in S' occur at the same time so that $t_1' = t_2'$ then we have $s_{12}'^2 = -[(x_2' - x_1')^2 + (y_2' - y_1')^2 + (z_2' - z_1')^2] < 0$

i.e. $s_{12}^2 < 0$ and hence s_{12} is imaginary. Such imaginary intervals are known as *space like intervals*, because s_{12}' contains only space coordinates.

Hence the condition for space like intervals is

$$c^2(t_2 - t_1)^2 < (x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2$$

Conclusively if an interval is space like, then it corresponds to a frame of reference in which the interval between two events is imaginary and the two events occur at the same instant of time.

Problem 18. The rest mass of an electron is 9×10^{-28} gm., what will be its mass if it were moving with velocity $0.8c$. (Kanpur, 1968)

$$\text{Given } v = 0.8c \text{ so that } \frac{v^2}{c^2} = \left(\frac{0.8c}{c}\right)^2 = 0.64.$$

Also given that $m_0 = 9 \times 10^{-28}$ gm.

$$\therefore m = \frac{m_0}{\left(1 - \frac{v^2}{c^2}\right)^{1/2}} = \frac{9 \times 10^{-28}}{(1 - 0.64)^{1/2}}$$

$$= \frac{9 \times 10^{-28}}{0.6} = 15 \times 10^{-28} \text{ gm.}$$

Problem 19. Find the velocity that an electron must be given so that its momentum is 10 times its rest mass times the speed of light. What is the energy at this speed. (Agra, 1970)

We have

$$p = 10m_0c = mv = \frac{m_0v}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\text{or } 1 - \frac{v^2}{c^2} = \frac{v^2}{100c^2}$$

$$\text{i.e. } \left(1 + \frac{1}{100}\right) \frac{v^2}{c^2} = 1 \quad \text{or } \frac{v^2}{c^2} = \frac{100}{101}$$

$$\text{or } v = c \sqrt{\frac{100}{101}} = 0.995c$$

$$\text{or } v = 0.995 \times 3 \times 10^{10} \text{ cm./sec.}$$

$$= 2.985 \times 10^{10} \text{ cm./sec.}$$

which is the required velocity.

$$\text{Again, } m = \frac{m_0}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{9 \times 10^{-28}}{\left(1 - \frac{100}{101}\right)^{1/2}} = 90.36 \times 10^{-28} \text{ gm.}$$

$$\text{Also } E = mc^2 = 90.36 \times 10^{-28} \times (3 \times 10^{10})^2$$

$$= 8.13 \times 10^{-6} \text{ ergs.}$$

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 20. Apply the law of relativistic composition of velocities to derive the elementary formula for aberration of light. (Agra, 1964; Banaras, 1970)

[Hint. Taking θ, θ' as angle between X-axis and the direction of motion of a light signal, in S, S' take $u_x = c \cos \theta, u_y = -c \sin \theta, u_z = 0, u_x' = c \cos \theta', u_y' = -c \sin \theta',$

$$u_z' = 0 \text{ and use Lorentz transformation to prove } \tan \theta' = \frac{\tan \theta \left(1 - \frac{v^2}{c^2}\right)^{1/2}}{1 + \frac{v}{c} \sec \theta}$$

which in classical treatment takes the shape $\delta\theta = \frac{v}{c} \sin \theta$]

Problem 21. Give relativistic treatment of Doppler's effect and distinguish it from the classical treatment. (Kanpur, 1971; Agra, 1965)

Problem 22. How much electric energy could theoretically be obtained by annihilation of 1 gm. of matter. (Agra, 1961)

[Ans. 5.885×10^{32} e.v.]

Problem 23. What is the annual loss in the mass of the sun, if approximately 2 calories of radiated energy are received by each square cm. of the earth's surface per minute? (Agra, 1954)

[Take distance of sun from the earth 150×10^6 k.m.]

[Ans. 6.915×10^{16} kgm/year]

Problem 24. Suppose that total mass of 1 kg is transformed into energy, how large is this energy in kilo-watt hours. (Agra, 1959)

[Ans. 2.5×10^{10} K.W.H.]

Problem 25. Calculate the Kinetic energy of an electron moving with a velocity of 0.98 times the velocity of light in the laboratory system. (Vikram, 1967)

[Ans. 3.25×10^{-6} ergs.]

CHAPTER 15

STATISTICAL PROBABILITY

15.1. INTRODUCTION

While reading newspapers and magazines in our daily life, we come across the word 'statistics', the use of which is not new, but its use in the present meaning is older than the year 1839 when the American Statistical Association was founded. The words 'statist', 'statistics' and 'statistical' seem to be derived more or less indirectly from the Latin word 'status' means a political state.

The dictionary meaning of statistics is 'numerical data collected systematically', or 'the science of collecting and interpreting such information'. According to its conventional use the word statistics may be defined as follows:

"By 'statistics' we mean quantitative data affected to a marked extent by multiplicity of causes, the 'statistical methods' mean elucidation of quantitative data affected by a multiplicity of causes and 'theory of statistics' means the exposition of statistical methods."

As a historical account Mr W. Hooper, M.D. (1770) while translating 'The Elements of Universal Erudition' written by Baron J.F. Von Bielfeld, defines the word 'statistics' as "the science that teaches us what is the political arrangement of all modern states of the known world." The German philosopher E. A. W. Zimmermann (1787) defines the word 'statistik' (statistics) in the preface to 'A Political Survey of the Present State of Europe' as "that branch of the political knowledge, which has for its object the actual and relative power of the several modern states, the power arising from their natural advantages, the industry and civilisation of their inhabitants and the wisdom of their Governments"

In early years statistics was supposed to be the science of kings, used for the purpose of administration while in later stage it was considered as necessarily a branch of economics. Now-a-days the meanings of statistics are interpreted in different ways by different classes. To a layman 'statistics' is nothing but a collection of figures and a statistician is no more than a computer who always counts the number of things. To an economist the field of statistics lies in quantitative analysis. To a physicist statistics is a probability distribution which forms the basis of the theory of errors.

The 'statistics' which was primarily regarded as a branch of economics has now become so popular and its application has become so wide that no branch of human knowledge escapes its approach. The prediction of H.G. Wells that "statistical thinking will one day

be as necessary for efficient citizenship as the ability to read and write now seems to become exactly true. In fact the statistical knowledge is going to become an unavoidable part of general education which provides the student according to George W. Snedecor 'an awareness of, and harmony with, the statistical content of the society' as mentioned in his article 'A Proposed Basic Course in Statistics'.

Today the naturalists, the biologists, the astronomers, the administrators, the businessmen, the economists, the chemists, the physicists, the photographers, all make frequent use of statistical methods of which the *probability*, because of the nature of statistical data and models, is the fundamental tool in statistical theory. Regarding probability as an idealization of the proportion of times that a certain result will occur in repeated trials of experiment, a probability model is the type of mathematical model. Consequently an astronomer uses statistical methods in making predictions about eclipses, a biologist utilizes them to generalize the laws of variations and heredity, a meteorologist uses them for weather forecasts, regarding temperature pressure and rainfall, etc. Due to the wide scope of statistics, it is almost impossible for any statistician to be expert in all branches.

The *statistical probability* can be successfully employed in finding the possible configurations of the birthdays of people in a year, in classifying the accidents according to the weekdays, in firing to get the number of hits at a number of targets, in sampling to classify people according to age or profession, in irradiation in biology when the cells in retina of the eye are exposed to light, in cosmic ray experiments to find the number of particles hitting the Geiger counters, in an elevator to find the different arrangements of discharging the passengers, in dice-rolling to find the possible outcomes of throw, in chemistry when a long chain polymer reacts with oxygen, in theory of photographic emulsions to find whether a grain reacts if it is hit by a certain number of quanta and in finding the possible distributions of misprints in a certain number of pages of a book.

The dictionary meaning of *probability* is 'likelihood' or 'anything that has appearance of truth..'. It seems that it has been derived from the Latin word '*probare*' meaning 'to prove'. For instance if we draw 99 balls out of an urn containing 100 balls and it is per chance that all the 99 balls are of green colour, then it is always possible or probable that the remaining one ball may be of some other colour say red, white or black. Though the uniformity of the colour of 99 balls is unable to confirm that the remaining ball is also green in colour unless we are told that all the hundred balls are of the same colour, but the conclusion that the balls are all green is not based on certainty rather than it is based on 'likelihood' or 'probability'.

In dealing with the mathematical theory of probability we give a numerical measure to the probability. For example if we toss a coin, then the probability of falling the head or the tail up is equal, or mathematically speaking, the probability of falling the head or the tail up each is half i.e., $\frac{1}{2}$. As another example, the probability of drawing a heart from a pack of cards is $\frac{1}{4}$ as there are four colours

(heart, diamond, spade and club) in all.

In fact the probability plays the same role in mathematics as the mass in mechanics e.g., the motion of the planetary system can be discussed without knowing their individual masses. But in statistics we are concerned much with physical or statistical probabilities which do not refer to judgements but the possible outcomes of a conceptual experiment.

Initially R. A. Fisher and R. Von Mises developed the statistical or empirical attitude towards probability.

15.2. DEFINITIONS OF PROBABILITY

Based on classical concepts, we give two definitions of probability.

[A] The **mathematical or 'a priori' probability**. If there are q number of exhaustive, mutually exclusive and equally likely cases of an event and suppose that p of them are favourable to the happenings of an event A under the given set of conditions, then the mathematical probability of the event A is defined as

$$P(A) = \frac{p}{q}.$$

We sometimes put this definitions in the words 'the odds in favour of A are p to $(q-p)$ ' or 'the odds against A are $(q-p)$ to p '. More precisely if we assume that the odds in favour of the event A are m to n (or n to m against A), the probability of happening the event A is defined as

$$P(A) = \frac{m}{m+n}.$$

e.g. The probability of drawing a white ball from a bag containing 3 white and 4 red balls is $\frac{3}{3+4}$ i.e. $\frac{3}{7}$.

Note 1. The word '*exhaustive*' used in the definition assures the happening of an event either in favour or against and rules out the possibility of happening of neither (in favour or against) in any trial. The word '*mutually exclusive*' is a safeguard against the probability of two simultaneous happenings in a trial, e.g., in tossing a coin, the head and tail cannot fall together, but falling of either excludes the other. The word '*equally likely*' means equally probable, i.e., no happening is biased or partially bound to occur.

[B] The **statistical or empirical or 'a posteriori' probability**. If in a large number of trials performed under the same conditions, the limit of the ratio of the number of happenings of an event A to the total number of trials is unique and finite when the number of trials tends to infinity, then this limit measures the probability of the happening of the event A .

Thus if in a large number of trials performed under the same set of conditions, p is the probability of happening of an event A and q that of its failure, then the probability of its happening in the next trial

is $\frac{p}{p+q}$, being assumed to determine the empirical probability that there is no known information relative to the probability of the happening of the event other than the past trials.

In other words, if an event A happens on pN occasions when a large number N is taken out of a series of trials, then the probability $P(A)$ of the event A is p defined as

$$P(A) = \lim_{N \rightarrow \infty} \frac{pN}{N} = p.$$

Precisely if m is the number of times in which the event A occurs in a series of n trials, then $P(A) = \lim_{n \rightarrow \infty} \frac{m}{n}$.

Note 2. If p is the probability of happening of an event A , i.e., $P(A)=p$ and q that of not happening of that event denoted by $P(\bar{A})$ is given by $P(\bar{A})=q=1-p$, so that

$$P(A) + P(\bar{A}) = 1.$$

Conclusions. (i) The probability $P(A)$ of an event A lies between 0 and 1, i.e., $0 \leq P(A) \leq 1$.

(ii) The probability of an impossible event is zero, i.e.,

$$P(0) = 0.$$

(iii) The probability of a certain event E is one, i.e.,

$$P(E) = 1.$$

Problem 1. Discuss the meaning and scope of statistics and show how it can be applied to social and physical sciences.

It seems that the word 'statistics' was derived from the latin word 'status' which means a political state. The dictionary gives the meaning of statistics as 'numerical data collected systematically' or 'the science of collecting and interpreting such information'. According to its popular use, by statistics we mean quantitative data affected to a marked extent by multiplicity of causes.

Primarily statistics was supposed to be the science of kings used for the purpose of administration, but later on it was regarded as a branch of economics. Nowadays the meanings of statistics are interpreted in different ways by different classes, e.g., for a layman statistics is nothing but a collection of figures, to an economist the field of statistics lies in quantitative analysis and to a physicist statistics is a probability distribution which forms the basis of the theory of errors.

Statistics, which was accepted for some time as necessarily a branch of economics has now become so popular and its application so wide that no branch of human knowledge escapes its approach. Today the naturalists, the biologists, the astronomers, the administrators, the businessmen, the economists, the chemists, the physicists, the photographers, all make frequent use of statistical methods of which probability is the fundamental tool. An astronomer uses statistical

methods in making predictions about eclipses, a biologist utilizes them to generalize the laws of variations and heredity, a meteorologist uses them for wheather forecasts, regarding temperature pressure and rainfall, etc.

So far as the applications of statistics to social and physical sciences are concerned, the statistical probability can be successfully employed in finding certain inferences in social and physical fields. Dealing with probability, we are able to show that the distribution of r balls in n cells has n^r different ordered samples with replacement of size r and without replacement it has ${}^nP_r = \frac{n!}{(n-r)!}$ different ordered samples of size r . Thus to find the possible configuration of birthdays of people in a year, take r the number of people and n the number of days in a year; to classify accidents according to the week days, take r the number of accidents and n the number of days in a week; to classify people according to age or profession, take a group of r people and n will be the number of classes; to observe irradiation in biology when the cells in the retina of the eye are exposed to light r is the number of light particles and n the number of cells; in cosmic ray experiments, r is the number of particles hitting the Geiger counters and n is the number of counters function: in an elevator to find the possible arrangements of discharging the passengers, r is the number of passengers and n is the number of floors; in dice throw r is the number of dice and n is the number of faces i.e. six; in tossing coins r is the number of coins and $n=2$; in coupon collecting r is the number of coupons collected and n are the kinds of coupons; in chemistry when a long chain polymer reacts with oxygen, r is the number of the reacting oxygen molecules and n is the number of polymer chains, etc.

Problem 2. From a pack of 52 cards two are drawn at random; find the chance that one is a knave and other a queen.

Total number of ways of drawing 2 cards from 52 cards $= {}^{52}C_2$.

The required cards a knave and a queen appear in different four colours, therefore each card can be drawn in 4 different ways. But the two events happen simultaneously and hence the number of favourable ways $= 4 \times 4 = 16$.

$$\therefore \text{required probability} = \frac{16}{{}^{52}C_2} = \frac{16}{52.51} = \frac{8}{663}.$$

Problem 3. What is the chance that a leap year, selected at random, will contain 53 Sundays?

There are 366 days in a leap year. Dividing 366 by 7, the number of days of a week, we conclude that the leap year consists of 52 complete weeks and 2 days more. These two days can be combined in 7 different ways as under:

(i) Sunday and Monday, (ii) Monday and Tuesday, (iii) Tuesday and Wednesday, (iv) Wednesday and Thursday, (v) Thursday and

Friday, (iv) Friday and Saturday, (viii) Saturday and Sunday.

Of these seven combinations only (i) and (vii) are favourable so that the required chance = $\frac{2}{7}$.

Problem 4. From a bag containing 4 white and 5 black balls a man draws 3 at random; what are the odds against these being all black?

Total number of ways in which 3 balls can be drawn = 9C_3 .

Number of ways in which 3 black balls can be drawn = 5C_3 .

$$\text{Required chance} = \frac{{}^5C_3}{{}^9C_3} = \frac{5.4.3}{1.2.3} \cdot \frac{9.8.7}{5} = \frac{5}{42}.$$

Problem 5. A card is drawn from an ordinary pack and a gambler bets that it is a spade or an ace. What are the odds against his winning this bet?

Number of ways in which a card can be drawn from 52 cards = ${}^{52}C_1 = 52$.

There are four aces so that the number of ways in which a card can be an ace is ${}^4C_1 = 4$.

Now there are 13 spade cards of which one is an ace. Out of the remaining 12 spade cards, a spade card can be drawn in ${}^{12}C_1$, i.e. 12 number of ways.

$\therefore$ the number of ways in which the drawn card may be a spade or an ace = $12 + 4 = 16$.

$$\text{Hence the required probability} = \frac{16}{52} = \frac{4}{13} = \frac{4}{9+4},$$

i.e. the odds against the gambler's winning are as 9 to 4.

Problem 6. The chance of an event happening is the square of the chance of a second event but the odds against the first are the cube of the odds against the second. Find the chance of each.

Let the chance of the happenings of the first and the second events be p and p' respectively. Then according to the first condition, we have

$$p = p'^2. \quad \dots (1)$$

According to the second condition, we have

$$\frac{1-p}{p} = \left(\frac{1-p'}{p'} \right)^3. \quad \dots (2)$$

Substituting the value of p from (1) to (2), we get

$$\frac{1-p'^2}{p'^2} = \left(\frac{1-p'}{p'} \right)^3$$

$$\text{or} \quad \frac{(1-p')(1+p')}{p'^2} = \frac{(1-p')(1-2p'+p'^2)}{p'^2}$$

$$\text{or} \quad p'(1+p') = 1-2p'+p'^2$$

$$\text{or} \quad p' + p'^2 = 1-2p'+p'^2$$

$$\text{or} \quad 3p' = 1, \text{ i.e. } p' = \frac{1}{3}$$

$$\text{and then from (1),} \quad p = \frac{1}{9}.$$

Problem 7. Three cards are drawn at random from an ordinary pack. Find the chance that they are a king, a queen and a knave.

Total number of ways in which 3 cards can be drawn from 52 cards = ${}^{52}C_3$.

The pack of cards consists of 4 kings, 4 queens and 4 knaves and therefore each of a king, a queen and a knave can be drawn in 4C_1 , i.e. 4 ways. But all the three events happen together, hence the number of ways in which a king, a queen and a knave can be drawn

$$= 4 \times 4 \times 4 = 64.$$

$$\therefore \text{required probability} = \frac{64}{{}^{52}C_3} = \frac{64.1.2.3}{52.51.50} = \frac{16}{5525}.$$

Problem 8. Eight letters, to each of which corresponds an envelope, are placed in the envelopes at random. What is the probability that all letters are not placed in the right envelopes.

Total number of ways in which 8 letters can be placed in 8 envelopes = $8!$.

Also there is only one way in which all the letters are placed in their right envelopes.

$$\therefore \text{Probability that all the letters are placed in the right envelopes} = \frac{1}{8!}.$$

Hence the required probability that all the letters are not placed in their right envelopes = $1 - \frac{1}{8!}$.

Problem 9. A and B stand in a ring with 10 other persons. If the arrangement of the persons is at random, find the chance that there are exactly 3 persons between A and B.

In a ring 12 persons can stand in $11!$ ways and 3 persons between A and B can be selected in ${}^{10}C_3$ ways. A and B interchange their positions in 2! ways. Also 3 persons between A and B can stand in 3! ways and the other 7 persons in 7! ways.

$$\begin{aligned} \therefore \text{Number of favourable ways} &= 2! 3! 7! \cdot {}^{10}C_3 \\ &= 2! 3! 7! \cdot \frac{10!}{7! 3!} \\ &= 2! 10!. \end{aligned}$$

$$\text{Required probability} = \frac{2! 10!}{11!} = \frac{2}{11}.$$

Problem 10. A number is chosen from each of two sets (1, 2, 3, 4, 5, 6, 7, 8, 9); (1, 2, 3, 4, 5, 6, 7, 8, 9).

If p_1 denotes the probability that the sum of the two numbers be 10 and p_2 the probability that their sum is 8, find $p_1 + p_2$.

Each set consists of 9 numbers and hence the total number of ways of choosing one from each = ${}^9C_1 \times {}^9C_1 = 81$.

A sum of 10 may be found in 9 different ways, like (1, 9), (2, 8), (3, 7), (4, 6), (5, 5), (6, 4), (7, 3), (8, 2), (9, 1) so that

$$p_1 = \frac{9}{81} = \frac{1}{9}.$$

Similarly a sum of 8 can be found in 7 ways so that $p_2 = \frac{7}{81}$.

$$\therefore p_1 + p_2 = \frac{1}{9} + \frac{7}{81} = \frac{16}{81}.$$

15.3. EVENTS

A collection of all possible outcomes of an experiment is said to be an event.

If such a collection contains the outcome of an event, then that event is said to have occurred.

An event is said to be *simple* or *compound* according as it cannot or can be decomposed.

e.g. if we toss a coin, it will turn up either a head or a tail. Thus there are only two possible outcomes giving a simple event.

If we throw a pair of dice, then to have a sum of 5 is a compound event as a sum of 5 can be obtained as (1, 4), (2, 3), (3, 2), (4, 1). Thus this compound event consists of 4 elementary events.

Mutually exclusive events. Two or more events are said to be mutually exclusive if the happening or occurrence of any one of them excludes the happening of the others.

For example, if we toss a coin, and it falls with head up, then the falling of the head up excludes the simultaneous happening of the tail up i.e., the two events of the falling head and tail up with a coin cannot happen together, but the happening of one excludes the happening of the other. So the two events are mutually exclusive.

Compound events (Joint occurrence). The simultaneous occurrence of two or more events in connection with each other is said to be a compound event.

For example, if we have an urn containing 100 balls of different colours say red and green and suppose 60 are red and 40 are green. If then it is proposed to draw 10 balls each of red colour, it is a simple event. But if it is proposed to draw first 20 balls of red colour and then 10 balls of green colour, then it is a compound event.

Dependent and independent events. Two or more events are said to be *dependent* or *independent* according as the occurrence of one does or does not affect the occurrence of the other or others. The dependent events are sometimes known as *contingent*.

For example, if from an urn containing 10 balls, it is proposed to draw 2 balls, then if a ball is drawn and it is not replaced unless the second ball is drawn, the event of the drawing of the second ball is dependent on that of the first. But if first ball is replaced and then the second ball is drawn, the event of drawing the second ball is independent.

15.4. THEOREM OF TOTAL PROBABILITY, i.e. ADDITIVE LAW OF PROBABILITY

If there are n mutually exclusive events $A_1, A_2, A_3, \dots, A_n$ whose probabilities are $P(A_1), P(A_2), P(A_3), \dots, P(A_n)$ respectively, then the probability that one of them will happen is the sum of their separate probabilities, i.e.,

$P(A_1 + A_2 + A_3 + \dots + A_n) = P(A_1) + P(A_2) + P(A_3) + \dots + P(A_n)$, where $P(A_1 + A_2 + \dots + A_n)$ denotes the probability of occurrence of at least one of the n events $A_1, A_2, A_3, \dots, A_n$.

Suppose there are N total number of exhaustive, mutually exclusive and equally likely cases of which $m_1, m_2, \dots, m_n$ are favourable to the events $A_1, A_2, \dots, A_n$ respectively. Then the total number of cases favourable to either A_1 or A_2 or A_3 or...or A_n is $m_1 + m_2 + m_3 + \dots + m_n$ so that the probability of happening of at least one of these events is

$$P(A_1 + A_2 + A_3 + \dots + A_n) = \frac{m_1 + m_2 + m_3 + \dots + m_n}{N}$$

$$= \frac{m_1}{N} + \frac{m_2}{N} + \dots + \frac{m_n}{N}$$

$$= P(A_1) + P(A_2) + P(A_3) + \dots + P(A_n).$$

Conclusions. (i) In case an event A is comprised by n mutually exclusive forms $A_1, A_2, \dots, A_n$, i.e.

$$A = A_1 + A_2 + \dots + A_n,$$

then probability A , i.e. $P(A)$ is the sum of the probabilities of $A_1, A_2, \dots, A_n$ separately, i.e.,

$$P(A) = P(A_1) + P(A_2) + \dots + P(A_n).$$

(ii) In case the n mutually exclusive events are exhaustive also, so that there is certainty of happening of at least one, i.e., $P(A_1 + A_2 + \dots + A_n) = 1$, we have

$$P(A_1) + P(A_2) + \dots + P(A_n) = 1.$$

Note. We generally use the following notations:

$P(A)$ denotes the probability for an event A to happen,

$P(\bar{A})$	„	„	„	„	A not to happen,
$P(A+B)$	„	„	„	„	occurrence of at least one of the events A and B
$P(AB)$	„	„	„	„	occurrence of both the events A and B ,
$P(A\bar{B})$	„	„	„	„	happening of A and not of B ,
$P(\bar{A}B)$	„	„	„	„	B „ A ,
$P(\bar{A}\bar{B})$	„	„	„	„	happening of neither of A and B ,

If A and B are two events such that AB and $A\bar{B}$ are two exhaustive and mutually exclusive forms in which A can occur, then we have

Thus A and B can be placed in 7 ways so as to have four letters in between. Also A and B can interchange their positions in $2!$ ways. The four letters between A and B can be chosen in ${}^{10}C_4$ ways out of remaining 10 letters when A and B have already been placed. Moreover, these four letters can be arranged in $4!$ ways and the remaining 6 letters in $6!$ ways.

$\therefore$ Number of favourable ways $= 7 \cdot 2! \cdot {}^{10}C_4 \cdot 4! \cdot 6!$
and total number of arrangements in which 12 letters can be put $= 12!$

$$\begin{aligned}\text{Hence the required probability} &= \frac{7 \cdot 2! \cdot {}^{10}C_4 \cdot 4! \cdot 6!}{12!} \\ &= \frac{7 \cdot 2! \cdot 10! \cdot 4! \cdot 6!}{4! \cdot 6! \cdot 12!} = \frac{14}{12 \cdot 11} \\ &= \frac{7}{66}.\end{aligned}$$

Problem 14. One of two events must happen; given that the chance of the one is two-thirds that of the other, find the odds in favour of the other.

Let p_1, p_2 be the probabilities of the two events and suppose that x is the chance of happening of either, say

$$p_1 = x \text{ so that } p_2 = \frac{2}{3}x.$$

But we have $p_1 + p_2 = 1$ for a sure event,

$$\text{i.e., } x + \frac{2}{3}x = 1 \text{ or } \frac{5}{3}x = 1 \text{ giving } x = \frac{3}{5}.$$

$$\therefore p_1 = \frac{3}{5} \text{ and as such } p_2 = \frac{2}{5} \cdot \frac{3}{5} = \frac{2}{5}.$$

Thus odds in favour of the other $= \frac{2}{5} : \frac{3}{5}$, i.e., 2:3.

15.5. THEOREM OF COMPOUND PROBABILITY OR MULTIPLICATIVE LAW OF PROBABILITY

If there are two events A and B , probabilities of their happening being $P(A)$ and $P(B)$ respectively, then the probability $P(AB)$ of the simultaneous occurrence of the events A and B is equal to the probability of A multiplied by the conditional probability of B (i.e., the probability of B when A has occurred) or the probability of B multiplied by the conditional probability of A , i.e.,

$$\begin{aligned}P(AB) &= P(A) P(B/A) \\ &= P(B) P(A/B),\end{aligned}$$

where $P(B/A)$ denotes conditional probability of B and $P(A/B)$ that of A and that if the two events are independent, then the theorem of compound probability is

$$P(AB) = P(A) P(B).$$

Suppose there are N total number of mutually exclusive and equally likely cases of which m are favourable to A . Let m_1 be the number of cases favourable to A and B both, while m_1 is included in m . Thus

$$P(B/A) = \frac{m_1}{m} \text{ and } P(A) = \frac{m}{N}.$$

Now $P(AB)$ = the probability of happening of A and B both

$$\begin{aligned}&= \frac{m_1}{N} = \frac{m_1}{m} \cdot \frac{m}{N} \\ &= P(A) P(B/A).\end{aligned} \quad \dots (1)$$

The interchange of A and B will yield a similar result

$$P(AB) = P(BA) = P(B) (A/B). \quad \dots (2)$$

In case the two events A and B are independent, i.e., the occurrence of one does not affect the other, $P(B/A)$ is the same as $P(B)$ and $P(A/B)$ is the same as $P(A)$, so that the results (1) and (2) both become $P(AB) = P(A) P(B)$.

Generalization. The result (3) may be generalized as, if there are $A_1, A_2, \dots, A_n$, n mutually independent events, then the compound probability is given by

$$P(A_1, A_2, A_3, \dots, A_n) = P(A_1) P(A_2) P(A_3) \dots P(A_n) \quad \dots (4)$$

In case of n mutually exclusive events $A_1, A_2, \dots, A_n$, the result (1) may be generalised as

$$P(A_1 A_2 \dots A_n) = P(A_1) P(A_2/A_1) P(A_3/A_1 A_2) \dots P(A_n/A_1 A_2 \dots A_{n-1}).$$

Conclusions. (i) If p be the chance that an event will happen in one trial, the chance that it will happen in any assigned succession of r trials is p^r ; for in this case

$$P(A_1) = P(A_2) = \dots = P(A_r) = p$$

$$\begin{aligned}\therefore \text{required probability} &= P(A_1) P(A_2) \dots P(A_r) \\ &= p \cdot p \dots r \text{ times} = p^r.\end{aligned}$$

(ii) If $p_1, p_2, p_3, \dots, p_n$ are the probabilities that n events happen, then probability that all the events fail is

$$(1-p_1)(1-p_2)(1-p_3) \dots (1-p_n)$$

Hence the chance that at least one of these events happens

$$= 1 - (1-p_1)(1-p_2) \dots (1-p_n) \quad (\text{IAS, 51})$$

Problem 15. In shuffling a pack of cards, three are accidentally dropped; find the chance that the missing cards should be from different suits:

The pack consists of 52 cards.

$$\text{The chance of dropping a card} = \frac{{}^{52}C_1}{{}^{52}C_1} = 1.$$

When one card is dropped, there remain 51 cards of which 39 cards are of suits different from that of dropped one. Thus the chance of dropping a card of different suit in second draw

$$= \frac{{}^{39}C_1}{{}^{51}C_1} = \frac{39}{51}.$$

When two cards are dropped, there remain 40 cards of which 26 cards are of suits different from those of dropped cards. Thus the chance of dropping a card of different suit in third draw

$$= \frac{{}^{26}C_1}{{}^{40}C_1} = \frac{26}{40}.$$

The events being dependent, the required chance

$$= 1 \times \frac{39}{51} \times \frac{38}{50} = \frac{16}{25}.$$

Problem 16. The face cards (three from each suit) are removed from a full pack. Out of the 40 remaining cards, 4 are drawn at random:

- (a) What is the probability that they belong to different suit?
 (b) What is the probability that the 4 cards drawn belong to different suits and different denominations.

Having removed 12 face cards, the remaining 40 consist of 10 cards of each suit:

(a) Chance of drawing a card in first draw $= \frac{40C_1}{40C_1} = 1.$

Having drawn 1 card, there remain 39 cards of which 30 are of suits different from the drawn one.

$\therefore$ the chance of drawing a card of different suit in second draw

$$= \frac{30C_1}{39C_1} = \frac{30}{39}.$$

Having drawn two cards, three remain 38 cards of which 20 are of suits different from the drawn cards.

$\therefore$ chance of drawing a card in third draw $= \frac{20C_1}{38C_1} = \frac{20}{38}$

Having drawn three cards, there remain 37 cards of which 10 are of suits different from drawn cards.

$\therefore$ chance of drawing a card in fourth draw $= \frac{10C_1}{37C_1} = \frac{10}{37}.$

All the events being dependent, the required probability

$$= 1 \times \frac{30}{39} \times \frac{20}{38} \times \frac{10}{37} = \frac{1000}{13323}.$$

(b) Chance of drawing a card in first draw $= \frac{40C_1}{40C_1} = 1.$

Having drawn one card, there remain 39 cards of which 9 cards are of the same suit and 3 of the same denomination (value) so that 27 cards out of 39 are such that they are of different colours and different denominations from the drawn one.

$\therefore$ Chance of drawing a card in second draw $= \frac{27C_1}{39C_1} = \frac{27}{39}.$

Similarly chance of drawing a card in third draw $= \frac{16C_1}{38C_1} = \frac{16}{38}$

and chance of drawing a card in fourth draw $= \frac{7C_1}{37C_1} = \frac{7}{37}.$

All events are mutually dependent.

$\therefore$ The required probability $= 1 \times \frac{27}{39} \times \frac{16}{38} \times \frac{7}{37} = \frac{504}{13323}.$

Problem 17. From a pack of cards two cards are drawn, the first being replaced before the second is drawn. Find the probability that

the first is a diamond and the second is a king.

Let A denote the event of drawing a diamond and B denote the event of drawing a king in the second draw, when the first card has been replaced. Then

$P(A)$ = Probability of drawing a diamond

$$= \frac{13C_1}{52C_1} = \frac{1}{4}.$$

$P(B)$ = Probability of drawing a king

$$= \frac{4C_1}{52C_1} = \frac{1}{13}.$$

The two events being independent, we have

$$P(AB) = P(A)P(B)$$

$$= \frac{1}{4} \times \frac{1}{13} = \frac{1}{52}.$$

Problem 18. Find the chance of throwing a 6 at least once in two throws of a single die.

Let A denote the event of throwing a six in the first throw and B that in the second throw. Then probability of throwing a 6 at least once in two throws may be represented by $P(A+B)$.

Now $P(A)$ = Prob. of throwing a 6 in first throw.

$$= \frac{1}{6}$$

and $P(B)$ = Prob. of throwing a 6 in second throw

$$= \frac{1}{6}.$$

But we have

$$P(A+B) = P(A) + P(B) - P(AB)$$

where $P(AB) = P(A)P(B)$, A , B being independent events

$$\therefore P(A+B) = \frac{1}{6} + \frac{1}{6} - \frac{1}{6} \cdot \frac{1}{6} \\ = \frac{1}{3} - \frac{1}{36} = \frac{11}{36}.$$

Problem 19. A coin is tossed three times. Find the probability of getting head and tail alternately.

Let $P(A)$ and $P(B)$ represent the probability of getting head and tail respectively. Then

$$P(A) = \frac{1}{2} = P(B).$$

The alternate occurring of head and tail may happen in two ways:

(i) starting with head,

(ii) starting with tail.

In case of first, the probability of the event

$$= P(A)P(B)P(A)$$

$$= \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2}$$

$$= \frac{1}{8}.$$

In second case the probability of the event

$$\begin{aligned} &= P(B)P(A)P(B) \\ &= \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} \\ &= \frac{1}{8}. \end{aligned}$$

But the two events being mutually exclusive, the total probability of happening any one of them $= \frac{1}{8} + \frac{1}{8} = \frac{1}{4}$.

Problem 20. Four persons are chosen at random from a group containing 3 men, 2 women and 4 children. Show that the chance that exactly two of them will be children is $\frac{1}{2}$.

Given Men	Women	Children	Total no. of persons		
3	+	2	+	4	= 9.

4 persons out of 9 can be selected in 9C_4 ways,

2 children out of 4 can be selected in 4C_2 ways.

When 2 children have to be selected, we are left with 5 persons (3 men and 2 women). To make the company of four including the two already selected children we can choose 2 persons out of 5 in 5C_2 ways.

$\therefore$ Number of ways of selecting 4 persons $= {}^4C_2 \times {}^5C_2$, the two events being independent.

Hence the required probability $= \frac{{}^4C_2 \times {}^5C_2}{{}^9C_4} = \frac{1}{2}$.

Problem 21. In a bag there are 6 balls of which 3 are white and 3 are black. They are drawn successively without replacement. What is the chance that the colours alternate?

Suppose the first drawn ball is white. Then probability of its drawing is ${}^3C_1/{}^6C_1 = \frac{3}{6}$. In the second draw, probability of the ball being black is then ${}^3C_1/{}^5C_1 = \frac{3}{5}$ (because there remain only 5 balls after first draw). Thus

$$\text{probability of white ball in third draw} = \frac{{}^2C_1}{{}^4C_1} = \frac{2}{4}.$$

$$,, \quad \text{black} \quad ,, \quad \text{fourth} \quad ,, \quad = \frac{{}^2C_1}{{}^3C_1} = \frac{2}{3}.$$

$$,, \quad \text{white} \quad ,, \quad \text{fifth} \quad ,, \quad = \frac{{}^1C_1}{{}^2C_1} = \frac{1}{2},$$

$$,, \quad \text{black} \quad ,, \quad \text{sixth} \quad ,, \quad = \frac{{}^1C_1}{{}^1C_1} = \frac{1}{1}.$$

$$\therefore \text{Total probability of this event} = \frac{3}{6} \cdot \frac{3}{5} \cdot \frac{2}{4} \cdot \frac{1}{2} \cdot \frac{1}{1} = \frac{1}{10}.$$

Similarly the probability of the event with a start of black ball $= \frac{3}{6} \cdot \frac{3}{5} \cdot \frac{2}{4} \cdot \frac{1}{2} \cdot \frac{1}{1} = \frac{1}{10}$.

Hence the two events being mutually exclusive, the probability of happening of any one of these events $= \frac{1}{10} + \frac{1}{10} = \frac{1}{5}$.

Problem 22. Three groups of children contain 3 girls and 1 boy, 2 girls and 2 boys, 1 girl and 3 boys. One child is selected at random from each group. Show that the chance that the three selected consist of 1 girl and 2 boys is $\frac{1}{3}$.

One girl and 2 boys can be selected in three different ways as discussed below:

probability of selecting girl from 1st and boys from 2nd and 3rd groups $= \frac{3}{4} \times \frac{2}{4} \times \frac{3}{4} = \frac{9}{64}$,

probability of selecting girl from 2nd and boys from 1st and 3rd groups $= \frac{2}{4} \times \frac{1}{4} \times \frac{3}{4} = \frac{3}{64}$,

probability of selecting girl from 3rd and boys from 1st and 2nd groups $= \frac{1}{4} \times \frac{3}{4} \times \frac{2}{4} = \frac{1}{8}$.

All these events being mutually exclusive, the required probability $= \frac{9}{64} + \frac{3}{64} + \frac{1}{8} = \frac{1}{8}$.

Problem 23. A can hit a target 2 times in 5 shots; B, 2 times in 3 shots; C, 3 times in 4 shots. They fire a volley. What is the probability that 2 shots hit?

There arise three cases, probabilities of which are as follows:

(i) Probability that A hits, B hits and C does not hit

$$= \frac{2}{5} \cdot \frac{2}{3} \cdot \frac{1}{4} = \frac{1}{15},$$

(ii) Probability that A hits, B does not hit, C hits

$$= \frac{2}{5} \cdot \frac{1}{3} \cdot \frac{3}{4} = \frac{1}{10},$$

(iii) Probability that A does not hit, B hits, C hits

$$= \frac{1}{5} \cdot \frac{2}{3} \cdot \frac{3}{4} = \frac{1}{10}.$$

All these events being mutually exclusive, the required probability $= \frac{1}{15} + \frac{1}{10} + \frac{1}{10} = \frac{4}{15} = \frac{2}{5}$.

15.6. PROBABILITY: A NEW STAND POINT

(A) Sample space or outcome space. A set consisting of the elementary events as its elements is said to be a sample space. It is generally denoted by S .

A sample space provides a mathematical model of an ideal experiment in the sense that every conceivable outcome of the experiment is completely described by one and only one sample point.

In fact an element in S is known as a *sample point* or *sample* and an event e.g. event A is a subset of the sample space S . The event $A = \{a\}$ consisting of a single sample point $a \in S$ is said to be an *elementary event*. The null sets ϕ and S itself are events. The null set ϕ is said to be an *impossible event* while S is said to be a *sure* or *certain event*.

(B) Correspondence between sets and events. Let there be two events A and B . Then

(i) $A \cup B$ denotes an event which occurs *iff* (if and only if) A occurs or B occurs (or both occur).

(ii) $A \cap B$ denotes an event which occurs *iff* A occurs and B occurs.

(iii) The complementary event of A denoted by A' or A is an event which occurs *iff* A does not occur.

$B-A$ will be the event consisting of all points not contained in the event A but contained in B , i.e.,

$$B-A = B \cap A'$$

$$\therefore A-A' = \phi = A \cap A'.$$

$$\text{Also } A \cup A' = S.$$

A complementary event A' is always mutually exclusive and exhaustive.

(iv) Two events A and B are called *mutually exclusive* or *disjoint* if $A \cap B = \phi$.

Problem 24. A coin is tossed and it is observed whether a head or tail is up. Describe the suitable sample space of the experiment.

Denoting by H the event in which the coin turns up head and by T , the event in which the coin turns up tail, the sample space of the experiment consists of only two elements i.e., $S = \{H, T\}$.

Problem 25. If a die is thrown and the number of spots on the uppermost face is observed, describe, the suitable sample space for this experiment.

The spots on the six faces of a die are 1, 2, 3, 4, 5, 6. When the die is thrown any of the six numbers will appear on the uppermost face; therefore

$$S = \{1, 2, 3, 4, 5, 6\}.$$

Problem 26. Let there be a pack of cards. Describe the suitable sample space for drawing a red card.

There are 52 cards in all out of which 26 are red and other 26 are black. Denoting by $e_1, e_2, e_3, \dots, e_{26}$ the events of drawing a red card, we have

$$S = \{e_1, e_2, \dots, e_{26}\}.$$

Problem 27. There is a box containing 4 chits numbered 1, 2, 3, 4. Describe the sample space of drawing two different numbered chits one after another.

Every outcome will be an ordered pair such that $1 \leq x \leq 4$, $1 \leq y \leq 4$ but $x \neq y$ where x denotes the number of first drawn chit and y that of the second.

Possible outcomes may be tabulated thus:

$x \backslash y$	1	2	3	4
1	$\times$	1, 2	1, 3	1, 4
2	2, 1	$\times$	2, 3	2, 4
3	3, 1	3, 2	$\times$	3, 4
4	4, 1	4, 2	4, 3	$\times$

$\therefore$ the sample space consists of 2 ordered pairs.

Hence, $S = \{(x, y) ; 1 \leq x \leq 4, 1 \leq y \leq 4 \text{ and } x \neq y\}$.

Problem 28. An urn contains three red and two white balls. Two balls are drawn and their colour is noted. Set up the sample space of this experiment.

Let the red ball be numbered as R_1, R_2, R_3 and white balls as W_1, W_2 .

Denoting the event of drawing ball R_1 in first draw and R_2 in second draw by $R_1 R_2$, the sample space consists of 20 outcomes as given below :

$$S = \{R_1 R_2, R_1 R_3, R_1 W_1, R_1 W_2, R_2 R_1, R_2 R_3, R_2 W_1, R_2 W_2, R_3 R_1, R_3 R_2, R_3 W_1, R_3 W_2, W_1 R_1, W_1 R_2, W_1 R_3, W_1 W_2, W_2 R_1, W_2 R_2, W_2 W_3, W_2 W_1\}.$$

(C) **The modern concept of probability.** It is a well known fact that the concept of probability developed from evil habits of games of chance or gambling used in France in 17th century. In this connection the French nobleman and gambler Chevalier de méré consulted the well known mathematician Blaise Pascal (1623-1662) who began to think over the problem how and to what degree of accuracy a gambler can be assured of his chance of success. Pascal solved the problem of de méré and had a correspondence with Pierre de Fermat (1601-1665) who became interested in this and other similar problems.

The phenomena occurring in nature or any secrets can be either *Deterministic* or *Probabilistic*, e.g., if a train moves at the rate of 20 km./hr., it is deterministic that it will travel 100 km. in 5 hours, but if a coin is tossed, then it is probabilistic to say that the chance of each either 'head up' or 'tail up' is equal.

Actually the theory of probability deals with the things likely to occur and their chance or probable values. It is a measure of degree of uncertainty rather than accuracy.

The concepts of probability are connected with the events or occurrences and the Repeated trials.

(D) **Probability of an event.** If in a series of n trials all made under the same conditions an event A is observed m times, then m is said to be the frequency of success and the ratio $\frac{m}{n}$ is said to be the relative frequency of success.

The probability of the event A is defined to be the limit p of the ratio $\frac{m}{n}$ when n tends to infinity (if it exists), i.e.,

$$P(B) = p = \lim_{n \rightarrow \infty} \frac{m}{n}.$$

This is sometimes known as the *frequency definition of probability*. In other words, if the event A consists of r clear events out of n , then

$$P(A) = \frac{x_r}{n} \frac{\text{Number of elementary events in } A}{\text{Number of elementary events in sample space } S} = \frac{N(A)}{N(S)}$$

In terms of sample space S associated with Σ the class of events if P be a real-valued function defined on Σ , then P is said to be the *probability function* and $P(A)$ the *probability* of the event A which satisfies the following conditions:

(i) For every event A , $0 \leq P(A) \leq 1$.

(ii) $P(S) = 1$ i.e. $\sum_{i=1}^n P(A_i) = 1$

where

$$S = \{A_1\} \cup \{A_2\} \cup \dots \cup \{A_n\}.$$

If $S = \phi$ (a null set), then $\sum_{i=1}^n P(A_i) = 0$ i.e. $P(\phi) = 0$.

(iii) If A and B are two mutually exclusive (i.e. disjoint) events i.e. $A \cap B = \phi$, then $P(A \cup B) = P(A) + P(B)$.

(iv) If $A_1, A_2, A_3, \dots, A_n$ are n mutually exclusive events then $P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$.

[E] Some theorems. **THEOREM 1.** The law of total probability.

If A and B are two mutually exclusive events, then

$$P(A \cup B) = P(A) + P(B)$$

or in general if $A_1, A_2, A_3, \dots, A_n$ are n mutually exclusive events, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n).$$

Out of m mutually exclusive and equally likely cases, let a correspond to the occurrence of event A and b corresponds to the occurrence of event B , so that the event $A \cup B$ occurs in $(a+b)$ cases due to the nature of the events A and B . Then,

$$P(A) = \frac{a}{m}, P(B) = \frac{b}{m}$$

and

$$\begin{aligned} P(A \cup B) &= \frac{a+b}{m} \\ &= \frac{a}{m} + \frac{b}{m} \\ &= P(A) + P(B). \end{aligned} \quad \dots (1)$$

To generalize the theorem by mathematical induction, since the theorem is true for $n=2$, let us assume that it is true for any positive integral value n i.e.

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n). \quad \dots (2)$$

Let $A_1, A_2, A_3, \dots, A_n, A_{n+1}$ be $(n+1)$ mutually exclusive events, then

$$\begin{aligned} P(A_1 \cup A_2 \cup \dots \cup A_n \cup A_{n+1}) &= P(A_1 \cup A_2 \cup \dots \cup A_n) \\ &\quad + P(A_{n+1}) \text{ by (1)} \\ &= P(A_1) + P(A_2) + \dots + P(A_n) \\ &\quad + P(A_{n+1}) \text{ by (2)} \end{aligned}$$

This follows that the theorem is true for $n+1$ and hence the theorem is true for any integral value n .

THEOREM 2. For any two events A and B , the probability that either A or B or both occur is given by

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

$A \cup B$ can be decomposed into two mutually exclusive events $A-B$ and B i.e.,

$$A \cup B = (A-B) \cup B.$$

Then from theorem 1, we have

$$\begin{aligned} P(A \cup B) &= P[(A-B) \cup B] \\ &= P(A-B) + P(B) \end{aligned}$$

Now A can be decomposed into two mutually exclusive events $A-B$ and $A \cap B$ i.e.

$$A = (A-B) \cup (A \cap B),$$

so that $P(A) = P(A-B) + P(A \cap B)$ by theorem 1

or $P(A-B) = P(A) - P(A \cap B)$.

$$\begin{aligned} \text{Thus } P(A \cup B) &= P(A) - P(A \cap B) + P(B) \\ &= P(A) + P(B) - P(A \cap B). \end{aligned} \quad \dots (1)$$

Aliter. $A \cup B = A \cup (B - A \cap B)$ where A and $B - A \cap B$ are disjoint.

$$\therefore P(A \cup B) = P(A) + P[B - A \cap B]$$

we have $B = [A \cap B] \cup [B - A \cap B]$ where $A \cap B$ and $B - A \cap B$ are disjoint.

$$\therefore P(B) = P(A \cap B) + P(B - A \cap B).$$

$$\text{Hence } P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

COROLLARY. For any events A, B, C applying this theorem twice we may easily get

$$\begin{aligned} P(A \cup B \cup C) &= P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) \\ &\quad - P(B \cap C) + P(A \cap B \cap C) \end{aligned} \quad \dots (2)$$

THEOREM 3. Let A be an event then its complementary event $\bar{A}$ will be disjoint to A , then $P(A) = 1 - P(\bar{A})$.

Since $A, \bar{A}$ are mutually exclusive events,

$$P(A \cup \bar{A}) = P(A) + P(\bar{A}).$$

But we know that $P(S) = 1 = P(A) + P(\bar{A})$

giving $P(A) = 1 - P(\bar{A})$.

THEOREM 4. If A and B are two events such that $A \subset B$, then

$$P(A) \leq P(B).$$

Since $A \subset B$, therefore B can be decomposed into two mutually exclusive events A and $B-A$ i.e.,




Fig. 15.1




Fig. 15.2

$B = A \cup (B - A)$,
so that by theorem 1,

$$P(B) = P(A) + P(B - A)$$

$$\therefore P(B) \geq P(A) \text{ as } 0 \leq P(B - A) \leq 1.$$

THEOREM 5. If A and B are two events, then

$$P(A \cup B) \leq P(A) + P(B).$$

By theorem 2, we have

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\therefore P(A \cup B) \leq P(A) + P(B) \text{ as } 0 \leq P(A \cap B) \leq 1.$$

COROLLARY. This result may be generalised in case of n events

$$A_1, A_2, A_3, \dots, A_n \text{ as}$$

$$P(A_1 \cup A_2 \cup \dots \cup A_n) \leq P(A_1) + P(A_2) + \dots + P(A_n).$$

[F] **Borel-field.** We assume that a family B of certain subsets of S (outcome space) is given, which will be called events and that this family satisfies the following axioms:

B₁. The outcome space S and the empty set ϕ are in B i.e.

$$S \in B, \phi \in B.$$

B₂. If each set of finite or countable sequence $a_1, a_2, \dots, a_i, \dots$ is in B , then their union and intersection are in B i.e.,

$$A_i \in B \text{ for } i=1, 2, \dots, \Rightarrow \bigcup_i A_i \in B$$

$$A_i \in B \text{ for } i=1, 2, \dots, \Rightarrow \bigcap_i A_i \in B.$$

B₃. $A \in B \Rightarrow \bar{A} = S - A \in B$.

The family B satisfying these three axioms is defined to be **Borel-field** B on the outcome space S .

[G] **Probability measure of a Borel-field.** Given a outcome space S and a Borel-field B , we consider a set-function $P(A)$ on B i.e. a rule which ascribes to every $A \in B$ a real number $P(A)$, then this $P(A)$ is called as **probability measure** on B provided it satisfies the following axioms:

P₁. $\forall$ (for every) $A \in B$, $P(A) \geq 0$ i.e. a non-negative real number.

P₂. $P(S) = 1$

P₃. If $A_1, A_2, \dots, A_j, \dots$ is a finite or countable sequence of mutually exclusive events, then

$$P(\bigcup_j A_j) = \sum_j P(A_j)$$

Illustration. $P(A) + P(\bar{A}) = 1$ and $A \cup \bar{A} = S$.

Axiom P_3 gives $P(A \cup \bar{A}) = P(S) = 1$

„ P_2 gives $0 \leq P(A) \leq 1$ from P_1 .

Also $P(\phi) = 0$, $S \cup \phi = S$, $P(S) + P(\phi) = P(S)$ i.e., $P(\phi) = 0$.

Problem 29. Let A and B be events with $P(A \cup B) = \frac{7}{8}$, $P(A \cap B) = \frac{1}{8}$ and $P(\bar{A}) = \frac{5}{8}$. Find $P(A)$, $P(B)$ and $P(A \cap \bar{B})$.

We have $P(A) = 1 - P(\bar{A})$
 $= 1 - \frac{5}{8} = \frac{3}{8}$.

Again we have

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\text{i.e., } \frac{7}{8} = \frac{3}{8} + P(B) - \frac{1}{8}.$$

$$\text{i.e., } P(B) = \frac{7}{8} - \frac{3}{8} + \frac{1}{8} = \frac{5}{8}.$$

Further we have $A - B = A \cap \bar{B}$, so that

$$P(A \cap \bar{B}) = P(A - B)$$

$$\text{i.e., } P(A - B) = P(A) - P(A \cap B) \text{ by theorem 2}$$

$$= \frac{3}{8} - \frac{1}{8} = \frac{2}{8}.$$

$$\therefore P(A \cap \bar{B}) = P(A - B) = \frac{1}{4}.$$

Problem 30. In an experiment of tossing two dice, if A denotes the event that the sum of the spots on uppermost faces is 7, find $P(A)$.

A seven can be found in following 6 ways

$$(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1).$$

While two dice can be thrown in 6^2 i.e., 36 ways.

$$\therefore \text{chance of throwing a 7 with two dice} = \frac{6}{36} = \frac{1}{6}.$$

If $e_1, e_2, e_3, e_4, e_5, e_6$ denote the events of throwing a seven as (1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1) respectively, then the event $A = \{e_1, e_2, e_3, e_4, e_5, e_6\}$,

$$\text{so that } P(A) = P(e_1) + P(e_2) + P(e_3) + P(e_4) + P(e_5) + P(e_6)$$

$$\text{But } P(e_1) = P(e_2) = \dots = P(e_6) = \frac{1}{36}.$$

$$\therefore P(A) = \frac{1}{36} + \frac{1}{36} + \dots 6 \text{ times.} \\ = \frac{1}{6}.$$

Problem 31. For any two events A and B prove that

$$P[(A \cap \bar{B}) \cup (B \cap \bar{A})] = P(A) + P(B) - 2P(A \cap B)$$

A can be decomposed into two mutually exclusive events $A \cap \bar{B}$ and $A \cap B$ i.e.,

$$A = (A \cap \bar{B}) \cup (A \cap B)$$

Similarly

$$B = (\bar{A} \cap B) \cup (A \cap B)$$

$$\therefore P(A) = P(A \cap \bar{B}) + P(A \cap B)$$

$$\text{and } P(B) = P(\bar{A} \cap B) + P(A \cap B).$$

Adding the last two results

$$P(A) + P(B) = P(A \cap \bar{B}) + P(\bar{A} \cap B) + 2P(A \cap B) \\ = P[(A \cap \bar{B}) \cup (B \cap \bar{A})] + 2P(A \cap B)$$




Fig. 15.3

as $A \cap \bar{B}$ and $B \cap \bar{A}$ are mutually exclusive events,

$$\therefore P[A \cap \bar{B}] \cup [B \cap \bar{A}] = P(A) + P(B) - 2P(A \cap B).$$

Problem 32. From a pack of cards one card is selected at random. What is the probability that the card is a spade, an honour or a deuce?

Denoting by A, B, C respectively the events of drawing a spade, an honour and a deuce, we have

$$P(A) = \frac{13}{52} = \frac{13}{52}, P(B) = \frac{12}{52} = \frac{12}{52}, P(C) = \frac{4}{52} = \frac{4}{52}$$

$$P(A \cap B) = \frac{3}{52}, P(A \cap C) = \frac{1}{52}, P(B \cap C) = 0 \text{ and } P(A \cap B \cap C) = 0.$$

Required probability

$$\begin{aligned} P(A \cup B \cup C) &= P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) \\ &\quad - P(B \cap C) + P(A \cap B \cap C) \\ &= \frac{13}{52} + \frac{12}{52} + \frac{4}{52} - \frac{3}{52} - \frac{1}{52} - 0 + 0 = \frac{25}{13}. \end{aligned}$$

Problem 33. An integer is chosen at random from the first 200 positive integers. What is the probability that the integer chosen is divisible by 6 or by 8?

Let A denote the event that an integer selected is divisible by 6 and B denote the event that an integer selected is divisible by 8. Then the sample space is

$$\begin{aligned} S &= \{1, 2, 3, \dots, 200\}. \\ P(A) &= \frac{33}{200}, P(B) = \frac{25}{200}, \text{ and } P(A \cap B) = \frac{8}{200}. \end{aligned}$$

Required probability $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

$$\begin{aligned} &= \frac{33}{200} + \frac{25}{200} - \frac{8}{200} \\ &= \frac{50}{200} = \frac{1}{4}. \end{aligned}$$

Problem 34. A coin and a die are thrown together. Find the chance of throwing a head and 5 or a tail and 6.

Let A denote an event of throwing a head and 5; and let B " " " " a tail and 6.

The two events being mutually exclusive, we have

$$P(A \cup B) = P(A) + P(B).$$

The sample space consists of 12 points as follows:

$$S = \{(H, 1), (H, 2), \dots, (H, 6); (T, 1), (T, 2), \dots, (T, 6)\}.$$

For the event $A = \{(H, 5)\}$, there is one sample point, so that $P(A) = \frac{1}{12}$ and for the event $B = \{(T, 6)\}$, there is also one sample point, so that $P(B) = \frac{1}{12}$.

$$\text{Hence } P(A \cup B) = \frac{1}{12} + \frac{1}{12} = \frac{1}{6}.$$

Problem 35. A has 3 shares in a lottery in which there are 3 prizes and 6 blanks and B has 1 share in lottery in which there is one prize

and two blanks. Show that A 's chance of success is to B 's as 16 to 7.

$$\text{We have to show that } \frac{P(A)}{P(B)} = \frac{16}{7}.$$

We have $P(A) + P(\bar{A}) = 1$, $\bar{A}$ being the event that A fails to win a prize.

$$\text{Sample points} = {}^9C_3.$$

$$\text{Event points} = {}^6C_3.$$

$$\therefore P(\bar{A}) = \frac{{}^6C_3}{{}^9C_3} = \frac{5}{21}, \text{ so that } P(A) = 1 - \frac{5}{21} = \frac{16}{21}.$$

$$\text{Similarly } P(\bar{B}) = \frac{{}^2C_1}{{}^3C_1} = \frac{2}{3}, \text{ so that } P(B) = 1 - \frac{2}{3} = \frac{1}{3}.$$

$$\text{Required ratio} = \frac{P(A)}{P(B)} = \frac{16}{21} \cdot \frac{1}{\frac{1}{3}} = \frac{16}{7}.$$

Aliter: Let

$A_1 = A$ wins a prize when 1 share wins but other two fails,

$A_2 =$ " " 2 shares win " one fails,

$A_3 =$ " " 3 " " 0 fail.

$$\text{Then } P(A_1) = \frac{{}^3C_1 \times {}^6C_2}{{}^9C_3}, P(A_2) = \frac{{}^3C_2 \times {}^6C_1}{{}^9C_3} \text{ and } P(A_3) = \frac{{}^3C_3 \times {}^6C_0}{{}^9C_3}$$

$$\begin{aligned} \therefore P(A) &= P(A_1) + P(A_2) + P(A_3) \\ &= \frac{{}^3C_1 \times {}^6C_2 + {}^3C_2 \times {}^6C_1 + {}^3C_3 \times {}^6C_0}{{}^9C_3} \\ &= \frac{16}{21}. \end{aligned}$$

Similarly $P(B)$ can be found to be $\frac{1}{3}$ and hence the required ratio.

[H] Conditional probability. If A and B are two dependent events in the sample space S , then the conditional probability of A given B (i.e. the probability of occurring of the event A on the assumption that B has already occurred) is defined as

$$P(A|B) = \frac{P(A \cap B)}{P(B)} \quad \dots (1)$$

provided $P(B) \neq 0$.

Similarly the conditional probability of B given A is defined as

$$P(B|A) = \frac{P(B \cap A)}{P(A)} \quad \dots (2)$$

provided $P(A) \neq 0$.

Note. In first case if $P(B) = 0$ or in second case $P(A) = 0$, the conditional probability is undefined.

[I] The law of compound probability. In the previous article we have

defined the conditional probability of A given B as

$$P(A/B) = \frac{P(A \cap B)}{P(B)}.$$

Multiplying both sides by $P(B)$, this becomes

$$P(B)P(A/B) = P(A \cap B),$$

$$\text{i.e. } P(A \cap B) = P(B) P(A/B).$$

$$\text{Similarly } P(B \cap A) = P(A) P(B/A).$$

$$\text{But } B \cap A = A \cap B.$$

$$\therefore P(B \cap A) = P(A \cap B),$$

$$\text{so that } P(A \cap B) = P(A) \cdot P(B/A) = P(B) \cdot P(A/B).$$

This gives the law of compound probability.

COROLLARY 1. In case of three events A, B, C , this law becomes

$$P(A \cap B \cap C) = P(A) \cdot P(B/A) \cdot P(C/A \cap B).$$

COROLLARY 2. In case of $n (\geq 2)$ events $A_1, A_2, A_3, \dots, A_n$ for which

$P(A_1 \cap A_2 \cap \dots \cap A_n) > 0$, then the compound probability

$$P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1) \cdot P(A_2/A_1) \cdot P(A_3/A_1 \cap A_2) \dots P[A_n/A_1 \cap A_2 \cap \dots \cap A_{n-1}].$$

This may be proved by the method of induction as follows:

Denoting by S_n the statement given above and denoting by N the set of those integers n for which S_n is true, we have by the method of induction for $N > 1$,

(i) $2 \in N$ for S_2 is the statement which is true,

(ii) $k \in N$, k being assumed an integer > 1 .

If we prove that $k+1 \in N$, then the statement becomes universally true.

For S_k the statement is

$$P(A_1 \cap A_2 \cap \dots \cap A_k) = P(A_1) P(A_2/A_1) \dots P(A_k/A_1 \cap A_2 \cap \dots \cap A_{k-1}). \dots (1)$$

It may be verified by using the definition of conditional probability and properties of set-intersection, that

$$\frac{P(A_1 \cap A_2 \cap \dots \cap A_{k+1})}{P(A_1 \cap A_2 \cap \dots \cap A_k)} = P(A_{k+1}/A_1 \cap A_2 \cap \dots \cap A_k). \dots (2)$$

Multiplying (1) and (2) together, we get

$$P(A_1 \cap A_2 \cap \dots \cap A_{k+1}) = P(A_1) \cdot P(A_2/A_1) \dots P(A_{k+1}/A_1 \cap A_2 \cap \dots \cap A_k),$$

which shows that if $k \in N$, then $k+1 \in N$, i.e. the statement is true for all values of k and hence the result holds good for any integral value n .

COROLLARY 3. Since B can be decomposed into two mutually exclusive events $A \cap B$ and $\bar{A} \cap B$ i.e.,

$$B = (A \cap B) \cup (\bar{A} \cap B)$$

we have,

$$\begin{aligned} P(B) &= P[(A \cap B) \cup (\bar{A} \cap B)] \\ &= P(A \cap B) + P(\bar{A} \cap B) \text{ as } P(E \cup F) = P(E) + P(F) \\ &= P(A) \cdot P(B/A) + P(\bar{A}) \cdot P(B/\bar{A}) \end{aligned}$$

Problem 36. Prove the following cases, assuming that in each case the conditional probabilities are defined.

(i) For any event A , $0 \leq P(A/B) \leq 1$, where $P(B) > 0$, B being another event.

(ii) For a certain event S , $P(S/B) = 1$, where $P(B) > 0$.

(iii) $P(A_1 \cup A_2/B) = P(A_1/B) + P(A_2/B) - P(A_1 \cap A_2/B)$.

But A_1 and A_2 are mutually exclusive and $P(B) > 0$,

$$P(A_1 \cup A_2/B) = P(A_1/B) + P(A_2/B)$$

or in general if $A_1, A_2, \dots, A_n$ is a sequence of n mutually exclusive events, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n/B) = P(A_1/B) + P(A_2/B) + \dots + P(A_n/B)$$

(iv) If A and B are mutually exclusive and $P(A \cup B) \neq 0$; then

$$P(A/A \cup B) = \frac{P(A)}{P(A) + P(B)}.$$

(i) We have $A \cap B \subset B$, so that B can be decomposed into two mutually exclusive events $A \cap B$ and $B - A \cap B$ i.e.

$$B = (A \cap B) \cup (B - A \cap B)$$

$$\begin{aligned} \therefore P(B) &= P(A \cap B) + P(B - A \cap B) \\ &\geq P(A \cap B) \text{ as } P(B - A \cap B) \geq 0 \end{aligned}$$

or

$$P(A \cap B) \leq P(B).$$

$$\text{Thus, } P(A/B) = \frac{P(A \cap B)}{P(B)} \leq 1$$

Also

$$P(A/B) \geq 0.$$

$\therefore$

$$0 \leq P(A/B) \leq 1.$$

(ii) We have $S \cap B = B$, S being a certain event

$$\text{i.e., } P(S \cap B) = P(B).$$

$$\therefore P(S/B) = \frac{P(S \cap B)}{P(B)} = \frac{P(B)}{P(B)} = 1.$$

(iii) We have

$$\begin{aligned} P(A_1 \cup A_2/B) &= \frac{P[(A_1 \cup A_2) \cap B]}{P(B)} \\ &= \frac{P[(A_1 \cap B) \cup (A_2 \cap B)]}{P(B)} \end{aligned}$$




Fig. 15.4

$$\begin{aligned}
 &= \frac{P(A_1 \cap B) + P(A_2 \cap B) - P[(A_1 \cap B) \cap (A_2 \cap B)]}{P(B)} \\
 &= \frac{P(B) \cdot P(A_1/B) + P(B) \cdot P(A_2/B) - P[A_1 \cap A_2 \cap B]}{P(B)}
 \end{aligned}$$

$$\begin{aligned}
 \text{or } P(A_1 \cup A_2/B) &= P(A_1/B) + P(A_2/B) - \frac{P(B) \cdot P(A_1 \cap A_2/B)}{P(B)} \\
 &= P(A_1/B) + P(A_2/B) - P(A_1 \cap A_2/B).
 \end{aligned}$$

In case A_1 and A_2 are mutually exclusive events, then $A_1 \cap B$ and $A_2 \cap B$ are also mutually exclusive, so that

$$\begin{aligned}
 P[(A_1 \cup A_2) \cap B] &= P[A_1 \cap B \cup (A_2 \cap B)] \\
 &= P(A_1 \cap B) + P(A_2 \cap B) \\
 \therefore P(A_1 \cup A_2/B) &= \frac{P[(A_1 \cup A_2) \cap B]}{P(B)}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{P(A_1 \cap B) + P(A_2 \cap B)}{P(B)} \\
 &= \frac{P(A_1 \cap B)}{P(B)} + \frac{P(A_2 \cap B)}{P(B)} \\
 &= P(A_1/B) + P(A_2/B).
 \end{aligned}$$

Again if $A_1, A_2, \dots, A_n$ are mutually exclusive events, then $A_1 \cap B, A_2 \cap B, \dots, A_n \cap B$ are also mutually exclusive, so that

$$\begin{aligned}
 P[(A_1 \cup A_2 \cup \dots \cup A_n) \cap B] &= P[(A_1 \cap B) \cup (A_2 \cap B) \cup \dots \cup (A_n \cap B)] \\
 &= P(A_1 \cap B) + P(A_2 \cap B) + \dots + P(A_n \cap B) \\
 \therefore P(A_1 \cup A_2 \cup \dots \cup A_n/B) &= \frac{P[(A_1 \cup A_2 \cup \dots \cup A_n) \cap B]}{P(B)}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{P(A_1 \cap B) + P(A_2 \cap B) + \dots + P(A_n \cap B)}{P(B)} \\
 &= \frac{P(A_1 \cap B)}{P(B)} + \frac{P(A_2 \cap B)}{P(B)} + \dots + \frac{P(A_n \cap B)}{P(B)} \\
 &= P(A_1/B) + P(A_2/B) + \dots + P(A_n/B).
 \end{aligned}$$

(iv) Since A and B are mutually exclusive events, we have

$$A \cup B = \phi$$

and

$$\begin{aligned}
 A \cap (A \cup B) &= (A \cup A) \cap (A \cup B) \\
 &= A \cap \phi = A
 \end{aligned}$$

so that

$$P[A \cap (A \cup B)] = P(A)$$

and also

$$P(A \cup B) = P(A) + P(B)$$

$$\begin{aligned}
 \text{Hence } P(A | A \cup B) &= \frac{P[A \cap (A \cup B)]}{P(A \cup B)} \\
 &= \frac{P(A)}{P(A) + P(B)}
 \end{aligned}$$

Problem 37. If A and B are two events such that $P(A) = \frac{2}{3}$, $P(B) = \frac{5}{6}$ and $P(A \cup B) = \frac{3}{4}$, find $P(A | B)$ and $P(B | A)$.

We have $P(A \cup B) = P(A) + P(B) - P(A \cap B)$.

$$\begin{aligned}
 \text{i.e.,} \quad \frac{3}{4} &= \frac{2}{3} + \frac{5}{6} - P(A \cap B) \\
 \text{giving} \quad P(A \cap B) &= \frac{1}{4}.
 \end{aligned}$$

$$\text{Now } P(A | B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{1}{4}}{\frac{5}{6}} = \frac{3}{10}$$

$$\text{and } P(B | A) = \frac{P(B \cap A)}{P(A)} = \frac{\frac{1}{4}}{\frac{2}{3}} = \frac{3}{8} \text{ as } B \cap A = A \cap B.$$

Problem 38. If A and B are two events such that $P(A) = \frac{1}{3}$, $P(B) = \frac{1}{4}$ and $P(A \cup B) = \frac{1}{2}$, find $P(A | B)$, $P(B | A)$, $P(A \cap \bar{B})$ and $P(A | \bar{B})$.

We have $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

$$\text{i.e.,} \quad \frac{1}{2} = \frac{1}{3} + \frac{1}{4} - P(A \cap B) \text{ giving } P(A \cap B) = \frac{1}{12}$$

$$\therefore P(A | B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{1}{12}}{\frac{1}{4}} = \frac{1}{3}$$

$$P(B | A) = \frac{P(B \cap A)}{P(A)} = \frac{\frac{1}{12}}{\frac{1}{3}} = \frac{1}{4}.$$

$$\text{Now, } P(\bar{B}) = 1 - P(B) = 1 - \frac{1}{4} = \frac{3}{4}.$$

A can be decomposed into two mutually exclusive events $A \cap B$ and $A \cap \bar{B}$, so that

$$A = (A \cap B) \cup (A \cap \bar{B})$$

$$\begin{aligned}
 \therefore P(A) &= P[(A \cap B) \cup (A \cap \bar{B})] \\
 &= P(A \cap B) + P(A \cap \bar{B}),
 \end{aligned}$$

$$\begin{aligned}
 \text{so that } P(A \cap \bar{B}) &= P(A) - P(A \cap B) \\
 &= \frac{1}{3} - \frac{1}{12} = \frac{1}{4}
 \end{aligned}$$

$$\text{also } P(A | \bar{B}) = \frac{P(A \cap \bar{B})}{P(\bar{B})} = \frac{\frac{1}{4}}{\frac{3}{4}} = \frac{1}{3}.$$

Problem 39. A bag contains 3 black and 4 white balls. Two balls are drawn at random one at a time without replacement.

(i) What is the probability that the second ball selected is white?

(ii) What is the conditional probability that the first ball selected is white if the second ball is known to be white?

Let W_1, W_2 denote the events of drawing a white ball in first and second draw, then

$$P(W_1) = \frac{{}^4C_1}{{}^7C_1} = \frac{4}{7}, \quad P(W_2) = \frac{{}^4C_1}{{}^7C_1} = \frac{4}{7}$$

$$\therefore P(\bar{W}_1) = 1 - P(W_1) = 1 - \frac{4}{7} = \frac{3}{7}$$

$$P(W_2 | W_1) = \frac{{}^3C_1}{{}^6C_1} = \frac{3}{6} = \frac{1}{2}$$

$$P(W_2 | \bar{W}_1) = \frac{{}^4C_1}{{}^6C_1} = \frac{4}{6} = \frac{2}{3}.$$

Also
$$P(W_2 | W_1) = \frac{P(W_2 \cap W_1)}{P(W_1)}$$

i.e.,
$$\frac{1}{2} = \frac{P(W_2 \cap W_1)}{\frac{4}{7}}$$

or
$$P(W_2 \cap W_1) = \frac{4}{7} \cdot \frac{1}{2} = \frac{2}{7}.$$

(i)
$$\begin{aligned} P(W_2) &= [(W_1 \cap W_2) \cup (\bar{W}_1 \cap W_2)] \\ &= P(W_1 \cap W_2) + P(\bar{W}_1 \cap W_2) \\ &= P(W_1) \cdot P(W_2 | W_1) + P(\bar{W}_1) \cdot P(W_2 | \bar{W}_1) \\ &= \frac{4}{7} \cdot \frac{1}{2} + \frac{3}{7} \cdot \frac{4}{6} = \frac{2}{7} + \frac{2}{7} = \frac{4}{7}. \end{aligned}$$

(ii)
$$\begin{aligned} P(W_1 | W_2) &= \frac{P(W_1 \cap W_2)}{P(W_2)} \\ &= \frac{\frac{2}{7}}{\frac{4}{7}} = \frac{1}{2}. \end{aligned}$$

[J] **Partition of a set.** Let there be a set A ;

$$A = \{A_1, A_2, \dots, A_n\}.$$

Then, the partition of the set A is the set $\{A_1, A_2, \dots, A_n\}$ provided

(i) $A_j \subseteq A$ for $j=1, 2, \dots, n$ i.e., $A_1, A_2, \dots, A_n$ are subsets of A (i.e., they are inclusive).

(ii) $A_j \cap A_k = \phi$ for $j=1, 2, \dots, n, k=1, 2, \dots, n$ and $j \neq k$ i.e. they are disjoint.

(iii) $A_1 \cup A_2 \cup \dots \cup A_n = A$ i.e. they are exhaustive.

We can see that every element of A is a member of one and only one of the subsets in the partition.

Further we can see that if S be a sample space, $S = \{A, \bar{A}\}$,

where

$$A = \{A \cap \bar{B}, A \cap B\}$$

and

$$A \cup B = \{A \cap \bar{B}, A \cap B, \bar{A} \cap B\}$$

then

$$S = \{A, \bar{A}\} = \{A \cap \bar{B}, A \cap B, \bar{A} \cap B, A \cap \bar{B}\}$$

Illustration. Suppose there is a pack of cards, then

$$S = (52 \text{ outcomes})$$

$$= \{A_S, A_H, A_D, A_C\}.$$

where S, H, D, C stand for spade, heart, diamond and club respectively and $A_S = A_H = A_D = A_C = 13$.

It is clear that

(i) $A_j \subseteq S$ for $j=S, H, D, C$, i.e. A_S, A_H, A_D, A_C are subsets of S .

(ii) $A_j \cap A_k = \phi$ for $j=S, H, D, C, k=S, H, D, C$, but, $j \neq k$, i.e., they are disjoint.

(iii) $A_S \cup A_H \cup A_D \cup A_C = S$ i.e. they are exhaustive.

All the three conditions being satisfied the set $\{A_S, A_H, A_D, A_C\}$ represents the partition of S .

[K] **Baye's theorem.** In order to prove the theorem given by Thomas Baye, let us first introduce a Lemma, required for its proof.

Lemma. Let $\{A_1, A_2, \dots, A_n\}$ be a partition of the sample space S and suppose that each of the events $A_1, A_2, \dots, A_n$ has non-zero probability, i.e., $P(A_j) > 0$ for $j=1, 2, \dots, n$. Then for any event A , we have

$$\begin{aligned} P(A) &= P(A_1) \cdot P(A | A_1) + P(A_2) \cdot P(A | A_2) + \dots + P(A_n) \cdot P(A | A_n) \\ &= \sum_{j=1}^n P(A_j) P(A | A_j). \end{aligned}$$

Its proof. As $A_1, A_2, \dots, A_n$ are partitions of S , therefore

$$\{A \cap A_1, A \cap A_2, \dots, A \cap A_n\}$$

will represent the partition of A .

$$\text{Thus } A = (A \cap A_1) \cup (A \cap A_2) \cup \dots \cup (A \cap A_n).$$

so that $P(A) = P(A \cap A_1) + P(A \cap A_2) + \dots + P(A \cap A_n)$

$$= \sum_{j=1}^n P(A \cap A_j)$$

Applying the result of the conditional probability, i.e.,

$$\begin{aligned} P(A | A_j) &= \frac{P(A \cap A_j)}{P(A_j)} \text{ giving } P(A \cap A_j) \\ &= P(A_j) \cdot P(A | A_j) \end{aligned}$$

$$\text{we have } P(A) = \sum_{j=1}^n P(A_j) \cdot P(A | A_j), \quad \dots (1)$$

which proves the lemma.

Statement of Baye's theorem. If an event A can occur only if one of the mutually exclusive events $A_1, A_2, \dots, A_n$, i.e., $A \subseteq \bigcup_{k=1}^n A_k, A_k \cap A_j = \phi$ when $k \neq j$ and suppose we are given the probabilities $P(A_k)$, $k=1, 2, \dots, n$ and the conditional probabilities $P(A | A_k)$, then we are required to find the probability of A_k when it is given that A has already occurred and $P(A) > 0$ for each integer k ($1 \leq k \leq n$) then Baye's formula is

$$P(A_k | A) = \frac{P(A_k) P(A | A_k)}{\sum_{j=1}^n P(A_j) P(A | A_j)}.$$

Proof. By the definition of conditional probability, we have

$$P(A_k | A) = \frac{P(A \cap A_k)}{P(A)}$$

i.e.,

$$\begin{aligned} P(A \cap A_k) &= P(A) P(A_k | A) \\ &= P(A_k) P(A | A_k) \end{aligned}$$

and by (1),

$$P(A) = \sum_{j=1}^n P(A_j) \cdot P(A | A_j).$$

$\therefore$

$$P(A_k | A) = \frac{P(A_k) \cdot P(A | A_k)}{\sum_{j=1}^n P(A_j) \cdot P(A | A_j)}.$$

Problem 40. There are three coins, identical in appearance, one of which is ideal and the other two biased with probabilities $\frac{1}{3}$ and $\frac{1}{4}$ respectively for a head. One coin is taken at random and tossed twice. If a head appears both the times, what is the probability that the ideal coin was chosen.

Let A_1, A_2, A_3 denote the events of choosing the 1st (ideal) 2nd and 3rd coins respectively. Then

$$P(A_1) = P(A_2) = P(A_3) = \frac{1}{3}.$$

Let A be the event of obtaining 2 heads in two tosses of the selected coin.

Probability of getting a head in a toss = $\frac{1}{2}$.

$\therefore$

$$P(A | A_1) = \left(\frac{1}{2}\right)^2 = \frac{1}{4}.$$

Probability of turning head up with 2nd coin is $\frac{1}{3}$ and that with 3rd coin is $\frac{2}{3}$; therefore in two tosses,

$$P(A | A_2) = \left(\frac{1}{3}\right)^2 = \frac{1}{9} \text{ and } P(A | A_3) = \left(\frac{2}{3}\right)^2 = \frac{4}{9}.$$

Using Baye's formula,

$$\begin{aligned} P(A_1 | A) &= \frac{P(A_1) P(A | A_1)}{\sum_{j=1}^3 P(A_j) \cdot P(A | A_j)} \\ &= \frac{\frac{1}{3} \cdot \frac{1}{4}}{\frac{1}{3} \cdot \frac{1}{4} + \frac{1}{3} \cdot \frac{1}{9} + \frac{1}{3} \cdot \frac{4}{9}} \\ &= \frac{\frac{1}{12}}{\frac{1}{12} + \frac{1}{27} + \frac{4}{27}} = \frac{\frac{1}{12}}{\frac{1}{12} + \frac{5}{27}} = \frac{1}{12} \times \frac{27}{27+20} = \frac{9}{47}. \end{aligned}$$

Problem 41. In a bolt factory, machines A, B, C manufacture respectively 25, 35 and 40% of the total. Of their output, 5, 4 and 2% are defective bolts. A bolt is drawn at random from the procedure and is found defective. What are the probabilities that it was manufactured by machine A, B or C ?

Here $P(A) = \frac{25}{100}, P(B) = \frac{35}{100}, P(C) = \frac{40}{100}$.

Let E denote the event of drawing a defective bolt.

$$P(E | A) = \frac{5}{100}, P(E | B) = \frac{4}{100}, P(E | C) = \frac{2}{100}.$$

Using Baye's formula,

$$\begin{aligned} P(A | E) &= \frac{P(A) P(E | A)}{P(A) P(E | A) + P(B) P(E | B) + P(C) P(E | C)} \\ &= \frac{\frac{25}{100} \times \frac{5}{100}}{\frac{25}{100} \times \frac{5}{100} + \frac{35}{100} \times \frac{4}{100} + \frac{40}{100} \times \frac{2}{100}} \end{aligned}$$

$$= \frac{125}{125 + 140 + 180} = \frac{125}{445}.$$

Similarly $P(B | E) = \frac{140}{445}$ and $P(C | E) = \frac{180}{445}$.

[L] **Independent events.** The two events A and B are said to be stochastically independent if and only if $P(A | B) = P(A)$, i.e., the happening of the event B does not affect the happening of A .

We have

$$P(A | B) = \frac{P(A \cap B)}{P(B)} \text{ and } P(B | A) = \frac{P(B \cap A)}{P(A)}$$

The two events are independent to each other when

$$P(A | B) = P(A), P(A) > 0,$$

$$P(B | A) = P(B), P(B) > 0.$$

Thus the two events A and B are stochastically independent if and only if

$$P(A \cap B) = P(A) \cdot P(B).$$

In general, $m(m > 2)$ events $A_1, A_2, \dots, A_m$ are independent if

$$P(A_1 \cap A_2 \cap \dots \cap A_m) = P(A_1) \cdot P(A_2) \dots P(A_m).$$

Problem 42. If A and B are two independent events in a sample space S , then prove that

(i) $\bar{A}$ and $\bar{B}$ are independent,

(ii) $\bar{A}$ and B are independent,

(iii) A and $\bar{B}$ are independent,

(iv) $P(A \cup B) = 1 - P(\bar{A}) P(\bar{B})$.

(i) We have $P(\bar{A} \cap \bar{B}) = 1 - P(A \cup B)$

$$= 1 - [P(A) + P(B) - P(A \cap B)]$$

$$= 1 - P(A) - P(B) + P(A) \cdot P(B)$$

as A and B are independent

$$= [1 - P(A)] [1 - P(B)]$$

$$= P(\bar{A}) \cdot P(\bar{B})$$

showing that $\bar{A}$ and $\bar{B}$ are independent.

(ii) We have $P(\bar{A} \cap B) = P(B) - P(A \cap B)$

$$= P(B) - P(A) \cdot P(B)$$

$$= P(B) [1 - P(A)]$$

$$= P(B) \cdot P(\bar{A})$$

showing that $\bar{A}$ and B are independent.

(iii) We have $A = A \cap \bar{B} \cup A \cap B$.

$$\therefore P(A) = P(A \cap \bar{B}) + P(A \cap B)$$

$$= P(A \cap \bar{B}) + P(A) \cdot P(B)$$

or
$$P(A \cap \bar{B}) = P(A) [1 - P(B)]$$

$$= P(A) \cdot P(\bar{B})$$

showing that A and $\bar{B}$ are independent.

(iv) We have $(A \cup B) \cup (\bar{A} \cap \bar{B}) = S$.

$$\therefore P(A \cup B) + P(\bar{A} \cap \bar{B}) = P(S) = 1$$

or $P(A \cup B) = 1 - P(\bar{A} \cap \bar{B})$

$$= 1 - P(\bar{A}) \cdot P(\bar{B}) \text{ as } \bar{A} \text{ and } \bar{B} \text{ are independent by (i).}$$

Problem 43. From a pack of cards, if the event of drawing a spade card is denoted by A and that of drawing an honour card by B , then show that A and B are independent events.

We have

$$P(A) = \frac{{}^{13}C_1}{{}^{52}C_1} = \frac{1}{4},$$

$$P(B) = \frac{{}^{12}C_1}{{}^{52}C_1} = \frac{1}{13} = \frac{4}{52},$$

$$P(A \cap B) = \frac{{}^3C_1}{{}^{52}C_1} = \frac{3}{52}.$$

Then

$$P(A | B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{3}{52}}{\frac{4}{52}} = \frac{3}{4} = P(A).$$

Since $P(A | B) = P(A)$, the events A and B are independent.

15.7. REPEATED TRIALS: BINOMIAL AND MULTINOMIAL EXPANSIONS

[A] Binomial theorem. If the probability of the happening of an event in one trial is known, then it is required to find the probability of its happening once, twice, thrice . . . exactly in n trials.

Let p be the probability of the happening of the event in a single trial and let q be the probability of its failure, such that $q = 1 - p$. Suppose we have to find the probability of exactly r successes in n trials.

There are n trials in all. Out of these n trials r can be taken in nC_r ways.

Now the chance that the event happens in r trials and fails in the remaining $(n-r)$ trials $= p.p. \dots r \text{ times} \times q.q \dots (n-r) \text{ times} = p^r q^{n-r}$.

$\therefore$ the chance of exactly r successes in n trials $= {}^nC_r \cdot p^r q^{n-r}$.

Putting $r = 1, 2, 3, \dots$ in succession we get the probabilities of exactly 1, 2, 3, . . . successes.

But we know that ${}^nC_r p^r q^{n-r}$ is the $(r+1)$ th term in the binomial expansion of $(q+p)^n$; hence the probability that the event will happen exactly r times in n trials is the $(r+1)$ th term in the binomial expansion of $(q+p)^n$.

[B] Multinomial theorem. If there are n dice with faces marked from 1 to f and these are thrown at random, then we have to find the chance that the sum of the numbers exhibited on the uppermost faces is equal to p .

Since a die has f faces, it can fall in f ways. Similarly the second die can fall in f ways. Thus the two dice can fall together in $f \times f$ i.e., f^2 ways. Similarly we can show that when the n dice fall together, they can fall in f^n ways.

Now the number of ways in which the numbers thrown will have p for their sum is equal to the coefficient of x^p in the expansion of

$$(x^1 + x^2 + x^3 + \dots + x^f)^n$$

because this coefficient arises out of the different ways in which n of the indices 1, 2, 3, . . . , f can be taken so as to form p by addition.

Hence the required probability

$$= \frac{\text{Coef. of } x^p \text{ in } (x^1 + x^2 + \dots + x^f)^n}{f^n}$$

$$= \frac{\text{Coef. of } x^p \text{ in } x^n \cdot \left[\frac{1-x^f}{1-x} \right]^n}{f^n}$$

$$= \frac{\text{Coef. of } x^{p-n} \text{ in } (1-x^f)^n (1-x)^{-n}}{f^n}$$

Problem. 44. What is the chance that a person with two dice (the faces of each being numbered 1 to 6) will throw aces exactly 4 times in 6 trials.

Probability of throwing an ace with a die $= \frac{1}{6}$.

$\therefore$ probability of throwing an ace with two dice in a single throw

$$= \frac{1}{6} \times \frac{1}{6} = \frac{1}{36}.$$

so that the probability of not throwing an ace in a throw

$$= 1 - \frac{1}{36} = \frac{35}{36}.$$

The required chance of throwing exactly four aces in 6 trials is $(4+1)$ th i.e., 5th term in expansion of $(\frac{35}{36} + \frac{1}{36})^6$.

$$\therefore \text{the required chance} = {}^6C_4 \left(\frac{35}{36}\right)^2 \left(\frac{1}{36}\right)^4.$$

Problem 45. An experiment succeeds twice as often as it fails. Find the chance that in the next six trials there will be at least 4 successes.

Let p be the probability of success and q that of failure. Then

$$p = 2q \text{ and } p + q = 1$$

which give

$$p = \frac{2}{3}, q = \frac{1}{3}.$$

The probabilities of 4, 5 and 6 successes are the 5th, 6th and 7th terms in the expansion of $(\frac{2}{3} + \frac{1}{3})^6$ i.e., they are ${}^6C_4 (\frac{2}{3})^2 (\frac{1}{3})^4$, ${}^6C_5 (\frac{2}{3})^1 (\frac{1}{3})^5$ and ${}^6C_6 (\frac{2}{3})^0 (\frac{1}{3})^6$.

The three events being mutually exclusive, the required chance

$$= {}^6C_4 \left(\frac{2}{3}\right)^2 \left(\frac{1}{3}\right)^4 + {}^6C_5 \left(\frac{2}{3}\right)^1 \left(\frac{1}{3}\right)^5 + {}^6C_6 \left(\frac{1}{3}\right)^6$$

$$= \frac{2^4}{729} (15 + 6 \times 2 + 4) = \frac{496}{729}.$$

Problem 46. An ordinary six-faced die is thrown 4 times. What are the probabilities of obtaining 4, 3, 2, 1, 0 aces?

Let p be the chance of obtaining an ace in a single throw and q that of not obtaining an ace. Then

$$p = \frac{1}{6} \text{ and } q = 1 - \frac{1}{6} = \frac{5}{6}.$$

When a die is thrown 4 times, the chances of 4, 3, 2, 1, 0 aces are the 5th, 4th, 3rd, 2nd and 1st terms in the binomial expansion of $(q+p)^4$ i.e., $(\frac{5}{6} + \frac{1}{6})^4$.

$$\therefore \text{Probability of 4 aces} = {}^4C_4 \left(\frac{1}{6}\right)^4 = \frac{1}{1296}.$$

$$,, \quad 3 \quad ,, = {}^4C_3 \left(\frac{1}{6}\right)^3 \left(\frac{5}{6}\right) = \frac{5}{1296}.$$

$$,, \quad 2 \quad ,, = {}^4C_2 \left(\frac{1}{6}\right)^2 \left(\frac{5}{6}\right)^2 = \frac{15}{1296}.$$

$$,, \quad 1 \quad ,, = {}^4C_1 \left(\frac{1}{6}\right)^1 \left(\frac{5}{6}\right)^3 = \frac{125}{1296}.$$

$$,, \quad 0 \quad ,, = {}^4C_0 \left(\frac{5}{6}\right)^4 = \frac{625}{1296}.$$

Problem 47. If m things are distributed among a men and b women, show that the chance that the number of things received by men is odd,

$$\text{is } \frac{1}{2} \frac{(b+a)^m - (b-a)^m}{(b+a)^m}.$$

The probability that a men get a thing = $\frac{a}{a+b}$

,, a women ,, = $\frac{b}{a+b}$.

If out of m things ' a ' men get only one thing and ' b ' women get the remaining things, then the probability for men is 2nd term in the expansion of $\left(\frac{b}{a+b} + \frac{a}{a+b}\right)^m$ i.e., it is ${}^mC_1 \left(\frac{b}{a+b}\right)^{m-1} \frac{a}{a+b}$.

Similarly the probabilities that ' a ' men get three things, five things, ... are respectively

$${}^mC_3 \left(\frac{b}{a+b}\right)^{m-3} \left(\frac{a}{a+b}\right)^3, {}^mC_5 \left(\frac{b}{a+b}\right)^{m-5} \left(\frac{a}{a+b}\right)^5, \dots$$

Hence the probability that the number of things received by men is odd, is

$$= {}^mC_1 \left(\frac{b}{a+b}\right)^{m-1} \frac{a}{a+b} + {}^mC_3 \left(\frac{b}{a+b}\right)^{m-3} \left(\frac{a}{a+b}\right)^3 + {}^mC_5 \left(\frac{b}{a+b}\right)^{m-5} \left(\frac{a}{a+b}\right)^5 + \dots$$

$$= \frac{1}{(a+b)^m} [{}^mC_1 a b^{m-1} + {}^mC_3 a^3 b^{m-3} + {}^mC_5 a^5 b^{m-5} + \dots]$$

$$= \frac{1}{2} \frac{(b+a)^m - (b-a)^m}{(b+a)^m}.$$

Problem 48. If n biscuits be distributed at random among N beggars, what is the chance that a particular beggar receives r ($< n$) biscuits.

The total number of ways in which n biscuits can be distributed among N beggars = N^n .

The specified beggar getting r biscuits in nC_r ways, the remaining $(n-r)$ biscuits among $(N-1)$ beggars can be distributed in $(N-1)^{n-r}$ ways.

$$\therefore \text{Number of favourable ways} = {}^nC_r (N-1)^{n-r}.$$

$$\text{Hence the required probability} = \frac{{}^nC_r (N-1)^{n-r}}{N^n}.$$

Problem 49. A , B and C in order toss a coin. The first one to throw a head wins. What are their respective chances of winning? Assume that the game may continue indefinitely.

Probability of throwing a head with a coin = $\frac{1}{2}$.

$\therefore$ Probability of not throwing a head with a coin

$$= 1 - \frac{1}{2} = \frac{1}{2}.$$

A will have to win if he throws head in 1st, 4th, 7th, ... throws. The probabilities of these events are given by $\frac{1}{2}$, $(\frac{1}{2})^3 \cdot \frac{1}{2}$, $(\frac{1}{2})^6 \cdot \frac{1}{2}$, ... respectively.

$$\therefore A's \text{ chance of winning} = \frac{1}{2} + (\frac{1}{2})^3 \cdot \frac{1}{2} + (\frac{1}{2})^6 \cdot \frac{1}{2} + \dots \infty$$

$$= \frac{\frac{1}{2}}{1 - (\frac{1}{2})^3} = \frac{4}{7}.$$

B will have to win if he throws head in 2nd, 5th, 8th, ... throws.

$$\therefore B's \text{ chance of winning} = (\frac{1}{2}) \cdot \frac{1}{2} + (\frac{1}{2})^4 \cdot \frac{1}{2} + (\frac{1}{2})^7 \cdot \frac{1}{2} + \dots \infty$$

$$= \frac{\frac{1}{2} \cdot \frac{1}{2}}{1 - (\frac{1}{2})^3} = \frac{2}{7}.$$

Similarly C will have to win if he throws head in 3rd, 6th, 9th, ... throws.

$$\therefore C's \text{ chance of winning} = (\frac{1}{2})^2 \cdot \frac{1}{2} + (\frac{1}{2})^5 \cdot \frac{1}{2} + (\frac{1}{2})^8 \cdot \frac{1}{2} + \dots \infty$$

$$= \frac{(\frac{1}{2})^2 \cdot \frac{1}{2}}{1 - (\frac{1}{2})^3} = \frac{1}{7}.$$

Hence the chances of A , B , C are $\frac{4}{7}$, $\frac{2}{7}$ and $\frac{1}{7}$ respectively.

Problem 50. A , B , C and D cut a pack of cards successively in the order mentioned. What are their respective chances of first cutting a spade?

The chance of cutting a spade = $\frac{1}{4}$.

$\therefore$ the chance of not cutting a spade = $1 - \frac{1}{4} = \frac{3}{4}$.

A wins if he cuts a spade in 1st, 5th, 9th, ... throws.

$$\therefore A's \text{ chance of cutting a spade}$$

$$= \frac{1}{4} + (\frac{3}{4})^4 \cdot \frac{1}{4} + (\frac{3}{4})^8 \cdot \frac{1}{4} + \dots \infty$$

$$= \frac{\frac{1}{4}}{1 - (\frac{3}{4})^4} = \frac{64}{165}.$$

B wins if he cuts a spade in 2nd, 6th, 10th, ... throws.

$$\therefore B's \text{ chance of cutting a spade}$$

$$= \frac{3}{4} \cdot \frac{1}{4} + (\frac{3}{4})^5 \cdot \frac{1}{4} + (\frac{3}{4})^9 \cdot \frac{1}{4} + \dots \infty$$

$$= \frac{\frac{3}{4} \cdot \frac{1}{4}}{1 - (\frac{3}{4})^4} = \frac{48}{175}.$$

Similarly C 's chance of cutting a spade

$$= \left(\frac{3}{4}\right)^2 \cdot \frac{1}{4} + \left(\frac{3}{4}\right)^6 \cdot \frac{1}{4} + \left(\frac{3}{4}\right)^{10} \cdot \frac{1}{4} + \dots \infty$$

$$= \frac{\left(\frac{3}{4}\right)^2 \cdot \frac{1}{4}}{1 - \left(\frac{3}{4}\right)^4} = \frac{36}{175}$$

and D 's chance of cutting a spade

$$= \left(\frac{3}{4}\right)^3 \cdot \frac{1}{4} + \left(\frac{3}{4}\right)^7 \cdot \frac{1}{4} + \left(\frac{3}{4}\right)^{11} \cdot \frac{1}{4} + \dots$$

$$= \frac{\left(\frac{3}{4}\right)^3 \cdot \frac{1}{4}}{1 - \left(\frac{3}{4}\right)^4} = \frac{27}{175}$$

Problem 51. Find the chance of throwing 10 exactly in one throw with three dice.

Three dice can be thrown in 6^3 , i.e. 216 ways.

The number of ways of getting 10 by throwing 3 dice

$$= \text{coeff. of } x^{10} \text{ in } (x+x^2+\dots+x^6)^3$$

$$= \text{coeff. of } x^{10} \text{ in } \frac{x^3(1-x^6)^3}{(1-x)^3}$$

$$= \text{coeff. of } x^{10} \text{ in } x^3(1-3x^6+3x^{12}-x^{18})(1-x)^{-3}$$

$$= \text{coeff. of } x^{10} \text{ in } x^3(1-3x^6+3x^{12}-x^{18})(1+3x+6x^2-10x^3+15x^4+21x^5+24x^6+36x^7+\dots)$$

$$= 36 - 9 = 27.$$

$$\therefore \text{the required probability} = \frac{27}{216} = \frac{1}{8}.$$

Aliter. Favourable number of ways may be found out as below.

Dice			Number of ways
1st	2nd	3rd	
6	2	2	$\frac{3!}{2!} = 3$
6	3	1	$3! = 6$
5	3	2	$3! = 6$
5	4	1	$3! = 6$
4	3	3	$\frac{3!}{2!} = 3$
4	4	2	$\frac{3!}{2!} = 3$
			<u>Total 27</u>

$$\therefore \text{required chance} = \frac{27}{216} = \frac{1}{8}.$$

Problem 52. Four dice are thrown, what is the probability that the sum of the numbers appearing on the dice is 18?

4 dice can be thrown in 6^4 ways.

Favourable number of ways to give a sum of 18 with 4 dice

$$= \text{coeff. of } x^{18} \text{ in } (x+x^2+\dots+x^6)^4$$

$$= \text{coeff. of } x^{18} \text{ in } x^4 \left(\frac{1-x^6}{1-x} \right)^4$$

$$= \text{coeff. of } x^{18} \text{ in } x^4 (1-x^6) (1-x)^{-4}$$

$$= \text{coeff. of } x^{18} \text{ in } x^4 (1-4x^6+6x^{12}-\dots) (1+4x+10x^2+\dots+165x^8+\dots+680x^{14}+\dots)$$

$$= 680 - 660 + 60 = 80.$$

$$\therefore \text{required probability} = \frac{80}{6^4} = \frac{5}{81}.$$

Problem 53. Find the chance of throwing 10 with 4 dice.

Proceeding just as above, the required probability is

$$\frac{104}{6^4} = \frac{13}{162}$$

Problem 54. Determine the probability of throwing more than 8 with perfectly symmetrical dice.

Total number of ways in which 3 dice can be thrown $= 6^3 = 216$.

The favourable number of getting the sum as 3, 4, 5, 6, 7, 8 will be equal to the sum of coefficients of $x^3, x^4, x^5, x^6, x^7, x^8$ in the expansion of $(x+x^2+\dots+x^6)^3$,

$$\text{i.e., of } x^2(1-x^6)^3(1-x)^{-3},$$

$$\text{i.e., of } x^3(1-3x^6+\dots)(1+3x+6x^2+10x^3+15x^4+21x^5+\dots)$$

$$= 1+3+6+10+15+21=56.$$

$$\therefore \text{The probability of getting the sum} \leq 8 = \frac{56}{6^3} = \frac{7}{27}.$$

$$\text{Hence the probability of getting the sum} > 8 = 1 - \frac{7}{27} = \frac{20}{27}.$$

Problem 55. Counters marked 1, 2, 3 are placed in a bag; one is withdrawn and replaced three times. What is the chance of obtaining a total of 6?

Total number of ways of drawing a counter three times $= 3^3 = 27$.

Favourable number of getting a sum of 6

$$= \text{the coeff. of } x^6 \text{ in } (x+x^2+x^3)^3$$

$$= \text{coeff. of } x^6 \text{ in } x^3(1-x^3)^3(1-x)^{-3}$$

$$= \text{coeff. of } x^3 \text{ in } (1-3x^3+\dots)(1+3x+6x^2+10x^3+\dots)$$

$$= 10 - 3 = 7.$$

$$\therefore \text{Required chance} = \frac{7}{27}.$$

Problem 56. Nine cards are drawn at random from a set of cards. Each card is marked with one of the numbers 1, 0 or -1 and it is equally likely that any of the three numbers will be drawn. Find the chance that the sum of the numbers drawn is zero.

The total number of ways in which 9 cards can be drawn $= 3^9$.

The favourable number of ways of getting a sum of zero

$$\begin{aligned}
&= \text{the coeff. of } x^0 \text{ in } (x^{-1} + x^0 + x^1)^9 \\
&= \quad \quad \quad (x^{-1} + 1 + x)^9 \\
&= \quad \quad \quad \frac{1}{x^9} (1 + x + x^2)^9 \\
&= \quad \quad \quad \frac{1}{x^9} (1 - x^3)^9 (1 + x)^{-9} \\
&= \quad \quad \quad \frac{1}{x^9} (1 - 9x^3 + 36x^6 - 84x^9 + \dots) \times (1 + 9x \\
&\quad \quad \quad + 45x^2 + 165x^3 + \dots + 3003x^6 + \dots + 24310x^9 + \dots) \\
&= 24310 - 9 \times 3003 + 165 \times 36 - 84 \\
&= 3139 \\
\therefore \text{ Required chance} &= \frac{3139}{(3)^9}.
\end{aligned}$$

15.8. MEASURES OF CENTRAL TENDENCY

[A] **Arithmetic mean or simply mean.** Let $x_1, x_2, x_3, \dots, x_n$ be the n values of a variate (or variable) x ; then their arithmetic mean denoted by m or M or $\bar{x}$ is defined to be

$$\bar{x} = \frac{x_1 + x_2 + x_3 + \dots + x_n}{n}$$

or using the sigma (Σ) notation,

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n} \text{ or simply } \frac{\Sigma x}{n} \quad \dots (1)$$

In case the 'weights' $w_1, w_2, w_3, \dots, w_n$ are attached to the n variate-values $x_1, x_2, x_3, \dots, x_n$ respectively, then the 'weighted mean' is defined to be

$$\bar{x} = \frac{w_1 x_1 + w_2 x_2 + \dots + w_n x_n}{w_1 + w_2 + w_3 + \dots + w_n} = \frac{\sum_{i=1}^n w_i x_i}{\sum_{i=1}^n w_i} = \frac{\Sigma w x}{\Sigma w} \quad \dots (2)$$

In other words if the value x_1 occurs f_1 times, x_2 occurs f_2 times, and so on, then

$$\bar{x} = \frac{f_1 x_1 + f_2 x_2 + \dots + f_n x_n}{f_1 + f_2 + \dots + f_n} = \frac{\sum_{i=1}^n f_i x_i}{\sum_{i=1}^n f_i} = \frac{\Sigma f x}{\Sigma f} \quad \dots (3)$$

$$\text{or} \quad \bar{x} = \frac{1}{N} \sum_{i=1}^n f_i x_i = \frac{\Sigma f x}{N} \quad \dots (4)$$

$$\text{where } N = f_1 + f_2 + \dots + f_n = \text{total frequency.} \quad \dots (5)$$

If we assume that A is any assumed mean, whose deviation from the variate x is ξ , then if M is the arithmetic mean, we have

$$\xi = x - A \text{ and } M = \frac{1}{N} \Sigma f x \text{ by (4).}$$

$$\text{Consider } \frac{1}{N} \Sigma f \xi = \frac{1}{N} \Sigma f (x - A)$$

$$= \frac{1}{N} \Sigma f x - \frac{1}{N} \Sigma f A$$

$$= M - A \text{ since from (5) } \Sigma f = N.$$

$$\therefore M = A + \frac{1}{N} \Sigma f \xi \quad \dots (6)$$

This result used as a short cut method for finding the arithmetic mean, often facilitates calculation.

Again if h be the width of equal class-intervals in a frequency table and u be the new variate defined as

$$u = \frac{x - A}{h} \text{ i.e., } x = A + uh.$$

then, we have

$$\begin{aligned} \Sigma f x &= \Sigma f (A + uh) \\ &= \Sigma f A + h \Sigma f u. \end{aligned}$$

Dividing either side by Σf , we find

$$\frac{\Sigma f(x)}{\Sigma f} = A + h \frac{\Sigma f u}{\Sigma f}$$

$$\text{i.e., } \bar{x} \text{ or } M = A + h \bar{u} \quad \dots (7)$$

where $\bar{u} = \frac{\Sigma f u}{\Sigma f}$, the arithmetic mean for values of the variate u .

Properties of the Arithmetic Mean

I. The algebraic sum of the deviations of the variate-values from their mean is zero.

Let ξ be the deviation of the variate x from the mean $\bar{x}$, then the sum of the deviations of the variate-values from the mean

$$\begin{aligned}
&= \Sigma f (x - \bar{x}) \\
&= \Sigma f x - \Sigma f \bar{x} \\
&= N \bar{x} - N \bar{x} \text{ from (4) and (5)} \\
&= 0.
\end{aligned}$$

II. If $\bar{x}_1, \bar{x}_2, \dots, \bar{x}_r$ are the arithmetic means of r distributions with respective frequencies $N_1, N_2, \dots, N_r$, then the mean $\bar{x}$ of the whole distribution with total frequency $N = \sum_{i=1}^r N_i$ is given by

$$\bar{x} = \frac{1}{N} \sum_{i=1}^r N_i \bar{x}_i.$$

Let f_{ij} denote the frequency of the observation x_{ij} of the i th distribution, so that $N_i = \sum_{j=1}^n f_{ij}$; then

$$\bar{x}_i = \frac{\sum_{j=1}^n f_{ij} x_{ij}}{\sum_{j=1}^n f_{ij}} = \frac{\sum_{j=1}^n f_{ij} x_{ij}}{N_i}.$$

Now $\bar{x}$ being the mean of the whole distribution, we have

$$\bar{x} = \frac{\sum_{ij} f_{ij} x_{ij}}{\sum_{ij} f_{ij}} = \frac{\sum_i N_i \bar{x}_i}{\sum_i N_i} = \frac{1}{N} \sum_{i=1}^i N_i \bar{x}_i.$$

Problem 57. The following table gives the population of males at different age groups of the U.K. and India at the time of the census of 1931.

Age group (years)	U.K. (in lakhs)	India (in lakhs)
0-5	18	214
5-10	19	258
10-15	20	222
15-20	18	157
20-25	16	145
25-30	14	161
30-40	27	257
40-50	25	184
50-60	19	120
above 60	17	100

compare the average age of males in the two countries and account for the difference, if any.

Take the assumed mean as 27.5.

Age group (years)	Mid-values (x)	Deviation from assumed mean 27.5 (ξ)	U.K.		India	
			Population in lakhs (f_1)	$f_1 \xi$	Population in lakhs (f_2)	$f_2 \xi$
0-5	2.5	-25.0	18	-450	214	-5350.0
5-10	7.5	-20.0	19	-380	258	-5160.0
10-15	12.5	-15.0	20	-300	222	-3330.0
15-20	17.5	-10.0	18	-180	157	-1570.0
20-25	22.5	-5.0	16	-80	145	-725.0
25-30	27.5	0	14	0	161	0
30-40	35.0	7.5	27	202.5	257	1927.5
40-50	45.0	17.5	25	437.5	180	3220.0
50-60	55.0	27.5	19	422.5	124	3300.0
above 60	65.0	37.5	17	637.5	100	3750.0
Total			193 = N_1 = $\sum f_1$	410 = $\sum f_1 \xi$	1818 = N_2 = $\sum f_2$	-3937.5 = $\sum f_2 \xi$

For U.K., average age of the people

$$= A + \frac{\sum f_1 \xi}{\sum f_1} = 27.5 + \frac{410}{1818} = 27.5 + 2.12 = 29.62.$$

For India, average age of the people

$$= A + \frac{\sum f_2 \xi}{\sum f_2} = 27.5 - \frac{3937.5}{1818} = 27.5 - 2.17 = 25.33.$$

These averages show that the average age of the people of U.K. is higher than that of the people in India.

Problem 58. If a variate x is expressed as a linear function of two variates u and v in the form $x = au + bv$, show that

$$\bar{x} = a\bar{u} + b\bar{v}.$$

Let the number of variables for each u and v be n . Then

$$x_i = au_i + bv_i,$$

so that

$$\sum_{i=1}^n x_i = a \sum_{i=1}^n u_i + b \sum_{i=1}^n v_i$$

or

$$\frac{\sum x_i}{n} = a \frac{\sum u_i}{n} + b \frac{\sum v_i}{n}$$

i.e.,

$$\bar{x} = a\bar{u} + b\bar{v}.$$

Problem 59. Show that if $\bar{x}$ is the arithmetic mean of the values x_i , $i=1, 2, \dots, n$, then

$$\sum_{i=1}^n f_i (x_i - \bar{x})^2 = \sum_{i=1}^n f_i x_i^2 - N\bar{x}^2$$

where $N = \sum_{i=1}^n f_i$.

$$\frac{\sum f_i x_i}{\sum f_i} = \frac{\sum f_i x_i^2}{N} \quad \dots (1)$$

We have

$$\begin{aligned} \text{Now } \sum_i f_i (x_i - \bar{x})^2 &= \sum_i f_i (x_i^2 + \bar{x}^2 - 2x_i \bar{x}) \\ &= \sum_i f_i x_i^2 + \bar{x}^2 \sum_i f_i - 2\bar{x} \sum_i f_i x_i \\ &= \sum_i f_i x_i^2 + \bar{x}^2 N - 2\bar{x} N\bar{x} \text{ from (1)} \\ &= \sum_i f_i x_i^2 - N\bar{x}^2. \end{aligned}$$

Problem 60. A distribution consists of 3 components with frequencies 45, 40, 65 having their means 2, 2.5 and 2 respectively. Prove that the mean of the combined distribution is 2.13 approximately.

Using the property II of arithmetic mean, the required mean is given by

$$\bar{x} = \frac{1}{N} \sum_{i=1}^3 N_i \bar{x}_i.$$

where $N = 45 + 40 + 65 = 150$.

$$\begin{aligned} \text{and } \sum_{i=1}^3 N_i \bar{x}_i &= N_1 \bar{x}_1 + N_2 \bar{x}_2 + N_3 \bar{x}_3 \\ &= 45 \times 2 + 40 \times 2.5 + 65 \times 2 \\ &= 90 + 100 + 130 \\ &= 320. \end{aligned}$$

$$\therefore \bar{x} = \frac{320}{150} = \frac{32}{15} = 2.13 \text{ approx.}$$

Problem 61. Supposing the frequencies of values $0, 1, 2, \dots, n$ of a variable to be given by the terms of the binomial series

$$q^n, \frac{n}{1} \cdot q^{n-1} p, \frac{n(n-1)}{1 \cdot 2} q^{n-2} p^2, \dots, p^n,$$

where $p+q=1$, find the mean.

Let M be the required mean; then

$$\begin{aligned} M &= \frac{\sum fx}{\sum f} = \frac{0 \cdot q^n + 1 \cdot {}^nC_1 q^{n-1} p + 2 \cdot {}^nC_2 q^{n-2} p^2 + \dots + n \cdot p^n}{q^n + {}^nC_1 q^{n-1} p + {}^nC_2 q^{n-2} p^2 + \dots + np^n} \\ &= \frac{1 \cdot n C_1 q^{n-1} p + 2 \cdot {}^nC_2 q^{n-2} p^2 + \dots + np^n}{(q+p)^n} \end{aligned}$$

But $(q+p)^n = 1$ as $q+p=1$ (given)

$$\therefore M = {}^nC_1 q^{n-1} p + {}^nC_2 q^{n-2} p^2 + \dots + np^n.$$

Now, we have

$$(q+p)^n = q^n + {}^nC_1 q^{n-1} p + {}^nC_2 q^{n-2} p^2 + \dots + p^n.$$

Differentiating both sides w.r.t. p and then multiplying through-out by p we get

$$\begin{aligned} n(p+q)^{n-1} p &= {}^nC_1 q^{n-1} p + {}^nC_2 q^{n-2} p^2 \cdot 2 + \dots + np^n \\ &= M \text{ from (1).} \end{aligned}$$

Thus $M = np$ as $p+q=1$.

Problem 62. Show that the arithmetic mean of the series $1, 2, 4, 8, 16, \dots, 2^n$ is $\frac{2^{n+1}-1}{n+1}$

$$\begin{aligned} \text{Required mean} &= \frac{1+2+2^2+\dots+2^n}{n+1} \\ &= \frac{2^{n+1}-1}{n+1}. \end{aligned}$$

Problem 63. Show that the weighted arithmetic mean of first n natural numbers whose weights are equal to the corresponding numbers is equal to $\frac{1}{3}(2n+1)$.

If $\bar{x}$ be the required arithmetic mean, then

$$\bar{x} = \frac{1 \cdot 1 + 2 \cdot 2 + 3 \cdot 3 + \dots + n \cdot n}{1 + 2 + 3 + \dots + n}$$

$$\begin{aligned} &= \frac{\sum n^2}{\sum n} = \frac{n(n+1)(2n+1)/6}{n(n+1)/2} \\ &= \frac{2n+1}{3}. \end{aligned}$$

Problem 64. The arithmetic mean of n numbers of a series is $\bar{x}$. The sum of the first $(n-1)$ terms is k . Show that the n th number is $n\bar{x} - k$.

Let t_n be the n th term of the given series; then

$$\bar{x} = \frac{k + t_n}{n}$$

giving

$$t_n = n\bar{x} - k.$$

[B] Geometric mean. Let $x_1, x_2, x_3, \dots, x_n$ be the n values of a variate x ; then their geometric mean denoted by G is defined to be

$$G = (x_1 \cdot x_2 \cdot x_3 \dots x_n)^{1/n} \quad \dots (1)$$

or

$$\log G = \frac{\sum_{i=1}^n \log x_i}{n} = \frac{\sum \log x}{n}$$

If the value x_1 occurs f_1 times, x_2 occurs f_2 times and so on, then $G = (x_1^{f_1} \cdot x_2^{f_2} \dots x_n^{f_n})^{1/N}$ where $N = f_1 + f_2 + \dots + f_n$

or

$$\log G = \frac{\sum_{i=1}^n f_i \log x_i}{N} = \frac{\sum f \log x}{N} \quad \dots (2)$$

which show that the logarithm of the geometric mean of a series of values is the arithmetic mean of their logarithms.

Properties of Geometric Mean

I. The geometric mean of a series is less than its arithmetic mean, i.e., if A be the arithmetic mean and G the geometric mean of a series, then $A > G$

We prove the result for a series having two numbers only and the result may be extended to any number.

Let x_1, x_2 be the values of a variate x . Then

$$A = \frac{x_1 + x_2}{2} \text{ and } G = \sqrt{x_1 x_2}.$$

Consider

$$A - G = \frac{x_1 + x_2}{2} - \sqrt{x_1 x_2}$$

$$\begin{aligned} &= \frac{1}{2} [\sqrt{x_1} - \sqrt{x_2}]^2 \\ &= a \text{ +ve quantity as } (\sqrt{x_1} - \sqrt{x_2})^2 > 0. \end{aligned}$$

showing that $A > G$.

II. If $G_1, G_2, \dots, G_r$ be the geometric means of r distributions with respective frequencies $N_1, N_2, \dots, N_r$, then the geometric mean G of the whole distribution with total frequency $N = \sum_{i=1}^r N_i$ is given by

$$N \log G = N_1 \log G_1 + N_2 \log G_2 + \dots + N_r \log G_r$$

$$\text{i.e.,} \quad \log G = \frac{1}{N} \sum_{i=1}^r N_i \log G_i.$$

Let f_{ij} denote the frequency of the observation x_{ij} of the i th distribution so that $N_i = \sum_{j=1}^n f_{ij}$, then

$$\log G_i = \frac{\sum_{j=1}^n f_{ij} \log G_{ij}}{\sum_{j=1}^n f_{ij}} = \frac{\sum_{j=1}^n f_{ij} \log G_{ij}}{N_i}$$

But G being the geometric mean of the whole distribution, we have

$$\begin{aligned} \log G &= \frac{\sum_{ij} f_{ij} \log G_{ij}}{\sum_{ij} f_{ij}} = \frac{\sum_i N_i \log G_i}{\sum_i N_i} \\ &= \frac{1}{N} \sum_{i=1}^r N_i \log G_i. \end{aligned}$$

III. If G_1, G_2 are the geometric means of two series of observations and G the geometric mean of the ratios of corresponding observations, then G is equal to the ratio of their geometric means,

$$\text{i.e.,} \quad G = \frac{G_1}{G_2}.$$

Let x_1, x_2 be the two variates corresponding to the two series of observations with frequency n each, and let x be the ratio of the two variates; then we have

$$x = \frac{x_1}{x_2}$$

$$\text{so that} \quad \log x = \log x_1 - \log x_2.$$

$$\therefore \quad \sum \log x = \sum \log x_1 - \sum \log x_2.$$

Dividing throughout by n ,

$$\frac{\sum \log x}{n} = \frac{\sum \log x_1}{n} - \frac{\sum \log x_2}{n}$$

$$\text{i.e.,} \quad \log G = \log G_1 - \log G_2$$

$$\text{giving} \quad G = \frac{G_1}{G_2}$$

IV. The geometric mean G of the product of r sets of observations with geometric means $G_1, G_2, \dots, G_r$ respectively, is the product of the geometric means of the component series, i.e.,

$$G = G_1 G_2 \dots G_r.$$

Let $x_1, x_2, \dots, x_r$ be the variates corresponding to r sets of observations and x their product, i.e.,

$$x = x_1 x_2 \dots x_r;$$

$$\text{then} \quad \log x = \log x_1 + \log x_2 + \dots + \log x_r$$

$$\text{or} \quad \sum \log x = \sum \log x_1 + \sum \log x_2 + \dots + \sum \log x_r$$

$$\text{or} \quad \frac{\sum \log x}{n} = \frac{\sum \log x_1}{n} + \frac{\sum \log x_2}{n} + \dots + \frac{\sum \log x_r}{n}$$

$$\text{i.e.,} \quad \log G = \log G_1 + \log G_2 + \dots + \log G_r$$

$$\text{giving} \quad G = G_1 G_2 \dots G_r.$$

Problem 65. Find the geometric mean of the series.

$$1, 2, 4, 8, 16 \dots 2^n.$$

$$G = \{1.2.2^2.2^3 \dots 2^n\}^{1/(n+1)}$$

$$= \{2^{1+2+3+\dots+n}\}^{1/(n+1)}$$

$$= \{(2)^{n(n+1)/2}\}^{1/(n+1)} = 2^{n/2}.$$

[C] **Harmonic mean.** Let $x_1, x_2, x_3, \dots, x_n$ be the n values of a variate x ; their harmonic mean denoted by H is defined to be the reciprocal of the arithmetic mean of their reciprocals, i.e.,

$$\frac{1}{H} = \frac{\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n}}{n} = \frac{1}{n} \sum_{i=1}^n \left(\frac{1}{x_i} \right) = \frac{1}{n} \sum \frac{1}{x} \quad \dots (1)$$

$$\text{or} \quad H = \frac{n}{\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n}} = \frac{n}{\sum_{i=1}^n \frac{1}{x_i}} = \frac{n}{\sum \frac{1}{x}}.$$

If the value x_1 occurs f_1 times, x_2 occurs f_2 times and so on,

$$\begin{aligned} \text{then } \frac{1}{H} &= \frac{\frac{f_1}{x_1} + \frac{f_2}{x_2} + \dots + \frac{f_n}{x_n}}{N} \quad \text{where } N = \sum_{i=1}^n f_i \\ &= \frac{\sum_{i=1}^n \left(f_i \times \frac{1}{x_i} \right)}{\sum_{i=1}^n f_i} = \frac{\sum f_i \cdot \frac{1}{x}}{N} \quad \dots (2) \end{aligned}$$

Property of harmonic mean. The harmonic mean of a series of quantities is less than the geometric mean of the same quantities and a fortiori (Latin phrase, means 'with stronger or greater reason') less than the arithmetic mean of the same quantities, i.e., if A, G, H be the arithmetic, geometric, and harmonic means of a series, then $A > G > H$.

In property I of geometric mean, we have already proved for two quantities x_1, x_2 that $A > G$.

$$\text{Now consider } G - H = \sqrt{(x_1 x_2)} - \frac{2x_1 x_2}{x_1 + x_2}, \text{ as } H = \frac{2}{\frac{1}{x_1} + \frac{1}{x_2}}$$

$$= \frac{\sqrt{(x_1 x_2)}}{x_1 + x_2} [x_1 + x_2 - 2\sqrt{(x_1 x_2)}]$$

or

$$G-H = \frac{\sqrt{(x_1 x_2)}}{x_1 + x_2} \{ \sqrt{(x_1)} - \sqrt{(x_2)} \}^2$$

$$= a + \text{ve quantity.}$$

 $\therefore G > H$; also $A > G$. $\therefore A > G > H$.This result can be extended to any number of values of x .

Problem 66. A variate takes values $a, ar, ar^2, \dots, ar^{n-1}$ each with frequency unity. Show that the A.M., A is $\frac{a(1-r^n)}{n(1-r)}$, the G.M., G is $ar^{(n-1)/2}$ and the H.M., H is $\frac{a \cdot n(1-r)r^{n-1}}{1-r^n}$. Prove that $AH = G^2$. Prove also that $A > G > H$ unless $n=1$ when all the three means coincide.

We have

$$A = \frac{a + ar + ar^2 + \dots + ar^{n-1}}{n}$$

$$= \frac{a}{n} [1 + r + r^2 + \dots + r^{n-1}]$$

$$= \frac{a}{n} \cdot \frac{1-r^n}{1-r} \text{ if } r < 1.$$

$$G = (a \cdot ar \cdot ar^2 \dots ar^{n-1})^{1/n}$$

$$= [a^n \cdot r^{1+2+\dots+(n-1)}]^{1/n}$$

$$= a \cdot \{r^{(n-1)/2 \cdot n}\}^{1/n} = ar^{(n-1)/2}$$

and

$$H = \frac{n}{\frac{1}{a} + \frac{1}{ar} + \frac{1}{ar^2} + \dots + \frac{1}{ar^{n-1}}}$$

$$= \frac{n}{\frac{1}{a} \left[1 + \frac{1}{r} + \frac{1}{r^2} + \dots + \frac{1}{r^{n-1}} \right]}$$

$$= \frac{an}{\left(\frac{1}{r^n} - 1 \right) / \left(\frac{1}{r} - 1 \right)} \text{ if } r < 1, \text{ i.e., } \frac{1}{r} > 1$$

$$= \frac{an(1-r)r^{n-1}}{1-r^n}$$

Now $AH = \frac{a(1-r^n)}{n(1-r)} \times \frac{an(1-r)r^{n-1}}{1-r^n}$

$$= a^2 r^{n-1} = \{ar^{(n-1)/2}\}^2$$

$$= G^2.$$

Again to prove $A > G > H$, consider

$$A - G = \frac{a}{n} \frac{1-r^n}{1-r} - ar^{(n-1)/2}$$

$$= \frac{a}{n} (1 + r + r^2 + \dots + r^{n-1}) - ar^{(n-1)/2}$$

$$= \frac{a}{n} [(1 + r + r^2 + \dots + r^{n-1}) - n \cdot r^{(n-1)/2}]$$

$$= \frac{a}{n} [(1 - r^{(n-1)/2}) + (r - r^{(n-1)/2}) + (r^2 - r^{(n-1)/2}) + \dots$$

$$+ (r^{n-1} - r^{(n-1)/2})]$$

$$= \frac{a}{n} [(1 - r^{(n-1)/2}) + r(1 - r^{(n-3)/2}) + r^2(1 - r^{(n-5)/2})$$

$$+ \dots + r^{n-1}(1 - r^{(1-n)/2})]$$

$$= a + \text{ve quantity, since every factor in the bracket of R.H.S. expression is +ve when } r < 1.$$

 $\therefore A > G$.But if $n=1$, $A-G=0$, i.e., $A=G$.

Also, $G-H = ar^{(n-1)/2} - \frac{an(1-r)r^{n-1}}{1-r^n}$

$$= ar^{(n-1)/2} - \frac{anr^{n-1}}{1+r+r^2+\dots+r^{n-1}}$$

$$\text{as } \frac{1-r^n}{1-r} = 1+r+r^2+\dots+r^{n-1}$$

$$= \frac{ar^{(n-1)/2}}{1+r+r^2+\dots+r^{n-1}} [1+r+r^2+\dots+r^{n-1} - nr^{(n-1)/2}]$$

$$= \frac{ar^{(n-1)/2}}{1+r+r^2+\dots+r^{n-1}} [(1 - r^{(n-1)/2}) + r(1 - r^{(n-3)/2})$$

$$+ \dots + r^{n-1}(1 - r^{(1-n)/2})]$$

$$\text{as above}$$

$$= a + \text{ve quantity as above.}$$

 $\therefore G > H$.But if $n=1$, $G-H=0$, i.e., $G=H$.Hence $A > G > H$ unless $n=1$ in which case $A=G=H$, i.e., all the three means coincide.

Problem 67. Find the average rates of (a) motion in the case of a person who rides the first mile at 10 m.p.h., the next mile at 8 m.p.h. and the third mile at 6 m.p.h.; (b) increase in population which in the first decade has increased 20% in the next 25% and in the third 44%.

(a) For this case, the harmonic mean being the suitable average, we have

$$H = \frac{3}{\frac{1}{10} + \frac{1}{8} + \frac{1}{6}} = \frac{3 \times 120}{47} = \frac{360}{47} = 7.66 \text{ m.p.h. approx}$$

(b) For this case, the geometric mean being the suitable average, we have

$$G = (20 \times 25 \times 44)^{1/3}.$$

$$\begin{aligned}\therefore \log G &= \frac{1}{3} [\log 20 + \log 25 + \log 44] \\ &= \frac{1}{3} [1.3010 + 1.3979 + 1.6435] \\ &= \frac{1}{3} \times 4.3424 = 1.4475 = \log 28.02. \\ \therefore G &= 28.02\%.\end{aligned}$$

[D] Median and quartiles. The median is defined as the middle-most or central value of the variate when the variate-values are arranged in ascending or descending order of magnitude i.e., it is the value of the variate for which greater and smaller values occur with equal frequency or in other words the total frequency above and below this value is divided into two equal halves.

In case the total frequency n is an odd number, then the value of $\left(\frac{n+1}{2}\right)$ th item gives the median, but if it is an even number then $\left(\frac{n}{2}\right)$ th and $\left(\frac{n}{2} + 1\right)$ th are the central items so that their arithmetic mean gives the median.

For grouped data, the median is formulated as

$$\text{Median} = l_1 + \frac{\frac{N}{2} - f}{f} \times h$$

where l_1 = the lower limit of the median class.

N = the total frequency,

f = the frequency of the median class.

f_1 = the cumulative frequency before entering the median class.

h = the size of the class-interval of the median class.

For a continuous series, the median is formulated as

$$\text{Median} = l_1 + \frac{l_2 - l_1}{f} (m - f_1)$$

where l_1 = the lower limit of the median-class,

l_2 = the upper limit of the median-class,

f = the frequency of the median-class.

f_1 = the cumulative frequency before entering the median class,

m = the size of the middle item.

The *quantiles* or *partition values* are defined to be the values of the variate which divide the total frequency into a number of equal parts. *Quartiles*, *deciles* and *percentiles* are the worth-considerable among partition values.

The *quartiles* are the values of the variate that divide the total frequency into four equal parts. 1st, 2nd, 3rd quartiles are denoted by Q_1 , Q_2 , Q_3 respectively, while Q_2 is the median. We have

$$Q_i = l_1 + \frac{\left(\frac{iN}{4} - f_1\right)}{f} \times h, i = 1, 2, 3$$

where l_1 = the lower limit of the class,

N = the total frequency,

h = the size of the class,

f_1 = the cumulative frequency upto and including the class preceding the class in which the particular quartiles lies.

f = the frequency of this class.

$$\text{Inter-quartile range} = Q_3 - Q_1.$$

The *deciles* are the values of the variate which divide the total frequency into ten equal parts and given by

$$D_j = l_1 + \frac{\left(\frac{jN}{10} - f_1\right)}{f} \times h, j = 1, 2, \dots, 9$$

the other quantities having the same meanings as in quartiles.

The *Percentiles* are the values of the variate which divide the total frequency into hundred equal parts and given by

$$P_k = l_1 + \frac{\left(\frac{kN}{100} - f_1\right)}{f} \times h, k = 1, 2, \dots, 99$$

the other quantities having the same meanings as above.

Problem 68. Show that for *J-shaped* distribution with the maximum frequency towards the lower values of the variate, the median is nearer to Q_1 than Q_3 .

In the adjoining diagram, let M_i stand for the median; then we have to show that M_i is nearer to Q_1 than to Q_3 i.e.

$$M_i - Q_1 < Q_3 - M_i$$

Let $y = f(x)$ be the equation of the curve (*J-shaped*) under consideration and Q_1 , Q_3 , M_i locate the positions of first quartile, third quartile and median respectively.

Now according to the definition of quartiles, the area between Q_1 and median is equal to the area between Median and Q_3 and each is equal to $\frac{1}{4}$ th of the area of the entire curve i.e.,

$$\begin{aligned}\text{Area} &= \int_{Q_1}^{M_i} f(x) dx = \frac{1}{4} \text{th of the entire area} \\ &= \int_{M_i}^{Q_3} f(x) dx.\end{aligned}$$

The curve $y = f(x)$ being continuous, we have

$$f(x) > f(M_i) \text{ for } Q_1 \leq x \leq M_i$$

$$\text{and } f(x) < f(M_i) \text{ for } M_i \leq x \leq Q_3.$$

Also $f(x)$ is positive as y is positive.




Fig. 15.5

$$A. \quad f(M_i) \int_{Q_1}^{M_i} dx < \int_{Q_1}^{M_i} f(x) dx < f(M_i) \int_{M_i}^{Q_3} dx,$$

$$i.e. \quad f(M_i) (M_i - Q_1) < f(M_i) (Q_3 - M_i)$$

$$\text{or} \quad M_i - Q_1 < Q_3 - M_i$$

which shows that M_i is nearer to Q_1 than to Q_3 .

Problem 69. Determine the quartiles and the median for the following table.

Income		No. of persons	Cumulative frequency
Below	Rs 30	69	69
Between	Rs 30 and below Rs 40	167	236
"	Rs 40 " " Rs 50	207	443
"	Rs 50 " " Rs 60	65	508
"	Rs 60 " " Rs 70	58	566
"	Rs 70 " " Rs 80	27	593
"	Rs 80 and over	10	603
Total		603	

Median = size of the $\frac{603+1}{2}$ th, i.e. 302nd item

= between Rs 40 and below Rs 50.

Applying the formula,

$$\text{Median} = l_1 + \frac{\frac{N}{2} - f_1}{f} \times h, \text{ where } l_1 = 40, f_1 = 236, f = 207, h = 10$$

and $N = 603$, we have

$$\text{Median} = 40 + \frac{\frac{603}{2} - 236}{207} \times 10$$

$$= 40 + \frac{301.52 - 236}{207} \times 10$$

$$= 40 + \frac{65.5 \times 10}{207} = 40 + \frac{655}{207}$$

$$= 40 + 3.2 = 43.2 \text{ approx.}$$

$$\text{Also, } Q_1 = l_1 + \frac{\frac{N}{4} - f_1}{f} \times h, \text{ where } l_1 = 30, f_1 = 69, f = 167, h = 10, N = 603 \text{ and } N/4 \text{ lies in second group}$$

$$= 30 + \frac{\frac{603}{4} - 69}{167} \times 10$$

$$= 30 + \frac{151 - 69}{167} \times 10, \text{ taking } \frac{N}{4} = 151 \text{ approx.}$$

$$= 30 + \frac{820}{167} = 30 + 4.9 \text{ approx.}$$

$$= 34.9$$

$$\text{and } Q_3 = l_1 + \frac{\frac{3N}{4} - f_1}{f} \times h, \text{ where } l_1 = 50, f_1 = 443, f = 65, h = 10, N = 603 \text{ and } \frac{3}{4}N \text{ lies in 4th group}$$

$$= 50 + \frac{\frac{3}{4} \times 603 - 443}{65} \times 10$$

$$= 50 + \frac{452 - 443}{65} \times 10 \text{ approx.}$$

$$= 50 + \frac{100}{65} = 50 + 1.7 \text{ approx.}$$

$$= 51.7.$$

[E] **Mode (or Modal value).** The mode is that variate-value of a distribution for which the frequency is maximum.

It may be found by following three methods:

(i) **By inspection.** When the measures of all the items are given in a frequency table, then the mode is the size of the item which occurs most frequently or in a frequency curve it is situated on the x-axis at the position of greatest ordinate, i.e. the peak of the curve.

(ii) **By grouping.** The items are grouped and regrouped until the point of greatest frequency is unaltered by adjustments of grouping. The mode is then situated at the smaller group common to each of the larger groups. Such a grouping is affected by writing the actual frequencies before their respective size, adding the frequencies in two's, adding again in two's leaving out the first frequency, adding in three's leaving out the first frequencies and then adding in three's leaving out the first two frequencies.

(iii) **By using the formula** (in a continuous series).

$$\text{Mode} = l_1 + \frac{f_2}{f_1 + f_2} \times h.$$

where l_1 = the lower limit of the modal-class (i.e. class having maximum frequency),

f_1 = the frequency of the class preceding the modal class,

f_2 = the frequency of the class following the modal class,

h = the size of the modal class.

Note. Sometimes the formula used is

$$\text{Mode} = l_1 + \frac{f - f_1}{2f - f_1 - f_2} \times h$$

where l_1 = the lower limit of the modal class (i.e. class having maximum frequency),

f_1 = the frequency of the class preceding the modal class,

f_2 = the frequency of the class following the modal class,

h = the size of the modal class.

f = the maximum frequency.

Problem 70. Evaluate the values of mean, mode and median for the following grouped cumulative data:

No. of days absent	No. of students (F)
Less than 5	29
" 10	124
" 15	349
" 20	442
" 25	478
" 30	487
" 35	493
" 40	497
" 50	500

These results can be tabulated as follows:

No. of days absent (class)	Mid-value (x)	No. of students (frequency) (f)	Cumulative frequency	Deviation from assumed mean (225) (ξ)	$f\xi$
0-5	2.5	29	29	-20	-580
5-10	7.5	95	124	-15	-1425
10-15	12.5	225	349	-10	-2250
15-20	17.5	93	442	-5	-465
20-25	22.5	36	478	0	0
25-30	27.5	9	487	+5	45
30-35	32.5	6	493	+10	60
35-40	37.5	4	497	+15	60
40-50	45	3	500	+22.5	67.5
Total		500 = $\Sigma f = N$			-4488.5 = $\Sigma f\xi$

$$\begin{aligned}\therefore \text{Mean} &= A + \frac{1}{N} \Sigma f\xi \\ &= 22.5 + \frac{-4488.5}{500} = 22.5 - 8.977 \\ &= 13.523 \text{ i.e. } 13.5 \text{ approx.}\end{aligned}$$

Median = A.M. of sizes of $\frac{N}{2}$ th and $(\frac{N}{2} + 1)$ th items
= A.M. of sizes of 250th and 251st items
= 12.5 as both items lie in the same group.

This gives a rough value.

We calculate it by using the formula

$$\text{Median} = l_1 + \frac{\frac{N}{2} - f_1}{f - f_1} \times h$$

where $l_1 = 10$, $\frac{N}{2} = 250$, $f_1 = 124$, $f = 225$, $h = 5$.

$$\begin{aligned}\therefore \text{Median} &= 10 + \frac{250 - 124}{225} \times 5 = 10 + \frac{630}{225} \\ &= 10 + 2.8 = 12.8.\end{aligned}$$

This method gives rather accurate value.

By inspection, the value of the mode is 12.5 which is the size of the items of maximum frequency.

By grouping method this can be calculated as follows:

Size of the item (mid-value)	Frequency I	II	III	IV	V	VI
2.5	29					
7.5	95	124				
12.5	225		320	349		354
17.5	93	318			413	
22.5	36		129	138		
27.5	9	45				19
32.5	6		15		51	
37.5	4					
45	3	10	7	13		

Analysis table:

Columns	Size of item having maximum frequency	Here
I	12.5	12.5 occurs 6 times
II	12.5, 17.5	17.5 " 3 "
III	7.5, 12.5	7.5 " 3 "
IV	2.5, 7.5, 12.5	2.5 " 1 "
V	7.5, 12.5, 17.5	Since 12.5 occurs maximum number of times, hence
VI	12.5, 17.5, 22.5	Mode = 12.5.

By using the formula,

$$\text{mode} = l_1 + \frac{f_2}{f_1 + f_2} \times h$$

where $l_1 = 10$, $f_1 = 95$, $f_2 = 93$, $h = 5$

$$\begin{aligned}\text{we have } \text{mode} &= 10 + \frac{93}{95 + 93} \times 5 \\ &= 10 + \frac{465}{188} = 10 + 2.47 \\ &= 12.47.\end{aligned}$$

Using the other formula,

$$\text{mode} = l_1 + \frac{f - f_1}{2f - f_1 - f_2} \times h$$

where $l_1 = 10$, $f_1 = 95$, $f_2 = 93$, $f = 225$, $h = 5$,

$$\begin{aligned}\text{we have } \text{mode} &= 10 + \frac{225 - 95}{2 \times 225 - 95 - 93} \times 5 \\ &= 10 + \frac{130 \times 5}{250 - 188} = 10 + \frac{650}{62} = 20.5 \text{ approx.}\end{aligned}$$

which gives very high result as compared to the previous results.

[F] **Empirical relation between mean, median and mode.** There exists an approximate relation between mean, median and mode for a moderately asymmetrical distribution and this is

$$\text{mode} = \text{mean} - 3(\text{mean} - \text{median}).$$

15.9. MEASURES OF DISPERSION (OR VARIATION)

The measurement of the scatter of the size of the items of a series about the average is said to be a measure of variation, or scatter or spread or dispersion. (Agra, 75)

[A] **The range.** This measure of dispersion known as range is the simplest possible measure to compute and the easiest to understand, but it is the least useful and informative.

The range is the difference between the greatest and the least values in the series.

If x_g and x_l denote the greatest and the least measurements of a series, then

$$\text{Its range} = x_g - x_l$$

and the coefficient of the range or the scatter or simply the ratio of the range is defined to be the ratio $\frac{x_g - x_l}{x_g + x_l}$.

Problem 71. Find the range and coefficient of range for the following set of observations:

10, 15, 20, 37, 58, 60, 90.

$$\text{Range} = x_g - x_l = 90 - 10 = 80.$$

$$\text{Coefficient of range} = \frac{x_g - x_l}{x_g + x_l} = \frac{90 - 10}{90 + 10} = \frac{80}{100} = .8.$$

[B] **The quartile deviation or the semi-interquartile range.** We have introduced the interquartile range as the difference of upper and lower quartiles, i.e., Interquartile range = $Q_3 - Q_1$.

Half of this difference is said to be semi-interquartile range, i.e.

$$\text{Semi-interquartile range} = \frac{Q_3 - Q_1}{2}.$$

The quartile deviation or semi-interquartile range is a better measure of dispersion than the range. Its coefficient is defined by

$$\text{Coefficient of quartile deviation} = \frac{\frac{Q_3 - Q_1}{2}}{\frac{Q_3 + Q_1}{2}} = \frac{Q_3 - Q_1}{Q_3 + Q_1}.$$

Note. The difference between the ninth and first deciles are similarly called as interdecile range which contains 80% of the total frequency while inter-quartile range contains 50% of the total frequency. Thus

the ranges give a fairly good idea of the scatteredness of the distribution and are commonly used in elementary descriptive statistics.

Problem 72. Calculate quartile deviation for the following data:

Farm size (acres)	No. of Farms
0—40	394
41—80	461
81—120	391
121—160	334
161—200	169
201—240	113
241 and over	148

The given data may be arranged in a cumulative frequency table as follows:

Farm size (acres) (x)	No. of Farms (frequency) (f)	Cumulative frequency
0—40	394	394
41—80	461	855
81—120	391	1246
121—160	334	1580
161—200	169	1749
201—240	113	1862
241 and over	148	2010

$$\therefore \frac{1}{4}N = \frac{2010}{4} = 502.5.$$

$\therefore Q_1$ lies in the class (41-80).

$$\text{Thus, } Q_4 = l_1 + \frac{\frac{1}{4}N - f_1}{f} \times h \text{ where } l_1 = 41, f_1 = 394, f = 461, h = 40$$

$$\text{and } \frac{1}{4}N = \frac{2010}{4} = 502.5$$

$$= 41 + \frac{502.5 - 394}{461} \times 40$$

$$= 41 + \frac{108.5 \times 40}{461} = 41 + \frac{4340}{461} = 41 + 9.4 = 50.4$$

$$\text{Again } \frac{3}{4}N = 3 \times 502.5 = 1507.5.$$

$\therefore Q_3$ lies in the class (121-160).

$$\text{Thus } Q_3 = l_1 + \frac{\frac{3}{4}N - f_1}{f} \times h, \text{ where } l_1 = 121, f_1 = 1246, f = 334.$$

$$h = 40, \frac{3}{4}N = 1507.5$$

$$= 121 + \frac{1507.5 - 1246}{334} \times 40$$

$$= 121 + \frac{261.5 \times 40}{334} = 121 + \frac{10460}{334}$$

$$= 121 + 31.02$$

$$= 152.02.$$

$$\begin{aligned}\text{Quartile deviation} &= \frac{1}{2}(Q_3 - Q_1) \\ &= \frac{1}{2}(152.02 - 50.4) \\ &= \frac{1}{2} \times 101.62 = 50.81.\end{aligned}$$

[C] **The mean (or average) deviation.** The mean deviation is defined as the sum of the absolute values of the deviations from an average (median mode or arithmetic mean) divided by the number of items i.e., if x_i ($i=1, 2, 3, \dots, n$) are the variate-values of x with frequencies f_i ($i=1, 2, 3, \dots, n$) such that $N = \sum f_i$ and M be the average (mean, median or mode), then

$$\text{Mean deviation} = \frac{1}{N} \sum f_i |x_i - M| \text{ or simply } \frac{\sum f |x - M|}{N}.$$

If we denote the difference $x - M$ by ξ , then

$$\text{Mean deviation} = \frac{\sum f |\xi|}{N}.$$

Problem 73. Find the mean deviations from the median and the mean of the following data:

Size of items:	4	6	8	10	12	14	16
Frequency:	2	4	5	3	2	1	4

These data in tabulated form are as follows:

Size (x)	Frequency	Cumulative frequency	f. x	Deviations from		f ξ	
				median i.e., 8 ξ ₁	mean i.e., 9.7 ξ ₂	f ξ ₁	f ξ ₂
4	2	2	8	4	5.7	8	11.4
6	4	6	24	2	3.7	8	14.8
8	5	11	40	0	1.7	0	8.5
10	3	14	30	2	0.3	6	0.9
12	2	16	24	4	2.3	8	4.6
14	1	17	14	6	4.3	6	4.3
16	4	21	64	8	6.3	22	25.2
Total	21 = N		204 = Σfx			68 = Σf ξ ₁	69.7 = Σf ξ ₂

$$\text{Median} = \text{size of } \left(\frac{N+1}{2} \right) \text{th item}$$

$$= \text{size of } \frac{21+1}{2} \text{th i.e., 11th item} = 8$$

$$\text{and mean} = \frac{\sum fx}{N} = \frac{204}{21} = 9.7 \text{ approx.}$$

$$\therefore \text{Mean deviation from the median} = \frac{\sum f |\xi_1|}{N} = \frac{68}{21} = 3.238$$

$$\text{and mean deviation from the mean} = \frac{\sum f |\xi_2|}{N} = \frac{69.7}{21} = 3.319$$

Note. Median coefficient of dispersion

$$= \frac{\text{mean deviation from the median}}{\text{median}}$$

$$= \frac{3.238}{8} = 0.40475$$

and mean coefficient of dispersion

$$= \frac{\text{Mean deviation from the mean}}{\text{mean}}$$

$$= \frac{3.319}{9.7} = 0.34 \text{ approx.}$$

[D] **The standard deviation.** We define the standard deviation of a variate x as the square root of the arithmetic mean of the squares of all deviations of x from the arithmetic mean of the observations and denote it by σ . (Agra, 1975)

Thus if M be the arithmetic mean of variate values $x_1, x_2, \dots, x_n$ with frequencies $f_1, f_2, \dots, f_n$ so that $N = \sum f_i$, then

$$\sigma = \sqrt{\left(\frac{1}{N} \sum f_i (x_i - M)^2 \right)} \text{ or simply } \sqrt{\left(\frac{1}{N} \sum f (x - M)^2 \right)}$$

$$\text{giving } \sigma^2 = \frac{1}{N} \sum f (x - M)^2.$$

The quantity σ^2 i.e., the square of the standard deviation is termed as **variance** and denoted by μ_2 , which is the second moment about the mean. (Agra, 1975)

$$\therefore \mu_2 = \sigma^2 = \frac{1}{N} \sum f (x - M)^2$$

The ratio $\frac{\sigma}{M} \times 100$ is known as the coefficient of variation.

When the deviations of x are measured from an assumed mean A , then the square root of the arithmetic mean of the squares of all deviations from x is termed as the **root-mean square deviation** and denoted by s i.e.,

$$s = \sqrt{\left(\frac{1}{N} \sum f_i (x_i - A)^2 \right)} \text{ or simply } \sqrt{\left[\frac{1}{N} \sum f (x - A)^2 \right]}.$$

The square of this quantity s i.e., s^2 is termed as **mean square deviation** and denoted by μ_2' which is the second moment about the assumed mean A , i.e.,

$$\mu_2' = s^2 = \frac{1}{N} \sum f (x - A)^2.$$

Relation between standard and root-mean square deviations.

Let M be the mean and A the assumed mean.

Also let $M-A=d$ and $x-A=\xi$; then

$$x-M=x-A+A-M=(x-A)-(M-A)=\xi-d$$

$$\text{Now, } \sigma^2 = \frac{1}{N} \sum f(x-M)^2 = \frac{1}{N} \sum f(\xi-d)^2 \text{ as } x-M=\xi-d$$

$$= \frac{1}{N} \sum f(\xi^2 - 2\xi d + d^2) = \frac{1}{N} \sum f \xi^2 - 2d \cdot \frac{1}{N} \sum f \xi + d^2 \cdot \frac{1}{N} \sum f$$

$$= \frac{1}{N} \sum f(x-A)^2 - 2d \cdot \frac{1}{N} \sum f(x-A) + d^2 \text{ as } \sum f = N$$

$$= s^2 - 2d \left(\frac{\sum fx}{N} - A \frac{\sum f}{N} \right) + d^2 \text{ as } s^2 = \frac{1}{N} \sum f(x-A)^2$$

$$= s^2 - 2d(M-A) + d^2 = s^2 - 2d \cdot d + d^2 = s^2 - d^2$$

$$\text{i.e., } s^2 = \sigma^2 + d^2 \text{ or } \mu_2' = \mu_2 + d^2$$

which gives relation between s and σ or between μ_2' and μ_2 .

Calculation of standard deviation.

(i) By short-cut method.

$$\text{We have, } \sigma^2 = \frac{1}{N} \sum f(x-M)^2 = \frac{1}{N} \sum f\{x-A-(M-A)\}^2$$

$$= \frac{1}{N} \sum f\{\xi-(M-A)\}^2 \text{ where } \xi=x-A$$

$$= \frac{1}{N} \sum f\{\xi^2 - 2(M-A) \cdot \xi + (M-A)^2\}$$

$$= \frac{1}{N} \sum f \xi^2 - 2(M-A) \cdot \frac{1}{N} \sum f \xi + 2(M-A)^2 \cdot \frac{1}{N} \sum f$$

$$= \frac{1}{N} \sum f \xi^2 - 2d \cdot \frac{1}{N} \sum f \xi + d^2 \text{ as } \sum f = N.$$

$$\text{But } d^2 = s^2 - \sigma^2 = \frac{1}{N} \{\sum f(x-A)^2 - \sum f(x-M)^2\}$$

$$= \frac{1}{N} [\sum f\{(x-A)^2 - (x-M)^2\}] \text{ for the same distribution.}$$

$$= \frac{1}{N} [\sum f\{(x-A+x-M)(x-A-x+M)\}]$$

$$= \frac{1}{N} \sum f\{(\xi+\xi-d) \cdot d\} \text{ as } M-A=d \text{ and } x-M=\xi-d$$

$$= \frac{1}{N} \sum f\{2\xi d - d^2\} = 2d \cdot \frac{1}{N} \sum f \xi - d^2 \cdot \frac{1}{N} \sum f$$

$$\text{or } 2d^2 = 2d \cdot \frac{\sum f \xi}{N} \text{ as } \sum f = N$$

$$\text{giving } d = M-A = \frac{\sum f \xi}{N}.$$

$$\text{Hence } \sigma^2 = \frac{1}{N} \sum f \xi^2 - 2 \left(\frac{\sum f \xi}{N} \right)^2 + \left(\frac{\sum f \xi}{N} \right)^2 = \frac{1}{N} \sum f \xi^2 - \left(\frac{\sum f \xi}{N} \right)^2$$

$$\sigma = \sqrt{\left\{ \frac{1}{N} \sum f \xi^2 - \left(\frac{\sum f \xi}{N} \right)^2 \right\}}.$$

(ii) By step deviation method. Taking $u = \frac{x-A}{h} = \frac{\xi}{h}$, i.e.,

hu , we have

$$\sigma^2 = \frac{1}{N} \sum f \xi^2 - \left(\frac{\sum f \xi}{N} \right)^2 = \frac{1}{N} \sum f h^2 u^2 - \left(\frac{\sum f hu}{N} \right)^2$$

$$= h^2 \left[\frac{1}{N} \sum f u^2 - \left(\frac{\sum f u}{N} \right)^2 \right],$$

$$\text{or } \sigma_x^2 = h^2 \sigma_u^2,$$

$$\text{i.e., } \sigma_x = h \sigma_u.$$

Properties of the standard deviation. The standard deviations of two sets containing N_1 and N_2 members are σ_1 and σ_2 respectively, being measured from their respective means M_1 and M_2 . If the two sets are grouped together as one set of N_1+N_2 members, then the standard deviation σ , of this set, measured from its mean is given by

$$\sigma^2 = \frac{N_1 \sigma_1^2 + N_2 \sigma_2^2}{N_1 + N_2} + \frac{N_1 N_2}{(N_1 + N_2)^2} (M_1 - M_2)^2.$$

If s_1^2 and s_2^2 be the mean square deviations of the two sets, s^2 be the mean square deviation of the combined set and A be the assumed mean; then

$$\begin{aligned} s^2 &= \frac{1}{N_1 + N_2} \sum_{i=1}^{N_1 + N_2} f(x_i - A)^2 \\ &= \frac{1}{N_1 + N_2} \left[\sum_{i=1}^{N_1} f(x_i - A)^2 + \sum_{i=N_1+1}^{N_1+N_2} f(x_i - A)^2 \right] \\ &= \frac{1}{N_1 + N_2} \left[N_1 \cdot \frac{1}{N_1} \sum_{i=1}^{N_1} f(x_i - A)^2 + N_2 \cdot \frac{1}{N_2} \sum_{j=1}^{N_2} (x_j - A)^2 \right] \\ &= \frac{1}{N_1 + N_2} [N_1 \cdot s_1^2 + N_2 \cdot s_2^2] \\ &= \frac{1}{N_1 + N_2} [N_1 (\sigma_1^2 + d_1^2) + N_2 (\sigma_2^2 + d_2^2)] \text{ as } s^2 = \sigma^2 + d^2 \\ &= \frac{N_1 \sigma_1^2 + N_2 \sigma_2^2}{N_1 + N_2} + \frac{N_1 d_1^2 + N_2 d_2^2}{N_1 + N_2} \\ &= \frac{N_1 \sigma_1^2 + N_2 \sigma_2^2}{N_1 + N_2} + \frac{N_1 (M_1 - A)^2 + N_2 (M_2 - A)^2}{N_1 + N_2} \text{ as } d = M - A. \end{aligned}$$

If the assumed mean A be regarded as the mean M of the combined set, then by the properties of arithmetic mean, we have

$$A = M = \frac{N_1 M_1 + N_2 M_2}{N_1 + N_2}$$

and then s becomes identical with σ , i.e. $s = \sigma$.

$$\therefore \sigma^2 = \frac{N_1 \sigma_1^2 + N_2 \sigma_2^2}{N_1 + N_2} + \frac{1}{N_1 + N_2} \left[N_1 \left\{ M_1 - \frac{N_1 M_1 + N_2 M_2}{N_1 + N_2} \right\}^2 + N_2 \left\{ M_2 - \frac{N_1 M_1 + N_2 M_2}{N_1 + N_2} \right\}^2 \right]$$

$$\begin{aligned}
&= \frac{N_1\sigma_1^2 + N_2\sigma_2^2}{N_1 + N_2} + \frac{1}{N_1 + N_2} \left[\frac{N_1 \cdot N_2^2 (M_1 - M_2)^2}{(N_1 + N_2)^2} \right. \\
&\quad \left. + \frac{N_2 \cdot N_1^2 (M_2 - M_1)^2}{(N_1 + N_2)^2} \right] \\
&= \frac{N_1\sigma_1^2 + N_2\sigma_2^2}{N_1 + N_2} + \frac{N_1 N_2}{N_1 + N_2} (M_1 - M_2)^2 \cdot \frac{(N_2 + N_1)}{(N_1 + N_2)^2} \\
&= \frac{N_1\sigma_1^2 + N_2\sigma_2^2}{N_1 + N_2} + \frac{N_1 N_2}{(N_1 + N_2)^2} (M_1 - M_2)^2. \quad \dots (1)
\end{aligned}$$

which is the required result.

This result can be extended to a combined series of observations consisting of r component series containing $N_1, N_2, \dots, N_r$ members with standard deviations $\sigma_1, \sigma_2, \dots, \sigma_r$ and means deviating from the mean of the combined series by $d_1, d_2, \dots, d_r$ whence the standard deviation σ of the whole series with component series having means $M_1, M_2, \dots, M_r$ respectively, is given by

$$\begin{aligned}
\sigma^2 &= \frac{N_1\sigma_1^2 + N_2\sigma_2^2 + \dots + N_r\sigma_r^2}{N_1 + N_2 + \dots + N_r} + \frac{1}{(N_1 + N_2 + \dots + N_r)^2} \\
&\quad \times \{N_1 N_2 (M_1 - M_2)^2 + N_2 N_3 (M_2 - M_3)^2 + \dots \\
&\quad \dots + N_{r-1} N_r (M_{r-1} - M_r)^2\} \dots (2)
\end{aligned}$$

Using Σ notation, the result (2) can be expressed as

$$\sigma^2 = \frac{\sum_{i=1}^r N_i \sigma_i^2}{\sum_{i=1}^r N_i} + \frac{1}{\left(\sum_{i=1}^r N_i \right)^2} \sum_{j, k=1}^r N_j N_k (M_j - M_k)^2. \quad \dots (3)$$

In case $M_1 = M_2 = M_3 = \dots$,

$$\sigma^2 = \frac{N_1\sigma_1^2 + N_2\sigma_2^2 + \dots + N_r\sigma_r^2}{N_1 + N_2 + \dots + N_r} \quad \dots (4)$$

Problem 74. Show that if the variable takes the values $0, 1, 2, \dots, n$ with frequencies given by the terms of the binomial series $q^n, {}^nC_1 q^{n-1} p, {}^nC_2 q^{n-2} p^2, \dots, p^n$ where $p+q=1$, then the mean square deviation is $n^2 p^2 + npq$ and the variance is npq .

By Problem 61, the mean M for such a distribution is given by $M=np$

Now $s^2 = \frac{\Sigma f \xi^2}{\Sigma f}$ where $\xi = x - A$, A being assumed mean.

Assuming that $A=0$, we have

$$s^2 = q^n \cdot 0^2 + {}^nC_1 q^{n-1} p \cdot 1^2 + {}^nC_2 q^{n-2} p^2 \cdot 2^2 + \dots + p^n \cdot n^2 \quad \dots (1)$$

But $(q+p)^n = q^n + {}^nC_1 q^{n-1} p + {}^nC_2 q^{n-2} p^2 + \dots + p^n$

Differentiating w.r.t. p and multiplying both sides by p , we find

$$n(q+p)^{n-1} p = {}^nC_1 q^{n-1} \cdot p + {}^nC_2 q^{n-2} \cdot 2p^2 + \dots + np^n$$

Again differentiating w.r.t. p and multiplying throughout by p , we have

$$\begin{aligned}
&p[{}^nC_1 q^{n-1} p \cdot 1^2 + {}^nC_2 q^{n-2} p^2 \cdot 2^2 + \dots + p^n \cdot n^2] = s^2 \text{ by (1)} \\
&\therefore s^2 = p[n(n-1)p + n] \text{ as } q+p=1 = np[np-p+1] \\
&= np(np+q) \text{ as } q+p=1 \text{ gives } 1-p=q \\
&= n^2 p^2 + npq.
\end{aligned}$$

Also $\sigma^2 = s^2 - d^2 = n^2 p^2 + npq - (np)^2$ as $M=d=np$ when $A=0$ $=npq$.

Problem 75. Show that if deviations are small compared with the mean M so that $\left(\frac{x}{M}\right)^3$ and higher powers of $\frac{x}{M}$ may be neglected,

$$(i) \quad G = M \left(1 - \frac{\sigma^2}{2M^2} \right)$$

$$(ii) \quad M^2 - G^2 = \sigma^2.$$

$$(iii) \quad H = M \left(1 - \frac{\sigma^2}{M^2} \right).$$

$$(iv) \quad MH = G^2$$

$$(v) \quad H + M = 2G.$$

$$(vi) \quad \text{Mean } \sqrt{x} = \sqrt{M} \left\{ 1 - \frac{\sigma^2}{8M^2} \right\}.$$

where σ is the standard deviation, A, G, M respectively arithmetic, geometric and harmonic means of the variate X .

We have $X-M=x$ so that $X=M+x$... (1)

Let N be the total frequency of the distribution. Then

$$(i) \quad \therefore G = (X_1 f_1, X_2 f_2, \dots, X_n f_n)^{1/N} \text{ where } N = \Sigma f.$$

$$\therefore \log G = \frac{1}{N} \Sigma f \log X = \frac{1}{N} \Sigma f \log (M+x) \text{ by (1)}$$

$$= \frac{1}{N} \Sigma f \log M \left(1 + \frac{x}{M} \right) = \frac{1}{N} \left[\Sigma f \log M + \Sigma f \log \left(1 + \frac{x}{M} \right) \right]$$

$$= \frac{1}{N} \Sigma f \log M + \frac{1}{N} \Sigma f \left\{ \frac{x}{M} - \frac{x^2}{2M^2} + \dots \right\} \quad \text{expanding the log-function}$$

$$= \log M \cdot \frac{\Sigma f}{N} = \frac{1}{N} \Sigma f \left(\frac{x}{M} - \frac{x^2}{2M^2} \right) \text{ neglecting higher powers of } x/M$$

$$= \log M + \left[\frac{1}{M} \frac{\Sigma f x}{N} - \frac{1}{2M^2} \frac{\Sigma f x^2}{N} \right] = \log M - \frac{1}{2M^2} \cdot \sigma^2$$

as $\sigma^2 = \frac{\Sigma f x^2}{N}$ and $\Sigma f x = 0$ by first property of A.M.

$$\therefore \log \frac{G}{M} = - \frac{\sigma^2}{2M^2} \text{ i.e. } \frac{G}{M} = e^{-\sigma^2/2M^2} \quad \dots (2)$$

$$= \left(1 - \frac{\sigma^2}{2M^2} + \dots \right)$$

or $G = M \left(1 - \frac{\sigma^2}{2M^2} \right)$ neglecting higher order terms.

$$\begin{aligned} \text{(ii) By (2),} \quad G^2 &= M^2 e^{-\sigma^2/M^2} \\ &= M^2 \left\{ 1 - \frac{\sigma^2}{M^2} \right\} \text{ neglecting higher order terms} \\ &= M^2 - \sigma^2 \text{ i.e. } M^2 - G^2 = \sigma^2 \end{aligned}$$

$$\begin{aligned} \text{(iii) We have } \frac{1}{H} &= \frac{1}{N} \sum f \left(\frac{1}{X} \right) = \frac{1}{N} \sum f \left(\frac{1}{M+x} \right) \\ &= \frac{1}{N} \sum f \left\{ \frac{1}{M \left(1 + \frac{x}{M} \right)} \right\} = \frac{1}{N \cdot M} \sum f \left(1 + \frac{x}{M} \right)^{-1} \\ &= \frac{1}{M} \cdot \frac{1}{M} \sum f \left(1 - \frac{x}{M} + \frac{x^2}{M^2} \right) \text{ neglecting higher order terms} \\ &= \frac{1}{M} \left[\frac{\sum f}{N} - \frac{1}{M} \cdot \frac{\sum fx}{N} + \frac{1}{M^2} \cdot \frac{\sum x^2}{N} \right] \\ &= \frac{1}{M} \left[1 + \frac{\sigma^2}{M^2} \right] \text{ as } \sum fx = 0 \text{ and } \frac{\sum x^2}{N} = \sigma^2, \\ \text{i.e., } H &= M \left[1 + \frac{\sigma^2}{M^2} \right]^{-1} = M \left[1 - \frac{\sigma^2}{M^2} \right] \text{ neglecting higher order terms.} \end{aligned}$$

(iv) From the above relation $H = M \left(1 - \frac{\sigma^2}{M^2} \right)$, we have

$$MH = M^2 \left(1 - \frac{\sigma^2}{M^2} \right) = M^2 - \sigma^2 = G^2 \text{ by (ii)}$$

$$\begin{aligned} \text{(v) We have } M+H &= M + M \left(1 - \frac{\sigma^2}{M^2} \right) = 2M - \frac{\sigma^2}{M} \\ &= 2M \left[1 - \frac{\sigma^2}{2M^2} \right] = 2G \text{ by (i)} \end{aligned}$$

$$\begin{aligned} \text{(vi) Mean } \sqrt{X} &= \frac{1}{N} \sum f \sqrt{X} = \frac{1}{N} \sum f \sqrt{M+x} \\ &= \frac{1}{N} \sum f \left\{ \sqrt{M} \left(1 + \frac{x}{M} \right)^{1/2} \right\} \\ &= \sqrt{M} \cdot \frac{1}{N} \sum f \left(1 + \frac{x}{2M} - \frac{x^2}{8M^2} \right) \text{ (neglecting higher powers of } \frac{x}{M} \text{)} \\ &= \sqrt{M} \left[\frac{\sum f}{N} + \frac{1}{2M} \cdot \frac{\sum fx}{N} - \frac{1}{8M^2} \cdot \frac{\sum fx^2}{N} \right] \\ &= \sqrt{M} \left[1 - \frac{\sigma^2}{8M^2} \right] \text{ as } \sum fx = 0 \text{ and } \frac{\sum fx^2}{N} = \sigma^2. \end{aligned}$$

Problem 76. Show that for any discrete distribution the standard deviation is not less than the mean deviation from the mean.

With usual notation we have to show that

$$\begin{aligned} \sqrt{\left\{ \frac{1}{N} \sum f(x-M)^2 \right\}} &\geq \left\{ \frac{1}{N} \sum f |x-M| \right\}, \\ \text{i.e.,} \quad \frac{1}{N} \sum f(x-M)^2 &\geq \left\{ \frac{1}{N} \sum f |x-M| \right\}^2 \\ \text{or} \quad N \sum f \xi^2 &\geq (\sum f |\xi|)^2 \quad \text{where } \xi = x-M \\ \text{or} \quad (f_1 + f_2 + \dots + f_n) (f_1 \xi_1^2 + f_2 \xi_2^2 + \dots + f_n \xi_n^2) &\geq \{ f_1 |\xi_1| + f_2 |\xi_2| + \dots + f_n |\xi_n| \}^2 \\ &\geq \{ f_1 |\xi_1| + f_2 |\xi_2| + \dots + f_n |\xi_n| \}^2 \\ \text{or} \quad f_1 f_2 (\xi_1^2 + \xi_2^2) + \dots &\geq 2 f_1 f_2 \xi_1 \xi_2 + \dots \\ \text{or} \quad f_1 f_2 (\xi_1 - \xi_2)^2 + \dots &\geq 0, \end{aligned}$$

which is true, since being the sum of perfect squares, the L.H.S. is never negative. Hence the proposition.

Problem 77. From a sample of n observations, the arithmetic mean and variance are calculated. It is then found that one of the values, x_1 , is in error and should be replaced by x_1' , show that the adjustment to the variance to correct this error is

$$\frac{1}{n} (x_1' - x_1) \left(x_1' + x_1 - \frac{x_1' - x_1 + 2T}{n} \right),$$

where T is the total of the original results.

If V be the variance, then V being the square of the standard deviation, we have

$$\begin{aligned} V &= \frac{x_1^2 + x_2^2 + \dots + x_n^2}{n} - \left(\frac{x_1 + x_2 + \dots + x_n}{n} \right)^2 \\ &= \frac{x_1^2 + x_2^2 + \dots + x_n^2}{n} - \left(\frac{T}{n} \right)^2 = \frac{1}{n} \left[\sum_{i=1}^n x_i^2 - \frac{T^2}{n} \right]. \end{aligned}$$

If V' be the corrected variance, then we have

$$\begin{aligned} V' &= \frac{x_1'^2 + x_2^2 + \dots + x_n^2}{n} - \left(\frac{x_1' + x_2 + \dots + x_n}{n} \right)^2 \\ &= \frac{x_1'^2 + (\sum x_i^2 - x_1^2)}{n} - \left(\frac{x_1' + T - x_1}{n} \right)^2 \\ &= \frac{1}{n} \left[(\sum x_i^2 + x_1'^2 - x_1^2) - \frac{(T - x_1 + x_1')^2}{n} \right] \end{aligned}$$

The required adjustment to correct the error is $= V' - V$

$$\begin{aligned} &= \frac{1}{n} \left[\{ (\sum x_i^2 + x_1'^2 - x_1^2) - \sum x_i^2 \} - \frac{1}{n} \{ (T - x_1 + x_1')^2 - T^2 \} \right] \\ &= \frac{1}{n} \left[(x_1'^2 - x_1^2) - \frac{1}{n} \{ (-x_1 + x_1')^2 + 2T(x_1' - x_1) \} \right] \\ &= \frac{1}{n} \left[(x_1'^2 - x_1^2) - \frac{(x_1' - x_1)}{n} \{ x_1' - x_1 + 2T \} \right] \\ &= \frac{1}{n} \left[(x_1' - x_1) \left(x_1' + x_1 - \frac{x_1' - x_1 + 2T}{n} \right) \right]. \end{aligned}$$

Problem 78. A distribution consists of three components with frequencies 200, 250, 300, having means of 25, 10 and 15, standard deviations, of 3, 4 and 5 respectively. Show that the mean of the combined distribution is 16 and its standard deviation is 7.2 approximately. Find also C.V.

The mean M of the combined distribution, is given by

$$M = \frac{1}{N} \sum_{i=1}^3 N_i \bar{x}_i = \frac{N_1 \bar{x}_1 + N_2 \bar{x}_2 + N_3 \bar{x}_3}{N_1 + N_2 + N_3}$$

where $N_1=200$, $N_2=250$, $N_3=300$, $\bar{x}_1=25$, $\bar{x}_2=10$, $\bar{x}_3=15$.

$$\therefore M = \frac{200 \times 25 + 200 \times 10 + 300 \times 15}{200 + 250 + 300} = \frac{5000 + 2500 + 4500}{750} = \frac{12000}{750} = 16.$$

Now, the standard deviation σ of the combined distribution is given by

$$\sigma^2 = \frac{N_1 \sigma_1^2 + N_2 \sigma_2^2 + N_3 \sigma_3^2}{N_1 + N_2 + N_3} + \frac{1}{(N_1 + N_2 + N_3)^2} \{N_1 N_2 (M_1 - M_2)^2 + N_2 N_3 (M_2 - M_3)^2 + N_3 N_1 (M_3 - M_1)^2\}$$

where $N_1=200$, $N_2=250$, $N_3=300$,

$$\sigma_1=3, \sigma_2=4, \sigma_3=5$$

$$M_1=25, M_2=10, M_3=15,$$

$$\therefore \sigma^2 = \frac{200 \times 9 + 250 \times 16 + 300 \times 25}{750} + \frac{1}{(750)^2} \{50000 \times 225 + 75000 \times 25 + 6000 \times 100\}$$

$$= \frac{1800 + 4000 + 7500}{750} + \frac{1}{750 \times 750} [11250000 + 1875000 + 6000000]$$

$$= \frac{13300}{750} + \frac{1}{750 \times 750} \times 19125000$$

$$= \frac{1330}{75} + \frac{191250}{75 \times 75} = \frac{266}{15} + \frac{510}{15} = \frac{776}{15} = 51.73$$

giving $\sigma = 7.2$ approx.

The coefficient of variation (C.V.) = $\frac{\sigma}{M} \times 100$

$$= \frac{7.2}{16} \times 100 = 45$$

Problem 79. The first of two samples has 100 items with mean 15 and standard deviation 3. If the whole group has 250 items with mean 15.6 and standard deviation $\sqrt{13.44}$, find the standard deviation of the second group.

We have $N_1=100$, $N_2=250-100=150$, $M_1=15$, $\sigma_1=3$,
 $M=15.6$, $\sigma=\sqrt{13.44}$

$$\sigma^2 = \frac{N_1 \sigma_1^2 + N_2 \sigma_2^2}{N_1 + N_2} + \frac{N_1 N_2}{(N_1 + N_2)^2} (M_1 - M_2)^2$$

$$13.44 = \frac{100 \times 9 + 150 \times \sigma_2^2}{250} + \frac{100 \times 150}{(250)^2} (15 - M_2)^2 \quad \dots (1)$$

$$\text{Also } M = \frac{N_1 M_1 + N_2 M_2}{N_1 + N_2}$$

$$15.6 = \frac{100 \times 15 + 150 \times M_2}{250}$$

$$150 M_2 = 250 \times 15.6 - 1500 = 2400 \text{ giving } M_2 = 16$$

Substituting this value of M_2 in (1), we get

$$13.44 = \frac{900 + 150 \sigma_2^2}{250} + \frac{6}{25} \times (15 - 16)^2$$

$$= \frac{900 + 150 \sigma_2^2}{250} + (.24)$$

$$900 + 150 \sigma_2^2 = 250 [13.44 - .24] = 250 \times 13.20 = 3300$$

$$150 \sigma_2^2 = 2400 \text{ i.e. } \sigma_2^2 = 16 \text{ or } \sigma_2 = 4.$$

Problem 80. Calculate (a) the quartile, (b) the mean, and (c) the standard deviation wages from the following data:

Weekly wage in dollars	-35	-36	-37	-38	-39	-40	-41	-42
No. of wage-earners	14	20	42	54	45	18	7	

Weekly wage in dollars class	Mid-value x	Frequency f	Cumulative frequency	Deviation from assumed mean = 38.5 (ξ)	ξ^2	$f\xi$	$f\xi^2$
35-36	35.5	14	14	-3	9	-42	126
36-37	36.5	20	34	-2	4	-40	80
37-38	37.6	42	76	-1	1	-42	42
38-39	38.5	54	130	0	0	0	0
39-40	39.5	45	175	1	1	45	45
40-41	40.5	18	193	2	4	18	72
41-42	41.5	7	200	3	9	7	63
Total		$200 = \sum f = N$				$-54 = \sum f\xi$	$428 = \sum f\xi^2$

$\therefore \frac{1}{4}N = \frac{200}{4} = 50 \therefore Q_1$ lies in group 37-38

and $\therefore \frac{3}{4}N = 150, \therefore Q_3$ lies in group 39-40

$Q_1 = \text{Median} = \text{Mean of the sizes of } \{\frac{200}{4} \text{th and } (\frac{200}{4} + 1) \text{th}\} \text{ items}$
 $= 38.5$

$$Q_1 = l_1 + \frac{\frac{1}{4}N - f_1}{f} \times h$$

where $l_1=37$, $\frac{1}{4}N=50$, $f_1=34$, $f=42$, $h=1$

$$\begin{aligned}
 &= 37 + \frac{50-34}{42} \times 41 \\
 &= 37 + \frac{5}{3} = 37 + .38 \\
 &= 37.38 \\
 Q_3 &= l_1 + \frac{\frac{3}{4}N - f_1}{f} \times h \text{ where } l_1 = 39, \frac{3N}{4} = 150, f_1 = 130, f = 45, h = 1 \\
 &= 39 + \frac{150-130}{45} \times 1 \\
 &= 39 + \frac{4}{9} = 39 + .44 = 39.44. \\
 \text{Mean} &= A + \frac{1}{N} \sum f\xi \text{ when; } A = 38.5, \\
 &= 38.5 + \frac{-54}{200} \\
 &= 38.5 - .27 = 38.23 \\
 \sigma &= \left[\frac{1}{N} \sum f\xi^2 - \left(\frac{\sum f\xi}{N} \right)^2 \right] \\
 &= \sqrt{\left\{ \frac{4 \cdot 2 \cdot 3}{2 \cdot 0 \cdot 0} - \left(\frac{5 \cdot 4}{2 \cdot 0 \cdot 0} \right)^2 \right\}} = \sqrt{(2.14 - .0729)} = \sqrt{(2.0671)} \\
 &= 1.4 \text{ approx.}
 \end{aligned}$$

15.10. MOMENTS

For a distribution having variate-values $x_1, x_2, \dots, x_n$ with frequencies $f_1, f_2, \dots, f_n$ respectively such that $N = \sum_{i=1}^n f_i$ (or simply Σf), the r th moment about any point (or assumed mean) A is defined as

$$\mu_r' = \frac{1}{N} \sum_{i=1}^n f_i(x_i - A)^r \text{ or simply } \frac{1}{N} \sum f(x - A)^r, \quad \dots (1)$$

where μ_r' represents the moment about any point A (other than the mean).

As such r th moment can be regarded as the arithmetic mean of the variate $(x-A)^r$.

If we now consider the r th moment about the mean M of the distribution, then it is denoted by μ_r and given by

$$\mu_r = \frac{1}{N} \sum_{i=1}^n f_i(x_i - M)^r, \text{ i.e. } \frac{1}{N} \sum f(x - M)^r. \quad \dots (2)$$

The definitions (1) and (2) give that

$$\mu_0' = \mu_0 = \frac{1}{N} \sum f = 1. \quad \dots (3)$$

$$\mu_1' = \frac{1}{N} \sum f(x - A)$$

$$\left. \begin{aligned} &= \frac{1}{N} \sum f x - A \frac{\sum f}{N}, \text{ where } \sum f = N \quad \text{and} \quad \frac{\sum f x}{N} = M \\ &= M - A = d \text{ (say)} \\ \mu_1' = M = \bar{x} \end{aligned} \right\} \dots (4)$$

$$\mu_1 = \frac{1}{N} \sum f(x-M) = \frac{1}{N} \sum f - M \cdot \frac{1}{N} \sum f = M - M = 0 \quad \dots (5)$$

$$\begin{aligned}\mu_2' &= \frac{1}{N} \sum f(x-A)^2 \\ &= \frac{1}{N} \sum f\xi^2, \text{ when } x-A=\xi \\ &= s^2 = \sigma^2 + d^2. \quad \dots (6)\end{aligned}$$

$$\mu_2 = \frac{1}{N} \sum f(x-M)^2 = d^2. \quad \dots (7)$$

Thus, $\mu_0' - \mu_0 = s^2 - \sigma^2 = d^2$ (8)

[A] Moments about the mean in terms of moments about any point and conversely.

Taking $x-A=\xi$ and $M-A=d$, we have

$$x - M = (x - A) - (M - A) = \xi - d$$

$$\begin{aligned} \mu_r &= \frac{1}{N} \Sigma f(x-M)^r = \frac{1}{N} \Sigma f(\xi-d)^r \\ &= \frac{1}{N} [\Sigma f(\xi^r - {}^r C_1 \xi^{r-1} d + {}^r C_2 \xi^{r-2} d^2 - \dots + (-d)^r)] \\ &= \frac{1}{N} \Sigma f \xi^r - d \cdot {}^r C_1 \frac{1}{N} \Sigma f \xi^{r-1} + d^2 \cdot {}^r C_2 \frac{1}{N} \Sigma f \xi^{r-2} - \dots \\ &\quad + (-1)^{r-1} \cdot d^{r-1} \cdot {}^r C_{r-1} \frac{1}{N} \Sigma f \xi + (-1)^r d^r \frac{1}{N} \Sigma f \\ &= \mu_r' - {}^r C_1 d \mu_{r-1}' + {}^r C_2 d^2 \mu_{r-2}' - \dots \\ &\quad + (-1)^{r-1} d^{r-1} \mu_1' + (-1)^r d^r \dots (1) \end{aligned}$$

But we have $d = \mu_1'$ by (4)

$$\therefore \mu_r = \mu_r' - {}^r C_1 \mu_1' \mu_{r-1}' + {}^r C_2 \mu_1'^2 \mu_{r-2}' - \dots + (-1)^{r-1} (r-1) \mu_1'^r \dots (2)$$

Putting $r=2, 3, 4, \dots$ successively, we get

$$\mu_2 = \mu_2' - \mu_1'^2 = \mu_2' - d^2 \quad \dots (3)$$

$$\mu_3 = \mu_3' - 3\mu_1'\mu_2' + 2\mu_1'^2\mu_1' = \mu_3' - 3d\mu_2' + 2d^3 \dots (4)$$

$$\left. \begin{aligned} \mu_4 &= \mu_4' - 4\mu_3'\mu_1' + 6\mu_2'\mu_1'^2 - 3\mu_1'^3 \\ &= \mu_4' - 4d\mu_3' + 6d^2\mu_2' - 3d^3\mu_1' \end{aligned} \right\}, \quad \dots (5)$$

... .. etc.

Conversely, $\mu_r' = \frac{1}{N} \sum f(x-A)^r = \frac{1}{N} \sum f(x-M+M-A)^r$

$$\begin{aligned}
 &= \frac{1}{N} \sum f(X+d)^r, \text{ when } X=x-M \\
 &= \frac{1}{N} [\sum fX^r + {}^rC_1 d \sum fX^{r-1} + {}^rC_2 d^2 \sum fX^{r-2} + \dots + d^r \sum f] \\
 &= \mu_r + {}^rC_1 \mu_{r-1} d + {}^rC_2 \mu_{r-2} d^2 + \dots + d^r. \quad \dots (6)
 \end{aligned}$$

Putting $r=2, 3, 4, \dots$ successively, we get

$$\mu_2' = \mu_2 + d^2 \text{ as } \mu_1 = 0, \quad \dots (7)$$

$$\mu_3' = \mu_3 + 3d\mu_2 + d^3, \quad \dots (8)$$

$$\mu_4' = \mu_4 + 4d\mu_3 + 6d^2\mu_2 + d^4, \quad \dots (9)$$

$$\dots \dots \dots \text{etc.}$$

[B] **Change of origin and scale of moments.** If we introduce a new variate u related to x , such that $u = \frac{x-A}{h}$, where h is the unit of scale of moments, then

$$x-A=hu \text{ and } \bar{x}-A=h\bar{u}, \quad \dots (1)$$

where $\bar{x}$ and $\bar{u}$ denote the means of variates x and u respectively.

$$\begin{aligned}
 \text{Now, } \mu_r &= \frac{1}{N} \sum f(x-\bar{x})^r \\
 &= \frac{1}{N} \sum f\{(A+hu)-(A+h\bar{u})\}^r \text{ by (1)} \\
 &= \frac{h^r}{N} \sum f(u-\bar{u})^r. \quad \dots (2)
 \end{aligned}$$

$$\text{Also } \mu_r' = \frac{1}{N} \sum f(x-A)^r = \frac{1}{N} \sum f(hu)^r = \frac{h^r}{N} \sum fu^r. \quad \dots (3)$$

Here (2) and (3) show that the r th moment of the variate x is h^r times the corresponding moment of the variate u .

$$\text{If } h=1, \text{ then (2) becomes } \mu_r = \frac{1}{N} \sum f(u-\bar{u})^r$$

which follows that "a change of origin does not affect the moments about the mean."

Again if we take $A=\bar{x}$ and $h=\sigma_x$ then the distribution of x has zero mean and unit variance and $\frac{x-\bar{x}}{\sigma_x}$ is said to be the standard form of the variate.

[C] **Sheppard's corrections for moments of grouped data.** The approximation of assuming the frequencies to be concentrated at the mid-values of class-intervals in a grouped frequency distribution is corrected in case of moments by W.F. Sheppard as follows:

$$\mu_2 (\text{corrected}) = \mu_2 - \frac{h^2}{12}$$

$$\mu_3 (\text{corrected}) = \mu_3$$

$$\mu_4 (\text{corrected}) = \mu_4 - \frac{1}{2}h^2\mu_2 + \frac{7}{240}h^4$$

where h is the width of the class-interval.

These results may be used in cases where the frequency distribution is continuous and the distribution tapers off to zero in both directions.

Problem 81. The first three moments of a distribution about the value 2 of the variable are 1, 16 and -40 . Show that the mean is 3, the variance 15 and $\mu_3 = -86$. Also show that the first three moments about $x=0$ are 3, 24 and 76.

Moments about $A=2$, are $\mu_1' = 1, \mu_2' = 16, \mu_3' = -40$

We have $\mu_1' = \frac{1}{N} \sum f(x-A)$, i.e. $1 = \frac{1}{N} \sum fx - 2$ as $\sum f = N$

$$\frac{1}{N} \sum fx = \text{mean} = 3$$

$$\therefore \mu_2 = \sigma^2 = \mu_2' - \mu_1'^2 = 16 - 1 = 15$$

$$\begin{aligned}
 \text{and } \mu_3 &= \mu_3' - 3\mu_1'\mu_2' + 2\mu_1'^3 \\
 &= -40 - 3 \times 1 \times 16 + 2 \cdot 1^3 = -86.
 \end{aligned}$$

Again when $A=0$, let the three moments about $x=0$ be ν_1', ν_2', ν_3' , then

$$\nu_1' = \frac{1}{N} \sum f(x-0) = \frac{1}{N} \sum fx = \text{mean} = 3$$

$$\nu_2' = \mu_2 + d^2.$$

Here $\mu_2 = 15$ and $d = M - A = 3$

$$\therefore \nu_2' = 15 + 9 = 24.$$

$$\begin{aligned}
 \text{Also } \nu_3' &= \mu_3 + 3d\mu_2 + d^3 \\
 &= -86 + 3 \times 3 \times 15 + (3)^3 = 76.
 \end{aligned}$$

[D] **Factorial and absolute moments.** About the origin i.e. $x=0$, of a distribution, the r th factorial moment is defined as

$$\mu'_{(r)} = \frac{1}{N} \sum_i f_i x_i^{(r)} \text{ or simply } \frac{1}{N} \sum f x^{(r)} \quad \dots (1)$$

where $N = \sum f_i = \sum f$ and $x^{(r)} = x(x-1)(x-2)\dots(x-r+1)$

and the r th absolute moment is defined as

$$\nu_r = \frac{1}{N} \sum_i f_i |x_i|^r \text{ or simply } \frac{1}{N} \sum f |x|^r \quad \dots (2)$$

where $N = \sum f_i = \sum f$

[E] **Moments for bivariate distribution.** If there are two measurable characters say x and y corresponding to each member of the population e.g. heights and weights of students in the survey of the population of a class of students, then each member has a number of pair of values like $(x_1, y_1), (x_2, y_2), \dots, (x_i, y_i), \dots$. The total assemblage of

such pairs of values of x and y along with their frequencies constitutes what is called the *Bivariate frequency distribution* of x and y .

If f_i ($i=1, 2, \dots, n$) be the frequency of the pair (x_i, y_i) , $i=1, 2, \dots, n$, then the moment about the origin of the distribution, of the order r, s with regard to x and y respectively (i.e. order r in x and order s in y) is defined as

$$\mu_{rs}' = \frac{1}{N} \sum_{i=1}^n f_i x_i^r y_i^s = \frac{1}{N} \sum f x^r y^s \quad \dots (1)$$

where $N = \sum_{i=1}^n f_i = \sum f$.

Thus, $\mu_{10}' = \frac{1}{N} \sum f_i x_i = \bar{x}$, the mean for x }
 $\mu_{01}' = \frac{1}{N} \sum f_i y_i = \bar{y}$, the mean for y } \quad \dots (2)

$\mu_{20}' = \frac{1}{N} \sum f_i x_i^2 = \mu_2'$ }
 $\mu_{02}' = \frac{1}{N} \sum f_i y_i^2 = \mu_2'$ } \quad \dots (3)

$\mu_{11}' = \frac{1}{N} \sum f_i x_i y_i$ \quad \dots (4)
, etc.

In a similar manner the moment μ_{rs} about the means $\bar{x}, \bar{y}$ may be defined as

$$\mu_{rs} = \frac{1}{N} \sum_{i=1}^n f_i (x_i - \bar{x})^r (y_i - \bar{y})^s \quad \dots (5)$$

So that $\mu_{10} = \mu_{01} = 0$

$$\begin{aligned} \mu_{11} &= \frac{1}{N} \sum f_i (x_i - \bar{x}) (y_i - \bar{y}) \\ &= \frac{1}{N} \sum f_i x_i y_i - \bar{x} \cdot \frac{1}{N} \sum f_i y_i - \bar{y} \cdot \frac{1}{N} \sum f_i x_i + \bar{x} \bar{y} \\ &= \mu_{11}' - \bar{x} \bar{y} - \bar{x} \bar{y} + \bar{x} \bar{y} = \mu_{11}' - \bar{x} \bar{y}. \end{aligned} \quad \dots (6)$$

The quantity μ_{11} is known as the **Covariance** between the variates x and y and denoted by $\text{Cov}(x, y)$, i.e.

$$\text{Cov}(x, y) = \mu_{11} = \mu_{11}' - \bar{x} \bar{y} \quad \dots (7)$$

Also $\mu_{20} = \sigma_x^2 = \frac{1}{N} \sum f_i (x_i - \bar{x})^2 = \mu_{20}' - \bar{x}^2$ }
 $\mu_{02} = \sigma_y^2 = \frac{1}{N} \sum f_i (y_i - \bar{y})^2 = \mu_{02}' - \bar{y}^2$ } \quad \dots (8)

Problem 82. Show that the first four factorial moments are related to ordinary moments by the relations

$$\begin{aligned} \mu_{(1)}' &= \mu_1' = \bar{x} \\ \mu_{(2)}' &= \mu_2' - \bar{x} \\ \mu_{(3)}' &= \mu_3' - 3\mu_2' + 2\bar{x} \\ \mu_{(4)}' &= \mu_4' - 6\mu_3' + 11\mu_2' - 6\bar{x}. \end{aligned}$$

For assumed mean $A=0$, $\mu_1' = \text{mean} = \bar{x}$ by (4) of §15.10

Now $\mu_{(r)}' = \frac{1}{N} \sum f_i x_i^{(r)}$ gives

$$\begin{aligned} \mu_{(1)}' &= \frac{1}{N} \sum f_i x_i^{(1)} = \frac{1}{N} \sum f_i x_i = \mu_1' = \bar{x} \\ \mu_{(2)}' &= \frac{1}{N} \sum f_i x_i^{(2)} = \frac{1}{N} \sum f_i x_i (x_i - 1) \\ &= \frac{1}{N} \sum f_i x_i^2 - \frac{1}{N} \sum f_i x_i = \mu_2' - \mu_1' = \mu_2' - \bar{x} \\ \mu_{(3)}' &= \frac{1}{N} \sum f_i x_i^{(3)} = \frac{1}{N} \sum f_i x_i (x_i - 1) (x_i - 2) \\ &= \frac{1}{N} \sum f_i x_i^3 - 3 \cdot \frac{1}{N} \sum f_i x_i^2 + 2 \cdot \frac{1}{N} \sum f_i x_i \\ &= \mu_3' - 3\mu_2' + 2\bar{x}. \end{aligned}$$

Similarly, $\mu_{(4)}' = \mu_4' - 6\mu_3' + 11\mu_2' - 6\bar{x}$.

Problem 83. Show that, for bivariate distributions, the moments about the mean are invariant for the change of origin but not for the change of scale.

Let the origin for x and y be transferred to A and A' respectively such that the unit or scale for both remains the same say h . Then,

$$x = A + hu \quad \text{so that} \quad \bar{x} = A + h\bar{u}$$

$$y = A' + hv \quad \quad y = A' + h\bar{v}$$

u, v being the new variates.

$$\begin{aligned} \therefore \mu_{11} &= \frac{1}{N} \sum f_i (x_i - \bar{x}) (y_i - \bar{y}) \text{ when } N = \sum f_i \\ &= \frac{h^2}{N} \sum f_i (u_i - \bar{u}) (v_i - \bar{v}) \\ &= h^2 \left[\frac{1}{N} \sum f_i u_i v_i - \frac{1}{N} \bar{u} \sum f_i v_i - \frac{1}{N} \sum f_i \bar{u} + \bar{u} \bar{v} \frac{1}{N} \sum f_i \right] \\ &= h^2 \left[\frac{1}{N} \sum f_i u_i v_i - \bar{u} \bar{v} - \bar{u} \bar{v} + \bar{u} \bar{v} \right] = h^2 \left[\frac{1}{N} \sum f_i u_i v_i - \bar{u} \bar{v} \right] \end{aligned}$$

$$h=1, \text{ yields, } \mu_{11} = \frac{1}{N} \sum_i f_i u_i v_i - \bar{u} \bar{v}$$

$$\text{By [E], } \mu_{11} = \frac{1}{N} \sum_i f_i x_i y_i - \bar{x} \bar{y} = \frac{1}{N} \sum_i f_i u_i v_i - \bar{u} \bar{v}.$$

Conclusively the moment of order 1 in x and order 1 in y about the mean is invariant for the change of origin but not for the change of scale. The similar results may be shown to hold for other moments also.

[F] β and γ coefficients of Karl Pearson. It is observed that in statistical work, there are certain quantities calculated from moments about the mean, which are found very useful. Such quantities are β and γ coefficients introduced by Karl Pearson. He defined β -coefficients as follows:

$$\beta_1 = \frac{\mu_3^2}{\mu_2^3} = \frac{\mu_3^2}{\sigma^6}, \quad \beta_2 = \frac{\mu_4}{\mu_2^2} = \frac{\mu_4}{\sigma^4} \quad \dots (1)$$

The other two coefficients γ_1 and γ_2 are defined as

$$\gamma_1 = +\sqrt{\beta_1} = \frac{\mu_3}{\sigma^3}; \quad \gamma_2 = \beta_2 - 3 = \frac{\mu_4 - 3\mu_2^2}{\mu_2^2} = \frac{\mu_4 - 3\sigma^4}{\sigma^4} \quad \dots (2)$$

All these coefficients are pure numbers, since μ_n has the dimension n . Their importance lies in giving information regarding the shape of the curve obtained from the frequency distribution.

β_1 gives a measure of departure from symmetry i.e. of *Skewness*.

β_2 or its derivative γ_2 gives a measure of flatness or peakedness of a distribution and known as *Kurtosis*.

[G] **Cumulants.** T.N. Thiele introduced a set of quantities similar to moments, which have a comparatively little theoretical and practical importance. Such quantities are known as cumulants. Their definition constitutes a complicated mathematical structure and hence for the elementary purposes, we mention the first four cumulants as follows:

$$\left. \begin{aligned} \kappa_1 &= \mu_1' \\ \kappa_2 &= \mu_2' - \mu_1'^2 \\ \kappa_3 &= \mu_3' - 3\mu_1'\mu_2' + 2\mu_1'^3 \\ \kappa_4 &= \mu_4' - 4\mu_1'\mu_3' - 3\mu_2'^2 + 12\mu_1'^2\mu_2' - 6\mu_1'^4 \end{aligned} \right\} \quad \dots (1)$$

As a particular case, about the mean, they are defined as

$$\kappa_1 = 0, \quad \kappa_2 = \mu_2, \quad \kappa_3 = \mu_3, \quad \kappa_4 = \mu_4 - 3\mu_2^2 \quad \dots (2)$$

15.11. EXPECTATION

[A] **Univariate probability distribution.** If a variate x assumes $x_1, x_2, \dots, x_n$ (denoted by x_i for $i=1, 2, \dots, n$) values, with their respective probabilities $p_1, p_2, \dots, p_n$ (denoted by p_i for $i=1, 2, \dots, n$) corresponding to the n exhaustive and mutually exclusive cases which may result from a trial, then since the values taken by x depend on chance, x is called a *chance* or *random variable* (variate) or *stochastic variate* (a greek term) and the set of values x_i together with their probabilities p_i constitutes what is called the

univariate probability distribution of the variate of that trial. In such cases since the variate takes a finite or enumerably infinite set of values (i.e. discrete set of values), the distribution is sometimes known as *Discrete Probability distribution*.

[B] Mathematical expectation or expected values.

Suppose $\phi(x)$ is a function of variate x such that it takes values $\phi(x_i)$ ($i=1, 2, \dots, n$) i.e. $\phi(x_1), \phi(x_2), \dots, \phi(x_n)$ when x takes values x_i ($i=1, 2, \dots, n$) with probabilities p_i ($i=1, 2, \dots, n$), the expected or probable value of $\phi(x)$ denoted by $E\{\phi(x)\}$ is defined as

$$E\{\phi(x)\} \text{ or simply } E(\phi) = p_1\phi(x_1) + p_2\phi(x_2) + \dots + p_n\phi(x_n) \\ = \sum_{i=1}^n p_i\phi(x_i) \text{ where } \sum_{i=1}^n p_i = 1 \quad \dots (1)$$

Assuming $\phi(x) = x^r$, we have

$$E(x^r) = p_1x_1^r + p_2x_2^r + \dots + p_nx_n^r = \sum_{i=1}^n p_ix_i^r \quad \dots (2)$$

which is said to be the r th moment of the discrete probability distribution about $x=0$ and denoted by μ_r i.e.

$$\mu_r = E(x^r) = \sum_i p_ix_i^r \quad \dots (3)$$

$$\text{This gives } \mu_1' = E(x) = \sum_i p_ix_i = \bar{x} \text{ by (4) of §15.10} \quad \dots (4)$$

which is said to define the *expected value* or the *expectation* of the variate x .

Again the r th moment of the probability distribution about the mean $\bar{x}$ is defined as

$$\mu_r = E(x_i - \bar{x})^r = \sum_i p_i(x_i - \bar{x})^r, \quad \dots (5)$$

so that μ_2 , the variance of the distribution of x denoted by $\text{Var}(x)$ is given by

$$\mu_2 = \text{Var}(x) = E\{x - E(x)\}^2 \text{ as } \bar{x} = E(x) \text{ from (4).} \quad \dots (6)$$

Problem 81. For a stochastic variable x , prove that

$$\text{Variance} = E(x^2) - \{E(x)\}^2$$

and deduce that $E(x^2) \geq \{E(x)\}^2$.

We have, $\text{Var}(x) = E\{x - E(x)\}^2$

$$= \sum_{i=1}^n p_i (x_i - \bar{x})^2 \text{ where } \bar{x} = E(x) \text{ and using (5)} \\ \text{of [B]}$$

$$= \sum_i p_i (x_i^2 - 2\bar{x}x_i + \bar{x}^2)$$

$$= \sum_i p_ix_i^2 - 2\bar{x} \sum_i p_ix_i + \bar{x}^2 \text{ as } \sum p_i = 1$$

$$= E(x^2) - 2\bar{x} E(x) + \bar{x}^2$$

$$= E(x^2) - 2\{E(x)\}^2 + \{E(x)\}^2 \text{ as } \bar{x} = E(x)$$

$$= E(x^2) - \{E(x)\}^2$$

Problem 88. If n fair dice are tossed and x denotes the sum of the numbers on the n dice, then find the expectation of x .

Denoting by x_i , the number of i th dice, we have

$$x = x_1 + x_2 + \dots + x_n,$$

so that $E(x) = E(x_1 + x_2 + \dots + x_n)$

$$= E(x_1) + E(x_2) + \dots + E(x_n) \text{ by (4) of [C].}$$

But for the i th dice, the variate being the number of points on the dice its values are 1, 2, 3, 4, 5, 6 each having the probability $\frac{1}{6}$, therefore

$$E(x_i) = \frac{1}{6} \cdot 1 + \frac{1}{6} \cdot 2 + \frac{1}{6} \cdot 3 + \frac{1}{6} \cdot 4 + \frac{1}{6} \cdot 5 + \frac{1}{6} \cdot 6 = \frac{21}{6} = \frac{7}{2}.$$

$$\therefore E(x_1) = E(x_2) = \dots = E(x_n) = \frac{7}{2}.$$

$$\text{Hence } E(x) = \frac{7}{2} + \frac{7}{2} + \frac{7}{2} + \dots n \text{ terms} = \frac{7n}{2}.$$

Problem 89. If an unbiased coin is tossed n times, find the mathematical expectation of the number of heads in all the n tosses.

Let x denote the total number of heads in all the n tosses and let x_i be the variate-value which takes 1 when at the i th toss head turns up and takes the value zero when it is not so. Then,

$$x = x_1 + x_2 + \dots + x_n,$$

so that

$$E(x) = E(x_1 + x_2 + \dots + x_n)$$

$$= E(x_1) + E(x_2) + \dots + E(x_n)$$

Now x_i takes the values 1, 0 with probabilities $\frac{1}{2}, \frac{1}{2}$ respectively.

$$\therefore E(x_i) = 1 \cdot \frac{1}{2} + 0 \cdot \frac{1}{2} = \frac{1}{2},$$

$$\text{i.e., } E(x_1) = E(x_2) = \dots = E(x_n) = \frac{1}{2}.$$

$$\text{Thus } E(x) = \frac{1}{2} + \frac{1}{2} + \dots n \text{ times} = \frac{n}{2}.$$

Problem 90. Thirteen cards are drawn simultaneously from a deck of 52. If aces count 1, face cards 10 and others according to denomination, find the expectation of the total score on 13 cards.

If x_i is the number corresponding to the i th card, then x_i takes values 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 10, 10, 10 each having the probability $\frac{1}{13}$. Hence

$$E(x_i) = \frac{1}{13} [1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10 + 10 + 10 + 10] = \frac{85}{13}.$$

$\therefore$ required expectation

$$E(x) = E(x_1) + E(x_2) + \dots + E(x_{13})$$

$$= \frac{85}{13} + \frac{85}{13} + \dots 13 \text{ times} = 85.$$

Problem 91. A box contains a white and b black balls; c balls are drawn. Find the expectation of the number of white balls drawn.

Let s be the number of white balls among the c drawn. Then defining a variate x_i , such that

$$x_i = 1 \text{ if } i\text{th ball drawn is white, } i = 1, 2, \dots, c$$

$$= 0 \text{ if } i\text{th ball drawn is black.}$$

We have

$$s = x_1 + x_2 + \dots + x_c$$

so that

$$E(s) = E(x_1 + x_2 + \dots + x_c).$$

$$= E(x_1) + E(x_2) + \dots + E(x_c).$$

$$\text{But } E(x_i) = 1 \cdot \frac{a}{a+b} + 0 \cdot \frac{b}{a+b} = \frac{a}{a+b},$$

since the probability of $x_i = 1$ is $\frac{a}{a+b}$ and that of $x_i = 0$ is $\frac{b}{a+b}$.

$$\text{Hence } E(s) = \frac{a}{a+b} + \frac{a}{a+b} + \dots c \text{ times} = \frac{ac}{a+b}.$$

Problem 92. A box contains 2^n tickets among which nC_i tickets bear the number i ($i = 0, 1, 2, \dots, n$). A group of m tickets is drawn. Let S denote the sum of their numbers. Find $E(S)$.

Let the variates $x_1, x_2, \dots, x_m$ represent the numbers on the first, second, ..., m th tickets.

Now, x_i being a stochastic variate assume values 0, 1, 2, ..., n and have probabilities $\frac{{}^nC_0}{2^n}, \frac{{}^nC_1}{2^n}, \dots, \frac{{}^nC_n}{2^n}$ respectively, so that

$$E(x_i) = \frac{1}{2^n} [1 \cdot {}^nC_1 + 2 \cdot {}^nC_2 + \dots + n \cdot {}^nC_n]$$

$$= \frac{n}{2^n} [1 + {}^{n-1}C_1 + {}^{n-1}C_2 + \dots + {}^{n-1}C_{n-1}]$$

$$= \frac{n}{2^n} (1 + 1)^{n-1} = \frac{n}{2^n} \cdot 2^{n-1} = \frac{n}{2}.$$

$\therefore S = x_1 + x_2 + \dots + x_m$ gives

$$E(S) = E(x_1 + x_2 + \dots + x_m)$$

$$= E(x_1) + E(x_2) + \dots + E(x_m) = \sum_{i=1}^m E(x_i) = m \cdot \frac{n}{2} = \frac{mn}{2}.$$

[D] **Mathematical expectation of a product of independent stochastic variates.** Two stochastic variates are said to be independent if the probability assumed by either of the variates does not depend upon the value taken by the other.

The expectation of the product of two independent stochastic variates is the product of their expectations, i.e. if x, y be two independent stochastic variates, then

$$E(xy) = E(x) E(y).$$

Suppose the variate x assumes the m values $x_1, x_2, \dots, x_m$ and y assumes the n values $y_1, y_2, \dots, y_n$. Let the probabilities of $x = x_i$ and $y = y_j$ be p_i and p_j' respectively. Now the variates x and y being independent, the compound probability of x taking the value x_i and y taking $y_j = p_i p_j'$,

$$\text{Now, } E(xy) = \sum_{i=1}^m \sum_{j=1}^n p_i p_j' x_i y_j \text{ by definition}$$

$$= \sum_{i=1}^m x_i p_i \sum_{j=1}^n p_j' y_j = E(x) \cdot E(y) \quad \dots (1)$$

This result can be extended to any number n of variates $x, y, z, \dots$ as

$$E(xyz\dots) = E(x) E(y) E(z)\dots, \quad \dots (2)$$

where $x, y, z, \dots$ are n independent stochastic variates.

Problem 93. Find the expected value of the product of points on n dice tossed all together.

Denoting by x_i the number of points on i th dice, we have

$$E(x_1 x_2 \dots x_n) = E(x_1) E(x_2) \dots E(x_n).$$

But $E(x_i) = \frac{7}{2}$ by Problem 89.

$$\therefore E(x_1 x_2 \dots x_n) = \frac{7}{2} \cdot \frac{7}{2} \dots n \text{ times} = \left(\frac{7}{2}\right)^n.$$

[E] **Covariance.** The covariance between two variates x and y whose expected values (or means) are $\bar{x}$ and $\bar{y}$, is defined as

$$\begin{aligned} \text{cov}(x, y) &= E[(x - \bar{x})(y - \bar{y})] \\ &= E[(x - E(x))(y - E(y))] \end{aligned} \quad \dots (1)$$

i.e. the covariance of two variates x and y is the expectation of the product of their deviations from their means.

Problem 94. Show that the covariance of two independent variates is always zero.

Let x and y be two independent variates with their expected values (or means) $\bar{x}$ and $\bar{y}$ respectively. Then we have

$$\begin{aligned} \text{cov}(x, y) &= E[(x - \bar{x})(y - \bar{y})] \\ &= E(x - \bar{x}) \cdot E(y - \bar{y}), \text{ the variates and so their means being independent.} \\ &= \{E(x) - E(\bar{x})\} \cdot \{E(y) - E(\bar{y})\} \\ &= (\bar{x} - \bar{x})(\bar{y} - \bar{y}) = 0. \end{aligned}$$

Problem 95. Show that the converse of Problem 94 is not true, i.e. if the covariance of two variates is zero, it is not necessary that they are independent.

Let us introduce two variates u and v with same variance, such that

$$\begin{aligned} x &= u + v, \text{ so that } \bar{x} = \bar{u} + \bar{v}, \\ y &= u - v, \text{ so that } \bar{y} = \bar{u} - \bar{v}. \end{aligned}$$

$$\text{Now } \text{cov}(x, y) = E[(x - \bar{x})(y - \bar{y})]$$

$$= E[(u - \bar{u} + v - \bar{v})(u - \bar{u} - v + \bar{v})]$$

$$\text{or } \text{cov}(x, y) = E[(\bar{u} - u)^2 - (v - \bar{v})^2]$$

$$= E(u - \bar{u})^2 - E(v - \bar{v})^2$$

$$= \text{var}(u) - \text{var}(v)$$

$$= 0, \text{ since } u \text{ and } v \text{ have the same variance.}$$

Here though $\text{cov}(x, y) = 0$, but x and y are not necessarily independent; for, if u and v are taken as the number of points on two dice, then their sum and difference, i.e., $u + v = x$ and $u - v = y$ are either both even or both odd and thus are dependent to each other. Hence the converse of problem 94 is not true.

Problem 96. Show that $\text{cov}(x, y) = E(xy) - E(x) E(y)$.

$$\begin{aligned} \text{cov}(x, y) &= E[(x - \bar{x})(y - \bar{y})] = E(xy - \bar{x}y - x\bar{y} + \bar{x}\bar{y}) \\ &= E(xy) - \bar{x}E(y) - \bar{y}E(x) + \bar{x}\bar{y} \text{ as } E(x\bar{y}) = \bar{x}\bar{y} \\ &= E(xy) - \bar{x}\bar{y} - \bar{y}\bar{x} + \bar{x}\bar{y} = E(xy) - \bar{x}\bar{y} \\ &= E(xy) - E(x) E(y). \end{aligned}$$

Problem 97. Prove the following:

$$(i) \text{cov}(x + a, y + b) = \text{cov}(x, y),$$

$$(ii) \text{cov}(ax, by) = ab \text{cov}(x, y),$$

$$(iii) \text{cov}\left(\frac{x - \bar{x}}{\sigma_x}, \frac{y - \bar{y}}{\sigma_y}\right) = \frac{1}{\sigma_x \sigma_y} \text{cov}(x, y),$$

where a and b are any numbers.

$$\begin{aligned} (i) \text{cov}(x + a, y + b) &= E[(x + a) - E(x + a)] \{ (y + b) - E(y + b) \} \\ &= E[(x - E(x)) \{ y - E(y) \}] \text{ as } E(a) = a \text{ etc.} \\ &= E\{(x - \bar{x})(y - \bar{y})\} \\ &= \text{cov}(x, y). \end{aligned}$$

$$\begin{aligned} (ii) \text{We have } \text{cov}(ax, by) &= E[\{ax - E(ax)\} \{by - E(by)\}] \\ &= E[\{ax - aE(x)\} \{by - bE(y)\}] \\ &= abE\{(x - \bar{x})(y - \bar{y})\} \\ &= ab \text{cov}(x, y). \end{aligned}$$

(iii) We have

$$\begin{aligned} \text{cov}\left(\frac{x - \bar{x}}{\sigma_x}, \frac{y - \bar{y}}{\sigma_y}\right) &= \frac{1}{\sigma_x \sigma_y} \text{cov}(x - \bar{x}, y - \bar{y}) \\ &= \frac{1}{\sigma_x \sigma_y} \cdot E[\{(x - \bar{x}) - E(x - \bar{x})\} \{(y - \bar{y}) - E(y - \bar{y})\}] \\ &= \frac{1}{\sigma_x \sigma_y} E\{(x - \bar{x})(y - \bar{y})\} \\ &= \frac{1}{\sigma_x \sigma_y} \text{cov}(x, y). \end{aligned}$$

Problem 98. Prove that $\text{cov}(x, x) = \text{var}(x)$.

$$\begin{aligned} \text{We have } \text{cov}(x, x) &= E[(x - \bar{x})(x - \bar{x})] = E(x - \bar{x})^2 \\ &= E\{x - E(x)\}^2 = \text{var}(x). \end{aligned}$$

[F] **Variance of n variates.** The variance of two independent variates x and y is equal to the sum of their variances, i.e.,

$$\text{var}(x + y) = \text{var}(x) + \text{var}(y).$$

$$\begin{aligned} \text{We have } \text{var}(x + y) &= E[(x + y) - E(x + y)]^2 \\ &= E[(x + y) - (\bar{x} + \bar{y})]^2 \\ &= E[(x - \bar{x})^2 + (y - \bar{y})^2 + 2(x - \bar{x})(y - \bar{y})] \\ &= E(x - \bar{x})^2 + E(y - \bar{y})^2 + 2E(x - \bar{x})(y - \bar{y}) \\ &= \text{var}(x) + \text{var}(y) + \text{cov}(x, y) \quad \dots (1) \end{aligned}$$

But by Problem 93, the covariance of two independent variates is zero, i.e., $\text{cov}(x, y) = 0$.

$$\therefore \text{var}(x+y) = \text{var}(x) + \text{var}(y). \quad \dots (2)$$

COROLLARY 1. It is easy to show that

$$\text{var}(x-y) = \text{var}(x) + \text{var}(y). \quad \dots (3)$$

COROLLARY 2. If $x_1, x_2, \dots, x_n$ be n independent variates then the result (2) can be extended as

$\text{var}(x_1 + x_2 + \dots + x_n) = \text{var}(x_1) + \text{var}(x_2) + \dots + \text{var}(x_n)$,
i.e., the variance of the sum of any number of independent variates is equal to the sum of their variances.

COROLLARY 3. If $x_1, x_2, \dots, x_n$ be n random variates with finite variances $\sigma_1^2, \sigma_2^2, \dots, \sigma_n^2$, then the variance of a variate u defined as

$$u = a_1x_1 + a_2x_2 + \dots + a_nx_n,$$

$a_1, a_2, \dots, a_n$ being constants, can be found out.

$$\begin{aligned} \text{We have } E(u) &= E[a_1x_1 + a_2x_2 + \dots + a_nx_n] \\ &= E(a_1x_1) + E(a_2x_2) + \dots + E(a_nx_n) \\ &= a_1E(x_1) + a_2E(x_2) + \dots + a_nE(x_n), \end{aligned}$$

so that

$$u - E(u) = a_1[x_1 - E(x_1)] + a_2[x_2 - E(x_2)] + \dots + a_n[x_n - E(x_n)].$$

Squaring both sides,

$$\begin{aligned} [u - E(u)]^2 &= a_1^2 [x_1 - E(x_1)]^2 + a_2^2 [x_2 - E(x_2)]^2 + \dots \\ &\quad + a_n^2 [x_n - E(x_n)]^2 + 2a_1a_2 [x_1 - E(x_1)][x_2 - E(x_2)] + \dots \\ &\quad + 2a_{n-1}a_n [x_{n-1} - E(x_{n-1})][x_n - E(x_n)], \end{aligned}$$

Taking expected values, we get

$$\begin{aligned} E[u - E(u)]^2 &= a_1^2 E\{x_1 - E(x_1)\}^2 + \dots + a_n^2 E\{x_n - E(x_n)\}^2 \\ &\quad + 2a_1a_2 E\{x_1 - E(x_1)\} \{x_2 - E(x_2)\} + \dots \\ &\quad + 2a_{n-1}a_n E\{x_{n-1} - E(x_{n-1})\} \{x_n - E(x_n)\}, \end{aligned}$$

$$\begin{aligned} \text{i.e., var}(u) &= a_1^2 \text{var}(x_1) + a_2^2 \text{var}(x_2) + \dots + a_n^2 \text{var}(x_n) \\ &\quad + 2a_1a_2 \text{cov}(x_1, x_2) + \dots + 2a_{n-1}a_n \text{cov}(x_{n-1}, x_n) \\ &= \sum_{i=1}^n a_i^2 \text{var}(x_i) + 2 \sum_{\substack{i,j=1 \\ i \neq j}}^n a_i a_j \text{cov}(x_i, x_j). \quad \dots (4) \end{aligned}$$

Deductions

(i) If $a_1 = a_2 = 1, a_3 = a_4 = \dots = a_n = 0$, (4) reduces to (1), i.e.,
 $\text{var}(x_1 + x_2) = \text{var}(x_1) + \text{var}(x_2) + 2 \text{cov}(x_1, x_2).$ $\dots (5)$

(ii) If $a_1 = 1, a_2 = -1, a_3 = a_4 = \dots = a_n = 0$, (4) reduces to
 $\text{var}(x_1 - x_2) = \text{var}(x_1) + \text{var}(x_2) - 2 \text{cov}(x_1, x_2).$ $\dots (6)$

(iii) If x_1 and x_2 are independent variates, then $\text{cov}(x_1, x_2) = 0$ so that results (5) and (6) reduce to (2) and (3), i.e.,
 $\text{var}(x_1 \pm x_2) = \text{var}(x_1) \pm \text{var}(x_2).$ $\dots (7)$

(iv) If $a_1 = a_2 = \dots = a_n = \frac{1}{n}$ so that $x = \bar{x}$, and if x_i 's be the independent variates and also if $\text{var}(x_i) = \sigma^2$, then (4) reduces to

$$\text{var}(\bar{x}) = \frac{\sigma^2}{n^2} + \frac{\sigma^2}{n^2} + \dots n \text{ times} = \frac{\sigma^2}{n} \quad \dots (8)$$

Problem 99. An urn contains pN white and qN black balls, the total number of balls being $N, p+q=1$. Balls are drawn one by one (without being returned to the urn) until a certain number n of balls is reached. What is the dispersion of the number of white balls drawn?

Let x_i ($i=1, 2, 3, \dots, n$) be n variates such that

$$\begin{aligned} x_i &= 1 \text{ if the } i\text{th ball drawn is white} \\ &= 0 \text{ if the } i\text{th ball drawn is black.} \end{aligned}$$

Then, $E(x_i) = p \cdot 1 + q \cdot 0 = p$.

If m be the number of white balls drawn, then

$$m = x_1 + x_2 + \dots + x_n$$

$$\begin{aligned} \therefore E(m) &= E(x_1 + x_2 + \dots + x_n) = E(x_1) + E(x_2) + \dots + E(x_n) \\ &= p + p + \dots n \text{ terms} = np; \end{aligned}$$

and the dispersion of the number of white balls drawn is

$$\begin{aligned} \text{var}(m) &= E\{[m - E(m)]^2\} = E\{(m - np)^2\} \\ &= E\{[(x_1 + x_2 + \dots + x_n) - np]^2\} \\ &= E\{(x_1 + x_2 + \dots + x_n)^2 - 2np(x_1 + x_2 + \dots + x_n) + n^2p^2\} \\ &= E\{x_1^2 + x_2^2 + \dots + x_n^2 \\ &\quad + 2(x_1x_2 + x_1x_3 + \dots + x_ix_j + \dots)\} - 2np \cdot np + n^2p^2 \\ &= \sum_{i=1}^n E(x_i^2) + 2 \sum_{\substack{i,j=1 \\ i \neq j}}^n E(x_ix_j) - n^2p^2 \quad \dots (1) \end{aligned}$$

where $E(x_i^2) = p \cdot 1 + q \cdot 0 = p$, as the possible values of x_i^2 are 1 and 0 with probabilities p and q .

$P(x_ix_j) = P(x_i) \cdot P(x_j|x_i)$ as x_ix_j assumes values 1 and 0 according as i th and j th balls are both white or not. The probability to attain the value 1 is given by $P(x_ix_j)$ which is the product of probability of x_i and the conditional probability of x_j when x_i has already happened.

$$\text{Here } P(x_i) = p \text{ and } P(x_j|x_i) = \frac{pN-1}{N-1}$$

$$\text{so that } P(x_ix_j) = p \cdot \frac{pN-1}{N-1}$$

$$\text{and therefore } E(x_ix_j) = p \cdot \frac{pN-1}{N-1} \cdot 1 + 0 = \frac{p(Np-1)}{N-1}.$$

$$\text{Thus } \sum_{i=1}^n E(x_i^2) = p + p + \dots n \text{ times} = np$$

$$\sum_{\substack{i,j=1 \\ i \neq j}}^n E(x_i x_j) = \frac{p(pN-1)}{N-1} + \dots + {}^nC_2 \text{ times}$$

$$= \frac{n(n-1)}{1.2} \frac{p(pN-1)}{N-1}$$

With these values (1), yields

$$\begin{aligned} \text{var}(m) &= np + 2 \cdot \frac{(n-1)}{1.2} \frac{p(pN-1)}{N-1} - n^2 p^2 \\ &= \frac{np}{N-1} [(N-1) + (n-1)(pN-1) - np(N-1)] \\ &= \frac{np}{N-1} [N\{1+p(n-1)-np\} + \{-1-(n-1)+np\}] \\ &= \frac{np}{N-1} [N(1-p) - n(1-p)] \\ &= \frac{np}{N-1} (1-p)(N-n) = \frac{npq(N-n)}{N-1} \text{ as } 1-p=q. \end{aligned}$$

15.12. CONTINUOUS UNIVARIATE PROBABILITY DISTRIBUTION

The distributions considered so far are the discrete distributions in which the variate assumes a finite or enumerably infinite set of values, but there are the distributions in which a variate like height or weight can take a non-enumerably infinite set of values in a given interval $a \leq x \leq b$. Such variates are said to be *continuous variates* and then probability distributions are known as *continuous probability distributions* or simply *continuous distributions*.

[A] **Probability density function.** If $f(x)$ is a continuous function of x defining the probability distribution of the stochastic variate x such that the probability of the value of the variate lies in the infinitesimal interval $(x - \frac{1}{2}dx, x + \frac{1}{2}dx)$ and is expressible in the form $f(x) dx$ or symbolically

$$P(x - \frac{1}{2}dx \leq X \leq x + \frac{1}{2}dx) = f(x) dx$$

then the function $f(x)$ is said to be the *probability density function* or simply *density function* and $f(x) dx$ is known as *probability differential*.

The continuous curve given by $y=f(x)$ is said to be the *probability density curve* or simply *probability curve*. In case this curve is symmetrical, the distribution is said to be symmetrical.

Though the range or interval of the variate may be finite or infinite, but it is convenient to consider it always infinite even if it is finite whence the density function outside the given range may be assumed zero. Suppose that X assumes values in the interval (a, b) and let its density function there be $\phi(x)$; then its distribution is that of a variate with density function defined as below

$$f(x) = 0 \quad \text{for } x < a,$$

$$\begin{aligned} &= \phi(x) \text{ for } a \leq x \leq b \\ &= 0 \quad \text{for } x > b. \end{aligned}$$

The density function $f(x)$ satisfies the following two properties:

- (i) $f(x) \geq 0$ for every x , as negative probability is meaningless
(ii) $\int_{-\infty}^{\infty} f(x) dx = 1$, i.e., the probability of a sure event is unity. ... (1)

The probability that the variate X falls in an interval (a_1, b_1) is given by

$$P(a_1 \leq X \leq b_1) = \int_{a_1}^{b_1} f(x) dx. \quad \dots (2)$$

Note. 1. The geometrical interpretation of property (ii) of density function namely $\int_{-\infty}^{\infty} f(x) dx = 1$ is that the total area under the curve is unity. In case it is not unity, it can be so by multiplying a suitable constant, e.g.

$$\int_0^1 x(1-x) dx = \frac{1}{6}.$$

Multiplying both sides by 6, this gives

$$\int_0^1 6x(1-x) dx = 1.$$

Thus the probability density function may be so defined

$$\begin{aligned} f(x) &= 0 \text{ for } x < 0 \\ &= 6x(1-x) \text{ for } 0 \leq x \leq 1 \\ &= 0 \text{ for } x > 1. \end{aligned}$$

Note 2. The variate is said to have a *rectangular distribution* of probability when $f(x)$ is constant throughout.

[B] **Cumulative distribution function.** Let the function $F(x)$ be the probability that the value of the variate X is $\leq x$, i.e.

$$F(x) = P(X \leq x) = \int_{-\infty}^x f(x) dx,$$

$$\text{such that } F(b) = P(X \leq b) = \int_{-\infty}^b f(x) dx$$

$$\text{and } F(b) - F(a) = P(a \leq X \leq b) = \int_a^b f(x) dx.$$

Then $F(x)$ is said to be the *cumulative distribution function* of x or simply the *distribution function*. It has the following properties:

- (i) $F(x) = f(x) \geq 0$, so that $F(x)$ is non-decreasing function.
(ii) $F(-\infty) = 0$.
(iii) $F(\infty) = 1$.

Note. Density function $f(x)$ is related to distribution $F(x)$ by $f(x) = F'(x)$.

Problem 100. If $f(x)$ has probability density kx^2 , $0 < x < 1$, determine k and find the probability that $\frac{1}{3} < x < \frac{1}{2}$.

Since $f(x)$ has probability density kx^2 , therefore

$$\int_0^1 f(x) dx = 1,$$

$$\text{i.e.,} \quad \int_0^1 kx^2 dx = 1$$

$$\text{or} \quad \left[k \frac{x^3}{3} \right]_0^1 = 1 \text{ giving } k=3, \text{ so that } f(x)=3x^2.$$

$$\text{Again, } P\left(\frac{1}{3} < x < \frac{1}{2}\right) = \int_{1/3}^{1/2} f(x) dx = \int_{1/3}^{1/2} 3x^2 dx = 3 \left[\frac{x^3}{3} \right]_{1/3}^{1/2} \\ = \frac{1}{8} - \frac{1}{27} = \frac{1}{216}.$$

Problem 101. If a function $f(x)$ of x is defined as follows:

$$f(x)=0 \text{ for } x < 2$$

$$= \frac{1}{18} (3+2x) \text{ for } 2 \leq x \leq 4$$

$$= 0 \text{ for } x > 4,$$

then show that it is a density function and find the probability that a variate having this density will be within the interval $2 \leq x \leq 3$.

In order that $f(x)$ is a density function, it must satisfy the two conditions:

$$(i) f(x) \geq 0 \text{ for every } x.$$

$$(ii) \int_{-\infty}^{\infty} f(x) dx = 1.$$

Here (i) follows from the definition of $f(x)$.

Now for (ii), consider

$$\begin{aligned} \int_{-\infty}^{\infty} f(x) dx &= \int_{-\infty}^2 f(x) dx + \int_2^4 f(x) dx + \int_4^{\infty} f(x) dx \\ &= \int_{-\infty}^2 0 \cdot dx + \int_2^4 \frac{3+2x}{18} dx + \int_4^{\infty} 0 \cdot dx \\ &= 0 + \frac{1}{18} \left[3x + x^2 \right]_2^4 + 0 \\ &= \frac{1}{18} [3(4-2) + (16-4)] = \frac{1}{18} (6+12) = 1 \end{aligned}$$

which satisfies the second condition.

Hence the function $f(x)$ is a density function.

$$\begin{aligned} \text{Now, } P(2 \leq x \leq 3) &= \int_2^3 f(x) dx = \int_2^3 \frac{3+2x}{18} dx \\ &= \frac{1}{18} \left[3x + x^2 \right]_2^3 = \frac{1}{18} [3+5] = \frac{8}{18} = \frac{4}{9} \end{aligned}$$

Problem 192. Verify that the following is a distribution function,

$$F(x) = \begin{cases} 0, & x < -a \\ \frac{1}{2} \left(\frac{a}{x} + 1 \right), & -a \leq x \leq a \\ 1, & x > a. \end{cases}$$

The function $F(x)$ to be a density function, must satisfy the following conditions.

$$(i) F(x) \geq 0,$$

$$(ii) F(-\infty) = 0,$$

$$(iii) F(+\infty) = 1.$$

Here the first condition is obviously satisfied, i.e., $F(x) \geq 0$ from the given definition of $F(x)$.

From $F(x)=0$ for $x < -a$, it is clear that $F(-\infty)=0$ and from $F(x)=1$ for $x > a$, it follows that $F(+\infty)=1$.

Problem 103. Supposing the life in hours of a certain kind of radio tube has the density $f(x) = \frac{100}{x^2}$ when $x \geq 100$ and zero, when $x < 100$, what is the probability that none of three such tubes in a given radio set will have to be replaced during the first 150 hours of operation? What is the probability that all three of the original tubes will have been replaced during the first 150 hours.

$$\begin{aligned} \text{We have} \quad f(x) &= \frac{100}{x^2}, \quad x \geq 100 \\ &= 0, \quad x < 100. \end{aligned}$$

The function $f(x)$ will be a probability density function if

$$\int_{-\infty}^{\infty} f(x) dx = 1$$

$$\text{i.e., if} \quad \int_{-\infty}^{100} f(x) dx + \int_{100}^{\infty} f(x) dx = 1$$

$$\text{or if} \quad \int_{-\infty}^{100} 0 \cdot dx + \int_{100}^{\infty} \frac{100}{x^2} dx = 1$$

$$\text{or if} \quad 0 + 100 \left[-\frac{1}{x} \right]_{100}^{\infty} = 1$$

which is so and hence $f(x)$ is a density function.

$$\text{Now } P(100 \leq x \leq 150) = \int_{100}^{150} \frac{100}{x^2} dx$$

$$= 100 \left[-\frac{1}{x} \right]_{100}^{150} = 100 \left[\frac{1}{100} - \frac{1}{150} \right]$$

$$= 100 \cdot \frac{50}{100 \times 150} = \frac{1}{3}.$$

(since the probability of failure during the first 100 hours is zero).

∴ Probability that none of the three bulbs shall have to be replaced $= \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{8}$.

Again, the probability that one tube fails during the first 150 hours $= 1 - \frac{1}{2} = \frac{1}{2}$.

Hence the probability that all three tubes fail during the first 150 hours $= \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{8}$.

Problem 104. A bomb-plane carrying three bombs flies directly above a rail road track. If a bomb falls within 40 feet of the track, the track will be sufficiently damaged to disrupt traffic. With a certain bomb sight, the density of the points of impact of a bomb is

$$\begin{aligned} f(x) &= \frac{100+x}{10,000}, \quad -100 \leq x \leq 0 \\ &= \frac{100-x}{10,000}, \quad 0 < x \leq 100 \\ &= 0, \text{ elsewhere.} \end{aligned}$$

If all three bombs are used what is the probability that the track will be damaged where x represents the vertical deviation from the aiming point which is the track in this case.

Since a bomb falls within 40 feet of the track, it therefore follows that in order to disrupt traffic the track will be damaged when the bomb falls within 40 feet either side of the track. Thus the probability of one bomb damaging the track is given by

$$\begin{aligned} P(-40 < x < 40) &= \int_{-40}^{40} f(x) dx \\ &= \int_{-40}^0 f(x) dx + \int_0^{40} f(x) dx \\ &= \int_{-40}^0 \frac{100+x}{10,000} dx + \int_0^{40} \frac{100-x}{10,000} dx \\ &= \frac{1}{10,000} \left[100x + \frac{x^2}{2} \Big|_{-40}^0 + \left[100x - \frac{x^2}{2} \Big|_0^{40} \right] \right] \\ &= \frac{1}{10,000} [4,000 - 800 + 4,000 - 80] = \frac{6400}{10,000} = \frac{16}{25}. \end{aligned}$$

So that the probability that the track is not damaged by one bomb $= 1 - \frac{16}{25} = \frac{9}{25}$.

∴ The probability that the track is not damaged by any of three bombs $= \left(\frac{9}{25}\right)^3$.

Hence the probability that the track is damaged by at least one bomb $= 1 - \left(\frac{9}{25}\right)^3 = 1 - \frac{729}{15625} = \frac{14896}{15625}$.

[C] Mean, median, mode, moments etc. for a continuous univariate probability distribution. (i) The Mathematical Expectation.

The expected value (or mean value) of any function $\phi(x)$ of a random variate x having probability density function $f(x)$ and cumulative distribution function $F(x)$, is given by

$$E[\phi(x)] = \int_{-\infty}^{\infty} \phi(x) f(x) dx = \int_{-\infty}^{\infty} \phi(x) dF(x) \quad \dots (1)$$

(ii) The Arithmetic Mean

The mean (arithmetic mean) is obtained from (1) by putting $\phi(x) = x$ whence

$$M = \bar{x} = E(x) = \mu_1' = \int_{-\infty}^{\infty} x f(x) dx \quad \dots (2)$$

which represents the x -coordinate of the centre of gravity of the area of the curve bounded by x -axis.

(iii) The Geometric Mean

The geometric mean G is defined as

$$\log G = \int_{-\infty}^{\infty} \log x f(x) dx = E(\log x) \quad \dots (3)$$

(iv) The Harmonic Mean

The harmonic mean H is defined as

$$\frac{1}{H} = \int_{-\infty}^{\infty} \frac{1}{x} f(x) dx = E\left(\frac{1}{x}\right) \quad \dots (4)$$

(v) The Median

The median say 'a' being that value of a variate x which divides the total frequency into two equal parts i.e.,

$$P(x \leq a) = P(x \geq a)$$

is defined as

$$\int_{-\infty}^a f(x) dx = \int_a^{\infty} f(x) dx = \frac{1}{2} \int_{-\infty}^{\infty} f(x) dx = \frac{1}{2} \quad \dots (5)$$

(vi) The Quartiles

The lower quartile Q_1 is defined as

$$\int_{-\infty}^{Q_1} f(x) dx = \frac{1}{4} \quad \dots (6)$$

and the upper quartile Q_3 is defined as

$$\int_{Q_3}^{\infty} f(x) dx = \frac{1}{4} \quad \dots (7)$$

(viii) The Mode

The mode or modal value being that value of a variate x for which the probability density $f(x)$ is maximum, is defined as

$$f'(x) = 0 \text{ and } f''(x) < 0, \quad \dots (8)$$

provided that the values of x given by $f'(x) = 0$ be within the range of the variate x .

(viii) The Mean Deviation

The mean deviation from the mean ($\bar{x}$) is defined as

$$\text{Mean deviation} = \int_{-\infty}^{\infty} |x - \bar{x}| \cdot f(x) dx \quad \dots (9)$$

(ix) The Moments

The r th moment about any arbitrary point 'a' is given by

$$\mu_r' = \int_{-\infty}^{\infty} (x-a)^r f(x) dx \quad \dots (10)$$

and the r th moment about the mean $\bar{x}$ is defined as

$$\mu_r = \int_{-\infty}^{\infty} (x-\bar{x})^r f(x) dx. \quad \dots (11)$$

Problem 105. If $f(x) = e^{-x}$, $0 \leq x < \infty$ show that it is a probability density function and hence evaluate mean μ_2 , μ_3 , μ_4 .

It is clear that $f(x) \geq 0$ for every x

$$\begin{aligned} \text{and } \int_{-\infty}^{\infty} f(x) dx &= \int_{-\infty}^0 f(x) dx + \int_0^{\infty} f(x) dx \\ &= 0 + \int_0^{\infty} -e^{-x} dx = 0 + \left[-e^{-x} \right]_0^{\infty} = 1. \end{aligned}$$

Both the conditions being satisfied, $f(x)$ is a probability density functions.

$$\begin{aligned} \text{Now mean} &= \int_0^{\infty} x e^{-x} dx = \left[-x e^{-x} \right]_0^{\infty} - \int_0^{\infty} e^{-x} dx \quad (\text{integrating by parts}) \\ &= 0 + \left[-e^{-x} \right]_0^{\infty} = 1. \end{aligned} \quad \dots (1)$$

$$\begin{aligned} \mu_2 &= \int_0^{\infty} (x-1)^2 e^{-x} dx \\ &= \left[-(x-1)^2 e^{-x} \right]_0^{\infty} + 2 \int_0^{\infty} (x-1) e^{-x} dx \\ &\quad (\text{integrating by parts}) \\ &= 1 + 2 \left[\left\{ -(x-1) e^{-x} \right\}_0^{\infty} + \int_0^{\infty} e^{-x} dx \right] \\ &= 1 + 2 [0 - 1 + 1] = 1 \text{ from (1)} \end{aligned} \quad \dots (2)$$

$$\begin{aligned} \mu_3 &= \int_0^{\infty} (x-1)^3 e^{-x} dx \\ \mu_3 &= \left[-(x-1)^3 e^{-x} \right]_0^{\infty} + 3 \int_0^{\infty} (x-1)^2 e^{-x} dx \\ &= -1 + 3 = 2 \text{ from (2)} \end{aligned} \quad \dots (3)$$

$$\begin{aligned} \text{and } \mu_4 &= \int_0^{\infty} (x-1)^4 e^{-x} dx \\ &= \left[-(x-1)^4 e^{-x} \right]_0^{\infty} + 4 \int_0^{\infty} (x-1)^3 e^{-x} dx \\ &= 1 + 4(2) = 9. \end{aligned} \quad \text{from (3)}$$

Problem 106. A frequency function in the range $(-3, 3)$ is defined

$$\begin{aligned} \text{by } y &= \frac{1}{16} (3+x)^2, -3 < x < -1, \\ y &= \frac{1}{16} (6-2x^2), -1 < x < 1, \\ \text{and } y &= \frac{1}{16} (3-x)^2, 1 < x < 3. \end{aligned}$$

Find the mean and the standard deviation of the distribution.

We have first to test whether the given function $y=f(x)$ say) is a density function and it will be so if $\int_{-3}^3 f(x) dx = 1$.

$$\begin{aligned} \text{Now } \int_{-3}^3 f(x) dx &= \int_{-3}^{-1} \frac{1}{16} (3+x)^2 dx + \int_{-1}^1 \frac{1}{16} (6-2x^2) dx \\ &\quad + \int_1^3 \frac{1}{16} (3-x)^2 dx \\ &= \frac{1}{16} \left[\frac{(3+x)^3}{3} \right]_{-3}^{-1} + \frac{1}{16} \left[6x - \frac{2}{3} x^3 \right]_{-1}^1 \\ &\quad + \frac{1}{16} \left[\frac{(3-x)^3}{-3} \right]_1^3 \\ &= \frac{1}{8} + \frac{2}{3} + \frac{1}{8} = 1. \end{aligned}$$

Hence $f(x)$ is a density function.

$$\begin{aligned} \text{Now, mean} &= \mu_1' = \int_{-3}^3 (x-0) \cdot f(x) dx \quad (\text{about the origin}) \\ &= \frac{1}{16} \left[\int_{-3}^{-1} x \cdot (3+x)^2 dx + \int_{-1}^1 x \cdot (6-2x^2) dx \right. \\ &\quad \left. + \int_1^3 x \cdot (3-x)^2 dx \right] \\ &= \frac{1}{16} \left\{ \left[\frac{3}{2} x^2 + 2x^3 + \frac{x^4}{4} \right]_{-3}^{-1} + \left[3x^2 - \frac{x^4}{2} \right]_{-1}^1 \right. \\ &\quad \left. + \left[\frac{3}{2} x^2 - 2x^3 + \frac{x^4}{4} \right]_1^3 \right\} \\ &= \frac{1}{16} \left\{ \left(\frac{9}{2} - 2 + \frac{1}{4} \right) - \left(\frac{81}{2} - 54 + \frac{81}{4} \right) \right\} + \{0 - 0\} \\ &\quad + \left\{ \left(\frac{81}{2} - 54 + \frac{81}{4} \right) - \left(\frac{9}{2} - 2 + \frac{1}{4} \right) \right\} \right\} \\ &= 0 \end{aligned}$$

$$\begin{aligned} \text{and } \mu_2' &= \frac{1}{16} \left[\int_{-3}^{-1} x^2 (3+x)^2 dx + \int_{-1}^1 x^2 (6-2x^2) dx \right. \\ &\quad \left. + \int_1^3 x^2 (3-x)^2 dx \right] \\ &= 1 \quad (\text{integrating and simplifying}). \end{aligned}$$

$$\therefore \text{ standard deviation } \mu_2 = \mu_2' - \mu_1'^2 = 1.$$

15.13. TCHEBYCHEFF'S INEQUALITY

If x be a random variate with expected value $E(x) = \bar{x}$ (i.e. mean) and standard deviation σ , then for any positive number ϵ , we have the inequality

$P(x < \epsilon^2 \bar{x}) \geq 1 - \frac{1}{\epsilon^2}$ (known as Tchebycheff's lemma) and the probability that the chosen value does not differ from the mean by more than $\epsilon\sigma$ does not exceed $\frac{1}{\epsilon^2}$ is represented by Tchebycheff's inequality as

$$P(|x - \bar{x}| \geq \epsilon\sigma) \leq \frac{1}{\epsilon^2}.$$

If x be a non-negative variate, then we have

$$\begin{aligned} \bar{x} &= \int_0^\infty x f(x) dx \\ &= \int_0^{\epsilon^2 \bar{x}} x f(x) dx + \int_{\epsilon^2 \bar{x}}^\infty x f(x) dx \\ &\geq \int_{\epsilon^2 \bar{x}}^\infty x f(x) dx \quad \because \int_0^{\epsilon^2 \bar{x}} x f(x) dx \geq 0 \\ &\geq \epsilon^2 \bar{x} \int_{\epsilon^2 \bar{x}}^\infty f(x) dx = \epsilon^2 \bar{x} P(x \geq \epsilon^2 \bar{x}) \end{aligned}$$

$$\text{i.e., } \bar{x} \geq \epsilon^2 \bar{x} P(x \geq \epsilon^2 \bar{x})$$

$$\text{or } \frac{1}{\epsilon^2} \geq P(x \geq \epsilon^2 \bar{x}) = 1 - P(x < \epsilon^2 \bar{x})$$

$$\therefore P(x < \epsilon^2 \bar{x}) \geq 1 - \frac{1}{\epsilon^2} \quad \dots (1)$$

which is Tchebycheff's lemma.

If we replace x by $(x - \bar{x})^2$, then $\bar{x}$ {i.e. $E(x)$ } will be replaced by variance of x i.e. σ^2 . As such (1) becomes

$$P\{(x - \bar{x})^2 < \epsilon^2 \sigma^2\} \geq 1 - \frac{1}{\epsilon^2}$$

$$\text{or } P(|x - \bar{x}| < \epsilon\sigma) \geq 1 - \frac{1}{\epsilon^2} \quad \dots (2)$$

$$\text{or } 1 - P(|x - \bar{x}| < \epsilon\sigma) \leq \frac{1}{\epsilon^2}$$

$$\text{or } P(|x - \bar{x}| \geq \epsilon\sigma) \leq \frac{1}{\epsilon^2} \quad \dots (3)$$

This proves Tchebycheff's inequality, which restricts the upper limit of the probability.

COROLLARY 1. Law of large numbers.

If we put in (2),

$$x = \frac{x_1 + x_2 + \dots + x_n}{n}, \quad \bar{x} = \frac{\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_n}{n} \text{ and } \epsilon \sqrt{\left(\frac{E_n}{n^2}\right)} = \lambda,$$

where λ is an arbitrary positive number and E_n is the mathematical expectation of the variate,

$$u = (x_1 + x_2 + \dots + x_n - \bar{x}_1 - \bar{x}_2 - \dots - \bar{x}_n)^2.$$

Then, we have

$$\begin{aligned} P\left[\left|\frac{x_1 + x_2 + \dots + x_n}{n} - \frac{\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_n}{n}\right| < \lambda\right] &\geq 1 - \frac{E_n}{n^2 \lambda^2} \\ &\geq 1 - \delta \text{ provided } \frac{E_n}{n^2 \lambda^2} < \delta. \end{aligned}$$

This gives the law of large numbers, stated as below:

Under the assumption that the probability approaches unity or certainty as near as we please, it is expected that the mean (arithmetic) of the values actually taken by n random variates will differ from the mean of their expectations or less than any arbitrary positive small number when the number of variates is sufficiently large and

$$\frac{E_n}{n^2} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

COROLLARY 2. Bernoulli's theorem

For given positive numbers λ and δ , however small, if m be the number of successes in n independent trials, constant probability of success being p , then a number N depending on λ and δ , can be found such that the probability in order that the inequality $\left|\frac{m}{n} - p\right| < \lambda$ holds good, is greater than $1 - \delta$, provided $n > N$.

Let us associate with trials, 1, 2, 3, ..., n random (stochastic) variates $x_1, x_2, \dots, x_n$ such that

$$\begin{aligned} x_i &= 1 \text{ when } i\text{-th trial is a success} \\ &= 0, \text{ when } i\text{-th trial is a failure,} \end{aligned}$$

then

$$m = x_1 + x_2 + \dots + x_n.$$

Now trials being independent, the variates $x_1, x_2, \dots, x_n$ are also independent.

$$\therefore E(x_i) = 1 \cdot p + 0 \cdot q = p \text{ where } p = 1 - q$$

$$E(m) = 1 + 1 + \dots + n \text{ times} = np$$

$$E(x_i^2) = p \cdot 1 + q \cdot 0 = p \text{ as the possible values of } x_i^2 \text{ are } 1 \text{ and } 0 \text{ with probabilities } p \text{ and } q.$$

$$\begin{aligned} \text{Var } (x_i) &= E\{[x_i - E(x_i)]^2\} \\ &= E[(x_i - p)^2] = E[x_i^2 - 2px_i + p^2] \\ &= \Sigma(x_i^2) - 2p\Sigma(x_i) + p^2 \\ &= p - 2p^2 + p^2 = p - p^2 \\ &= p(1 - p) = pq \text{ as } p + q = 1 \end{aligned}$$

and

$$\begin{aligned} \text{var } (m) &= \sum_{i=1}^n \text{var } (x_i) \\ &= \sum_{i=1}^n pq = npq. \end{aligned}$$

$$\text{As such } E\left(\frac{m}{n}\right) = \frac{1}{n} E(m) = p$$

[See Problem 87 (ii)]

$$\text{Var} \left(\frac{m}{n} \right) = \frac{1}{n^2} \text{Var} (m) = \frac{pq}{n} \text{ by Problem 87 (iii)} = \sigma^2 \text{ (say)}$$

Also let $\lambda = \epsilon \sigma$.

Then an application of form (2) of Tchebycheff's inequality for $\frac{m}{n}$ variates, yields

$$P \left(\left| \frac{m}{n} - p \right| < \lambda \right) \geq 1 - \frac{\sigma^2}{\lambda^2} = 1 - \frac{pq}{n\lambda^2} \\ \geq 1 - \delta \text{ when } n > N = \frac{pq}{\delta\lambda^2}.$$

This completes Bernoulli's theorem.

Problem 107. For the number of points x on a dice, prove that Tchebycheff's inequality gives

$$P[|x - E(x)| > 2.5\sigma] < .47$$

while the actual probability is nearly zero.

$$\therefore E(x) = \frac{1}{6} \cdot 1 + \frac{1}{6} \cdot 2 + \frac{1}{6} \cdot 3 + \frac{1}{6} \cdot 4 + \frac{1}{6} \cdot 5 + \frac{1}{6} \cdot 6 \text{ by Problem 88} \\ = \frac{21}{6} = \frac{7}{2} = 3.5$$

$$\therefore \text{Var} (x) = \sigma^2 = E[(x - E(x))^2] = E(x^2) - \{E(x)\}^2 \\ = \frac{1}{6} [1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2] - \left(\frac{7}{2} \right)^2 \\ = \frac{1}{6} [1 + 4 + 9 + 16 + 25 + 36] - \frac{49}{4} \\ = \frac{91}{6} - \frac{49}{4} = \frac{182 - 147}{12} = \frac{35}{12} \\ = 2.9167$$

$$\therefore \sigma = 1.7 \text{ approx.}$$

The maximum deviation of x from $E(x)$ is $6 - 3.5 = 2.5 \approx \frac{5}{2}\sigma$. The probability of greater deviation is zero, whereas Tchebycheff's inequality asserts that this probability is smaller than 0.47.

Problem 108. Let x_i assume values i and $-i$ with equal probabilities, show that the law of large numbers cannot be applied to variables $x_1, x_2, x_3, \dots, x_n$. But if x_i assumes values i^α and $-i^\alpha$, the law of large numbers can be applied to $x_1, x_2, \dots, x_n$ provided $\alpha < \frac{1}{2}$.

We have

$$P(x_i = i) = \frac{1}{2} = P(x_i = -i)$$

$$E(x_i) = \frac{1}{2} \cdot i + \frac{1}{2} \cdot (-i) = 0, i = 1, 2, \dots, n$$

$$E(x_i^2) = \frac{1}{2} i^2 + \frac{1}{2} i^2 = i^2, i = 1, 2, \dots$$

$$E(x) = E(x_1 + x_2 + \dots + x_n)$$

$$= E(x_1) + E(x_2) + \dots + E(x_n)$$

$$= 0 + 0 + \dots + 0 = 0$$

$$E_n = E(x_1 + x_2 + \dots + x_n)^2 = \sum_{i=1}^n E(x_i^2)$$

$$= \sum_{i=1}^n i^2 = 1^2 + 2^2 + \dots + n^2$$

$$= \frac{1}{6} n(n+1)(2n+1)$$

so that

$$\frac{E_n}{n^2} = \frac{1}{6n} (n+1)(2n+1)$$

$\therefore \lim_{n \rightarrow \infty} \frac{E_n}{n^2} = \infty \neq 0$ and hence the law of large numbers is not applicable to this case.

Further when x_i assumes values $i^\alpha, -i^\alpha$, we have

$$E(x_i) = \frac{1}{2} \cdot i^\alpha + \frac{1}{2} \cdot (-i^\alpha) = 0 \text{ as } P(x_i = i^\alpha) = \frac{1}{2} = P(x_i = -i^\alpha)$$

$$E(x_i^2) = \frac{1}{2} \cdot i^{2\alpha} + \frac{1}{2} \cdot i^{2\alpha} = i^{2\alpha}, i = 1, 2, \dots, n$$

$$\text{So that } E_n = \sum_{i=1}^n E(x_i^2) = 1^{2\alpha} + 2^{2\alpha} + \dots + n^{2\alpha}$$

$$\approx \frac{n^{2\alpha+1}}{2\alpha+1} \text{ by Euler's summation formula}$$

$$\therefore \frac{E_n}{n^2} = \frac{n^{2\alpha-1}}{2\alpha+1}$$

$$\text{i.e., } \lim_{n \rightarrow \infty} \frac{E_n}{n^2} = \lim_{n \rightarrow \infty} \frac{n^{2\alpha-1}}{2\alpha+1} = 0 \text{ iff } 2\alpha-1 < 0 \text{ i.e., } \alpha < \frac{1}{2}.$$

Hence in this case the law of large numbers holds only if $\alpha < \frac{1}{2}$.

15.14. CHARACTERISTIC FUNCTION

The characteristic function for a continuous probability distribution having $f(x)$ as its density function is defined as

$$\phi(t) = E(e^{itx}) = \int_{-\infty}^{\infty} e^{itx} f(x) dx = \int_{-\infty}^{\infty} e^{itx} dF(x),$$

where t is a parameter.

$\phi(t)$ has the properties:

- (i) $\phi(t)$ is continuous in t ,
- (ii) $\phi(t)$ is defined in every finite t interval,
- (iii) $\phi(0) = 1$,
- (iv) $\phi(t)$ and $\phi(-t)$ are conjugate quantities,
- (v) $|\phi(t)| \leq \int_{-\infty}^{\infty} |e^{itx}| f(x) dx \leq 1 = \phi(0)$.

15.15. FOURIER'S INVERSION THEOREM

If $F(x)$ and $\phi(t)$ denote the distribution and characteristic functions respectively of a random variate x , then

$$\begin{aligned} F(x) - F(0) &= \lim_{t \rightarrow \infty} \frac{1}{2\pi} \int_{-t}^t \frac{1 - e^{-itx}}{it} \phi(t) dt \\ &= P_r \left[\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{1 - e^{-itx}}{it} \phi(t) dt \right], \end{aligned}$$

where P_r denotes the principal value of the integral.

In case $F(x)$ is everywhere continuous and $dF(x) = f(x) dx$, we can express

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} \phi(t) dt, \quad \dots (1)$$

which is equivalent to Fourier's integral

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dt \int_{-\infty}^{\infty} e^{-it(x-y)} f(y) dy$$

and can be easily deduced from (1) by the substitution

$$\phi(t) = \int_{-\infty}^{\infty} e^{-ity} f(y) dy. \quad \dots (2)$$

Problem 109. Show that the distribution

$$dF = \frac{1}{\pi} \left(\frac{1 - \cos x}{x^2} \right) dx, \quad -\infty \leq x \leq \infty,$$

has the characteristic function

$$\begin{aligned} \phi(t) &= 1 - |t|, \quad |t| \leq 1, \\ &= 0, \quad |t| > 1. \end{aligned}$$

It is evident from calculus of residues that

$$\int_0^{\infty} \frac{1 - \cos x}{x^2} dx = \frac{\pi}{2} \quad \text{or} \quad \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{1 - \cos x}{x^2} dx = 1.$$

$$\therefore f(x) = \frac{1}{\pi} \cdot \frac{1 - \cos x}{x^2}, \quad -\infty \leq x \leq \infty.$$

$\therefore$ the characteristic function is given by

$$\begin{aligned} \phi(t) &= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{1 - \cos x}{x^2} e^{itx} dx \\ &= \frac{2}{\pi} \int_0^{\infty} \frac{1 - \cos x}{x^2} \cos tx dx \\ &= 1 - |t| \quad |t| \leq 1 \\ &= 0 \quad |t| > 1. \end{aligned}$$

Problem 110. Show that the distribution for which the characteristic function $e^{-|t|}$ has the density function

$$f(x) = \frac{1}{\pi} \cdot \frac{dx}{1+x^2}, \quad -\infty \leq x \leq \infty.$$

$$\text{We have } f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-|t|} e^{-itx} dt$$

$$\begin{aligned} &= \frac{1}{2\pi} \left[\int_{-\infty}^0 e^t e^{-itx} dt + \int_0^{\infty} e^{-t} e^{-itx} dt \right] \\ &= \frac{1}{2\pi} \left[\int_0^{\infty} e^{-t} e^{itx} dt + \int_0^{\infty} e^{-t} e^{-itx} dt \right] \text{ by replacing } t \text{ by } -t \text{ in the first integral or putting } t = -z \text{ etc.} \\ &= \frac{1}{2\pi} \int_0^{\infty} e^{-t} (e^{itx} + e^{-itx}) dt \\ &= \frac{1}{\pi} \int_0^{\infty} e^{-t} \cos tx dt \\ &= \frac{1}{\pi} \left[-e^{-t} \cos tx \right]_0^{\infty} - \frac{x}{\pi} \int_0^{\infty} e^{-t} \sin tx dx \\ &\quad \text{(integrating by parts)} \\ &= \frac{1}{\pi} + \frac{x}{\pi} \left[e^{-t} \sin tx \right]_0^{\infty} - \frac{x^2}{\pi} \int_0^{\infty} e^{-t} \cos tx dt \\ &= \frac{1}{\pi} - x^2 f(x) \end{aligned}$$

$$\text{giving} \quad f(x) = \frac{1}{\pi} \cdot \frac{1}{1+x^2}, \quad -\infty \leq x \leq \infty.$$

15.16. THEORETICAL DISTRIBUTIONS

Having started with certain general hypotheses, if it is possible to deduce mathematically the frequency distributions of certain populations or universes, then such distributions are said to be *theoretical distributions*, which constitute three types of distributions.

(i) **Binomial distribution.** It was discovered by James Bernoulli ((1654-1705) in the year 1700 and was first published in 1713, eight years after the death of Bernoulli.

(ii) **Poisson distribution.** It was discovered by the French mathematician and physicist Siméon Denis Poisson (1781-1840) who published it in 1837.

(iii) **Normal distribution.** Though it was first discovered by De Moivre as early as 1733, but associated with the names of the distinguished French mathematician Pierre Simon, Marquis de Laplace (1749-1827) and the German mathematician and physicist Carl Friedrich Gauss (1777-1855) who discussed it independently at the close of 18th and the beginning of 19th centuries.

Here below we shall discuss these three distributions taking one by one.

[A] The Binomial Distribution

If we toss a coin which is a uniform, homogeneous circular disc, then nothing is biased to make it to tend to fall more often on the

one side than on the other. It is therefore expected that in a series of throws the coin will fall heads-up and tails-up an approximately equal number of times and so the chance of throwing heads or tails with a coin is $\frac{1}{2}$. Similarly the chance of throwing an ace with a fair die is $\frac{1}{6}$. Instead of considering the particular instances we generally use to say that the chance of success of an event is p and chance of its failure is q such that $p+q=1$. Assuming the events in a number of trials to be independent, the chances p and q may be supposed to remain constant throughout.

If we take a number of sets of n trials and count the number of successes in each set, then there will be some sets with no success, some with one success, some with two successes, some with three successes and so on. The classification of the sets according to the number of successes which they contain, will give us a frequency distribution.

Suppose there are N sets of n trials in which the chance of the success and failure are respectively p and q . We have to find the frequencies of 0, 1, 2, 3,...successes in cases of one event, two events, three events and so on.

When $n=1$, i.e., in case of a single event, out of N sets of 1 trial each we expect Np successes and Nq failures.

When $n=2$, i.e., there are N sets of two events or N sets of two trials each, the event which has taken place once is repeated again. We have Nq failure of the first event or trial and the events being independent among these Nq there will be $Nq \times q$ failures and $Nq \times p$ successes of the second event on the average. Similarly among the Np successes of the first event there are $Np \times p$ successes and $Np \times q$ failures of the second event on the average. Hence there are in total Nq^2 failures of both events, $2Npq$ cases of the two events with one success and one failure and Np^2 successes of both events. Thus the frequencies of 0, 1, 2 successes are respectively

$$Nq^2, 2Npq, Np^2$$

When $n=3$ i.e., there are N sets of three events, we see that of the Nq^2 cases in which the first two events were failures, we have $Nq^2 \times q$ a third failure and $Nq^2 \times p$ one success, of the $2Npq$ cases, $2Npq^2$ will give two failures and one success and $2Np^2q$ one failure and two successes and of the Np^2 cases, Np^2q will give one failure and two successes and Np^3 a third success. Hence the frequencies of 0, 1, 2, 3 successes are respectively

$$Nq^3, 3Nq^2p, 3Nqp^2, Np^3.$$

From the foregoing discussions we conclude that the frequencies 0, 1, 2,...successes are given

for one event by the binomial expansion of $N(q+p)$

for two " " " " " $N(q+p)^2$

for three " " " " " $N(q+p)^3$.

In general for N sets of n -events (trials) the frequencies of

0, 1, 2, 3,...successes are given by the successive terms in the binomial expansion of $N(q+p)^n$ i.e.,

$$N \left[q^n + nq^{n-1} + \frac{n(n-1)}{1.2} q^{n-2}p^2 + \frac{n(n-1)(n-2)}{1.2.3} q^{n-3}q^3 + \dots \right].$$

This is called the *Binomial Frequency Distribution* or simply *Binomial Distribution* and the quantities n, p (or q) are said to be parameters of Binomial distribution.

Characteristics of Binomial Distribution

- (1) Its general form depends on parameters p, q and n .
- (2) The probability that there are r successes in n independent trials is given by ${}^nC_r p^r q^{n-r}$ and hence in all the N sets, this is given by $N {}^nC_r p^r q^{n-r}$.
- (3) The numerical coefficients of the binomial expansion can be found by Pascal's triangle.

Exponent power of $(p+q)$ i.e. n	Coefficients of terms										Sum of coefficients
1	1	1									2
2	1	2	1								4
3	1	3	3	1							8
4	1	4	6	4	1						16
5	1	5	10	10	5	1					32
...	...	...	...	...	...	...	...	...	...	...	...
10	1	10	45	120	210	252	210	120	45	10	1024
11	1	11	55	...	...	...	...	...	...	...	...
etc.											

(4) It is chiefly applied when the population being sampled is infinite so that ' p ' remains unchanged by sampling.

(5) It can be applied to finite populations also if they are not too small.

(6) It is used under the conditions:

(i) The variable is discrete.

(ii) A dichotomy (i.e. process of classification of collected individuals into two classes according to whether they do or do not possess a particular attribute) exists.

(iii) Statistical independence is assumed.

(iv) The exponent power n is finite and small.

(v) For symmetrical distribution $p=q$ and for asymmetrical $p \neq q$.

Constants of the Binomial Distribution

Let us take an arbitrary origin at 0 (zero) successes so that the successive deviations are 0, 1, 2, 3,... n .

1. The Mean.

We have

the mean $= \mu_1'$ (about the origin)

$$= \sum_{r=0}^n {}^nC_r p^r q^{n-r} \cdot r$$

$$\begin{aligned}
&= (q \cdot 0) + ({}^nC_1 q^{n-1} p \cdot 1) + ({}^nC_2 q^{n-2} p^2 \cdot 2) + \dots + (p^n \cdot n) \\
&= p [nq^{n-1} + n(n-1) q^{n-2} p + \dots + np^{n-1}] \\
&= np [q^{n-1} + (n-1) q^{n-2} p + \dots + p^{n-1}] \\
&= np (q+p)^{n-1} \\
&= np \text{ since } q+p=1.
\end{aligned}$$

2. The Variance and Standard Deviation

We have, μ_2' (about the origin)

$$\begin{aligned}
&= \sum_0^n {}^nC_r p^r q^{n-r} \cdot r^2 \\
&= (q^n \cdot 0) + ({}^nC_1 q^{n-1} p \cdot 1^2) + ({}^nC_2 q^{n-2} p^2 \cdot 2^2) + \dots + (p^n \cdot n^2)
\end{aligned}$$

$$\begin{aligned}
\text{or } \mu_2' &= np \left[q^{n-1} + 2(n-1) q^{n-2} p + \frac{3(n-1)(n-2)}{2} q^{n-3} p^2 + \dots + np^{n-1} \right] \\
&= np [(n-1)p + 1]
\end{aligned}$$

since the bracketed expression is the first moment of $(q+p)^{n-1}$ about origin -1 and hence is equal to $(n-1)p + 1$.

As an alternative,

$$\begin{aligned}
\mu_2' &= \sum_0^n {}^nC_r p^r q^{n-r} \cdot r^2 = \sum_0^n {}^nC_r p^r q^{n-r} [r(r-1) + r] \\
&= n(n-1) p^2 \sum_0^n {}^nC_{r-2} p^{r-2} q^{n-r} + np \sum_0^n {}^nC_{r-1} p^{r-1} q^{n-r} \\
&= n(n-1) p^2 (p+q)^{n-2} + np (p+q)^{n-1} \\
&= n(n-1) p^2 + np \text{ as } p+q=1 \\
&= np [(n-1)p + 1].
\end{aligned}$$

$$\begin{aligned}
\therefore \text{The variance} &= \sigma^2 = \mu_2' - \mu_1'^2 \\
&= np [(n-1)p + 1] - (np)^2 \\
&= np (1-p) \\
&= npq,
\end{aligned}$$

so that standard deviation $= \sigma = \sqrt{\mu_2} = \sqrt{npq}$.

3. Third Moments about the Origin and about the Mean

$$\begin{aligned}
\mu_3' &= \sum_0^n {}^nC_r p^r q^{n-r} \cdot r^3 \\
&= \sum_0^n {}^nC_r p^r q^{n-r} \{r(r-1)(r-2) + 3r(r-1) + r\} \\
&= n(n-1)(n-2) p^3 \sum_0^n {}^nC_{r-3} p^{r-3} q^{n-r} \\
&\quad + 3n(n-1) p^2 \sum_0^n {}^nC_{r-2} p^{r-2} q^{n-r} + np \sum_0^n {}^nC_{r-1} p^{r-1} q^{n-r} \\
&= n(n-1)(n-2) p^3 (p+q)^{n-3} + 3n(n-1) p^2 (p+q)^{n-2} \\
&\quad + np (p+q)^{n-1} \\
&= n(n-1)(n-2) p^3 + 3n(n-1) p^2 + np \text{ as } p+q=1 \\
&= np \{(n-1)(n-2) p^2 + 3(n-1)p + 1\}
\end{aligned}$$

$$\begin{aligned}
\text{and } \mu_3 &= \mu_3' - 3\mu_2'\mu_1' + 2\mu_1'^3 \\
&= npq (q-p).
\end{aligned}$$

4. Fourth Moments About the origin and About the Mean

$$\begin{aligned}
\mu_4' &= \sum_0^n {}^nC_r p^r q^{n-r} \cdot r^4 \\
&= \sum_0^n {}^nC_r p^r q^{n-r} \{r(r-1)(r-2)(r-3) + 6r(r-1)(r-2) \\
&\quad + 7r(r-1) + r\} \\
&= np \{(n-1)(n-2)(n-3) p^3 + 6(n-1)(n-2) p^2 \\
&\quad + 7(n-1)p + 1\} \text{ as above}
\end{aligned}$$

$$\begin{aligned}
\text{and } \mu_4 &= \mu_4' - 4\mu_3'\mu_1' + 6\mu_2'\mu_1'^2 - 3\mu_1'^4 \\
&= 3p^2 q^2 n^2 + npn (1-6pq).
\end{aligned}$$

Problem 111. A perfect cubic die is thrown a large number of times in sets of 8. The occurrence of a 5 or a 6 is called a success. In what proportion of the sets would you expect 3 successes.

Number of faces in a die = 6.

a '5' + a '6' = 2 successes.

$\therefore p = \frac{2}{6} = \frac{1}{3}$, so that $q = 1 - p = \frac{2}{3}$ and $n = 8$.

The binomial distribution is therefore $N(\frac{2}{3} + \frac{1}{3})^8$.

Probability of 3 successes in one set of 8 = ${}^8C_3 p^3 q^5$

$$= \frac{8 \cdot 7 \cdot 6}{1 \cdot 2 \cdot 3} (\frac{1}{3})^3 (\frac{2}{3})^5 = \frac{8 \times 7 \times 32}{81 \times 81}.$$

$$\begin{aligned}
\therefore \text{Probability of 3 successes in 100 sets} &= \frac{8 \times 7 \times 32}{81 \times 81} \times 100 \\
&= \frac{179200}{6561} = 27.31\%.
\end{aligned}$$

Problem 112. An irregular six-faced die is thrown and the expectation that in 10 throws it will give 5 even numbers is twice the expectation that it will give four even numbers. How many times in 10,000 sets of 10 throws would you expect it to give no even members?

If p is the expectation of getting an even number then the probability of 5 even numbers in 10 throws of a die is twice the probability of 4 even numbers in 10 throws of the die, i.e.

$${}^{10}C_5 p^5 q^5 = 2 \times {}^{10}C_4 p^4 q^6,$$

$$\frac{2}{5} p = q = 1 - p \text{ giving } p = \frac{5}{8} \text{ and } q = \frac{3}{8}.$$

$\therefore$ Frequency for no even number in 10,000 sets of 10 throws = $10,003 \cdot q^{10} = 10,000 (\frac{3}{8})^{10} = 1$ nearly.

Problem 113. Show the results of throwing 12 dice, 4096 times, a throw of 4, 5 or 6 being called a success.

Find the expected frequencies and compare the actual mean with those of the expected distribution. Calculate the standard deviation.

Probability of success of falling 4, 5 or 6 = $\frac{3}{6} = \frac{1}{2} = p$ (say).

$$\therefore q = 1 - p = \frac{1}{2}.$$

Success	Frequency	Success	Frequency
0	—	7	847
1	7	8	536
2	60	9	257
3	198	10	71
4	430	11	11
5	731	12	—
6	948		
Total 4,096			

Binomial distribution is $4096 \left(\frac{1}{2} + \frac{1}{2}\right)^{12}$.

Frequency of 0 successes = $4096 \left(\frac{1}{2}\right)^{12} = 1$,

∴ 1 success = $4096 \cdot {}^{12}C_1 q^{11} \cdot p = 12 \times \frac{4096}{2^{12}} = 12$.

∴ 2 successes = $4096 \cdot {}^{12}C_2 q^{10} p^2 = 66$

∴ 3 successes = ${}^{12}C_3 q^9 p^3 = 220$

∴ 4 successes = $4096 \cdot {}^{12}C_4 q^8 p^4 = 495 \dots \text{etc.}$

Now, expected mean = $np = 12 \cdot \frac{1}{2} = 6$,

$$\sigma^2 = npq = 12 \cdot \frac{1}{2} \cdot \frac{1}{2} = 3.$$

∴ standard deviation $\sigma = \sqrt{3} = 1.732$.

Problem 114. If a coin is tossed N times, where N is very large even number, show that the probability of getting exactly $\frac{1}{2}N - p$ heads and $\frac{1}{2}N + p$ tails is approximately

$$\left(\frac{2}{\pi N}\right)^{1/2} e^{-2p^2/N}$$

Since the coin is unbiased $p = q = \frac{1}{2}$.

∴ Required probability = ${}^N C_{\frac{1}{2}N - p} p^{\frac{n}{2} - p} q^{\frac{n}{2} + p}$

$$= \frac{N!}{\left(\frac{1}{2}N - p\right)! \left(\frac{1}{2}N + p\right)!} \frac{1}{2^N} \text{ as } p = q = \frac{1}{2}$$

Applying Stirling's theorem i.e.,

$$n! = \sqrt{(2\pi n)} \cdot n^n e^{-n} \text{ approx.}$$

$$= e^{-n} \cdot n^{n + \frac{1}{2}} \sqrt{(2\pi)}, \text{ } n \text{ being large,}$$

we have the required probability

$$\begin{aligned} & \frac{\sqrt{(2\pi)} \cdot N^{N + \frac{1}{2}} e^{-N}}{[\sqrt{(2\pi)}]^2 \left(\frac{1}{2}N - p\right)^{\frac{1}{2}N - p + \frac{1}{2}} e^{-(\frac{1}{2}N - p)} \left(\frac{1}{2}N + p\right)^{\frac{1}{2}N + p + \frac{1}{2}} e^{-(\frac{1}{2}N + p)}} \cdot \frac{1}{2^N} \\ &= \frac{1}{\sqrt{(2\pi)}} \cdot \frac{N^{N + \frac{1}{2}}}{\left(\frac{1}{2}N\right)^{N + \frac{1}{2}} \left(1 - \frac{2p}{N}\right)^{\frac{1}{2}N - p + \frac{1}{2}} \left(1 + \frac{2p}{N}\right)^{\frac{1}{2}N + p + \frac{1}{2}}} \cdot \frac{1}{2^N} \\ &= \frac{2}{\sqrt{(2\pi)}} \cdot \frac{1}{N^{\frac{1}{2}}} \left(1 + \frac{2p}{N}\right)^{\frac{1}{2}N - p + \frac{1}{2}} \left(1 - \frac{2p}{N}\right)^{\frac{1}{2}N + p + \frac{1}{2}} \end{aligned}$$

$$= \left(\frac{2}{\pi N}\right)^{\frac{1}{2}} \left(1 - \frac{4p^2}{N^2}\right)^{N/2} \text{ neglecting other terms}$$

$$= \left(\frac{2}{\pi N}\right)^{\frac{1}{2}} e^{-2p^2/N} \text{ approx. when } N \text{ is large, for then}$$

$$\lim_{N \rightarrow \infty} \left(1 - \frac{4p^2}{N^2}\right)^{N/2} = \lim_{N \rightarrow \infty} \left(1 - \frac{2p^2/N}{\frac{1}{2}N}\right)^{\frac{1}{2}N} = e^{-2p^2/N}$$

[B] Limiting form of the binomial distribution when n is large.

We know that for large values of n , each term of the binomial becomes small, but we consider the sum of the terms lying within certain ranges. It is observed that as n increases and becomes infinitely large, the curve representing the binomial distribution becomes smooth and continuous and approaches the *normal curve* which is a symmetrical curve such that the maximum frequencies are clustered around the mean and the deviation below the mean are equal in number and magnitude to the deviations above the mean. Thus normal curve is a standardized and special case of the binomial and the distribution giving this particular curve is said to be the *normal distribution*. We consider it in two cases:

Case I. Normal distribution as a limiting case of binomial distribution where $p = q$.

When $p = q = \frac{1}{2}$, the terms of the series $N\left(\frac{1}{2} + \frac{1}{2}\right)^n$ are

$$N \left(\frac{1}{2}\right)^n \left[1 + n + \frac{n(n-1)}{1 \cdot 2} + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3} + \dots \right]$$

The frequency of m successes is

$$N \left(\frac{1}{2}\right)^n \cdot {}^N C_m \text{ i.e., } N \left(\frac{1}{2}\right)^n \frac{n!}{m! (n-m)!}$$

and the frequency of $(m+1)$ successes is $N \left(\frac{1}{2}\right)^n \cdot {}^N C_{m+1}$

$$= N \left(\frac{1}{2}\right)^n \frac{n!}{(m+1)! (n-m-1)!}$$

$$= \left\{ N \left(\frac{1}{2}\right)^n \frac{n!}{m! (n-m)!} \right\} \frac{n-m}{m+1}$$

$$= \left(\frac{n-m}{m+1}\right) \text{th times the frequency of } m \text{ successes.}$$

So that the frequency of $(m+1)$ successes is greater than that of m successes if

$$\frac{n-m}{m+1} > 1$$

or if

$$n - m > m + 1$$

or if

$$m < \frac{n-1}{2}$$

For the sake of convenience let us suppose that $n = 2k$; then the frequency of k successes say $y_0 = N \left(\frac{1}{2}\right)^{2k} \frac{(2k)!}{k! k!}$ and the frequency of $(k+x)$ successes $y_x = N \left(\frac{1}{2}\right)^{2k} \frac{(2k)!}{(k+x)! (k-x)!}$.

$$\therefore \frac{y_x}{y_0} = \frac{k! k!}{(k+x)! (k-x)!} = \frac{k! k (k-1) (k-2) \dots (k-x+1) (k-x)!}{k! (k+1) (k+2) \dots (k+x) (k-x)!}$$

$$= \frac{k^x \left(1 - \frac{1}{k}\right) \left(1 - \frac{2}{k}\right) \dots \left(1 - \frac{x-1}{k}\right)}{k^x \left(1 + \frac{1}{k}\right) \left(1 + \frac{2}{k}\right) \dots \left(1 + \frac{x-1}{k}\right) \left(1 + \frac{x}{k}\right)}$$

$$\text{or } \log \frac{y_x}{y_0} = \log \left(1 - \frac{1}{k}\right) + \log \left(1 - \frac{2}{k}\right) + \dots + \log \left(1 - \frac{x-1}{k}\right)$$

$$- \log \left(1 + \frac{1}{k}\right) - \log \left(1 + \frac{2}{k}\right) - \dots - \log \left(1 + \frac{x-1}{k}\right) - \log \left(1 + \frac{x}{k}\right)$$

Supposing k to be very large as compared with x so that $\left(\frac{x}{k}\right)$ may be neglected, expanding the logarithmic functions as $\log(1+z) = z - \frac{z^2}{2} + \frac{z^3}{3} \dots$ and $\log(1-z) = -z - \frac{z^2}{2} - \frac{z^3}{3} \dots$ and then neglecting the squares of $\frac{x}{k}$ i.e., $\left(\frac{x}{k}\right)^2$, we find

$$\log \frac{y_x}{y_0} = -\frac{1}{k} - \frac{2}{k} - \frac{3}{k} - \dots - \frac{x-1}{k} - \frac{1}{k} - \frac{2}{k} - \dots - \frac{x-1}{k} - \frac{x}{k}$$

$$= -\frac{2}{k} (1+2+3+\dots+x-1) - \frac{x}{k}$$

$$= -\frac{2}{k} \cdot \frac{x(x-1)}{2} - \frac{x}{k}$$

$$= -\frac{x^2}{y_0}$$

$$\text{or } \frac{y_x}{y_0} = e^{-x^2/k}$$

i.e., $y_x = y_0 e^{-x^2/k}$.

But for a binomial distribution with $p=q=\frac{1}{2}$, $n=2k$, we have

$$\sigma^2 = npq = 2k \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{2}k \text{ i.e., } k = 2\sigma^2$$

$$\text{Hence } y_x = y_0 e^{-(x^2/2\sigma^2)}$$

which gives the normal distribution and the equation $y = y_0 e^{-x^2/2\sigma^2}$ represents normal curve.

Case II. Normal distribution as a limiting case of binomial distribution $p \neq q$ but $p \approx q$.

The frequency of m successes is

$$N \cdot {}^nC_m q^{n-m} p^m = N \cdot \frac{n!}{m! (n-m)!} \cdot q^{n-m} p^m$$

and the frequency of $(m+1)$ successes is $N \cdot {}^nC_{m+1} q^{n-m-1} p^{m+1}$

$$= \frac{n!}{(m+1)! (n-m-1)!} q^{n-m-1} p^{m+1}$$

$$= \left\{ N \cdot \frac{n!}{m! (n-m)!} q^{n-m} p^m \right\} \cdot \frac{n-m}{m+1} \cdot \frac{p}{q}$$

$$= \left(\frac{n-m}{m+1} \cdot \frac{p}{q} \right) \text{th times the frequency of } m \text{ successes.}$$

So that the frequency of $(m+1)$ successes is greater than that of m successes if $\frac{n-m}{m+1} \cdot \frac{p}{q} > 1$ or if $(n-m)p > (m+1)q$ i.e. if $m < n p - q$.

Assuming that np is a whole number, since there is no loss of generality as n ultimately tends to infinity, the frequency of np successes may be taken as maximum frequency.

The frequency of np successes (say)

$$y_0 = N \cdot \frac{n!}{np! (n-np)!} \cdot q^{n-np} p^{np}$$

$$= N \cdot \frac{n!}{np! nq!} q^{nq} p^{np} \text{ as } 1-p=q$$

and the frequency of $(np+x)$ successes (say)

$$y_x = N \cdot \frac{n!}{(np+x)! (np-x)!} q^{nq-x} p^{np+x}$$

$$= N \cdot \frac{n!}{np! nq!} q^{nq} p^{np} \text{ as } 1-p=q$$

and the frequency of $(np+p)$ successes (say)

$$y_x = N \cdot \frac{n!}{(np+x)! (nq-x)!} q^{nq-x} p^{np+x}$$

$$\therefore \frac{y_x}{y_0} = \frac{(np)! (nq)!}{(np+x)! (nq-x)!} q^{-x} p^x.$$

Applying the Stirling's theorem that when n is large

$$n! = \sqrt{(2\pi n)} n^n e^{-n} = e^{-n} n^{n+1/2} \sqrt{(2\pi)} \text{ approx.}$$

we have

$$\frac{y_x}{y_0} = \frac{e^{-np} (np)^{np+1/2} \sqrt{(2\pi)} \times e^{-nq} (nq)^{nq+1/2} \sqrt{(2\pi)}}{e^{-(np+x)} (np+x)^{np+x+1/2} \sqrt{(2\pi)} \times e^{-(nq-x)} (nq-x)^{nq-x+1/2} \sqrt{(2\pi)}} \cdot \left(\frac{np}{nq}\right)^x$$

$$= \frac{(np)^{np+1/2} (nq)^{nq+1/2}}{(np)^{np+x+1/2} \left\{1 + \frac{x}{np}\right\}^{np+x+1/2} \times \left\{1 - \frac{x}{nq}\right\}^{nq-x+1/2}} \left(\frac{np}{nq}\right)^x$$

$$= \frac{1}{\left(1 + \frac{x}{np}\right)^{np+x+1/2} \left(1 - \frac{x}{nq}\right)^{nq-x+1/2}}$$

$$\text{or } \log \frac{y_x}{y_0} = -(np+x+\frac{1}{2}) \log \left(1 + \frac{x}{np}\right) - (nq-x+\frac{1}{2}) \log \left(\frac{x}{nq}\right)$$

$$\begin{aligned}
 &= -(np + x + \frac{1}{2}) \left[\frac{x}{np} - \frac{x^2}{2n^2 p^2} + \dots \right] \\
 &\quad + (nq - x + \frac{1}{2}) \left[\frac{x}{nq} + \frac{x^2}{2n^2 q^2} + \dots \right] \\
 &= -x + \frac{x^2}{2np} - \frac{x^2}{np} - \frac{x}{2np} + x + \frac{x^2}{2nq} - \frac{x^2}{nq} + \frac{x}{2nq} \\
 &\quad \text{neglecting the terms containing } \frac{1}{n^2} \\
 &= \frac{x^2}{n} \left\{ \frac{1}{2p} - \frac{1}{p} + \frac{1}{2q} - \frac{1}{q} \right\} - \frac{x}{2n} \left\{ \frac{1}{p} - \frac{1}{q} \right\} \\
 &= -\frac{x^2(p+q)}{2npq} - \frac{x(q-p)}{2npq} = -\frac{x^2}{2npq} - \frac{(q-p)}{2npq} x \text{ as } p+q=1
 \end{aligned}$$

Since q and p both are less than 1 and very nearly equal, therefore $\frac{q-p}{2npq}$ can be neglected and then we are left with

$$\log \frac{y_x}{y_0} = -\frac{x^2}{2npq} = -\frac{x^2}{2\sigma^2} \text{ as } \sigma^2 = npq$$

or

$$\frac{y_x}{y_0} = e^{-x^2/2\sigma^2}$$

i.e.,

$$y_x = y_0 e^{-x^2/2\sigma^2}$$

Note. The curve given by $y = y_0 e^{-x^2/2\sigma^2}$ is said to be the normal curve.

A normal curve is symmetrical about the point $x=0$ where the ordinate has its maximum value. In a normal curve, the mean, the median and mode coincide.

[C] The Normal Distribution

We have just introduced that the equation to the normal curve is

$$y = y_0 e^{-x^2/2\sigma^2} \quad \dots (1)$$

$$\text{Its area} = \int_{-\infty}^{\infty} y_0 e^{-x^2/2\sigma^2} dx$$

$$= 2y_0 \int_0^{\infty} e^{-x^2/2\sigma^2} dx, \text{ being symmetrical}$$

$$= 2y_0 \cdot \frac{\sqrt{\pi}}{2 \cdot \frac{1}{\sqrt{2\sigma^2}}} \text{ as } \int_0^{\infty} e^{-a^2 x^2} dx = \frac{\sqrt{\pi}}{2a} \text{ if } a > 0$$

$$= \sigma y_0 \sqrt{(2\pi)} \quad \dots (2)$$

$$= 2.506627 y_0 \sigma \text{ by putting the value of } \sqrt{(2\pi)}.$$

In order to make (1), the normal probability curve, the value of y_0 be so determined that the total frequency is one i.e., the area of the normal curve is unity and this will be so if from (2)

$$\sigma y_0 \sqrt{(2\pi)} = 1$$




Fig. 15.5

$$\text{i.e., } y_0 = \frac{1}{\sigma \sqrt{(2\pi)}} \quad \dots (3)$$

Substituting this value of y_0 in (1), the standard form of the normal curve becomes

$$y = \frac{1}{\sigma \sqrt{(2\pi)}} e^{-x^2/2\sigma^2} \quad \dots (4)$$

which is the normal distribution.

In deriving the form (1) of normal curve, we have taken the mean at the origin, but if however we take another point as the origin such that the excess of the mean over the arbitrary origin is m , the form of the normal curve is

$$y = \frac{1}{\sigma \sqrt{(2\pi)}} e^{-(x-m)^2/2\sigma^2} \quad \dots (5)$$

which represents the standard form of the normal curve with origin at $(m, 0)$.

Physical conditions leading a normal curve

(i) The casual forces that affect individual events are mutually independent.

(ii) The casual forces of equal magnitude are very large in number.

(iii) The casual forces operate in such a way that the maximum frequencies are clustered around the mean value thereby giving a symmetrical curve. Conclusively the deviations below the mean are equal in number and magnitude to those above the mean.

(iv) The normal distribution can be used as an error distribution by inquiring what law of distribution errors of observation should obey in order to make the mean of a set of measurements the most probable value of the 'true' magnitude.

Hence according to Gauss, if we call the 'precision' h , such that $h^2 = \frac{1}{2\sigma^2}$, the form (4) becomes

$$y = \frac{h}{\sqrt{\pi}} e^{-h^2 x^2} \quad \dots (6)$$

It is clear from (6), that as h increases, the normal curve would

become narrower and as such h is a measure of closeness of the mass of observations to the true value.

Definition of a normal distribution. A normal distribution is a continuous distribution given by

$$y = \frac{1}{\sigma\sqrt{(2\pi)}} \cdot e^{-\frac{1}{2}[(x-m)/\sigma]^2},$$

where x is a continuous normal variate distributed with probability density function $f(x) = \frac{1}{\sigma\sqrt{(2\pi)}} e^{-\frac{1}{2}[(x-m)/\sigma]^2}$, with mean m and standard deviation σ .

Properties of a Normal Distribution

(1) When $p=q$ or $p \approx q$ (i.e., p is very nearly equal to q), the distribution fitted is symmetrical. But a normal curve is distinguished from the other symmetrical curves in a remarkable point that a normal curve is symmetrical not only with regard to skewness as are all symmetrical curves; but it is also symmetrical with regard to peakedness (i.e., kurtosis).

(2) The normal curve is a mathematical abstraction, not found in practical work, but used to describe the form of distribution that would be obtained by some continuous data in very large numbers.

(3) The normal curves are based upon regular variation and uniformity conjoined so that no single force playing on each item of the distribution is dominating.

(4) In a normal distribution, items differing from the mean (or median or mode which coincide in a symmetrical distribution) by the same amount in either direction occur with the same frequency. As

such above and below the mean at equal distances, there are same number of measurements.

(5) Normal curve is a single peaked.

(6) Normal curve is asymptotic to the horizontal base as y decreases rapidly when x increases numerically.

(7) The mean, median and mode coincide and lower and upper quartiles are equidistant from the median.




Fig. 15.6

tiles are equidistant from the median.

(8) The curve can be completely specified by mean (i.e., origin of x) and the standard deviation along with the value of y_0 found as in equation (3).

(9) The points of inflexion of the normal curve are obtained by

putting $\frac{d^2y}{dx^2} = 0$ provided $\frac{d^3y}{dx^3} \neq 0$ for these points.

These are found to be

$$x = \pm \sigma, y = \frac{1}{\sigma\sqrt{(2\pi)}} e^{-1/2}.$$

Note. The points of inflexion are the points where the curvature changes its direction.

Constants of Normal Distribution

1. The Mean

Mean $= \mu_1'$ (about the origin)

$$= \frac{1}{\sigma\sqrt{(2\pi)}} \int_{-\infty}^{\infty} x \cdot e^{-[(x-m)^2/2\sigma^2]} dx$$

$$\text{Put } \frac{x-m}{\sqrt{2}\sigma} = z, \text{ i.e. } dx = \sqrt{2}\sigma dz$$

$$= \frac{1}{\sigma\sqrt{(2\pi)}} \int_{-\infty}^{\infty} (m + \sqrt{2}\sigma z) e^{-z^2} dz \cdot \sqrt{2}\sigma$$

$$= \frac{m}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-z^2} dz + \sigma \sqrt{\left(\frac{2}{\pi}\right)} \int_{-\infty}^{\infty} z \cdot e^{-z^2} dz$$

$$= \frac{2m}{\sqrt{\pi}} \int_0^{\infty} e^{-z^2} dz, \text{ the second integral vanishes, being an odd function of } z$$

$$= \frac{2m}{\sqrt{\pi}} \cdot \frac{\sqrt{\pi}}{2} = m.$$

2. The Standard Deviation and Variance

We have μ_2 (about the origin)

$$= \frac{1}{\sigma\sqrt{(2\pi)}} \int_{-\infty}^{\infty} x^2 \cdot e^{-[(x-m)^2/2\sigma^2]} dx$$

$$\text{or } \mu_2' = \frac{1}{\sigma\sqrt{(2\pi)}} \int_{-\infty}^{\infty} (m + \sqrt{2}\sigma z)^2 e^{-z^2} \sqrt{2}\sigma dz \text{ Put } \frac{x-m}{\sqrt{2}\sigma} = z \text{ i.e. } dx = \sqrt{2}\sigma dz$$

$$= \frac{m^2}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-z^2} dz + \frac{2\sqrt{2}\sigma}{\sqrt{\pi}} \int_{-\infty}^{\infty} z \cdot e^{-z^2} dz + \frac{2\sigma^2}{\sqrt{\pi}} \int_{-\infty}^{\infty} z^2 \cdot e^{-z^2} dz$$

$$= \frac{2m^2}{\sqrt{\pi}} \int_0^{\infty} e^{-z^2} dz + 0 + \frac{4\sigma^2}{\sqrt{\pi}} \int_0^{\infty} z^2 e^{-z^2} dz$$

the second integral vanishes by the property of definite integrals

$$= \frac{2m^2}{\sqrt{\pi}} \cdot \frac{\sqrt{\pi}}{2} + \frac{4\sigma^2}{\sqrt{\pi}} \int_0^{\infty} z^2 e^{-z^2} dz$$

$$= m^2 - \frac{2\sigma^2}{\sqrt{\pi}} \int_0^{\infty} z (-2z) e^{-z^2} dz$$

$$= m^2 - \frac{2\sigma^2}{\sqrt{\pi}} \left[\left(z e^{-z^2} \right)_0^\infty - \int_0^\infty e^{-z^2} dz \right] \text{integrating by parts}$$

$$= m^2 + \frac{2\sigma^2}{\sqrt{\pi}} \cdot \frac{\sqrt{\pi}}{2} = m^2 + \sigma^2.$$

$$\therefore \text{The variance } \mu_2 = \mu_2' - \mu_1'^2$$

$$= m^2 + \sigma^2 - m^2 = \sigma^2$$

$$\therefore \text{Standard deviation} = \sqrt{\mu_2} = \sigma.$$

3. The Mean Deviation from the Mean

The mean deviation about the mean

$$m = \int_{-\infty}^{\infty} |x-m| \cdot \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left[\frac{(x-m)}{\sigma}\right]^2} dx$$

$$= \frac{\sigma\sqrt{2}}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} |\sqrt{2}\sigma z| \cdot e^{-z^2} dz \quad \text{Put } \frac{x-m}{\sqrt{2}\sigma} = z$$

$$= \sigma \sqrt{\frac{2}{\pi}} \left[\int_{-\infty}^0 -ze^{-z^2} dz + \int_0^{\infty} ze^{-z^2} dz \right], \text{ i.e., } dx = \sqrt{2}\sigma dz$$

$$= \sigma \sqrt{\frac{2}{\pi}} \left\{ \left[\frac{e^{-z^2}}{-2} \right]_{-\infty}^0 + \left[\frac{e^{-z^2}}{-2} \right]_0^{\infty} \right\}$$

$$= \sigma \sqrt{\frac{2}{\pi}} \left\{ \frac{1}{2} + \frac{1}{2} \right\} = \sigma \sqrt{\frac{2}{\pi}} = 0.7979 \sigma = \frac{4}{5} \sigma \text{ approx.}$$

4. Moments about the Mean

Let us first consider the odd moments about the mean

$$\mu_{2n+1} = \int_{-\infty}^{\infty} (x-m)^{2n+1} \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-m)^2/2\sigma^2} dx.$$

$$\text{Put } \frac{x-m}{\sqrt{2}\sigma} = z, \text{ i.e., } dx = \sqrt{2}\sigma dz$$

$$= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} (\sqrt{2}\sigma z)^{2n+1} e^{-z^2} dz$$

$$= 0 \text{ being an odd function of } z, \text{ by the properties of definite integral.}$$

$$\therefore \mu_3 = \mu_5 = \mu_7 = \dots = 0$$

i.e., all odd moments about the mean are zero.

Let us now consider the even moments about the mean

$$\mu_{2n} = \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} (x-m)^{2n} e^{-(x-m)^2/2\sigma^2} dx$$

$$\text{Put } \frac{x-m}{\sqrt{2}\sigma} = z, dx = \sqrt{2}\sigma dz$$

$$= \frac{2^n \sigma^{2n}}{\sqrt{\pi}} \int_{-\infty}^{\infty} z^{2n} \cdot e^{-z^2} dz$$

$$= \frac{2^{n+1} \cdot \sigma^{2n}}{\sqrt{\pi}} \int_0^{\infty} e^{-z^2} z^{2n-1} \cdot z dz \quad \text{Put } z^2 = t, 2z dz = dt$$

$$= \frac{2^n \sigma^{2n}}{\sqrt{\pi}} \int_0^{\infty} e^{-t} t^{(2n-1)/2} dt = \frac{2^n \sigma^{2n}}{\sqrt{\pi}} \int_0^{\infty} e^{-t} t^{(n+1/2)-1} dt$$

$$= \frac{2^n \sigma^{2n}}{\sqrt{\pi}} \Gamma(n + \frac{1}{2}) \text{ by the definition of Gamma integral.}$$

In particular,

$$\mu_k = \frac{2^{k/2} \cdot \sigma^k}{\sqrt{\pi}} \Gamma\left(\frac{k+1}{2}\right) \text{ when } k=2n$$

$$\text{Now we have } \mu_2 = \frac{2\sigma^2}{\sqrt{\pi}} \Gamma\left(\frac{3}{2}\right) = \frac{2\sigma^2}{\sqrt{\pi}} \cdot \frac{1}{2} \sqrt{\pi} = \sigma^2,$$

$$\mu_4 = \frac{2^2 \sigma^4}{\sqrt{\pi}} \Gamma\left(\frac{5}{2}\right) = \frac{4\sigma^4}{\sqrt{\pi}} \cdot \frac{3}{2} \cdot \frac{1}{2} \sqrt{\pi} = 3\sigma^4.$$

5. The Normal Probability Integral or Error Functions

It has been shown that the total area of the normal curve being unity, it is given by

$$y = \frac{1}{\sigma\sqrt{2\pi}} e^{-x^2/2\sigma^2}$$

or putting $\frac{1}{2\sigma^2} = h^2$, this becomes

$$y = \frac{h}{\sqrt{\pi}} e^{-h^2 x^2}$$

where h is known as *precision*.

Thus the probability that a deviation lies between x and $-x$ is

$$P = \frac{h}{\sqrt{\pi}} \int_{-x}^x e^{-h^2 x^2} dx$$

$$= \frac{2h}{\sqrt{\pi}} \int_0^x e^{-h^2 x^2} dx$$

$$= \frac{2}{\sqrt{\pi}} \int_0^x e^{-h^2 x^2} h dx$$

$$\text{or } \phi(hx) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-h^2 x^2} h dx$$

$$\text{so that } \phi(y) = \frac{2}{\sqrt{\pi}} \int_0^y e^{-y^2} dy$$

which is known as *error function* or the *probability integral*.

Note. The probable error λ or quartile deviation is defined to be the error such that the chance of an error lying within the limits $m-\lambda$ and $m+\lambda$ is exactly the same as chance of an error lying outside these limits i.e.


$$\frac{1}{\sigma\sqrt{2\pi}} \int_{m-\lambda}^{m+\lambda} e^{-\frac{1}{2}\left[\frac{(x-m)}{\sigma}\right]^2} dx = \frac{1}{2}. \text{ Put } \frac{x-m}{\sigma} = z$$

$$\text{or } \frac{1}{\sqrt{2\pi}} \int_0^{\lambda/\sigma} e^{-z^2/2} dz = \frac{1}{2}$$

whence from table $\frac{\lambda}{\sigma} = 0.6745$ i.e. $\lambda = 0.6745\sigma = \frac{2}{3}\sigma$ approx.

$$\therefore Q_1 = m - \frac{2}{3}\sigma \text{ and } Q_3 = m + \frac{2}{3}\sigma.$$

TABLE I—AREA UNDER THE NORMAL CURVE



Area being measured from mean
($x=0$) to distance x/σ .

Fig. 15.7

x/σ	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0754
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
0.6	.2258	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2518	.2549
0.7	.2580	.2612	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
0.8	.2881	.2910	.2939	.2966	.2996	.3023	.3051	.3078	.3106	.3133
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4626	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857

AREA UNDER THE NORMAL CURVE (Contd.)

x/σ	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4983	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.49865	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990
3.1	.49903	.4991	.4991	.4991	.4992	.4992	.4992	.4992	.4993	.4993
3.2	.499313	.4993	.4994	.4994	.4994	.4994	.4994	.4995	.4995	.4995
3.3	.499517	.4995	.4995	.4996	.4996	.4996	.4996	.4996	.4996	.4997
3.4	.499663	.4997	.4997	.4997	.4997	.4997	.4997	.4997	.4997	.4998
3.5	.499767	.4998	.4998	.4998	.4998	.4998	.4998	.4998	.4998	.4998
3.6	.499841	.4998	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999
3.7	.499892	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999
3.8	.499928	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999
3.9	.499952	.5000	.5000	.5000	.5000	.5000	.5000	.5000	.5000	.5000
4.0	.499968	.5000	.5000	.5000	.5000	.5000	.5000	.5000	.5000	.5000

Table for ordinates. Normal curve is given by

$$y = \frac{1}{\sigma\sqrt{2\pi}} e^{-x^2/2\sigma^2} \quad \dots (1)$$

So the tables are to be prepared to give the values of $\frac{1}{\sqrt{2\pi}} e^{-x^2/2\sigma^2}$, the origin of x being taken at the origin whence division of these values by σ will yield the ordinates y as required by (1).

TABLE II—ORDINATES OF THE STANDARD NORMAL CURVE




Fig. 15.8

Ordinates (y) being obtained on dividing by σ , the values of $\frac{1}{\sqrt{2\pi}} e^{-x^2/2\sigma^2}$ given by the following table, origin of x being at the mean.

$z = x/\sigma$	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	.3989	.3989	.3989	.3988	.3986	.3984	.3982	.3980	.3977	.3973
0.1	.3970	.3965	.3961	.3956	.3951	.3945	.3939	.3932	.3925	.3918
0.2	.3910	.3902	.3894	.3885	.3876	.3867	.3857	.3847	.3836	.3825
0.3	.3814	.3802	.3790	.3778	.3765	.3752	.3739	.3725	.3712	.3697
0.4	.3683	.3668	.3653	.3637	.3621	.3605	.3589	.3572	.3555	.3538
0.5	.3521	.3503	.3485	.3467	.3448	.3429	.3410	.3391	.3372	.3352
0.6	.3332	.3312	.3292	.3271	.3251	.3230	.3209	.3187	.3166	.3144
0.7	.3123	.3101	.3079	.3056	.3034	.3011	.2989	.2966	.2943	.2920
0.8	.2897	.2874	.2850	.2827	.2803	.2780	.2756	.2732	.2709	.2685
0.9	.2661	.2637	.2613	.2589	.2565	.2541	.2516	.2492	.2468	.2444
1.0	.2420	.2396	.2371	.2347	.2323	.2299	.2275	.2251	.2227	.2203
1.1	.2179	.2155	.2131	.2107	.2083	.2059	.2036	.2012	.1989	.1965
1.2	.1942	.1919	.1895	.1872	.1849	.1826	.1804	.1781	.1758	.1736
1.3	.1714	.1691	.1669	.1647	.1626	.1604	.1582	.1561	.1539	.1518
1.4	.1497	.1476	.1456	.1435	.1415	.1394	.1374	.1354	.1334	.1315
1.5	.1295	.1276	.1257	.1238	.1219	.1200	.1182	.1163	.1145	.1127
1.6	.1109	.1092	.1074	.1057	.1040	.1023	.1006	.0989	.0973	.0957
1.7	.0940	.0925	.0909	.0893	.0878	.0863	.0848	.0833	.0818	.0804
1.8	.0790	.0775	.0761	.0748	.0734	.0721	.0707	.0694	.0681	.0669
1.9	.0656	.0644	.0632	.0620	.0608	.0596	.0584	.0573	.0562	.0551
2.0	.0540	.0529	.0519	.0508	.0498	.0488	.0478	.0468	.0459	.0449
2.1	.0440	.0431	.0422	.0413	.0404	.0396	.0387	.0379	.0371	.0363

$z = x/\sigma$	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
2.2	.0355	.0347	.0339	.0332	.0325	.0317	.0310	.0303	.0297	.0290
2.3	.0283	.0277	.0270	.0264	.0258	.0252	.0246	.0241	.0235	.0229
2.4	.0224	.0219	.0213	.0208	.0203	.0198	.0194	.0189	.0184	.0180
2.5	.0175	.0171	.0167	.0163	.0153	.0154	.0151	.0147	.0143	.0139
2.6	.0136	.0132	.0129	.0126	.0122	.0119	.0116	.0113	.0110	.0107
2.7	.0104	.0101	.0099	.0096	.0093	.0091	.0088	.0086	.0084	.0081
2.8	.0079	.0087	.0075	.0073	.0071	.0069	.0067	.0065	.0063	.0061
2.9	.0060	.0058	.0056	.0055	.0053	.0051	.0050	.0048	.0047	.0046
3.0	.0044	.0043	.0042	.0040	.0039	.0038	.0037	.0036	.0035	.0034
3.1	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026	.0025	.0025
3.2	.0024	.0023	.0022	.0022	.0021	.0020	.0020	.0019	.0018	.0018
3.3	.0017	.0017	.0016	.0016	.0015	.0015	.0014	.0014	.0013	.0013
3.4	.0012	.0012	.0012	.0011	.0011	.0015	.0010	.0010	.0009	.0009
3.5	.0009	.0008	.0008	.0008	.0008	.0007	.0007	.0007	.0007	.0006
3.6	.0006	.0006	.0006	.0005	.0005	.0005	.0005	.0005	.0005	.0004
3.7	.0004	.0004	.0004	.0004	.0004	.0004	.0003	.0003	.0003	.0003
3.8	.0003	.0003	.0003	.0003	.0003	.0002	.0002	.0002	.0002	.0002
3.9	.0002	.0002	.0002	.0002	.0002	.0002	.0002	.0002	.0001	.0001

Proportions of items included within $\pm\sigma$, $\pm2\sigma$, $\pm3\sigma$ of the mean in normal curve. The total area of a normal curve being treated as unity, the probability corresponding to any interval in the range of the variate is measured by the area under the curve within that interval given by Table I. Hence if m is the mean of the normal distribution, then P , the probability from m to any value x of the variate is given by

$$P = \frac{1}{\sigma\sqrt{2\pi}} \int_m^x e^{-(x-m)^2/2\sigma^2} dx$$

$$\text{Put } \frac{x-m}{\sigma} = z \text{ i.e., } dx = \sigma dz$$

$$= \frac{1}{\sqrt{2\pi}} \int_0^z e^{-z^2/2} dz$$

This value of P is known as the *Probability integral* or *Error function*. Thus

$$P[m - \sigma < x < m + \sigma] = \frac{1}{\sigma \sqrt{2\pi}} \int_{m-\sigma}^{m+\sigma} e^{-\frac{1}{2}[(x-m)/\sigma]^2} dx$$

$$\text{Put } \frac{x-m}{\sigma} = z, dx = \sigma dz$$

$$\therefore P[-1 < z < 1] = \frac{1}{\sqrt{2\pi}} \int_{-1}^1 e^{-z^2/2} dz$$

$$= \frac{2}{\sqrt{2\pi}} \int_0^1 e^{-z^2/2} dz$$

$$= 2 \times .3413 \text{ since from table I, for } z=1,$$

$$\frac{1}{\sqrt{2\pi}} \int_0^1 e^{-z^2/2} dz = .3413$$

$$= .6826$$

which follows that 68.26% of the items in the normal distribution fall between the range $\pm \sigma$ of the mean.

Now

$$P[m - 2\sigma < x < m + 2\sigma] = \frac{1}{\sigma \sqrt{2\pi}} \int_{m-2\sigma}^{m+2\sigma} e^{-\frac{1}{2}[(x-m)/\sigma]^2} dx$$

$$\text{Put } \frac{x-m}{\sigma} = z, \text{ i.e., } dx = \sigma dz$$

$$= \frac{1}{\sqrt{2\pi}} \int_{-2}^2 e^{-z^2/2} dz$$

$$= \frac{2}{\sqrt{2\pi}} \int_0^2 e^{-z^2/2} dz$$

$$= 2 \times .4772 \text{ from Table I for } z=2$$

$$= .9544.$$

So that 95.44% of the items in the normal distribution fall within the range $\pm 2\sigma$ of the mean.

Again

$$P[m - 3\sigma < x < m + 3\sigma] = \frac{1}{\sigma \sqrt{2\pi}} \int_{m-3\sigma}^{m+3\sigma} e^{-\frac{1}{2}[(x-m)/\sigma]^2} dx$$

$$\text{Put } \frac{x-m}{\sigma} = z, \therefore dx = \sigma dz$$

$$= \frac{2}{\sqrt{2\pi}} \int_0^3 e^{-z^2/2} dz$$

$$= 2 \times .49865$$

$$= .99720,$$

i.e., 99.72% of the items in the normal distribution fall within the range $\pm 3\sigma$ of the mean.

Conclusively a normal curve can be used to find

- (i) The number of cases at any given distance from the mean;
- (ii) the number of cases lying within certain range of values in the distribution;
- (iii) the probability that a case selected at random lies above or below a given point.

Problem 115. If n is large and neither of p and q is too close to zero, then show that the binomial distribution can be closely approximated by a normal distribution with standardized variable given by $z = \frac{x - np}{\sqrt{npq}}$, where x is the binomial variate with mean np and standard deviation $\sqrt{npq}$.

From the given properties of x , we conclude that z is a variate with mean zero and variance unity. Also as x goes from 0 to n , z takes

values $\frac{-np}{\sqrt{npq}}$ to $\frac{nq}{\sqrt{npq}}$ so that the total range of z is

$$\frac{nq}{\sqrt{npq}} - \left(\frac{-np}{\sqrt{npq}} \right) = \frac{n(p+q)}{\sqrt{npq}} = \frac{n}{\sqrt{npq}}, \text{ i.e., } z \text{ jumps } \frac{1}{\sqrt{npq}} \text{ at each stage.}$$

Now

$$\frac{1}{\sqrt{npq}} \rightarrow 0 \text{ as } n \rightarrow \infty$$

$$\frac{nq}{\sqrt{npq}} \rightarrow +\infty \text{ as } n \rightarrow \infty, p, q \neq 0$$

$$\frac{-np}{\sqrt{npq}} \rightarrow -\infty \text{ as } n \rightarrow \infty, p, q \neq 0.$$

As such the distribution of z is a continuous distribution from $-\infty$ to $+\infty$, with mean zero and variance unity.

If $P(n, x)$ denotes the probability for the variate taking the value x , then $P(n, x) = {}^nC_x p^x q^{n-x}$

$$= \frac{n!}{x!(n-x)!} p^x q^{n-x}$$

Applying Stirling's theorem i.e., $n! = \sqrt{2\pi} \cdot e^{-n} n^{n+1/2}$, we get

$$P_n = \lim_{n \rightarrow \infty} P(n, x) = \lim_{n \rightarrow \infty} \frac{\sqrt{2\pi} \cdot e^{-n} n^{n+1/2} \cdot p^x q^{n-x}}{\sqrt{2\pi} \cdot e^{-x} x^{x+1/2} \cdot \sqrt{2\pi} \cdot e^{-(n-x)} (n-x)^{n-x+1/2}}$$

$$= \lim_{n \rightarrow \infty} \left(\frac{1}{B} \cdot \frac{1}{\sqrt{2\pi npq}} \right)$$

where

$$B = \left(\frac{x}{np} \right)^{x+1/2} \left(\frac{n-x}{nq} \right)^{n-x+1/2} \dots (1)$$

Now given substitution is $z = \frac{x - np}{\sqrt{npq}}$

or

$$x = np + z\sqrt{npq}$$

$$i.e., \frac{x}{np} = 1 + z \sqrt{\left(\frac{q}{np}\right)} \text{ and } \frac{n-x}{nq} = \frac{1}{nq} [n - np - z\sqrt{npq}]$$

$$= \frac{1}{nq} [nq - z\sqrt{npq}]$$

$$= 1 - z \sqrt{\left(\frac{p}{nq}\right)} \quad \dots (2)$$

Taking logarithms of both sides of (1), we have

$$\log B = (x + \frac{1}{2}) \log \frac{x}{np} + (n - x + \frac{1}{2}) \log \frac{n-x}{nq}.$$

Making substitutions from (2), this becomes

$$\log B = [np + z\sqrt{npq}] \log \left[1 + z \sqrt{\left(\frac{q}{np}\right)} \right]$$

$$+ [nq - z\sqrt{npq}] \log \left[1 - z \sqrt{\left(\frac{p}{nq}\right)} \right]$$

$$= [np + z\sqrt{npq}] \left\{ z \sqrt{\left(\frac{q}{np}\right)} - \frac{1}{2} \frac{z^2 q}{np} + \dots \right\}$$

$$+ [nq - z\sqrt{npq}] \left\{ -z \sqrt{\left(\frac{p}{nq}\right)} - \frac{1}{2} \frac{z^2 p}{nq} - \dots \right\}$$

$$= \frac{z}{2\sqrt{n}} \left(\sqrt{\frac{q}{p}} - \sqrt{\frac{p}{q}} \right) + \frac{z^2}{2} - \frac{z^2}{4n} \left(\frac{q}{p} + \frac{p}{q} \right) + \text{terms}$$

containing higher power of $(1/n)$

when $n \rightarrow \infty$, $\log B \rightarrow \frac{z^2}{2}$ i.e. $B \rightarrow e^{+z^2/2}$

Now as x takes integral values, z jumps through $\frac{1}{\sqrt{npq}}$ i.e., $(npq)^{-1/2}$ so that increment in z i.e., $dz = (npq)^{-1/2}$ when $n \rightarrow \infty$. Thus, if dP represents the probability for the variate z to lie within the range $z - \frac{1}{2} dz$ to $z + \frac{1}{2} dz$, then

$$dP = \frac{1}{\sqrt{2\pi}} e^{-z^2/2} dz, \quad -\infty \leq z \leq \infty$$

which is the required normal distribution for z .

$$\text{Hence } f(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2} \text{ as } dP = f(z) dz.$$

If m be the mean and σ the standard deviation of the normal distribution, then we can replace z by $\frac{x-m}{\sigma}$ and dz by $\frac{1}{\sigma} dx$,

whence we have

$$dP = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}[(x-m)/\sigma]^2} dx$$

giving

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}[(x-m)/\sigma]^2}$$

so that $y = f(x)$ i.e. $y = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}[(x-m)/\sigma]^2}$

gives the normal probability curve.

Problem 116. If skulls are classified A, B, C according as the length, breadth index as under 75, between 75 and 80 or over 80 find approximately (assuming that the distribution is normal) the mean and standard deviation of a series in which A are 58 per cent, B are 38 per cent and C are 4 per cent being given that if

$$f(t) = \frac{1}{\sqrt{2\pi}} \int_0^t e^{-(x^2/2)} dx,$$

then $f(0.20) = 0.08$ and $f(1.75) = 0.46$.

Let m be the mean and σ the standard deviation for the given distribution.

As given the area between $t=0$ and $t=0.20$ is 0.08, so that the area to the left of this ordinate is $0.5 + 0.08 = 0.58$ which corresponds to $x=75$.




Fig. 15.9

$$\therefore \frac{75-m}{\sigma} = 0.20. \quad \dots (1)$$

Also the area to the right of the ordinate at $x=80$ is 0.04, so that the area to the left of this ordinate is $1 - 0.04 = 0.96$ i.e. the area from 0 to $t = \frac{80-m}{\sigma}$ is $0.96 - 0.5 = 0.46$, which corresponds to $t=1.75$.

$$\therefore \frac{80-m}{\sigma} = 1.75. \quad \dots (2)$$

(1) and (2) are

$$0.20\sigma = 75 - m, \quad 1.75\sigma = 80 - m.$$

$$\text{Subtracting, } 1.55\sigma = 5, \text{ i.e., } \sigma = \frac{5}{1.55} = 3.2 \text{ approx.}$$

and then

$$m = 75 - 0.20 \times 3.2$$

$$= 75 - 0.64 = 74.4 \text{ approx.}$$

Problem 117. In a normal distribution 31 per cent of the items are under 45 and 8 per cent are over 64. Find the mean and standard deviation of the distribution.

Let m be the mean and σ the standard deviation of the distribution. Since 31 per cent of the items are under 45; therefore 19 per cent of the items lie between 45 and m . Thus if

$$\phi(t) = \frac{1}{\sqrt{2\pi}} e^{-t^2/2} \quad \text{where } t = \frac{x-m}{\sigma},$$

then $\int_{(45-m)/\sigma}^0 \phi(t) dt = .19$, i.e. $\int_0^{(m-45)/\sigma} \phi(t) dt = .19$.

From Table I, we have $\int_0^{0.5} \phi(t) dt = .1915$.

So $\frac{m-45}{\sigma} = .5$ approx. ... (1)

Again since 8 per cent of the items are over 64, therefore 42 per cent lie between m and 64, so that $\int_0^{(64-m)/\sigma} \phi(t) dt = .42 = \int_0^{1.4} \phi(t) dt$ nearly from Table I.

$\therefore \frac{64-m}{\sigma} = 1.4$ (2)

Solving (1) and (2), we get $m=50$, $\sigma=10$.

Problem 118. If x is a normal variate with mean $\bar{x}$ and S.D. σ , find the mean and variance of variate y defined by

$$y = \frac{1}{2} \left(\frac{x - \bar{x}}{\sigma} \right)^2$$

For the normal variate x , we have

$$f(x) dx = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\{(x-\bar{x})/\sigma\}^2} dx. \quad \dots (1)$$

If $y = \frac{1}{2} \left(\frac{x - \bar{x}}{\sigma} \right)^2$, i.e., $dy = \frac{x - \bar{x}}{\sigma} \cdot \frac{dx}{\sigma}$ or $dx = \frac{\sigma}{\sqrt{2y}} dy$,

then (1) becomes

$$f(y) dy = \frac{1}{\sigma\sqrt{2\pi}} e^{-y} \cdot \frac{\sigma}{\sqrt{2y}} dy = \frac{1}{2\sqrt{\pi}} y^{-1/2} e^{-y} dy,$$

so that

mean $= \mu_1'$ (about the origin), for a normal distribution

$$= 2 \int_0^\infty y f(y) dy = \frac{1}{\sqrt{\pi}} \int_0^\infty y^{1/2} e^{-y} dy$$

$$= \frac{1}{\sqrt{\pi}} \cdot \Gamma_{\frac{3}{2}} \text{ (by gamma integrals)} = \frac{1}{\sqrt{\pi}} \cdot \frac{1}{2} \sqrt{\pi} = \frac{1}{2}$$

and

$$\begin{aligned} \mu_2' &= 2 \int_0^\infty y^2 f(y) dy \\ &= \frac{1}{\sqrt{\pi}} \int_0^\infty y^{3/2} e^{-y} dy = \frac{1}{\sqrt{\pi}} \Gamma_{\frac{5}{2}} = \frac{1}{\sqrt{\pi}} \cdot \frac{3}{2} \cdot \frac{1}{2} \sqrt{\pi} = \frac{3}{4}; \end{aligned}$$

$$\therefore \sigma^2 = \mu_2' - \mu_1'^2 = \frac{3}{4} - \frac{1}{4} = \frac{1}{2} = \text{variance.}$$

Problem 119. If a normal distribution is grouped in intervals of total frequency N and S is the sum of squares of the frequencies, an estimate of the standard deviation σ is given by $\frac{N^2}{2S\sqrt{\pi}}$.

The normal frequency distribution with mean m and standard deviation σ is given by

$$dF = f(x) dx = y dx = \frac{N}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}[(x-m)/\sigma]^2}$$

where the ordinate of the normal curve is

$$y = \frac{N}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}[(x-m)/\sigma]^2}.$$

$\therefore$ square of the ordinate is

$$y^2 = \frac{N^2}{\sigma^2 \cdot 2\pi} e^{-[(x-m)/\sigma]^2},$$

as such the distribution function for y^2 is

$$\begin{aligned} dP = y^2 dx &= \frac{N^2}{2\pi\sigma^2} e^{-[(x-m)/\sigma]^2} dx \\ &= \frac{N^2}{2\sigma\sqrt{\pi}} \cdot \frac{1}{\left(\frac{\sigma}{\sqrt{2}}\right)\sqrt{2\pi}} e^{-\frac{1}{2}(x-m)^2/(\sigma/\sqrt{2})^2} dx, \end{aligned}$$

which gives a normal distribution with S.D. $\sigma/\sqrt{2}$ and total frequency $N^2/(2\sigma\sqrt{\pi})$. Hence taking S as equivalent to this value, we have

$$S = \frac{N^2}{2\sigma\sqrt{\pi}}, \text{ i.e. } \sigma = \frac{N^2}{2S\sqrt{\pi}}.$$

Problem 120. The local authorities in a certain city instal 2000 lamps in the streets of the city. If the lamps have an average life of 1000 burning hours with a standard deviation of 200 hours, (i) what number of lamps might be expected to fail in first 700 burning hours, and (ii) after what period of burning hours would we expect that 10 per cent of the lamps would have failed? Assume that the lives of the lamps are normally distributed. You are given that

$$F(1.50) = 0.933, F(1.28) = .900 \text{ where } F(z) = \int_{-\infty}^z \frac{1}{\sigma\sqrt{2\pi}} e^{-z^2/2} dz$$

(i) We have the average life of the lamp i.e., $m=1000$ hour and standard deviation $\sigma=200$ hours.

$\therefore$ The normal distribution being symmetrical, the area of the normal curve to the left of $x=700$ is equal to that to the right of $x=(2000-700)=1300$.

$$\therefore z = \frac{x-m}{\sigma} = \frac{1300-1000}{200} = 1.50$$

$$\text{and area to the left of } x=1300 \text{ is } = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^z e^{-z^2/2} dz = F(z)$$

where $z=1.50$ and also $F(1.50)=0.933$ which gives the probability of a value of x lying to the left of $x=1300$ or right of $x=700$.

$\therefore$ probability of its failure $= 1 - .933 = .067$.

So that the number of lamps expected to fail in first 700 hours $= 2000 \times .067 = 134$.

(ii) The failure of 10 per cent of lamps gives that a value of x is to be so found that the area of the standard normal curve to the left of it is $\frac{10}{100}$ i.e., 0.1. But the area to the left of $z=1.28$ is .900, hence by symmetry in the distribution of a normal curve it follows that the area to the left of $z=-1.28$ is equal to the area to the right of $z=1.28$, i.e., 0.1.

$$\text{Hence } z = -1.28 = \frac{x-m}{\sigma} = \frac{x-1000}{200}$$

giving $x = 1000 - 256 = 744$.

Hence after 744 hours, it is expected that 10 per cent of the lamps fail.

[D] The Poisson Distribution

We have shown in the preceding sections that the normal curve is the limit to the binomial, whether p is or is not equal to q provided that n becomes sufficiently large in order to make $(q-p)$ negligibly small as compared to $\sqrt{npq}$. Now we have to consider the limit to the same series if p (or q) becomes sufficiently small and n is increased sufficiently to keep the product np (or nq) a finite number say m i.e., $np=m$.

The probability of r successes in the binomial distribution is given by the $(r+1)$ th term in the binomial expansion $(q+p)^n$, i.e.,

$$P(r) = {}^nC_r p^r q^{n-r} = \frac{n!}{r!(n-r)!} p^r q^{n-r}$$

$$= \frac{n!}{r!(n-r)!} \left(\frac{m}{n}\right)^r \left(1 - \frac{m}{n}\right)^{n-r}$$

$$\left(\because np=m, \text{ i.e., } p=\frac{m}{n} \text{ and so } q=1-p=1-\frac{m}{n} \text{ as } p+q=1\right)$$

$$= \frac{m^r}{r!} \left(1 - \frac{m}{n}\right)^n \cdot \frac{n}{(n-r)!} \left(1 - \frac{m}{n}\right)^{n-r}$$

$$\text{Here } \lim_{n \rightarrow \infty} \left(1 - \frac{m}{n}\right)^n = \lim_{n \rightarrow \infty} \left\{ \left(1 - \frac{m}{n}\right)^{-n/m} \right\}^{-m} = e^{-m}$$

and applying Stirling's approximation i.e., $n! = n^n e^{-n} \sqrt{(2\pi n)}$, when n is large,

$$P(r) = \frac{m^r e^{-m}}{r!} \cdot \frac{n^n e^{-n} \sqrt{(2\pi n)}}{(n-r)^{n-r} e^{-(n-r)} \sqrt{2\pi(n-r)} \times n^r \left(1 - \frac{m}{n}\right)^r}$$

$$= \frac{m^r e^{-m} e^{-r}}{r!} \left(1 - \frac{r}{n}\right)^{-n} \left(1 - \frac{r}{n}\right)^{r-\frac{1}{2}} \left(1 - \frac{m}{n}\right)^{-r}$$

$$= \frac{m^r e^{-m} e^{-r}}{r!} e^r \because \lim_{n \rightarrow \infty} \left(1 - \frac{r}{n}\right)^{-n} = \lim_{n \rightarrow \infty} \left\{ \left(1 - \frac{r}{n}\right)^{-n/r} \right\}^r = e^r;$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{r}{n}\right)^{r-\frac{1}{2}} = 1 \text{ and } \lim_{n \rightarrow \infty} \left(1 - \frac{m}{n}\right)^{-r} = 1$$

$$= \frac{m^r e^{-m}}{r!}$$

Hence successive terms in the expansion of $(q+p)^n$ are

$$e^{-m}, e^{-m} \cdot m, e^{-m} \frac{m^2}{2!}, e^{-m} \frac{m^3}{3!}, \text{ etc.}$$

and the limiting value of $(q+p)^n$ is

$$e^{-m} \left[1 + m + \frac{m^2}{2!} + \frac{m^3}{3!} + \dots \right]$$

This expression is known as *Poisson's distribution* or *Poisson's Exponential limit*.

Note 1. The limiting form of $N(q+p)^n$ is

$$N e^{-m} \left[1 + m + \frac{m^2}{2!} + \frac{m^3}{3!} + \dots \right]$$

Note 2. The quantity m introduced in Poisson distribution is said to be the parameter of binomial distribution.

Note 3. Characteristics of the Poisson distribution.

(i) This is the limiting form of binomial distribution when n is large and p (or q) is small.

(ii) Here p or q is very close to zero or unity, but if p is close to zero, the distribution is J-shaped or unimodal.

(iii) Since it consists of a single parameter m , the entire distribution can therefore be obtained by knowing the mean only.

Note 4. Some examples of Poisson distribution are:

(i) The number of defective screws per box of 100 screws.

(ii) The number of typographical errors per page in typed material.

(iii) The number of cars passing through a certain street in time t .

Constants of the Poisson Distribution

Assume that origin is located at the first term of the distribution, so that the values of the deviation from the assumed origin are 0, 1, 2,

1. The Mean

$$\text{Mean} = \mu_1' \text{ (about the origin)} = \sum_0^{\infty} e^{-m} \frac{m^r}{r!} \cdot r$$

$$= e^{-m} \left[0 + m + \left(\frac{m^2}{2!} \times 2\right) + \left(\frac{m^3}{3!} \times 3\right) + \dots \right]$$

$$= m e^{-m} \left(1 + \frac{m}{1!} + \frac{m^2}{2!} + \dots \right) = m e^{-m} \cdot e^m = m$$

2. The Variance

We have μ_2' (about the origin)

$$= \sum_0^{\infty} e^{-m} \frac{m^r}{2!} \cdot r^2$$

$$= e^{-m} \left[0 + m + \left(\frac{m^2}{2!} \times 2^2\right) + \left(\frac{m^3}{3!} \times 3^2\right) + \dots \right]$$

$$\begin{aligned}
&= me^{-m} \left[1 + \left(\frac{m}{1!} \times 2 \right) + \left(\frac{m^2}{2!} \times 3 \right) + \dots \right] \\
&= me^{-m} \left[1 + \frac{m}{1!} (1+1) + \frac{m^2}{2!} (2+1) + \dots \right] \\
&= me^{-m} \left[\left(1 + \frac{m}{1!} + \frac{m^2}{2!} + \dots \right) + \left(m + \frac{m^2}{1!} + \frac{m^3}{2!} + \dots \right) \right] \\
&= me^{-m} \left[e^m + m \left(1 + \frac{m}{1!} + \frac{m^2}{2!} + \dots \right) \right] \\
&= me^{-m} [e^m + me^m] = m(m+1)
\end{aligned}$$

$\therefore$ variance $= \mu_2 = \sigma^2 = \mu_2' - \mu_1'^2 = m^2 + m - m^2 = m$
so that standard deviation, $\sigma = \sqrt{m}$.

3. The Moments

μ_3' (about the origin)

$$\begin{aligned}
&= \sum e^{-m} \frac{m^r}{r!} \cdot r^3 \\
&= e^{-m} \sum m^r \cdot \left\{ \frac{r(r-1)(r-2) + 3r(r-1) + r}{r!} \right\} \\
&= e^{-m} m^3 \sum \frac{m^{r-3}}{(r-3)!} + e^{-m} \cdot 3m^2 \sum \frac{m^{r-2}}{(r-2)!} + e^{-m} \cdot m \sum \frac{m^{r-1}}{(r-1)!}
\end{aligned}$$

or $\mu_3' = m^3 + 3m^2 + m$

so that $\mu_3 = \mu_3' - 3\mu_2' \cdot \mu_1' + 2\mu_1'^3$

$$= m^3 + 3m^2 + m - 3m^2 - m^3 + 2m^3 = m$$

and

$$\begin{aligned}
\mu_4' &= \sum e^{-m} \cdot \frac{m^r}{r!} \cdot r^4 \\
&= e^{-m} \sum m^r \cdot \left\{ \frac{r(r-1)(r-2)(r-3) + 6r(r-1)(r-2) + 7(r-1)r + r}{r!} \right\} \\
&= e^{-m} \cdot m^4 \sum \frac{m^{r-4}}{(r-4)!} + e^{-m} \cdot 6m^3 \sum \frac{m^{r-3}}{(r-3)!} + e^{-m} \cdot 7m \sum \frac{m^{r-2}}{(r-2)!} \\
&\quad + e^{-m} \cdot m \sum \frac{m^{r-1}}{(r-1)!}
\end{aligned}$$

$$= m^4 + 6m^3 + 7m^2 + m$$

so that $\mu_4 = \mu_4' - 4\mu_3' \mu_1' + 6\mu_2' \mu_1'^2 - 3\mu_1'^4$

$$= m^4 + 6m^3 + 7m^2 - 4m(m^3 + 3m^2 + m) + 6m^2(m^2 + m) - 3m^4$$

$$= 3m^2 + m.$$

Problem 121. Define a Poisson random variable and give some physical situations illustrating it. Find out its mean and variance.

If x and y are independently distributed as Poisson random variates with parameters λ and μ respectively find the probability distribution of $X+Y$.

We know that a random variable is a function defined as a sample space. A random variable x assuming values $0, 1, 2, \dots, r, \dots$ with probabilities

$$e^{-m}, me^{-m}, \frac{m^2}{2!} e^{-m}, \dots, \frac{m^r e^{-m}}{r!}, \dots$$

i.e.,

$$P(x=r) = \frac{m^r e^{-m}}{r!}, r=0, 1, 2, \dots$$

is said to be a Poisson random variable.

In order to illustrate it with the help of a physical situation, let us consider a random variate X denoting the number of calls during a time t at a telephone switch board. Let us assume that the calls are independent and the probability of a call in time dt is λdt .

Denoting by $P_x(t)$ and $P_x(t+\delta t)$, the chance of x calls in times t and $t+\delta t$ respectively we have two mutually exclusive possibilities, (i) there are x -calls in t and no call in time dt , (ii) there are $(x-1)$ calls in t and one call in dt , neglecting the possibility of more than one call in dt , for it would be of order 2 (i.e., dt^2) and higher.

Thus,

$$\begin{aligned}
P_x(t+\delta t) &= P_x(t) \cdot \{1 - \lambda \delta t\} + P_{x-1}(t) \cdot \lambda \delta t \\
&= P_x(t) - P_x(t) \cdot \lambda \delta t + P_{x-1}(t) \cdot \lambda \delta t
\end{aligned}$$

or

$$P_x(t+\delta t) - P_x(t) = \{P_{x-1}(t) - P_x(t)\} \cdot \lambda \delta t$$

or

$$\frac{P_x(t+\delta t) - P_x(t)}{\delta t} = \lambda [P_{x-1}(t) - P_x(t)].$$

Proceeding to the limit as $\delta t \rightarrow 0$, we have

$$\lim_{\delta t \rightarrow 0} \frac{P_x(t+\delta t) - P_x(t)}{\delta t} = \lim_{\delta t \rightarrow 0} \lambda [P_{x-1}(t) - P_x(t)]$$

i.e.,

$$\frac{dP_x(t)}{dt} = \lambda [P_{x-1}(t) - P_x(t)].$$

Putting

$$P_x(t) = \frac{(\lambda t)^x}{x!} \cdot f(t)$$

so that

$$P_0(t) = f(t) \text{ and } P_0(0) = f(0) = 1, \dots (1)$$

We have $\frac{d}{dt} \left[\frac{(\lambda t)^x}{x!} f(t) \right] = \lambda \left[\frac{(\lambda t)^{x-1}}{(x-1)!} f(t) - \frac{(\lambda t)^x}{x!} f(t) \right]$

or $\frac{(\lambda t)^x}{x!} f'(t) + \frac{\lambda^x}{x!} f(t) \cdot x t^{x-1} = \frac{\lambda^x t^{x-1}}{(x-1)!} f(t) - \frac{\lambda t}{x} \cdot \frac{\lambda^x t^{x-1}}{(x-1)!} f(t).$

Dividing throughout by $\frac{\lambda^x t^{x-1}}{(x-1)!} f(t)$, this reduces to

$$\frac{\lambda t}{x} \cdot \frac{f'(t)}{f(t)} + 1 = 1 - \frac{\lambda t}{x} \text{ or } \frac{f'(t)}{f(t)} = -\lambda.$$

Integrating with regard to t , this yields

$$\log f(t) = -\lambda t + A, A \text{ being constant of integration.}$$

Initially when $t=0$, $f(0)=1$, from (1), giving $A=0$

$$\log f(t) = -\lambda t \text{ or } f(t) = e^{-\lambda t}$$

or

$$\frac{(\lambda t)^x}{x!} f(t) = P_x(t) = e^{-\lambda t} \cdot \frac{(\lambda t)^x}{x!}$$

where $\sum_{x=0}^{\infty} P_x(t) = e^{-\lambda t} \sum_{x=0}^{\infty} \frac{(\lambda t)^x}{x!} = e^{-\lambda t} \cdot e^{\lambda t} = 1.$

It follows that the number of calls in a fixed time t is a Poisson variate with parameter λt .

For the second part, we have

$$P(X=x) = e^{-\lambda} \cdot \frac{\lambda^x}{x!}, \quad P(Y=y) = e^{-\mu} \cdot \frac{\mu^y}{y!}$$

so that $P(X=x, Y=y) = P(X=x) \cdot P(Y=y)$

$$= e^{-\lambda} \frac{\lambda^x}{x!} \cdot e^{-\mu} \frac{\mu^y}{y!} = e^{-(\lambda+\mu)} \frac{\lambda^x \mu^y}{\left| \begin{smallmatrix} x & y \end{smallmatrix} \right|}$$

Since the variable take values 0, 1, 2, 3, ...; let us find the Probability that their sum i.e. $x+y$ takes value r so that $y=r-x$. Summing for all values of x from 0 to r , we have

$$\begin{aligned} P(x+y=r) &= e^{-(\lambda+\mu)} \sum_{x=0}^r \frac{\lambda^x \mu^{r-x}}{\left| \begin{smallmatrix} x & r-x \end{smallmatrix} \right|} = e^{-(\lambda+\mu)} \cdot \frac{\mu^r}{\left| \begin{smallmatrix} r \end{smallmatrix} \right|} \\ &\quad \sum_{x=0}^r \frac{\left| \begin{smallmatrix} r \\ x & r-x \end{smallmatrix} \right|}{\left| \begin{smallmatrix} r \end{smallmatrix} \right|} \left(\frac{\lambda}{\mu} \right)^x \\ &= e^{-(\lambda+\mu)} \cdot \frac{\mu^r}{\left| \begin{smallmatrix} r \end{smallmatrix} \right|} \sum_{x=0}^r {}^r C_x \left(\frac{\lambda}{\mu} \right)^x \\ &= e^{-(\lambda+\mu)} \cdot \frac{\mu^r}{\left| \begin{smallmatrix} r \end{smallmatrix} \right|} \left[{}^r C_0 + {}^r C_1 \cdot \frac{\lambda}{\mu} + {}^r C_2 \left(\frac{\lambda}{\mu} \right)^2 + \dots + {}^r C_r \left(\frac{\lambda}{\mu} \right)^r \right] \\ &= e^{-(\lambda+\mu)} \cdot \frac{\mu^r}{\left| \begin{smallmatrix} r \end{smallmatrix} \right|} \left(1 + \frac{\lambda}{\mu} \right)^r = e^{-(\lambda+\mu)} \cdot \frac{(\lambda+\mu)^r}{\left| \begin{smallmatrix} r \end{smallmatrix} \right|} \therefore r=x+y \end{aligned}$$

which is a Poisson distribution with mean $(\lambda+\mu)$.

Problem 122. Find the probability that at most 5 defective fuses will be found in a box of 200 fuses if experience shows that 2 per cent of such fuses are defective.

Here $n=200$, $q=\frac{2}{100}=0.02$, so that $m=nq=4$.

Hence $P(x \leq 5) = \sum_{x=0}^5 e^{-4} \cdot \frac{4^x}{x!}$

$$= e^{-4} \left[1 + 4 + \frac{4^2}{2!} + \frac{4^3}{3!} + \frac{4^4}{4!} + \frac{4^5}{5!} \right]$$

$$= e^{-4} \left[1 + 4 + 8 + \frac{64}{3} + \frac{64}{3} + \frac{128}{15} \right]$$

$$= e^{-4} \cdot \frac{649}{15} = 0.0183 \times \frac{649}{15} = .785 \text{ approx.}$$

$\therefore e^{-4} = 0.0183$

Problem 123. Fit a Poisson's distribution to the set of observations

Deaths	0	1	2	3	4
Frequency	122	60	15	2	1

and calculate the theoretical frequencies.

We have, Mean = $\frac{0 \times 122 + 1 \times 60 + 2 \times 15 + 3 \times 2 + 4 \times 1}{122 + 60 + 15 + 2 + 1}$

$$= \frac{100}{200} = 0.5$$

$\therefore e^{-0.5} = 1 - (0.5) + \frac{1}{2} (0.5)^2 - \frac{1}{6} (0.5)^3 + \dots$

$$= 1 - 0.5 + 0.125 - 0.0208 + \dots = .61 \text{ approx.}$$

The theoretical frequency of r deaths is given by

$$Ne^{-m} \cdot \frac{m^r}{r!} = 200 \cdot e^{-0.5} \cdot \frac{(0.5)^r}{r!} = 122 \cdot \frac{(0.5)^r}{r!}$$

which gives for $r=0, 1, 2, 3, 4$ the theoretical frequencies as 122, 61, 15, 2 and 0 respectively.

Problem 124. In a certain factory turning out razor blades, there is a small chance $\frac{1}{5000}$ for any blade to be defective. The blades are supplied in packets of 10. Use Poisson distribution to calculate the approximate number of packets containing no defective, one defective and two defective blades respectively in a consignment of 10,000 packets, given that $e^{-0.02} = 0.9802$.

Here $N=10,000$, $m=np=10 \times \frac{1}{5000} = 0.02$ and $e^{-0.02} = .9802$.

$\therefore$ Required frequencies are given by Ne^{-m} , $Ne^{-m} \cdot m$, $Ne^{-m} \cdot \frac{m^2}{2!}$

i.e., $10,000 \times .9802$; $10,000 \times .9802 \times 0.02$; $10,000 \times .9802 \times \frac{(0.02)^2}{2}$

i.e., 9802; 196, 2 packets.

Problem 125. If x is a Poissonian variate with mean m ,

(a) What would be the expectation of $e^{-kx} \cdot kx$ where k is a constant.

(b) Show that, expectation of $(e^{-kx}) = e^{-m(1-e^{-k})}$.

(a) We have, $E(kxe^{-kx}) = \sum_{x=0}^{\infty} kx \cdot e^{-kx} \cdot \frac{e^{-m} m^x}{x!}$

$$= mke^{-(m+k)} \sum_{x=0}^{\infty} e^{-k(x-1)} \frac{m^{x-1}}{(x-1)!}$$

$$= mke^{-(m+k)} \sum_{x=0}^{\infty} \frac{(me^{-k})^{x-1}}{(x-1)!}$$

$$= mke^{-(m+k)} e^{me^{-k}} = mke^{m(e^{-k}-1)-k}$$

and $E(e^{-kx}) = \sum_{x=0}^{\infty} e^{-kx} \cdot \frac{e^{-m} m^x}{x!} = e^{-m} \sum_{x=0}^{\infty} \frac{(me^{-k})^x}{x!}$

$$= e^{-m} \cdot e^{me^{-k}} = e^{-m(1-e^{-k})}$$

15.17. OTHER DISTRIBUTIONS

(A) Casual distribution. This is a distribution with one parameter say u like Poisson distribution and found casually in exact sciences. In this type of distribution the variate takes with certainty one value, i.e. $x=u$, such that $P(x=u)=1$,

$$\begin{aligned} P(x=t) &= 0 \text{ when } t \neq u, \\ P(x \leq t) &= 0 \text{ when } t < u, \\ &= 1 \text{ when } t \geq u. \end{aligned}$$

Problem 126. Show that the moment generating function for a casual distribution is e^{ut} , i.e., $M_0(t) = e^{ut}$.

$$\begin{aligned} \text{We have } M_0(t) &= \sum_{u=0}^{\infty} P(x=u) e^{ut} \\ &= \sum_{u=0}^{\infty} e^{ut} \text{ as } P(x=u) = 1 \\ &= e^{ut} \text{ as } P(x=t) = 0, t \neq u. \end{aligned}$$

[B] Rectangular or uniform distribution. A distribution in which the variate takes values $x_1, x_2, \dots, x_n$ (or 1, 2, ..., n in particular) each having the same probability $1/n$, is said to be a rectangular or uniform distribution. Thus,

$$P(x=x_i) = \frac{1}{n}, i=1, 2, \dots, n.$$

$$\therefore \sum_{i=1}^n P(x=x_i) = \frac{1}{n} + \frac{1}{n} + \dots n \text{ times} = 1$$

In tossing a coin or throwing a die, this type of distribution appears.

Problem 127. If $f(x)$ be the probability law of continuous random variable x in the interval $a \leq x \leq b$, so that $f(x) = 0$ for $x < a$ and $x > b$ and $\int_a^b f(x) dx = 1$, show that the variable $P = \int_a^x f(x) dx$ has a rectangular distribution.

$$\text{Given } P = \int_a^x f(x) dx = P(X \leq x), \text{ so that } dP = f(x) dx.$$

Assuming $u = F(x)$, where $F'(x) = f(x)$, we have

$$\begin{aligned} dP &= f(x) dx = f(x) \frac{dx}{du} du \\ &= f(x) \frac{1}{f(x)} du = du, 0 \leq u \leq 1. \end{aligned}$$

Since u is a distribution with unit range, therefore $u = P$ is a rectangular distribution.

Problem 128. A variate x has uniform distribution over the unit interval. Find the function of x having the distribution

$$\begin{aligned} dP &= e^{-x} dx, 0 \leq x \leq \infty. \\ dP &= e^{-x} dx. \end{aligned}$$

Here

Suppose $u = F(x)$ where $F'(x) = f(x) = e^{-x}$

$$\begin{aligned} &= \int_0^x e^{-x} dx \\ &= \left[-e^{-x} \right]_0^x = 1 - e^{-x} \end{aligned}$$

$$\text{giving } x = \log_e \frac{1}{1-u}.$$

[C] The negative binomial distribution. In a succession of Bernoulli trial (Binomial distribution) let $P(r)$ denote the probability that exactly $r+k$ ($k > 0$), trials are required to produce k successes. This will so happen when the last trial, i.e., $(r+k)$ th trial is a success with probability p and the previous $(r+k-1)$ trials must have $(k-1)$ successes with probability ${}^{r+k-1}C_{k-1} p^{k-1} q^r$, where $q = 1-p$.

$\therefore P(r) = \text{prob. of } (k-1) \text{ successes in } (x+k-1) \text{ trials}$
 $\times \text{probability of } (x+k) \text{th success}$

$$\begin{aligned} &= {}^{r+k-1}C_{k-1} p^{k-1} q^r \cdot p \\ &= {}^{r+k-1}C_{k-1} p^k q^r \end{aligned} \quad \dots (1)$$

$$\begin{aligned} &= \frac{p^k (k+r-1) (k+r-2) \dots \{k+r+1-(r+1)\}}{r!} q^r \\ &= \frac{p^k (k+r-1) (k+r-2) \dots (k+1) \cdot k}{r!} q^r \\ &= p^k (-1)^r \frac{(-k) (-k-1) \dots (-k-r+1)}{r!} q^r \end{aligned}$$

$$\begin{aligned} &= p^k (-1)^r \cdot {}^rC_r q^r \\ &= {}^rC_r p^k (-q)^r, \end{aligned} \quad \dots (2)$$

$$\begin{aligned} \text{whence } \sum_{r=0}^{\infty} P(r) &= p^k \sum_{r=0}^{\infty} {}^rC_r (-q)^r \\ &= p^k [1-q]^{-k} = p^k p^{-k} = 1. \end{aligned} \quad \dots (3)$$

The distributions given by (1) and (2) for $k \geq 0$ even if k is not an integer are known as negative binomial distribution.

Deductions

(1) Pascal's distribution. The distribution

$$P(r) = {}^rC_r p^k (-q)^r \quad \dots (4)$$

when regarded as one having two parameters p and k is said to be the Pascal-distribution.

(2) Geometric distribution. In (1) if we put $k=1$, this becomes

$$\begin{aligned} P(r) &= {}^rC_0 p q^r \\ &= q^r p, r=0, 1, 2, \dots \text{ and } q=1-p. \end{aligned} \quad \dots (5)$$

This is known as geometrical distribution, which can also be written directly by considering that r is the number of failures preceding the first success in a Bernoulli-sequence of trials, probability of success being p , so that r may be regarded as a random variate with probability distribution

$$P(r) = q^r \cdot p, r=0, 1, 2, \dots \text{ and } q=1-p,$$

(3) Polya's distribution. If we put $k = \frac{1}{\beta}$, $p = \frac{1}{1+\beta\mu}$, $q = \frac{\beta\mu}{1+\beta\mu}$ in (1), we get

$$P(r) = \frac{(1+\beta)(1+2\beta)\dots[1+(r-1)\beta]}{r!} \left(\frac{1}{1+\beta\mu} \right)^{1/\beta} \left(\frac{\mu}{1+\beta\mu} \right)^r, \dots (1)$$

which is known as Polya's distribution with two parameters β and μ .

(4) **Second form of geometric distribution.** In Polya's distribution given by (6), if we put $\beta=1$, we get

$$P(r) = \frac{1}{1+\mu} \left(\frac{\mu}{1+\mu} \right)^r. \dots (7)$$

(5) **General term in the binomial expansion of $(Q-P)^{-k}$.** Putting $p = \frac{1}{Q}$, $q = \frac{P}{Q}$, so that $Q-P=1$ as $p+q=1$, (1) gives

$$\begin{aligned} P(r) &= Q^{-k} \cdot r+1-k-1 C_{k-1} \left(\frac{P}{Q} \right)^r \\ &= Q^{-k} \cdot {}^{-k}C_r \cdot \left(\frac{-P}{Q} \right)^r, \quad r=0, 1, 2, \dots \text{ for the form (2), which} \\ &\text{gives the general term of } (Q-P)^{-k}. \end{aligned}$$

[D] **Hypergeometric distribution.** From an urn containing m white and n red balls if r balls are drawn at a time without replacement, then the probability that x out of r are white is given by

$$P(x) = \frac{{}^m C_x {}^n C_{r-x}}{{}^{m+n} C_r}, \quad n=0, 1, 2, \dots, r, \quad r \leq m, \quad r \leq n \quad \dots (1)$$

where $\sum_{x=0}^r P(x)=1$ since $\sum_{x=0}^r {}^m C_x {}^n C_{r-x} = {}^{m+n} C_r$

which is evident by equating the coefficients of x^r on either side of $(1+x)^m (1+x)^n = (1+x)^{m+n}$.

The distribution given by (1) is said to be the hypergeometric distribution.

Another form of hypergeometric distribution. From an urn containing Np white and Nq red balls such that $p+q=1$, if n balls containing x red and $(n-x)$ white are drawn, then their probability is given by

$$f(x) = \frac{{}^{Np} C_x {}^{Nq} C_{n-x}}{{}^N C_n} \quad \dots (2)$$

where $f(x)$ represents a probability density function known as hypergeometric function.

As above it is evident that $\sum_{x=0}^n f(x)=1$.

$$\begin{aligned} \text{Consider } \lim_{N \rightarrow \infty} \frac{{}^{Np} C_x {}^{Nq} C_{n-x}}{{}^N C_n} \\ &= \lim_{N \rightarrow \infty} \frac{(Np)!}{x! (Np-x)!} \cdot \frac{(Nq)!}{(n-x)! (Nq-n+x)!} \cdot \frac{n! (N-n)!}{N!} \\ &\quad \cdot \frac{\{Np(Np-1)\dots(Np-x+1)\} Nq(Nq-1)\dots}{\dots (Nq-n+x+1)} \\ &= \lim_{N \rightarrow \infty} \frac{n!}{x! (n-x)!} \cdot \frac{\dots (Nq-n+x+1)}{N(N-1)\dots(N-n+1)} \end{aligned}$$

$$\begin{aligned} & {}^n C_x \left\{ p \left(p - \frac{1}{N} \right) \left(p - \frac{2}{N} \right) \dots \left(p - \frac{x-1}{N} \right) \right\} \\ & \times \left\{ q \left(q - \frac{1}{N} \right) \left(q - \frac{2}{N} \right) \dots \left(q - \frac{n-x-1}{N} \right) \right\} \\ &= \lim_{N \rightarrow \infty} \frac{{}^n C_x p^x q^{n-x}}{\left(1 - \frac{1}{N} \right) \left(1 - \frac{2}{N} \right) \dots \left(1 - \frac{n-1}{N} \right)} \end{aligned}$$

which is binomial frequency distribution.

Problem 129. Find the mean and variance of hypergeometric distribution.

Case I. Using the form (1), we have

$$\begin{aligned} \text{Mean} = E(x) &= \sum_{x=0}^r x \cdot P(x) = \sum_{x=0}^r \frac{x \cdot {}^m C_x {}^n C_{r-x}}{{}^{m+n} C_r} \\ &= \frac{m \sum_{x=1}^r {}^{m-1} C_{x-1} {}^n C_{r-x}}{{}^{m+n} C_r} = \frac{m}{m+n} \cdot {}^{m-1} C_{r-1} {}^n C_{r-1} \quad (\text{if } x-1=y) \\ &= \frac{m}{m+n} \cdot {}^{m-1+n} C_{r-1} \\ &= \frac{m(r)! (m+n-r)!}{(m+n)!} \times \frac{(m-1+n)!}{(r-1)! (m+n-r)!} = \frac{mr}{m+n} \end{aligned}$$

$$\begin{aligned} \mu_2' = E(x^2) &= \sum_{x=0}^r x^2 P(x) = \sum_{x=0}^r x(x-1) P(x) + \sum_{x=0}^r x P(x) \\ &\quad \because x^2 = \{x(x-1) + x\} \end{aligned}$$

$$= \frac{m(m-1) \cdot r \cdot (r-1)}{(m+n)(m+n-1)} + \frac{mr}{m+n} \quad (\text{solving as above.})$$

$$\begin{aligned} \mu_2 = \mu_2' - \mu_1'^2 &= \frac{m(m-1)r(r-1)}{(m+n)(m+n-1)} + \frac{mr}{m+n} - \left(\frac{mr}{m+n} \right)^2 \\ \text{variance} &= \frac{mnr(m+n-r)}{(m+n)^2(m+n-1)} \quad (\text{simplifying}) \end{aligned}$$

Case II. Using the form (2), we have

$$\begin{aligned} \text{Mean} = \mu_1' &= \sum_{x=0}^n \left\{ x \cdot \frac{{}^{Np} C_x {}^{Nq} C_{n-x}}{{}^N C_n} \right\} = \sum_{x=1}^n \left\{ \frac{x \cdot {}^{Np} C_x {}^{Nq} C_{n-x}}{{}^N C_n} \right\} \\ &= Np \sum_{x=1}^n \frac{\{ {}^{Np-1} C_{x-1} {}^{Nq} C_{n-x} \}}{{}^N C_n} \\ &= Np \cdot \frac{{}^{Np+Nq-1} C_{n-1}}{{}^N C_n} = np \quad (\text{simplifying}) \\ \mu_2' &= \sum_{x=0}^n x^2 f(x) = \sum_{x=0}^n \{x(x-1) + x\} f(x) \\ &= \frac{np(n-1)(Np-1)}{N-1} \quad (\text{on simplification}) \end{aligned}$$

Normal equations given by $\frac{\partial S}{\partial x}=0$, $\frac{\partial S}{\partial y}=0$ and $\frac{\partial S}{\partial z}=0$ are on simplification

$$27x+6y=88, 6x+15y+z=70, -y+54z=107$$

which give on solving, $x=2.47$, $y=3.55$ and $z=1.92$.

These are the most plausible values

Note 2. Here it may be shown that $\frac{\partial^2 S}{\partial x^2}$, $\frac{\partial^2 S}{\partial y^2}$, $\frac{\partial^2 S}{\partial z^2}$ all are positive for these values of x, y, z .

Curve-fitting by method of least squares

The principle of least squares enables us to fit a polynomial of any degree i.e., to get a close functional relation between the variables x and y . Besides a polynomial, this relationship may be any of algebraic, exponential or logarithmic.

Assuming that we are given n paired observations $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$, we have to fit a polynomial of degree p such as

$$y=a_0+a_1x+a_2x^2+\dots+a_px^p \quad \dots (5)$$

where constants a 's can be determined by the method of least squares.

Taking (5) to be true, the estimated value of y for (x_1, y_1) is $a_0+a_1x_1+a_2x_1^2+\dots+a_px_1^p$ whereas the observed value is y_1 , so that the error of estimation say E_1 is

$$E_1=(y_1-a_0-a_1x_1-a_2x_1^2-\dots-a_px_1^p)$$

Similarly if $E_2, E_3, \dots, E_n$ be the errors of estimation for the other paired observations, then we have

$$E_2=y_2-a_0-a_1x_2-a_2x_2^2-\dots-a_px_2^p$$

$$\dots\dots\dots$$

$$E_n=y_n-a_0-a_1x_n-a_2x_n^2-\dots-a_px_n^p$$

$\therefore S$, the sum of squares of the errors of estimation is

$$S=E_1^2+E_2^2+\dots+E_n^2=\sum_{\alpha=1}^n E_{\alpha}^2$$

$$=\Sigma(y-a_0-a_1x-a_2x^2-\dots-a_px^p)$$

The normal equations given by

$$\frac{\partial S}{\partial a_0}=0, \frac{\partial S}{\partial a_1}=0, \frac{\partial S}{\partial a_2}=0, \dots, \frac{\partial S}{\partial a_n}=0 \text{ are on simplification}$$

$$\left. \begin{aligned} \Sigma y &= na_0 + a_1 \Sigma x + a_2 \Sigma x^2 + \dots + a_p \Sigma x^p \\ \Sigma xy &= a_0 \Sigma x + a_1 \Sigma x^2 + a_2 \Sigma x^3 + \dots + a_p \Sigma x^{p+1} \\ \Sigma x^2 y &= a_0 \Sigma x^2 + a_1 \Sigma x^3 + a_2 \Sigma x^4 + \dots + a_p \Sigma x^{p+2} \\ &\dots\dots\dots \\ \Sigma x^p y &= a_0 \Sigma x^p + a_1 \Sigma x^{p+1} + a_2 \Sigma x^{p+2} + \dots + a_p \Sigma x^{2p} \end{aligned} \right\} \quad \dots (6)$$

These $(p+1)$ normal equations can uniquely determine the $(p+1)$ constants $a_0, a_1, a_2, \dots, a_p$.

COROLLARY 1. In case the polynomial (5) is of degree 1 i.e., it is a straight line, the normal equations are (on putting $p=1$)

$$\left. \begin{aligned} \Sigma y &= na_0 + a_1 \Sigma x \\ \Sigma xy &= a_0 \Sigma x + a_1 \Sigma x^2 \end{aligned} \right\} \quad \dots (7)$$

COROLLARY 2. In case the polynomial (5) is of degree 2 i.e. it is a parabola the normal equations (on putting $p=2$) are

$$\left. \begin{aligned} \Sigma y &= na_0 + a_1 \Sigma x + a_2 \Sigma x^2 \\ \Sigma xy &= a_0 \Sigma x + a_1 \Sigma x^2 + a_2 \Sigma x^3 \\ \Sigma x^2 y &= a_0 \Sigma x^2 + a_1 \Sigma x^3 + a_2 \Sigma x^4 \end{aligned} \right\} \quad \dots (8)$$

Note 3. In order to ease the calculations a change of scale and origin may be suggested by putting $u = \frac{x-A}{h}$, $v = \frac{y-B}{k}$.

Problem 133. Fit a straight line to the following data regarding x as the independent variable

$x :$	0	1	2	3	4
$y :$	0	1.8	3.3	4.5	6.3

we have to fit a straight line say $y=a_0+a_1x$ $\dots (1)$

Normal equations are $\Sigma y=na_0+a_1 \Sigma x$, $\Sigma xy=a_0 \Sigma x+a_1 \Sigma x^2$ $\dots (2)$

As given, $x :$	0	1	2	3	4	$\therefore \Sigma x=10$
$y :$	0	1.8	3.3	4.5	6.3	$\therefore \Sigma y=16.9$
$xy :$	0	1.8	6.6	13.5	25.2	$\therefore \Sigma xy=47.1$
$x^2 :$	0	1	4	9	16	$\therefore \Sigma x^2=30$

Substituting these values in (2), we get

$$16.9=5a_0+10a_1; 47.1=10a_0+30a_1$$

which give on solving, $a_0=0.72$, $a_1=1.33$.

Their substitution in (1) yields the required straight line as $y=0.72+1.33x$.

Problem 134. Fit a second degree parabola to the following data:

$x :$	1	2	3	4	5
$y :$	1090	1220	1390	1625	1915

Using the change of scale and origin as $u=x-3$, $v=\frac{y-1450}{5}$, let the required parabola be $v=a_0+a_1u+a_2u^2$ $\dots (1)$

The normal equations are

$$\Sigma v=na_0+a_1 \Sigma u+a_2 \Sigma u^2, \Sigma uv=a_0 \Sigma u+a_1 \Sigma u^2+a_2 \Sigma u^3 \text{ and}$$

$$\Sigma u^2 v=a_0 \Sigma u^2+a_1 \Sigma u^3+a_2 \Sigma u^4 \quad \dots (2)$$

Now, $u :$	-2	-1	0	1	2	$\therefore \Sigma u=0$
$v :$	-72	-46	-12	35	93	$\therefore \Sigma v=-2$
$uv :$	144	46	0	35	186	$\therefore \Sigma uv=411$
$u^2 :$	4	1	0	1	4	$\therefore \Sigma u^2=10$
$u^2 v :$	-288	-46	0	35	372	$\therefore \Sigma u^2 v=73$
$u^3 :$	-8	-1	0	1	8	$\therefore \Sigma u^3=0$
$u^4 :$	16	1	0	1	16	$\therefore \Sigma u^4=34$

Substituting these values in (2), we get

$$-2 = 5a_0 + 10a_2; 411 = 10a_1; 73 = 10a_0 + 34a_2$$

which give on solving $a_0 = -11.4$, $a_1 = 41.1$, $a_2 = 5.5$.

Hence (1) gives $v = -11.4 + 41.1u + 5.5u^2$

$$\text{i.e., } \frac{y-1450}{5} = -11.4 + 41.1(x-3) + 5.5(x-3)^2$$

or $y = 1024 + 40.5x + 27.5x^2$

which is the required parabola.

Problem 135. Fit the curve $y = ae^{bx}$ to the following data, when $e = 2.71828$

$x :$	0	2	4
$y :$	5.012	10	31.62

Equation to the curve is $y = ae^{bx}$

Taking logarithms, $\log_{10} y = \log_{10} a + (b \log_{10} e) x$

say $y = A + Bx \quad \dots (1)$

Normal equations are

$$\Sigma Y = nA + B \Sigma x, \Sigma xY = A \Sigma x + B \Sigma x^2 \quad \dots (2)$$

Now, $x : 0 \quad 2 \quad 4$ So that $\Sigma x = 6$

$y : 5.012 \quad 10 \quad 31.62$

$Y = \log_{10} y : 0.7 \quad 1.0 \quad 1.5$ (by log table) $\Sigma y = 3.2$

$x^2 : 0 \quad 4 \quad 16$ $\Sigma x^2 = 20$

$xY : 0 \quad 2 \quad 6$ $\Sigma xY = 8$

$\therefore (2)$ give $3.2 = 3A + 6B$, $8 = 6A + 20B$

Solving $A = 0.666$, $B = 0.2$.

So that $a = (10)^A = (10)^{0.666} = 4.642$ and $b = \frac{B}{\log_{10} e} = 0.46$ approx. (by log table)

Hence the required equation (1) yields

$$y = 4.642 e^{0.46x}$$

15.19. CORRELATION AND REGRESSION

The two variables x and y are said to be correlated when they are so related or linked that a change in one is accompanied by a change in other such that an increase in one brings an increase or decrease in the other. In case an increase in one is accompanied by an increase in the other, then the two variables are said to be *positively correlated* e.g. volume and temperature of a gas. But if an increase in one is linked with the decrease in other, then the two variables are said to be *negatively correlated* e.g. pressure and volume of a gas.

In 1890, Karl Pearson defined the 'Product moment correlation coefficient' or simply 'coefficient of correlation' between the two variables x and y such that

$$r_{xy} = \frac{E(x-\bar{x})(y-\bar{y})}{\sqrt{E(x-\bar{x})^2} \sqrt{E(y-\bar{y})^2}} = \frac{\text{Cov}(x, y)}{\sqrt{\text{Var}(x)} \sqrt{\text{Var}(y)}} = \frac{\mu_{11}}{\sigma_x \sigma_y} \dots (1)$$

since $E(x-\bar{x})^2$ gives us a measure for the variation in x and $E(y-\bar{y})^2$ gives that in y while $E(x-\bar{x})(y-\bar{y})$ gives us the measure for simultaneous variation in x and y .

Also $\text{cov}(x, y) = E(x-\bar{x})(y-\bar{y}) = \mu_{11}$ and σ_x and σ_y are the standard deviations for x and y respectively.

Properties of r , the Coefficient of Correlation

I. It is purely a number (being ratio) and hence has no unit of measurement.

II. It is independent of the origin and scale.

Suppose that we introduce two variables u and v such that

$$u = \frac{x-a}{h}, v = \frac{y-b}{k} \quad \dots (2)$$

where (a, b) is the new origin instead of $(0, 0)$.

(2) can be written as $x = a + hu$, $y = b + kv$ giving

$$\bar{x} = a + h\bar{u}, \bar{y} = b + k\bar{v}$$

where $\bar{u}$, $\bar{v}$ are respectively the means of the u and v variates.

Thus, $\text{Cov}(x, y) = E(x-\bar{x})(y-\bar{y})$ by §15.11

$$= E\{(a+hu)-(a+h\bar{u})\} \{(b+kv)-(b+k\bar{v})\}$$

$$= E(hu-h\bar{u})(kv-k\bar{v}) = hk E(u-\bar{u})(v-\bar{v})$$

$$\text{Var}(x) = E(x-\bar{x})^2 = E\{(a+hu)-(a+h\bar{u})\}^2$$

$$= E(hu-h\bar{u})^2 = h^2 E(u-\bar{u})^2 = h^2 \text{Var}(u)$$

$$\text{Similarly } \text{Var}(y) = k^2 E(v-\bar{v})^2 = k^2 \text{Var}(v)$$

Hence from (1),

$$r_{xy} = \frac{\text{Cov}(x, y)}{\sqrt{\text{Var}(x)} \sqrt{\text{Var}(y)}} = \frac{hk}{\sqrt{h^2 k^2}} \frac{E(u-\bar{u})(v-\bar{v})}{\sqrt{\text{Var}(u)} \sqrt{\text{Var}(v)}} \\ = \frac{hk}{\sqrt{h^2 k^2}} r_{uv}$$

In case h and k both are positive or both negative, we get

$$r_{xy} = r_{uv} \quad \dots (3)$$

showing that r_{uv} is independent of a, b (i.e. change of origin) and of h, k (i.e. change of scale).

But if h and k are of opposite sign, then we have $r_{xy} = -r_{uv}$.

III. When x and y are independent i.e. uncorrelated then $r_{xy} = 0$

$$\therefore r_{xy} = \frac{\text{Cov}(x, y)}{\sigma_x \sigma_y} = \frac{E(x-\bar{x})(y-\bar{y})}{\sqrt{E(x-\bar{x})^2} \sqrt{E(y-\bar{y})^2}}$$

But x, y being independent, $E(x, y) = E(x) E(y)$ by §15.11.

$$\begin{aligned} \text{So that } E(x-\bar{x})(y-\bar{y}) &= E(x-\bar{x})E(y-\bar{y}) \\ &= \{E(x)-E(\bar{x})\}\{E(y)-E(\bar{y})\} \\ &= (\bar{x}-\bar{x})(\bar{y}-\bar{y})=0 \end{aligned}$$

$$\text{Hence } r_{xy}=0. \quad \dots (4)$$

Note. Its converse is not true i.e. if two variates are uncorrelated, it is not necessary that they are independent too, e.g. if x is a variate with constant density function say

$f(x)=\frac{1}{2}, -1 \leq x \leq 1$ and $y=x^2$, then we have

$$E(x) = \int_{-1}^1 x \cdot f(x) dx = \frac{1}{4} \int_{-1}^1 x dx = \frac{1}{4} \left[\frac{x^2}{2} \right]_{-1}^1 = 0$$

$$\therefore E(x) \cdot E(y) = 0 \cdot E(y) = 0.$$

$$\text{Also } E(xy) = E(x^3) = \int_{-1}^1 x^3 \cdot f(x) dx = \frac{1}{4} \int_{-1}^1 x^3 dx = \frac{1}{4} \left[\frac{x^4}{4} \right]_{-1}^1 = 0$$

So that $\text{Cov}(x, y) = E(x-\bar{x})(y-\bar{y}) = E(xy) - E(x)E(y) = 0 - 0 = 0$ showing that x and y are uncorrelated but y and x are not independent as is obvious by the relation $y=x^2$.

IV. Limits for the coefficient of correlation are $-1 \leq r \leq 1$.

Let us assume that,

$$X_i = \frac{x_i - \bar{x}}{\sigma_x} \text{ so that } \sum_{i=1}^n X_i^2 = \frac{1}{\sigma_x^2} \sum_{i=1}^n (x_i - \bar{x})^2 = \frac{n\sigma_x^2}{\sigma_x^2} = n \quad \dots (5)$$

$$\text{and } Y_i = \frac{y_i - \bar{y}}{\sigma_y} \text{ so that } \sum_{i=1}^n Y_i^2 = \frac{1}{\sigma_y^2} \sum_{i=1}^n (y_i - \bar{y})^2 = \frac{n\sigma_y^2}{\sigma_y^2} = n \quad \dots (6)$$

$$\begin{aligned} \therefore r &= \frac{E(x-\bar{x})(y-\bar{y})}{\sigma_x \sigma_y} = \frac{1}{n\sigma_x \sigma_y} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) \\ &= \frac{1}{n} \sum_{i=1}^n X_i Y_i \quad \dots (7) \end{aligned}$$

Now, since a perfect squared quantity is never negative, therefore

$$\sum_{i=1}^n (X_i + Y_i)^2 \geq 0 \text{ i.e. } \frac{1}{n} \sum_{i=1}^n (X_i + Y_i)^2 \geq 0$$

$$\text{or } \frac{1}{n} \sum_{i=1}^n X_i^2 + \frac{1}{n} \sum_{i=1}^n Y_i^2 + \frac{2}{n} \sum_{i=1}^n X_i Y_i \geq 0$$

$$\text{or } 1 + 1 + 2r \geq 0 \text{ by (5), (6) and (7)}$$

$$\text{giving } 2r \geq -2 \text{ i.e. } r \geq -1 \quad \dots (8)$$

$$\text{Again, } \sum_{i=1}^n (X_i - Y_i)^2 \geq 0 \text{ and hence } \frac{1}{n} \sum_{i=1}^n (X_i - Y_i)^2 \geq 0$$

$$\therefore \frac{1}{n} \sum_{i=1}^n X_i^2 + \frac{1}{n} \sum_{i=1}^n Y_i^2 - \frac{2}{n} \sum_{i=1}^n X_i Y_i \geq 0$$

$$\text{i.e. } 1 + 1 - 2r \geq 0 \text{ by (5), (6) and (7)}$$

$$\text{giving } 2 - 2r \geq 0 \text{ or } 2r \leq 2 \text{ i.e. } r \leq 1 \quad \dots (9)$$

$$\text{Combining (8) and (9), } -1 \leq r \leq 1 \quad \dots (10)$$

Various other Forms of r

If α and β be the deviations of the two variates x and y from their respective means and σ_x, σ_y be the standard deviations of these series, then we define

$$r = -\frac{\sum \alpha \beta}{n \sigma_x \sigma_y} = \frac{p}{\sigma_x \sigma_y} \text{ where } p = \frac{\sum \alpha \beta}{n} \quad \dots (11)$$

Let M_x, M_y be the true means; A_x, A_y be the assumed means for the two series and ξ, η be the deviations from the assumed means i.e.

$$\alpha = x - M_x, \beta = y - M_y$$

$$\xi = x - A_x, \eta = y - A_y$$

Also let $d_x = M_x - A_x$ and $d_y = M_y - A_y$.

Then, $\xi = x - A_x = x - M_x + M_x - A_x = \alpha + d_x$ and similarly $\eta = \beta + d_y$

$\therefore \sum \xi = \sum \alpha + \sum d_x = n d_x \therefore \sum \xi = 0$ by Problem I of Arithmetic-mean

$$\text{giving } d_x = \frac{\sum \xi}{n} \text{ and similarly } d_y = \frac{\sum \eta}{n}$$

$$\begin{aligned} \text{Now, } \sum \xi \eta &= \sum (\alpha + d_x)(\beta + d_y) = \sum \alpha \beta + d_y \sum \alpha + d_x \sum \beta + \sum d_x d_y \\ &= \sum \alpha \beta + \eta d_x d_y \therefore \sum \alpha = 0 = \sum \beta \end{aligned}$$

$$\therefore \sum \alpha \beta = \sum \xi \eta - n d_x d_y = \sum \xi \eta - \frac{1}{n} \sum \xi \sum \eta$$

$$\text{and } \sigma_x = \sqrt{\frac{\sum \xi^2}{n} - \left(\frac{\sum \xi}{n}\right)^2} \text{ and } \sigma_y = \sqrt{\frac{\sum \eta^2}{n} - \left(\frac{\sum \eta}{n}\right)^2} \text{ by §15.9 (D)}$$

With these substitutions (11) yields

$$r = \frac{\sum \xi \eta - \frac{1}{n} \sum \xi \sum \eta}{\sqrt{\sum \xi^2 - \frac{1}{n} \left(\sum \xi\right)^2} \sqrt{\sum \eta^2 - \frac{1}{n} \left(\sum \eta\right)^2}} \quad \dots (12)$$

Further if we use the change of scale and origin such that

$$u = \frac{x - A_x}{h} = \frac{\xi}{h} \text{ and } v = \frac{y - A_y}{k} = \frac{\eta}{k}, \text{ then (12) reduces to}$$

$$r = \frac{\sum uv - \frac{1}{n} \sum u \sum v}{\sqrt{\sum u^2 - \frac{1}{n} \left(\sum u\right)^2} \sqrt{\sum v^2 - \frac{1}{n} \left(\sum v\right)^2}} \quad \dots (13)$$

which for a bivariate frequency distribution yields

$$r = \frac{\sum f uv - \frac{\sum u \sum v}{\sum f}}{\sqrt{\sum f u^2 - \frac{1}{\sum f} (\sum f u)^2} \sqrt{\sum f v^2 - \frac{1}{\sum f} (\sum f v)^2}} \quad \dots (14)$$

where f is the frequency of a particular rectangle in the correlation table, whose mention is not needed in the present volume.

Problem 136. x and y are two variates with variances σ_x^2 and σ_y^2 respectively and r is the coefficient of correlation between them. If

$$u = x + ky, \quad v = x + \frac{\sigma_y}{\sigma_x} y,$$

find the value of k so that u and v may be uncorrelated.

$$\therefore u \text{ and } v \text{ are uncorrelated, } \text{Cov}(u, v) = 0.$$

$$\therefore 0 = \text{Cov}(u, v) = E(u - \bar{u})(v - \bar{v})$$

$$= E(x + ky - \bar{x} + k\bar{y}) \left(x + \frac{\sigma_y}{\sigma_x} y - \bar{x} + \frac{\sigma_y}{\sigma_x} \bar{y} \right)$$

$$= E\{(x - \bar{x}) + k(y - \bar{y})\} \left\{ (x - \bar{x}) + \frac{\sigma_y}{\sigma_x} (y - \bar{y}) \right\}$$

$$= E(x - \bar{x})^2 + kE(x - \bar{x})(y - \bar{y})$$

$$+ \frac{\sigma_y}{\sigma_x} E(x - \bar{x})(y - \bar{y}) + k \frac{\sigma_y}{\sigma_x} E(y - \bar{y})^2$$

$$\text{i.e., } \sigma_x^2 + \left(k + \frac{\sigma_y}{\sigma_x} \right) r \sigma_x \sigma_y + k \frac{\sigma_y}{\sigma_x} \sigma_y^2 = 0$$

$$\text{or } k(-\sigma_x \sigma_y - r \sigma_x \sigma_y) = \sigma_x^2 + r \sigma_x^2$$

$$\text{or } -k \sigma_x \sigma_y (1 + r) = \sigma_x^2 (1 + r) \text{ giving } k = -\frac{\sigma_x}{\sigma_y}.$$

Problem 137. Show that the coefficient of correlation r between two variables x and y is given by

$$r = (\sigma_x^2 + \sigma_y^2 - \sigma_{x-y}^2) / 2\sigma_x \sigma_y,$$

where σ_x^2 , σ_y^2 and σ_{x-y}^2 are the variances of x , y and $x - y$ respectively.

We have $\sigma_{x-y}^2 = \sigma_x^2 + \sigma_y^2 - 2 \text{Cov}(x, y)$.

$$\therefore \text{Cov}(x, y) = \{\sigma_x^2 + \sigma_y^2 - \sigma_{x-y}^2\} / 2.$$

$$\therefore r = \frac{\text{Cov}(x, y)}{\sigma_x \sigma_y} = \frac{(\sigma_x^2 + \sigma_y^2 - \sigma_{x-y}^2)}{2\sigma_x \sigma_y}.$$

Probable Error of r

According to Secrist, 'the probable error of the correlation coefficient is an amount which if added to and subtracted from the mean correlation coefficient, produces amounts within which the chances are even that a coefficient of correlation from a series selected at random will fall'

Since in a normal distribution $m \pm .6745 \sigma$ covers 50 per cent of the total area, therefore Probable Error (P.E.) is 0.6745 times the Standard Error (S.E.) defined as

$$\text{S.E. of } r = \frac{1-r^2}{\sqrt{N}}, \quad (N \text{ being total number of observation}) \quad \dots (15)$$

$$\therefore \text{P.E. of } r = 0.6745 (\text{S.E.}) = 0.6745 \frac{1-r^2}{\sqrt{N}} \quad \dots (16)$$

The significance of correlation is decided as follows:

$$\left. \begin{array}{l} \text{If } r < \text{P.E., there is no correlation} \\ \text{But if } r > \text{P.E., there is correlation} \end{array} \right\} \quad \dots (17)$$

Actually the two limits within which the coefficient of correlation always lies, are

$$r \pm \text{P.E.} \quad \dots (18)$$

Rank Correlation

Suppose a group of n individuals is ranked according to two characters or attributes A and B (say) such as $x_1, x_2, \dots, x_n$ and $y_1, y_2, \dots, y_n$ respectively where n ranks for each variable ranges from 1 to n . Then using the notations introduced in this section,

$$\text{we have, } M_x = M_y = \frac{1+2+\dots+n}{n} = \frac{n(n+1)}{2n} = \frac{n+1}{2}$$

$$\text{So that } \sigma_x = \sqrt{\frac{1}{n} \sum (x - M_x)^2} = \sqrt{\frac{1}{n} \sum \left(x - \frac{n+1}{2} \right)^2}$$

$$= \sqrt{\frac{1}{n} \left\{ \sum x^2 - (n+1) \sum x + \sum \left(\frac{n+1}{2} \right)^2 \right\}}$$

$$\text{Here } \sum x^2 = 1^2 + 2^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\text{and } \sum x = 1+2+\dots+n = \frac{n}{2} (n+1)$$

$$\therefore \sigma_x = \sqrt{\frac{1}{n} \left\{ \frac{n(n+1)(2n+1)}{6} - \frac{n}{2} (n+1)^2 + n \left(\frac{n+1}{2} \right)^2 \right\}} = \sqrt{\frac{n^2-1}{12}}.$$

$$\text{Similarly } \sigma_y = \sqrt{\frac{n^2-1}{12}}$$

Let $d = \alpha - \beta$, then

$$\sum d^2 = \sum (\alpha - \beta)^2 = \sum \alpha^2 + \sum \beta^2 - 2 \sum \alpha \beta$$

$$= \sum (x - M_x)^2 + \sum (y - M_y)^2 - 2 \sum \alpha \beta$$

$$= \frac{n(n^2-1)}{12} + \frac{n(n^2-1)}{12} - 2 \sum \alpha \beta \text{ solving as above}$$

$$\text{giving } \sum \alpha \beta = \frac{1}{12} n(n^2-1) - \frac{1}{2} \sum d^2.$$

Hence by (11), we have

$$r = 1 - \frac{6 \sum d^2}{n(n^2-1)} \quad \dots (19)$$

$$\text{where } d = \alpha - \beta = \left(x - \frac{n+1}{2} \right) - \left(y - \frac{n+1}{2} \right) = x - y \quad \dots (20)$$

Problem 138. Ten students got the following percentage of marks in Principles of Economics and statistics

Students:	1	2	3	4	5	6	7	8	9	10
Marks in Economics:	78	36	98	25	75	82	90	62	65	39
Marks in Statistics:	84	51	91	60	68	62	86	58	53	47

Calculate rank correlation coefficient.

Denoting by x and y the ranks in Economics and Statistics respectively, we have

Students:	1	2	3	4	5	6	7	8	9	10	Total
x :	4	9	1	10	5	3	2	7	6	8	
y :	3	9	1	6	4	5	2	7	8	10	
$ x-y = d $:	1	0	0	4	1	2	0	0	2	2	
d^2 :	1	0	0	16	1	4	0	0	4	4	30

Putting $\sum d^2=30$ and $n=10$ in (19), we find

$$r=1-\frac{6 \times 30}{10(100-1)}=1-\frac{2}{11}=\frac{9}{11}=.82 \text{ approx.}$$

Regression and Lines of Regression

Assuming that there exists an association or relationship between two variables x and y for which n paired values $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ are observed, if we plot these points on a graph paper with proper choice of scale then these points are found more or less concentrated round a curve termed as *curve of regression* and the relationship is said to be expressed by means of *curvilinear regression*. In case the curve is a straight line, then it is known as the *line of regression* and the regression is known to be linear. In fact the line of regression gives the best fit or best estimate in the least square sense to an assigned probability distribution.

Whenever such a straight line falls to the choice that sum of squares of deviations parallel to y -axis is minimum then it is called the *line of regression of y on x* which gives the best estimate of y for any assigned value of x . Similarly if the sum of squares of deviations parallel to x -axis is minimum, then the line is called the *line of regression of x on y* which gives the best estimate of x for given y .

Let the form of the line of best fit of y for given x be $y=ax+b$ and let $\bar{x}, \bar{y}$ be the means of two series.

Also let $R_1, R_2, \dots, R_n$ be the residuals for the points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with frequencies $f_1, f_2, \dots, f_n$ respectively, then we have $R_i=f_i(y_i-ax_i-b)$, so that

$$S=\sum_{i=1}^n f_i(y_i-ax_i-b).$$

Applying the principle of least squares, S will be a minimum

if $\frac{\partial S}{\partial a}=0, \frac{\partial S}{\partial b}=0$ i.e. if $\sum f_i x_i (y_i-ax_i-b)=0$;

$$\sum f_i (y_i-ax_i-b)=0$$

where $\sum f_i=N, \sum f_i x_i=N\bar{x}, \sum f_i y_i=N\bar{y}, \sum f_i x_i^2=N(\sigma_x^2+\bar{x}^2)$
 $\sum f_i y_i^2=N(\sigma_y^2+\bar{y}^2)$ and $\sum f_i x_i y_i=N(\mu_{11}+\bar{x}\bar{y})$ by § 15.10 (E).
 $\therefore \bar{x}\bar{y}+\mu_{11}=a(\bar{x}^2+\sigma_x^2)+b\bar{x}$ and $\bar{y}=a\bar{x}+b$.

Solving these give $a=\frac{\mu_{11}}{\sigma_x^2}, b=\bar{y}-\frac{\mu_{11}}{\sigma_x^2}\bar{x}$.

Putting these values in $y=a x+b$, the line of regression of y on x is

$$y-\bar{y}=\frac{\mu_{11}}{\sigma_x^2}(x-\bar{x}) \quad \dots (21)$$

Similarly the line of regression of x on y is

$$x-\bar{x}=\frac{\mu_{11}}{\sigma_y^2}(y-\bar{y}) \quad \dots (22)$$

Here the coefficients $\frac{\mu_{11}}{\sigma_x^2}$ and $\frac{\mu_{11}}{\sigma_y^2}$ are known as the *regression coefficients* of y on x and of x on y respectively and denoted by

$$b_{yx}=\frac{\mu_{11}}{\sigma_x^2}, b_{xy}=\frac{\mu_{11}}{\sigma_y^2} \quad \dots (23)$$

In view of (1), we have $\mu_{11}=r\sigma_x\sigma_y$

$$\therefore b_{yx}=r\frac{\sigma_y}{\sigma_x} \text{ and } b_{xy}=r\frac{\sigma_x}{\sigma_y} \quad \dots (24)$$

$$\text{Clearly } b_{yx} \cdot b_{xy}=r^2 \quad \dots (25)$$

which shows that the correlation-coefficient r is the geometric mean between the regression coefficients.

Using the change of scale and origin such that

$$u=\frac{x-a}{h} \text{ and } v=\frac{y-b}{k}, \text{ we may have}$$

$$b_{yx}=\frac{k}{h} b_{vu} \text{ and } b_{xy}=\frac{k}{h} b_{uv} \text{ whence } \sigma_x=h\sigma_u \text{ etc.} \quad \dots (26)$$

Problem 139. Show that θ , the acute angle between two lines of regression, is given by

$$\tan \theta=\frac{1-r^2}{r} \frac{\sigma_x\sigma_y}{\sigma_x^2+\sigma_y^2}.$$

Interpret the case when $r=0, r=\pm 1$.

$$\therefore \text{Regression line of } y \text{ on } x \text{ is } y-\bar{y}=r\frac{\sigma_y}{\sigma_x}(x-\bar{x})$$

$$\text{Its slope}=r\frac{\sigma_y}{\sigma_x}=m_1 \text{ (say)}$$

$$\text{And regression line of } x \text{ on } y \text{ is } x-\bar{x}=r\frac{\sigma_x}{\sigma_y}(y-\bar{y}).$$

$$\text{Its slope}=\frac{\sigma_y}{\sigma_x}=m_2 \text{ (say)}$$

$$\therefore \tan \theta = \pm \frac{m_1 \sim m_2}{1 + m_1 m_2} = \pm \frac{r \frac{\sigma_y}{\sigma_x} \sim \frac{\sigma_y}{r \sigma_x}}{1 + r \frac{\sigma_y}{\sigma_x} \cdot \frac{\sigma_y}{r \sigma_x}} = \pm \frac{r^2 \sigma_x \sigma_y \sim \sigma_x \sigma_y}{r^2 \sigma_x^2 + r \sigma_y^2}$$

$$\text{For } \theta, \text{ acute only, } \tan \theta = + \frac{\sigma_x \sigma_y (r^2 \sim 1)}{(\sigma_x^2 + \sigma_y^2) r} = \frac{r^2 \sim 1}{r} \frac{\sigma_x \sigma_y}{\sigma_x^2 + \sigma_y^2}$$

$$\text{or } \tan \theta = \frac{1 - r^2}{r} \frac{\sigma_x \sigma_y}{\sigma_x^2 + \sigma_y^2}$$

($\because r^2 \leq 1$, hence $1 - r^2$ is also positive).

Case I. When $r=0$, $\tan \theta = \infty$. $\therefore \theta = 90^\circ$, i.e., both the regression lines are mutually perpendicular. Thus estimated value of x (or y) will be the same for all values of y (or x).

Case II. When $r = \pm 1$, $\tan \theta = 0$. $\therefore \theta = 0$ or π , i.e., both the regression lines coincide and so there is perfect correlation (either positive when $r=1$ or negative when $r=-1$) between the variables involved.

Problem 140. Given the following data, find what will be (a) the height of a policeman whose weight is 200 lbs., (b) the weight of a policeman who is 5 ft. tall.

Average height = 68 inches, average weight 150 lbs.

Coefficient of correlation between height and weight = .6.

S.D. of heights = 2.5 inches.

S.D. of weights = 20 lbs.

Suppose variable x indicates height (in inches) and y indicates weight (in lbs.). Then we have

$$\bar{x} = 68 \text{ inches, } \bar{y} = 150 \text{ lbs., } \sigma_x = 2.5 \text{ inches, } \sigma_y = 20 \text{ lbs., } r_{xy} = .6.$$

Now, we want to estimate y for $x=5$ ft. or for 60 inches and x for $y=200$ lbs.

$$(i) \therefore \text{Regression line of } y \text{ on } x \text{ is } y - \bar{y} = r \frac{\sigma_y}{\sigma_x} (x - \bar{x})$$

$$\therefore y - 150 = .6 \times \frac{20}{2.5} (x - 68).$$

$$\text{or } y - 150 = 4.8 (x - 68). \quad \dots (1)$$

$$(ii) \therefore \text{Regression line of } x \text{ on } y \text{ is } x - \bar{x} = r \frac{\sigma_x}{\sigma_y} (y - \bar{y})$$

$$\therefore x - 68 = .6 \frac{2.5}{20} (y - 150)$$

$$\text{or } x - 68 = .75 (y - 150). \quad \dots (2)$$

(a) Since (1) is the regression line of y on x so we shall estimate y .

Putting $x=60$ in (1), we get $y=150+4.8(60-68)$

$$= 150 - 38.4 = 111.6 \text{ lbs.}$$

Thus policeman of height 5 feet will have 111.6 lbs. weight.

(b) Since (2) is the regression line of x on y so we shall estimate x .

Putting $y=200$ in (2), we get $x=68+.75(200-150)$

$$= 68 + .75(50) = 105.5 \text{ inches.}$$

15.20. SAMPLING DISTRIBUTIONS

A *Population* or *Universe* is defined as the collection or class or aggregate of objects or a set of results of an operation. It is *finite* if the number of objects is finite and it is *infinite* if the number of objects is infinite.

A *sample* is defined as a part of the population selected from it in order to study its properties. The sample is to be *random* when the selection is made at random. A unit of sample or a sampling unit included in the sample is known as *sample unit*. The difference between a value of the population and its estimate derived from sample is said to be a *sampling error*, which may arise due to human bias or due to some other reasons. Actually the sampling procedure is based on the theory of probability as every sampling unit has a definite probability of being included in the sample.

A *simple random sampling* is one in which every sampling unit has an equal chance of being included in the samples e.g. if N be the size of a population, then ${}^N C_n$ samples of size n are possible so that each sample has a chance $\frac{1}{{}^N C_n}$ of being selected. Also the probability of some r th unit being selected at r th draw is

$$\frac{N-1}{N} \cdot \frac{N-2}{N-1} \cdots \frac{N-(r-1)}{N-(r-1)+1} \cdot \frac{1}{N-(r-1)} = \frac{1}{N}.$$

Properties of random sampling

I. The sample mean is an unbiased estimate of population mean. Taking $X_1, X_2, \dots, X_n$ as the population values and $x_1, x_2, x_n, \dots$ as the sample values, we have

$$\text{Population mean } x = \frac{1}{n} \sum_{i=1}^n X_i \text{ and sample mean}$$

$$\bar{X} = \frac{1}{N} \sum_{i=1}^n x_i \quad \dots (1)$$

Let the population total $\sum_{i=1}^N X_i = X$ and the sample total

$$\sum_{i=1}^n x_i = x \quad \dots (2)$$

Then we have to prove that $E(\bar{x}) = \bar{X}$... (3)

$$\text{Now, } E(\bar{x}) = E\left\{ \frac{1}{n} (x_1 + x_2 + \dots + x_n) \right\} = \frac{1}{n} E(x_1 + x_2 + \dots + x_n)$$

$$= \frac{1}{n} \{E(x_1) + E(x_2) + \dots + E(x_n)\} \text{ by (4) of § 15.11 (C),} \quad \dots (4)$$

Since x_i can be either $X_1, X_2, \dots, X_n$ each with probability $\frac{1}{N}$,
therefore

$$E(x_i) = \frac{X_1}{N} + \frac{X_2}{N} + \dots + \frac{X_n}{N} = \sum_{i=1}^n \frac{X_i}{N} = \bar{X} \text{ by (1)}$$

But $\bar{X}$ being a term free from index i

$$\therefore E(x_1) = E(x_2) = \dots = E(x_n) = \bar{X}$$

Thus (4) gives, $E(\bar{x}) = \frac{1}{n} [\bar{X} + \bar{X} + \dots n \text{ terms}] = \frac{1}{n} \cdot n\bar{X} = \bar{X}$
which proves (3).

COROLLARY 1. we have $E(N\bar{x}) = NE(\bar{x}) = N\bar{X} = X$ by (1) and (2) ... (5)

This follows that $N\bar{x}$ is an unbiased estimate of X and we denote it by

$$\hat{X} = N\bar{x} \quad \dots (6)$$

II. The variance of the sample mean $\bar{x}$ from a simple random sample is

$$\text{Var}(\bar{x}) = E(\bar{x} - \bar{X})^2 = \frac{S^2}{n} \frac{N-n}{N} \quad \dots (7)$$

where
$$S^2 = \frac{1}{N-1} \sum_{i=1}^N (X_i - \bar{X})^2 \quad \dots (8)$$

We have $\text{Var}(\bar{x}) = E\{\bar{x} - E(\bar{x})\}^2 = (\bar{x} - \bar{X})^2$

$$\begin{aligned} &= E\left\{\frac{1}{n} (x_1 + x_2 + \dots + x_n) - \bar{X}\right\}^2 \\ &= \frac{1}{n^2} E\{(x_1 + x_2 + \dots + x_n) - n\bar{X}\}^2 \\ &= \frac{1}{n^2} [E(x_1 - \bar{X})^2 + E(x_2 - \bar{X})^2 + \dots + E(x_n - \bar{X})^2] \\ &\quad + 2 \sum_{i=1}^{n-1} \sum_{j=2}^n E(x_i - \bar{X})(x_j - \bar{X})] \quad \dots (9) \end{aligned}$$

But $E(x_i - \bar{X})^2 = \frac{1}{N} (X_1 - \bar{X})^2 + \frac{1}{N} (X_2 - \bar{X})^2 + \dots$
 $+ \frac{1}{N} (X_N - \bar{X})^2$ as in I.

$$= \frac{1}{N} \sum_{i=1}^N (X_i - \bar{X})^2 = \frac{1}{N} (N-1)S^2 \text{ by (8)}$$

and $E\{(x_i - \bar{X})(x_j - \bar{X})\}$
$$= \frac{1}{N} \cdot \frac{1}{N-1} \sum_{i=1}^{N-1} \sum_{j=2}^N (X_i - \bar{X})(X_j - \bar{X})$$

so that
$$\sum_{i=1}^{n-1} \sum_{j=2}^n E\{(x_i - \bar{X})(x_j - \bar{X})\}$$

$$= \frac{n(n-1)}{N(N-1)} \sum_{i=1}^{N-1} \sum_{j=2}^N (X_i - \bar{X})(X_j - \bar{X})$$

Hence (9) becomes

$$\begin{aligned} \text{Var}(\bar{x}) &= \frac{1}{n^2} \left\{ n \cdot \frac{N-1}{N} S^2 + \frac{2n(n-1)}{N(N-1)} \right. \\ &\quad \left. \sum_{i=1}^{N-1} \sum_{j=2}^N (X_i - \bar{X})(X_j - \bar{X}) \right\} \quad \dots (10) \end{aligned}$$

But $N\bar{X} = \sum_{i=1}^N X_i$ gives $(X_1 - \bar{X}) + (X_2 - \bar{X}) + \dots + (X_n - \bar{X}) = 0$

or $\{(X_1 - \bar{X}) + (X_2 - \bar{X}) + \dots + (X_n - \bar{X})\}^2 = 0$

or $\sum_{i=1}^N (X_i - \bar{X})^2 + 2 \sum_{i=1}^{N-1} \sum_{j=2}^N (X_i - \bar{X})(X_j - \bar{X}) = 0$

i.e. $2 \sum_{i=1}^{N-1} \sum_{j=2}^N (X_i - \bar{X})(X_j - \bar{X}) = - \sum_{i=1}^N (X_i - \bar{X})^2 = -(N-1)S^2$

With its substitution, (10) yields

$$\text{Var}(\bar{x}) = \frac{1}{n^2} \left[n \frac{N-1}{N} S^2 - \frac{n(n-1)}{N(N-1)} (N-1)S^2 \right] = \frac{N-n}{Nn} S^2$$

which proves the result (7).

COROLLARY 2. With the notation of Cor. 1, we have

$$\text{Var}(\hat{X}) = \text{Var}(N\bar{x}) = N^2 \text{Var}(\bar{x}) = \frac{N(N-n)}{n} S^2 \quad \dots (11)$$

COROLLARY 3. By (7), $\sigma_{\bar{x}}^2 = \frac{N-n}{N} \cdot \frac{S^2}{n}$

gives
$$\sigma_{\bar{x}} = \frac{S}{\sqrt{n}} \sqrt{\frac{N-n}{N}} \quad \dots (12)$$

which gives standard error of $\bar{x}$.

COROLLARY 4. When $N \rightarrow \infty$ i.e., the population becomes to be infinite, then

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{N-n}{N} &= \lim_{N \rightarrow \infty} \left(1 - \frac{n}{N}\right) = 1 \text{ and } \lim_{N \rightarrow \infty} \sqrt{\frac{N-n}{N}} \\ &= \lim_{N \rightarrow \infty} \sqrt{1 - \frac{n}{N}} = 1. \end{aligned}$$

$$\therefore \text{ by (7) or (12), } \text{Var}(\bar{x}) = \frac{S^2}{n} \text{ i.e., } \sigma_{\bar{x}} = \frac{S}{\sqrt{n}} \quad \dots (13)$$

COROLLARY 5. When $N \rightarrow \infty$, then $N-1 \approx N$.

$$\text{So that } \frac{1}{N-1} \sum_{i=1}^N (X_i - \bar{X})^2 \approx \frac{1}{N} \sum_{i=1}^N (X_i - \bar{X})^2 \text{ i.e., } S^2 \approx \sigma^2 \text{ or } S \approx \sigma$$

$$\text{Hence by (13), } \sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}} \quad \dots (14)$$

which gives the standard error of $\bar{x}$, the sample mean.

III. Taking s^2 and S^2 , as the sample variance and population variance respectively, s^2 is an unbiased estimate of S^2 i.e. $E(s^2) = S^2$.

$$\text{We have, } s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 \text{ and } S^2 = \frac{1}{N-1} \sum_{i=1}^N (X_i - \bar{X})^2$$

$$\therefore s^2 = \frac{1}{n-1} \sum_{i=1}^n \{(x_i - \bar{X}) - (\bar{x} - \bar{X})\}^2 = \frac{1}{n-1} \left\{ \sum_{i=1}^n (x_i - \bar{X})^2 + n(\bar{x} - \bar{X})^2 - 2(\bar{x} - \bar{X}) \sum_{i=1}^n (x_i - \bar{X}) \right\}$$

$$= \frac{1}{n-1} \left\{ \sum_{i=1}^n (x_i - \bar{X})^2 + n(\bar{x} - \bar{X})^2 - 2(\bar{x} - \bar{X}).n(\bar{x} - \bar{X}) \right\}$$

$$= \frac{1}{n-1} \left\{ \sum_{i=1}^n (x_i - \bar{X})^2 - n(\bar{x} - \bar{X})^2 \right\}$$

$$\text{so that } E(s^2) = \frac{1}{n-1} E \left\{ \sum_{i=1}^n (x_i - \bar{X})^2 - n(\bar{x} - \bar{X})^2 \right\}$$

$$= \frac{1}{n-1} \left\{ \sum_{i=1}^n E(x_i - \bar{X})^2 - n E(\bar{x} - \bar{X})^2 \right\}$$

$$= \frac{1}{n-1} \left\{ n \sum_{i=1}^n \frac{(X_i - \bar{X})^2}{N} - n \text{ var}(\bar{x}) \right\}$$

$$\therefore E(x_i - \bar{X})^2 = \sum_{i=1}^n \frac{1}{N} (X_i - \bar{X})^2$$

$$= \frac{1}{n-1} \left[\frac{n}{N} \cdot (N-1) S^2 - n \cdot \frac{N-n}{\sqrt{n}} S^2 \right] \text{ by II}$$

$\dots (15)$

COROLLARY 6. Unbiased estimate of the variance of $\bar{x}$ is denoted by

$$v(\bar{x}) \text{ and given as } v(\bar{x}) = \frac{N-n}{N} \cdot \frac{s^2}{n} \quad \dots (16)$$

$$\text{In limit } v(\bar{x}) = \frac{s^2}{n} \quad \dots (17)$$

Similarly unbiased estimate of the variance of $\hat{X}$ denoted by $v(\hat{X})$

$$\text{is given as } v(\hat{X}) = \frac{N(N-n)}{n} s^2 \quad \dots (18)$$

Difference between the means of two large samples

Let there be two random samples of size n_1 and n_2 respectively, taken from the same population of standard deviation σ . Assuming that the samples are independent, the standard error say ϵ of the difference of their means is given by

$$\epsilon^2 = \epsilon_1^2 + \epsilon_2^2 \quad \dots (19)$$

where ϵ_1, ϵ_2 are the standard errors of the means of two samples

$$\text{given by } \epsilon_1^2 = \frac{\sigma^2}{n_1} \text{ and } \epsilon_2^2 = \frac{\sigma^2}{n_2}$$

$$\text{Hence } \epsilon^2 = \sigma^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \quad \dots (20)$$

COROLLARY 7. In case the samples are drawn from two different populations with standard deviations σ_1 and σ_2 respectively, then

$$(20) \text{ yields } \epsilon^2 = \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2} \quad \dots (21)$$

COROLLARY 8. The values $\bar{x} \pm 1.96 \frac{\sigma}{\sqrt{n}}$ are known as 95 per cent *Fiducial limits* or *confidence limits* for the mean of the population from which the sample has been taken.

Problem 141. Assuming that N is large, show that the error in writing $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$ is approximately $\frac{50(n-1)}{N}$ per cent of the value of $\sigma_{\bar{x}}$.

$$\text{Reqd. error} = \left\{ \frac{\sigma}{\sqrt{n}} - \frac{\sigma}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}} \right\} \frac{\sigma}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}}$$

$$= \sqrt{\frac{N-1}{N-n}} - 1 = \left\{ 1 + \frac{n-1}{N-n} \right\}^{1/2} - 1$$

$$= 1 + \frac{1}{2} \cdot \frac{n-1}{N-n} - \frac{1.3}{2.4} \left(\frac{n-1}{N-n} \right)^2 + \dots - 1$$

$$= \frac{n-1}{2(N-n)} \text{ approx. } \frac{n-1}{2N}$$

$$\therefore \text{Percentage of error} = \frac{n-1}{2N} \cdot 100 = \frac{n-1}{N} \cdot 50 \text{ approx.}$$

15.21. THEORY OF ERRORS

We know that an *error* is a quantity which must be added to the true value in order to get the observed value in performing a physical experiment *i.e.*

$$\text{Observed value} = \text{True value} + \text{error} \quad \dots (1)$$

$$\text{or Observed value} - \text{error} = \text{True value}$$

which is generally expressed by saying that

$$\text{Observed value} + \text{correction} = \text{True value} \quad \dots (2)$$

It means that a *correction* is the error with reversed sign.

In testing a statistical hypothesis, there arise two types of possible errors: (1) by rejecting the hypothesis when we ought to accept it *i.e.* when it is true and (2) by accepting the hypothesis when we ought to reject it *i.e.* when it is false. On the other hand in order to understand what we mean by true value of a physical quantity we first estimate the true values from observations (measurements), then try to seek what certainty can be attributed to these estimates and finally compare estimates found from different sets of observations.

Since any number formed empirically by expressing it in units of the smallest possible unit say ϵ that can be measured by a physical apparatus, is an integer, therefore any real number lying in an interval of length ϵ may be termed as *true value of the quantity* *e.g.*, probable values or probabilities may be regarded as the true values of the relative frequencies. Since no measuring instrument is perfect and also there arise several other disturbing factors, every observation is liable to be a random variable as different observations are capable of yielding different results. We thus conclude that all such observations made, are full of errors and deviate from the true values to a certain degree of accuracy. We may classify errors into three types: (1) *Coarse or gross errors*, (2) *systematic errors* and (3) *random or statistical errors*. The *coarse errors* are mainly caused by the mis-handling of the apparatus due to carelessness of an observer. The *systematic errors* are caused in a certain direction due to one or more reasons governed by a definite rule. Consequently in a repeated set of observations made under constant conditions, the same systematic errors are bound to occur. Such errors can be rectified by knowing the governing rules of these errors. The *Random Errors* are the uncertainties not showing any irregularities and having equal chances for positive and negative values for such errors.

In order to find the best estimated value for an observed quantity, let us assume that t_1 and t_2 are two independent observations for an unknown physical quantity t and $\phi(t_1, t_2)$ is the best estimate for t . If t_1, t_2 are increased by an amount α , then it may be assumed that the estimate $\phi(t_1, t_2)$ is also increased by α *i.e.*

$$\phi(t_1 + \alpha, t_2 + \alpha) = \phi(t_1, t_2) + \alpha \quad \dots (3)$$

Also assuming that multiplication of t_1, t_2 by a constant β , results in multiplication of the estimate by β we have

$$\phi(\beta t_1, \beta t_2) = \beta \phi(t_1, t_2) \quad \dots (4)$$

Again the two observations being made under the same conditions, it is immaterial which observation yields t , and which one t_2 , we have

$$\phi(t_1, t_2) = \phi(t_2, t_1) \quad \dots (5)$$

Now keeping t_1, t_2 fixed and setting $\alpha = -t_2$, (3) gives

$$\phi(t_1 - t_2, 0) = \phi(t_1, t_2) - t_2 \text{ i.e. } \phi(t_1, t_2) = \phi(t_1 - t_2, 0) + t_2 \quad \dots (6)$$

with its substitution (4) yields

$$\phi(\beta t_1, \beta t_2) = \beta \phi(t_1 - t_2, 0) + \beta t_2 = \beta t_2 + \phi\{\beta(t_1 - t_2), 0\} \text{ by (4)} \quad \dots (7)$$

$$\text{or } \beta \phi(t_1, t_2) = \beta t_2 + \phi\{\beta(t_1 - t_2), 0\} \text{ by (4)}$$

If we now assume that $\beta = \frac{1}{t_1 - t_2}$ and $t_1 \neq t_2$, then (7) becomes

$$\frac{1}{t_1 - t_2} \phi(t_1, t_2) = \frac{t_2}{t_1 - t_2} + (1, 0) \quad \dots (8)$$

$$\text{giving } t_2 + (t_1 - t_2) \phi(1, 0) = \phi(t_1, t_2)$$

Proceeding similarly, we can find

$$t_1 + (t_2 - t_1) \phi(1, 0) = \phi(t_2, t_1) \quad \dots (9)$$

In view of (5), (8) and (9) yield

$$t_2 + (t_1 - t_2) \phi(1, 0) = t_1 + (t_2 - t_1) \phi(1, 0) \text{ giving } \phi(1, 0) = \frac{1}{2} \quad \dots (10)$$

$\therefore$ (8) gives $t_2 + \frac{1}{2}(t_1 - t_2) = \phi(t_1, t_2)$ *i.e.* $\phi(t_1, t_2) = \frac{1}{2}(t_1 + t_2) \dots (11)$

Had we taken $t_1 = t_2$, (6) would have given

$$\phi(t_1, t_1) = t_1 + \phi(0, 0) \quad \dots (12)$$

with the choice $\beta = 0$, (8) would give $\phi(0, 0) = 0$

So that in view of (12), (11) gives $\phi(t_1, t_1) = t_1 = \frac{1}{2}(t_1 + t_1) = \frac{1}{2}(t_1 + t_1)$ which is the same result as (10).

Now we know that the best estimate of the true value say v made by n observations $t_1, t_2, \dots, t_n$ is defined as

$$m \approx v = \bar{t} = \frac{t_1 + t_2 + \dots + t_n}{n} = \frac{[t]}{n} \quad \dots (13)$$

where m is the expected value of t and $[t] = \sum_{i=1}^n t_i$.

Now we define, true error as $x_i = t_i - v$, $i = 1, 2, \dots, n$ and best errors as $\epsilon_i = t_i - t$, $i = 1, 2, \dots, n$

Further to calculate the probability of best estimate to lie within an assigned range of true value take $f(x_i)$ as the probability density of the random variable x_i , $i = 1, 2, \dots, n$. By the theorem of compound probability, we have

$$f(x_1, x_2) = f(x_1) f(x_2) \text{ and } f(x_1, x_2, x_3) = f(x_1) \cdot f(x_2) \cdot f(x_3) \quad \dots (14)$$

where x_1, x_2, x_3 etc. are errors made in first, second and third experiments.

Thus true value of observed quantity i.e. $v = \frac{1}{3} (t_1 + t_2 + t_3) \dots (17)$ in case of three observations only.

As such $f(x_1, x_2, x_3) = f(x_1) \cdot f(x_2) \cdot f(x_3)$
 $= f(t_1 - v) \cdot f(t_2 - v) \cdot f(t_3 - v)$ is maximum and so its logarithm will also be maximum i.e.

$\log f(t_1 - v) + \log f(t_2 - v) + \log f(t_3 - v)$ is maximum.

For maxima, its differential w.r.t. v must be equated to zero i.e.,

$$\frac{f'(t_1 - v)}{f(t_1 - v)} + \frac{f'(t_2 - v)}{f(t_2 - v)} + \frac{f'(t_3 - v)}{f(t_3 - v)} = 0 \quad \dots (18)$$

Setting $F(x) = \frac{f'(x)}{f(x)}$, we have $F(x_1) = F(t_1 - v) = \frac{f'(t_1 - v)}{f(t_1 - v)}$

and similarly $F(x_2) = \frac{f'(t_2 - v)}{f(t_2 - v)}$, $F(x_3) = \frac{f'(t_3 - v)}{f(t_3 - v)}$

$$\therefore (18) \text{ gives } F(x_1) + F(x_2) + F(x_3) = 0 \quad \dots (19)$$

which is true when (17) is true i.e. $3v = t_1 + t_2 + t_3$

$$\text{or } (t_1 - v) + (t_2 - v) + (t_3 - v) = 0 \text{ or } x_1 + x_2 + x_3 = 0 \quad \dots (20)$$

In case of two variables x_1 and x_2 ; (19) and (20) reduce to

$$F(x_1) + F(x_2) = 0 \text{ provided } x_1 + x_2 = 0 \quad \dots (21)$$

In case of a single variable x_1 ; this condition reduces to

$$F(x_1) = 0 \text{ when } x_1 = 0 \quad \dots (22)$$

Setting $x_1 = x_3$ and $x_2 = -x_3$ in (21), we find

$$F(x_2) = -F(-x_3) \quad \dots (23)$$

So that (19) yields,

$$F(x_1) + F(x_2) = -F(x_3) = F(-x_3) \text{ by (4)} \\ = F(x_1 + x_2) \text{ by (20)} \quad \dots (24)$$

Partial Differentiation of (24) w.r.t. x_1 and x_2 in succession gives

$$F'(x_1) = F'(x_1 + x_2) \text{ and } F'(x_2) = F'(x_1 + x_2)$$

giving $F'(x_1) = F'(x_2)$

Treating x_2 as constant, $F'(x_1) = F'(x_1 + x_2)$ will only hold if

$$F'(x_1) = \text{constant} = k' (\text{say}), \text{ so that}$$

$$F(x_1) = k' x_1 = \frac{f'(x_1)}{f(x_1)} \text{ implying } \frac{f'(x)}{f(x)} = F(x) = kx$$

which gives an integration $f(x) = A e^{\frac{1}{2} k x^2}$, A being constant of integration.

Now $f(x)$ being probability density function, we have by §15.12 (A),

$$\int_{-\infty}^{\infty} f(x) dx = 1 = A \int_{-\infty}^{\infty} e^{\frac{1}{2} k x^2} dx$$

Setting $k = -2h^2$, we have $1 = A \int_{-\infty}^{\infty} e^{-h^2 x^2} dx$ Put $h^2 x^2 = \frac{p^2}{2}$
i.e. $dx = \frac{1}{\sqrt{2}h} dp$

$$\text{or } 1 = \frac{A}{h\sqrt{2}} \int_{-\infty}^{\infty} e^{-p^2/2} dp \\ = \frac{A}{h\sqrt{2}} \sqrt{2\pi} \text{ i.e. } A = \frac{h}{\sqrt{\pi}} \quad \dots (25)$$

$$\text{Hence } f(x) = \frac{h}{\sqrt{\pi}} e^{-h^2 x^2}$$

which is called *Gaussian law of error*, and asserts that the probability for an error x to be $p_1 < \sqrt{2}hx < p_2$ is given by

$$\int_{p_1/\sqrt{2}h}^{p_2/\sqrt{2}h} \frac{h}{\sqrt{\pi}} e^{-h^2 x^2} dx \quad \dots (26)$$

$$\text{with } p = \sqrt{2}hx, \text{ this yields } \frac{1}{\sqrt{2\pi}} \int_{p_1}^{p_2} e^{-p^2/2} dp = \phi(p_2) - \phi(p_1) \quad \dots (27)$$

$$\text{where } \phi(p) = \frac{1}{\sqrt{2\pi}} \int_0^p e^{-p^2/2} dp.$$

Here h is called *precision constant* and it measures the accuracy of the observer. An error with probability $\frac{1}{2}$ is known as the *probable error* and may be determined by the value of x given by

$$\frac{h}{\sqrt{\pi}} e^{-h^2 x^2} = \frac{1}{2} \text{ i.e. } x = \frac{0.4769}{h} \quad \dots (28)$$

This renders the most probable error.

We define the *absolute error* $E(|x|)$ as

$$E(|x|) = \int_{-\infty}^{\infty} (x) f(x) dx = \frac{2h}{\sqrt{\pi}} \int_0^{\infty} x e^{-h^2 x^2} dx = \frac{1}{h\sqrt{\pi}} \\ = \frac{0.5642}{h} \quad \dots (29)$$

and the *mean square error* $E(x^2)$ as

$$E(x^2) = \int_{-\infty}^{\infty} x^2 f(x) dx = \frac{2h}{\sqrt{\pi}} \int_0^{\infty} x^2 e^{-h^2 x^2} dx = \frac{1}{2h^2} = \frac{0.5}{h^2} \quad \dots (30)$$

Lastly to discuss the effect of increasing the number of observations n , we have

$$x_i = t_i - v \text{ giving } \bar{x} = \bar{t} - v \quad \dots (31)$$

$$\text{where } \bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \text{ and } \bar{t} = \frac{1}{n} \sum_{i=1}^n t_i.$$

Conclusively, the error in the mean is the mean of the errors. Also the positive and negative errors tending to cancel in computing $2x_i$,

as n the number of observations increases, the value of error decreases.

ADDITIONAL MISCELLANEOUS PROBLEMS

Problem 142. State and Prove the normal law of errors, and find an expression for the measure of precision and the probable error of the arithmetic mean.

A cylinder has a length " l " cm., which is measurable with a probable error $\pm a$ and has a radius " a " cm., which is measurable with a probable error $\pm b$. Find the area of its curved surface and determine the probable error of this value.

(Nagpur, 1965)

Problem 143. Derive the normal law of errors and calculate the probable error of an observation.

(Vikram, 1969)

Problem 144. (a) Write short notes on the binomial, the Poissonian and normal distributions.

(b) Derive the normal law of errors from first principles and discuss some of its applications.

(Agra, 1969)

Problem 145. (a) Define Poisson's distribution and discuss its importance in Physics.

(b) Calculate the mean and the standard deviation for the Poisson distribution.

(Agra, 1970)

Problem 146. Write a comprehensive note on the theory of errors. (Agra, 1971)

Problem 147. If the probability function $P(x)$ of events x is given by the Gaussian function

$$P(x) = Ae^{-kx^2}$$

prove that $k = \frac{1}{2\sigma^2}$ and $A = \frac{1}{\sigma\sqrt{2\pi}}$ where σ is the standard deviation.

Assuming the errors in a series of observations to have the Gaussian distribution obtain expressions for the probable error of a single observation and that in the arithmetic mean of the observations.

(Agra, 1972)

Problem 148. Define the error function $\text{erf}(x)$ and obtain a convergent expansion for it in ascending powers of x correct upto x^5 . (See Appendix B)

Use the approximate expression to calculate the percentage of molecules of an ideal gas which have speeds less than $\frac{1}{2\sqrt{b}}$. The speed distribution of the mole-

cules is given to be $\frac{dN_c}{N} = \frac{2}{\sqrt{\pi}} b^3 c^2 e^{-bc^2} dc$; $\frac{dN_c}{N}$ being the fraction of molecules having speeds in the range c to $c+dc$. Given $\pi=3.4$ and $e^{1/4}=1.28$.

(Agra, 1973)

Problem 149. (a) Explain what you mean by a binomial distribution. Find its mean and standard deviation.

(b) The mean number of particles emitted in one second by a radioactive sample is 4.5. Write an expression for the probability of just 3 particles being emitted in a particular second.

(Agra, 1974)

Problem 150. Explain the meanings of the following terms in the theory of errors: dispersion, variance, standard deviation, regression and correlation.

(Agra, 1975)

Problem 151. (a) Explain normal distribution and derive an expression for the same.

(b) Explain the Principle of least squares.

(Rohilkhand, 1976)

APPENDICES

[A] SOME FORMULATED RESULTS IN BASIC MATHEMATICS

(a) Laws of Exponent Powers

$$x^a \cdot x^b = x^{a+b}; x^a \cdot y^a = (xy)^a; (x^a)^b = x^{ab}; \frac{x^a}{x^b} = x^{a-b}; x^{a/b} = \sqrt[b]{x^a},$$

$$x^0 = 1, x^{-a} = \frac{1}{x^a} \text{ provided } x \neq 0.$$

(b) Factors and Expansions

Taking n and r both as positive integers, we have

$${}^nC_r = \frac{{}^nC_0}{{}^nC_r} = \frac{n!}{r!(n-r)!} \text{ and } |r| = r(r-1)(r-2)\dots 3.2.1.$$

$${}^nC_0 = 1 = {}^nC_n; {}^nC_r = {}^nC_{n-r} \quad |1| = |0| = 1 \text{ and } |-r| = 0$$

$$(x+y)^n = x^n + {}^nC_1 x^{n-1}y + {}^nC_2 x^{n-2}y^2 + \dots + {}^nC_n y^n$$

$$= \sum_{r=0}^n {}^nC_r x^{n-r} y^r$$

$$x^n + y^n = (x+y)(x^{n-1} - x^{n-2}y + \dots + y^{n-1}), n \text{ being an odd integer}$$

$$x^n + y^n = (x+y)(x^{n-1} - x^{n-2}y + \dots - y^{n-1}), n \text{ being an even integer}$$

$$x^n - y^n = (x-y)(x^{n-1} + x^{n-2}y + \dots + y^{n-1}).$$

$$(x+y+z+\dots)^n = \sum_{\text{over all } r_i} \frac{n!}{|r_1|! |r_2|! |r_3|! \dots} x^{r_1} y^{r_2} z^{r_3} \dots$$

where $\sum r_i = n$.

In particular:

$$x^2 - y^2 = (x-y)(x+y); x^3 \pm y^3 = (x \pm y)(x^2 \mp xy + y^2)$$

(c) Summations

$$\Sigma n = 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

$$\Sigma n^2 = 1^2 + 2^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\Sigma n^3 = \frac{n^2(n+1)^2}{4}, \Sigma n^4 = \frac{n^5}{5} + \frac{n^4}{2} + \frac{n^3}{3} - \frac{n}{30}$$

$$\Sigma n^p = 1^p + 2^p + \dots + n^p = \frac{n^{p+1}}{p+1} + \frac{n^p}{2} + \frac{1}{2} {}^pC_1 B_1 n^{p-1}$$

$$- \frac{1}{4} {}^pC_3 B_2 n^{p-3} + \frac{1}{5} {}^pC_5 B_3 n^{p-5} + \dots$$

where B 's are Bernoulli's numbers and the series ends with n when p is even and with n^2 when p is odd. The Bernoulli's numbers are

$$B_1 = \frac{1}{6}, B_2 = \frac{1}{30}, B_3 = \frac{1}{42}, B_4 = \frac{1}{30}, B_5 = \frac{5}{66}, B_6 = \frac{691}{2730}, B_7 = \frac{7}{6} \text{ etc. } \dots$$

Taking a as the first term, d common difference and l the last term of an arithmetical progression, the sum

$$S = a + (a+d) + (a+2d) + \dots + \{a + (n-1)d\}$$

$$= \frac{n}{2} [2a + (n-1)d] = \frac{n}{2} [a+l]$$

where $l = a + (n-1)d = n$ th term of the series.

If a be the first term and r the common ratio of a geometrical progression, then the sum

$$S = a + ar + ar^2 + \dots + ar^{n-1} = \frac{a(r^n - 1)}{r - 1} \text{ if } r > 1$$

$$= \frac{a(1 - r^n)}{1 - r} \text{ if } r < 1$$

$$= \frac{a}{1 - r} \text{ if } n \text{ is infinite.}$$

(d) Ratio and Proportions

$$\text{If } \frac{p}{q} = \frac{r}{s} \text{ then } \frac{p+q}{q} = \frac{r+s}{s} \text{ and } \frac{p-q}{q} = \frac{r-s}{s}.$$

$$\text{Also } \frac{p-q}{p+q} = \frac{r-s}{r+s} \text{ (componendo and dividendo)}$$

$$\text{In general if } \frac{p}{q} = \frac{r}{s} = \frac{t}{u} = \dots = k \text{ then}$$

$$k = \frac{p+r+t+\dots}{q+s+u+\dots} = \frac{xp+yr+zt+\dots}{xq+ys+zu+\dots}$$

(e) Logarithms

We express $10^2 = 100$ by saying $\log_{10} 100 = 2$ and read 'logarithm of 100 to the base 10 is 2'.

In general, taking the base of the logarithm as e , we express $e^x = y$ by saying 'logarithm of y to the base ' e ' is x ' i.e. $\log_e y = x$

$$\log_e x + \log_e y = \log_e xy; \log_e x - \log_e y = \log_e \frac{x}{y}$$

$$\log_e x^m = m \log_e x; \log_e x^{-n} = \log_e \frac{1}{x^n} = -n \log_e x$$

$$\log_a x = \log_b x \log_a b = \frac{\log_b x}{\log_b a} \text{ (change of base)}$$

(f) Special Series

(i) *Binomial series* for $|x| < 1$ and for any n , positive, negative, integral or fractional,

APPENDICES

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1.2} x^2 + \frac{n(n-1)(n-2)}{1.2.3} x^3 + \dots$$

$$(1+x)^{-n} = 1 - nx + \frac{n(n+1)}{1.2} x^2 - \frac{n(n+1)(n+2)}{1.2.3} x^3 + \dots$$

In particular,

$$(1+x)^{-1} = 1 - x + x^2 - x^3 + x^4 - x^5 + \dots$$

(ii) Exponential series

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right)^n = 1 + \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots = 2.71828 \dots$$

$$e^x = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n} \right)^n = 1 + \frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{3} + \dots$$

$$e^{-x} = \lim_{n \rightarrow \infty} \left(1 - \frac{x}{n} \right)^n = 1 - x + \frac{x^2}{2} - \frac{x^3}{3} + \dots$$

$$a^x = 1 + x \log_e a + \frac{(x \log_e a)^2}{2} + \dots$$

(iii) Logarithmic series

$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \quad |x| < 1$$

$$\log(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \dots \quad |x| < 1$$

(iv) Other series

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$$

$$\frac{\pi^2}{6} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\frac{\pi^2}{12} = 1 - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$$

$$\frac{\pi^2}{8} = 1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots$$

$$\frac{\pi^2}{24} = \frac{1}{2^2} + \frac{1}{4^2} + \frac{1}{6^2} + \dots$$

(v) Taylor's series

$$(a+h) = f(a) + hf'(a) + \frac{h^2}{2} f''(a) + \dots + \frac{h^n}{n!} f^n(a) + \dots$$

$$f(x) = f(a) + (x-a)f'(a) + \frac{(x-a)^2}{2} f''(a) + \dots + \frac{(x-a)^n}{n!} f^n(a) + \dots$$

(vi) Maclaurin's series

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2} f''(0) + \dots + \frac{x^n}{n!} f^n(0) + \dots$$

(g) Sterling's formula

For very large values of n ,

$$|n| \approx \sqrt{2\pi n} \cdot n^n e^{-n} \text{ approx.} = e^{-n} n^{n+1/2} \sqrt{2\pi}$$

$$\log |n| = (n + \frac{1}{2}) \log n - n \log e + \log \sqrt{2\pi}$$

where $\pi = 3.14159\dots$ and $e = 2.71828\dots$

(h) Trigonometric functions

(i) In a right-angled triangle ABC where $B = 90^\circ$, and $\angle BAC = \theta$ (say) AB is the base, BC is the perpendicular in respect to θ and AC is the hypotenuse.

$$\sin \theta = \frac{\text{perpendicular}}{\text{hypotenuse}} = \frac{BC}{AC}; \cos \theta = \frac{\text{base}}{\text{hypotenuse}} = \frac{AB}{AC}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{BC}{AB}, \cot \theta = \frac{\cos \theta}{\sin \theta} = \frac{AB}{BC}$$

$$\sec \theta = \frac{1}{\cos \theta} \text{ and } \operatorname{cosec} \theta = \frac{1}{\sin \theta}$$

(ii) Signs of these functions in different quadrants

Quadrant	sin	cos	tan	cot	sec	cosec
I	+	+	+	+	+	+
II	+	-	-	-	-	+
III	-	-	+	+	-	-
IV	-	+	-	-	+	-

(iii) Values of these functions (sine and cosine only others can be calculated)

	0°	30°	45°	60°	90°	180°	270°	360°
Sin	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1	0	-1	0
Cos	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0	-1	0	1

$$\sin(90^\circ \pm \theta) = \cos \theta; \cos(90^\circ \pm \theta) = \pm \sin \theta$$

$$\sin(180^\circ \pm \theta) = \pm \sin \theta; \cos(180^\circ \pm \theta) = -\cos \theta$$

$$\sin(360^\circ \pm \theta) = \pm \sin \theta; \cos(360^\circ \pm \theta) = \cos \theta$$

$$\sin^2 \theta + \cos^2 \theta = 1; 1 + \tan^2 \theta = \sec^2 \theta; 1 + \cot^2 \theta = \operatorname{cosec}^2 \theta$$

$$\sin(\theta \pm \phi) = \sin \theta \cos \phi \pm \cos \theta \sin \phi$$

$$\cos(\theta \pm \phi) = \cos \theta \cos \phi \mp \sin \theta \sin \phi$$

$$\sin 2\theta = 2 \sin \theta \cos \theta; \cos 2\theta = \cos^2 \theta - \sin^2 \theta = 1 - 2 \sin^2 \theta = 2 \cos^2 \theta - 1$$

$$\sin 2\theta = \frac{2 \tan \theta}{1 + \tan^2 \theta}; \cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}$$

$$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}, \tan(\theta \pm \phi) = \frac{\tan \theta \pm \tan \phi}{1 \mp \tan \theta \tan \phi}$$

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta, \cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$$

$$\sin \theta + \sin \phi = 2 \sin \frac{\theta + \phi}{2} \cos \frac{\theta - \phi}{2}; \sin \theta - \sin \phi = 2 \cos \frac{\theta + \phi}{2} \sin \frac{\theta - \phi}{2}$$

$$\sin \frac{\theta - \phi}{2}; \cos \theta + \cos \phi = 2 \cos \frac{\theta + \phi}{2} \cos \frac{\theta - \phi}{2};$$

$$\cos \theta - \cos \phi = 2 \sin \frac{\theta + \phi}{2} \sin \frac{\phi - \theta}{2}$$

$$(iv) \sin^{-1} x = \cos^{-1} \sqrt{1 - x^2} = \tan^{-1} \frac{x}{\sqrt{1 - x^2}} = \operatorname{cosec}^{-1} \frac{1}{x}$$

$$\cos^{-1} x = \sin^{-1} \sqrt{1 - x^2} = \tan^{-1} \frac{\sqrt{1 - x^2}}{x} = \sec^{-1} \frac{1}{x}$$

$$\tan^{-1} x = \sin^{-1} \frac{x}{\sqrt{1 + x^2}} = \cos^{-1} \frac{1}{\sqrt{1 + x^2}} = \cot^{-1} \frac{1}{x}$$

$$\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$$

(v) Hyperbolic functions

$$\sinh x = \frac{e^x - e^{-x}}{2}, \cosh x = \frac{e^x + e^{-x}}{2}, \tanh x = \frac{e^x - e^{-x}}{e^x + e^{-x}} = \frac{\sinh x}{\cosh x}$$

$$\coth x = \frac{1}{\tanh x}; \operatorname{sech} x = \frac{1}{\cosh x}, \operatorname{cosech} x = \frac{1}{\sinh x}$$

$$\sinh^{-1} x = \log \{x + \sqrt{x^2 + 1}\}; \cosh^{-1} x = \log \{x + \sqrt{x^2 - 1}\}$$

$$\tanh^{-1} x = \frac{1}{2} \log \left(\frac{1+x}{1-x} \right)$$

$$\cosh^2 x - \sinh^2 x = 1; \operatorname{sech}^2 x + \tanh^2 x = 1,$$

$$\sinh 2x = 2 \sinh x \cosh x;$$

$$\cosh 2x = \cosh^2 x + \sinh^2 x = 1 + 2 \sinh^2 x = 2 \cosh^2 x - 1$$

$$\sin ix = i \sinh x; \cos ix = \cosh x$$

$$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta \text{ for all values of } n.$$

(vi) Trigonometric series

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$\tan x = x + \frac{x^3}{3} + \frac{2}{15}x^5 + \dots$$

$$\sin^{-1} x = x + \frac{1}{2} \frac{x^3}{3} + \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{x^5}{5} + \dots \quad |x| < 1$$

$$\tan^{-1} x = x - \frac{1}{3} x^3 + \frac{1}{5} x^5 - \dots \quad |x| < 1$$

$$\sinh x = x + \frac{x^3}{3} + \frac{x^5}{5} + \dots$$

$$\cosh x = 1 + \frac{x^2}{2} + \frac{x^4}{4} + \dots$$

$$\tanh x = x - \frac{x^3}{3} + \frac{2}{15}x^5 - \frac{17}{315}x^7 + \dots$$

$$\tanh^{-1} x = x + \frac{x^3}{3} + \frac{x^5}{5} + \frac{x^7}{7} + \dots \quad |x| < 1$$

(i) Analytical Geometry

If Cartesian coordinates of a point P be (x, y) and its polar coordinates be (r, θ) , where x is called abscissa, y is called ordinate, r is called radius vector and θ is called vectorial angle then,

$$x = r \cos \theta, \quad y = r \sin \theta$$

or
$$r = \sqrt{x^2 + y^2} \text{ and } \theta = \tan^{-1} \frac{y}{x}$$

Distance between two points $P(x_1, y_1)$ and $Q(x_2, y_2)$ is given by

$$PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

The slope (gradient) of the line is given by $\frac{y_2 - y_1}{x_2 - x_1}$ and denoted by m .

If m_1, m_2 be the gradients of two straight lines, the angle between them is

$$\tan^{-1} \frac{m_1 - m_2}{1 + m_1 m_2}$$

The two lines will be perpendicular or parallel according as,

$$m_1 m_2 = -1 \text{ and } m_1 = m_2.$$

Rectangular transformation (1) New axes parallel to old axes: If (x, y) and (x', y') respectively be the coordinates of a point w.r.t. old and new axes and (h, k) be the coordinates of the new origin w.r.t. old axes, then

$$x = x' + h, \quad y = y' + k$$

(2) New axes rotated through an angle θ about the origin from old axes: If (x, y) and (x', y') respectively be the coordinates of a point w.r.t. old and new axes, then

$$x = x' \cos \theta - y' \sin \theta, \quad y = x' \sin \theta + y' \cos \theta$$

or conversely,

$$x' = x \cos \theta + y \sin \theta, \quad y' = -x \sin \theta + y \cos \theta$$

(j) Differential Calculus

$$d(ax) = a dx, \quad a \text{ being a constant, } d(a) = 0$$

$$d(2x - 3y + z) = 2dx - 3dy + dz$$

$$d(xy) = x dy + y dx$$

$$d\left(\frac{x}{y}\right) = \frac{y dx - x dy}{y^2}$$

$$d(x^n) = nx^{n-1} dx; \quad d(x^y) = yx^{y-1} dx + x^y (\log_e x) dy$$

$$d(e^x) = e^x dx; \quad d(e^{\pm ax}) = \pm ae^{\pm ax} dx$$

$$d(a^x) = (a^x \log_e a) dx; \quad d(x^x) = x^x (1 + \log_e x) dx$$

$$d(\log_e x) = \frac{1}{x} dx; \quad d(\log_a x) = \frac{1}{x} \log_a dx = \frac{dx}{x \log_e a}$$

$$d(\sin x) = \cos x dx; \quad d(\cos x) = -\sin x dx$$

$$d(\tan x) = \sec^2 x dx; \quad d(\cot x) = -\operatorname{cosec}^2 x dx$$

$$d(\sec x) = \sec x \tan x dx; \quad d(\operatorname{cosec} x) = -\operatorname{cosec} x \cot x dx$$

$$d(\sinh x) = \cosh x dx; \quad d(\cosh x) = \sinh x dx$$

$$d(\tanh x) = \operatorname{sech}^2 x dx; \quad d(\coth x) = -\operatorname{cosech}^2 x dx$$

$$d(\operatorname{sech} x) = -\operatorname{sech} x \tanh x dx; \quad d(\operatorname{cosech} x) = -\operatorname{cosech} x \coth x dx$$

$$d(\sin^{-1} x) = \frac{dx}{\sqrt{1-x^2}}; \quad d(\cos^{-1} x) = -\frac{dx}{\sqrt{1-x^2}}$$

$$d(\tan^{-1} x) = \frac{dx}{1+x^2}; \quad d(\cot^{-1} x) = -\frac{dx}{1+x^2}$$

$$d(\sec^{-1} x) = \frac{dx}{x\sqrt{x^2-1}}; \quad d(\operatorname{cosec}^{-1} x) = -\frac{dx}{x\sqrt{x^2-1}}$$

$$d(\sinh^{-1} x) = \frac{dx}{\sqrt{x^2+1}}; \quad d(\cosh^{-1} x) = \frac{dx}{\sqrt{x^2-1}}$$

$$d(\tanh^{-1} x) = \frac{dx}{1-x^2}; \quad d(\coth^{-1} x) = -\frac{dx}{x^2-1}$$

$$d(\operatorname{sech}^{-1} x) = \frac{dx}{-x\sqrt{(1-x^2)}}; \quad d(\operatorname{cosech}^{-1} x) = -\frac{dx}{x\sqrt{(x^2+1)}}$$

Taking $D \equiv \frac{d}{dx}$, we have

$$D^n(ax+b)^m = m(m-1)(m-2)\dots(m-n+1)a^n(ax+b)^{m-n}$$

$$D^n(ax+b)^{-1} = (-1)^n \frac{n!}{(n-1)!} a^n(ax+b)^{-n-1}$$

$$D^n e^{ax} = a^n e^{ax}; \quad D^n a^x = (\log a)^n a^x$$

$$D^n \log(ax+b) = \frac{(-1)^{n-1} (n-1)! a^n}{(ax+b)^n}$$

$$D^n \sin(ax+b) = a^n \sin\left(ax+b + \frac{n\pi}{2}\right)$$

$$D^n \cos(ax+b) = a^n \cos\left(ax+b + \frac{n\pi}{2}\right)$$

$$D^n \left\{ e^{ax} \frac{\sin}{\cos}(bx+c) \right\} = r^n e^{ax} \frac{\sin}{\cos}(bx+c+n\phi)$$

where $r = \sqrt{a^2 + b^2}$ and $\phi = \tan^{-1} \frac{b}{a}$

$$D^n(uv) = (D^n u) \cdot v + nC_1 D^{n-1} u \cdot Dv + nC_2 D^{n-2} u \cdot D^2 v + \dots + nC_r D^{n-r} u \cdot D^r v + \dots + u \cdot D^n v$$

(k) Integral Calculus :

$$\int x^n dx = \frac{x^{n+1}}{n+1}, n \neq -1; \int \frac{1}{x} dx = \log x; \int e^x dx = e^x; \int a^x dx = \frac{a^x}{\log a}$$

$$\int \sin x dx = -\cos x; \int \cos x dx = \sin x; \int \tan x dx = -\log \cos x$$

$$\int \cot x dx = \log \sin x; \int \sec x dx = \log \tan \left(\frac{x}{2} + \frac{\pi}{4} \right)$$

$$= \log (\sec x + \tan x)$$

$$\int \operatorname{cosec} x dx = \log \tan \frac{x}{2}; \int \frac{dx}{a^2+x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a}; \int \frac{dx}{x^2-a^2}$$

$$= \frac{1}{2a} \log \frac{x-a}{x+a}$$

$$\int \frac{dx}{\sqrt{a^2-x^2}} = \sin^{-1} \frac{x}{a}; \int \frac{dx}{\sqrt{a^2+x^2}} = \sinh^{-1} \frac{x}{a}$$

$$= \log \left\{ x + \sqrt{x^2+a^2} \right\}$$

$$\int \frac{dx}{\sqrt{x^2-a^2}} = \cosh^{-1} \frac{x}{a} = \log \left\{ x + \sqrt{x^2-a^2} \right\}; \int \frac{dx}{x\sqrt{x^2-a^2}}$$

$$= \frac{1}{a} \sec^{-1} \frac{x}{a}$$

$$\int \sqrt{a^2-x^2} dx = \frac{x}{2} \sqrt{a^2-x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a}$$

$$\int \sqrt{a^2+x^2} dx = \frac{x}{2} \sqrt{a^2+x^2} + \frac{a^2}{2} \sinh^{-1} \frac{x}{a}$$

$$\int \sqrt{x^2-a^2} dx = \frac{x}{2} \sqrt{x^2-a^2} - \frac{a^2}{2} \cosh^{-1} \frac{x}{a}$$

$$\int e^{ax} \sin bx dx = \frac{e^{ax}}{\sqrt{a^2+b^2}} \sin \left(bx - \tan^{-1} \frac{b}{a} \right)$$

$$\int e^{ax} \cos bx dx = \frac{e^{ax}}{\sqrt{a^2+b^2}} \cos \left(bx - \tan^{-1} \frac{b}{a} \right)$$

$$\int \sinh x dx = \cosh x; \int \cosh x dx = \sinh x; \int \operatorname{sech}^2 x dx = \tanh x$$

$$\int \operatorname{cosech}^2 x dx = -\coth x$$

$$\int (ax+b)^n dx = \frac{(ax+b)^{n+1}}{(n+1)a}; \int \frac{dx}{ax+b} = \frac{1}{a} \log (ax+b)$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}; \int \sin (ax+b) dx = -\frac{1}{a} \cos (ax+b)$$

$$\int \sin^n x dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} \int \sin^{n-2} x dx, n \text{ positive integer}$$

$$\int \frac{dx}{\sin ax} = \frac{1}{a} \log \tan \frac{ax}{2} = \frac{1}{a} \log (\operatorname{cosec} ax - \cot ax)$$

$$\int \frac{dx}{1 \pm \sin ax} = \mp \frac{1}{a} \tan \left(\frac{\pi}{4} \mp \frac{ax}{2} \right)$$

$$\int \sin ax \sin bx dx = \frac{\sin (a-b)x}{2(a-b)} - \frac{\sin (a+b)x}{2(a+b)}, a^2 \neq b^2$$

$$\int \cos^n x dx = \frac{\cos^{n-1} x \sin x}{n} + \frac{n-1}{n} \int \cos^{n-2} x dx, n \text{ positive integer}$$

$$\int \frac{dx}{\cos ax} = \frac{1}{a} \log \tan \left(\frac{ax}{2} + \frac{\pi}{4} \right) = \frac{1}{a} \log (\tan ax + \sec ax)$$

$$\int \frac{dx}{1+\cos ax} = \frac{1}{a} \tan \frac{ax}{2}, \int \frac{dx}{1-\cos ax} = -\frac{1}{a} \cot \frac{ax}{2}$$

$$\int \sin^m ax \cos^n ax dx = -\frac{\sin^{m-1} ax \cos^{n+1} ax}{a(m+n)} + \frac{m-1}{m+n}$$

$$\int \sin^{m-2} ax \cos^n ax dx; m, n > 0$$

$$= \frac{\sin^{m+1} ax \cos^n ax}{a(m+n)} + \frac{n-1}{m+n} \int \sin^m ax \cos^{n-2} ax dx; m, n > 0$$

$$\int \tan^n ax dx = \frac{1}{a(n-1)} \tan^{n-1} ax - \int \tan^{n-2} ax dx, \text{ integral } n > 1$$

$$\int \cot^n ax dx = -\frac{1}{a(n-1)} \cot^{n-1} ax - \int \cot^{n-2} ax dx, \text{ integral } n > 1$$

$$\int \sec^n ax dx = \frac{1}{a(n-1)} \frac{\sin ax}{\cos^{n-1} ax} + \frac{n-2}{n-1} \int \sec^{n-2} ax dx, \text{ integral } n > 1$$

$$\int \operatorname{cosec}^n ax dx = -\frac{1}{a(n-1)} \frac{\cos ax}{\sin^{n-1} ax} + \frac{n-2}{n-1} \int \operatorname{cosec}^{n-2} ax dx, \text{ integral } n > 1$$

$$\int x^n e^{ax} dx = \frac{1}{a} x^n e^{ax} - \frac{n}{a} \int x^{n-1} e^{ax} dx, \text{ positive } n$$

$$\int e^{ax} \sin bx dx = \frac{e^{ax}}{a^2+b^2} (a \sin bx - b \cos bx)$$

$$\int e^{ax} \cos bx dx = \frac{e^{ax}}{a^2+b^2} (a \cos bx + b \sin bx)$$

$$\int \log ax dx = x \log ax - x; \int \frac{dx}{x \log ax} = \log (\log ax)$$

$$\int x^n (\log ax)^m dx = \frac{x^{n+1}}{n+1} (\log ax)^m - \frac{m}{n+1} \int x^n (\log ax)^{m-1} dx$$

$$\int e^{ax} \log bx dx = \frac{1}{a} e^{ax} \log bx - \frac{1}{a} \int \frac{e^{ax}}{x} dx$$

$$\int x^m (ax+b)^n dx = \frac{1}{a(m+n+1)} \left[x^m (ax+b)^{n+1} - mb \int x^{m-1} (ax+b)^n dx \right]$$

$$= \frac{1}{m+n+1} \left[x^{m+1} (ax+b)^n + nb \int x^m (ax+b)^{n-1} dx \right],$$

$$m > 0, m+n+1 \neq 0$$

$$\int \frac{dx}{(ax+b)(cx+d)} = \frac{1}{bc-ad} \log \frac{cx+d}{ax+b}, bc-ad \neq 0$$

$$\int \frac{dx}{(cx+d)\sqrt{ax+b}} = \frac{2}{\sqrt{c}\sqrt{ad-bc}} \tan^{-1} \frac{\sqrt{c(ax+b)}}{ad-bc},$$

$$c > 0, ad > bc$$

Definite Integrals

$$\int_0^\infty \frac{a dx}{a^2+x^2} = \begin{cases} \pi/2, & a > 0 \\ 0, & a = 0 \\ -\pi/2, & a < 0 \end{cases}$$

$$\int_0^\infty x^{n-1} e^{-x} dx = \left[\log \frac{1}{x} \right]^{n-1} dx = \sqrt{n}$$

$$\int_0^\infty x^{m-1} (1-x)^{n-1} dx = \int_0^\infty \frac{x^{m-1} dx}{(1+x)^{m+n}} = B(m, n) = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}$$

$$\int_0^\infty \frac{x^{n-1} dx}{1+x} = \frac{\pi}{\sin n\pi}, 0 < n < 1$$

$$\int_0^\infty e^{-ax} \cos bx dx = \frac{a}{a^2+b^2}, a > 0$$

$$\int_0^\infty e^{-ax} \sin bx dx = \frac{b}{a^2+b^2}, a > 0$$

$$\int_0^\pi \frac{dx}{a+b \cos x} = \frac{\pi}{\sqrt{a^2-b^2}}, a > b > 0$$

$$\int_0^\infty \frac{\cos ax dx}{1+x^2} = \frac{\pi}{2} e^{-a}, a > 0 \text{ and } \frac{\pi}{2} e^a, a < 0$$

$$\int_0^\infty \frac{\cos x}{\sqrt{x}} dx = \int_0^\infty \frac{\sin x dx}{\sqrt{x}} = \sqrt{\frac{\pi}{2}}; \int_0^{\pi/2} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x}$$

$$= \frac{\pi}{2ab}$$

$$\int_0^\infty e^{-ax} \cosh bx dx = \frac{a}{a^2-b^2}, a > 0, a^2 \neq b^2$$

$$\int_0^\infty e^{-ax} \sinh bx dx = \frac{b}{a^2-b^2}, a > 0, a^2 \neq b^2$$

$$\int_0^\infty e^{-x} \log x dx = \int_0^\infty \left(\frac{1}{1-e^{-x}} - \frac{1}{x} \right) e^{-x} dx$$

$$= \int_0^\infty \frac{1}{x} \left(\frac{1}{1+x} - e^{-x} \right) dx$$

$$= \int_0^1 \left(\frac{1-e^{-x}-e^{-1/x}}{x} \right) dx = \gamma = .5772152$$

where γ is Euler's constant given by

$$r = \lim_{t \rightarrow \infty} \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{t} - \log t \right)$$

(I) Elliptic Integrals

Elliptic integrals of first, second and third kinds are respectively

$$F(k, \phi) = \int_0^\phi \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}}; 0 < k < 1$$

$$E(k, \phi) = \int_0^\phi \sqrt{1-k^2 \sin^2 \theta} d\theta; 0 < k < 1$$

$$H(k, \phi, a) = \int_0^\phi \frac{d\theta}{(\sqrt{1-k^2 \sin^2 \theta})(1+a^2 \sin^2 \theta)}, 0 < k < 1, a \neq k$$

If ϕ , the upper limit is replaced by $\frac{\pi}{2}$, these become complete

elliptic integrals:

$$K(k) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}}$$

$$E(k) = \int_0^{\pi/2} \sqrt{1-k^2 \sin^2 \theta} d\theta$$

$$H(k, a) = \int_0^{\pi/2} \frac{d\theta}{(\sqrt{1-k^2 \sin^2 \theta})(1+a^2 \sin^2 \theta)}$$

(m) Probability functions

If $\phi(x)$ be a normal function s.t. $\phi(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$

and $\alpha = \int_{-x}^x \phi(x) dx$ i.e. area under $\phi(x)$ from $-x$ to x , then

$$\frac{1}{2}(1+\alpha) = \int_{-\infty}^x \phi(x) dx$$

Second derivative if $\phi(x)$ i.e. $\phi^{(2)}(x) = (x^2-1)\phi(x)$

Third ,, ,, i.e. $\phi^{(3)}(x) = (3x-x^3)\phi(x)$

Fourth ,, ,, i.e. $\phi^{(4)}(x) = (x^4-6x^2+3)\phi(x)$

Now if $(p+q)^n = \sum_{r=0}^n {}^nC_r p^{n-r} q^r$, $p+q=1$, then

$$\sum_{r=a}^b {}^nC_r p^{n-r} q^r = \int_{x_1}^{x_2} \phi(x) dx + \left[\frac{q-p}{6\sigma} \phi^{(2)}(x) + \frac{1}{24} \left(\frac{1}{\sigma^2} - \frac{6}{n} \right) \phi^{(3)}(x) \right]_{x_1}^{x_2} \text{ approx.}$$

where $x_1 = \frac{(a - \frac{1}{2} - nq)}{\sigma}$; $x_2 = \frac{(b + \frac{1}{2} - nq)}{\sigma}$, $\sigma = \sqrt{npq}$ and $a \leq r \leq b$.

$$\sum_{r=0}^r {}^nC_r p^{n-r} q^r = \int_x^\infty \phi(x) dx + \frac{q-p}{6\sigma} \phi^{(2)}(x) - \frac{1}{24} \left(\frac{1}{\sigma^2} - \frac{6}{n} \right) \phi^{(3)}(x) \quad \text{approx.}$$

where $x = \frac{(n-r-\frac{1}{2}-np)}{\sigma}$

$$\sum_{r=r}^n {}^nC_r p^{n-r} q^r \int_x^\infty \phi(x) dx - \frac{q-p}{6\sigma} \phi^{(2)}(x) - \frac{1}{24} \left(\frac{1}{\sigma^2} - \frac{6}{n} \right) \phi^{(3)}(x) \quad \text{approx.}$$

where $x = \frac{(r-\frac{1}{2}-nq)}{\sigma}$.

[B] Asymptotic Expansion of Error-Function

The Error function is defined as

$$\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt \quad \dots (1) \text{ where } \int_0^\infty e^{-t^2} dt = \frac{1}{2} \sqrt{\pi} \quad \dots (2)$$

$$(1) \text{ and } (2) \text{ yield } \operatorname{erf} \infty = \lim_{x \rightarrow \infty} \operatorname{erf} x = 1 \quad \dots (3)$$

Using Maclaurin series of e^{-x^2} after integrating term by term, for small values of $|x|$, $\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \right) \quad \dots (4)$

For large values of x , the asymptotic expansion of $\operatorname{erf} x$ given below is used.

The Complementary error function is defined as

$$\operatorname{erfc} x = \frac{2}{\sqrt{\pi}} \int_x^\infty e^{-t^2} dt \quad \dots (5)$$

The relation between error and complementary error functions is given by

$$\operatorname{erfc} x = 1 - \operatorname{erf} x$$

$$\text{or } \operatorname{erf} x = 1 - \operatorname{erfc} x \quad \dots (6)$$

Now to find the asymptotic expansion of $\operatorname{erf} x$, let us

put $t^2 = p$ i.e. $dt = \frac{1}{2\sqrt{p}} dp$ so that we have

$$\operatorname{erfc} x = \frac{2}{\sqrt{\pi}} \int_x^\infty e^{-t^2} dt = \frac{1}{\sqrt{\pi}} \int_{x^2}^\infty \frac{e^{-p}}{x^2} p^{-\frac{1}{2}} dp$$

If we go on integrating the R.H.S. by parts, then we shall have after n trials, an integral of the form

$$(\text{say}) I_n(x) = \int_{x^2}^\infty e^{-p} p^{-\frac{2n+1}{2}} dp, n=0, 1, 2, \dots \quad \dots (7)$$

$$\text{whence } \operatorname{erfc} x = \frac{1}{\sqrt{\pi}} I_0(x) \quad \dots (8)$$

Now integrating R.H.S. of (7) by parts, we get

$$I_n(x) = \left[p^{-\frac{2n+1}{2}} (-e^{-p}) \right]_{x^2}^\infty - \frac{2n+1}{2} \int_{x^2}^\infty p^{-\frac{2n+3}{2}} e^{-p} dp \\ = \frac{1}{x^{2n+1}} e^{-x^2} - \frac{2n+1}{2} I_{n+1}(x), n=0, 1, 2, \dots$$

Multiplying both sides by e^{x^2} , we get the recurrence relation

$$e^{x^2} I_n(x) = \frac{1}{x^{2n+1}} - \frac{2n+1}{2} e^{x^2} I_{n+1}(x) \quad \dots (9)$$

Applying this result repeatedly, we find

$$e^{x^2} I_0(x) = \frac{1}{x} - \frac{1}{2} e^{x^2} F_1(x) \quad (\text{for } n=0) \\ = \frac{1}{x} - \frac{1}{2x^3} + \frac{1}{2} \cdot \frac{3}{2} e^{x^2} F_2(x) \quad (\text{for } n=1) \\ \dots \dots \dots \\ = \left\{ \frac{1}{x} - \frac{1}{2x^3} + \frac{1.3}{2^2 x^5} - \dots + (-1)^{n-1} \frac{1.3.5 \dots (2n-3)}{2^{n-1} x^{2n-1}} \right\} \\ + (-1)^n \frac{1.3.5 \dots (2n-1)}{2^n} e^{x^2} I_n(x) \quad \dots (10)$$

In order to show that this series is an asymptotic expansion,

$$\text{consider, } e^{x^2} I_0(x) \sim \frac{1}{x} - \frac{1}{2x^3} + \frac{1.3}{2^2 x^5} \dots \quad \dots (11)$$

$$\text{and denote } T_{2n-1} = \frac{1}{x} - \frac{1}{2x^3} + \frac{1.3}{2^2 x^5} - \dots + (-1)^{n-1} \frac{1.3.5 \dots (2n-3)}{2^{n-1} x^{2n-1}} \quad \dots (12)$$

As such (10) can be rewritten as

$$\left(e^{x^2} I_0(x) - T_{2n-1} \right) = (-1)^n \frac{1.3.5 \dots (2n-1)}{2^n} e^{x^2} I_n(x)$$

Multiplying both sides by x^{2n-1} and putting

$$f_n = (-1)^n \frac{1.3.5 \dots (2n-1)}{2^n}, \text{ the last relation yields}$$

$$\left(e^{x^2} I_0(x) - T_{2n-1} \right) x^{2n-1} = f_n e^{x^2} x^{2n-1} I_n(x) \quad \dots (13)$$

The series in (11) will be an asymptotic expansion, if we can show that for each fixed $n=1, 2, \dots$ $f_n \rightarrow 0$ as $x \rightarrow \infty$.

Now in (7), we have

$$\frac{1}{p^{2n+1/2}} \leq \frac{1}{x^{2n+1}} \quad \text{for all } p \geq x^2$$

$$\text{So that, } I_n(x) = \int_{x^2}^\infty \frac{e^{-p}}{p^{2n+1/2}} dp < \frac{1}{x^{2n+1}} \int_{x^2}^\infty e^{-p} dp$$

$$= \frac{e^{-x^2}}{x^{2n+1}} \quad \dots (14)$$

Thus R.H.S. of (13) i.e.

$$|f_n| e^{x^2} x^{2n-1} I_n(x) < \frac{|f_n|}{x^2} \rightarrow 0 \text{ as } x \rightarrow \infty.$$

Hence the series on the right hand side of (10) is an asymptotic expansion of the function $e^{x^2} I_0(x)$.

Thus $\operatorname{erf} x = 1 - \operatorname{erfc} x = 1 - \frac{I_0(x)}{\sqrt{\pi}}$ renders the required asymptotic expansion of the error function as

$$\operatorname{erf} x \sim 1 - \frac{1}{\sqrt{\pi}} e^{-x^2} \left(\frac{1}{x} - \frac{1}{2x^3} + \frac{1.3}{2^2 x^5} - \frac{1.3.5}{2^3 x^7} + \dots \right) \dots (15)$$

For large values of x , this reduces to

$$\operatorname{erf} x \approx 1 - \frac{1}{\sqrt{\pi}} \frac{e^{-x^2}}{x} \dots (16)$$

Note 1. $\frac{1.3.5 \dots (2n-1)}{2^n \sqrt{\pi}} |I_n(x)|$ is known as the absolute value of the error.

Note 2. Here $\sim$ read as asymptotically equal.

[C] Character Tables in Group Theory

Denoting a reducible or irreducible representation of a group $G = \{E, A, B, \dots\}$ by Γ , its character χ may be defined as the set of traces of all the matrices of representation Γ i.e.,

$$\chi(A) = \sum_i \Gamma_{ii}(A) \dots (1)$$

where Γ_{ii} are the elements of the matrix corresponding to A , the matrix of representation.

It is notable that the character is the same as the representation in case of one-dimensional representation and also the characters are the same in case of representations of conjugate elements since the trace of a matrix under a similarity transformation is invariant.

Taking A and B as conjugate elements, there may be found an element C such that

$$A = C^{-1} B C \dots (2)$$

which yields on taking traces on either side

$$\Gamma(A) = \Gamma(C^{-1}) \Gamma(B) \Gamma(C) \dots (3)$$

But the cyclic property of traces gives for any three matrices P, Q, R ,

$$\operatorname{tr}(PQR) = \operatorname{tr}(QRP) = \operatorname{tr}(RPQ) \dots (4)$$

$$\operatorname{tr}(\Gamma(A)) = \operatorname{tr}(\Gamma(B)) \Rightarrow \chi(A) = \chi(B) \dots (5)$$

Now if we denote an irreducible representation of the group G by $\Gamma^{(i)}$ and assume that an irreducible representation may occur two or more times in reduction of a reducible representation Γ , then the matrices of representation Γ are the direct sum of the matrices of

the components of irreducible representations i.e.

$$\Gamma = \sum_i a_i \Gamma^{(i)} \dots (6)$$

where a_i 's are non-negative integers.

Taking traces of either side of (6), for A , the matrix of representation,

$$\operatorname{tr} \Gamma(A) = \sum_i a_i \operatorname{tr} \Gamma^{(i)}(A) \quad \forall A \in G$$

which in view of (5) yields

$$\chi(A) = \sum_i a_i \chi^{(i)}(A) \dots (7)$$

Now if we call $\Gamma_{jk}^{(i)}$ as the representation vectors in the group space, then the orthogonality theorem for an irreducible representation in the present notation can be stated as

$$\sum_{A \in G} \Gamma_{jk}^{(i)}(A) \Gamma_{j'k'}^{(i')*}(A^{-1}) = 0$$

$$\text{i.e.,} \quad \sum_{A \in G} \Gamma_{jk}^{(i)}(A) \Gamma_{k'j'}^{(i')*}(A) = 0 \dots (8)$$

which means that the product of (j, k) th element of the irreducible representation $\Gamma^{(i)}$ with the complex conjugate of the (k', j') th element of the irreducible representation $\Gamma^{(i')}$ summed over all the group elements is zero.

Also taking g as the order of the group G and m as the dimension of the representation, in case of the elements of a single unitary irreducible representation, we have

$$\sum_{A \in G} \Gamma_{jk}^{(i)}(A) \Gamma_{j'k'}^{(i)}(A^{-1}) = \frac{g}{m} \delta_{jk'} \delta_{kj'}$$

$$\text{i.e.,} \quad \sum_{A \in G} \Gamma_{jk}^{(i)}(A) \Gamma_{k'j'}^{(i)*}(A) = \frac{g}{m} \delta_{jk'} \delta_{kj'} \dots (9)$$

Combining (8) and (9) we can write

$$\sum_{A \in G} \Gamma_{jk}^{(i)}(A) \Gamma_{k'j'}^{(i')*}(A) = \frac{g}{m} \delta_{ii'} \delta_{jk'} \delta_{kj'} \dots (10)$$

In order to transform (10) into an orthogonality relation for the characters of irreducible representation, putting $j=k$ and $j'=k'$ and summing over j and j' we get with the help of (1),

$$\sum_{A \in G} \chi^{(i)}(A) \chi^{(i')*}(A) = \frac{2}{m} \delta_{ii'} m = g \delta_{ii'} \dots (11)$$

where $\chi^{(i)}(A)$ represents the character of the element A in the irreducible representation $\Gamma^{(i)}$

Now multiplying both sides of (7) by $\chi^{(i')*}(A)$ and summing over all the elements of G , we find

$$\sum_{A \in G} \chi^{(i')*}(A) \chi(A) = \sum_i a_i \sum_{A \in G} \chi^{(i')*} \chi^{(i)}(A) = a_i g \text{ by (11)}$$

giving $a_i' = \frac{1}{g} \sum_{A \in G} \chi^{(i)*}(A) \chi(A)$

or replacing i' by i , this becomes

$$a_i = \frac{1}{g} \sum_{A \in G} \chi^{(i)*}(A) \chi(A) \quad \dots (12)$$

which gives coefficient a_i as introduced in (6).

It should be noted that the characters of the irreducible representations are known as *Primitive* or *simple character* whereas those of reducible ones are known as *compound character*. Actually a compound character may be decomposed in to simple characters.

Again multiplying (7) by its complex conjugate i.e., $\chi^*(A)$ and summing over all the elements of G and then dividing by g , we get

$$\begin{aligned} \frac{1}{g} \sum_{A \in G} \chi^*(A) \chi(A) &= \frac{1}{g} \sum_{i, j} a_i^* a_j \sum_{A \in G} \chi^{(i)*}(A) \chi^{(j)}(A) \\ &= \sum_i |a_i|^2 \text{ by (12)} \end{aligned} \quad \dots (13)$$

If the R.H.S. of (13) is unity for any representation Γ , then it follows that all a_i 's except one must be zero. Let $a_k \neq 0$, then it must be unity i.e., $a_k = 1$, in which case Γ must be equivalent (identical) with the irreducible representation $\Gamma^{(k)}$ and hence for a representation to be irreducible the *criterion* (i.e., the necessary and sufficient condition) is

$$\sum_{A \in G} \chi^*(A) \chi(A) = g \quad \dots (14)$$

Now to discuss some character tables, let us use the following notations.

Let the entire group consist of s classes say $c_1, c_2, c_3 \dots c_s$ with elements $h_1, h_2, h_3, \dots, h_s$ then clearly

$$g = h_1 + h_2 + \dots + h_s.$$

Also, take

E = Identity operation

C_n = rotation about an axis of symmetry through an angle equal to $\frac{2\pi}{n}$.

C_2', C_2'' = rotation through 180° about two axes perpendicular to C_n .

σ_v = reflection in a plane of symmetry, containing the principal axis i.e. the axis having the largest n value in C_n .

σ_d = reflection in a plane of symmetry, containing the principal axis and bisecting the angle between C_2' and C_2'' .

σ_h = reflection in a plane perpendicular to the principal axis.

i = inversion in a centre of symmetry.

S_n = rotation about an axis by $\frac{2\pi}{n}$, followed by a reflection in a plane perpendicular to the axis of rotation.

Here $\sigma_d^2 = \sigma_d$, σ_d and $C_n^3 = C_n$, C_n , C_n etc...

Let R, T denote the irreducible representations of the rotational and translational coordinates respectively in addition to x, y, z , coordinates. The reflections are also denoted by m , with subscripts if necessary, in places of σ_v (vertical reflection), σ_h (horizontal reflection) and σ_d (diagonal reflection).

To clearly understand these notations, we give two particular examples.

[1] Symmetry Transformations of an Equilateral Triangle.

Let A, B, C be the vertices of an equilateral triangle with D, E, F as mid-points of sides BC, CA, AB respectively and O the centroid.




Fig. C.1

To discuss the operations (transformations) of the triangle such as rotations, reflections, inversion, translation etc., leaving the triangle invariant, if we take z -axis as an axis through O normal to the plane of triangle, then rotations through $\frac{2\pi}{3}$ about

z -axis and its multiples leave the triangle invariant.

The six symmetry operations of an equilateral triangle are :




Fig. C.2

$E \rightarrow$ identity operation.




Fig. C.3

$C_3 \rightarrow$ anticlockwise rotation through $\frac{2\pi}{3}$ about z -axis.




Fig. C.4

$C_3^2 \rightarrow$ anticlockwise rotation through $\frac{4\pi}{3}$ about z -axis.




Fig. C.5

$m_1 \rightarrow$ reflection in the vertical plane passing through AD .




Fig. C.6

$m_2 \rightarrow$ reflection in the vertical plane passing through BE .




Fig. C.7

$m_3 \rightarrow$ reflection in the vertical plane passing through CF .

It is easy to see that the operation $C_3 m_1$ is the same as that of m_3 i.e. $C_3 m_1 = m_3$ etc. Also since $(C_3)^{-1} = C_3^2$ gives $C_3 (C_3)^{-1} = C_3 \cdot C_3^2$ i.e. $E = C_3 \cdot C_3^2$ therefore inverse of C_3 is C_3^2 and vice versa. E is the identity element.

Such a set being a group of order six is denoted by C_{3v} and known as the *symmetry group of an equilateral triangle*.

Multiplication table for group C_{3v} .

Second operation ↓	First operation →					
	E	C_3	C_3^2	m_1	m_2	m_3
E	E	C_3	C_3^2	m_1	m_2	m_3
C_3^2	C_3^2	E	C_3	m_2	m_3	m_1
C_3	C_3	C_3^2	E	m_3	m_1	m_2
m_1	m_1	m_3	m_2	E	C_3	C_3^2
m_2	m_2	m_1	m_3	C_3^2	E	C_3
m_3	m_3	m_2	m_1	C_3	C_3^2	E

The ordering of rows and columns being immaterial, we have chosen an ordering such that principal diagonal contains the identity element E only and this can be effected by choosing an order such that an element in the first column (second operation) is inverse of the corresponding element in first row (i.e. first operation).

Now if $G = \{E, A, B, C, \dots\}$ be a finite group of order 'g' with identity element E and $T = \{T(E), T(A), T(B), \dots\}$ be a set of square matrices all of the same order such that

$$T(A)T(B) = T(AB) \quad \dots (15)$$

then there arise two cases: (1) All the matrices of the set T are distinct in which case G and T are isomorphic to each other and so the representation generated by the elements of T is known as a faithful representation of G , (2) when the elements of T are not all distinct, G and T being homomorphic to each other, the representation is known as an unfaithful representation of G .

An identity representation which is actually an unfaithful one is the simplest representation obtained by associating unity (which is a square matrix of order one) with every element of the group e.g. for the group C_{3v} , we have

$$\left. \begin{array}{l} \text{Element: } E \quad C_3 \quad C_3^2 \quad m_1 \quad m_2 \quad m_3 \\ \text{Representation: } 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \end{array} \right\} \dots (16)$$

It should be noted that every group has at least one faithful representation.

Now to prepare the character table for group C_{3v} , we are required to have the following points in consideration:

Note I If C be the total number of irreducible representations of a finite group $G = \{E, A, B, \dots\}$ of order g and dimension l_i for $i=1, 2, 3, \dots, c$ then $\sum_{i=1}^c l_i^2 \leq g$. $\dots (17)$

Note II. The character being a function of the classes just as a representation is a function of the group elements, we have number of irreducible representation of $G \leq$ number of classes of G . $\dots (17a)$

Note III. Taking n_k as the number of elements in a class, C_k of group G , and $\phi^{(i)}$ as the character vectors in the class space, the orthogonality of characters yields for different classes,

$$\sum_{i=1}^c \phi_k^{(i)*} \phi_l^{(i)} = \frac{g}{n_k} \delta_{kl} \quad \dots (18)$$

whereas the product of two classes is defined as

$$C_i C_j = \sum_k a_{ijk} C_k \quad \dots (19)$$

a_{ijk} being non-negative integers.

Denoting by P_i^α the resulting matrix obtained by adding the matrices representing the elements of C_i in the irreducible representation $\Gamma^{(\alpha)}$, we have

$$P_i^\alpha = \sum_{A \in C_i} \Gamma^{(\alpha)}(A) \quad \dots (20)$$

Constructing the matrices for all the classes of G in a similar way, we have the normalization condition for any $B \in G$.

$$[\Gamma^{(\alpha)}(B)]^{-1} P_i^\alpha \Gamma^{(\alpha)}(B) = \sum_{A \in C_i} [\Gamma^{(\alpha)}(B)]^{-1} \Gamma^{(\alpha)}(A) \Gamma^{(\alpha)}(B) = \sum_{A \in C_i} \Gamma^{(\alpha)}(B^{-1}AB) = \sum_{A \in C_i} \Gamma^{(\alpha)}(A) \quad \dots (21)$$

Note IV. Conjugate elements and classes. If A, B and C be elements of a group such that $A^{-1}BA = C$, then B and C are said to be conjugate elements and the operation is said to be a similarity transformation of B by A . Clearly $ACA^{-1} = B$. $\dots (22)$

If we split the elements of a group into sets such that all the elements of a set are conjugate to each other, but no two elements of different sets are conjugate to each other, then such sets of elements are said to be the conjugacy class or simply classes of a group.

How to prepare a character table for a group say C_{3v} .

1. According to the preceding note, we conclude that:

(E) Constitutes a class by itself in any group, since for any element A of the group, $A^{-1}EA = E$.

Also (C_3, C_3^2) is a class of C_{3v} since $m_1^{-1} C_3 m_1 = C_3^2$ as $m_1^{-1} = m_1$ renders, $(m_1^{-1} C_3) m_1 = (m_1 C_3) m_1 = m_3 m_1 = C_3^2$ from the table.

Similarly (m_1, m_2, m_3) is a class of C_{3v} .

Hence the classes of C_{3v} are (E) , (C_3, C_3^2) , (m_1, m_2, m_3) say, $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3$ respectively. Then $\mathcal{C}_1 = (E)$; $\mathcal{C}_2 = (C_3, C_3^2) = 2C_3$ (say) and $\mathcal{C}_3 = (m_1, m_2, m_3) = 3\sigma_v$ (say).

2. Since C_{3v} has three classes, it must have three irreducible representations say $\Gamma^{(1)}, \Gamma^{(2)}, \Gamma^{(3)}$ whose dimensions may be taken as l_1, l_2, l_3 respectively.

In view of (17), here $g=6$ since C_{3v} consists of 6 elements,
 $\therefore I_1^2 + I_2^2 + I_3^2 = 6$.

With integral I_i , the only possible solution is that two of I_i 's are unity while the remaining one is 2 say $I_1 = I_2 = 1$ and $I_3 = 2$.

3. In view of (16), the first row is obtained by writing unity for character of each class. The first column is obtained in view of the facts that the matrix of E in any representation is a unit matrix having its trace or character as I_i which is the dimension of the representation viz. $I_1 = 1 = I_2$, $I_3 = 2$.

4. The character table for C_{3v} .

Characters ↓		Classes →		
		E_1 (E)	E_2 (C_3, C_3^2) i.e. $2C_3$	E_3 (m_1, m_2, m_3) i.e. $3\sigma_v$
$\Gamma^{(1)} \rightarrow$	$\chi^{(1)}$	1	1	1
$\Gamma^{(2)} \rightarrow$	$\chi^{(2)}$	1	1	-1
$\Gamma^{(3)} \rightarrow$	$\chi^{(3)}$	2	-1	0

Explanation. The Character being identical with one-dimensional representation, for $\Gamma^{(1)}$ and $\Gamma^{(2)}$ the characters themselves must satisfy the multiplication table so that for elements whose square equals E such as m_1, m_2, m_3 the only allowed characters are ± 1 .

Now the elements in the same class having the same character as the character of (C_3, C_3^2) in $\Gamma^{(2)}$ can be determined.

Also since for C_3, C_3^2 with $(C_3^2)^3 = (C_3^2)^3 = E$, the one-dimensional representation could be the power of $i = \sqrt{-1}$, for, if n is the order of an element A so that $A^n = E$, then its only one-dimensional representation can be powers of $e^{2\pi i/n}$ as these are only numbers whose n th power yields unity because such numbers are unitary as their inverses are equal to their complex conjugate. Again since $(C_3)^2 = (C_3^2)^3 = C_3^2$, $\chi(C_3)$ and $\chi(C_3^2)$ can only be ± 1 for $\Gamma^{(2)}$. Now the normalization condition (21) can only be achieved by taking $+1$ in $\Gamma^{(2)}$. Finally the third row is determined by using the orthogonality relations (18) for the columns.




Fig. C.8

[2] Symmetry Transformations of a Square.

Let A, B, C, D be the vertices of a square with E, F, G, H as mid-points of sides DA, AB, BC and CD respectively and O the centroid. Take OG as x -axis, OF as y -axis and an axis through O normal to the plane of the square as z -axis. Then the eight symmetry operations of the square are as follows:

APPENDICES




Fig. C.9

$E \rightarrow$ identity operation




Fig. C.10

$C_4 \rightarrow$ clockwise rotation through 90° about z -axis.




Fig. C.11

$C_4^2 \rightarrow$ clockwise rotation through 180° about z -axis




Fig. C.12

$C_4^3 \rightarrow$ clockwise rotation through 270° about z -axis




Fig. C.13

$m_x \rightarrow$ reflection in the vertical plane passing through EG i.e. x -axis




Fig. C.14

$m_y \rightarrow$ reflection in the vertical plane passing through FH i.e. y -axis




Fig. C.15

$\sigma_v \rightarrow$ reflection in the vertical plane passing through AC




Fig. C.16

$\sigma_v' \rightarrow$ reflection in the vertical plane passing through BD

It is easy to see that $\sigma_u C_4 = m_x$ etc.

Also $(C_4)^{-1} = C_4^3$ i.e. $C_4 C_4^3 = C_4^3 C_4 = E$ i.e. C_4 is the inverse of C_4^3 etc.

We also have $C_4 m_x = \sigma_u$, $\sigma_v C_4^3 = m_x$, $\sigma_u \sigma_v = C_4^2$ etc.

Group multiplication table for C_{4v} .

Second operation ↓	First operation →							
	E	C_4	C_4^2	C_4^3	m_x	m_y	σ_u	σ_v
E	E	C_4	C_4^2	C_4^3	m_x	m_y	σ_u	σ_v
C_4^3	C_4^3	E	C_4	C_4^2	σ_v	σ_u	m_x	m_y
C_4^2	C_4^2	C_4^3	E	C_4	m_y	m_x	σ_v	σ_u
C_4	C_4	C_4^2	C_4^3	E	σ_u	σ_v	m_y	m_x
m_x	m_x	σ_v	m_y	σ_u	E	C_4^2	C_4^3	C_4
m_y	m_y	σ_u	m_x	σ_v	C_4^2	E	C_4	C_4^3
σ_u	σ_u	m_x	σ_v	m_y	C_4	C_4^3	E	C_4^2
σ_v	σ_v	m_y	σ_u	m_x	C_4^3	C_4	C_4^2	E

In view of (22), the five classes of C_{4v} are

(E) , $(C_4)^2$, (C_4, C_3) , (m_x, m_y) , (σ_u, σ_v)

(since $C_4^{-1} m_x C_4 = m_y$ as $(C_4^{-1} m_x) C_4 = C_4^3 m_x C_4 = \sigma_u C_4 = m_y$ etc.)

These classes can also be justified by the following simple rules:

1. Since rotations through different angles must be put in different classes, C_4 and C_4^2 belong to different classes.
2. Since rotations through an angle in clockwise and anticlockwise sense about an axis belong to a class only when there exists a transformation in the group, such that it reverses the sense of the coordinate system, therefore C_4 and C_4^3 are put in the same class, for, m_x or σ_u operation changes the sense of the coordinate system.
3. Since rotations through the same angle about two different axes or reflection in two different planes belong to the same class only when there exists some element of group, which is capable of bringing the two axes or two planes into each other, therefore m_x, m_y belong to the same class but σ_u, m_x do not belong to the same class, for in the former case C_u can bring the line EF into the line FH .

The character table for C_{4v}

Characters ↓	classes →				
	$\mathcal{C}_1$ (E)	$\mathcal{C}_2$ (C_4^2)	$\mathcal{C}_3$ (C_4, C_4^3) i.e., $2 C_4$	$\mathcal{C}_4$ (m_x, m_y) i.e., $2m$	$\mathcal{C}_5$ (σ_u, σ_v) i.e., 2σ
$\Gamma^{(1)} \rightarrow \chi^{(1)}$	1	1	1	1	1
$\Gamma^{(2)} \rightarrow \chi^{(2)}$	1	1	1	-1	-1
$\Gamma^{(3)} \rightarrow \chi^{(3)}$	1	1	-1	1	-1
$\Gamma^{(4)} \rightarrow \chi^{(4)}$	1	1	-1	-1	1
$\Gamma^{(5)} \rightarrow \chi^{(5)}$	2	-2	0	0	0

Explanation. Since C_{4v} has five classes, it must have five irreducible representations (by 17(a) say $\Gamma^{(1)}, \Gamma^{(2)}, \Gamma^{(3)}, \Gamma^{(4)}, \Gamma^{(5)}$, whose dimensions are l_1, l_2, l_3, l_4, l_5 respectively, such that by (17)

$$l_1^2 + l_2^2 + l_3^2 + l_4^2 + l_5^2 = 8,$$

since C_{4v} consists of 8 elements i.e. $g=8$

with integral l_i , the only possible solution is that $l_1 = l_2 = l_3 = l_4 = 1$ and $l_5 = 2$. Now in view of (16), the first row is obtained by writing unity for character of each class and the first column is obtained by the fact that the matrix of E in any representation is a unit matrix having its trace or character as l_i , the dimension of representation viz $l_1 = l_2 = l_3 = l_4 = 1, l_5 = 2$.

Now the character being identical with one dimensional representation for $\Gamma^{(1)}$ through $\Gamma^{(4)}$ the characters must themselves satisfy the multiplication table so that for elements whose square equals E such as C_4^2, m_x, σ_u , the characters to be allowed are ± 1 . But since $m_x m_y = C_4^2$ or $\sigma_u \sigma_v = C_4^2$, therefore m_x and m_y both are represented by $+1$ or -1 , because elements in the same class have the same characters and hence the characters of C_4^2 in $\Gamma^{(2)}, \Gamma^{(3)}, \Gamma^{(4)}$ are determined. Also since for C_4 and C_4^3 , with $(C_4)^4 = (C_4^3)^4 = E$, the one-dimensional representation could be the power of $i = \sqrt{-1}$. Further $(C_4)^2 = (C_4^3)^2 = C_4^2$ imply that $\phi(C_4^3)$ can only be ± 1 for $\Gamma^{(2)}, \Gamma^{(3)}, \Gamma^{(4)}$. Now the normalization condition (21) can be achieved by taking $+1$ for one of the classes $\mathcal{C}_3, \mathcal{C}_4, \mathcal{C}_5$ in representations $\Gamma^{(2)}, \Gamma^{(3)}, \Gamma^{(4)}$ and -1 for the remaining ones. We thus determine characters for the first four representations and the fifth one can be obtained in view of (18) for the columns.

Character tables for some Symmetry Point Groups

C_1	E
	1

C_2	E	C_2
	1	1
	1	-1

C_{2v}	E	C_2	σ_v	σ_v
	1	1	1	1
	1	1	-1	-1
	1	-1	1	-1
	1	-1	-1	1

C_3	E	C_3	C_3^2	$\omega = e^{2\pi i/3}$
	1	1	1	
	1	ω	ω^2	
	1	ω^2	ω	

C_4	E	C_4	C_4^2	C_4^3
	1	1	1	1
	1	-1	1	-1
	1	-i	-1	i
	1	i	-1	-i

D_2	E	$C_2(x)$	$C_2(y)$	$C_2(z)$
	1	1	1	1
	1	1	-1	-1
	1	-1	1	-1
	1	-1	-1	1

D_3	E	$2C_3$	$3C_2'$
	1	1	1
	1	1	-1
	2	-1	0

D_4	E	C_4^2	$2C_4$	$2C_2'$	$2C_2''$
	1	1	1	1	1
	1	1	1	-1	-1
	1	1	-1	1	-1
	1	1	-1	-1	1
	2	-2	0	0	0

INDEX

- Absolute
 motion, 1027
 rest, 1027
 space, 1027
 time, 1027
 Addition modula m , 385
 Adjoint of an operator, 440
 Ampere's circuital relation, 1020
 Analytic continuation, 566
 function, 474, 477
 Angular velocity, 48
 Approximate integration, 653
 Arc
 continuous, 497
 Jordan, 497
 regular, 497
 Argand diagram, 451
 plane, 451
 Arithmetic mean, 1092
 Associated function
 first kind, 711
 second kind, 711
 Associated polynomials
 Laguerre, 777
 Legendre, 707
 Augmented matrix, 259
 Ausdehnungslehre, 183
 Automorphism, 411
 inner, 416
 outer, 416
 Axiom of extension, 368
 pairing, 369
 power set, 369
 specification, 370
 Basis, 270
 linear space, 434
 Baye's theorem, 1083
 Bei functions, 727
 Ber functions, 727
 Bernoulli's theorem, 1145
 Bessel's equation, 714
 formula, 650
 functions 758, 878
 functions of 1st kind, 920
 functions of 2nd kind, 921
 modified functions, 727
 Beta coefficient, 1126
 function, 577
 Bianchi's identity, 357
 Binary operations, 379
 associative, 379
 commutative, 379
 Binomial theorem, 1086
 Borel field, 1074
 Boundary point, 474
 Bromwich contour, 913
 Canonical matrix, 241
 Cauchy's
 integral formula, 505
 residue theorem, 522
 theorem, 499
 Cauchy-Riemann equations, 478
 polar form, 480
 Cayley-Hamilton theorem, 268, 279
 Cayley's theorem, 414, 446
 Central difference formula, 648
 Character, 452
 compound, 1222
 primitive, 1222
 simple, 1222
 tables, 452, 1220
 Characteristic
 equation, 268
 frequencies, 1002
 functions, 995, 1147
 matrix, 268
 polynomials, 268
 roots, 277
 values, 277, 995, 1002
 vectors, 277
 Christoffel's symbols, 335, 341
 Circulation, 125
 Circular harmonics, 965
 motion, 47
 Class of the group, 418
 Clock paradox, 1038
 Closure property, 6
 Coefficient matrix, 259
 Column matrix, 209
 rank, 270
 vector, 209
 Combination of vectors, 19
 Complex, 421
 argument, 451
 differential operators, 494
 equality, 458
 modulus, 451, 458
 numbers, 458
 Composition table, 387
 Concurrent forces, 45
 Condition for orthogonality, 108
 Conditional probability, 1077
 Conduction current, 1015

Conditions

- boundary, 603
- initial, 603
- Dirichlet, 795
- Confluent hypergeometric equation, 753
- function, 755
- Congruence, 292
- Congruent transformation, 292
- Conjugacy, 416
- class, 1225
- Conjugate elements, 416, 1225
- functions, 482
- Conservative field, 127
- Consistent equations, 263
- Constant of separation, 631
- Continuity, 796
- Contour integration, 522
- indented, 530
- Contravariant tensor, 307, 308
- vector, 306
- Convolution properly, 890
- theorem, 848
- Correlation, 1188
- Cosets, 397
- left, 398
- right, 398
- Covariant
- curvature tensor, 355
- tensor, 307
- vector, 306
- Cramer's rule, 261
- Cross cut, 503
- Cross product, 22, 24
- characteristics, 24
- Curl, 85, 113, 361
- of sum, 86
- of product, 87
- Curvature tensor, 353
- Curvilinear coordinates, 106
- Cylindrical coordinates, 115
- functions, 758, 920
- harmonic, 920
- D'Alembert's method, 630, 995
- Deciles, 1102
- Derogatory matrix, 244
- Diagonalization method, 288
- Differentiation
- operators, 494
- partial, 66
- total, 65
- Differentiability, 476
- Difference table, 633
- Diffusion equation, 943
- one dimensional, 945
- two dimensional, 955
- three dimensional, 959
- diffusivity, 943
- Dimensions linear space, 435
- representation, 448
- Dirac-delta function, 784

- Direction cosines, 11
- Directional derivatives, 70, 71, 72
- Dirichlet's conditions, 795
- Discontinuities, 796
- first kind, 796
- infinite, 796
- mixed, 796
- ordinary, 796
- removable, 796
- second kind, 796
- Displacement current, 1015, 1016
- Distributions
- binomial, 1149
- casual, 1179
- Cauch's, 1184
- continuous, 1136
- geometric, 1181, 1182
- hypergeometric, 1182
- multinomial, 1184
- negative binomial, 1181
- normal, 1158
- Pascal, 1181
- Poisson, 1174
- Polya's, 1181
- probability, 1136
- sampling, 1197
- theoretical, 1149
- Divergence, 78, 112, 361
- of sum, 80
- of product, 81
- Divided differences, 642
- Domain, 475
- closed, 475
- open, 475
- Dot product, 22
- characteristics, 23
- Dummy suffix, 304
- Dyadic, 302
- Eigen frequencies, 1002
- functions, 995
- roots, 277
- values, 277, 995, 1002
- Vectors, 277
- Electro-magnetic induction law, 1020
- Elementary functions, 489
- exponential, 489
- inverse trigonometric, 492
- logarithmic, 491
- trigonometric, 490
- Elliptic integrals, 750
- Equations
- differential, 602
- diffusion, 628
- elliptic, 628
- Fourier's, 628
- homogeneous, 624
- hyperbolic, 628
- Laplace's, 628
- non-homogeneous, 626
- ordinary, 603
- parabolic, 628

INDEX

- partial, 616
- wave, 629
- Equivalence, 239
- of matrices, 239
- Error
- absolute, 1205
- coarse, 1202
- functions, 597, 757, 880, 1218
- function (asymptotic expansion), 1218
- Gaussian law, 1205
- mean square, 1205
- random, 1202
- statistical, 1202
- systematic, 1202
- theory, 1202
- Ether displacement current, 1016
- Everett's formula, 652
- E-waves, 990
- Expectation, 1126, 1140
- Extension of rank, 322
- Extrapolation, 636
- Faltung theorem, 848
- Faraday, 183
- Fermat's theorem, 401
- Fields
- lameller, 179
- scalar, 69
- solenoidal, 179
- vector, 69, 179
- Final value theorem, 874
- Finite differences, 632
- Fourier
- constants, 793
- equation, 943
- expansion, 792
- integral, 821
- inversion formula, 1148
- series, 793
- transform, 836
- Fresnel's Coefficient of Drag, 1041
- integral, 757
- Frobenius method, 674
- Functionals, 439
- linear, 439
- Functions, 373
- analytic, 477
- beta, 577
- bijective, 375
- complex, 374
- conjugate, 482
- complementary, 603
- domain, 373
- elementary, 602
- epimorphism, 374
- explicit, 602
- extension, 374
- gamma, 577
- harmonic, 482
- holomorphic, 477
- implicit, 602
- into, 374
- injective, 374
- inverse, 375
- many one, 374
- monogesis, 477
- monomorphism, 374
- onto, 374
- one-one, 374
- range, 373
- real, 374
- restriction, 374
- surjective, 374
- scalar, 70
- vector, 70
- Fundamental tensors, 327
- velocity, 1041
- Galilean transformations, 1027, 1028
- Gamma
- coefficient, 1126
- function, 577, 878
- graph, 580
- transformation, 579
- Gauss
- backward formula, 648
- divergence theorem, 146, 148, 150
- equation, 739
- forward formula, 647
- formula, 745
- Pi function, 578
- theorem, 150, 187, 745, 1020
- Generalized Fourier-Bessel Series, 933, 1006
- Generating functions, 690, 716
- Geodesic, 338
- equation, 340
- Geometric mean, 1097
- Goursat lemma, 500
- Gradient, 73, 74, 75, 111, 361
- Grassman, 183
- Green's formula, 163, 983
- functions, 982, 984
- identity, 151
- theorem, 159, 161
- Groups, 381
- abelian, 381
- alternating, 407, 419
- automorphism, 412
- commutative, 381
- centre, 397
- cosets, 397
- cyclic, 401
- class, 418
- continuous, 442
- diedral, 432
- endo-morphism, 413
- finite, 382
- factor, 420
- homomorphism, 410
- Hamiltonian, 441
- infinite, 382
- improper, 385

- isomorphism, 410
- isometries, 427
- left coset, 398
- mixed, 441
- order, 382
- proper, 385
- per mutation, 407
- point, 444
- p-primary, 441
- p-prufer, 441
- quotient, 420
- quaternion, 445
- right coset, 398
- regular permutation, 413
- rotation (n -D), 443
- semi, 384
- sub, 385
- simple, 441
- table, 386
- transference, 413
- torsion, 441
- torsion free, 441
- unitary, 441
- Hamilton, W.R., 183
- Hankel function, 727
 - transforms, 921, 922
- Harmonic functions, 482, 978
 - mean, 1099
- Harmonics, 668
 - cylindrical, 670
 - spherical, 669
 - solid, 669
 - surface, 669
 - zonal, 668
- Heaviside unit function, 881
 - expansion-formula, 896
- Helmholtz equation, 983, 988
 - theorem, 180
- Hermite equation, 759
 - functions, 765
 - generating function, 764
 - polynomials, 761
- Hermitian forms, 275
- Homomorphism, 410
 - Canonical, 425
 - natural, 425
- H -waves, 990
- Hypergeometric equations, 698, 739
 - functions, 742
- Identity
 - additive, 194
 - multiplicative, 194
 - permutation, 405
 - Inconsistent equations, 263
 - Independent events, 1085
- Index, 294
- Inertial systems, 1026
- Initial value theorem, 874
- Inner product, 197
 - vector space, 440
- Interdecile range, 1108
- Interquartile range, 1103, 1108
- Interpolation, 636
- Interior point, 474
- Integration of vectors, 121
- Integrals
 - complete, 616
 - Eulerian, 597
 - infinite, 584
 - improper, 584
 - particular, 602, 616
 - Riemann, 584
 - singular, 617
 - transforms, 856
- Invariant tensors, 314
- Inverse
 - additive, 7, 194
 - multiplicative, 7
- Inverses 238
- Inversion theorem, 911
- Irrational vector function, 177
- Isomorphism, 438
- Jordan's
 - inequality, 531
 - lemma, 530
- Ker, Kei functions, 729
- Kernel, 421, 856
- Kronecker delta, 304, 703
 - symbol, 318
- Kummer's theorem, 746
- Lagrange's
 - formula, 645
 - method, 617
 - subsidiary equation, 617
 - theorem, 400
- Laguerre
 - equation, 770
 - polynomial, 771
 - poly (integral prop.), 774
 - recurrence relation, 772
- Lameller field, 179
- Lami's theorem, 45
- Laplace's
 - equation, 165, 182, 482, 668, 959, 960
 - equation, (2-D), 960
 - equation, (3-D), 970, 972
 - integral, 693, 694
 - transform, 856
- Laplacian operator, 82, 114, 187, 361
- Laplace-Everett formula, 651
- Laurent's expansion, 511
- Law
 - associative, 5, 6, 193, 199, 372
 - cancellation, 194
 - commutative, 4, 6, 193, 198, 372
 - De Morgan, 372
 - distributive, 7, 193, 199, 373
 - idempotent, 371